Linking the *Salmonella enterica* 1,2-propanediol utilization bacterial

microcompartment shell to the enzymatic core via the shell protein PduB

Nolan W. Kennedy[a], Carolyn E. Mills[b], Charlotte H. Abrahamson[b], Andre Archer[c],

Michael C. Jewett[b,d], Niall M. Mangan[c,d], Danielle Tullman-Ercek[b,d,#]

[a]Interdisciplinary Biological Sciences Program, Northwestern University, Evanston,

Illinois, USA

[b]Department of Chemical and Biological Engineering, Northwestern University,

Evanston, Illinois, USA

[c]Department of Engineering Sciences and Applied Mathematics, Northwestern

University, Evanston, Illinois, USA

[d]Center for Synthetic Biology, Northwestern University, Evanston, Illinois, USA

Running Title: PduB links the microcompartment shell and core

#Address correspondence to Danielle Tullman-Ercek, ercek@northwestern.edu

Nolan W. Kennedy and Carolyn E. Mills contributed equally to this work. Author order

was determined via a best-of-three arm wrestling match.

Abstract word count: 250

Importance word count: 110

Manuscript word count: 4711

Abstract

Bacterial microcompartments (MCPs) are protein-based organelles that house the enzymatic machinery for metabolism of niche carbon sources, allowing enteric pathogens to outcompete native microbiota during host colonization. While much progress has been made toward understanding MCP biogenesis, questions still remain regarding the mechanism by which core MCP enzymes are enveloped within the MCP protein shell. Here we explore the hypothesis that the shell protein PduB is responsible for linking the shell of the 1,2-propanediol utilization (Pdu) MCP from *Salmonella enterica* serovar Typhimurium LT2 to its enzymatic core. Using fluorescent reporters, we demonstrate that all members of the Pdu enzymatic core are encapsulated in Pdu MCPs. We also demonstrate that PduB is the sole protein responsible for linking the entire Pdu enzyme core to the MCP shell. Using MCP purifications, transmission electron microscopy, and fluorescence microscopy we find that shell assembly can be decoupled from the enzymatic core, as apparently empty MCPs are formed in *Salmonella* strains lacking PduB. Mutagenesis studies also reveal that PduB is incorporated into the Pdu MCP shell via a conserved, lysine-mediated hydrogen bonding mechanism. Finally, growth assays and systems-level pathway modeling reveal that unencapsulated pathway performance is strongly impacted by enzyme concentration, highlighting the importance of minimizing polar effects when conducting these functional assays. Together, these results provide insight into the mechanism of enzyme encapsulation within Pdu MCPs and demonstrate that the process of enzyme encapsulation and shell assembly are separate processes in this system, a finding that will aid future efforts to understand MCP biogenesis.

Importance

MCPs are unique, genetically encoded organelles used by many bacteria to survive in resource-limited environments. There is significant interest in understanding the biogenesis and function of these organelles, both as potential antibiotic targets in enteric pathogens and also as useful tools for overcoming metabolic engineering bottlenecks. However, the mechanism by which these organelles are formed natively is still not completely understood. Here we provide evidence of a potential mechanism in *S. enterica* by which a single protein, PduB, links the MCP shell and metabolic core. This finding is critical for those seeking to disrupt MCPs during pathogenic infections or for those seeking to harness MCPs as nanobioreactors in industrial settings.

Introduction

Subcellular compartmentalization into organelles is an important feature of cellular life (1). Organelles allow cells to organize chemical transformations in space, enabling diverse metabolic processes to occur simultaneously within single cells. This functionality, sequestering diverse processes into membrane delimited organelles, is most frequently associated with the eukaryotic domain (1). However, prokaryotes have diverse modes of subcellular compartmentalization as well, including both lipid (2) and protein (3) bound organelles.

A prime example of a protein-delimited organelle is the bacterial microcompartment (MCP) (4). These protein-based organelles come in numerous varieties, including carbon-fixing carboxysomes found in cyanobacteria and carbon-harvesting metabolosomes often found in enteric pathogens (Figure 1A) (5, 6). Metabolosomes consist of a proteinaceous shell surrounding an encapsulated enzymatic core (Figure 1B) (6). Typically the enzymatic core consists of a specific metabolic pathway that utilizes a niche carbon source (Figure 1B), often one that must move through a toxic aldehyde intermediate (Figure 1B) (7). The MCP shell is thought to provide a diffusional barrier between the cytosol and enzymatic core, which both helps to increase flux through the encapsulated pathway and also protects the cell from the toxic intermediate (8, 9). In this way, MCPs are thought to be important for providing a growth advantage to pathogenic microbes during propagation within hosts (10).

One of the most well-studied types of metabolosomes is the 1,2-propanediol utilization (Pdu) MCP found in *Salmonella enterica* serovar Typhimurium LT2 (hereafter referred to as "LT2"). The Pdu MCP provides an excellent model for metabolosome

study as Pdu MCPs natively exist in a number of model bacterial hosts such as the

aforementioned LT2 as well as strains of *Escherichia coli* (11). Pdu MCPs are broadly

distributed throughout many bacterial phyla and numerous Pdu subtypes exist (12–14).

The LT2 Pdu MCP is also a valuable model metabolosome because it contains

representative shell proteins of all but one of the known shell protein subtypes (Figure

1C), making Pdu-based studies broadly applicable to a variety of MCP systems (13).

Generally, Pdu MCP shells are irregular and polyhedral in geometry and roughly

140 nm in diameter (15). They are composed of small, assembled shell proteins which

tile together around the enzymatic core into a perforated, mosaic surface, containing

small pores to allow metabolite diffusion (Figure 1B-C) (16). The assembled shell

proteins come in a number of basic archetypal categories (Figure 1C) (13). These

varying architectures include six-sided hexameric or pseudohexameric proteins

composed of pfam00936 domains, as well as five-sided pentameric proteins (BMC-P or

BMV) composed of pfam03319 domains (Figure 1C) (17, 18). The hexagonal

pfam00936 proteins can be further subdivided into simple hexamers (BMC-H), circularly

permutated hexamers (BMC-H^P), pseudohexameric trimers (BMC-T), circularly

permutated pseudohexameric trimers (BMC-T^P), and circularly permutated

pseudohexameric trimer dimers (BMC-T^{DP}) (Figure 1C) (13). The LT2 Pdu operon

contains eight total shell proteins, with representatives from all but the BMC-T^{DP}

subcategory (19).

Although tremendous progress has been made toward understanding the

biogenesis and assembly of the Pdu MCP, questions remain regarding the mechanism

linking the shell and the enzymatic core. Numerous studies exist throughout the

literature proposing nearly every single Pdu shell protein as responsible for

encapsulating one or more of the enzymes that localize to the MCP core (Figure 2A)

(20–24). In this study we expand on the hypothesis set forth by Lehman et al., which

posited that the BMC-T[P] protein PduB is responsible enzyme encapsulation in Pdu

MCPs (20). To build on this study, we created a suite of fluorescently-tagged enzyme

reporters which report on the encapsulation state of all enzyme interaction partners (i.e.

at least one enzyme from sets of enzymes that are known to interact) (Figure 2B). This

reveals that PduB is responsible for encapsulating all members of the enzymatic core.

Surprisingly, we demonstrate that PduW, an enzyme previously thought to be cytosolic

(25), is also encapsulated within MCPs. We also show that signal sequences are

sufficient for targeting to the enzymatic core, indicating that signal sequences interact

with other members of the enzymatic core and not necessarily the MCP shell as

previously thought (21, 23). Interestingly, MCP purifications from strains with the PduB

open reading frame knocked out demonstrate that the MCP shell still assembles in the

absence of the enzymatic core. This provides the first conclusive evidence that shell

assembly and enzymatic core assembly are separate processes that can be decoupled.

Through mutagenesis, we show that PduB is incorporated into the MCP shell via a

conserved hydrogen bonding mechanism, similar to the mechanism for incorporation of

the BMC-H proteins PduA and PduJ (26–28). Finally, our work shows decoupling the

enzymatic core from the MCP shell results in a surprising growth phenotype, differing in

growth rate and metabolic profile from other disrupted MCPs. Together, these results

provide clarity to a field crowded with conflicting evidence regarding MCP enzyme

encapsulation and organelle biogenesis.

Results

Fluorescent reporters demonstrate PduB is critical for enzyme encapsulation

Results

Fluorescent reporters demonstrate PduB is critical for enzyme encapsulation

In 2017, Lehman et al. reported that deletion of the PduB N-terminus reduced the

apparent enzyme content of purified MCPs as indicated by sodium dodecyl sulphate-

polyacrylamide gel electrophoresis (SDS-PAGE). While groundbreaking, this study did

not directly demonstrate effects of PduB deletion on enzyme localization to MCPs, as

malformed MCPs or mistargeted enzymes are co-purified with MCPs in the differential

centrifugation process (27). To build on this landmark study, we created a suite of green

fluorescent protein (GFP) reporters to inform on which members of the enzymatic core

are directly impacted by the loss of PduB. Numerous studies have indicated that many

Pdu shell proteins directly bind and encapsulate various core enzymes. Thus, we

reasoned that PduB may only be responsible for binding and encapsulating a select

subset of Pdu enzymes. For example, studies have suggested that PduK (21), PduT

(22), PduA (23), PduJ (23), and PduU (24) are responsible for binding and targeting one

or more Pdu enzymes to the MCP core (Figure 2A). However, the work by Lehman et

al. is the only study to date to directly show that a genetic alteration (deletion of the

PduB N-terminus in the *pdu* operon) leads to changes in enzyme localization. We

therefore hypothesized that PduB is the most likely shell protein to be responsible for

enzyme targeting (20).

To test the hypothesis that PduB is responsible for linking the Pdu MCP shell and

enzymatic core, we created GFP protein fusions to report on each Pdu enzyme (Figure

1B, 2B). Some enzymes or sets of enzymes contain known, short, N-terminal signal

sequences that are sufficient for targeting enzymes to the MCP lumen. The signal

sequence of PduD (ssD) is sufficient for targeting PduC, PduD, and PduE (29); the

signal sequence for PduP (ssP) is sufficient for targeting PduP (30); and the signal

sequence for PduL (ssL) is sufficient for targeting PduL (31). Since these are the

minimal components necessary for targeting these proteins, they were fused to GFP

and used to report on the enzyme's change in encapsulation as a result of genetic

manipulations in this study (Figure 2B). Since PduD interacts with PduC and PduE, only

ssD-GFP was necessary to report on these three proteins. In the same way, since PduP

interacts with PduQ (32), ssP-GFP was used to report on this enzyme set (PduP and

PduQ). PduL has been hypothesized to interact with PduS (33), but this has not been

confirmed experimentally. However, PduS interacts with PduO (34), and since PduO is

the more abundant member of this pair (25), it was fused to GFP and used as a reporter

for PduS and PduO (O-GFP) (Figure 2B). Similarly, PduG and PduH are subunits of a

diol dehydratase reactivase enzyme and PduG is the more abundant member of this

protein pair, so it was fused with GFP (G-GFP) and used for this study (Figure 2B) (25,

35–37). Finally, PduW, which is thought to be cytosolic (25), was fused C-terminally with

GFP (W-GFP) to report on its encapsulation state (Figure 2B).

To test the hypothesis that PduB is necessary for the encapsulation of all

members of the enzymatic core listed above, each fluorescent reporter was expressed

in four strains of LT2: (1) wild type (WT) LT2, (2) a strain with the genes encoding the

BMC-H proteins PduA and PduJ knocked out (ΔA ΔJ), (3) a strain with the

transcriptional activator PocR knocked out (ΔpocR), and (4) an experimental strain with

PduB knocked out (ΔB) to inform on the effect of PduB loss on enzyme encapsulation

(Figure 3A). The WT strain serves as a positive control and enables visualization of

which enzymes are normally encapsulated within MCPs (Figure 3A). The ΔA ΔJ strain

serves as a negative control, as this strain is incapable of forming assembled MCPs but

still forms proto-MCP aggregates, which appear as polar bodies (Figure 3A) (27). The

ΔpocR strain serves as a negative control for MCP expression, as MCP proteins cannot

be expressed from the genome in the presence of the native inducer 1,2-propanediol

(1,2-PD) in this strain (Figure 3A) (27, 38–41). This allows us to determine which GFP-

tagged constructs formed aggregates when expressed in the absence of MCPs and to

what degree.

When each fluorescent reporter is expressed in the WT strain they are localized

to MCPs as indicated by the presence of bright fluorescent puncta (Figure 3B-C). This

assay has been used as a readout of protein localization to MCPs and has been

validated in numerous studies (27, 29, 30, 42). However, the relative encapsulation

level of each enzyme set has not been quantitatively assessed using this assay before.

Our results reveal that ssD-GFP and ssP-GFP are the most efficiently encapsulated,

which is expected based on quantitative mass spectrometry analyses of purified MCPs

(25). ssL-GFP is efficiently targeted to a relatively high number of MCPs, but puncta

appear much dimmer qualitatively relative to ssD-GFP and ssP-GFP, as expected

(Figure 3B). G-GFP and O-GFP are also encapsulated into MCPs, albeit less efficiently

than any of the signal sequence tagged reporters. We hypothesize that this is due to

their propensity to aggregate, as demonstrated by the formation of polar bodies in the

ΔpocR strain (Figure 3B-C). Surprisingly, W-GFP is targeted to MCPs as well, in spite of

the fact that it has been reported to be primarily cytosolic (Figure 3B-C) (25). It should

be noted that it is only targeted to MCPs in a subset of cells, as indicated by the bimodal

distribution in the MCP counts for this construct (Figure 3C). Together these results

demonstrate that all of the enzymatic members of the Pdu metabolic pathway are likely

targeted to some degree to MCPs, and that the GFP reporter constructs used here are

adequate reporters of this targeting.

When the fluorescent reporter constructs are co-expressed with MCPs in the ΔB

strain, only polar bodies are observed, similar to the ΔA ΔJ strain (Figure 3B-C).

Quantitatively, significantly fewer puncta are observed per cell for each fluorescent

reporter in both the ΔB and ΔA ΔJ strains compared with the WT strain. Significantly

more puncta are present in the WT strain than in the ΔpocR strain as well, as expected

(Figure 3C). This indicates that either MCP formation or enzyme encapsulation is

significantly impacted by the loss of PduB.

The PduB knockout strain forms assembled but empty MCPs

The encapsulation assay reveals that PduB loss dramatically alters the

encapsulation state of all fluorescently tagged enzymes tested. However, this assay

cannot distinguish between changes in encapsulation or changes in MCP formation. To

test the hypothesis that PduB impacts encapsulation but not assembly, MCPs were

purified from the ΔB strain using differential centrifugation (43, 44). When purified

samples are analyzed by SDS-PAGE, the ΔB strain appears dramatically different from

the standard MCP banding pattern apparent in the WT samples (Figure 4A). The ΔB

samples more closely resemble the ΔA ΔJ samples, which are known to contain

malformed MCP aggregates (Figure 4A-B) (27). An important distinction is that the ΔB

sample has high apparent levels of PduA and PduJ (Figure 4A).

Purified samples were then analyzed by transmission electron microscopy (TEM)

to determine if MCPs or partially formed aggregates were produced in the ΔB strain

(Figure 4B). To our surprise, the ΔB sample shows many apparently well-formed MCPs.

TEM micrographs from this sample contain structures with flat, distinguishable

boundaries that appear roughly polyhedral in morphology (Figure 4B). Interestingly, the

ΔB shells appear smaller than the WT shells (83 ± 21 nm vs 121 ± 19 nm, respectively

(mean ± standard deviation)) and have qualitatively less electron density in their center,

possibly indicating that they are empty (Figure 4B-C). The structures in the ΔB sample

are clearly distinct from the aggregates observed in the ΔA ΔJ sample and the

background structures present in the ΔpocR sample. These results, coupled with the

SDS-PAGE and encapsulation assay results, indicate that the ΔB strain likely forms

empty MCP shells as well as separate aggregates of Pdu enzymes.

To confirm this surprising finding, co-localization assays were performed in which

the shell protein PduA was fused with a fluorescent mCherry reporter protein and

expressed from the PduA locus in the WT, ΔB, and ΔpocR strain backgrounds. As

expected, PduA-mCherry is localized to MCPs as indicated by the presence of bright

fluorescent puncta (Figure 4D) (45). PduA is known to be important for shell formation

and is one of the most abundant members of the Pdu MCP shell (25, 27, 46). When

expressed in the WT background with the tagged GFP reporter constructs, GFP and

mCherry colocalize, indicating reporter targeting to MCPs, as expected (Figure 4D).

When expressed in the ΔpocR background, no mCherry puncta are present, as none of

the proteins in the *pdu* operon, including PduA-mCherry, are expressed in this control

(Figure 4D). In the ΔB strain background, mCherry puncta are still present, verifying the

finding that MCP shells were still formed in the absence of PduB. As expected based on

the results described in Figure 3, GFP reporter constructs are only found in polar bodies

in the ΔB strain (Figure 4D). These results conclusively demonstrate that PduB deletion

impacts protein encapsulation but not MCP shell formation, and that these are two,

distinct processes that can be decoupled upon PduB deletion. Furthermore, this

suggests that signal sequences are sufficient for targeting to the enzymatic core,

challenging the idea that primary Pdu MCP signal sequence function is to bind to MCP

shell proteins, enabling enzymatic core encapsulation (21, 23).

PduB is incorporated into MCPs by a conserved hydrogen bonding mechanism

While PduB appears to be important for linking the MCP shell and enzymatic

core, the mechanism by which PduB is incorporated into the shell (Figure 5A) is not

known. A groundbreaking structural study by Sutter et al. showed that BMC-T^{PD}

proteins similar to PduB appear to incorporate into the shell of a smaller, more regular,

empty MCP from *Haliangium ochraceum* via a lysine-lysine hydrogen bonding motif

(Figure 5B) (26, 47). In this motif, lysine side chains on adjacent BMC proteins (in this

case, a BMC-H protein and BMC-T^{PD} protein) arrange antiparallel to each other and

hydrogen bond with the backbone carbonyl oxygen of the adjacent lysine (Figure 5B)

(26, 47). Interestingly, analogous lysine residues (Figure 5C) are also important for the

self-assembly of BMC-H proteins PduA and PduJ, and are necessary for their ability to

drive shell assembly (27). For these reasons, we hypothesized that analogous lysine

residues at these positions in the BMC-T^{P} protein PduB (Figure 5C) are important for

PduB incorporation into the MCP shell, bridging the gap between shell and enzymatic

core.

To test this hypothesis, a double mutant version of PduB was cloned where the

lysine residues at position 102 and 207 were mutated to alanine residues. These

residues appear to be well-conserved among PduB homologs (26), so we predicted that

mutations to alanine should be disruptive. We expect that this protein is well-folded, as

analogous mutations in other BMC proteins increase solubility and are used frequently

in crystallographic studies (48, 49). The double mutant PduB (B-K102A-K207A) was

then placed at the *pduB* locus in the *pdu* operon in place of WT *pduB*, generating the

ΔB::B-K102A-K207A strain (Figure 3A). The encapsulation assay, in which each

fluorescent reporter is expressed simultaneously with MCPs, reveals that the ΔB::B-

K102A-K207A strain has a similar phenotype to both the ΔB and ΔA ΔJ strains, where

polar bodies are primarily observed rather than the scattered puncta indicative of MCPs

(Figure 3B-C). This indicates that the PduB double mutant is likely unable to incorporate

into MCP shells, effectively recapitulating the ΔB phenotype.

To confirm the conclusion based on the encapsulation assay, MCPs were

purified from the ΔB::B-K102A-K207A strain. Samples from this strain appear similar to

the ΔB strain by SDS-PAGE, where bands corresponding to enzymes are greatly

reduced, but bands for PduA and PduJ are still visible (Figure 4A). Purified structures

were then imaged by TEM, which reveals that apparently well-formed MCPs are present

and similar in size and morphology to the MCPs purified from the ΔB strain (Figure 4B-

C). The PduB double mutant strain was also grown with PduA-mCherry expressed from

the *pduA* locus, and this reveals bright puncta throughout the cytoplasm, similar to the

ΔB strain, indicating the formation of MCP shells (Figure 4D). As in the ΔB strain, GFP-

tagged cargo appears only in polar bodies, again indicating well-formed shells

decoupled from the enzymatic cargo (Figure 4D). Together, these results strongly

support the hypothesis that PduB is incorporated into the MCP shell by a similar

mechanism to other BMC proteins, and that disruption of the conserved lysine hydrogen

bonding network disrupts this interaction.

Loss of PduB impacts LT2 growth on 1,2-propanediol as a carbon source

Having established that PduB is essential for linking the core to the shell of the

Pdu MCP, we next explored how decoupling of the core and shell impacts the function

of the native Pdu pathway. To this end, we compared growth and metabolite profiles in

culture conditions that differentiate strains with well-formed MCPs and broken MCPs.

Specifically, these growth conditions encourage buildup of the toxic propionaldehyde

intermediate, leading to a growth lag in strains with broken MCPs that cannot sequester

this intermediate away from the cytosol (9, 16, 27). Under these growth conditions, we

hypothesized that our strains with decoupled core and shell assembly would exhibit a

growth phenotype similar to that of a strain expressing broken Pdu MCPs (ΔA ΔJ).

We tested this hypothesis by comparing the growth and metabolite profiles of

three control strains (WT, ΔpocR, ΔA ΔJ) to two experimental strains (ΔB, ΔB::B-

K102A-K207A). Our three control strains were the same as those used in the

encapsulation assay—WT, which harbors functional MCPs, ΔpocR, which cannot

express the *pdu* operon, and ΔA ΔJ, which forms proto-MCP aggregates but no shell

and thus serves as our broken MCP control. Our two experimental strains (ΔB, ΔB::B-

K102A-K207A), as described above, both have decoupled MCP core and shell

assembly. Our control strains behave as expected. The ΔpocR strain, which is unable to

express any Pdu pathway enzymes, does not grow nor does it consume 1,2-PD (Figure

6A, B). Our broken MCP control strain (ΔA ΔJ) initially grows and consumes 1,2-PD

faster than the MCP-containing strain (WT), as expected—the absence of a MCP shell

accelerates apparent pathway kinetics by providing more direct access to 1,2-PD

(Figure 6A, B). However, propionaldehyde build up in the ΔA ΔJ strain eventually leads

to a lag in growth from 12-30 hours (Figure 6A, B). During this timeframe, the WT strain

begins to outgrow the ΔA ΔJ strain, as the MCPs in the WT strain successfully

sequester the toxic propionaldehyde intermediate away from the cytosol (Figure 6A, B).

Surprisingly, we find that our two experimental strains (ΔB, ΔB::B-K102A-

K207A), both of which have decoupled MCP core and shell assembly, exhibit distinct

growth and metabolite profiles. The growth and metabolite profiles of the PduB double

mutant strain (ΔB::B-K102A-K207A) are similar to those of the broken MCP control (ΔA

ΔJ), in agreement with our initial hypothesis. This strain (ΔB::B-K102A-K207A) grows

rapidly at first, and then suffers a growth lag after propionaldehyde accumulates after 9

hours (Figure 6A, B). Interestingly, the growth lag is longer (9-40 hours) and the peak

propionaldehyde level is higher (2.5 ± 0.6 mM) in the ΔB::B-K102A-K207A strain than in

the ΔA ΔJ strain. More perplexing is the behavior of the ΔB strain—not only does this

strain exhibit minimal propionaldehyde buildup, it also eventually outgrows the WT

strain (Figure 6A, B). Correspondingly, the downstream metabolites propionate and 1-

propanol are consumed more rapidly in the ΔB strain than in other strains (Figure 6A,

B). Given that ΔB and ΔB::B-K102A-K207A strains exhibit essentially identical behavior

with respect to assembly phenotype (shown above), we turned to kinetic modeling to

generate hypotheses that could explain this observed discrepancy in growth behavior

and differences in propionaldehyde accumulation.

Using a systems-level kinetic model of the Pdu pathway modified from previous

work (8) (see Materials and Methods), we examined which features of the pathway or

polar body geometry had the strongest impact on propionaldehyde buildup in cells

containing polar bodies. We considered MCPs as a control case, modeling these MCPs

as spheres with a diffusive barrier at their surface that limits metabolite transport to the

enzymatic core. Polar bodies, then, are modeled as one large sphere with free diffusion

of metabolites at the surface (Figure 7A). The volume of this polar body sphere was set

to equal the total volume of MCPs in the cell (*i.e.* polar body volume = number of MCPs

per cell x volume of a single MCP). We used this polar body model to assess the

sensitivity of maximum propionaldehyde buildup to parameters affecting kinetics,

geometry and transport. Comparison of the MCP (Supplementary Figure S1) and polar

body (Supplementary Figure S2) models reveals that the maximum propionaldehyde

buildup in the polar body case is much more sensitive to the maximum velocity (V_{max}) of

the PduCDE enzyme than in the MCP case. As the concentration of PduCDE in the

polar body is decreased (leading to an equivalent fractional decrease in V_{max}), the

predicted propionaldehyde profile for the polar body approaches that of the MCP

(Figure 7A). Indeed, decreasing the PduCDE concentration in the polar bodies

decreases the maximum propionaldehyde concentration outside the cell (Figure 7B).

Specifically, we find that the maximum propionaldehyde concentration observed in the

polar body model matches that of the MCP model if the PduCDE concentration is

decreased by 76%, suggesting that variation in PduCDE concentration between strains

can impact propionaldehyde buildup as strongly as Pdu pathway encapsulation in

MCPs (Figure 7B). We thus hypothesized that the difference between our strains

containing polar bodies was the level of PduCDE expression. We hypothesized that this

was the result of polar effects in which different alterations to the *pduB* locus

differentially impact the expression of downstream *pduCDE* genes.

We tested our hypothesis that PduCDE expression differed across strains

containing polar bodies by measuring GFP expression at the *pduD* locus. Specifically,

we replaced the *pduD* gene at the *pduD* locus with a gene encoding for ssD-GFP in our

five growth strains (WT, ΔpocR, ΔA ΔJ, ΔB, ΔB::B-K102A-K207A) and used flow

cytometry to quantify GFP expression. We find that the ΔB strain has the lowest

expression at the *pduD* locus, with 49 ± 7% lower GFP fluorescence than the WT strain

($p < 0.05$). By contrast, the ΔB::B-K102A-K207A strain had higher apparent GFP

expression at the *pduD* locus than the WT strain ($p < 0.05$). Finally, we find that our

broken MCP control strain, ΔA ΔJ, has just slightly lower expression at the *pduD* locus

than WT ($p < 0.05$) (Figure 7C). Importantly, we note that increased GFP expression at

the *pduD* locus correlates with peak propionaldehyde level in the strains with

unencapsulated enzymes (Pearson's correlation coefficient = 0.92), supporting our

hypothesis that PduCDE expression causes differences in propionaldehyde buildup in

these strains.

Together, these results highlight two important considerations for future work in

the MCP field. First, modeling is a powerful tool for generating hypotheses and

exploring parameter space, enabling rapid identification of the key governing

parameters for a given subcellular organizational strategy. Second, polar effects can

impact downstream enzyme expression, which, in turn, can dramatically alter observed

pathway performance. Thus, quantifying these effects and minimizing them whenever

possible is essential when conducting genetic studies of MCP function.

Discussion

Metabolosomes are bacterial organelles confirmed to exist in a number of enteric

bacteria, where their expression and activation has been linked to pathogenesis (10). Of

the metabolosomes investigated to-date, the Pdu MCP is the best studied, and has

served as a model system for understanding the basics of metabolosome assembly and

metabolic function. However, even in this well-studied model system, a large body of

conflicting evidence exists regarding the mechanism by which enzymatic cargo is

loaded into the MCP shell (20–24).

Here, we used a comprehensive suite of fluorescent reporters to show that the

shell protein PduB is responsible for linking the enzymatic core to the MCP shell—when

PduB is absent or unable to incorporate into the shell, no Pdu enzymes are loaded into

the MCP. This interaction between PduB and the enzymatic core likely happens via the

N-terminus of PduB, as previously suggested in Lehman et al., since deletion of this

unique structural domain reduces enzyme loading into MCPs (20). Instead, enzymes

localize to proto-MCP aggregates, termed polar bodies, in the absence of PduB.

Importantly, we find that N-terminal signal sequence tags such as those identified for

PduD and PduP are sufficient for loading cargo to these proto-MCP aggregates (29,

30). This is significant as these signal sequence tags were previously thought to exist

primarily to link the shell and core of the MCP. We also report that although PduB is

required for cargo loading, it is not necessary for MCP shell assembly, as empty MCP

structures form in the absence of PduB. The fact that empty MCP shells and proto-MCP

aggregates form separately in the absence of PduB suggests that MCP shell and core

assembly are independent processes in the Pdu MCP system. Combined with previous

work demonstrating that a self-assembling proteins PduA or PduJ are required for Pdu

MCP formation (27), this discovery brings us one step closer to realizing the

components necessary to construct a minimal Pdu MCP. Specifically, two of the

requisite components are: (1) a hexameric protein with a strong propensity for self-

assembly (PduA or PduJ) and (2) a protein that links the core and shell by interacting

with both the enzymatic core and the self-assembling hexameric proteins via a

conserved hydrogen bonding mechanism. We expect that computational models of this

assembly process will provide insight into the specific roles that these different proteins

play in MCP assembly (46, 50).

We also investigated how LT2 growth on 1,2-propanediol was impacted by the

decoupling of MCP core and shell assembly. As expected, when enzymes are localized

to polar bodies, buildup of the toxic aldehyde intermediate is not well-controlled as it is

in MCPs. However, using a combination of systems-level kinetic modeling and

experiments, we also discovered that strains that contain polar bodies rather than MCPs

exhibit dramatically different growth profiles depending on PduCDE expression level. In

our strains, we believe the differences in PduCDE expression we observed are a

consequence of polar effects on gene expression, as the genes encoding for *pduCDE*

are immediately downstream of the *pduB* gene. Specifically, we find that while our *pduB*

knockout strain (ΔB) and our *pduB* double mutant strain (ΔB::B-K102A-K207A) both

result in a similar decoupling of MCP core and shell assembly, the strain with the more

dramatic genetic alteration at the *pduB* locus (ΔB, the knockout) decreases expression

at the *pduD* locus substantially, whereas the strain with minimal genetic alteration at the

pduB locus (ΔB::B-K102A-K207A) does not. This result emphasizes an important point

of concern for studies in this field—namely that polar effects on downstream gene

expression can confound interpretation of experiment results, and thus must be

considered. Indeed, it has been shown that alterations to the *pduL* locus, which

encodes for enzymatic cargo, can disrupt Pdu MCP shell assembly, likely by modifying

expression levels of downstream shell proteins like PduN (42). Conversely, we show

here that modifications to the *pduB* locus, which encodes for a shell protein, can alter

enzyme concentrations in the core, altering Pdu pathway performance. Thus, the

identification of point mutants that minimally disrupt operon sequence, but prevent

specific protein functions, will be essential in future MCP studies.

Materials and Methods

Plasmid and strain creation

Materials and Methods

Plasmid and strain creation

All plasmids used for this study were created using the Golden Gate cloning (51)

method into a Golden Gate-compatible pBAD33t parent vector (chloramphenicol

resistance, p15a origin of replication). All strains, plasmids, and primers are listed and

described in Supplemental Tables S1-3, respectively. For constructs in which a signal

sequence was appended to the N-terminus of GFPmut2, a BsaI cut site and the DNA

sequence encoding the signal sequence were added to GFP using PCR (see Table S3

for primer sequence information). For strains in which GFPmut2 was appended to the

C-terminus of a full enzyme, a two-piece Golden Gate reaction was carried out. This is

true for all GFP-enzyme fusions with the exception of PduG-GFP, which was cloned

using SacI and XbaI restriction sites. First, forward and reverse primers were used to

amplify the enzyme and GFPmut2 open reading frames. Each primer encoded

compatible sticky ends to enable ligation into the pBAD33t parent vector in the proper

orientation. A glycine-serine (GS) linker was also encoded between the enzyme and

GFPmut2 open reading frames. Golden Gate reactions were carried out using

FastDigest Eco31I (Thermo Fisher Scientific) and T4 DNA ligase (New England

Biolabs), transformed into *E. coli* DH10b cells, purified using Zyppy™ Plasmid Miniprep

kit (Zymo Research), and sequence verified by Sanger sequencing (Genewiz).

All modified LT2 strains used in this study are listed in Table S1 and were

generated using λ red recombineering as described in previous reports (27, 52). Briefly,

a marker encoding dual selection (*cat/sacB*) was PCR amplified containing flanking

homologous overhangs matching the target gene locus and inserted into the *pdu*

operon. The *cat/sacB* selectable marker was then either replaced with a gene of interest

or knocked out, leaving only the C-terminus of the target gene in order to reduce polar

effects on downstream gene expression. For the PduB-K102A-K207A double mutant

strain, QuikChange was used (with KOD Hot Start DNA Polymerase (Sigma-Aldrich)) to

perform site directed mutagenesis on the two lysine residues on a plasmid backbone

first before it was amplified and inserted into the *pdu* operon. Modified strains were

sequence confirmed using Sanger sequencing (Genewiz).

Fluorescence and phase contrast microscopy

Fluorescence and phase contrast micrographs were collected on Nikon Eclipse

Ni-U upright microscope, 100X oil immersion objective, and an Andor Clara digital

camera with NIS Elements Software (Nikon). GFP fluorescence micrographs were

collected using a C-FL Endow GFP HYQ bandpass filter and mCherry fluorescence

micrographs were collected using a C-FL Y-2E/C filter. Exposure times of 100 ms were

used for ssD-GFP and ssP-GFP, 300 ms for ssL-GFP, 400 ms for G-GFP, O-GFP, and

W-GFP, and 500 ms for mCherry. Cell culture samples were placed onto Fisherbrand^{TM}

frosted microscope slides and covered using 22 mm x 22 mm, #1.5 thickness cover

slips. All images were equally adjusted within experiments and sample type for

brightness and contrast using ImageJ (53). Puncta counts were collected on brightness

and contrast adjusted images.

Microcompartment expression and purification

MCP expression and purification was done as described in detail previously (43).

Briefly, overnight starter cultures were started from single colonies streaked out onto

Lysogeny Broth, Miller (LB-M, Thermo Fisher Scientific) agar plates. These 5 mL LB-M

cultures (with 34 μg/mL chloramphenicol added when appropriate) were grown in 24-

well blocks for 15-18 hours at 37°C and shaking at 225 RPM. Saturated starter cultures

were used to subculture expression cultures. If cultures were used for microscopy

experiments, cultures were subcultured 1:500 in 5 mL LB-M supplemented with 0.02%

(w/v) final concentration of L-(+)-arabinose to induce fluorescent reporter expression,

0.4% (v/v) final concentration 1,2-propanediol to induce MCP expression, and 34 μg/mL

chloramphenicol. Cultures were then grown for a minimum of 6 hours at 37°C, 225 RPM

before imaging. For cultures used for MCP purification, saturated overnight cultures

were diluted 1:1000 into 200 mL No Carbon Essential (NCE) media supplemented with

50 μM ferric citrate, 42 mM succinate as a carbon source, 1 mM magnesium sulfate,

and 55 mM 1,2-propanediol to induce MCP expression. The cultures were grown in 1 L

glass Erlenmeyer flasks at 37°C, 225 RPM until cultures reached an OD_{600} of at least 1.

MCPs were purified using an established differential centrifugation method described

previously (43). Briefly, cells were pelleted at 5,000 x g and lysed chemically using an

octylthioglucoside (OTG) solution. Cell lysate was clarified at 12,000 x g (4°C, 5

minutes) and MCPs were pelleted at 21,000 x g (4°C, 20 minutes). Purified MCPs were

stored in a buffered saline solution containing 50 mM Tris (pH 8.0), 50 mM potassium

chloride, 5 mM magnesium chloride, and 1% (v/v) 1,2-propanediol at 4°C. Cultures used

for flow cytometry were subcultured from overnights 1:1000 into 5 mL of the same NCE

media used for compartment purifications and grown for 16 hours at 37 °C at 225 RPM

prior to analysis.

Transmission electron microscopy

Transmission electron microscopy (TEM) on purified MCPs was done as

described in detail in a prior study (15). Samples containing purified MCPs were applied

and fixed onto 400 mesh Formvar-coated copper grids (EMS Cat# FF400-Cu). Fixation

was done using 2% (v/v) glutaraldehyde and staining was done using 1% (w/v) uranyl

acetate. Samples were then imaged using a JEOL 1230 transmission electron

microscope with a Gatan 831 bottom-mounted CCD camera. MCP diameter

measurements were collected using the Ferret diameter tool in ImageJ (53).

SDS-PAGE

For analysis of purified MCPs using SDS-PAGE, samples were first normalized

by protein concentration (measured by bicinchoninic acid assay) before loading onto the

15 wt% polyacrylamide Tris-glycine minigels. SDS-PAGE was then run a second time

with sample loading normalized by densitometry on the PduA band to correct for

potential contaminants within the samples. Samples were boiled at 95°C for 5 minutes

in Laemmli buffer and then immediately run at 120 V for 90 minutes. Gels were stained

with Coomassie Brilliant Blue R-250 and imaged using the Bio-Rad ChemiDoc XRS +

System.

Growth assays and metabolite analysis

Overnight starter cultures of LT2 strains of interest were started from a single

colony and grown in test tubes in 5 mL of Terrific Broth (Dot Scientific, Inc.) without

glycerol. Overnights were grown 15-16 h at 37 °C at 225 RPM with orbital shaking.

These overnights were subcultured into 50 mL No Carbon Essential (NCE) media

supplemented with 50 µM ferric citrate, 1 mM magnesium sulfate, 150 nM

adenosylcobalamin (Santa Cruz Biotechnology), and 55 mM 1,2-propanediol to an

OD_{600} of 0.05. Cultures were grown in 250 mL, unbaffled Erlenmeyer flasks at 37 °C,

225 rpm. At each time point, 500 µL of cell culture was taken for OD_{600} measurement

and HPLC analysis. OD_{600} measurements were taken using a BioTek Synergy HTX

multi-mode plate reader. After 9 hours, cultures from all strains except the ΔpocR strain

were diluted 1:5 in fresh NCE to ensure measurements were within the linear range of

the instrument. Error bars on growth curves represent standard deviation over three

biological replicates. The remainder of the cell culture sample (that not used for OD_{600}

measurement) was centrifuged at 13,000 x g for 5 minutes to pellet cells. The

supernatant was collected and frozen at -20 °C. After completion of the growth

experiment, supernatant samples were thawed and filtered (Corning™ Costar™ Spin-X

LC filters) in preparation for HPLC analysis. Samples were run on an Agilent 1260

HPLC system. Metabolites were separated using a Rezex™ ROA-Organic Acid H+

(8%) LC Column (Phenomenex) at 35 °C, in 5 mM sulfuric acid flowing at 0.4 mL/min.

Metabolites were detected with a refractive index detector (RID) as previously described

(9). Peak areas were calculated using the Agilent ChemLab software and converted to

metabolite concentrations using standards of the metabolites of interest (1,2-

propanediol, propionaldehyde, propionate, 1-propanol) at 200 mM, 100 mM, 50 mM, 20

mM, and 5 mM. Error bars on metabolite concentrations represent standard deviation

over three biological replicates.

Modelling

The kinetic model used here is modified from previous work, which described

MCP function in the cell using a reaction-diffusion framework (8). Modifications include

an assumption that the cytosol is well-mixed to produce a compartmental model and the

incorporation of the effects of cell growth. We analyze how metabolite profiles evolve

over time, while previous studies focused on steady state. A detailed description of the

model used in this work is provided here and in Supplemental Document 1.

We model cells as capsule-shaped (cylinders with a hemisphere at either end)

and allow passive transport of 1,2-propanediol, propionaldehyde, propionyl-CoA, 1-

propanol, and propionate at the cell surface, specified by the permeabilities included in

Supplemental Table S4. MCPs are modeled as spheres 140 nm in diameter (15 per

cell) that have a specified permeability at the surface; this permeability is assumed to be

the same for all metabolites. Polar bodies are modeled as a single sphere per cell 345

nm in diameter (selected such that the volume of the polar body is equal to that of 15

MCPs), and allow free diffusion of all metabolites at their surface. Recognizing that

diffusion in the MCP, cytosol, and media are much faster than the transport processes

and enzyme reaction rates, we assume that the concentration is uniform in each of

these locations.

We assume that conversion of 1,2-propanediol to propionaldehyde by PduCDE is

irreversible. Similarly, we assume that conversion of propionyl-CoA to propionate by

PduL and PduW occurs in a single step, and that this reaction is also irreversible.

Reverse reactions are included for conversion of propionaldehyde to 1-propanol and

propionyl-CoA by PduQ and PduP respectively. Reactions catalyzed by PduCDE,

PduP, and PduQ are assumed to occur only in the MCP or polar body volume, whereas

the reaction catalyzed by PduL/PduW is assumed to occur only in the cytosol. All

enzymes are assumed to have Michaelis-Menten kinetics.

Cell concentration and growth are calculated using the data from the growth

curves. The model was built and implemented in Python (54). All codes used are

available on GitHub at https://github.com/cemills/MCP-vs-PolarBody. A detailed

description of the equations used in the model are available in Supplemental Document

1. The parameters used for calculations presented here are available in Supplemental

Table S4. Metabolite profiles over time of 1,2-propanediol, propionaldehyde, propionate,

1-propanol for the PduCDE concentrations shown in Figure 7A can be found in

Supplementary Figure S3.

Flow cytometry

Prior to measurement, cells were diluted to OD_{600} = 0.03 into 200 µL phosphate-

buffered saline (PBS) supplemented with 2 mg/mL kanamycin to stop translation.

Samples were prepared in a U-bottomed 96-well plate and kept away from light until

measurement. 10,000 events were collected per sample on an Attune NxT acoustic

focusing cytometer paired with an Attune NxT auto sampler. Cell populations were

gated using forward and side scatter channels, and average reported fluorescence was

calculated using the geometric mean of the population. Values were converted to

molecules of equivalent fluorescence (MEF) using BD Sphero™ Rainbow Calibration

Particles (Fisher Scientific catalog number BDB559123) according to manufacturer's

instructions. Analysis was performed using FlowJo software (www.FlowJo.com).

Acknowledgements

The authors would like to thank and acknowledge members of the Tullman-Ercek and

Mangan groups for insightful advice and discussion related to designing experiments

associated with this study as well as preparing the manuscript. The authors would like

to specifically acknowledge Dr. Taylor M. Nichols, Dr. Eddy Kim, Dr. Chris Jakobson,

and Mike Vincent for strains, plasmids, and primers used in this study. This work was

funded in part by the Army Research Office (grant W911NF-19-1-0298 to DTE), and the

Department of Energy (grant DE-SC0019337 to DTE and NMM). NWK was funded by

the National Science Foundation Graduate Research Fellowship Program (grant DGE-

1842165), and by the National Institutes of Health Training Grant (T32GM008449) via

the Northwestern University Biotechnology Training Program.

CRediT Contributor Roles: **NWK**: Conceptualization, formal analysis, funding

acquisition, investigation, methodology, project administration, validation, visualization,

writing – original draft, writing – review & editing. **CEM**: Conceptualization, formal

analysis, investigation, methodology, project administration, software, validation,

visualization, writing – original draft, writing – review & editing. **CHA**: Conceptualization,

investigation, methodology, writing – review & editing. **AA**: Conceptualization, formal

analysis, software, writing – review and editing. **MCJ**: Funding acquisition, resources,

supervision, writing – review & editing. **NMM**: Conceptualization, funding acquisition,

resources, supervision, writing – review & editing. **DTE**: Conceptualization, funding

acquisition, resources, supervision, writing – review & editing.

References

1. Gabaldón T, Pittis AA. 2015. Origin and evolution of metabolic sub-cellular

compartmentalization in eukaryotes. Biochimie 119:262–268.

2. Uebe R, Schüler D. 2016. Magnetosome biogenesis in magnetotactic bacteria.

Nature Reviews Microbiology 14:621–637.

3. Greening C, Lithgow T. 2020. Formation and function of bacterial organelles. Nat

Rev Microbiol 18:677–689.

4. Kennedy NW, Mills CE, Nichols TM, Abrahamson CH, Tullman-Ercek D. 2021.

Bacterial microcompartments: tiny organelles with big potential. Current Opinion in

Microbiology 63:36–42.

5. Kerfeld CA, Melnicki MR. 2016. Assembly, function and evolution of cyanobacterial

carboxysomes. Current Opinion in Plant Biology 31:66–75.

6. Kerfeld CA, Aussignargues C, Zarzycki J, Cai F, Sutter M. 2018. Bacterial

microcompartments. Nature Reviews Microbiology 16:277–290.

7. Asija K, Sutter M, Kerfeld CA. 2021. A Survey of Bacterial Microcompartment

Distribution in the Human Microbiome. Frontiers in Microbiology 12:1090.

8. Jakobson CM, Tullman-Ercek D, Slininger MF, Mangan NM. 2017. A systems-level

model reveals that 1, 2-Propanediol utilization microcompartments enhance

pathway flux through intermediate sequestration. PLoS computational biology

13:e1005525.

9. Sampson EM, Bobik TA. 2008. Microcompartments for B12-Dependent 1,2-

Propanediol Degradation Provide Protection from DNA and Cellular Damage by a

Reactive Metabolic Intermediate. Journal of Bacteriology 190:2966–2971.

10. Prentice MB. 2021. Bacterial microcompartments and their role in pathogenicity.

Current Opinion in Microbiology 63:19–28.

11. Stewart KL, Stewart AM, Bobik TA. 2020. Prokaryotic Organelles: Bacterial

Microcompartments in E. coli and Salmonella. EcoSal Plus 9.

12. Axen SD, Erbilgin O, Kerfeld CA. 2014. A Taxonomy of Bacterial Microcompartment

Loci Constructed by a Novel Scoring Method. PLoS Computational Biology

10:e1003898.

13. Sutter M, Melnicki MR, Schulz F, Woyke T, Kerfeld CA. 2021. A catalog of the

diversity and ubiquity of bacterial microcompartments. Nat Commun 12:3809.

14. McFarland AG, Kennedy NW, Mills CE, Tullman-Ercek D, Huttenhower C, Hartmann

EM. 2021. Density-based binning of gene clusters to infer function or evolutionary

history using GeneGrouper. bioRxiv 2021.05.27.446007.

15. Kennedy NW, Hershewe JM, Nichols TM, Roth EW, Wilke CD, Mills CE, Jewett MC,

Tullman-Ercek D. 2020. Apparent size and morphology of bacterial

microcompartments varies with technique. PLOS ONE 15:e0226395.

16. Slininger Lee MF, Jakobson CM, Tullman-Ercek D. 2017. Evidence for Improved

Encapsulated Pathway Behavior in a Bacterial Microcompartment through Shell

Protein Engineering. ACS Synth Biol 6:1880–1891.

17. Kerfeld CA, Sawaya MR, Tanaka S, Nguyen CV, Phillips M, Beeby M, Yeates TO.

2005. Protein structures forming the shell of primitive bacterial organelles. Science

309:936–938.

18. Wheatley NM, Gidaniyan SD, Liu Y, Cascio D, Yeates TO. 2013. Bacterial

microcompartment shells of diverse functional types possess pentameric vertex

proteins. Protein Science 22:660–665.

19. Bobik TA, Havemann GD, Busch RJ, Williams DS, Aldrich HC. 1999. The

Propanediol Utilization (pdu) Operon ofSalmonella enterica Serovar Typhimurium

LT2 Includes Genes Necessary for Formation of Polyhedral Organelles Involved in

Coenzyme B12-Dependent 1, 2-Propanediol Degradation. Journal of bacteriology

181:5967–5975.

20. Lehman BP, Chowdhury C, Bobik TA. 2017. The N Terminus of the PduB Protein

Binds the Protein Shell of the Pdu Microcompartment to Its Enzymatic Core. Journal

of Bacteriology 199:e00785-16.

21. Lawrence AD, Frank S, Newnham S, Lee MJ, Brown IR, Xue W-F, Rowe ML,

Mulvihill DP, Prentice MB, Howard MJ, Warren MJ. 2014. Solution Structure of a

Bacterial Microcompartment Targeting Peptide and Its Application in the

Construction of an Ethanol Bioreactor. ACS Synthetic Biology 3:454–465.

22. Parsons JB, Lawrence AD, McLean KJ, Munro AW, Rigby SEJ, Warren MJ. 2010.

Characterisation of PduS, the pdu Metabolosome Corrin Reductase, and Evidence

of Substructural Organisation within the Bacterial Microcompartment. PloS one

5:e14009.

23. Fan C, Cheng S, Sinha S, Bobik TA. 2012. Interactions between the termini of

lumen enzymes and shell proteins mediate enzyme encapsulation into bacterial

microcompartments. Proceedings of the National Academy of Sciences 109:14995–

15000.

24. Jorda J, Liu Y, Bobik TA, Yeates TO. 2015. Exploring Bacterial Organelle

Interactomes: A Model of the Protein-Protein Interaction Network in the Pdu

Microcompartment. PLOS Computational Biology 11:e1004067.

25. Yang M, Simpson DM, Wenner N, Brownridge P, Harman VM, Hinton JCD, Beynon

RJ, Liu L-N. 2020. Decoding the stoichiometric composition and organisation of

bacterial metabolosomes. 1. Nature Communications 11:1976.

26. Sutter M, Greber B, Aussignargues C, Kerfeld CA. 2017. Assembly principles and

structure of a 6.5-MDa bacterial microcompartment shell. Science 356:1293–1297.

27. Kennedy NW, Ikonomova SP, Slininger Lee M, Raeder HW, Tullman-Ercek D. 2020.

Self-assembling shell proteins PduA and PduJ have essential and redundant roles

in bacterial microcompartment assembly. Journal of Molecular Biology

https://doi.org/10.1016/j.jmb.2020.11.020.

28. Pang A, Frank S, Brown I, Warren MJ, Pickersgill RW. 2014. Structural Insights into

Higher Order Assembly and Function of the Bacterial Microcompartment Protein

PduA. Journal of Biological Chemistry 289:22377–22384.

29. Fan C, Bobik TA. 2011. The N-terminal region of the medium subunit (PduD)

packages adenosylcobalamin-dependent diol dehydratase (PduCDE) into the Pdu

microcompartment. Journal of bacteriology 193:5623–5628.

30. Fan C, Cheng S, Liu Y, Escobar CM, Crowley CS, Jefferson RE, Yeates TO, Bobik

TA. 2010. Short N-terminal sequences package proteins into bacterial

microcompartments. Proceedings of the National Academy of Sciences 107:7509–

7514.

31. Liu Y, Jorda J, Yeates TO, Bobik TA. 2015. The PduL Phosphotransacylase Is Used

To Recycle Coenzyme A within the Pdu Microcompartment. Journal of Bacteriology

197:2392–2399.

32. Cheng S, Fan C, Sinha S, Bobik TA. 2012. The PduQ Enzyme Is an Alcohol

Dehydrogenase Used to Recycle NAD+ Internally within the Pdu Microcompartment

of Salmonella enterica. PLoS ONE 7:e47144.

33. Kerfeld CA, Erbilgin O. 2015. Bacterial microcompartments and the modular

construction of microbial metabolism. Trends in Microbiology 23:22–34.

34. Sampson EM, Johnson CLV, Bobik TA. 2005. Biochemical evidence that the pduS

gene encodes a bifunctional cobalamin reductase. Microbiology 151:1169–1177.

35. Honda S, Toraya T, Fukui S. 1980. In situ reactivation of glycerol-inactivated

coenzyme B12-dependent enzymes, glycerol dehydratase and diol dehydratase.

Journal of bacteriology 143:1458-1465.

36. Toraya T, Mori K. 1999. A Reactivating Factor for Coenzyme B12-dependent Diol

Dehydratase. Journal of Biological Chemistry 274:3372–3377.

37. Walter D, Ailion M, Roth J. 1997. Genetic characterization of the pdu operon: use of

1, 2-propanediol in Salmonella typhimurium. Journal of bacteriology 179:1013–1022.

38. Chen P, Andersson DI, Roth JR. 1994. The control region of the pdu/cob regulon in

Salmonella typhimurium. Journal of bacteriology 176:5474–5482.

39. Bobik TA, Ailion M, Roth JR. 1992. A single regulatory gene integrates control of

vitamin B12 synthesis and propanediol degradation. Journal of bacteriology

174:2253–2266.

40. Rondon MR, Escalante-Semerena JC. 1992. The poc locus is required for 1, 2-

propanediol-dependent transcription of the cobalamin biosynthetic (cob) and

propanediol utilization (pdu) genes of Salmonella typhimurium. Journal of

bacteriology 174:2267–2272.

41. Kim EY, Jakobson CM, Tullman-Ercek D. 2014. Engineering Transcriptional

Regulation to Control Pdu Microcompartment Formation. PLOS ONE 9:e113814.

42. Nichols TM, Kennedy NW, Tullman-Ercek D. 2020. A genomic integration platform

for heterologous cargo encapsulation in 1,2-propanediol utilization bacterial

microcompartments. Biochemical Engineering Journal 156:107496.

43. Nichols TM, Kennedy NW, Tullman-Ercek D. 2019. Cargo encapsulation in bacterial

microcompartments: Methods and analysis. Methods Enzymol 617:155–186.

44. Sinha S, Cheng S, Fan C, Bobik TA. 2012. The PduM Protein Is a Structural

Component of the Microcompartments Involved in Coenzyme B12-Dependent 1,2-

Propanediol Degradation by Salmonella enterica. J Bacteriol 194:1912–1918.

45. Parsons JB, Frank S, Bhella D, Liang M, Prentice MB, Mulvihill DP, Warren MJ.

2010. Synthesis of Empty Bacterial Microcompartments, Directed Organelle Protein

Incorporation, and Evidence of Filament-Associated Organelle Movement. Molecular

Cell 38:305–315.

46. Li Y, Kennedy NW, Li S, Mills CE, Tullman-Ercek D, Olvera de la Cruz M. 2021.

Computational and Experimental Approaches to Controlling Bacterial

Microcompartment Assembly. ACS Cent Sci 7:658–670.

47. Greber BJ, Sutter M, Kerfeld CA. 2019. The Plasticity of Molecular Interactions

Governs Bacterial Microcompartment Shell Assembly. Structure 27:749-763.e4.

48. Chowdhury C, Chun S, Pang A, Sawaya MR, Sinha S, Yeates TO, Bobik TA. 2015.

Selective molecular transport through the protein shell of a bacterial

microcompartment organelle. Proceedings of the National Academy of Sciences

112:2990–2995.

49. Chowdhury C, Chun S, Sawaya MR, Yeates TO, Bobik TA. 2016. The function of

the PduJ microcompartment shell protein is determined by the genomic position of

its encoding gene. Mol Microbiol 101:770–783.

50. Mohajerani F, Sayer E, Neil C, Inlow K, Hagan MF. 2021. Mechanisms of Scaffold-

Mediated Microcompartment Assembly and Size Control. ACS Nano 15:4197–4212.

51. Engler C, Gruetzner R, Kandzia R, Marillonnet S. 2009. Golden Gate Shuffling: A

One-Pot DNA Shuffling Method Based on Type IIs Restriction Enzymes. PLoS ONE

4:e5553.

52. Datta S, Costantino N, Court DL. 2006. A set of recombineering plasmids for gram-

negative bacteria. Gene 379:109–115.

53. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nat Meth 9:671–675.

54. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D,

Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J,

Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ,

Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I,

Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P.

2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat

Methods 17:261–272.

55. Crowley CS, Cascio D, Sawaya MR, Kopstein JS, Bobik TA, Yeates TO. 2010.

Structural insight into the mechanisms of transport across the Salmonella enterica

Pdu microcompartment shell. Journal of Biological Chemistry 285:37838–37846.

56. Pang A, Liang M, Prentice MB, Pickersgill RW. 2012. Substrate channels revealed

in the trimeric Lactobacillus reuteri bacterial microcompartment shell protein PduB.

Acta Cryst D, Acta Cryst Sect D, Acta Crystallogr D, Acta Crystallogr Sect D, Acta

Crystallogr D Biol Crystallogr, Acta Crystallogr Sect D Biol Crystallogr 68:1642–

1652.

57. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin

TE. 2004. UCSF Chimera--a visualization system for exploratory research and

analysis. J Comput Chem 25:1605–1612.

Figure Legends

Figure 1 – Schematic representation of cells containing Pdu MCPs. (A) Cells harboring MCPs can scavenge niche carbon sources from their environment. (B) The Pdu MCP encapsulates enzymes responsible for the metabolism of 1,2-propanediol to propionate and 1-propanol via the toxic intermediate propionaldehyde, which is sequestered by the MCP shell. The Pdu MCP also encapsulates various enzymes responsible for cofactor recycling. (C) MCPs are composed of a protein shell made of a variety of different shell protein architectures, including hexamers (BMC-H), trimers (BMC-T), and pentamers (BMC-P) and their various subtypes.

Figure 2 – A suite of fluorescent reporters is used to investigate hypotheses about enzyme encapsulation within Pdu MCPs. (A) A number of hypotheses have been proposed in the Pdu MCP literature regarding the mechanism of enzyme encapsulation (20-25, 33). Six of the eight Pdu shell proteins have been suggested to have a role in enzyme encapsulation. In order to test the hypothesis proposed in Lehman et al., we created a suite of fluorescent reporters (B) to inform on the encapsulation state on the entire Pdu MCP core.

Figure 3 – Loss of PduB disrupts enzyme encapsulation. (A) Schematic representation of the *pdu* operon and modifications to various LT2 strains used in this study, including the wild type (WT) operon. ΔA ΔJ has the open reading frames encoding PduA and PduJ knocked out, disrupting MCP formation. The ΔpocR strain has the transcriptional activator for the *pdu* operon knocked out, disrupting Pdu MCP expression. The ΔB strain has PduB knocked out and the ΔB::B-K102A-K207A strain has WT PduB replaced with a double mutant that disrupts PduB incorporation into the

MCP shell. (B) Phase contrast and GFP fluorescence micrographs of cells expressing

different GFP reporter constructs (scale bar = 1 μm). Different LT2 strains are indicated

by the far left vertical axis label and the micrograph type is indicated by the near left

vertical axis label. The reporter construct is indicated by the top horizontal labels.

Representative bright, fluorescent puncta indicating MCPs are labelled with a blue

arrow, polar bodies are labelled with a purple arrow, and diffuse fluorescence is

indicated with a yellow arrow. (C) Quantification of puncta per cell for different

fluorescently-tagged enzyme reporter constructs and various LT2 strains. The strain is

indicated by the top, horizontal label and the encapsulation tag is indicated by the

bottom horizontal label. Note that each reporter construct is readily encapsulated to

varying degrees into the WT strain, whereas their encapsulation is significantly depleted

in strains lacking PduB ($p < 0.0001$, one-tailed t-test). Puncta were counted from three

biological replicates and >106 cells were counted per strain/construct.

Figure 4 – Loss of PduB decouples Pdu MCP shell assembly from the enzymatic

core. (A) Coomassie-stained SDS-PAGE gel of MCP purifications from various LT2

strains (top horizontal label). Bands corresponding to various Pdu MCP components

(right vertical labels) are present in the WT strain but absent in other strains. MWM =

molecular weight marker. (B) Representative negative-stained transmission electron

micrographs of Pdu MCP purifications from various LT2 strains. Well-formed MCP

shells are irregular and polyhedral in shape and have a defined boundary and are

present in the WT, ΔB, and ΔB::B-K102A-K207A samples. (C) Pdu MCPs purified from

WT are significantly larger in diameter than shells purified from the ΔB or ΔB::B-K102A-

K207A strains ($p < 0.0001$, one-tailed t-test). Measurements were collected on three

biological replicates and >120 MCPs were measured for each sample type. (D)

Representative phase contrast and fluorescence micrographs for various strains of LT2

expressing ssP-GFP and PduA-mCherry reporter constructs (scale bar = 1 µm). The

strain is labelled on the top horizontal axis and the microscopy type is labelled on the

near left vertical axis. Bright, fluorescent puncta indicative of MCPs are labelled with a

blue arrow, polar bodies are indicated with a purple arrow, and diffuse fluorescence is

indicated with a yellow arrow. mCherry fluorescence is pseudocolored in yellow and

GFP fluorescence is pseudocolored in cyan. Overlapping fluorescence is indicated in

the merged image.

Figure 5 – PduB is incorporated in the Pdu MCP shell via a conserved hydrogen

bonding motif. (A) Schematic representation of BMC-H (orange) and BMC-T (blue)

proteins tiling together to form the facets of the MCP shell. (B) Lysine residues at the

hexamer-hexamer and hexamer-trimer interfaces are predicted to play a role in facet

tiling in a MCP from *H. ochraceum* (PDB: 5V74) (26). BMC-H proteins are in orange and

BMC-T proteins are in blue. (C) The edge lysine residue important for PduA (orange)

self-assembly is also present in PduB (blue) (PduA PDB: 3NGK (55); PduB PDB: 4FAY

(56)). All structural visualization was done using Chimera (57).

Figure 6 – PduB alterations change LT2 growth and metabolite profiles. (A) LT2

control and experimental strains containing different MCP morphologies (WT MCPs,

polar bodies (ΔA ΔJ, ΔB, and ΔB::B-K102A-K207A), no protein expression (ΔpocR)).

These differing morphologies lead to various LT2 growth profiles, as measured by the

optical density at 600 nm (OD_{600}) at different time points. (B) Metabolite profiles of

different LT2 strains during growth on 1,2-propanediol as a sole carbon source with

excess adenosylcobalamin (150 nM). The concentration (mM) of four key metabolites

(1,2-propanediol, propionaldehyde, propionate, and 1-propanol) was measured over the

duration of the growth curve visualized in Fig. 6A.

Figure 7 – System-level kinetic modelling reveals how a polar effect alters growth

and metabolite profiles. (A) Two organizational strategies for Pdu enzymes (MCPs

and polar bodies) were tested in a systems-level kinetic model. Plotted here are

propionaldehyde concentrations in the external media. PduCDE enzyme concentration

was also varied in the polar body model case to account for the potential genetic polar

effects that alter gene expression. (B) The kinetic model demonstrates that changes to

PduCDE concentration (mM) impact the maximum propionaldehyde concentration (mM)

observed in the external media when pathway enzymes are contained in a polar body.

(C) Comparison of GFP fluorescence (MEF FITC) measured by flow cytometry as a

proxy for gene expression vs peak propionaldehyde concentration (mM) reveals that

PduCDE expression is likely reduced due to a genetic polar effect in the ΔB strain.

A Human gut flora

△ Carbon sources
○ Bacterial microcompartments

B Bacterial microcompartment

Pdu metabolic pathway

1-propanol

propionate

1,2-propanediol

PduQ   NAD+
NADH
PduP   propionyl-CoA
PduW

PduCDE
propionaldehyde
Ado-B₁₂   cob(I)alamin
PduS
PduL   propionyl phosphate

PduGH
cob(III)alamin   PduO

Shell protein components

C

Architecture: BMC-H BMC-H^P BMC-T BMC-T^P BMC-T^{PD} BMV/BMC-P

Pdu example: PduA, PduJ PduK PduU PduT PduBB' PduN

A Possible encapsulation mechanisms

PduW (Cytosolic)
(Yang, 2020[25])

PduA/J

PduT

PduP
PduQ
(Fan, 2012[23])

PduS
PduO
(Parsons, 2010[22])

PduU

PduV
(Jorda, 2015[24])

PduL
(Kerfeld, 2014[33])

PduP
PduQ
(Lawrence, 2014[21])

PduK

PduGH
(Unknown)

PduCDE
(Lehman, 2017[20])

PduB

B Fluorescent reporters to test encapsulation

Construct: ssD-GFP
Purpose: Report on CDE encapsulation

Construct: ssP-GFP
Purpose: Report on PQ encapsulation

Construct: W-GFP
Purpose: Report on W encapsulation

Construct: ssL-GFP
Purpose: Report on L encapsulation

GFP
ssP

ssD
GFP

PduB

ssL

GFP

PduW

GFP

ssL

GFP

PduG

PduO
GFP

Construct: O-GFP
Purpose: Report on SO encapsulation

Construct: G-GFP
Purpose: Report on GH encapsulation

Hypothesis: PduB is necessary for encapsulation of all members of enzymatic core

A

WT operon

Pdu operon modifications

pocR F P_pdu A BB' C D E G H J K L M N O P Q S T U V W X

ΔA ΔJ

ΔpocR

ΔB

ΔB::B-K102A-K207A

Expected phenotype

WT	ΔA ΔJ	ΔpocR	ΔB and ΔB::B-K102A-K207A
MCPs/puncta	Polar bodies	Diffuse	Unknown
Controls			**Experimental**

B

Representative micrographs of cargo encapsulation

N-terminal GFP encapsulation tags

Strains: ssD ssP ssL PduG PduO PduW

WT — Phase / GFP

ΔA ΔJ — Phase / GFP

ΔpocR — Phase / GFP

ΔB — Phase / GFP

ΔB::B-K102A-K207A — Phase / GFP

Microscopy type

1 μm

▶ MCPs/puncta ▶ Polar bodies ▶ Diffuse fluorescence

Phenotypes

C

Trends in cargo encapsulation

Strain

| WT | ΔA ΔJ | ΔpocR | ΔB | ΔB::B-K102A-K207A |

p < 0.0001

Puncta per cell (y-axis: 0, 3, 6, 9)

Encapsulation tag (x-axis: ssD ssP ssL G O W, repeated per strain)

A SDS-PAGE gel of purified MCPs

B Representative micrographs of purified MCPs

C Trends in MCP diameter

D Representative micrographs of MCP shell and cargo colocalization

A

Bacterial microcompartment

BMC-H   BMC-T

MCP shell facet

B

Assembly of BMC-H and BMC-T proteins (*H. ochraceum*)

Hexamer-hexamer H-bonding interface

Hexamer-trimer H-bonding interface

C

Structural alignment of PduA (*S. enterica*) and PduB (*L. reuteri*)

Conserved edge lysine residues

BMC-H   BMC-T   ---- H-bond   PduA   PduB