Title: Climate-driven variation in biotic interactions provides a narrow and

variable window of opportunity for an insect herbivore at its ecological margin

Running Head: Host availability shaped by seasonal variation

James E. Stewart[1], Ilya M.D. Maclean[2], Gara Trujillo[3], Jon Bridle[4,5] and Robert

J. Wilson[1,2,6]

[1]College of Life and Environmental Sciences, University of Exeter, United

Kingdom

[2]Environment & Sustainability Institute, University of Exeter, Penryn Campus,

United Kingdom

[3]International Institute for Industrial Environmental Economics (IIIEE),

Lund University P.O. Box 196, 22100 Lund, Sweden

[4]School of Biological Sciences, University of Bristol, United Kingdom

[5]Department of Genetics, Evolution, and Environment, University College

London, London

[6]Departmento de Biogeografía y Cambio Global, Museo Nacional de Ciencias

Naturales, Madrid E28006, Spain

Correspondence: James E. Stewart, College of Life and Environmental

Sciences, University of Exeter, Exeter, United Kingdom. E-mail:

j.e.stewart8@gmail.com

Abstract

Climate-driven geographic range shifts have been associated with transitions between dietary specialism and generalism at range margins. The mechanisms underpinning these often transient niche breadth modifications are poorly known, but utilisation of novel resources likely depends on phenological synchrony between the consumer and resource. We use a climate-driven range and host shift by the butterfly *Aricia agestis* to test how climate-driven changes in host phenology and condition affect phenological synchrony, and consider implications for host use.

Our data suggest that the perennial plant which was the primary host before range expansion is a more reliable resource than the annual Geraniaceae upon which the butterfly has become specialised in newly colonised parts of its range. In particular, climate-driven phenological variation in the novel host *Geranium dissectum* generates a narrow and variable 'window of opportunity' for larval productivity in summer. Therefore, although climatic change may allow species to shift hosts and colonise novel environments, specialisation on phenologically-limited hosts may not persist at ecological margins as climate change continues. We highlight the potential role for phenological (a)synchrony in determining lability of consumer-resource associations at range margins, and the importance of considering causes of synchrony and success in biotic interactions when predicting range shifts.

Keywords: brown argus, Lepidoptera, host shift, specialisation, asynchrony

Introduction

Climate change is causing widespread shifts in species' geographic range limits [1–3]. The extent of such shifts depends on species' life histories, potential for plastic responses and the quality of available habitat at the expanding margin [4–6]. Habitat availability may itself be determined by the process of range expansion: range shifts have recently been identified as a cause, rather than consequence, of increased dietary generalism at poleward range margins in herbivorous insects [7,8]. However, dietary generalism and incorporation of novel host plants can be transient in such systems, and the mechanisms

underlying gain or loss of hosts from insect diets is poorly known [7–9]. Here,
we highlight phenological (a)synchrony between insects and unpredictable host
resources as a potential mechanism for lability in insect-host associations at
range margins.

The phenology of many consumers is closely synchronised with the
development and availability of their resources, and their interactions occur
within an often narrow 'window of opportunity' for the consumer defined by the
phenology of both partners [10–14]. For herbivorous insects, the length of the
phenological window of opportunity will be partly determined by the specificity of
its interaction with the host, the host's growth form, and the broader
environmental context. For example, the window of opportunity for herbivory is
typically longer on perennials than on short-lived annual plants, the availability
of which may be defined by environmental drivers of their germination and
senescence [15,16]. The window is also typically longer for polyphagous than
obligately monophagous species (which can exploit fewer distinct phenological
windows), and for populations inhabiting topographically variable landscapes in
which heterogeneous microclimates provide diverse phenological windows
[16,17]. Differences in synchrony among nearby microclimates may also cause
local variation in host condition, quality and profitability for the herbivore,
thereby influencing local patterns of host selection and opportunities for dietary
change [7,8,18–20].

Robust evidence is therefore required on the drivers and vulnerability of host-
herbivore phenological synchrony (which may scale up to emergent patterns of
herbivore range and host shifts [10,19,21]), including the role of microclimate in
consumer persistence by potentially buffering asynchrony in biotic interactions
[18,20]. Here we address this knowledge gap, using as an exemplar the brown
argus butterfly (*Aricia agestis*; Lepidoptera: Lycaenidae) at its range margin in
the UK. In doing so, we highlight how phenological (a)synchrony could provide
an underlying mechanism for apparently high rates of host shifting near range
margins [7,22]. Brown argus butterflies have two generations per year; larval
offspring of the second generation emerge from mid-August and feed on leaves
of the host plant before overwintering [23]. The brown argus' UK range was
historically largely restricted to calcareous grassland where its host, the
perennial *Helianthemum nummularium* (Malvales: Cistaceae; hereafter

Helianthemum), grows. However, since the 1990s, the brown argus has

undergone a climate-driven range expansion associated with rapid evolution of

biotic interactions, including specialisation on annual Geraniaceae species

(*Erodium cicutarium*, *Geranium dissectum* and *G. molle*; Geraniales:

Geraniaceae) mainly in regions beyond the former range limit [21,24–27].

In this study, we test expectations of (a) greater temporal variation in annual

versus perennial host plant condition, and (b) more opportunity for asynchrony

between the consumer and its annual hosts than its perennial host, as a

consequence of (a). We expect asynchrony with the less-predictable annual

hosts to be more pronounced under warm, dry summer conditions, and we test

for such effects on the condition and phenology of annual Geraniaceae hosts

that have enabled the range expansion. We conduct these tests (c) across sites

and years, and (d) across a range of microclimates within a site. We then

consider the implications of asynchrony for host use, shifting host associations

at range margins, and the range dynamics of host-limited herbivores.

Methods

Study system

Brown argus butterflies prefer to lay their eggs on Geraniaceae, on which larvae

grow 10% larger and faster, than on *Helianthemum* (the ancestral host at the

range margin in Britain), and prefer to lay on better condition leaves regardless

of host species [19,21,24–26,28,29]. However, the annual, ephemeral growth

form of the Geraniaceae host plants may make them less reliable as a food

source than the evergreen perennial *Helianthemum*, especially under more

variable climatic conditions at the range margin [16]. To investigate this, we

surveyed ten sites fortnightly–monthly between July 2016 and October 2017, to

monitor phenology and condition of three host plant species. *Helianthemum*

was the dominant host at five calcareous grassland sites, while Geraniaceae

(*G. dissectum* and *E. cicutarium*; hereafter *Geranium* and *Erodium*) were

dominant at five grassland/ dune sites (Figure 1). The number of sites and

quadrats was chosen to maximise spatial coverage and replication within

logistical constraints. See SI-1.1 and SI-1.2 for survey dates and site profiles.

Quadrat surveys and analyses

Using an average of 33.7 (0.25 m^2) quadrats per survey (SI-1.3 for quadrat placement), we recorded percentage cover (SI-1.3.2), phenophase (phenological stage) and condition of each host species. Host phenophase and condition underpin brown argus egg-laying site choice in the field [19,30]. Phenophase was estimated on a four-point scale (in leaf L, bud B, flower F, or had set seed S). Condition was visually assessed on a scale of 0–3 (poor–high quality for egg-laying, following [19,26]; see SI-1.3.3 for details and justification). We also measured mean sward height and percentage cover of bare ground (SI-1.3) which can alter local microclimates [31,32].

Quadrat-level phenophase and condition were estimated based on the average for each host species in each quadrat. This approach can mask fine-scale changes in phenophase and host condition, particularly for the annual hosts, so we also recorded *plant-level* phenophase and condition of the smallest, earliest phenophase Geraniaceae plant in each quadrat. This approach allows inference as to whether new germination has occurred, and of the age/condition of the plant material likely to be available to overwintering larvae. By September, Geraniaceae plants in later phenophases are typically senescent, in poor condition and expected to die before the autumn, making them a poor resource for larvae [30].

Quantifying variation in host plant condition and phenophase

To test expectation (a), for greater temporal variation in annual than perennial host condition, we used a Kruskal-Wallis (KW) test for each sampling period to compare quadrat-level condition between host species. Bonferroni-corrected Dunn's tests were then used to identify which hosts differed significantly from one another in condition score within each sampling period. We then conducted interannual comparisons of Geraniaceae host condition (Mann-Whitney U tests, MWU) and phenophase (X^2 tests): these compared 2016 data with 2017 data, separately for each month between July to October. These months are the most relevant for host choice and larval feeding by second generation brown argus and their offspring.

Assessing plant-herbivore (a)synchrony

To contextualise host condition with reference to herbivore phenology, and address question (b), we overlaid plots of host condition indices (calculated using survey data, SI-1.5.1) from all July–October surveys with emergence phenology curves of adult second generation brown argus and their larval offspring. Adult phenology was described as Gaussian curves, following [30,33], based on the mean and standard deviation of the brown argus' peak flight date from UK Butterfly Monitoring Scheme count data across 293 regularly-surveyed sites between 1992–2016 [34]. The phenomenological models used to generate phenology estimates account for variation in phenology based on latitude and temperature [30,33].

In summary, brown argus second brood phenology in Britain varies with latitude (earlier further north) and between-brood temperature (earlier under warmer conditions between the first and second brood), and is related to the (latitude-dependent) phenology of the first brood [30,33]. See SI-1.6 for more information. Using model-derived estimates ([30]; SI-1.6), we plotted two adult phenology curves: 'early' phenology based on the latitude of the most northerly site and 95th percentile between-brood temperature, and 'late' phenology based on the latitude of the most southerly site and 5th percentile between-brood temperature. The larval phenology curves were assumed to track the adult curves, with an 11-day lag to account for mating and egg-laying (four days post-emergence) and larval emergence (one week) [24].

Testing climatic predictors of Geraniaceae recruitment and condition

To address our third question (c), using data on the youngest host within each quadrat, we tested for climatic drivers of recruitment and condition of each Geraniaceae species in early September, when most larval offspring were expected to have emerged to feed. We defined recruitment as the presence of at least one young, leaf-stage host plant in the focal quadrat. Plants in condition categories 1 and 2 were rarely observed during September surveys. We therefore reclassified condition 0/1 plants as poor condition (0) and condition 2/3 plants as good condition (1).

To test predictors of recruitment and condition, we used logistic regression with

the following putative predictors: year, site, northing, easting, vegetation height,

bare ground cover, and (linear and quadratic terms for) local weather estimates

based on the UK Meteorological Office's 5 km gridded weather data [35]. Using

daily weather data [35], we calculated the minimum, mean and maximum

temperature and mean rainfall for three periods in each year (justified in SI-1.7):

July, July–September and August–September (only including weather data up

to the day of quadrat sampling at each site in early September). We also used

daily weather data to calculate the Gaussen Aridity Index (GAI; precipitation/(2

× temperature)) for each period (e.g. [36]). Higher GAI values indicate cooler,

wetter conditions. Day of year was tested both as a putative fixed effect

predictor and as an offset term to account for day of sampling.

All continuous predictors were standardised and site ($n ≤ 5$ per host plant) was

included as a fixed effect (following [37]). We constructed candidate models by

considering all plausible parameter combinations (including temperature-rainfall

interactions), estimated parameters using maximum likelihood, and used AIC-

based model selection to establish the most parsimonious model(s) (see SI-

1.8–1.9 for details of model selection, validation and diagnostics).

Testing microclimate effects on Geraniaceae phenology and condition

In September 2017, we calculated condition and phenophase indices (SI-1.5.2)

for each of 31 quadrats at site G1, using the phenophase (L, B, F and S) and

condition (0–3) of all *Geranium* plants in each quadrat. The indices range

between 0 (quadrats contain only plants at condition 0/seed set stage) and 1

(plants at condition 3/leaf stage). To test expectation (d), we modelled the

indices (logistic regression) as a function of putative quadrat-specific predictors:

microclimate (mean, maximum and minimum temperature and soil moisture; SI-

1.10), percentage cover of bare ground and mean sward height. We also

considered plausible temperature-moisture interactions.

<u>**Results**</u>

Variation in host plant condition and phenophase

There was a substantial decline in quadrat-level condition of the annual host

G. dissectum over summer 2016 that was not observed in the perennial

Helianthemum or to the same extent in *Erodium* (Figure 2a; KW tests, Table

S4). However, by early November 2016, senesced *Geranium* had mostly been

replaced by recently-germinated, better condition recruits, and the quadrat-level

condition was at least as high as that of *Helianthemum* (Figure 2a-c; KW tests,

Table S4).

Overall, in summer 2016 both Geraniaceae species showed substantial

evidence of leaf senescence, including wilting and abscission, and little

germination or seedling establishment until early October (*pers. obs.*)

(Geranium, Figure 2b-c; Erodium, Figure S3). By contrast, in 2017, there were

many more seedlings and good condition plants throughout the summer months

and into the main sampling period in September (*pers. obs.*) (Geranium, Figure

2b-c; Erodium, Figure S3).

The condition of the youngest *Geranium* in each quadrat was significantly

higher in all 2017 survey periods compared to 2016 (Figure 2b; MWU tests,

Table S5). *Geranium* phenophases differed significantly between years (X^2

tests, Table S5): in 2016, the youngest plant in each *Geranium* quadrat was

typically an older plant in the seed set stage (Figure 2c) and new plants in the

leaf stage did not dominate until October; however, this younger form was

dominant throughout summer 2017 (Figure 2c).

Erodium showed similar patterns, though condition was approximately

equivalent between years for the August and September surveys (Figure S3;

Mann-Whitney U tests, Table S5). *Erodium* quadrats were also dominated by

young, leaf-stage plants in 2017: significantly more so than in 2016 during July

and August (Figure S3; X^2 tests, Table S5). Similar patterns of Geraniaceae

phenophase and condition were observed at quadrat-level as these plant-level

assessments (Figures S4 and S5).

Plant-herbivore (a)synchrony

We assessed potential for asynchrony between the butterfly and its hosts

(question (b)) by overlaying plots of host condition indices with curves

representing phenology of 'early' and 'late' emerging brown argus and their

offspring (Figure 3). At their respective sites, good-condition *Helianthemum* and

Erodium hosts appear to be available throughout egg-laying and early larval

feeding periods of the brown argus butterfly, with little variation between two
climatically different years (2016 and 2017), or among sites at each sampling
period (Figure 3a,c). By contrast, in 2016 *G. dissectum* condition declined over
the early and late peaks of adult brown argus emergence, and was lowest
during the period in which females would normally be laying eggs and most
larvae would be beginning to feed (Figure 3b). Among-site variation at each
sampling period appears to be more pronounced regarding *G. dissectum*
condition, but this does not mask the temporal variation within and between
years (Figure 3; Table S4). Among-site variation in host condition likely results
from local variation in temperature and water relations linked to factors including
weather, topography and geology.

Climatic predictors of Geraniaceae condition and recruitment

We integrated climate data with in-field host plant surveys to investigate
potential drivers of condition and phenology across the *Geranium* sites,
addressing question (c). The probability of the youngest *Geranium* plants being
in good condition increased with summer rainfall (MC_{final}; Figure 4a, Table 1a).
There was limited evidence that moister, cooler conditions in areas with shorter
vegetation were associated with better condition (Table S7). Forcing models of
Geranium condition to include an effect of year resulted in higher AIC values,
and inflated parameter estimates and standard errors by several orders of
magnitude, so these are reported only in the Supplementary Information for
context (SI-1.11.4 , Table S7), and there are no effects of year in the final model
set. The evidence suggests that probability of new *Geranium* recruitment was
higher in 2017 (MR_{best}; SI-1.11.4, Table S8), and lower following higher mean
daily temperatures during August–September (MR_{final}; Figure 4b, Table 1b). .
There were no detectable effects of site northing or easting, or day of sampling
on condition or recruitment.

We had low statistical power to detect relationships between weather and the
condition and recruitment of *Erodium*, models for which are outlined in SI1.11.4.

Microclimate effects on Geraniaceae phenology and condition

To address question (d), we assessed Geranium phenology and condition
across a range of microclimates at site G1 in September 2017. At this site,

areas of moister soil (where plants are less likely to dry out) were associated

with Geranium plants in better condition and earlier phenophases (Figure 4c,d;

Table 2). Candidate models suggested such plants were also more prevalent in

areas with warmer, moister microclimates (SI-1.11.5).

Discussion

Here, we describe temporal variation in condition and phenology of three plant

species and highlight the implications for their use as larval host plants by a

butterfly species that has recently expanded its geographic range. Our data,

from multiple sites across two years, suggest that the annual host species vary

more in condition and availability than a perennial species that has been used

as a long-standing host at the range margin, and do so in a way that differs

between years and with (micro-)climatic conditions. Such variation likely

generates narrow and unpredictable phenological 'windows of opportunity' for

exploitation of annual species that vary among sites and years under the

conditions of variable population sizes or phenology observed near the limits of

species' geographic ranges [38]. In this case, the perennial plant that was used

as the main pre-expansion host appears to be a more reliable resource, where

present, than the more widespread annual Geraniaceae species that have

acted as primary hosts during the climate-associated range expansion. These

results suggest that climatic effects on the phenological synchrony of biotic

interactions could act as a mechanism generating transient patterns of host

associations at range margins, and consequently of habitat availability in the

landscape and patterns of range shifting.

(Micro-)climatic variation and host plant phenology

Our data indicate that the phenology of two winter annuals used by the brown

argus butterfly – in particular *G. dissectum* – is sensitive to weather variability.

Greenhouse experiments and field observations suggest that summer

temperatures and moisture thresholds are crucial to dormancy breaking and

germination in *Geranium* [39,40]. A complementary interpretation of these data

is that *Geranium* plants may germinate early following a cool spell and early

summer rain, but will suffer high seedling mortality where the summer is

subsequently hot and dry (e.g. [41,42]). For example, our data show that hot,

dry conditions in 2016 were associated with early and pronounced senescence
of plants in July, as well as delayed germination and/or early seedling mortality.
By contrast, our study sites received relatively high rainfall throughout summer
2017 [30], which is likely to have overcome moisture-dependency in dormancy
breaking and/or promoted seedling survival. *Geranium* condition was also
higher following wetter (2017) summer conditions, which supports evidence that
drought and thermal stress cause premature senescence and declines in the
quality (for consumers) of herbaceous plants [43,44]. A higher proportion of
younger and better-condition *Erodium* were available in 2017 than in 2016,
although we lacked statistical power to associate this with climatic variables (SI-
1.11.4). However, our data support previous observations that reduced soil
water availability advances the reproductive stage and the end of the growing
season in *Erodium*, and reduce its investment in leaf biomass [45,46].

Trophic interactions at range margins

Host condition and phenological synchrony in biotic interactions appear to be
mediated by local (micro-)climatic variation, and are major determinants of
spatiotemporal variation in fecundity and population size of host-specialist
herbivores such as the brown argus [19,22,26,47,48]. By reducing temporal
overlap between suitable resources and key herbivore life stages (e.g. adult
egg-laying and early larval stages), adverse (micro-)climatic conditions may limit
egg-laying and feeding opportunities and reduce larval survival, particularly
where plants with limited temporal availability (such as the annual *Geraniaceae*
studied here) are the main hosts [18,49]. Climatic conditions that are set to
become more common (i.e. variable rainfall and longer, hotter summers
[50,51]), may therefore narrow or close the phenological window of opportunity
for this host-specialist herbivore to exploit these ephemeral annual resources in
late summer [47,48].

The widespread annual Geraniaceae species are the hosts primarily used by
brown argus at the expanding front of its recent range in Britain, whereas
Helianthemum represents the apparent ancestral host at most sites where the
species has been present for the past century or longer [26,30]. In the brown
argus, variation in butterfly-host synchrony (and the success of this relatively
novel interaction) may therefore be especially pronounced and unpredictable

near range limits, particularly if abiotic conditions are marginal and population

sizes small [22].

The relationships between weather, plant condition and phenology, that we

identify here for *Geranium*, are therefore crucial in mediating how climate

change, variability and unpredictability will affect synchrony in biotic

interactions. Given that climate change is causing widespread changes in

phenological synchrony, both existing and novel host interactions may be

vulnerable under climate change. However, recent evidence suggests that at

least as many biotic interactions are becoming more synchronous as are

becoming less synchronous [22]. This evidence therefore also highlights the

potential for novel biotic interactions to emerge, and further supports a role for

spatiotemporal variation in synchrony underlying transient host interactions.

Spatiotemporal variation in the synchrony of biotic interactions is likely to

generate transient mosaics of selection pressures for different diets, and

thereby influence patterns of dietary specialisation/generalisation that recent

research has shown to be an emergent and surprisingly common property of

range dynamics [7,8,21,24]. In range expansions, many new consumer-

resource interactions form and some may be lost [7,8]. For example, diet

breadths in populations of Edith's checkerspot butterfly (*Euphydryas editha*)

increased after colonisation events as individual host preferences diversified,

but populations subsequently reverted to monophagy [8].

Spatiotemporal variation in phenological synchrony may prove to be an

underlying mechanism not only for lability in insect-host associations at range

margins, but also for range limit stability under scenarios of temporal

environmental variability. Range limits are more stable (expansions less likely)

where environmental variance is too large for adaptation and colonisation [52].

Therefore, large environmental variance may preclude colonisation events that

depend on predictably synchronous biotic interactions.

Evolution during range expansion

Following the recent range expansion and incorporation of Geraniaceae into the

diet of the brown argus, Geraniaceae-feeding populations that were able to

persist became specialised on the novel hosts, losing the adaptive capacity to

use *Helianthemum* [21,24]. Though specialisation on Geraniaceae appears to

have become more reliable on average [29], our data suggest that interannual

variation in phenology could still alter the success of this interaction, which may

prove to be locally transient in the face of phenological asynchrony. In a

comparable example from Scandinavia, narrow oviposition preferences of the

Glanville fritillary (*Melitaea cinxia*) for phenologically-limited hosts risks high

larval mortality under severe drought conditions in some years [20,53].

Conversely, phenological asynchrony at recently colonised Geraniaceae sites

could lead to variable population dynamics, and selection favouring continued

dispersal from the natal site in search of suitable egg-laying locations [54,55].

During the range expansion, selection has apparently favoured more dispersive

phenotypes which have increased flight capacity and more readily accept the

geographically widespread Geraniaceae hosts [24,27,28]. Consequently, under

certain (micro-)climatic conditions, the dispersive, Geraniaceae-favouring

phenotype may represent an alternative life history strategy that drives

expansion at range margins and in-filling of the core range. Subsequent

migrants that colonise *Helianthemum* sites may need to regain the ability to use

Helianthemum (as shown in [21]; cf. [8]) in order to benefit from stability of (and

phenological synchrony with) the host resource.

Conservation and management implications

Understanding constraints and opportunities for species' distributions is central

to successful conservation practices. Our results highlight the importance of

considering drivers of synchrony and the outcomes of biotic interactions when

examining climate-driven range shifts, and recognising the crucial roles of

microclimate and individual behaviour in mediating these interactions [19].

Conservation strategies could seek to maximise habitat and microclimatic

heterogeneity to promote diversity in local phenologies across trophic levels

[56,57]. In some cases, microclimatic variation may generate sufficient fine-

scale spatial heterogeneity in relative phenology and host condition to buffer

local herbivore populations against phenological asynchrony [10,18,58,59].

However, to improve our predictions of ecological responses to climate change,

and of the critical levels of environmental change likely to cause rapid loss of

ecosystem outputs, more empirical data are needed on shifts in biotic

interactions across populations, climates and species ranges, and their rates

and patterns of evolution. In the present case, we have highlighted what might

be typical variation in phenological synchrony across two years, which

emphasises the potentially large indirect impacts of climate change on

herbivore success. However, it would be beneficial to expand sampling across

both time and space to better understand general patterns of interannual

variation and gain a more holistic understanding of the phenomena discussed.

Conclusions

Our results place a novel emphasis on the interactions between phenology,

resource use and climate change in a range-expanding herbivore whose

sensitivity to small changes in temperature might otherwise predict a positive

range expansion response under future climates. Instead, small changes in

temperature and moisture regime have the potential to disrupt the range

expansion, mediated by phenological and physiological changes in the

herbivore's larval resource that essentially fragment the herbivore's potential

range. This suggests that novel host interactions may only remain as transient

resources at shifting range margins. The mechanisms underlying insect-plant

interactions, and their responses to climate change, are likely to be more

complex than they appear, and we need more detailed knowledge of these

mechanisms to understand and predict species' interactions and responses to

climate change.

Acknowledgements

Research was funded by the Natural Environment Research Council (PhD

studentship grant NE/L002434/1 to JES) and by the Botanical Society of Britain

and Ireland (BSBI Science and Research Council grant to JES). Research

permits were kindly granted by Butterfly Conservation, the Forestry

Commission, Holkham Estate, the National Trust, Natural England, and Norfolk

Wildlife Trust. Special thanks to S. Henderson, J. Chapman and their respective

teams at Holkham Estate and Magdalen Hill Down for their friendly assistance

and repeated support. Thanks also R. Menéndez and J. Bennie for helpful

comments on the research, and to R. Butlin, A. Phillimore and an anonymous

reviewer whose comments greatly improved this work.

References

1. Louthan AM, Doak DF, Angert AL. 2015 Where and when do species
interactions set range limits? *Trends in Ecology & Evolution* **30**, 780–792.
(doi:10.1016/j.tree.2015.09.011)

2. Nadeau CP, Urban MC, Bridle JR. 2017 Coarse climate change projections
for species living in a fine-scaled world. *Global Change Biology* **23**, 12–24.
(doi:10.1111/gcb.13475)

3. Sexton JP, McIntyre PJ, Angert AL, Rice KJ. 2009 Evolution and ecology of
species range limits. *Annual Review of Ecology, Evolution, and Systematics*
40, 415–436. (doi:10.1146/annurev.ecolsys.110308.120317)

4. Macgregor CJ *et al.* 2019 Climate-induced phenology shifts linked to range
expansions in species with multiple reproductive cycles per year. *Nat*
Commun **10**, 1–10. (doi:10.1038/s41467-019-12479-w)

5. Platts PJ, Mason SC, Palmer G, Hill JK, Oliver TH, Powney GD, Fox R,
Thomas C. 2019 Habitat availability explains variation in climate-driven
range shifts across multiple taxonomic groups. *Scientific Reports* **9**, 15039.

6. Eriksson M, Rafajlović M. 2021 The role of phenotypic plasticity in the
establishment of range margins. , 2021.10.04.463099.
(doi:10.1101/2021.10.04.463099)

7. Lancaster LT. 2020 Host use diversification during range shifts shapes
global variation in Lepidopteran dietary breadth. *Nature Ecology & Evolution*
4, 963–969. (doi:10.1038/s41559-020-1199-1)

8. Singer MC, Parmesan C. 2021 Colonizations cause diversification of host
preferences: A mechanism explaining increased generalization at range
boundaries expanding under climate change. *Global Change Biology* **27**,
3505–3518. (doi:10.1111/gcb.15656)

9. de la Paz Celorio-Mancera M, Wheat CW, Vogel H, Söderlind L, Janz N,
Nylin S. 2013 Mechanisms of macroevolution: polyphagous plasticity in
butterfly larvae revealed by RNA-Seq. *Molecular Ecology* **22**, 4884–4895.
(doi:10.1111/mec.12440)

10. Ward SF, Moon RD, Herms DA, Aukema BH. 2019 Determinants and
consequences of plant–insect phenological synchrony for a non-native
herbivore on a deciduous conifer: implications for invasion success.
Oecologia **190**, 867–878. (doi:10.1007/s00442-019-04465-2)

11. Visser ME, Holleman LJM, Gienapp P. 2006 Shifts in caterpillar biomass
phenology due to climate change and its impact on the breeding biology of
an insectivorous bird. *Oecologia* **147**, 164–172. (doi:10.1007/s00442-005-
0299-6)

12. Fuentealba A, Pureswaran D, Bauce É, Despland E. 2017 How does
synchrony with host plant affect the performance of an outbreaking insect
defoliator? *Oecologia* **184**, 847–857. (doi:10.1007/s00442-017-3914-4)

13. Posledovich D, Toftegaard T, Wiklund C, Ehrlén J, Gotthard K. 2018
Phenological synchrony between a butterfly and its host plants:
Experimental test of effects of spring temperature. *Journal of Animal*
Ecology **87**, 150–161. (doi:10.1111/1365-2656.12770)

14. Toftegaard T, Posledovich D, Navarro-Cano JA, Wiklund C, Gotthard K,
Ehrlén J. 2019 Butterfly–host plant synchrony determines patterns of host
use across years and regions. *Oikos* **128**, 493–502. (doi:10.1111/oik.05720)

15. Schoonhoven LM, van Loon JJA, Dicke M. 2005 *Insect-Plant Biology.*
Second Edition. Oxford, New York: Oxford University Press.

16. Kemp RJ, Hardy PB, Roy DB, Dennis RLH. 2008 The relative exploitation of
annuals as larval host plants by European butterflies. *Journal of Natural*
History **42**, 1079–1093. (doi:10.1080/00222930801937608)

17. Ward SE, Schulze M, Roy B. 2018 A long-term perspective on microclimate
and spring plant phenology in the Western Cascades. *Ecosphere* **9**, e02451.
(doi:10.1002/ecs2.2451)

18. Hellmann JJ, Weiss SB, McLaughlin JF, Ehrlich PR, Murphy DD, Launer
AE. 2004 Structure and dynamics of Euphydryas editha populations. In *On*
the Wings of Checkerspots. A Model System for Population Biology (eds PR
Ehrlich, I Hanski), pp. 34–62. Oxford, United Kingdom: Oxford University
Press.

19. Stewart JE, Maclean IMD, Edney EJ, Bridle JR, Wilson RJ. 2021
Microclimate and resource quality determine resource use in a range-
expanding herbivore. *Biology Letters* **17**, 20210175.
(doi:10.1098/rsbl.2021.0175)

20. Rytteri S, Kuussaari M, Saastamoinen M. 2021 Microclimatic variability
buffers butterfly populations against increased mortality caused by
phenological asynchrony between larvae and their host plants. *Oikos* **130**,
753–765. (doi:https://doi.org/10.1111/oik.07653)

21. Buckley J, Bridle JR. 2014 Loss of adaptive variation during evolutionary
responses to climate change. *Ecology Letters* **17**, 1316–1325.
(doi:10.1111/ele.12340)

22. Kharouba HM, Ehrlén J, Gelman A, Bolmgren K, Allen JM, Travers SE,
Wolkovich EM. 2018 Global shifts in the phenological synchrony of species
interactions over recent decades. *Proc Natl Acad Sci USA* **115**, 5211–5216.
(doi:10.1073/pnas.1714511115)

23. Asher J, Warren MS, Fox R, Harding P, Jeffcoate G, Jeffcoate S. 2001 *The*
Millennium Atlas of Butterflies In Britain and Ireland. Oxford, United
Kingdom: Oxford University Press.

24. Bridle JR, Buckley J, Bodsworth EJ, Thomas CD. 2014 Evolution on the
move: specialization on widespread resources associated with rapid range
expansion in response to climate change. *Proceedings of the Royal Society*
B: Biological Sciences **281**, 20131800. (doi:10.1098/rspb.2013.1800)

25. Thomas CD, Bodsworth EJ, Wilson RJ, Simmons AD, Davies ZG, Musche
M, Conradt L. 2001 Ecological and evolutionary processes at expanding
range margins. *Nature* **411**, 577–581. (doi:10.1038/35079066)

26. Bourn NAD, Thomas JA. 1993 The ecology and conservation of the brown
argus butterfly Aricia agestis in Britain. *Biological Conservation* **63**, 67–74.
(doi:10.1016/0006-3207(93)90075-C)

27. Buckley J, Butlin RK, Bridle JR. 2012 Evidence for evolutionary change
associated with the recent range expansion of the British butterfly, Aricia
agestis, in response to climate change. *Mol. Ecol.* **21**, 267–280.
(doi:10.1111/j.1365-294X.2011.05388.x)

28. Bodsworth EJ. 2002 Dispersal and behaviour of butterflies in response to
their habitat. PhD thesis, University of Leeds, Leeds, United Kingdom.

29. Pateman RM. 2012 Climate change and habitat associations at species'
range boundaries. PhD thesis, University of York.
(doi:http://etheses.whiterose.ac.uk/2910/1/Thesis_RM_Pateman_2012.pdf)

30. Stewart JE. 2020 Linking Phenology to Population Dynamics and
Distribution Change in a Changing Climate. Unpublished PhD Thesis,
University of Exeter.

31. Suggitt AJ, Gillingham PK, Hill JK, Huntley B, Kunin WE, Roy DB, Thomas
CD. 2011 Habitat microclimates drive fine-scale variation in extreme
temperatures. *Oikos* **120**, 1–8. (doi:10.1111/j.1600-0706.2010.18270.x)

32. Bramer I *et al.* 2018 Advances in monitoring and modelling climate at
ecologically relevant scales. *Next Generation Biomonitoring, part 1* **58**, 101–
161. (doi:http://dx.doi.org/10.1016/bs.aecr.2017.12.005)

33. Dennis EB, Morgan BJT, Freeman SN, Roy DB, Brereton T. 2016 Dynamic
models for longitudinal butterfly data. *JABES* **21**, 1–21.
(doi:10.1007/s13253-015-0216-3)

34. UKBMS. 2017 UK Butterfly Monitoring Scheme (UKBMS) occurrence
dataset. (doi:10.15468/gmqvmk)

35. Met Office. 2017 UKCP09: Met Office gridded land surface climate
observations - daily temperature and precipitation at 5km resolution.
Updated to 2017.

36. Donoso I, Stefanescu C, Martinez-Abrain A, Treveset A. 2016 Phenological
asynchrony in plant-butterfly interactions associated with climate: a
community-wide perspective. *Oikos* **125**, 1434–1444.

37. Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin
CED, Robinson BS, Hodgson DJ, Inger R. 2018 A brief introduction to
mixed effects modelling and multi-model inference in ecology. *PeerJ* **6**,
e4794. (doi:10.7717/peerj.4794)

38. Oliver TH, Roy DB, Brereton T, Thomas JA. 2012 Reduced variability in
range-edge butterfly populations over three decades of climate warming.

Global Change Biology **18**, 1531–1539. (doi:10.1111/j.1365-
2486.2012.02659.x)

39. Caudle C, Baskin JM. 1968 The germination pattern of three winter annuals.
Bulletin of the Torrey Botanical Club **95**, 331–335. (doi:10.2307/2483867)

40. Gama-Arachchige NS, Baskin JM, Geneve RL, Baskin CC. 2012 The
autumn effect: timing of physical dormancy break in seeds of two winter
annual species of Geraniaceae by a stepwise process. *Ann Bot* **110**, 637–
651. (doi:10.1093/aob/mcs122)

41. Baskin JM, Baskin CC. 1971 Germination of winter annuals in July and
survival of the seedlings. *Bulletin of the Torrey Botanical Club* **98**, 272–276.
(doi:10.2307/2483627)

42. D'Aguillo MC, Edwards BR, Donohue K. 2019 Can the environment have a
genetic basis? A case study of seedling establishment in Arabidopsis
thaliana. *J Hered* **110**, 467–478. (doi:10.1093/jhered/esz019)

43. Sade N, del Mar Rubio-Wilhelmi M, Umnajkitikorn K, Blumwald E. 2018
Stress-induced senescence and plant tolerance to abiotic stress. *Journal of
Experimental Botany* **69**, 845–853. (doi:10.1093/jxb/erx235)

44. Chaves MM, Maroco J, Pereira JS. 2003 Understanding plant responses to
drought - from genes to the whole plant. *Functional Plant Biology* **30**, 239–
264.

45. Fresnillo Fedorenko DE, Fernández OA, Busso CA, Elia OE. 1996
Phenology of Medicago minima and Erodium cicutarium in semi-arid
Argentina. *Journal of Arid Environments* **33**, 409–416.
(doi:10.1006/jare.1996.0076)

46. Cox JA, Conran JG. 1996 The effect of water stress on the life cycles of
Erodium crinitum Carolin and Erodium cicutarium (L.) L'Hérit. ex Alton
(Geraniaceae). *Australian Journal of Ecology* **21**, 235–240.
(doi:10.1111/j.1442-9993.1996.tb00604.x)

47. Abarca M, Spahn R. 2021 Direct and indirect effects of altered temperature
regimes and phenological mismatches on insect populations. *Current
Opinion in Insect Science* (doi:10.1016/j.cois.2021.04.008)

48. Hamann E, Blevins C, Franks SJ, Jameel MI, Anderson JT. 2021 Climate
change alters plant–herbivore interactions. *New Phytologist* **229**, 1894–
1910. (doi:https://doi.org/10.1111/nph.17036)

49. Kemp R. 1998 Importance of larval foodplant lifespan to British butterflies -
with particular reference to the Brown Argus (Aricia agestis). *Bulletin of the
Amateur Entomologists' Society* **57**, 225–227.

50. IPCC. 2014 Climate Change 2014: Synthesis Report. Contribution of
Working Groups I, II and III to the Fifth Assessment Report of the
Intergovernmental Panel on Climate Change (Core Writing Team, R.K.
Pachauri and L.A. Meyer (eds.)). , 151.

51. Wang J, Guan Y, Wu L, Guan X, Cai W, Huang J, Dong W, Zhang B. 2021
Changing lengths of the four seasons by global warming. *Geophysical*
Research Letters **48**, e2020GL091753.
(doi:https://doi.org/10.1029/2020GL091753)

52. Benning JW, Hufbauer RA, Weiss-Lehman C. 2021 Increasing temporal
variance leads to stable species range limits. , 2021.08.09.455156.
(doi:10.1101/2021.08.09.455156)

53. Salgado AL, DiLeo MF, Saastamoinen M. 2020 Narrow oviposition
preference of an insect herbivore risks survival under conditions of severe
drought. *Functional Ecology* **34**, 1358–1369. (doi:10.1111/1365-
2435.13587)

54. Saastamoinen M *et al.* 2018 Genetics of dispersal. *Biol Rev Camb Philos*
Soc **93**, 574–599. (doi:10.1111/brv.12356)

55. Hanski IA. 2011 Eco-evolutionary spatial dynamics in the Glanville fritillary
butterfly. *PNAS* **108**, 14397–14404. (doi:10.1073/pnas.1110020108)

56. Suggitt AJ *et al.* 2018 Extinction risk from climate change is reduced by
microclimatic buffering. *Nature Clim Change* **8**, 713–717.
(doi:10.1038/s41558-018-0231-9)

57. Oliver T, Roy DB, Hill JK, Brereton T, Thomas CD. 2010 Heterogeneous
landscapes promote population stability. *Ecology Letters* **13**, 473–484.
(doi:10.1111/j.1461-0248.2010.01441.x)

58. Connor EF, Adams-Manson RH, Carr TG, Beck MW. 1994 The effects of
host plant phenology on the demography and population dynamics of the
leaf-mining moth, Cameraria hamadryadella (Lepidoptera: Gracillariidae).
Ecological Entomology **19**, 111–120. (doi:10.1111/j.1365-
2311.1994.tb00400.x)

59. Bennett NL, Severns PM, Parmesan C, Singer MC. 2015 Geographic
mosaics of phenology, host preference, adult size and microhabitat choice
predict butterfly resilience to climate warming. *Oikos* **124**, 41–53.
(doi:10.1111/oik.01490)

Table 1. Parameter estimates (with standard errors) for the final (*final*; selected, most
parsimonious) and null (N) logistic regression models investigating drivers of (a) the
condition of the youngest *Geranium* (M*C*), and (b) the probability of new recruitment of
Geranium (M*R*) in each quadrat during September surveys (2016–2017). Also
presented for comparison are the restricted models (M*X*$_R$) used to check for class bias
in M*X*$_{final}$. All models bar the null contain a subset of fixed effects from: site (S, using
sum contrasts: see SI-1.8), daily mean rainfall (R) or daily mean temperature (T) in the
period between August 1st–September sampling date. Other terms and time periods
were tested and detected in the final candidate set of models (Table S7 and S8). β$_0$ is
the intercept, which accounts for the mean of site effects in all but the null models, *k* is
the number of parameters, *LL* is the log-likelihood of the model and ΔAIC is the ΔAIC
relative to the model with the lowest AIC in each case (Table S7 and S8).

Model	Model parameters					*k*	*LL*	ΔAIC
	β$_0$	S_1	S_2	*R*	*T*			
(a) Condition of youngest G. dissectum								
MC$_{final}$	2.053 (0.535)	5.749 (0.985)	-1.590 (0.551)	5.871 (0.891)	–	4	-61.17	0.13
MC$_N$	0.545 (0.141)	–	–	–	–	1	-143.30	168.38
MC$_R$	1.565 (0.622)	5.071 (1.116)	-1.305 (0.630)	5.611 (1.128)	–	4	*	*
(b) Probability of G. dissectum recruitment								
MR$_{final}$	0.310 (0.354)	4.366 (0.794)	-3.558 (0.718)	–	-3.872 (0.555)	4	-49.07	1.43
MR$_N$	0.352 (0.138)	–	–	–	–	1	-147.78	192.84
MR$_R$	-0.044 (0.495)	4.770 (1.126)	-4.120 (1.017)	–	-4.096 (0.792)	4	*	*

Table 2. Parameter estimates (with standard errors) for the final ($_{final}$; selected, most
parsimonious) and null ($_N$) logistic regression models investigating microclimatic drivers
of (a) quadrat-specific condition indices (MCI), and (b) quadrat-specific phenophase
indices (MPI) of *Geranium* at site G1 in September 2017. Also presented for
comparison are the associated null models (MX_N). All models bar MX_N contain a subset
of fixed effects from: bare ground cover (G), soil moisture (M), mean (T) and minimum
temperature (t). Other terms were tested and detected in the final candidate sets of
models (Tables S9–10). β_0 is the intercept, k is the number of parameters, LL is the
log-likelihood of the model and ΔAIC is the ΔAIC relative to the model with the lowest
AIC in each case (Tables S9–10).

Model	Model parameters					k	LL	ΔAIC
	β_0	G	M	T	t			
(a) Quadrat-specific condition indices								
MCI_{final}	0.496 (0.413)	–	0.810 (0.448)	–	–	2	-15.59	1.88
MCI_N	0.428 (0.380)	–	–	–	–	1	-19.69	8.07
(b) Quadrat-specific phenophase indices								
MPI_{final}	0.366 (0.412)	–	0.889 (0.453)	–	–	2	-14.84	1.57
MPI_N	0.307 (0.376)	–	–	–	–	1	-19.98	9.85

660   **Figures**

662   Figure 1. Map of study sites in England. **+** denotes *Helianthemum* sites (H1–H5, Table
663   S1), **×** denotes Geraniaceae sites (*Geranium*, G1–G3; *Erodium,* E1–E2; Table S1).

Figure 2. (a) Mean (± SD) of site-averaged quadrat-level host plant condition
categorisations (0–3). Starred brackets indicate significant differences in condition
between hosts (Table S4). Condition (0–3; b) and phenophase (c) of the youngest *G.*
dissectum plant in each quadrat across all sites visited during late July–early October
in both 2016 and 2017. Phenophase L: leaf; B: in bud; F: in flower; S: set seed.
Equivalents of (b) and (c) for *Erodium* are available as Figure S3.

Figure 3. Condition index of (a) *Erodium*, (b) *Geranium*, and (c) *Helianthemum* in 2016

(orange shading, closed circles) and 2017 (blue shading, open circles), indicating the

timing of host condition changes relative to the UK-wide emergence of second

generation brown argus adults (solid curves; [33]) and their larvae (dashed curves).

Blue curves represent phenology at southerly sites under cooler (5[th] percentile)

conditions, red curves represent northerly sites under warmer (95^{th} percentile)

conditions. Error bars indicate among-site standard deviation of condition indices.

Figure 4. (a,b) Predicted probabilities with 95% confidence intervals (dashed lines) of

(a) a state of good condition (q = 2 or 3) in the youngest *Geranium* in each quadrat,

and (b) recruitment of *Geranium* in each quadrat. The probabilities (a) increase as a

function of late summer rainfall (Table 1) and (b) decrease as a function of late summer

temperature (MR_{final}, Table 1). Some points have been offset in the x and y planes to

show the raw data; mean daily rainfall ranges between -1.76 and 2.18 (1.29 mm–2.87

mm), and temperature ranges between -1.16 and 1.85 (15.61 °C–18.11 °C). The

condition (c) and phenophase (d) indices of *Geranium* at site G1 increase as a function

of soil moisture. Quadrats with higher soil moisture are more likely to contain high

proportions of good condition new recruits, and low proportions of poor-condition,

reproductive *Geranium*. Soil moisture ranges between -1.58 and 1.83 (14.6–34.4 %).

Soil moisture was recorded at a different time of year to plant condition, so is used as a

relative measure between quadrats.