Recommendations for improving statistical inference in population genomics

Parul Johri[1], Charles F. Aquadro[2], Mark Beaumont[3], Brian Charlesworth[4], Laurent Excoffier[5], Adam Eyre-Walker[6], Peter D. Keightley[7], Michael Lynch[1], Gil McVean[8], Bret A. Payseur[9], Susanne P. Pfeifer[1], Wolfgang Stephan[10], and Jeffrey D. Jensen[1,*]

[1] School of Life Sciences, Arizona State University, Tempe, US
[2] Department of Molecular Biology and Genetics, Cornell University, Ithaca, US
[3] School of Biological Sciences, University of Bristol, Bristol, UK
[4] Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK
[5] Institute of Ecology and Evolution, University of Berne, Berne, CH
[6] School of Life Sciences, University of Sussex, Brighton, UK
[7] Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, Edinburgh, UK
[8] Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK
[9] Laboratory of Genetics, University of Wisconsin-Madison, Madison, US
[10] Natural History Museum, Berlin, DE

[*] corresponding author: jeffrey.d.jensen@asu.edu

keywords: population genetics, population genomics, statistical inference, model-fitting, positive selection, purifying selection, background selection, selective sweeps, genome scans, demography

This paper is dedicated to the memories of Richard Lewontin (1929-2021) and Bill Hill (1940-2021).

ABSTRACT

The field of population genomics has grown rapidly in response to the recent advent of

affordable, large-scale sequencing technologies. As opposed to the situation during the majority

of the 20th century, in which the development of theoretical and statistical population-genetic

insights out-paced the generation of data to which they could be applied, genomic data are now

being produced at a far greater rate than they can be meaningfully analyzed and interpreted. With

this wealth of data has come a tendency to focus on fitting specific (and often rather

idiosyncratic) models to data, at the expense of a careful exploration of the range of possible

underlying evolutionary processes. For example, the approach of directly investigating models of

adaptive evolution in each newly sequenced population or species often neglects the fact that a

thorough characterization of ubiquitous non-adaptive processes is a prerequisite for accurate

inference. We here describe the perils of these tendencies, present our consensus views on

current best practices in population genomic data analysis, and highlight areas of statistical

inference and theory that are in need of further attention. Thereby, we argue for the importance

of defining a biologically relevant baseline model tuned to the details of each new analysis, of

skepticism and scrutiny in interpreting model-fitting results, and of carefully defining

addressable hypotheses and underlying uncertainties.

Introduction

A brief overview

Population genomic inference – the use of data on molecular variation within species and divergence between species to infer evolutionary processes – has become widely embraced and highly utilized in fields including evolutionary biology, ecology, anthropology, agriculture, and medicine. The underlying questions may be demographic in nature, be it estimating the timing of the peopling of the world [1] or of viral transmission in a congenitally infected newborn [2]; alternatively, they may concern the selective history of specific populations, be it identifying mutations that confer cryptic coloration in species adapting to major post-glacial climatic and geological changes [3] or viral drug-resistance to clinical therapeutics [4].

The foundational work allowing for the dissection of these evolutionary processes from levels and patterns of variation and divergence was conducted by Fisher, Wright, and Haldane nearly a century ago (*e.g.*, [5-7]; for a historical overview, see [8]). This work demonstrated the possibility of studying evolution at the genetic level, integrating the revolutionary ideas of Darwin [9] with the turn-of-the-century appreciation of Mendel's [10] research. However, as was famously described by Lewontin [11], this initial theoretical progress during the first half of the 20th century was "like a complex and exquisite machine, designed to process a raw material that no one had succeeded in mining". With the first 'mining' of population-level molecular variation in the 1960s (see [12]), this machine was put to work. The next major steps forward were provided by Kimura and Ohta, who offered a comprehensive framework for studying DNA and protein sequence variation based on these fundamental theoretical insights – the Neutral Theory of Molecular Evolution [13-15] – an advance for which molecular biology also provided support [16]. Despite some claims to the contrary [17], Kimura and Ohta's initial postulates have since

been largely validated [18-19], and have provided a means to interpret observed molecular

variation and divergence within the context of constantly occurring evolutionary processes

including mutation, genetic drift, and purifying selection. While ascribing an important role for

positive selection at the level of phenotypic evolution (consistent with Darwin's initial notions),

the Neutral Theory hypothesizes that at the genetic level beneficial mutations are rare compared

to the much larger input of neutral, nearly neutral, and deleterious mutations that are constantly

raining down on the genomes of all species. Accordingly, episodes of positive selection per

nucleotide are rare compared to genetic drift and purifying selection. However, the significant

effects on evolution at linked sites caused by fitness-altering mutations have been described in

detail in the decades since Kimura's initial formulation of the Neutral Theory [20-22].

With this framework and the availability of datasets to which it could be applied,

statistical approaches for analyzing molecular data began to proliferate, frequently employing

some form of neutral expectation as a null model. A wide range of rather sophisticated statistical

machinery is now available for reconstructing histories of population size change, population

subdivision and migration (*e.g.,* [23-24]), for identifying beneficial mutations based on patterns

associated with selective sweeps (*e.g.,* [25-26]), for quantifying the distribution of fitness effects

(DFE) of newly arising mutations (*e.g.,* [27-28]), as well as for estimating rates of mutation (*e.g.,*

[29-31]) and recombination (*e.g.,* [32-34]). These approaches operate in a variety of statistical

frameworks (see [35-37]), and utilize various aspects of the data – including the frequencies of

variants in a sample (the site frequency spectrum, SFS), associations between variants (linkage

disequilibrium, LD), and/or levels and patterns of between-species divergence at contrasted site

classes (*e.g.,* synonymous versus non-synonymous sites).

Challenges of model choice and parameter fitting

The growing variety of statistical approaches and associated software implementations presents a

dizzying array of choices for any given analysis; although many approaches share the same aims,

there also exist important differences. For example, some approaches require a relatively high

level of coding ability to implement while others may be applied in easy-to-use software

packages; while some are well-tested and justified by population-genetic theory, others are not.

Moreover, even the process of translating raw sequencing data into the allele calls and genotypes

used as input for these approaches is accompanied by uncertainty that depends on sequencing

quality and coverage, availability of a reference genome, and choice of variant calling and

filtering strategies [38-39]. Adding to this complexity, it has become increasingly clear that

demographic estimation may be highly biased when selection and recombination-associated

biased gene conversion are neglected [40-41], whereas estimates of selection intensity and

recombination rate may be highly biased when neglecting demographic effects [42-45]. This

creates a circular problem when commencing any new analysis: one needs information about the

demographic history to estimate parameters of recombination and selection, while at the same

time one needs information about recombination and selection to estimate the demographic

history. An additional challenge, and a frustration for many, is that there is no single 'best

approach'; the correct analysis tools to use, and indeed which questions can be answered at all,

depend entirely on the details of the organism under study [46]. Specifically, biological

parameters that vary among species – including evolutionary parameters (*e.g.*, effective

population size (N_e), mutation rates, recombination rates, and population structure and history),

genome structure (*e.g.*, the distribution of functional sites along the genome), and life history

traits (*e.g.*, mating system) – must all be considered in order to define addressable hypotheses

and optimal approaches.

Beyond these initial considerations, a more difficult issue often emerges. Namely, very

different models may be found to provide a good fit to the observed data (*e.g.*, [47]; see [48] for

a phylogenetic perspective on the topic). In other words, particular parameter combinations may

be found under competing models that are all capable of predicting the observed patterns of

variation. For example, assuming neutrality, one may match an empirical observation at a locus

by fitting the timing, severity, and duration of a population bottleneck; or, alternatively, when

assuming a constant population size, by fitting the rate and mean strength of selective sweeps.

This fact alone implies a simple truism: the ability to fit the parameters of one's preferred model

to data does not alone represent proof of biological reality. Rather, it suggests that this model is

one – out of potentially very many – that represents a viable hypothesis, which should be further

examined via subsequent analyses or experimentation.

Examples abound of enthusiastic promotion of a single preferred model, only to be

tempered by subsequent demonstrations of the fit of alternative and often simpler / more

biologically realistic models. For example, the view that segregating alleles may be commonly

maintained by balancing selection [49] was tempered by the realization that genetic drift is often

a sufficient explanation [14], and the view that genome-wide selective sweeps on standing

variation are pervasive [50-51] was tempered by the realization that neutral population histories

can result in similar patterns [47,52]. While one may readily find such examples of using

episodic or hypothesized processes to fit large-scale data patterns by neglecting to define

expectations arising from common and certain-to-be-occurring processes, determining which

models to evaluate, and how to interpret the fit of a model and its alternatives, are challenges for

all researchers. To better illustrate this point, Figure 1 presents three scenarios (constant

population size with background selection, constant population size with background selection

and selective sweeps, and a population bottleneck with background selection and selective

sweeps), and provides the fit of each of those scenarios to two incorrect models (population size

change assuming strict neutrality, and recurrent selective sweeps assuming constant population

size). As is shown in the figure, each scenario can be well fit by both incorrect models, with

selective sweeps and population bottlenecks generally being confounded, as well as background

selection and population growth, as has been described several times before (*e.g.*, [40, 53-55]).

Methodology

In order to provide a series of examples to accompany key points - as with the above Figure 1 -

both forward-in-time simulations and coalescent simulations were performed for (1) the

inference of demographic history assuming complete neutrality, (2) the inference of positive

selection assuming constant population size, and (3) obtaining test datasets representing different

evolutionary scenarios. While any statistical framework involving model / parameter exploration

and comparison may be consistent with our recommendations, we here utilize approximate

Bayesian computation (ABC) for our examples, as it is a particularly useful framework for

quantifying uncertainty and for exploring complex models.

In all simulations, a chromosomal segment of 99,012 bp was simulated with an intron-

exon-intergenic structure resembling the *D. melanogaster* genome. Each gene comprised five

exons (of 300 bp each) and four introns (of 100 bp each) separated by intergenic regions of

length 1,068 bp. Such a construct resulted in a total of 33 genes across the simulated segment.

Population parameters were chosen to resemble those from *D. melanogaster* populations

following Campos *et al.* [56], assuming an effective population size (N_e) of 10^6 individuals with

a mean mutation rate (μ) of 4.5×10^{-9} per base pair /generation and a mean recombination rate

(r) of 1×10^{-8} per base pair /generation. For computational efficiency, all parameters were

rescaled by a factor of 200.

Modeling and inference of demographic history

A simple demographic history was modeled in which a single population undergoes an

instantaneous change from an ancestral size (N_{anc}) to a current size (N_{cur}), τ generations ago.

Priors for both N_{anc} and N_{cur} were sampled from a loguniform distribution between 10 and

50,000, while priors for the time of change (τ) were sampled from a loguniform distribution

between 10 and N_{cur}. One hundred replicates were simulated for each parameter combination.

Simulations required for ABC were performed in *msprime* v. 0.7.3 [57] assuming complete

neutrality. Mutation and recombination rates were assumed to be constant across the genome and

across replicates.

Modeling and inference of positive selection

A recurrent selective sweep scenario was modeled in which only neutral and beneficial

autosomal mutations were allowed, with simulations performed using *SLiM* v. 3.1 [58]. Introns

and intergenic regions were assumed to be neutral, while exons experienced beneficial mutations

with fitness effects sampled from an exponential distribution with mean *s* for homozygotes,

assuming semi-dominance. The two parameters varied were the mean population-scaled strength

of selection, $\gamma = 2N_{anc}s$, and the proportion of new beneficial mutations, f_{pos}. Priors for these

parameters were sampled from a loguniform distribution such that $\gamma \in [0.1, 10000]$ and $f_{pos} \in$

$[0.00001, 0.01]$. For all parameter combinations, the true rate of beneficial substitutions per site

(d_a) and the true fraction of substitutions due to beneficial mutations (λ, which is related to the α

parameter of Eyre-Walker & Keightley [59]) were calculated using the total number of fixations

(as provided by *SLiM*), with λ observed to range from 0-0.85 depending on the underlying

parameters. Parameter inference was performed for γ and d_a and the corresponding λ was

inferred using $\lambda = \dfrac{d_a}{d_a + ((1 - f_{pos}) \times \mu \times num\ of\ generations)}$, where it was assumed that $1 - f_{pos} \sim 1$.

Populations had a constant size of 5000 diploid individuals with constant mutation and

recombination rates, as specified above. Simulations were run for 100,100 generations (*i.e.,* $20N_e$

+ 100 generations).

ABC

The sample size was set to 100 haploid genomes (or 50 diploid individuals). Under the

demographic and selection models described above, all exonic regions were masked and the

mean and variance (across replicates) of the following summary statistics were calculated:

number of segregating sites, nucleotide site diversity (π), Watterson's theta (θ_W), θ_H, H',

Tajima's D, number of singletons, haplotype number and frequency distribution, and statistics

summarizing LD (r^2, D, D'). All statistics were calculated in non-overlapping sliding windows of

2 kb using *pylibseq* v. 0.2.3 [60]. ABC was performed with the R package "abc" v. 2.1 [61] using

all summary statistics, with "neural net" to account for non-linearity between statistics and

parameters. A 100-fold cross-validation was used to identify the optimum tolerance level, which

was found to be 0.05 (*i.e.,* 5% of the simulations were accepted during ABC inference to

estimate the posterior probability of each parameter). Point estimates of each inferred parameter

were calculated as the weighted medians of the posterior estimates.

Simulations of different evolutionary scenarios as 'true scenarios'

To consider more biologically realistic models and evaluate model violations, a number of

evolutionary scenarios were simulated (using *SLiM*) as follows:

a) Background selection: Exons experienced deleterious mutations modeled by a discrete

244   DFE comprised of four non-overlapping uniform distributions, representing the

245   effectively neutral $(-1 < 2N_{anc}s \leq 0)$, weakly deleterious $(-10 < 2N_{anc}s \leq -1)$,

246   moderately deleterious $(-100 < 2N_{anc}s \leq -10)$, and strongly deleterious $(2N_{anc}s \leq$

247           $-100)$ classes of mutations. All four bins were assumed to contribute equally to new

248   mutations (*i.e.*, 25% of all new mutations belonged to each class of mutation).

249   b) Positive selection: Exons experienced beneficial mutations with $\gamma = 125$ and $f_{pos} =$

250   2.2×10^{-3} (modified from [56]), resulting in $\lambda \approx 0.35$.

251   c) Population size change: A population decline was simulated such that the population

252   declined from 5000 to 100 individuals instantaneously 100 generations ago. A population

253   expansion was similarly simulated with parameters $N_{anc} = 5000$ and $N_{cur} = 10000$. A

254   population bottleneck model was also simulated such that $N_{anc} = N_{cur} = 5000$, and a

255   bottleneck occurred 2000 generations ago with a reduction to 1% of the population size

256   for 100 generations.

257   d) SNP ascertainment: Genotype error was modeled as an inability to detect the true number

258   of singletons when using low-coverage population-genomic data to call variants [38]. To

model this scenario, a random set of singletons, representing a third of all singletons

present in the sample, were removed.

e) Progeny skew: A skew in the offspring distribution (ψ) was modeled such that 5% and

10% of the population was replaced by the offspring of a single individual each

generation ([62], and see [63-64]).

f) Variation in mutation and recombination rates across the genome (*e.g.*, [65-67]): Every

10 kb of the ≈100 kb genomic region considered was assumed to have a different

mutation and recombination rate. For every simulated replicate, these rates were sampled

from a Gaussian distribution with the same mean as above, and a coefficient of variation

of 0.5. Negative values were truncated to 0.

Posterior checks

For the purposes of illustration, an example of posterior checks are provided in Figure 1 (*i.e.,*

showing a simple evaluation of the fit of the inferred posteriors under the incorrect models to

the true scenarios under consideration). Specifically, the mean estimates of the inferred

parameters were used to simulate the "best-fitting model" in *SLiM* v. 3.1 [58]. Exons were

masked and summary statistics were calculated as above in windows of 2 kb using *pylibseq*

v.0.2.3 [60]. In order to simulate the inferred models of positive selection, f_{pos} was

calculated from λ assuming a Wright-Fisher diploid population of size N and a total mutation

rate of μ_{tot} (which for our purpose is the same as μ). Thus, $\mu_b = f_{pos} \times \mu_{tot}$ and $\mu_{neu} =$

$(1 - f_{pos}) \times \mu_{tot}$ where μ_b and μ_{neu} are the beneficial and neutral mutation rates,

respectively. Given a value of λ, and assuming that the distribution of fitness effects of

beneficial mutations is exponential (with mean $\bar{s}$), we calculate f_{pos} as follows:

given that

$$\lambda = \frac{\#of \ beneficial \ subs}{\#of \ beneficial \ subs + \#of \ neutral \ subs} \qquad (1)$$

285   where

$$\#of \ beneficial \ subs = P_{fix} \times L \times 2N\mu_b \qquad (2)$$

287   and

$$\#of \ neutral \ subs = \mu_{neu} \times L \qquad (3)$$

289   where L is the length of the region being considered and P_{fix} is the probability of fixation of

290   beneficial mutations, given by:

$$P_{fix} = \int_0^\infty \frac{(1 - e^{-x})}{(1 - e^{-2Nx})} \left(\frac{e^{-x/\bar{s}}}{\bar{s}}\right) dx \qquad (4)$$

292   Substituting (2) and (3) in (1), and rearranging we get

$$f_{pos} = \frac{\lambda}{(1 - \lambda)P_{fix}2N + \lambda} \qquad (5)$$

295   Integrating (4) in R and substituting it into (5) gives us values of f_{pos}.

297   Statistics were calculated in non-overlapping windows of 2 kb and intervals (CIs) were

298   calculated as the 0.025 and 0.975 quantiles of the distribution of the statistics.

Recommendations

Constructing an appropriate baseline model for population genomic analysis

The somewhat disheartening exercise of fitting incorrect models to data (as depicted in Figure 1)

naturally raises the questions of whether, and if so how, accurate evolutionary inferences can be

extracted from DNA sequences sampled from a population. The first point of importance is that

the starting point for any genomic analysis should be the construction of a biologically relevant

baseline model, which includes the processes that must be occurring and shaping levels and

patterns of variation and divergence across the genome. This model should include mutation,

recombination, and gene conversion (each as applicable), purifying selection acting on functional

regions and its effects on linked variants (*i.e.,* background selection [21, 68-69]), as well as

genetic drift as modulated by, amongst other things, the demographic history and geographic

structure of the population. Depending on the organism of interest, there may be other significant

biological components to include, such as mating system, progeny distributions, ploidy, and so

on (though, for certain questions of interest, some of these biological factors may simply be

included in the resulting effective population size). It is thus helpful to view this baseline model

as being built from the ground up for any new data analysis. Importantly, the point is not that

these many parameters need to be fully understood in a given population in order to perform any

evolutionary inference, but rather that they all require consideration, and that the effects of

uncertainties in their underlying values on downstream inference can be quantified.

However, even prior to considering any biological processes, it is important to investigate

the data themselves. Firstly, there exists an evolutionary variance associated with the myriad of

potential realizations of a stochastic process, as well as the statistical variance introduced by

finite sampling. Secondly, it is not advisable to compare one's empirical observations, which

may include missing data, variant calling or genotyping uncertainty (*e.g.*, effects of low

coverage), masked regions (*e.g.*, regions in which variants were omitted due to low mappability

and/or callability) and so on, against either an analytical or simulated expectation that lacks those

considerations and thus assumes optimal data resolution [70]. The dataset may also involve a

certain ascertainment scheme, either for the variants surveyed [71], or given some pre-defined

criteria for investigating specific genomic regions (*e.g.*, regions representing genomic outliers

with respect to a chosen summary statistic [72]). For the sake of illustration, Figure 2 follows the

same format as Figure 1, but considers two scenarios: population growth with background

selection and selective sweeps, and the same scenario together with data ascertainment (in this

case, an under-calling of the singleton class). As can be seen, due to the changing shape of the

frequency spectra, neglecting to account for this ascertainment can greatly affect inference,

considerably modifying the fit of both the incorrect demographic and incorrect recurrent

selective sweep models to the data.

Hence, if sequencing coverage is such that rare mutations are being excluded from

analysis, due to an inability to accurately differentiate genuine variants from sequencing errors,

the model used for subsequent testing should also ignore these variants. Similarly, if multiple

regions are masked in the empirical analysis due to problems such as alignment difficulties, the

expected patterns of LD that are observable under any given model may be affected.

Furthermore, while the added temporal dimension of time-series data has recently been shown to

be helpful for various aspects of population genetic inference [73-76], such data in no way

sidestep the need for an appropriate baseline model, but simply requires the development of a

baseline that matches the temporal sampling. In sum, as these factors can greatly affect the

power of planned analyses and may introduce biases, the precise details of the dataset (*e.g.*,

region length, extent and location of masked regions, the number of callable sites, and

ascertainment) and study design (*e.g.,* sample size and single time-point versus time-series data)

should be directly matched in the baseline model construction.

Once these concerns have been satisfied, the first biological addition will logically be the

mutation rate and mutational spectrum. For a handful of commonly studied species, both the

mean of, and genomic heterogeneity in, mutation rates have been quantified via mutation-

accumulation lines and/or pedigree studies [77]. However, even for these species, ascertainment

issues remain complicating [78], variation amongst individuals may be substantial [79], and

estimates only represent a temporal snapshot of rates and patterns that are probably changing

over evolutionary time-scales and may be affected by the environment [31, 80]. In organisms

lacking experimental information, often the best available estimates come either from a distantly

related species or from molecular clock-based approaches. Apart from stressing the importance

of implementing either of the experimental approaches in order to further refine mutation-rate

estimates for such a species of interest, it is noteworthy that this uncertainty can also be modeled.

Namely, if proper estimation has been performed in a closely related species, one may quantify

the expected effect on observed levels of variation and divergence of higher and lower rates. The

variation in possible data observations induced by this uncertainty is thus now part of the

underlying model.

The same logic follows for the next parameter addition(s): crossing over / gene

conversion, as applicable for the species in question. For example, for a subset of species, per-

generation crossover rates in cM per Mb have been estimated by comparing genetic maps based

on crosses or pedigrees with physical maps [81-83]. In addition, recombination rates scaled by

the effective population size have also been estimated from patterns of LD (*e.g.*, [84-85]) –

though this approach typically requires assumptions about evolutionary processes that may be

violated (*e.g.,* [42]). As with mutation, the effects on downstream inference arising from the

variety of possible recombination rates – whether estimated for the species of interest or a

closely related species – can be modeled.

The next additions to the baseline model construction are generally associated with the

greatest uncertainty – the demographic history of the population, and the effects of direct and

linked purifying selection. This is a difficult task given the virtually infinite number of potential

demographic hypotheses (*e.g.*, [86]); furthermore, the interaction of selection with demography

is inherently non-trivial and difficult to treat (*e.g.*, [55, 87-88]). This realization continues to

motivate attempts to jointly estimate the parameters of population history together with the DFE

of neutral, nearly neutral, weakly deleterious and strongly deleterious mutations – a distribution

which is often estimated in both continuous and discrete forms [89]. One of the first important

advances in this area used putatively neutral synonymous sites to estimate changes in population

size based on patterns in the SFS and conditioned on that demography to fit a DFE to non-

synonymous sites, which presumably experience considerable purifying selection [90-92]. This

stepwise approach may become problematic, however, for organisms in which synonymous sites

are not themselves neutral [93-95], or when the SFS of synonymous sites is affected by

background selection, which is probably the case generally given their close linkage to directly

selected non-synonymous sites ([41], and see [96-97]).

In an attempt to address some of these concerns, Johri *et al.* [44] recently developed an

ABC approach that relaxes the assumption of synonymous site neutrality and corrects for

background selection effects by simultaneously estimating parameters of the DFE alongside

population history. The posterior distributions of the parameters estimated by this approach in

any given data application (*i.e.*, characterizing the uncertainty of inference) represent a logical

treatment of population size change and purifying / background selection for the purposes of

inclusion within this evolutionarily relevant baseline model. That said, the demographic model in

this implementation is highly simplified, and extensions are needed to account for more complex

population histories. In particular, estimation biases that may be expected owing to the neglect of

cryptic population structure and migration, and indeed the feasibility of co-estimating population

size change and the DFE together with population structure and migration within this framework,

all remain in need of further investigation. While such simulation-based inference (see [98]),

including ABC, provides one promising platform for joint estimation of demographic history and

selection, progress on this front has been made using alternative frameworks as well [99-100],

and developing analytical expectations under these complex models should remain as the

ultimate, if distant, goal. Alternatively, in functionally sparse genomes with sufficiently high

rates of recombination, such that assumptions of strict neutrality are viable for some genomic

regions, multiple well-performing approaches have been developed for estimating the parameters

of much more complex demographic models (*e.g.*, [101-104]). In organisms for which such

approaches are applicable (*e.g.*, certain large, coding sequence sparse vertebrate and land-plant

genomes), this intergenic demographic estimation assuming strict neutrality may helpfully be

compared to estimates derived from data in or near coding regions that account for the effects of

direct and linked purifying selection [41, 44, 105]. For newly studied species lacking functional

annotation and information about coding density, following the joint estimation procedure would

remain as the more satisfactory strategy in order to account for possible background selection

effects.

Quantifying uncertainty in model choice and parameter estimation, investigating potential model

violations, and defining answerable questions

One of the useful aspects of these types of analyses is the ability to incorporate uncertainty in

underlying parameters under relatively complex models, in order to determine the impact of such

uncertainty on downstream inference. The computational burden of incorporating variability in

mutation and recombination rate estimates, or drawing from the confidence- or credibility-

intervals of demographic or DFE parameters, can be met with multiple highly flexible simulation

tools [58, 106-107]. These are also useful programs for investigating potential model violations

that may be of consequence. For example, if a given analysis for detecting population structure

assumes an absence of gene flow, it is possible to begin with one's constructed baseline model,

add migration parameters to the model in order to determine the effects of varying rates and

directions of migration on the summary statistics being utilized in the empirical analysis, and

thereby quantify how a violation of that assumption may affect the subsequent conclusions.

Similarly, if an analysis assumes the Kingman coalescent (*e.g.*, a small progeny distribution such

that at most one coalescent event occurs per generation), but the organism in question could

violate this assumption (*i.e.*, with the large progeny number distributions associated with many

plants, viruses, and marine spawners, or simply owing to the relatively wide array of

evolutionary processes that may similarly lead to multiple merger coalescent events), these

distributions may too be modeled in order to quantify potential downstream mis-inference.

To illustrate this point, Figure 3 considers two scenarios of constant population size and

strict neutrality but with differing degrees of progeny skew, to demonstrate that a violation of

this sort that is not corrected for may result in severely under-estimated population sizes as well

as the false inference of high rates of strong selective sweeps. In this case, the mis-inference

arises from the reduction in contributing ancestors under these models, as well as to the fact that

neutral progeny skew and selective sweeps can both generate multiple-merger events [63-64,

108-109]. Similarly, one may investigate the assumptions of constant mutation or recombination

rates when they are in reality variable. As shown in Figure 4, when these rates are assumed to be

constant as is common practice, but in reality vary across the genomic region under

investigation, the fit of the (incorrect) demographic and selection models considered may again

be substantially modified. Notably, this rate heterogeneity may inflate the inferred strength of

selective sweeps. While Figures 3 and 4 serve as examples, the same investigations may be made

for cases such as a fixed selective effect when there is in reality a distribution, independent

neutral variants when there is in reality linkage disequilibrium, panmixia when there is in reality

population structure, and so on. Simply put, even if a particular biological process / parameter is

not being directly estimated, its consequences can nonetheless be explored.

As detailed in Figure 5, with such a model incorporating both biological and stochastic

variance as well as statistical uncertainty in parameter estimates, and with an understanding of

the role of likely model violations, one may investigate which additional questions / hypotheses

can be addressed with the data at hand. By using a simulation approach starting with the baseline

model and adding hypothesized processes, it is possible to quantify the extent to which models,

and the parameters underlying those models, may be differentiated and which result in

overlapping or indistinguishable patterns in the data (*e.g.*, [110]). For example, if the goal of a

given study is to identify recent beneficial fixations in a genome – be they potentially associated

with high-altitude adaptation in humans, crypsis in mice, or drug-resistance in a virus – one may

begin with the baseline model and simulate selective sweeps under that model. As illustrated in

Figure 6, by varying the strengths, rates, ages, dominance and epistasis coefficients of beneficial

mutations, the patterns in the SFS, LD, and/or divergence that may differentiate the addition of

such selective sweep parameters from the baseline expectations can be quantified. Moreover, any

intended empirical analyses can be evaluated using simulated data (*i.e.*, the baseline, compared

to the baseline + the hypothesis) to define the power and false-positive rates associated. If the

differences in resulting patterns cannot be distinguished from the expected variance under the

baseline model (in other words, if the power and false-positive rate of the analyses are not

favorable), the hypothesis is not addressable with the data at hand (*e.g.*, [54]). If the results are

favorable, this analysis can further quantify the extent to which the hypothesis may be tested;

perhaps only selective sweeps from rare mutations with selective effects greater than 1% and that

have fixed within the last 0.1 N_e generations are detectable (see [111-112]), and any others could

not be statistically distinguished from expected patterns under the baseline model. Hence, such

an exercise provides a critically essential key for interpreting the resulting data analysis.

A consideration of alternative strategies

In this regard, it is worth mentioning two common approaches that may be viewed as alternatives

to the strategy that we recommend. The first tactic concerns identifying patterns of variation that

are uniquely and exclusively associated with one particular process, the presence of which could

support that model regardless of the various underlying processes and details composing the

baseline. For example, Fay & Wu's [113] *H*-statistic, capturing an expected pattern of high-

frequency derived alleles generated by a selective sweep with recombination, was initially

proposed as a powerful statistic for differentiating selective sweep effects from alternative

models. Results from the initial application of the *H*-statistic were interpreted as evidence of

widespread positive selection in the genome of *Drosophila melanogaster*. However, Przeworski

[112] subsequently demonstrated that the statistic was characterized by low power to detect

positive selection, and that significant values could readily be generated under multiple neutral

demographic models. The composite likelihood framework of Kim & Stephan [111] provided a

significant improvement by incorporating multiple predictions of a selective sweep model, and

was subsequently built upon by Nielsen *et al.* [114] in proposing the SweepFinder approach.

However, Jensen *et al*. [115] described low power and high false-positive rates under certain

neutral demographic models. The particular pattern of LD generated by a beneficial fixation with

recombination described by Kim & Nielsen [116] and Stephan *et al*. [117] (and see [118]), was

also found to be produced under an (albeit more limited) range of severe neutral population

bottlenecks [119-120].

The point here is that the statistics themselves represent important tools for studying

patterns of variation and are useful for visualizing multiple aspects of the data, but in any given

empirical application they are impossible to interpret without the definition of an appropriate

baseline model and related power and false-positive rates. Thus, the search for a pattern unique

to a single evolutionary process is not a work-around, and historically such patterns rarely turn

out to be process specific after further investigation. Even if a 'bullet-proof' test were to be

someday constructed, it would not be possible to establish its utility without appropriate

modeling, an examination of model violations, and extensive power / sensitivity-specificity

analyses. But in reality, the simple fact is that some test statistics and estimation procedures

perform well under certain scenarios, but not under others.

The second common strategy involves summarizing empirical distributions of a given

statistic, and assuming that outliers of that distribution represent the action of a process of

interest, such as positive selection (*e.g.,* [121]). However, such an approach is problematic. To

begin with, any distribution has outliers, and there will always exist a 5% or 1% tail for a chosen

statistic under a given model. Consequently, a fit baseline model remains necessary to determine

whether the observed empirical outliers are of an unexpected severity, and if the baseline model

together with the hypothesized process has, for example, a significantly improved likelihood.

Moreover, only by considering the hypothesized process within the context of the baseline model

can one determine whether affected loci (*e.g.*, those subject to recent sweeps) would even be

expected to reside in the tails of the chosen statistical distribution, which is far from a given [72,

122]. As such, approaches which may not necessarily require a defined baseline model in order

to perform the initial analyses (*e.g.,* [114]), nonetheless require such modeling to accurately

define expectations, power and false-positive rates, and thus to interpret the significance of

observed empirical outliers. For these reasons, the approach for which we advocate remains

essential. As the appropriate baseline evolutionary model may differ strongly by organism and

population, this performance must be carefully defined and quantified for each empirical analysis

in order to accurately interpret results.

Conclusions

When it comes to evolutionary analyses, wanting to answer a question is not necessarily

equivalent to being able to answer it. The ability of population genomics to address a hypothesis

of interest with a given dataset is something that must be demonstrated, and this may be achieved

by constructing a model composed of common biological and evolutionary processes, including

the uncertainty in those underlying parameters, as well as the specific features of the dataset at

hand. The variation in possible observational outcomes associated with a chosen baseline model,

and the ability to distinguish an hypothesized additional evolutionary process from such

'background noise', are both quantifiable. Furthermore, even if the model were correct, there

exists a limit on the precision of estimation imposed by the evolutionary variance in population

statistics that requires description, and which no amount of sampling can remove.

Demonstrating that multiple models, and/or considerable parameter space within a model,

are compatible with the data need not be viewed as a negative or weak finding. Quite the

contrary – the honest presentation of such results motivates future theoretical, experimental, and

empirical developments and analyses, which can further refine the list of competing hypotheses,

and this article contains many citations that have succeeded in doing this. At the same time, this

analysis can define which degrees of uncertainty are most damaging (*e.g.*, Figures 3 and 4), also

highlighting the simple fact that organisms in which basic biological processes have been better

characterized are amenable to a wider range of potential evolutionary analyses. The impact of

uncertainty in these parameters in non-model organisms may motivate taking a step back to first

better characterize the basic biological processes such as mutation rates and spectra via mutation-

accumulation lines or pedigree studies, in order to improve resolution on the primary question of

interest.

Importantly, the framework we describe will also generally identify many models and

parameter realizations that are in fact inconsistent with the observed data. This 'ruling-out'

process can often be just as useful as model-fitting, and rejecting possible hypotheses is

frequently the more robust exercise of the two. The value of this narrowing down, rather than the

enthusiastic promotion of individual scenarios, is worthy of heightened appreciation.

Nevertheless, all models should not be viewed equally. Decades of work supporting the central

tenets of the Neutral Theory [19], high-quality experimental and computational work quantifying

mutation and recombination rates [77-79, 83-84, 123], constantly improving experimental and

theoretical approaches to quantify the neutral and deleterious DFE from natural population,

mutation-accumulation, or directed mutagenesis data [44, 90, 124-126], and historical knowledge

(*e.g.*, anthropological, ecological, clinical) of population size change or structure – combined

with the fact that all of these factors may strongly shape observed levels and patterns of variation

and divergence – justify their role in comprising the appropriate baseline model for genomic

analysis.

Given this, and particularly after accounting for the inflation of variance contributed by

uncertainty in relevant parameters, potential model violations, as well as the quantity and quality

of data available in any given analysis, it will often be the case that many hypotheses of interest

may not be addressable with the dataset and knowledge at hand. However, recognizing that a

question cannot be accurately answered, and defining the conditions under which it could

become answerable, should be preferred over making unfounded and thus misleading claims.

Consistent with this call for caution however, it should equally be emphasized that the fit of a

baseline model to data is certainly not inherent evidence that the model encompasses all relevant

processes shaping the population. In reality, it is virtually guaranteed not to be all-encompassing,

and building these models involves simplifying more complex processes (for a helpful and more

general perspective, see [127]). When an additional process cannot be satisfactorily detected, that

may rather be viewed as a statement about statistical identifiability – the inability to distinguish a

hypothesized process from other processes that are known to be acting – and in such scenarios,

absence of evidence need not be taken as evidence of absence.

While the many considerations we describe may appear daunting, it is our hope that these

recommendations may serve as a useful roadmap for future data analyses in population

genomics, one that may inform not only the perspectives of authors, but also that of reviewers

and editors as well. Helpfully, these strategies can save considerable time, money, and effort

prior to the start of empirical data handling, by determining which questions are accessible to the

researcher. If a question is addressable, this preliminary analysis can additionally define what

types of data are needed; for example, the number of variants or sample size necessary to obtain

sufficient power, or how alternative data collections (*e.g.*, temporal samples) could improve

resolution. This further highlights the value of defining specific hypotheses and of studying

specific patterns as opposed to running a general suite of software on each new dataset in the

hopes of identifying something of interest – namely, one cannot define the power of a study to

address an unformulated question. Such hypothesis-driven population genomics has resulted in a

number of success stories over the past decade; systems in which specific hypotheses were

formed, data was collected for the purpose, detailed population genomic analyses were designed,

and ultimately important insights were gained about the evolutionary history of the population in

question (*e.g.*, the study of cryptic coloration has proven fruitful in this regard [3]). One feature

common to these studies is interdisciplinarity: the utilization of population genetic theory and

inference as described here, combined with classical genetic crosses, large-scale field studies,

and genetic manipulation in order to connect genotype to phenotype to fitness and to validate

statistical inference. Importantly however, without a population genetic framework for defining

hypotheses, quantifying processes contributing to observed variation and divergence, evaluating

and distinguishing amongst competing models, and defining uncertainty and potential biases, the

observations remain merely descriptive.

ACKNOWLEDGEMENTS

This work was funded by National Institutes of Health grants R01GM135899 and

R35GM139383 to JDJ. We would like to thank Nick Barton, Matt Dean, Fabian Freund, Ryan

Gutenkunst, Mark Kirkpatrick, Sarah Marion, Mohamed Noor, Sally Otto, Kevin Thornton, and

John Wakeley for helpful comments and suggestions.

607
608
609

CITATIONS

[1]   Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E. Tracing the peopling of the world through genomics. Nature 2017;541(7637): 302-310.

[2]   Renzette N, Gibson L, Jensen JD, Kowalik TF. Human cytomegalovirus intrahost evolution - a new avenue for understanding and controlling herpesvirus infections. Curr Opin Virol 2014;8: 109-115.

[3]   Harris RB, Irwin K, Jones MR, Laurent S, Barrett RDH, Nachman MW, et al. The population genetics of crypsis in vertebrates: recent insights from mice, hares, and lizards. Heredity 2020;124(1): 1-14.

[4]   Irwin KK, Renzette N, Kowalik TF, Jensen JD. Antiviral drug resistance as an adaptive process. Virus Evol 2016;2(1): vew014.

[5]   Fisher RA. The genetical theory of natural selection. Clarendon Press, Oxford, UK; 1930.

[6]   Wright S. Evolution in Mendelian populations. Genetics 1931;16(2): 97-159.

[7]   Haldane JBS. The causes of evolution. Longmans, London, UK; 1932.

[8]   Provine WB. The origins of theoretical population genetics. University of Chicago Press; 1971.

[9]   Darwin C. On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life. John Murray, London, UK; 1859.

[10]   Mendel G. Versuche über Pflanzenhybriden. Verh Naturforsch Ver Brünn 1866;4: 3-47.

[11]   Lewontin RC. The genetic basis of evolutionary change. Columbia Univ. Press, New York; 1974.

[12]   Lewontin RC. Twenty-five years ago in Genetics: electrophoresis in the development of evolutionary genetics: milestone or millstone? Genetics 1991;128(4): 657-662.

[13]   Kimura M. Evolutionary rate at the molecular level. Nature 1968;217(5129): 624-626.

[14]   Kimura M. The neutral theory of molecular evolution. Cambridge Univ. Press, Cambridge; 1983.

[15]   Ohta T. Slightly deleterious mutant substitutions in evolution. Nature 1973;246(5428): 96-98.

646   [16]   King JL, Jukes TH. Non-Darwinian evolution. Science 1969;164(3881): 788-798.

647   [17]   Kern AD, Hahn MW. The neutral theory in light of natural selection. Mol Biol Evol
648   2018;35(6): 1366-1371.

649   [18]   Walsh B, Lynch M. Evolution and selection of quantitative traits. Oxford University
650   Press, Oxford; 2018.

651   [19]   Jensen JD, Payseur BA, Stephan W, Aquadro CF, Lynch M, Charlesworth D, et al. The
652   importance of the Neutral Theory in 1968 and 50 years on: a response to Kern & Hahn
653   2018. Evolution 2019;73(1): 111-114.
654
655   [20]   Maynard Smith J, Haigh J. The hitch-hiking effect of a favourable gene. Genet Res
656   1974;23(1): 23-25.

657   [21]   Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on
658   neutral molecular variation. Genetics 1993;134(4): 1289-1303.

659   [22]   Charlesworth B, Jensen JD. The effects of selection at linked sites on patterns of genetic
660   variability. Annu Rev Ecol Evol Syst 2021;52: 177-197.

661   [23]   Ray N, Excoffier L. Inferring past demography using spatially explicit population genetic
662   models. Hum Biol 2009;81(2-3): 141-157.

663   [24]   Beichman AC, Huerta-Sanchez E, Lohmueller KE. Using genomic data to infer historic
664   population dynamics of non-model organisms. Annu Rev Ecol Evol Syst 2018;49: 433-
665   456.
666
667   [25]   Booker TR, Jackson BC, Keightley PD. Detecting positive selection in the genome. BMC
668   Biol 2017;15(1): 98.

669   [26]   Stephan W. Selective sweeps. Genetics 2019;211(1): 5-13.

670   [27]   Eyre-Walker A, Keightley PD. The distribution of fitness effects of new mutations. Nat
671   Rev Genet 2007;8(8): 610-618.
672
673   [28]   Bank C, Foll M, Ferrer-Admetlla A, Ewing G, Jensen JD. Thinking too positive?
674   Revisiting current methods in population genetic selection inference. Trends
675   Genet 2014;30(12): 540-546.
676
677   [29]   Keightley PD, Halligan DL. Analysis and implications of mutational variation. Genetica
678   2009;136(2): 359-369.
679
680   [30]   Keightley PD. Rates and fitness consequences of new mutations in humans. Genetics
681   2012;190(2): 295-304.

682   [31]   Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, et al. Genetic drift,
683   selection and the evolution of the mutation rate. Nat Rev Genet 2016;17(11): 704-714.
684

685   [32]   Stumpf MP, McVean GA. Estimating recombination rates from population-genetic data.
686   Nat Rev Genet 2003;4(12): 959-968.
687

688   [33]   Auton A, Fledel-Alon A, Pfeifer SP, Venn O, Ségurel L, Street T, et al. A fine-scale
689   chimpanzee genetic map from population sequencing. Science 2012;336(6078): 193-198.
690

691   [34]   Spence JP, Song YS. Inference and analysis of population-specific fine-scale
692   recombination maps across 26 diverse human populations. Sci Adv 2019;5(10):
693   eaaw9206.

694   [35]   Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population
695   genetics. Genetics 2002;162(4): 2025-2035.

696   [36]   Beaumont MA, Rannala B. The Bayesian revolution in genetics. Nat Rev Genet
697   2004;5(4): 251-261.
698

699   [37]   Schraiber JG, Akey JM. Methods and models for unravelling human evolutionary history.
700   Nat Rev Genet 2015;16(12): 727-740.
701

702   [38]   Han E, Sinsheimer JS, Novembre J. Characterizing bias in population genetic inferences
703   from low-coverage sequencing data. Mol Biol Evol 2014;31(3): 723-735.
704

705   [39]   Pfeifer SP. Studying mutation rate evolution in primates – the effects of computational
706   pipeline and parameter choices. GigaScience 2021;10(10): giab069.
707

708   [40]   Ewing G, Jensen JD. The consequences of not accounting for background selection in
709   demographic inference. Mol Ecol 2016;25(1): 135-141.
710

711   [41]   Pouyet F, Aeschbacher S, Thiery A, Excoffier L. Background selection and biased gene
712   conversion affect more than 95% of the human genome and bias demographic inferences.
713   eLife 2018;7: e36317.
714

715   [42]   Dapper AL, Payseur BA. Effects of demographic history on the detection of
716   recombination hotspots from linkage disequilibrium. Mol Biol Evol 2018;35(2): 335-353.
717

718   [43]   Rousselle M, Maeva M, Nabholz B, Bataillon T, Galtier N. Overestimation of the
719   adaptive substitution rate in fluctuating populations. Biol Lett 2018;14(5): 20180055.
720

721   [44]   Johri P, Charlesworth B, Jensen JD. Towards an evolutionarily appropriate null model:
722   jointly inferring demography and purifying selection. Genetics 2020;215(1): 173-192.
723

724   [45]   Samuk K, Noor MAF. Gene flow biases population genetic inference of recombination
725   rate. biorxiv 2021.
726   https://www.biorxiv.org/content/10.1101/2021.09.26.461846v1.full.pdf
727

728   [46]   Myers S, Fefferman C, Patterson N. Can one learn history from the allelic spectrum?
729   Theor Popul Biol 2008;73(3): 342-348.
730

731   [47]   Harris RB, Sackman A, Jensen JD. On the unfounded enthusiasm for soft selective
732   sweeps II: examining recent evidence from humans, flies, and viruses. PLoS Genet
733   2018;14(12): e1007859.

734   [48]   Louca S, Pennell MW. Extant timetrees are consistent with a myriad of diversification
735   histories. Nature 2020;580(7804): 502-505.

736   [49]   Ford EB. Ecological genetics. Chapman and Hall, London, UK; 1975.

737   [50]   Garud N, Messer P, Buzbas E, Petrov D. Recent selective sweeps in North American
738   *Drosophila melanogaster* show signatures of soft sweeps. PLoS Genet 2015;11(2):
739   e1005004.

740   [51]   Schrider DR, Kern AD. Soft sweeps are the dominant mode of adaptation in the human
741   genome. Mol Biol Evol 2017;34(8): 1863-1877.

742   [52]   Johri P, Stephan W, Jensen JD. Soft selective sweeps: addressing new definitions,
743   evaluating competing models, and interpreting empirical outliers. PLoS Genet 2022;
744   18(2): e1010022.

745   [53]   Barton NH. Genetic hitchhiking. Phil Trans R Soc B 2000;355(1403): 1553-1562.
746

747   [54]   Poh YP, Domingues V, Hoekstra HE, Jensen JD. On the prospect of identifying adaptive
748   loci in recently bottlenecked populations. PLoS One 2014;9(11): e110579.
749

750   [55]   Johri P, Riall K, Becher H, Excoffier L, Charlesworth B, Jensen JD. The impact of
751   purifying and background selection on the inference of population history: problems and
752   prospects. Mol Biol Evol 2021;38(7): 2986-3003.
753

754   [56]   Campos JL, Charlesworth B. The effects on neutral variability of recurrent selective
755   sweeps and background selection. Genetics 2019;212(1): 287-303.
756

757   [57]   Kelleher J, Etheridge AM, McVean G. Efficient coalescent simulation and
758   genealogical analysis for large sample sizes. PLoS Comput Biol 2016;12(5): e1004842.
759

760   [58]   Haller BC, Messer PW. SLiM 3: Forward genetic simulations beyond the Wright–Fisher
761   model. Mol Biol Evol 2019;36(3): 632-637.
762

763 [59] Eyre-Walker A, Keightley PD. Estimating the rate of adaptive molecular evolution in the
764 presence of slightly deleterious mutations and population size change. Mol Biol Evol
765 2009;26(9): 2097-2108.

766 [60] Thornton K. Libsequence: a C++ class library for evolutionary genetic analysis.
767 Bioinformatics 2003;19(17): 2325–2327.
768

769 [61] Csilléry K, François O, Blum M. abc: an R package for approximate Bayesian
770 computation (ABC). Methods Ecol Evol 2012;3: 475-479.
771

772 [62] Eldon B, Wakeley J. Coalescent processes when the distribution of offspring
773 number among individuals is highly skewed. Genetics 2006;172(4): 2621-2633.
774

775 [63] Matuszewski M, Hildebrandt ME, Achaz G, Jensen JD. Coalescent processes with
776 skewed offspring distributions and non-equilibrium demography. Genetics 2018;208(1):
777 323-338.
778

779 [64] Sackman A, Harris RB, Jensen JD. Inferring demography and selection in organisms
780 characterized by skewed offspring distributions. Genetics 2019;211(3): 1019-1028.
781

782 [65] McVean G, Myers S, Hunt S, Deloukas P, Bentley D, Donnelly P. The fine-scale
783 structure of recombination rate variation in the human genome. Science 2004;304(5670):
784 581-584.
785

786 [66] Chan AH, Jenkins P, Song Y. Genome-wide fine-scale recombination rate variation in
787 *Drosophila melanogaster*. PLoS Genet 2012;8(12): e1003090.
788

[67] Penalba JV, Wolf JB. From molecules to populations: appreciating and estimating
recombination rate variation. Nat Rev Genet 2020;21(8): 476-492.

[68] Charlesworth D, Charlesworth B, Morgan MT. The pattern of neutral molecular variation
under the background selection model. Genetics 1995;141(4): 1619-1632.

[69] Charlesworth B. Background selection 20 years on. The Wilhelmine E. Key 2012
invitational lecture. J Hered 2013;104(2): 161-171.

[70] Pfeifer SP. From next-generation resequencing reads to a high quality variant data
set. Heredity 2017;118(2): 111-124.
797

798 [71] Nielsen R. Population genetic analysis of ascertained SNP data. Hum Genomics
799 2004;1(3): 218-224.
800

801 [72] Thornton KR, Jensen JD. Controlling the false positive rate in multi-locus genome scans
802 for selection. Genetics 2007;175(2): 737-750.
803

804 [73] Malaspinas AS, Malaspinas O, Evans SN, Slatkin M. Estimating allele age and selection
805   coefficient from time-serial data. Genetics 2012;192(2): 599-607.
806

[74] Foll M, Shim H, Jensen JD. A Wright-Fisher ABC-based approach for inferring per-site
effective population sizes and selection coefficients from time-sampled data. Mol Ecol
Res 2015;15(1): 87-98.

[75] Ferrer-Admetlla A, Leuenberger C, Jensen JD, Wegmann D. An approximate Markov
model for the Wright-Fisher diffusion and its application to time series data. Genetics
2016;203(2): 831-846.

[76] Lynch M, Ho WC. The limits to estimating population-genetic parameters with temporal
data. Gen Biol Evol 2020;12(4): 443-455.
815

816 [77] Pfeifer SP. Spontaneous mutation rates. In *The Molecular Evolutionary Clock. Theory*
817   *and Practice.* Springer Nature; 2020.
818

819 [78] Smith TCA, Arndt PF, Eyre-Walker A. Large scale variation in the rate of germ-line *de*
820   *novo* mutations, base composition, divergence and diversity in humans. PLoS Genet
821   2018;14(3): e1007254.
822

823 [79] Ness RW, Morgan AD, Radhakrishnan V, Colegrave N, Keightley PD. Extensive de
824   novo mutation rate variation between individuals and across the genome of
825   *Chlamydomonas reinhardtii*. Genome Res 2015;25(11): 1739-1749.
826

[80] Maddamsetti R, Grant NA. Divergent evolution of mutation rates and biases in the long-
term evolution experiment with *Escherichia coli*. Gen Biol Evol 2020;12(9): 1591-1603.

[81] Kong A, Gudbjartsson DF, Sainz J, Jonsdottir G, Gudjonsson S, Richardsson B, et
al. 2002. A high-resolution recombination map of the human genome. Nat Genet
2002;31(3): 241-247.

[82] Cox A, Ackert-Bicknell C, Dumont B, Ding Y, Tzenova Bell J, Brockmann G, et al. A
new standard genetic map for the laboratory mouse. Genetics 2009;182(4): 1335-1344.

[83] Comeron JM, Ratnappan R, Bailin S. The many landscapes of recombination in
Drosophila melanogaster. PLoS Genet 2012;8(10): e1002905.

[84] Auton A, McVean G. Estimating recombination rates from genetic variation in humans.
Methods Mol Biol 2012;856: 217-237.
838

[85] Pfeifer SP. A fine-scale genetic map for vervet monkeys. Mol Biol Evol 2020;37(7):
1855-1865.

[86] Chikhi L, Sousa VC, Luisi P, Goossens B, Beaumont MA. The confounding effects of
population structure, genetic diversity and the sampling scheme on the detection and
quantification of population size change. Genetics 2010;186(3): 983-995.

[87] Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. On the accumulation of
deleterious mutations during range expansions. Mol Ecol 2013;22(24): 5972-5982.
847
848 [88] Peischl S, Kirkpatrick M, Excoffier L. Expansion load and the evolutionary
849   dynamics of a species range. Am Nat 2015;185(4): E81-E93.
850
[89] Johri P, Eyre-Walker A, Gutenkunst RN, Lohmueller KE, Jensen JD. On the prospect of
achieving accurate joint estimation of selection with population history. In revision, Gen
Biol Evol.

[90] Keightley PD, Eyre-Walker A. Joint inference of the distribution of fitness effects of
deleterious mutations and population demography based on nucleotide polymorphism
frequencies. Genetics 2007;177(4): 2251-2261.

[91] Schneider A, Charlesworth B, Eyre-Walker A, Keightley PD. A method for
inferring the rate of occurrence and fitness effects of advantageous mutations. Genetics
2011;189(4): 1427-1437.

[92] Lynch M. The origins of genome architecture. Sinauer Associates, Sunderland, MA;
2007.

[93] Singh ND, Bauer DuMont VL, Hubisz MJ, Nielsen R, Aquadro CF. Patterns of mutation
and selection at synonymous sites in *Drosophila*. Mol Biol Evol 2007;24(12): 2687-
2697.

[94] Zeng K, Charlesworth B. Studying patterns of recent evolution at synonymous sites and
intronic sites in *Drosophila melanogaster*. J Mol Evol 2010;70(1): 116-128.

[95] Choi JY, Aquadro CF. Recent and long term selection across synonymous sites in
Drosophila ananassae. J Mol Evol 2016;83(1-2): 50-60.

[96] Comeron JM. Background selection as baseline for nucleotide variation across the
Drosophila genome. PLoS Genet. 2014;10(6): e1004434.

[97] Comeron JM. Background selection as a null hypothesis in population genomics:
insights and challenges from *Drosophila* studies. Phil Trans R Soc B 2017;372(1736):
20160471.

[98] Cranmer K, Brehmer J, Louppe G. The frontier of simulation-based inference. Proc Natl
Acad Sci USA 2020;117(48): 30055-30062.
876

877   [99]   Williamson SH, Hernandez R, Fledel-Alon A, Zhu L, Nielsen R, Bustamante CD.
878   Simultaneous inference of selection and population growth from patterns of variation in
879   the human genome. Proc Natl Acad Sci USA 2005;102(22): 7882-7887.

880   [100] Ragsdale A, Moreau C, Gravel S. Genomic inference using diffusion models and the
881   allele frequency spectrum. Curr Opin Gen Deve 2018;53: 140-147.

882   [101] Gutenkunst R, Hernandez R, Williamson S, Bustamante CD. Inferring the joint
883   demographic history of multiple populations from multidimensional SNP data. PLoS
884   Genet 2009;5(10): e1000695.

885   [102] Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. Robust demographic
886   inference from genomic and SNP data. PLoS Genet 2013;9(10): e1003905.

887   [103] Kelleher J, Wong Y, Wohns AW, Fadil C, Albers PK, McVean G. Inferring whole-
888   genome histories in large population datasets. Nat Genet 2019;51(9): 1330-1338.

889   [104] Steinrücken M, Kamm J, Spence J, Song YS. Inference of complex population histories
890   using whole-genome sequences from multiple populations. Proc Natl Acad Sci USA
891   2019;116(34): 17115-17120.
892
893   [105] Torres R, Szpiech Z, Hernandez RD. Human demographic history has amplified the
894   effects of background selection across the genome. PLoS Genet 2018;14(6): e1007387.
895
896   [106] Thornton KR. A C++ template library for efficient forward-time population genetic
897   simulation of large populations. Genetics 2014;198(1): 157-166.
898
899   [107] Kelleher J, Thornton K, Ashander J, Ralph P. Efficient pedigree recording for fast
900   population genetics simulation. PLoS Comput. Biol 2018;14(11): e1006581.
901
902   [108] Durrett R, Schweinsberg J. Approximating selective sweeps. Theor Popul Biol
903   2004;66(2): 129-138.
904
905   [109] Hallatschek O. Selection-like biases emerge in population models with recurrent jackpot
906   events. Genetics 2018;210(3): 1053-1073.

907   [110] Lapierre M, Lambert A, Achaz G. Accuracy of demographic inference from the site
908   frequency spectrum: the case of the Yoruba population. Genetics 2017;206(1): 439-449.

909   [111] Kim Y, Stephan W. Detecting a local signature of genetic hitchhiking along a
910   recombining chromosome. Genetics 2002;160(2): 765-777.

911   [112] Przeworski M. The signature of positive selection at randomly chosen loci.
912   Genetics 2002;160(3): 1179-1189.

[113] Fay J, Wu CI. Hitchhiking under positive Darwinian selection. Genetics 2000;155(3):
1405-1413.

[114] Nielsen R., Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante CD. Genomic
scans for selective sweeps using SNP data. Genome Res 2005;15(11): 1566-1575.

[115] Jensen JD, Kim Y, DuMont VB, Aquadro CF, Bustamante CD. Distinguishing between
selective sweeps and demography using DNA polymorphism data. Genetics 2005;170(3):
1401-1410.

[116] Kim Y, Nielsen R. Linkage disequilibrium as a signature of selective sweeps. Genetics
2004;167(3): 1513-1524.

[117] Stephan W, Song YS, Langley CH. Hitchhiking effect on linkage disequilibrium between
linked neutral loci. Genetics 2006;172(4): 2647-2663.

[118] McVean G. The structure of linkage disequilibrium around a selective sweep.
Genetics 2007;175(3): 1395-1406.
927

928 [119] Jensen JD, Thornton KR, Bustamante CD, Aquadro CF. On the utility of linkage
929   disequilibrium as a statistic for identifying targets of positive selection in non-equilibrium
930   populations. Genetics 2007;176(4): 2371-2379.
931

[120] Crisci J, Poh YP, Mahajan S, Jensen JD. The impact of equilibrium assumptions on tests
of selection. Front Genet 2013; 4: 235.

[121] Garud N, Messer P, Petrov D. Detection of hard and soft selective sweeps from
Drosophila melanogaster population genomic data. PLoS Genet. 2021;17(2): e1009373.

[122] Teshima K, Coop G, Przeworski M. How reliable are empirical genome scans for
selective sweeps? Genome Res 2006;16(6): 702-712.
938

939 [123] Lynch M, Sung W, Morris K, Coffey N, Landry CR, Dopman EB, et al. A genome-wide
940   view of the spectrum of spontaneous mutations in yeast. Proc Natl Acad Sci USA
941   2008;105(27): 9272-9277.
942

943 [124] Bank C, Hietpas RT, Wong A, Bolon DNA, Jensen JD. A Bayesian MCMC approach to
944   assess the complete distribution of fitness effects of new mutations: uncovering the
945   potential for adaptive walks in challenging environments. Genetics 2014;196(3): 841-
946   852.
947

948 [125] Foll M, Poh YP, Renzette N, Ferrer-Admetlla A, Shim H, Malaspinas AS, et al. Influenza
949   virus drug resistance: a time-sampled population genetics perspective. PLoS
950   Genet 2014;10(2): e1004185.
951

952 [126] Böndel KB, Kraemer SA, Samuels TS, McClean D, Lachapelle J, Ness RW, et al.
953 Inferring the distribution of fitness effects of spontaneous mutations in *Chlamydomonas*
954 *reinhardtii*. PLoS Biol 2019;17(6): e3000192.
955
956 [127] Gelman A, Shalizi CR. Philosophy and the practice of Bayesian statistics. Br J Math Stat
957 Psychol 2013;66(1): 8-38.

977
978
Figure 1. Incorrect models may often readily be fit to a given dataset
Here we present three scenarios varying from simple to more complex: the first row presents a
constant-sized population experiencing background selection (denoted by 'Eqm + BGS'), the
second row is the same scenario with the addition of recurrent selective sweeps (denoted by
'Eqm +BGS + Pos'), and the final row adds a population bottleneck (denoted by 'Bottleneck +
BGS + Pos'). For each scenario, the resulting site frequency spectra (SFS, truncated to $n = 20$)
and linkage disequilibrium (r^2) distributions are given, together with mean pairwise (π) and
haplotype diversity. To these simulated data we fit two incorrect models; one assuming all sites
are neutral but including a change in population size, and a second model in which there are
recurrent selective sweeps, no change in population size, and all mutations are assumed to be
neutral or beneficial (with a population-scaled beneficial selection coefficient (γ) and the fraction
of beneficial substitutions (λ) being estimated from the data). For each inference panel, the red
cross gives the true value, the distribution presents the joint-posterior obtained from the ABC
analysis, and the summary statistics given above the posteriors represent the mean values, and
the range from the 95% CIs, obtained from posterior checks. In all cases, exonic sites (*i.e.*,
directly selected sites) were masked, and the summary statistic calculations as well as inference
is based only on neutral regions (see Methodology). As shown, demographic and selection
models can be fit to all datasets, often resulting in strong mis-inference when the assumptions
underlying the estimation procedure are violated.

998
999
1000
1001

1002

1003
1004
1005
Figure 2. Ascertainment errors may amplify mis-inference, if not corrected
As in Figure 1, the scenarios are given in the first column, here population growth with
background selection and recurrent selective sweeps ('Growth + BGS + Pos'), as well as the same
scenario in which the imperfections of the variant-calling processes are taken into account – in
this case, one-third of singletons are not called ('Growth + BGS + Pos + Ascertainment'). The
middle columns present the resulting SFS and LD distributions, and the final columns provide
the joint posterior distributions when the data are fit to two incorrect models: a demographic
model that assumes strict neutrality, and a recurrent selective sweep model that assumes a
constant population size. All exonic (*i.e.*, directly selected) sites were masked prior to analysis.
Red crosses indicate the true values. As shown, unaccounted for ascertainment errors may
contribute to mis-inference.

1030
1031
1032
Figure 3. The impact of potential model violations can be quantified
As in Figures 1 and 2, the scenarios are given in the first column, here equilibrium population
size together with a moderate degree of progeny skew ('Eqm + $\psi = 0.05$') as well as with a high
degree of progeny skew ('Eqm + $\psi = 0.1$') (see Methodology); the middle columns present the
resulting SFS and LD distributions, and the final columns provide the joint posterior distributions
when the data are fit to two incorrect models: a demographic model assuming neutrality, and a
recurrent selective sweep model assuming equilibrium population size. Red crosses indicate the
true values. As shown, this violation of Kingman coalescent assumptions can lead to drastic mis-
inference, but the biases resulting from such potential model violations can readily be described.
1042
1043
1044

Figure 4. The effects of not correcting for mutation and recombination rate heterogeneity
Three scenarios are here considered: equilibrium population size with background selection and recurrent selective sweeps ('Eqm +BGS + Pos'), declining population size together with background selection and recurrent selective sweeps ('Decline + BGS + Pos'), and growing population size together with background selection and recurrent selective sweeps ('Growth + BGS + Pos'). Inference is again made under an incorrect demographic model assuming neutrality, as well as an incorrect recurrent selective sweep model assuming equilibrium population size. However, within each category, inference is performed under two settings: mutation and recombination rates are constant and known, and mutation and recombination rates are variable across the region but assumed to be constant (see Methodology). Red crosses indicate the true values, and all exonic (*i.e.*, directly selected) sites were masked prior to analysis. As shown, neglecting mutation and recombination rate heterogeneity across the genomic region in question can have an important impact on inference, particularly with regards to selection models.

1068
Figure 5: **Diagram of important considerations in constructing a baseline model for**
genomic analysis
Considerations related to mutation rate are coded in red, recombination rate in blue, demographic
history in green, and the distribution of fitness effects in purple - as well as combinations thereof.
Beginning from the top with the source of data collected, the arrows suggest a path that is needed
to be considered. Dotted lines indicate a return to the starting point.

1075
1076 **Figure 6**: **Diagram of important considerations in detecting selective sweeps**
1077 The color scheme matches that in Figure 5, with 'selective sweeps' coded in orange.