WGS analysis of *Listeria monocytogenes* from rural, urban, and farm environments in Norway: Genetic diversity, persistence, and relation to clinical and food isolates

Annette Fagerlund[a]#, Lene Idland[b], Even Heir[a], Trond Møretrø[a], Marina Aspholm[b], Toril Lindbäck[b], and Solveig Langsrud[a]

[a] Nofima, Norwegian Institute of Food, Fisheries and Aquaculture Research, Ås, Norway

[b] Department of Paraclinical Sciences, Faculty of Veterinary Medicine, Norwegian University of Life Sciences, Ås, Norway

Running Head: WGS of environmental *L. monocytogenes* from Norway

Keywords: *Listeria monocytogenes*, whole genome sequencing, WGS, source tracking, persistence, molecular epidemiology, environmental pathogens, dairy farms.

Address correspondence to Annette Fagerlund, annette.fagerlund@nofima.no

Annette Fagerlund and Lene Idland contributed equally to this work. Author order was determined by mutual agreement.

Word count Abstract: 250 words. Word count Importance: 148 words.

Word count text (Introduction, Results, Discussion, and Materials & Methods): 5408 words

Abstract

Listeria monocytogenes is a ubiquitous environmental bacterium associated with a wide variety of natural and man-made environments, such as soil, vegetation, livestock, food processing environments, and urban areas. It is also among the deadliest foodborne pathogens, and knowledge about its presence and diversity in potential sources is crucial to effectively track and control it in the food chain. Isolation of *L. monocytogenes* from various rural and urban environments showed higher prevalence in agricultural and urban developments than in forest or mountain areas, and that detection was positively associated with rainfall. Whole genome sequencing (WGS) was performed for the collected isolates and for *L. monocytogenes* from Norwegian dairy farms and slugs, in total 218 isolates. The data was compared with available datasets from clinical and food associated sources in Norway collected within the last decade. Multiple examples of clusters of isolates with 0-8 wgMLST allelic differences were collected over time in the same location, demonstrating persistence of *L. monocytogenes* in natural, urban and farm environments. Furthermore, several clusters with 6-20 wgMLST allelic differences containing isolates collected across different locations, times and habitats were identified, including nine clusters harbouring clinical isolates. The most ubiquitous clones found in soil and other natural and animal ecosystems (CC91, CC11, and CC37) were distinct from clones predominating among both clinical (CC7, CC121, CC1) and food (CC9, CC121, CC7, CC8) isolates. The analyses indicated that ST91 was more prevalent in Norway than other countries and revealed a high proportion of the hypovirulent ST121 among Norwegian clinical cases.

Importance

Listeria monocytogenes is a deadly foodborne pathogen that is widespread in the environment. For effective management, both public health authorities and food producers need reliable tools for source tracking, surveillance, and risk assessment. For this, whole genome sequencing (WGS) is regarded as the present and future gold standard. In the current study, we use WGS to show that *L. monocytogenes* can persist for months and years in natural, urban and dairy farm environments. Notably, clusters of almost identical isolates, with genetic distances within the thresholds often suggested for defining an outbreak cluster, can be collected from apparently unrelated samples. The work highlights the need for a greater knowledge of the genetic relationships between clinical isolates and isolates of *L. monocytogenes* from a wide range of environments, including natural, urban, agricultural, livestock, food production, and food processing environments, in order to correctly interpret and use results from WGS analyses.

Introduction

Listeria monocytogenes is a bacterial pathogen responsible for the life-threatening disease listeriosis. The most common cause of listeriosis is considered to be ingestion of food contaminated by *L. monocytogenes* from unclean food production equipment (1, 2). *L. monocytogenes* is a ubiquitous environmental bacterium that has been associated with a wide variety of environments, such as rivers, soil, vegetation, wild and domesticated animals, food processing environments, and urban areas (3, 4). Consequently, a total absence of *L. monocytogenes* in non-heat-treated foods is difficult, perhaps impossible, to achieve. The literature is, however, not fully consistent about the main habitats of *L. monocytogenes* and the factors affecting its occurrence and spread to humans. It is therefore of importance to increase the understanding of the relationship between *L. monocytogenes* in natural and animal reservoirs, food processing environments, and human clinical disease.

The occurrence of *L. monocytogenes* in soil varies widely, from 0.7% to 45%, depending on the geographic area, season, and humidity (4-6). In comparative investigations, higher frequencies of *L. monocytogenes* have been found after rain, flooding and irrigation events (7, 8). Several studies have reported high incidence of *L. monocytogenes* in water from rivers and lakes, with frequencies from 10-62% of the samples depending on the area and detection method (9-13). A link has been found between the proximity to upstream dairy farms and cropped land and the presence of *L. monocytogenes* in river water (10, 12). An explanation for this could be high frequencies of *L. monocytogenes* in feces from farm animals, e.g., cattle, ducks, and sheep, leaking into surrounding soil and water (9, 14, 15). Dairy farms are, for example, known to hold a *L. monocytogenes* reservoir, and prevalences in environmental samples of 11-24% have been reported (6, 15-17). However, *L. monocytogenes* is not particularly linked to farm animals, and is frequently found in other animals and birds, such as game and urban birds, boars, garden slugs and rodents (9, 18-21). An association between dense populations of humans and occurrence of *L. monocytogenes* in the environment has been reported. A U.S. study showed that 4.4% of samples from urban or residential areas contained *L. monocytogenes*, while the pathogen was less frequently found in samples from forests and mountains (1.3%) (22).

L. monocytogenes comprises four separate deep-branching lineages, which from an evolutionary viewpoint could be considered separate species (23). These are further subdivided by multilocus sequence typing (MLST) into sequence types (STs) and clonal complexes (CCs or clones). The lineage I clones CC1, CC2, CC4, and CC6 are reported to be associated with human disease, while lineage II clones CC9 and CC121 are strongly associated with food and food processing environments (24-28).

While many studies have examined the molecular genotypes of *L. monocytogenes* isolates found in

food, food processing environments, and clinical disease, much less is known about the diversity

present in other environments. In the few published studies, the clonal diversity in environmental

samples from soil and water appears to be very high, sometimes dominated by CCs associated with

disease (CC1, CC4), although other dominating clones (e.g., CC37) have also been observed (5, 13,

29). Several clones are also found in wild animals, e.g., CC7 and CC37 found in moose, boars, slugs,

and game birds (18, 21, 30, 31). In environmental samples from dairy/cattle farms in Finland and

Latvia, the lineage II clones CC11 (ST451), CC14, CC18, CC20, CC37, and CC91 were most

predominant, while lineage I clones were rare (6, 32). Although there are some exceptions, e.g., CC1

being predominant in slugs collected in garden and farm environments in Norway (21), the majority

of the clones identified in natural and farm environments do not seem to belong to CCs dominant

among European food and clinical isolates.

Many studies have described persistence over time for *L. monocytogenes* clones in food processing

facilities (2, 33, 34) and in individual cattle herds or farm environments (15, 35, 36). Whether *L.*

monocytogenes can persist over long periods of time also in rural, urban, or agricultural

environments has rarely been investigated. Studies of genetic relationships between *L.*

monocytogenes isolates from natural and animal reservoirs and isolates from food and clinical

sources are scarce. High resolution molecular fingerprinting based on whole genome sequencing

(WGS) technology has revolutionized the ability to detect outbreaks and the presence of persistent

strains (37). Few studies have, however, carried out WGS analyses of *L. monocytogenes* isolates

collected from non-food associated locations over the span of months and years. The present study

aimed to use WGS to investigate the diversity and genetic relationships between *L. monocytogenes*

isolates from rural, agricultural, and urban environments in Norway, and to compare these with

available datasets containing genomes of *L. monocytogenes* from human clinical and food-associated

sources in Norway collected within the last decade.

Results

Prevalence of *L. monocytogenes* in rural and urban environments in Norway.

A total of 618 distinct environmental sites from rural and urban environments were sampled for *L.*

monocytogenes between April 2016 and April 2020. The overall sampling scheme was designed to

obtain an overview of the presence of *L. monocytogenes* in various habitats, and samples were

collected from several geographical regions in Norway (Figure 1A). To study potential persistence of

L. monocytogenes clones over time, some sites were sampled more than once. At the onset of the

study, we hypothesized that the presence of *L. monocytogenes* would be more strongly associated

with farm animals, agricultural activity, and urban areas than with natural forests and other wildlands

(22). During the first sampling occasion, 10% of sample sites were positive for *L. monocytogenes*

(Table 1). In addition, 13 samples of commercial bags of plant soil or compost were negative for *L.*

monocytogenes. In concordance with our hypothesis, the prevalence of *L. monocytogenes* was

significantly higher in urban areas and in areas associated with agriculture and livestock (agricultural

fields, grazelands and animal paths) than in forest/mountain areas and on footpaths (Fisher's exact

test *p*<0.02). Sampling locations classified as footpaths were generally from non-urban areas such as

woods or other areas used for hiking. While 14% of urban areas were positive for *L. monocytogenes*,

all samples from footpaths were negative for *L. monocytogenes*, and only 2% of the samples

collected in woodland or mountain areas were positive.

Detection of *L. monocytogenes* correlated with rain and sample humidity.

Previous studies have indicated that *L. monocytogenes* is more frequently isolated after recent

rainfall, irrigation, and flooding events (38, 39). In the present study, 271 out of 618 samples were

collected on days with rainfall and 347 samples on days with no rain within the previous 24 hours.

When collected on days with rain, 20% of samples were positive for *L. monocytogenes*, while on days

with no rain within the last 24 hours, only 3% of samples were positive. Thus, our data supports

previous studies suggesting that prevalence of *L. monocytogenes* is positively associated with rainfall

(Fisher's exact test $p=2\times10^{-12}$).

Upon sample collection, the humidity of the sampled material was categorized on a scale from 1

(completely dry) to 5 (liquid). Overall, the prevalence of *L. monocytogenes* in samples from the two

driest sample categories was 5.6% (4/70) and 5.7% (11/196), while it was 17% (27/164) and 14%

(12/86) in the more humid categories 3 and 4. The prevalence was significantly higher in the humid

samples (categories 3 and 4) than in the two driest sample categories (Fishers exact test, *p*<0.02).

The prevalence in liquid samples (category 5) was 10% (10/102). Among the samples collected in

urban environments, the sample humidity was not significantly associated (*p*>0.05) with the

prevalence of *L. monocytogenes*, with an overall prevalence of 10% in categories 1 and 2 (10/92), and

17% in categories 3 to 5 (14/85).

Persistent strains detected in rural and urban environments.

To examine whether environmental locations retained their status as *L. monocytogenes* positive or

negative over time, and whether the same clones were isolated repeatedly from the same location,

70 sites were subjected to one to three additional rounds of sampling the following years. In total,

115 *L. monocytogenes* isolates were collected in the current study (S1 Table). All isolates were

subjected to WGS, *in silico* MLST, and whole genome MLST (wgMLST) analysis. The distribution of

clones (CCs) among the identified isolates is presented in Figure 2.

Of the 44 sampling points positive for *L. monocytogenes* in the first round of sampling, 28 sites (64%)

were positive on at least one of the subsequent sampling occasions. Of the 26 initially negative sites,

five turned out positive during later sampling events (19%), and one of these was positive twice. In

total, 29 sampling sites were positive for *L. monocytogenes* more than once, and isolates belonging

to the same ST was collected repeatedly from seven sites (Table 2). In six cases, STs repeatedly

isolated from the same site were very closely related, with a maximum wgMLST allelic distance of 20.

When also adjacent or slightly more distant sampling sites (max 3 km) were included, a total of 14

clusters with genetic distances <20 were repeatedly collected from the same location over periods

ranging from 4 months to 3 years (S2 Table and S1 Text). When the commonly employed core

genome MLST (cgMLST) scheme described by Moura *et al*. (23) was employed, the isolates could not

be distinguished, except in one cluster with distances of 0 to 1 cgMLST alleles. Twelve clusters,

including two clusters each for CC91, CC11 (ST451) and CC37, represent clusters of highly similar

isolates with range 0 to 8 wgMLST allelic differences. Together, the results strongly indicate that *L.*

monocytogenes clones had persisted in the same environment or were repeatedly reintroduced

between sampling events in both rural and urban locations.

We also observed a case where a recent common contamination source was obscure: only 9 wgMLST

alleles (and no cgMLST alleles) separated a pair of CC6 isolates found 30 km and 3 years apart; one

isolate from a grazing pasture in Akershus county in 2020 and the other from soil by the root of a

tree in Oslo city centre in 2017.

Persistence and cross-contamination on Norwegian dairy farms.

In the next step, WGS was performed for a panel of 79 *L. monocytogenes* isolates collected from

Norwegian dairy farms (16). A total of 18 dairy herds from four different geographical areas within a

100 km radius from downtown Oslo (Figure 1B) had each been sampled four to six times between

August 2019 and July 2020. Out of the 556 analyzed samples, *L. monocytogenes* was detected in 12

milk filters (13% prevalence), 30 cattle feces samples (30%), 32 samples of cattle feed (silage or silage

mixture; 32%), and 5 teat swabs (5%). All bulk tank milk and teat milk samples were negative for *L.*

monocytogenes, and for one of the farms (farm 16), all 34 collected samples were negative. An

overview of the STs of the collected isolates (S1 Table) is presented in Table 3, and a phylogenetic

tree showing the genetic relationships between the individual isolates is shown in Figure 3.

Twelve clusters, each comprising two to four isolates, with pairwise genetic distances in the range 0

to 11 wgMLST alleles, were isolated from the same farm during repeated multiple visits, over periods

ranging from 2 to 10 months. These clusters involved 33 of the collected isolates and comprised ten

different CCs (S3 Table). These observations strongly support previous studies indicating that the

same *L. monocytogenes* clones can persist over time in individual cattle herds or farm environments

(15, 35, 36).

Out of 12 isolates from milk filters, four belonged to a persistent cluster, and one was closely related

to an isolate from a teat swab sample obtained on the same sampling occasion. When the same

clone was isolated from several sampling sites at the same farm, the pairwise genetic distances

separating milk filter isolates from fecal, feed or teat swab isolates ranged from 0 to 7 wgMLST allelic

differences (S3 Table). These links represent likely cross-contamination events where milk filters (and

consequently milk) have been contaminated with *L. monocytogenes* clones found in the farm

environment.

Detection of closely related isolates from different geographic areas.

In four cases, closely related isolates belonging to CC11 (ST451), CC226, and CC415 (ST394), were

collected from more than one dairy farm. The genetic differences between isolates from different

farms were somewhat greater than the diversity between isolates found on the same farm, with

between 9 and 20 pairwise wgMLST allelic differences. The number of cgMLST differences within

each cluster was 0 or 1 (S3 Table and S1 Text). These data indicate that farms located at different

geographical areas may host the same strain of *L. monocytogenes*.

Six clusters comprising *L. monocytogenes* from both dairy farms and isolates obtained from rural and

urban environments were detected. The genetic distances separating isolates from the two datasets

in these clusters ranged from 9 to 27 wgMLST allelic differences, and 0 or 1 cgMLST differences (S4

Table and S1 Text). The closest link was observed for a cluster of four CC37 isolates; two from grazing

land/pasture in the vicinity of Ås, and two from feed and teat swab samples obtained on two

different visits to farm 12, located about 50 km east of Ås. The two pairs of isolates were separated

by 9 to 14 wgMLST allelic differences and indistinguishable by cgMLST.

To further explore the occurrence of genetic links between Norwegian isolates from natural and

animal reservoirs, 24 of the 34 *L. monocytogenes* isolates collected from invading slugs (*Arion*

vulgaris) from garden and farm environments in Norway by Gismervik *et al*. in 2012 (21) were

subjected to WGS analysis (S1 Table). Interestingly, two pairs of slug isolates collected from different

geographic locations differed by only 2 (CC14) and 11 (CC1) wgMLST allelic differences. Furthermore,

five clusters with 10 to 21 wgMLST allelic differences comprised a slug isolate and one or more

isolates from either a rural/urban sampling site or from a dairy farm (S5 Table). The closest genetic

relationship concerned two CC1 isolates, in which a slug isolate from the west coast of Norway

(collected in 2012) showed only 10 wgMLST allelic differences compared to an isolate collected from

a street in a residential area in Oslo in 2017.

Thus, counting the previously mentioned pair of CC6 isolates collected in Akershus and Oslo, a total

of 17 close genetic links between isolates collected at relatively distant geographic areas in Norway

were detected in the set of 218 examined isolates. Presumably, not all clusters represent direct

epidemiological links, especially in cases where isolates were collected several years apart. The

observed genetic distances within the clusters; ≤21 wgMLST and ≤3 cgMLST allelic differences, are

within the thresholds often suggested as an appropriate guide for defining an outbreak cluster, which

is about 7-10 cgMLST differences (23, 40, 41) or about 20 single nucleotide polymorphisms (SNPs) in

SNP analyses (42, 43), which have a sensitivity comparable to wgMLST (34, 44).

Comparison with Norwegian clinical isolates.

The identification of close genetic links between isolates from different natural and animal-

associated sources without known connections led us to hypothesise that it would be possible to

identify clusters containing both environmental and clinical isolates with a similar level of genetic

relatedness. A dataset of Norwegian clinical isolates was identified, comprising 130 genomes from

2010-2015 (92% of all reported cases in these years) (45) and two genomes from 2018 (ST20 and

ST37), made publicly available by the European Centre for Disease Prevention and Control (ECDC)

and the Norwegian Institute of Public Health (NIPH), respectively. Sequencing data of sufficient

quality for wgMLST analysis was available for 111 of these isolates (S1 Table). An initial comparison

between the clinical isolates identified 15 pairs of isolates and nine larger clusters comprising 3 to 12

isolates showing genetic distances of ≤10 cgMLST allelic differences (S6 Table). Most clusters

contained isolates collected during a time span of several years and could potentially represent

listeriosis outbreaks or epidemiologically linked cases.

A wgMLST analysis showing the genetic relationships between isolates originating from rural and

urban environments, dairy farms, slugs, and clinical cases is shown in Figure 4. Nine clusters

contained clinical isolates differentiated from isolates sequenced in the current study by genetic

distances in the range 6 to 23 wgMLST allelic differences (0 to 7 cgMLST alleles) (S7 Table and S1

Text). The environmental *L. monocytogenes* closely related to clinical isolates comprised isolates

from soil samples from both urban and rural locations (belonging to CC4, CC7, CC11/ST451, CC220,

CC403, and CC415/ST394), three slug isolates obtained from garden and farm environments (CC7,

CC8, and CC9) and a group of CC11/ST451 isolates from dairy farms. The closest genetic link was

found between the single CC9 slug isolate (from 2012) and a clinical isolate from 2015, differentiated

by only 6 wgMLST alleles (and 0 cgMLST alleles). The analysis shows that *L. monocytogenes* isolates

that are genetically very closely related to clinical isolates can be detected in various natural and

agricultural environments, even when isolates are collected across timespans ranging several years.

Comparison of prevalence and diversity of MLST clones from different sources.

Most isolates from natural and agricultural environments belonged to *L. monocytogenes* lineage II,

comprising 89%, 94%, and 68% of isolates from rural/urban environments, dairy farms, and slugs,

respectively (Figure 5A). The remaining isolates belonged to *L. monocytogenes* lineage I, as lineage III

or IV isolates were not detected in the current study. The predominant clones among the rural/urban

isolates were CC91 (15%), CC19/ST398 (13%), CC37 (10%), and CC11/ST451 (9%). No specific niches

were found for these clones as isolates were spread geographically (3-5 counties), found in 3-5

different habitats/areas and in a range of humidity and weather conditions. CC91 appeared most

ubiquitous, as it was isolated from five different counties, from different sample types (soil, sand,

vegetation, and feces), from five different areas (agricultural fields, urban area, beach, grazeland and

forest), during all seasons and from all categories of humidity. CC11/ST451, CC91, and CC37 were

also the most frequently isolated clonal groups at the dairy farms (18%, 15%, and 11%, respectively);

each detected on seven different farms. Among the slug isolates, the most common clones were CC1

(15) and CC91 (12%) (21). A survey of previous studies indicated that CC1, CC7 and CC37 were the

clones most commonly detected in various natural and farm environments (S8 Table).

Among the examined Norwegian clinical isolates, CC7 was the most prevalent clonal group,

accounting for 23% (n=30) of the reported listeriosis cases, followed by CC121 (13%), CC8 (8%), and

CC1 (6%) (Figure 5B). In contrast to that observed in many other countries (27, 46), lineage I isolates

comprised a minority of the clinical isolates in this dataset (20%). The high prevalence of CC121

among the clinical isolates was unexpected, as this clone is commonly regarded as hypovirulent due

to the frequent occurrence of premature stop codons (PMSC) in the gene encoding the virulence

factor internalin A (*inlA*) (27), a characteristic also shared by the Norwegian CC121 clinical isolates.

Interestingly, the single *L. monocytogenes* CC121 isolated in the current study; MF7617 from soil at

the edge of a university campus pond in Ås, had an intact and presumably functional copy of *inlA*.

This isolate was only distantly related to the clinical CC121 isolates, separated by 195 wgMLST alleles

from the nearest clinical isolate. In contrast to CC121, the other three most commonly detected CCs

among the Norwegian clinical isolates – CC1, CC7 and CC8 – were relatively common also among the

isolates from natural and agricultural environments, with each CC having an average prevalence of

between 6% and 7.5% in the rural/urban, dairy farms, and slug isolate datasets (Figure 5A).

Since listeriosis is primarily acquired from food, the frequency distribution of CCs for *L.*

monocytogenes from Norwegian food processing industry (Figure 5C) was estimated from previous

work encompassing 680 isolates from five meat and four salmon processing plants, collected during 2011-2015 (47). The prevalence of lineage I isolates was <1% among the food processing industry isolates; represented by only two CC2 isolates from the meat industry. In meat processing environments, CC9 was by far the most prevalent clonal group, representing 70% of isolates. It must, however, be noted that most of the collected isolates were from two intensively sampled processing plants (34). One slug isolate and three clinical isolates (2011, 2012, 2015) but none from dairy farms or samples from rural and urban environments belonged to CC9. In salmon processing environments, CC14 was most prevalent (25%), followed by CC121 (22%), CC7 (ST7, ST732 and ST995; 18%), and CC8 (ST8 and ST551; 14%). CC14/ST14 was only represented by two clinical and two slug isolates, and not detected among isolates from rural/urban environments or dairy farms. The latter three were among the four most prevalent CCs among the Norwegian clinical isolates.

To examine the diversity of the most commonly detected *L. monocytogenes* clonal groups from Norwegian natural environments in an international context, a representative subset of reference genomes belonging to ST37, ST91, and ST451 were selected for comparative analysis using cgMLST (Figure 6). Of the >6000 examined publicly available genomes, 243 belonged to one of the three relevant STs. For ST91, a limited number of international reference sequences were available, and nearly 60% of the analysed isolates were Norwegian. This suggest that this ST could be more prevalent in Norway relative to in many other countries. For ST37, only limited clustering of Norwegian isolates relative to the international isolates was observed. In contrast, for ST91 and ST451, the Norwegian isolates appeared to cluster together with isolates from other countries, indicating that they represent internationally dispersed clones.

Discussion

It has long been acknowledged that *L. monocytogenes* clones predominating among human clinical isolates differ from those that dominate in food (23, 24, 26, 46, 48, 49), and that persistent clones of *L. monocytogenes* may establish in food processing environments (2, 33, 34). Here, we show that the most ubiquitous clones found in soil and other natural and animal ecosystems (CC91, CC11, and CC37) are distinct from clones predominating among both clinical and food isolates, and that *L. monocytogenes* may persist and spread also in urban and rural areas, grazeland, agricultural fields, and farm environments. The correspondence of major CCs was high for the three examined sets of environmental isolates (rural/urban, dairy farms, and slugs). CC37 appeared to be exceptionally widespread in natural environments, and was isolated from nine different counties and a wide variety of habitats. It was also found to persist for years both at a farm and on a bike path in the capital of Norway. The ubiquity of this clone is also reflected by its detection in a large proportion of

other studies investigating the identity of *L. monocytogenes* clones from natural and animal

reservoirs, including wildlife, forest areas and farms (5, 6, 13, 18, 21, 25, 29-31, 50).

The current study identified close genetic relationships between environmental isolates of *L.*

monocytogenes collected from geographically and temporally unrelated sources, despite a relatively

low number of analysed isolates. Although fixed clustering thresholds for defining outbreak clusters

are controversial (37, 51, 52), the genetic distances in the observed clusters were well within the

thresholds used to guide outbreak analyses (23, 40-43). In the majority of observed clusters, isolates

with no known likely association were indistinguishable using cgMLST analysis, which is the method

currently employed for surveillance of *L. monocytogenes* by many laboratories including the

Norwegian Institute of Public Health (53). This finding underscores the need for careful consideration

of additional evidence such as epidemiological data, traceback evidence, and phylogenetic tree

topology as part of WGS-based surveillance and outbreak investigations (54). Ideally, evaluation of

possible epidemiological links should consider the occurrence of closely related strains in the whole

food chain, including external contamination sources in urban and natural environments (52).

Currently, a lack of published genomic data on *L. monocytogenes* from various sources is a barrier for

effective management of this pathogen, both for public health authorities and for industrial actors.

During the last decade (2010-2020), an average of 24 yearly listeriosis cases have been reported in

Norway, and most of them (80%) were domestically acquired (http://www.msis.no/). The implicated

food is rarely identified. Only two outbreaks have been publicly reported during this period, both

associated with traditional fermented fish (rakfisk); one in 2013 (ST802, 4 cases in Norway) (55), and

one during the winter of 2018-2019 (ST20, 12 cases in Norway, one in Sweden (56)). A predominance

of lineage II was observed among the Norwegian clinical isolates – comprising 80% of isolates during

the years 2010-2015; an increase relative to the 56% observed during 1992-2005 (57). A distinct

feature among Norwegian clinical isolates was the large proportion of CC121 isolates lacking

functional internalin A. The only CC121 isolate collected from a natural environment did not have an

inlA PMSC, supporting the hypothesis that *inlA* mutations may constitute an adaptation to food

industry environments (58). The relatively high proportion of cases caused by clones of a

hypovirulent strain in Norway could potentially be linked to national consumption and storage

practices leading to sporadic ingestion of large numbers of the pathogen among high risk groups.

Worldwide, the hypervirulent clones CC1 and CC4 are significantly more prevalent among clinical

isolates than food isolates (27, 59, 60). Together, these two CCs constituted 8% of Norwegian clinical

isolates and 11% of the isolates from natural and farm environments. CC1 and CC4 also appear to be

prevalent in natural environments in other countries (13, 29). They were, however, not detected in a

study of *L. monocytogenes* in nine Norwegian food processing plants (47). Although at least 80% of

meat, cheese and fish consumed in Norway is produced domestically (61), imported processed foods

remain a potential source of infections. Notably, however, 45% of Norwegian households report that

they hunt, fish or collect bivalve molluscs, and about half of the population grow their own

vegetables, herbs or fruit, and collect berries in the wild (62). Furthermore, the current study

identified clusters comprising closely related isolates from both clinical sources and natural

environments, despite comparing temporally non-overlapping sets of isolates. Together, these

observations suggest that the relative contribution of industrially processed foods to listeriosis

infections could be lower in Norway than in other countries.

Materials and Methods

Sampling of *L. monocytogenes* from rural and urban environments.

Samples were taken to cover what was hypothesized as hot spots and cold spots for *L.*

monocytogenes in the outer environment. The sampling plan was designed to cover different

geographical regions of Norway and areas hypotesized to have respectively high (urban areas,

grazeland, animal paths and areas near food processing factories) and low (forests and mountain

areas, agricultural fields, beaches and sandbanks) occurrence of *L. monocytogenes*. Samples

classified as footpaths were generally from non-urban areas in woods or other areas used for hiking,

but separately categorized as we considered footpaths to be associated with human activities to a

greater extent than more pristine woodland or mountain areas. A detailed sampling scheme was

prepared and convenience sampling was performed by people living in or travelling to different areas

to cover Norway geographically and to get detailed results from specific areas (e.g., gardens) and

local information. The sampling was performed by trained microbiologists informed about the

objective of the study and which types of sites that should be sampled. When possible, several

different suspected hot and cold spots were sampled in the same geographical area, e.g., grazeland

and a forest nearby where the cattle did not have access. Sampling was performed year around

except for winter. For a selection of sampling sites, sampling was repeated once or more over a

period of three years.

The environmental samples (soil, sand, mud, decaying vegetation, surface water, animal dung, etc.)

were collected in sterile 50 mL Nunc tubes. All sampling locations were photographed, and GPS

coordinates, sample content, habitat/area, and weather conditions were recorded at the time of

sample collection. Specific information about the sample was also noted, such as which animals the

area was frequently exposed to (e.g., cattle, deer, sheep, doves) and local information (e.g., popular

area for hiking). The humidity of the collected samples was assessed on a scale from 1 to 5, ranging

from completely dry (1) to liquid (5). Samples were stored at 4°C for up to a week before processing,

and analysed according to ISO 11290-1 (63) with selective enrichment in half-Fraser and Fraser broth

(Oxoid) and final plating on RAPID'*L.mono* agar (Bio-Rad).

Whole genome sequencing.

For each *L. monocytogenes* isolate from rural/urban environments or from *Arion vulgaris* slugs (21), a

single colony was picked, inoculated in 5 mL brain heart infusion broth, and grown at 37°C overnight.

Culture samples (1 mL) were lysed using Lysing Matrix B and a FastPrep instrument (both MP

Biomedicals) and genomic DNA was isolated using the DNeasy Blood and Tissue Kit (Qiagen).

Libraries for genome sequencing were prepared using the Nextera XT DNA Sample Preparation Kit

(Illumina) and sequenced using 2×300 bp reads on a MiSeq instrument (Illumina). Raw reads were

filtered on q15 and trimmed of adaptors before *de novo* genome assembly was performed using

SPAdes v3.10.0 or v.3.13.0 (64) with the careful option and six k-mer sizes (21, 33, 55, 77, 99, and

127). Contigs with sizes of <500 bp and with coverage of <5 were filtered out.

Colonies from the *L. monocytogenes* isolates from dairy farms (16) were inoculated in 20 mL tryptone

soy broth and incubated at 37°C for 24 h before 1 mL was pelleted and DNA extracted using the

DNeasy Blood & Tissue kit (Qiagen). Libraries for genome sequencing were prepared using the

NEBNext Ultra DNA Library Prep Kit (New England Biolabs) with random fragmentation to 350 bp and

sequencing of 2×150 bp on a NovaSeq 6000 S4 flow cell (Illumina). *De novo* genome assembly was

performed using SPAdes v.3.14.1, incorporated in the software tool Shovill available at

https://github.com/tseemann/shovill. Shovill also performed adaptor trimming using Trimmomatic,

corrected assembly errors, and removed contigs with sizes <500 bp and coverage <2.

Phylogenetic analyses.

Classical MLST analysis followed the MLST scheme described by Ragon *et al*. (65) and the database

maintained at the Institute Pasteur's *L. monocytogenes* online MLST repository

(http://bigsdb.web.pasteur.fr/listeria/listeria.html). *In silico* MLST typing was performed for raw

sequencing data using the program available at https://bitbucket.org/genomicepidemiology/mlst

(66), and for genome assemblies using the program available at https://github.com/tseemann/mlst.

CCs are defined as groups of ST profiles sharing at least six of seven genes with at least one other

member of the group, except for CC14, which is divided into CC14 represented by ST14 and ST399 in

the current work, and CC91 represented by ST91, as isolates belonging to these two groups do not

cluster together in phylogenetic analyses of *L. monocytogenes* populations (27). The wgMLST analysis

was performed using a whole-genome schema containing 4797 coding loci from the *L.*

monocytogenes pan-genome and the assembly-based BLAST approach, implemented in BioNumerics

7.6 (http://www.applied-maths.com/news/listeria-monocytogenes-whole-genome-sequence-

typing). The cgMLST analysis was performed using the scheme described by Moura *et al*. (23), and

results were either extracted from the wgMLST data or obtained by sequence query at the BIGSdb-

Lm web server (https://bigsdb.pasteur.fr/listeria/listeria.html). Minimum spanning trees were

constructed using BioNumerics based on the categorical differences in the allelic cgMLST or wgMLST

profiles for each isolate. Loci with no allele calls were not considered in the pairwise comparison

between two genomes.

Publicly available genomes.

Available genomes of clinical isolates from human patients in Norway were identified by searching

the NCBI Pathogen Detection database (https://www.ncbi.nlm.nih.gov/pathogens) on the

30.08.2021. Available raw sequencing data from NCBI BioProjects submitted by ECDC and NPIH;

PRJEB26061 (45) and PRJEB25848, were subjected to *de novo* genome assembly as described for

isolates from rural/urban environments. *In silico* MLST genotyping was successful for all genomes

except one of the genomes published by the ECDC, and wgMLST analysis was successful for all except

21 of the ECDC genomes.

Reference genomes included in the cgMLST analysis of ST37, ST91 and ST451 genomes were

identified from the following selected sources on the 27.08.2021: (i) cgMLST profiles from the

BIGSdb-*Lm* database (http://bigsdb.pasteur.fr/listeria/listeria.html); 15 genomes belonging to

relevant STs were identified. (ii) Raw WGS data from the ListAdapt project

(https://onehealthejp.eu/jrp-listadapt/) comprising 1552 genomes (BioProject PRJEB38828). *De novo*

genome assembly was performed for the 165 genomes of relevant STs. (iii) Genome assemblies from

NCBI GenBank; among the 3926 *L. monocytogenes* genomes, 63 genomes belonged to the relvant

STs.

Data availability.

Data from this whole-genome shotgun project has been deposited at the NCBI GenBank database

under BioProjects PRJNA689486, PRJNA744724, and PRJNA689487. For GenBank and Sequence Read

Archive (SRA) accession numbers, see S1 Table. The assemblies were annotated using the NCBI

Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP) server

(http://www.ncbi.nlm.nih.gov/genome/annotation_prok/).

Acknowledgments

The authors wish to thank Merete Rusås Jensen, Anette Wold Åsli, Janina Berg, Tove Maugesten at

Nofima for excellent technical assistance. We also sincerely thank collaborators in the food and

salmon processing industry, colleagues at Nofima (Runar Gjerp Solstad, Rasmus Karstad, Halvor

Nygaard, Kristin Skei Nerdal) and Arild Hugo Solstad for contributing to sample collection. We thank

Ann-Katrin Llarena (Faculty of Veterinary Medicine, Norwegian University of Life Sciences) for help

during genome assembly of *L. monocytogenes* dairy farm isolates. We thank Ida Skaar (Norwegian

Veterinary Institute) for providing the *L. monocytogenes* isolates from slugs (21). We thank the team

of curators of the Institut Pasteur MLST system (Paris, France) for importing novel alleles, profiles

and/or isolates at http://bigsdb.pasteur.fr/listeria/. This work was funded by the Norwegian

Agriculture and Food Industry Research Funds, grant numbers 262306 and 314743.

References

1. Lopez-Valladares G, Danielsson-Tham ML, Tham W. 2018. Implicated food products for listeriosis
and changes in serovars of *Listeria monocytogenes* affecting humans in recent decades.
Foodborne Pathog Dis 15:387-397.

2. Ferreira V, Wiedmann M, Teixeira P, Stasiewicz MJ. 2014. *Listeria monocytogenes* persistence in
food-associated environments: Epidemiology, strain characteristics, and implications for public
health. J Food Prot 77:150-170.

3. Liao J, Guo X, Weller DL, Pollak S, Buckley DH, Wiedmann M, Cordero OX. 2021. Nationwide
genomic atlas of soil-dwelling *Listeria* reveals effects of selection and population ecology on
pangenome evolution. Nat Microbiol 6:1021-1030.

4. Vivant AL, Garmyn D, Piveteau P. 2013. *Listeria monocytogenes*, a down-to-earth pathogen.
Front Cell Infect Microbiol 3:87.

5. Linke K, Rückerl I, Brugger K, Karpiskova R, Walland J, Muri-Klinger S, Tichy A, Wagner M, Stessl
B. 2014. Reservoirs of *Listeria* species in three environmental ecosystems. Appl Environ
Microbiol 80:5583-5592.

6. Terentjeva M, Šteingolde Z, Meistere I, Elferts D, Avsejenko J, Streikiša M, Gradovska S, Alksne L,
Kibilds J, Bērziņš A. 2021. Prevalence, genetic diversity and factors associated with distribution
of *Listeria monocytogenes* and other *Listeria* spp. in cattle farms in Latvia. Pathogens 10:851.

7. Weller D, Wiedmann M, Strawn LK. 2015. Spatial and temporal factors associated with an
increased prevalence of *Listeria monocytogenes* in spinach fields in New York State. Appl Environ
Microbiol 81:6059-6069.

8.   Ivanek R, Grohn YT, Wells MT, Lembo AJ, Sauders BD, Wiedmann M. 2009. Modeling of spatially
referenced environmental and meteorological factors influencing the probability of *Listeria*
species isolation from natural environments. Appl Environ Microbiol 75:5893-5909.

9.   Fenlon DR, Wilson J, Donachie W. 1996. The incidence and level of *Listeria monocytogenes*
contamination of food sources at primary production and initial processing. J Appl Bacteriol
81:641-650.

10.   Lyautey E, Lapen DR, Wilkes G, McCleary K, Pagotto F, Tyler K, Hartmann A, Piveteau P, Rieu A,
Robertson WJ, Medeiros DT, Edge TA, Gannon V, Topp E. 2007. Distribution and characteristics
of *Listeria monocytogenes* isolates from surface waters of the South Nation River watershed,
Ontario, Canada. Appl Environ Microbiol 73:5401-5410.

11.   Stea EC, Purdue LM, Jamieson RC, Yost CK, Hansen LT. 2015. Comparison of the prevalences and
diversities of *Listeria* species and *Listeria monocytogenes* in an urban and a rural agricultural
watershed. Appl Environ Microbiol 81:3812-3822.

12.   Weller D, Belias A, Green H, Roof S, Wiedmann M. 2020. Landscape, water quality, and weather
factors associated with an increased likelihood of foodborne pathogen contamination of New
York streams used to source water for produce production. Front Sustain Food Syst 3:124.

13.   Raschle S, Stephan R, Stevens MJA, Cernela N, Zurfluh K, Muchaamba F, Nuesch-Inderbinen M.
2021. Environmental dissemination of pathogenic *Listeria monocytogenes* in flowing surface
waters in Switzerland. Sci Rep 11:9066.

14.   Nightingale KK, Schukken YH, Nightingale CR, Fortes ED, Ho AJ, Her Z, Grohn YT, McDonough PL,
Wiedmann M. 2004. Ecology and transmission of *Listeria monocytogenes* infecting ruminants
and in the farm environment. Appl Environ Microbiol 70:4458-4467.

15.   Castro H, Jaakkonen A, Hakkinen M, Korkeala H, Lindström M. 2018. Occurrence, persistence,
and contamination routes of *Listeria monocytogenes* genotypes on three Finnish dairy cattle
farms: a longitudinal study. Appl Environ Microbiol 84:e02000-17.

16.   Idland L, Granquist EG, Aspholm M, Lindbäck T. 2021. The occurrence of *Campylobacter* spp.,
Listeria monocytogenes and Shiga toxin-producing *Escherichia coli* in Norwegian dairy cattle
farms; a comparison between free stall and tie stall housing systems. bioRxiv
https://doi.org/10.1101/2021.10.25.465738.

17.   Fox E, O'Mahony T, Clancy M, Dempsey R, O'Brien M, Jordan K. 2009. *Listeria monocytogenes* in
the Irish dairy farm environment. J Food Prot 72:1450-1456.

18. Sauvala M, Woivalin E, Kivisto R, Laukkanen-Ninios R, Laaksonen S, Stephan R, Fredriksson-
Ahomaa M. 2021. Hunted game birds - Carriers of foodborne pathogens. Food Microbiol
98:103768.

19. Wang Y, Lu L, Lan R, Salazar JK, Liu J, Xu J, Ye C. 2017. Isolation and characterization of *Listeria*
species from rodents in natural environments in China. Emerg Microbes Infect 6:e44.

20. Hellström S, Kiviniemi K, Autio T, Korkeala H. 2008. *Listeria monocytogenes* is common in wild
birds in Helsinki region and genotypes are frequently similar with those found along the food
chain. J Appl Microbiol 104:883-888.

21. Gismervik K, Aspholm M, Rørvik LM, Bruheim T, Andersen A, Skaar I. 2015. Invading slugs (*Arion*
vulgaris) can be vectors for *Listeria monocytogenes*. J Appl Microbiol 118:809-816.

22. Sauders BD, Overdevest J, Fortes E, Windham K, Schukken Y, Lembo A, Wiedmann M. 2012.
Diversity of *Listeria* species in urban and natural environments. Appl Environ Microbiol 78:4420-
4433.

23. Moura A, Criscuolo A, Pouseele H, Maury MM, Leclercq A, Tarr C, Björkman JT, Dallman T,
Reimer A, Enouf V, Larsonneur E, Carleton H, Bracq-Dieye H, Katz LS, Jones L, Touchon M,
Tourdjman M, Walker M, Stroika S, Cantinelli T, Chenal-Francisque V, Kucerova Z, Rocha EPC,
Nadon C, Grant K, Nielsen EM, Pot B, Gerner-Smidt P, Lecuit M, Brisse S. 2016. Whole genome-
based population biology and epidemiological surveillance of *Listeria monocytogenes*. Nat
Microbiol 2:16185.

24. Painset A, Björkman JT, Kiil K, Guillier L, Mariet JF, Félix B, Amar C, Rotariu O, Roussel S, Perez-
Reche F, Brisse S, Moura A, Lecuit M, Forbes K, Strachan N, Grant K, Møller-Nielsen E, Dallman
TJ. 2019. LiSEQ - whole-genome sequencing of a cross-sectional survey of *Listeria*
monocytogenes in ready-to-eat foods and human clinical cases in Europe. Microb Genom
5:000257.

25. Félix B, Feurer C, Maillet A, Guillier L, Boscher E, Kerouanton A, Denis M, Roussel S. 2018.
Population genetic structure of *Listeria monocytogenes* strains isolated from the pig and pork
production chain in France. Front Microbiol 9:684.

26. Maury MM, Bracq-Dieye H, Huang L, Vales G, Lavina M, Thouvenot P, Disson O, Leclercq A,
Brisse S, Lecuit M. 2019. Hypervirulent *Listeria monocytogenes* clones' adaptation to mammalian
gut accounts for their association with dairy products. Nat Commun 10:2488.

27. Maury MM, Tsai YH, Charlier C, Touchon M, Chenal-Francisque V, Leclercq A, Criscuolo A,
Gaultier C, Roussel S, Brisabois A, Disson O, Rocha EPC, Brisse S, Lecuit M. 2016. Uncovering
Listeria monocytogenes hypervirulence by harnessing its biodiversity. Nat Genet 48:308-313.

28. Cabal A, Pietzka A, Huhulescu S, Allerberger F, Ruppitsch W, Schmid D. 2019. Isolate-based
surveillance of *Listeria monocytogenes* by whole genome sequencing in Austria. Front Microbiol
10:2282.

29. Papić B, Pate M, Félix B, Kušar D. 2019. Genetic diversity of *Listeria monocytogenes* strains in
ruminant abortion and rhombencephalitis cases in comparison with the natural environment.
BMC Microbiol 19:299.

30. Fredriksson-Ahomaa M, London L, Skrzypczak T, Kantala T, Laamanen I, Biström M, Maunula L,
Gadd T. 2020. Foodborne zoonoses common in hunted wild boars. Ecohealth 17:512-522.

31. Sauvala M, Laaksonen S, Laukkanen-Ninios R, Jalava K, Stephan R, Fredriksson-Ahomaa M. 2019.
Microbial contamination of moose (*Alces alces*) and white-tailed deer (*Odocoileus virginianus*)
carcasses harvested by hunters. Food Microbiol 78:82-88.

32. Castro H, Douillard FP, Korkeala H, Lindström M. 2021. Mobile elements harboring heavy metal
and bacitracin resistance genes are common among *Listeria monocytogenes* strains persisting on
dairy farms. mSphere 6:e0038321.

33. Carpentier B, Cerf O. 2011. Review - Persistence of *Listeria monocytogenes* in food industry
equipment and premises. Int J Food Microbiol 145:1-8.

34. Fagerlund A, Langsrud S, Møretrø T. 2020. In-depth longitudinal study of *Listeria monocytogenes*
ST9 isolates from the meat processing industry: Resolving diversity and transmission patterns
using whole-genome sequencing. Appl Environ Microbiol 86:e00579-20.

35. Haley BJ, Sonnier J, Schukken YH, Karns JS, Van Kessel JA. 2015. Diversity of *Listeria*
monocytogenes within a U.S. dairy herd, 2004-2010. Foodborne Pathog Dis 12:844-850.

36. Latorre AA, Van Kessel JA, Karns JS, Zurakowski MJ, Pradhan AK, Zadoks RN, Boor KJ, Schukken
YH. 2009. Molecular ecology of *Listeria monocytogenes*: evidence for a reservoir in milking
equipment on a dairy farm. Appl Environ Microbiol 75:1315-1323.

37. Pightling AW, Pettengill JB, Luo Y, Baugher JD, Rand H, Strain E. 2018. Interpreting whole-
genome sequence analyses of foodborne bacteria for regulatory applications and outbreak
investigations. Front Microbiol 9:1482.

38. Weller D, Wiedmann M, Strawn LK. 2015. Spatial and temporal factors associated with an
increased prevalence of *Listeria monocytogenes* in spinach fields in New York State. Appl Environ
Microbiol 81:6059-6069.

39. Ivanek R, Grohn YT, Wells MT, Lembo AJ, Jr., Sauders BD, Wiedmann M. 2009. Modeling of
spatially referenced environmental and meteorological factors influencing the probability of
Listeria species isolation from natural environments. Appl Environ Microbiol 75:5893-5909.

40. Zamudio R, Haigh RD, Ralph JD, De Ste Croix M, Tasara T, Zurfluh K, Kwun MJ, Millard AD,
Bentley SD, Croucher NJ, Stephan R, Oggioni MR. 2020. Lineage-specific evolution and gene flow
in *Listeria monocytogenes* are independent of bacteriophages. Environ Microbiol 22:5058-5072.

41. Ruppitsch W, Pietzka A, Prior K, Bletz S, Fernandez HL, Allerberger F, Harmsen D, Mellmann A.
2015. Defining and evaluating a core genome multilocus sequence typing scheme for whole-
genome sequence-based typing of *Listeria monocytogenes*. J Clin Microbiol 53:2869-2876.

42. Wang Y, Pettengill JB, Pightling A, Timme R, Allard M, Strain E, Rand H. 2018. Genetic diversity of
Salmonella and *Listeria* isolates from food facilities. J Food Prot 81:2082-2089.

43. Allard MW, Strain E, Rand H, Melka D, Correll WA, Hintz L, Stevens E, Timme R, Lomonaco S,
Chen Y, Musser SM, Brown EW. 2019. Whole genome sequencing uses for foodborne
contamination and compliance: Discovery of an emerging contamination event in an ice cream
facility using whole genome sequencing. Infect Genet Evol 73:214-220.

44. Jagadeesan B, Baert L, Wiedmann M, Orsi RH. 2019. Comparative analysis of tools and
approaches for source tracking *Listeria monocytogenes* in a food facility using whole-genome
sequence data. Front Microbiol 10:947.

45. Van Walle I, Bjorkman JT, Cormican M, Dallman T, Mossong J, Moura A, Pietzka A, Ruppitsch W,
Takkinen J, European Listeria Wgs Typing G. 2018. Retrospective validation of whole genome
sequencing-enhanced surveillance of listeriosis in Europe, 2010 to 2015. Euro Surveill
23:1700798.

46. Chenal-Francisque V, Lopez J, Cantinelli T, Caro V, Tran C, Leclercq A, Lecuit M, Brisse S. 2011.
Worldwide distribution of major clones of *Listeria monocytogenes*. Emerg Infect Dis 17:1110-
1112.

47. Møretrø T, Schirmer BCT, Heir E, Fagerlund A, Hjemli P, Langsrud S. 2017. Tolerance to
quaternary ammonium compound disinfectants may enhance growth of *Listeria monocytogenes*
in the food industry. Int J Food Microbiol 241:215-224.

48. Henri C, Leekitcharoenphon P, Carleton HA, Radomski N, Kaas RS, Mariet JF, Felten A, Aarestrup
FM, Gerner Smidt P, Roussel S, Guillier L, Mistou MY, Hendriksen RS. 2017. An assessment of
different genomic approaches for inferring phylogeny of *Listeria monocytogenes*. Front
Microbiol 8:2351.

49. Gray MJ, Zadoks RN, Fortes ED, Dogan B, Cai S, Chen Y, Scott VN, Gombas DE, Boor KJ,
Wiedmann M. 2004. *Listeria monocytogenes* isolates from foods and humans form distinct but
overlapping populations. Appl Environ Microbiol 70:5833-5841.

50. Kim SW, Haendiges J, Keller EN, Myers R, Kim A, Lombard JE, Karns JS, Van Kessel JAS, Haley BJ.
2018. Genetic diversity and virulence profiles of *Listeria monocytogenes* recovered from bulk
tank milk, milk filters, and milking equipment from dairies in the United States (2002 to 2014).
PLoS One 13:e0197053.

51. Schürch AC, Arredondo-Alonso S, Willems RJL, Goering RV. 2018. Whole genome sequencing
options for bacterial strain typing and epidemiologic analysis based on single nucleotide
polymorphism versus gene-by-gene-based approaches. Clin Microbiol Infect 24:350-354.

52. Gerner-Smidt P, Besser J, Concepción-Acevedo J, Folster JP, Huffman J, Joseph LA, Kucerova Z,
Nichols MC, Schwensohn CA, Tolar B. 2019. Whole genome sequencing: Bridging one-health
surveillance of foodborne diseases. Front Public Health 7:172.

53. Løvdal T, Brandal LT, Sundaram AYM, Naseer U, Roth B, Lunestad BT. 2021. Small-scale
comparative genomic analysis of *Listeria monocytogenes* isolated from environments of salmon
processing plants and human cases in Norway. Hygiene 1:43–55.

54. Brown E, Dessai U, McGarry S, Gerner-Smidt P. 2019. Use of whole-genome sequencing for food
safety and public health in the United States. Foodborne Pathog Dis 16:441-450.

55. Nielsen EM, Björkman JT, Kiil K, Grant K, Dallman T, Painset A, Amar C, Roussel S, Guillier L, Félix
B, Rotariu O, Perez-Reche F, Forbes K, Strachan N. 2017. Closing gaps for performing a risk
assessment on *Listeria monocytogenes* in ready-to-eat (RTE) foods: activity 3, the comparison of
isolates from different compartments along the food chain, and from humans using whole
genome sequencing (WGS) analysis. EFSA supporting publication EN-1151:170.

56. Public Health Agency of Sweden (Folkhälsomyndigheten). 2020. Listerios 2019. Sammanfattning
och bedömning. https://www.folkhalsomyndigheten.se/folkhalsorapportering-statistik/statistik-
a-o/sjukdomsstatistik/listeriainfektion/kommentarer-och-specialstatistik/2019/. Accessed
07.09.2021.

57. Rosef O, Klæboe H, Paulauskas A, Ambrasiene D. 2012. Diversity of *Listeria monocytogenes*
isolated from humans, food, and environmental sources in Norway. Vet Med Zoot 59:71-79.

58. Harrand AS, Jagadeesan B, Baert L, Wiedmann M, Orsi RH. 2020. Evolution of *Listeria*
monocytogenes in a food processing plant involves limited single-nucleotide substitutions but
considerable diversification by gain and loss of prophages. Appl Environ Microbiol 86:e02493-19.

59. Lee S, Chen Y, Gorski L, Ward TJ, Osborne J, Kathariou S. 2018. *Listeria monocytogenes* source
distribution analysis indicates regional heterogeneity and ecological niche preference among
serotype 4b clones. mBio 9:e00396-18.

60. Kwong JC, Mercoulia K, Tomita T, Easton M, Li HY, Bulach DM, Stinear TP, Seemann T, Howden
BP. 2016. Prospective whole-genome sequencing enhances national surveillance of *Listeria*
monocytogenes. J Clin Microbiol 54:333-342.

61. Helsedirektoratet. 2021. Utviklingen i norsk kosthold 2020. Matforsyningsstatistikk [*online*].
https://www.helsedirektoratet.no/rapporter/utviklingen-i-norsk-kosthold. Accessed 21.10.2121.

62. Bugge AB. 2015. Moralities of food and meals – how to eat properly [*online*]. Published by:
Statens Institutt for Forbruksforskning (SIFO), SIFO-report no. 3-2015.
https://hdl.handle.net/20.500.12199/927.

63. International Organization of Standardization. 2017. Microbiology of the food chain - Horizontal
method for the detection and enumeration of *Listeria monocytogenes* and of *Listeria* spp. —
Part 1: Detection method (ISO 11290-1). https://www.iso.org/standard/60313.html.

64. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Son
P, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012.
SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J
Comput Biol 19:455-477.

65. Ragon M, Wirth T, Hollandt F, Lavenir R, Lecuit M, Le Monnier A, Brisse S. 2008. A new
perspective on *Listeria monocytogenes* evolution. PLoS Pathog 4:e1000146.

66. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Ponten
T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-
sequenced bacteria. J Clin Microbiol 50:1355-1361.

Tables

Table 1: Prevalence of L. monocytogenes in rural and urban environments

Habitat or sampling area	No. of collected isolates	Prevalence of *L. monocytogenes* (%)
Grazeland or animal path	85	21 %
Urban or residential area	177	14 %
Agricultural field	70	11 %
Near food processing plant	106	10 %
Beach or sandbank	24	4 %
Forest or mountain area	121	2 %
Footpaths	35	0 %

Table 2: STs identified at sampling points positive for L. monocytogenes on repeated occasions

Site no.	Sampling point description	2016, Oct	2017, Jun, Oct, Nov	2018, Jun	2018, Sept, Oct	2019, Sept	2020, Jan
Urban or residential area							
33	Brook in residential area	ST451				ST4	
48	Garden compost heap	ST451 [a]				ST451 [a]	ST425
49	Garden compost heap	ST451				ST425 [b]	ST425 [b]
66	Puddle next to road	ST20				ST4	
120	Flowerbed in town centre		ST91		negative	ST1813	ST398
121	In front of park bench by flowerbed		ST399		ST451	ST398 [c]	ST398 [c]
123	Grass lawn in town centre		ST398 [d]		ST398 [d]	negative	negative
129	Roadside close to brook		ST37			ST91	negative
251	Decaying leaves/vegetation on bike path		ST18			ST37 [e]	ST37 [e]
252	Soil near horse paddock		ST204			ST398	ST7
253	Along sidewalk curb		ST1			negative	ST425
259	Flowerbed with pigeons, city centre		ST204 [f]			ST204 [f]	ST398
262	At foot of tree, city centre		ST6			ST204	ST120
268	Puddle on gravel path, city park		ST6			ST451	negative
Grazeland or animal path							
53	Decaying vegetation by feeding station	ST4				ST399	ST37
54	Soil close to cattle feeding station	ST91				ST399	ST451
55	Mud close to cattle enclosure	ST37				ST399	ST91
56	Puddle of mud close to cattle enclosure	ST91				ST398	ST6
98	Sheep grazing pasture		ST398		ST451	ST2343	ST91
99	Sheep manure		ST398		negative	negative	ST91
100	Animal tracks by feeding station		ST398		negative	ST37	negative
101	Animal tracks by feeding station		ST398		negative	ST91	negative
130	Soil at edge of pond		ST91		negative	ST121	negative
133	Decaying vegetation at edge of pond		ST398		negative	ST20 [g]	negative
134	Decaying vegetation at edge of pond		ST29		negative	ST4	ST20 [g]
Near food processing plant							
279	Grass next to cold storage entrance		ST732	ST732	ST732		
287	Storm drain outside plant		ST1	negative	ST647		negative
363	Gravel from quay outside factory			negative	ST732		ST647
406	Gravel from quay outside factory				ST1		ST647

Allelic distance between isolates with same superscript letter: [a] 3 wgMLST differences, [b] 2 wgMLST differences, [c] 2 wgMLST differences, [d] 34 wgMLST differences and 0 cgMLST differences, [e] 7 wgMLST differences, [f] 3 wgMLST differences, [g] 0 wgMLST differences (sites 133 and 134 are located 5 m apart), [h] 0-4 wgMLST differences.

Table 3: L. monocytogenes STs identified on dairy farms

Visit	Sample	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	17	18
Aug/Sept 2019	MF			ns														
	Feces	ns				ns			ST91		ST11	ST4				ST7	ST21	
	Feed				ST226	ST37							ST37				ST6	
	TS															ns		
Nov/Des 2019	MF	ns		ns	ST451			ST2760	ns		ST451	ST451						
	Feces	ST18				ST16	ST394		ST451			ST204		ST8				
	Feed	ST18				ST226	ST226	ST451	ST37				ST91					
	TS															ST37		
Jan 2020	MF	ST18								ns			ST37					
	Feces	ST18		ST2761									ST451				ST18	
	Feed	ST425	ST394	ST2761		ST7				ST224		ST4	ST451		ST91	ST91	ST18	
	TS			ST20														
Feb/Mar 2020	MF	ns						ns			ns	ns	ST177			ns		
	Feces	ST20		ST2761			ST451				ST21	ST451	ST177		ST91			
	Feed	ST37	ST394				ST451				ST412		ST177		ST91			
	TS	ST20																
May 2020	MF	ns	ns	ns									ST91	ST91		ns		
	Feces	ns	ns	ns									ST91	ST37		ns		
	Feed	ns	ns	ns									ST511	ST8		ns		ST91
	TS	ns	ns	ns									ST37	ST91		ns		
Jun 2020	MF	ns		ns			ST91		ST451	ST451						ns		
	Feces	ns		ns		ST37					ST21	ST412	ST4	ST451	ST124	ns		
	Feed	ns		ns							ST451	ST394	ST177			ns		
	TS	ns	ns	ns				ns								ns		

MF; milk filter, TS; teat swab, *ns*; not sampled, empty cells; negative for *L. monocytogenes*

Figures

Figure 1: Maps showing the geographic location of sampling sites. A) The location of sampling sites

in rural and urban environments in Norway, with red triangles representing *L. monocytogenes*

positive samples and blue circles negative sampling points. The area outlined by the dashed square in

a) is the area shown in **b)**. The green shaded areas in **b)** show the geographical origins of the dairy

cattle farms sampled for *L. monocytogenes* in Idland *et al*. (16).

L. monocytogenes identified in initial round of sampling (n=64)
L. monocytogenes identified during repeated sampling of previously sampled sites (n=51)

Figure 2: Distribution of CCs among identified isolates from rural and urban environments. The data is reported as percentage of the grand total number of isolates (n=115). STs represented within each clonal complex (CC) are given in parentheses.

Figure 3: Phylogeny for the _L. monocytogenes_ isolates from dairy farms. A minimum spanning tree based on wgMLST analysis. The area of each circle is proportional to the number of isolates represented, and the number of allelic differences between isolates is indicated on the edges connecting two nodes. The CCs and STs are indicated next to each node (the CC number is the same as the ST number unless indicated). Isolates separated from the nearest other isolate by >1700 wgMLST alleles (D011L, D058L, D080L, D084L, D144L, and D190L) were excluded from the figure for clarity.

Figure 4: wgMLST phylogeny for *L. monocytogenes* isolates from Norway. Shown is a minimum

spanning tree based on wgMLST analysis. The area of each circle is proportional to the number of

isolates represented, and the number of allelic differences between isolates is indicated on the edges

connecting two nodes. Edges shown as dashed lines separate clusters belonging to different clonal

complexes, which are indicated next to each node. Isolates separated from the nearest other isolate

by >900 wgMLST alleles (MF7614, MF6841, D084L, D144L, D190L, ERR2522251, ERR2522285,

ERR2522267, ERR2522291, and ERR2522298) were excluded from the figure for clarity.

Figure 5: Prevalence and distribution of *L. monocytogenes* MLST clonal complexes (CCs) from different sample types in Norway. The data are reported as percentages of isolates within a given CC in each source category. **a)** Prevalence in rural and urban environments (isolated during 2016-2020; n=115 isolates), dairy farms (2019-2020; n=87), and slugs (2012; n=34). The "other" category comprises one isolate each for CC31, CC121, CC475, and CC671. **b)** Prevalence in publicly available genomes from human cases of listeriosis in Norway (2010-2015; n=129 and 2018; n=2). The lineage I "other" category comprises a CC3 and a CC59 isolate, and the lineage II "other" category includes one isolate each for CC11, CC101, and CC177. **c)** Prevalence within food processing facilities in Norway. The CCs were inferred for isolates from five meat and four salmon processing facilities (meat: 2012-2015; n=293, and salmon: 2011-2014; n=358). The data used to predict the CC for each isolate were multiple-locus variable-number tandem repeat analysis (MLVA) obtained for all isolates and MLST data obtained for representative isolates from each obtained MLVA profile (47). The "unknown" category represents isolates with MLVA profiles identified only once and not subjected to MLST.

Figure 6: cgMLST phylogeny for the most common STs identified in the current study. Minimum

spanning trees based on cgMLST allelic profiles for: **a)** ST91, **b)** ST451, and **c)** ST37, showing the

relationship between the Norwegian isolates from natural environments, Norwegian clinical isolates,

and reference genomes obtained from public databases. Reference genomes were obtained from the

BIGSdb-*Lm* database hosted at the Pasteur Institute, WGS data from the EU project ListAdapt (also

including genomes from Norwegian sources, labelled in red), and genome assemblies from NCBI

GenBank. The area of each circle is proportional to the number of isolates represented, and the

number of allelic differences between isolates is indicated on the edges connecting two nodes.

List of supplemental information

S1 Text (*Supplemental_S1_Text.docx*): Additional information regarding clusters of closely related *L. monocytogenes* isolates.

Supplemental Tables (*Supplemental_Tables.xlsx*):

S1 Table: Isolates included in the current study

S2 Table: Clusters of isolates from rural or urban environments

S3 Table: Clusters of dairy farms isolates

S4 Table: Clusters containing isolates from both rural/urban environments and dairy farms

S5 Table: Clusters containing isolates from slugs or both slugs and rural/urban/farm environments

S6 Table: Clusters of clinical isolates

S7 Table: Clusters containing both clinical isolates and isolates from rural/urban/farm environments or slugs

S8 Table: Survey of STs and CCs found in previous studies of various natural and farm environments