GATA4 regionalizes intestinal metabolism and barrier function to prevent immunopathology

Zachary M. Earley[1,2], Wioletta Lisicka[1,2], Joseph J. Sifakis[3], Raúl Aguirre-Gamboa[2], Anita

Kowalczyk[2], Jacob T. Barlow[4], Dustin G. Shaw[1,2], Valentina Discepolo[5], Ineke L. Tan[6], Saideep

Gona[7], Jordan D. Ernest[2], Polly Matzinger[8], Luis B. Barreiro[1,2,7], Andriy Morgun[9], Albert

Bendelac[1,10], Rustem F. Ismagilov[4,11], Natalia Shulzhenko[12], Samantha J. Riesenfeld[13], Bana

Jabri[1,2,10,*]

Affiliations:

[1] Committee on Immunology, University of Chicago, Chicago, IL, USA

[2] Department of Medicine, University of Chicago, Chicago, IL, USA

[3] Department of Chemistry, University of Chicago, Chicago, IL, USA

[4] Division of Biology and Biological Engineering, California Institute of Technology, CA, USA

[5] Department of Medical Translational Sciences and European Laboratory for the Investigation

of Food Induced Diseases, University of Federico II, Naples, IT

[6] Department of Gastroenterology and Hepatology, University of Groningen and University of

Medical Center Groningen, Groningen, NL

[7] Genetics, Genomics, and Systems Biology, University of Chicago, Chicago, IL, USA

[8] Ghost Lab. National Institute of Allergy and Infectious Diseases, National Institutes of Health,

Bethesda, MD, USA

[9] College of Pharmacy, Oregon State University, Corvallis, OR, USA

[10] Department of Pathology, University of Chicago, Chicago, IL, USA

[11] Division of Chemistry and Chemical Engineering, California Institute of Technology, CA, USA

[12] Department of Biomedical Sciences, Oregon State University, Corvallis, OR, USA

[13] Pritzker School of Molecular Engineering, University of Chicago, Chicago, IL, USA

* Correspondence: bjabri@bsd.uchicago.edu (B.J.)

27
28

Summary

Different regions of the gastrointestinal tract have distinct digestive and absorptive

functions, which may be locally disrupted by infection or autoimmune disease. Yet, the

mechanisms underlying intestinal regionalization and its dysregulation in disease are not well

understood. Here, we used mouse models, transcriptomics, and immune profiling to show that

regional epithelial expression of the transcription factor GATA4 prevented adherent bacterial

colonization and inflammation in the proximal small intestine by regulating retinol metabolism

and luminal IgA. Loss of epithelial GATA4 expression increased mortality in mice infected with

Citrobacter rodentium. In active celiac patients with villous atrophy, low GATA4 expression was

associated with metabolic alterations, mucosal *Actinobacillus,* and increased IL-17 immunity.

This study reveals broad impacts of GATA4-regulated intestinal regionalization and highlights an

elaborate interdependence of intestinal metabolism, immunity, and microbiota in homeostasis

and disease.

One-sentence summary: Epithelial GATA4 regulates intestinal regionalization of bacterial

colonization, metabolic pathways, and tissue immunity.

Introduction

Each region of the gastrointestinal tract plays a distinct physiological role, featuring specific metabolic functions, host-microbe interactions, and immune responses. The duodenum and jejunum of the proximal intestine are optimized to digest and absorb critical nutrients. Despite having a large surface area in close contact with luminal contents, these regions also exhibit relatively low microbial growth and adhesion (*1*). In contrast, the ileum of the distal small intestine has shorter villi, less optimized for nutrient uptake, and instead specializes in reabsorption of bile acids and vitamin B12 (*1*). The ileum harbors more adherent commensal microbes, such as *segmented filamentous bacteria* (SFB), which promote local T_H17 differentiation and inflammatory IL-17 responses and protect the host from enteropathogenic infections (*2-4*). Finally, the colon is characterized by a thick mucus layer that permits absorption of water, electrolytes, and numerous microbial metabolites, while preventing the microbiota from invading the intestinal epithelium and driving inflammatory immunity (*1, 5, 6*). Together, these patterns suggest that each region uniquely balances its critical digestive and barrier immune functions, yet the regulatory mechanisms controlling regionalization remain elusive.

One known regulator of intestinal regionalization is the transcription factor GATA4, which is selectively expressed in the intestinal epithelial cells (IECs) of the duodenum and jejunum (*7, 8*). Deleting GATA4 in IECs was shown to alter lipid absorption in the jejunum and drive major transcriptional changes, including down-regulation of jejunal-specific genes and up-regulation of ileal-specific genes (*8*). These results suggest that epithelial GATA4 has a profound

role controlling numerous cell types in the tissue, a regulatory circuit that has not yet been

dissected.

Using GATA4-deficient mice, we investigated how GATA4 controls barrier function and

bacterial colonization in the jejunum through its key metabolic functions. We also assessed the

impact of disrupted intestinal regionalization on host susceptibility to pathology, both in the

context of enteric bacterial infection and human celiac disease, a dietary-induced autoimmune-

like disorder of the proximal small intestine.

Results

GATA4 regulates regionalization of intestinal immunity in a microbiota-dependent manner

Analysis of cytokine production in T cells from the intestinal track of specific-pathogen-

free (SPF) and germ-free (GF) mice, revealed regional-specific immune regulation (Fig. 1A). For

instance, despite the microbial load being considerably higher in the colon than in the ileum,

inflammatory microbiota-dependent IFN-γ^+ and IL-17$^+$ T cell responses were significantly higher

in the ileum. In contrast, regulatory IL-10$^+$ T cell responses were found at similar levels in the

ileum and colon, and were dependent on the presence of the microbiota in both locations.

Furthermore, the microbiota was required for the presence of intraepithelial IFN-γ^+ CD4+ T cells

in the ileum but not the proximal small intestine.

To determine how IECs shape regional tissue immunity, we analyzed mice in which

GATA4 was selectively depleted in IECs (GATA4fl/fl villin-cre+ or GATA4$^{\Delta IEC}$) (8). First, we

confirmed that GATA4 expression was restricted to duodenal and jejunal IECs, and that, in its

absence, jejunal IECs acquired an ileal-like transcriptional program (S1A, B). GATA4 strongly

repressed ileal-specific genes (*Fabp6*, *Slc10a2*) involved in the enterohepatic circulation of bile

acids (Fig. S1C). In contrast, GATA4 induced expression of lipid metabolic genes in the jejunum

involved in retinol metabolism (*Adh1*), and fat digestion and absorption (*Cd36*, *Fabp1*, *Dgat2*,

Apoa4). GATA4 also played an important role in controlling jejunal epithelial uptake of vitamins

and folate (Slc46a1, Pdxk) (Fig. S1C). To systematically assess the impact of GATA4 on regional

tissue functions, we performed total tissue RNA sequencing in GATA4$^{\Delta IEC}$ and littermate control

wild-type (WT) mice (GATA4fl/fl villin-cre-). Our analysis identified what we term "region-

specific, GATA4-regulated" genes, which are defined by expression that distinguishes "ileum-

like" tissue, *i.e.,* WT ileum and GATA4$^{\Delta IEC}$ jejunum, from WT jejunum (Fig. 1B, C). Intersecting

region-specific, GATA4-regulated genes with public databases of immune genes (*9*) revealed

distinct immune signatures of WT jejunum and ileum, a regionalization of tissue immunity that

was lost in GATA4$^{\Delta IEC}$ mice (Fig. 1B). Indeed, among the 625 regional-specific, GATA4-regulated

immune genes, more than a third (238) were published potential targets of IFNγ or IL-17

regulation (Fig. 1C).

Consistent with GATA4-regulated regionalization of IFNγ and IL-17 immune pathways, in

the absence of epithelial GATA4, the frequency of intraepithelial IFNγ^{+} cells among CD8αβ T

cells in the jejunum increased to the levels observed in the ileum (Fig. 1D). Furthermore, GATA4

deficiency led to a heightened T$_{H}$17 response in the jejunum, with frequencies of IL-17^{+} cells

among CD4 T cells surpassing those in the ileum (Fig. 1E). The heightened IFNγ and IL-17

immune responses observed in the jejunum of GATA4$^{\Delta IEC}$ mice were generally microbiota

dependent (Fig. 1F, S1D-F). Intriguingly, though, some particular immune genes, such as *Csf2,*

which encodes for GM-CSF, the anti-viral response genes *Ifnar2* and *Mavs*, and the tissue

alarmins IL-33 and IL-15, were GATA4 regulated in a microbiota-independent manner (Figure

1F). Among epithelial, region-specific GATA4-regulated genes, the microbiota-independent

subset was enriched, compared to the microbiota-dependent subset, in direct targets of GATA4

(Fig. S1G), as indicated by GATA4 binding of promoter regions in published ChIP-seq data

(hypergeometric distribution test; p-value = 3.21E-80) (Fig. S1G) (*10, 11*). Taken together, these

results demonstrate that GATA4 in IECs regulated regional tissue immunity and blocked

development of microbiota-dependent inflammatory immune responses in the proximal

intestine.

GATA4 shapes regional immunity by controlling colonization of *segmented filamentous*

bacteria

To investigate which specific microbiota may trigger inflammatory immune responses in

the absence of GATA4 in the proximal small intestine, we performed 16S ribosomal RNA

sequencing of luminal- and mucosal-associated bacterial communities in the jejunum and ileum

of littermate control WT and GATA4$^{\Delta IEC}$ mice. This analysis revealed that in the jejunum, SFB

(*Candidatus arthromitus*) loads were much higher in GATA4$^{\Delta IEC}$ relative to WT mice (Fig. 2A,

S2A). Furthermore, GATA4 was required to prevent SFB from colonizing (Fig. 2A, S2A) and

adhering (Fig. 2B) to IECs in the jejunum. In contrast, and in accordance with the lack of GATA4

expression in the ileum, no changes in bacterial composition were observed in the ileum of

GATA4$^{\Delta IEC}$ versus WT mice (Fig. S1A).

Previous studies showed that SFB triggers the upregulation of various adaptive immune

pathways (*2, 12-14*). To determine whether jejunal colonization by SFB was responsible for the

significant increase in inflammatory T cell immunity observed in the jejunum of SPF GATA4$^{\Delta IEC}$

mice (Fig.1 C-E), we performed a series of microbial transfers. First, transplantation of

GATA4$^{\Delta IEC}$ and littermate control WT microbiota into GF WT and GATA4$^{\Delta IEC}$ hosts, respectively,

showed that the host genotype determined the immune outcome irrespective of the input

microbial community (Figure S2B, C). Furthermore, transfer of either altered Schaedler flora

(ASF), a defined bacterial community that lacks SFB, or jejunal microbiota from a WT donor

within our colony with low-to-undetectable levels of SFB, failed to induce inflammatory T cells

in GATA4$^{\Delta IEC}$ mice. These results suggest that SFB is the causative microbe for dysregulated

immunity in GATA4$^{\Delta IEC}$ mice (Figure S2B, C). Finally, transfer of fecal homogenates from Jackson

C57BL/6J mice with or without SFB supplemented demonstrated that the increases in both

IFNγ^{+} CD8αβ T cells and T$_H$17 cells observed in the jejunum of GATA4$^{\Delta IEC}$ mice (Fig. 1D-E), were

dependent on the presence of SFB (Fig. 2C, D).

Since SFB colonization is known to induce an antigen-specific T$_H$17-cell response against

the 3340 epitope of SFB (*15*), we asked if the increased T$_H$17 cell response observed in the

jejunum was a consequence of altered T cell priming. To address that question, we transferred

congenically marked CD45.1^{+} 7B8^{+} CD4 T cells, specific to the 3340 epitope of SFB, into CD45.2

WT and GATA4$^{\Delta IEC}$ hosts. In the absence of GATA4, 7B8^{+} CD4 T cells differentiated into RORγt^{+}

Foxp3^{-} CD4 T cells in both the jejunal and ileal draining mesenteric lymph nodes (MLN) (Fig. 2E),

indicating a change in regional T cell priming against SFB in the MLN. Furthermore, nine days

after transfer, 7B8^{+} T cells expanded (Fig. 2F) and downregulated expression of their TCR in the

jejunum of GATA4$^{\Delta IEC}$, versus WT, mice (Fig. 2G). Together, these results indicate that, through

GATA4, IECs regulate regional colonization of the commensal microbe SFB and SFB-specific T-

cell priming.

To establish whether colonization of the jejunum by SFB depended on its ability to

adhere to IECs, we monocolonized GF WT and GATA4$^{\Delta IEC}$ mice with rat SFB, which is 86%

genetically identical to mouse SFB but unable to adhere to mouse IECs and thereby induce a

$T_H 17$ response (*12, 16*). Rat SFB was unable to expand and induce inflammatory T cell responses

in the jejunum of GATA4$^{\Delta IEC}$ mice (Figure 2H-J). To further test the hypothesis that GATA4

prevents colonization of the jejunum by adherent commensals, we transplanted GATA4$^{\Delta IEC}$ mice

with ASF containing the mucosal-associated commensal microbe *Mucispirillum ASF 457*.

Consistent with the hypothesis, ASF colonization of GF GATA4$^{\Delta IEC}$ and WT mice led to an

expansion *Mucispirillum ASF 457* in the jejunum of GATA4$^{\Delta IEC}$ but not WT mice (Fig. S2D).

GATA4 regulates luminal IgA levels to control colonization by SFB

We next sought to understand how GATA4 in IECs restricts adherent bacterial

colonization in the proximal intestine. Expansion of SFB was reported in the ileum and feces of

IgA deficient mice (*17, 18*). Furthermore, a previous report revealed that B-cell deficient mice

display lipid metabolic defects in the jejunum, as well as a gene expression signature associated

with GATA4$^{\Delta IEC}$ mice (*19*). Together, these studies suggest the possibility that an IgA defect in

GATA4$^{\Delta IEC}$ mice controls SFB colonization. In accordance with this hypothesis, SFB

monocolonization of GF B-cell deficient Jh$^{-/-}$ mice and GF IgA$^{-/-}$ mice substantially increased SFB

loads in the jejunum compared to those observed in the ileum of littermate control mice

indicating that IgA regulates regionalization of SFB colonization (Fig. 3A, B, S3A, B).

Given that B cells and IgA prevented SFB from colonizing the jejunum, we asked

whether GATA4 may control SFB colonization by regulating the regional distribution of IgA$^+$

plasma cells in the small intestine. We observed that the jejunum contained approximately

three times as many IgA$^+$ B220$^-$ plasma cells as the ileum (Fig. 3C), and that the higher numbers

of IgA-producing plasma cells were associated with a greater capacity to produce IgA in tissue

explants (Fig. S3C). This regionalization of IgA response was GATA4-dependent, as evidenced by

significantly reduced numbers of IgA$^+$ B220$^-$ plasma cells (Fig. 3C) and IgA production in the

jejunum of GATA4$^{\Delta IEC}$ versus WT mice (Fig. S3C). The overall result was a substantial decrease in

luminal secretory IgA in both the jejunum and the ileum of GATA4$^{\Delta IEC}$ mice (Fig. 3D). Reduced

luminal IgA was also observed in the jejunum and ileum of GF GATA4$^{\Delta IEC}$ mice, indicating that

the reduction is microbiota independent (Fig. S3D). Moreover, free IgA in the jejunal luminal

content from GATA4$^{\Delta IEC}$ mice had less capacity than that of littermate-control WT mice to coat

the microbiota present in the feces of immune-deficient (RAG$^{-/-}$) mice (Fig. 3E and S3E). Taken

together, these results indicate that GATA4 blocks the regionalization of adherent microbes,

such as SFB, by increasing luminal IgA.

We hypothesized that epithelial GATA4 mediated IgA levels by controlling region-

specific metabolic processes. To identify potential candidates, we performed gene set

enrichment analysis in our epithelial RNA-seq data (Fig. S1B) using the KEGG database, which

revealed retinol metabolism as a top GATA4-dependent, region-specific pathway (Fig. S3F).

Genes in the retinol metabolic pathway were expressed higher in the jejunum of WT, but not

GATA4$^{\Delta IEC}$, mice, compared to WT ileum (Fig. 3F), supporting previous reports that the proximal

intestine facilitates greater vitamin A uptake and metabolism (*20, 21*). Published ChIP-seq data

(*10, 11*) shows direct binding of GATA4 to the promoters of 23/35 (66%) of the differentially

expressed epithelial genes in the retinol metabolism pathway (Fig. 3F), suggesting direct

transcriptional regulation of retinol metabolism in IECs. Concordantly, in the absence of GATA4,

jejunal epithelial cells exhibited impaired aldehyde dehydrogenase (ALDH) activity, indicating a

decreased capacity to produce retinoic acid (RA) (Fig. 3G).

RA was previously shown to regulate intestinal B cell responses, including upregulating

intestinal homing receptors on B cells and IgA class switching (*22*). In addition, epithelial

RARα/β reportedly regulates the number of IgA-producing B cells (*23, 24*). Therefore, we asked

whether retinol deficiency was sufficient to enable the SFB expansion in the proximal small

intestine that we observed in GATA4$^{\Delta IEC}$ mice. We fed GF WT mice a control or vitamin-A

deficient diet for 4 weeks post-weaning and subsequently colonized them with SFB. SFB

colonized the jejunum of mice fed a vitamin-A deficient diet but failed to colonize the mice fed

a control diet (Fig 3H). To determine whether the defect in luminal IgA in the small intestine of

GATA4$^{\Delta IEC}$ mice could be rescued with exogenous RA, we injected GATA4$^{\Delta IEC}$ mice

intraperitoneally with RA for two weeks. RA significantly restituted luminal IgA in the jejunum

of GATA4$^{\Delta IEC}$ mice (Fig. 3I). We also tested whether exogenous luminal IgA was sufficient to

prevent colonization of the jejunum by SFB in GATA4$^{\Delta IEC}$ mice. Polyclonal luminal sIgA, capable

of strongly coating microbes from RAG-/- feces, was isolated from the intestinal contents of WT

mice with protein L magnetic beads (Figure S3G). This luminal polyclonal IgA or phosphate-

buffered saline (PBS) was gavaged to GF WT or GATA4$^{\Delta IEC}$ mice prior to and after colonization

with SFB (Fig. S3H). This IgA was indeed sufficient to prevent SFB from colonizing the jejunum

and ileum of GATA4$^{\Delta IEC}$ mice (Fig. 3J, S3I). These data indicate that, by regulating regional

retinol metabolism in IECs, GATA4 regulates luminal IgA, which in turn restricts bacterial

adhesion to the proximal small intestine.

Loss of GATA4 control of colonization by an adherent pathogen leads to severe

immunopathology

Since GATA4 controlled intestinal colonization by adherent commensal microbes, we

tested whether it also regulated colonization by adherent pathogens. *Citrobacter rodentium* is

an adherent pathogen that preferentially colonizes the colon but not the small intestine of WT

mice (Fig. 4A). In the absence of GATA4, the niche for *C. rodentium* was altered such that the

pathogen colonized the small intestine at levels approaching those in the colon of WT mice (Fig.

4A). Furthermore, the colonization in the small intestine of GATA4$^{\Delta IEC}$ mice depended on *C.*

rodentium's ability to adhere to IECs. Mutant ΔEAE *C. rodentium,* which lacks the gene intimin

required for adherence to IECs, did not colonize the jejunum and ileum of GF GATA4$^{\Delta IEC}$ mice

beyond levels observed in GF WT mice infected with WT *C. rodentium* (Fig. 4B). These results

suggest that GATA4 controls regional colonization of the small intestine not only by commensal

bacteria, but also by adherent pathogenic microbes.

We tested whether the expansion of *C. rodentiu*m colonization to the small intestine of

GATA4$^{\Delta IEC}$ mice impacted host immune response or survival. In accordance with the change in

colonization pattern, *C. rodentiu*m infection induced a striking expansion of IL-17$^+$ IFNγ$^-$, IL-17$^-$

IFNγ$^+$, and IL-17$^+$ IFNγ$^+$ double-positive CD4 T cells in the colon and ileum of GATA4$^{\Delta IEC}$ (Fig. 4C).

This contrasted with WT littermate control mice, which only showed a moderate T$_H$17 and T$_H$1

expansion in the colon (Figure 4C). The immune response in GATA4$^{\Delta IEC}$ mice was accompanied

by the development ten days after infection of severe colitis and villous atrophy in the ileum

(Fig. 4D), symptoms not normally associated with the infection in WT. Once the intestinal

barrier was compromised in GATA4$^{\Delta IEC}$ mice, *C. rodentium* translocated to systemic sites,

including the mesenteric lymph nodes, liver, and spleen (Fig. 4E), highlighting the substantial

impacts on distal sites that may be triggered by region-specific intestinal defects. By day 12

post-infection, 87.5% of GATA4$^{\Delta IEC}$ mice had died (Fig. 4F), punctuating the critical role of

GATA4-dependent intestinal regionalization in controlling host disease susceptibility to an

enteric pathogen.

Loss of GATA4 expression in celiac disease is associated with increased IL-17 immunity and

changes in bacterial colonization

Celiac disease (CeD) is an auto-immune–like T$_H$1-mediated enteropathy of the

duodenum caused by dietary gluten in genetically susceptible individuals (*25*). A long-standing

conundrum has been the increase of gluten-dependent, but not gluten-specific, duodenal T$_H$17

responses in CeD patients (*26-28*). Previous observations noted alterations in IECs (*25, 29*), as

well as lipid (*30*) and vitamin deficiencies (*31*) during active CeD (ACeD). We therefore

hypothesized that ACeD patients have decreased expression of GATA4 in IECs, and that this

decrease may be associated with reported changes in microbiota (*32, 33*) and increased IL-17

immunity. To investigate this possibility, we compared the transcriptional profiles of duodenal

biopsies from ACeD patients, CeD patients on a gluten free diet (GFD), and control patients with

non-inflammatory intestinal symptoms that required upper endoscopies. Intriguingly, our

analysis revealed significantly lower *Gata4* expression in ACeD patients versus controls, which

was restored by GFD (Fig. 5A). Immunohistochemistry staining demonstrated an ACeD-

associated loss of GATA4 protein production specifically in apical epithelial cells, whereas

intestinal crypts cells retained GATA4 production (Fig. 5B). To gain insights into the impact of

GATA4 in ACeD, we identified genes that were differentially expressed between "GATA4-hi"

individuals, defined by *Gata4* expression above the 70th percentile of the entire cohort, and

"GATA4-lo" ACeD patients, defined by *Gata4* expression below the 30th percentile of ACeD

patients (Fig. S4A). GATA4-lo specific genes were enriched in mouse genes specific to ileum-like

but not jejunum tissue. Furthermore, GATA4-hi specific genes were significantly enriched in

mouse genes specific to jejunum but not ileum-like tissue (Fig. 5C, S4B). Thus, during ACeD,

intestinal regionalization may be lost as the duodenum decreases GATA4-dependent jejunum-

specific gene expression and increases ileum-specific gene expression.

Many of the genes driving these enrichments are direct targets of GATA4 and involved

in lipid metabolic processes, such as cholesterol absorption and retinol metabolism (Fig. 5D).

Compared to GATA4-hi patients, GATA4-lo ACeD patients demonstrated decreased enrichment

in retinol metabolism genes and increased enrichment in IL-17 signaling genes (Fig. 5E). In fact,

expression of these pathways was correlated (or anti-correlated, in the case of IL-17 signaling

genes) with *GATA4* expression across all ACeD patients (Fig S4C-F). Together, these data reveal

that GATA4 expression can be lost in ACeD, and that this loss may play a role in regional

defects, such as low retinol metabolism, low plasma cholesterol (*34*), and increased IL-17

immunity.

Studies in mice revealed that the presence of T_H17 cells in the ileum is dependent on the

microbiota (*2*) and that bacteria can drive distinct immune responses depending on the host

context (*35*). Therefore, we asked whether a particular microbe was associated with the IL-17

response in ACeD patients. To answer this, we leveraged a quantitative framework to detect

absolute abundances of individual bacterial taxa in duodenal biopsies in which host

transcriptional analysis is performed concomitantly (*36*). Digital PCR anchoring of 16S rRNA

amplicon sequencing revealed no change in overall bacterial load in ACeD patients versus

controls (Fig. S4G) and only a trend of stratification between ACeD and control patients in the

second and third principal components (Fig. S4I). We did observe an expansion of *Neisseria* in

ACeD patients, the only microbial change significantly associated with disease state (Fig. S4H, J),

confirming a previous report (*37*). The increased abundance of *Neisseria* was not, however,

associated with decreased *Gata4* expression, metabolic defects, or IL-17 signaling in ACeD

patients (Fig. 5F).

In contrast, we found that *Actinobacillus* was associated with lower *Gata4* expression,

lower retinol metabolism, and higher IL-17 signaling in ACeD, but not control, patients (Fig. 5F,

G, S4K). Of note, we did not observe a significant association with the prototypical gluten-

specific T_H1 IFNγ pathway in ACeD patients, suggesting that *Actinobacillus* may play a specific

role in IL-17 immunity (Fig. 5F). Intriguingly, *Actinobacillus* was also associated with other

metabolic defects in ACeD patients, such as in xenobiotic, bile acid, and heme metabolism,

whereas other microbes, such as *Streptococcus,* had inverse patterns and positive associations

with bile and heme metabolism. The discovery of mucosal-associated bacteria, such as

Actinobacillus, associated with the loss of GATA4, metabolic dysfunction, and IL-17 immunity in

ACeD patients, highlights the potential importance in the pathogenesis of celiac disease of

GATA4 regulation of host microbial interactions.

Discussion

In this report, we identify mechanisms controlling regionalization of the proximal and

distal small intestine and reveal the importance of this segregation in homeostasis and disease.

We found that in part by regulating retinol metabolism and local IgA response, the epithelial

transcription factor GATA4 plays a critical role in limiting bacterial adhesion in the jejunum,

which may allow the host to focus on digestion and absorption of nutrients. On the other hand,

the ileum, lacking GATA4 expression and producing less IgA permits adherent microbes, such

as SFB, to induce inflammatory IL-17[+] T cell responses required for controlling pathogenic

infections (*2*). The intriguing association between loss of GATA4 expression in IEC and microbial-

associated dysregulated IL-17 immune responses in active CeD, poses the intriguing question of

whether a decrease in GATA4 in IECs of the proximal intestine may play a role in other complex

immune disorders by altering host-microbial interactions and triggering dysregulated immune

responses to bacteria.

Figure legends

Figure 1. Intestinal epithelial cells control microbiota-dependent and independent

regionalization of tissue immunity through GATA4.

(A) Percentage (y axis) of cells stained for cytokines IFNγ, IL-17, or IL-10 (panel titles), as

measured by intracellular flow cytometry, among all CD8αβ (leftmost panel) or CD4 (other

panels) T cells isolated from intraepithelial lymphocytes (IELs, left set) or the lamina propria (LP,

right set) of each intestinal segment (x axis) of SPF (orange circle) or GF (purple square) mice

and then stimulated by phorbol myristate acetate (PMA) and ionomycin. ****, $P<0.0001$,

effect due to region; ^{oooo} $P<0.0001$, ^{ooo} $P<0.001$, ^{oo} $P<0.01$, effect due to microbiota; two-way

ANOVA of microbiota and region impact on cytokine levels N= 5-7 mice/group.

(**B**) Top two principal components (PCs) (x, y axes) of the expression of region-specific, GATA4-

regulated genes, based on RNA-seq of tissue samples (dots) from the jejunum (dark colors) and

ileum (light colors) of WT (blues) and GATA4$^{\Delta IEC}$ (reds) mice. N= 8 mice/group.

(**C**) Heatmap of the z-scored expression (color) in tissue samples of region-specific, GATA4-

regulated immune genes (rows) from the jejunum and ileum (columns) of WT (left set) and

GATA4$^{\Delta IEC}$ (right set) mice. Of 625 total genes, 145 are in only the IFNγ module, 54 in only the IL-

17 module, 39 in both modules, and 387 in neither (annotation column) (Methods). N= 8

mice/group.

(**D**) Representative (left) and summary (right) plots of the frequencies of IFN-γ$^+$ cells among

CD8αβ T cells in the IELs of the jejunum and ileum of WT and GATA4$^{\Delta IEC}$ mice. **** $P<0.0001$,

paired t-test. N= 6 mice/group.

(**E**) Representative (left) and summary (right) plots of the frequencies of IL-17a$^+$ cells among

CD4 T cells in the LP of the jejunum and ileum of WT and GATA4$^{\Delta IEC}$ mice. *** $P<0.001$, paired t-

test. N= 6 mice/group

(**F**) Heatmap of the z-scored expression (color) in tissue samples of 50 selected microbiota

dependent and independent (right annotation column), region-specific, GATA4-regulated

immune genes (rows) in the jejunum (columns) of SPF (left set) and GF (right set) WT and

GATA4$^{\Delta IEC}$ mice. Gene modules (left annotation column) as in C.

Figure 1 supplemental. GATA4 controls regionalization of tissue metabolism and immunity

(**A**) Expression of GATA4 (y axis), as measured by qPCR relative to GAPDH, in the tissue of all intestinal regions (x axis).

(**B**) Top two principal components (x, y axes) of the expression of region-specific, GATA4-regulated genes in samples (dots) of sorted Epcam$^+$ CD45$^-$ IECs isolated from the jejunum (dark colors) and ileum (light colors) of WT (blues) and GATA4$^{\Delta IEC}$ (reds) mice.

(**C**) Heatmap of z-scored expression (color) in IECs of the jejunum and ileum of WT and GATA4$^{\Delta IEC}$ mice (columns) of selected DE genes (rows) from the following metabolic pathways: metabolism of steroids (Reactome), retinol metabolism (KEGG), fat digestion and absorption (KEGG), and metabolism of vitamins and cofactors (Reactome).

(**D**) Heatmap of the z-scored expression (color) of all region-specific, GATA4-regulated genes (rows) in jejunum tissue (columns) of SPF (left set) and GF (right set) WT and GATA4$^{\Delta IEC}$ mice.

(**E** and **F**) Frequency (y axis) of IFNγ$^+$ cells among CD8αβ T cells (**E**) or of IL-17a$^+$ cells among CD4 T cells (**F**) in the jejunum (left panel) and ileum (right panel) of SPF and GF WT and GATA4$^{\Delta IEC}$ mice (bar color). ****$P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$, ANOVA with Tukey multiple comparison test N= 6 mice/group.

(**G**) Frequency of genes bound (red) or not bound (gray) by GATA4, as determined by GATA4 ChIP-seq, among microbiota dependent (top) or microbiota independent (bottom) genes, as in D.

Figure 2. GATA4 controls the colonization of *segmented filamentous bacteria* to shape tissue immunity.

(**A**) SFB load (y axis), as measured by qPCR, relative to the amount of host DNA in mucosal scrapings (dots) of jejunum (dark colors) and ileum (light colors) from WT (blues) and GATA4$^{\Delta IEC}$ (reds) mice. **** $P<0.0001$, *** $Pp<0.001$, ** $P<0.01$,Kruskal-Wallis with Dunn multiple comparison test. N= 18-19 mice/group.

(**B**) Representative FISH staining of jejunal (left) and ileal (right) tissue of WT (top) and GATA4$^{\Delta IEC}$ (bottom) mice using universal 16s rRNA probes (Alexa 546) and a probe specific for SFB 16s (Alexa 488) and counterstained with DAPI. The overlay of the 16s probes represents SFB.

(**C** and **D**) Frequency (y axis) of IFNγ^+ cells among CD8$\alpha\beta$ T cells (**C**) or IL-17a$^+$ cells among CD4 T cells (**D**) in jejunum samples (dots) from SPF, GF, Jackson (JAX) microbiota transfer, and Jackson microbiota + SFB transfer (x axis) into WT and GATA4$^{\Delta IEC}$ mice (color, as in A). **** $P<0.0001$, ** $P<0.01$, t-test. N= 4-6 mice/group.

(**E**) Frequency (y axis) of RORγt$^+$ FOXP3$^-$ cells among transferred CD45.1$^+$ CD4$^+$ Vβ14$^+$ 7B8$^+$ T cells in the MLN (dots) draining the jejunum and ileum of CD45.2 WT and GATA4$^{\Delta IEC}$ mice (color, as in A), three days after transfer. **** $P<0.0001$, *** $P<0.001$, ANOVA with Tukey multiple comparison test. N= 5 mice/group.

(**F**) Number (y axis) of transferred T cells (as in E) in the jejunum LP(dots) 9 days after transfer into WT and GATA4$^{\Delta IEC}$ mice (color, as in A). ** $P<0.01$, t-test. N= 4 mice/group.

(**G**) Representative plot showing frequency of cells with vβ14+ TCR among transferred CD4+ CD45.1+ 7B8+ T cells in in the jejunum of WT (top) or GATA4$^{\Delta IEC}$ (bottom) mice.

(**H**) Rat SFB loads (y axis), as measured by qPCR relative to host DNA, in the jejunum (dots) of

monocolonized WT and GATA4$^{\Delta IEC}$ mice (color, as in A).

(**I** and **J**) Frequencies (y axis) of IFNγ$^+$ cells among CD8αβ T cells (**I**) and of IL-17$^+$ cells among CD4

T cells (**J**) in jejunum (dots) of GATA4$^{\Delta IEC}$ mice monocolonized with rat or mouse SFB (x axis).

**** P<0.0001, t-test. N= 5 mice/group.

Figure 2 supplemental. GATA4 controls the colonization of regional mucosal associated

bacteria to shape tissue immunity.

(**A**) Relative frequencies (x axis, log$_2$ fold change) in GATA4$^{\Delta IEC}$ versus WT mice of different types

of microbes (dots), and their statistical significance (y axis, −log$_{10}$ of the FDR-adjusted P value),

based on 16S rRNA sequencing of the luminal contents and mucosal scrapings of the jejunum

(left set) and ileum (right set) of WT and GATA4$^{\Delta IEC}$ mice. SFB are classified as *Candidatus*

arthromitus.

(**B** and **C**) Frequencies (y axis) of IFNγ$^+$ cells among CD8αβ T cells (**B**) and of IL-17a$^+$ cells among

CD4 T cells (**C**) in the jejunum (dots) of WT (blue) and GATA4$^{\Delta IEC}$ (red) mice with different

microbiota (x axis), *i.e.,* SPF, GF, or ASF, WT microbiota (no SFB), WT microbiota (with SFB),

GATA4$^{\Delta IEC}$ microbiota, or SFB. **** P<0.0001, *** P<0.001, ** P<0.01, t-test. N= 3-6

mice/group.

(**D**) *Mucispirillum schaedleri* load (y axis), as measured by qPCR, relative to host DNA in the

jejunum after ASF transfer to WT (blue) and GATA4$^{\Delta IEC}$ (red) mice. * P<0.05, Mann-Whitney.

N=4-5 mice/group.

Figure 3. GATA4 regulates regionalization of retinol metabolism and IgA to limit SFB

colonization in the proximal intestine.

(**A**) FISH staining of SFB from jejunal (left) and ileal (right) tissue of monocolonized IgA deficient

(IgA-/-) (bottom) and littermate control (IgA+/-) (top) mice. SFB is stained red with Cy5-

conjugated SFB-specific 16S probes. The DAPI counterstain is blue.

(**B**) SFB load (y axis), as measured by qPCR, relative to host DNA in mucosal scrapings (dots)

from the jejunum (violet) and ileum (light pink) of control (IgA+/-) mice and the jejunum

(fuchsia) of IgA deficient (IgA-/-) mice (x axis) mice. ** $P<0.01$, * $P<0.05$, Kruskal-Wallis with

Dunn multiple comparison test. N= 7-8 mice/group.

(**C**) Number of CD45$^+$ lin$^-$ IgA$^+$ B220$^-$ plasma cells (y axis), as determined by flow cytometry, in

tissue (dots) taken from the jejunum (dark colors) and ileum (light colors) of WT (blues) and

GATA4$^{\Delta IEC}$ (reds) mice. *** $P<0.001$, ** $P<0.01$, ANOVA with Tukey multiple comparison test.

N= 5 mice/group.

(**D**) Concentration of IgA (y axis), as determined by enzyme-linked immunoassay (ELISA), in the

contents (dots) of the jejunum (left) or ileum (right) of WT and GATA4$^{\Delta IEC}$ mice (color, as in C).

**** $P<0.0001$, t-test.

(**E**) Frequency of IgA coated bacteria (gated SYTOBC+) after staining of RAG-/- feces with

supernatant from WT and GATA4$^{\Delta IEC}$ jejunal contents, followed with labelled anti-IgA APC

antibody . *** $P<0.001$, t-test. N= 4-5 mice/group.

(**F**) Heatmap of z-scored expression (color) of epithelial, region-specific, GATA4-regulated genes

(rows) overlapping the KEGG retinol metabolism pathway, determined from RNA-seq of

epithelial cell samples (columns) from the jejunum and ileum of WT (left) and GATA4$^{\Delta IEC}$ (right)

mice.

(**G**) Top, representative histogram of ALDH expression in jejunal epithelial cells of WT (blue) and

GATA4$^{\Delta IEC}$ (red) mice revealed by ALDEFLUOR staining. WT epithelial cells treated with ALDH

inhibitor are shown as negative control for background fluorescence. Bottom, summary plots

show the normalized geometric mean fluorescence intensity (gMFI, y axis) of ALDEFLUOR

staining in samples (dots) of epithelial cells from the jejunum and ileum of WT and GATA4$^{\Delta IEC}$

mice (color, as in C). **** *P*<0.0001, ANOVA with Tukey multiple comparison test. *N*= 6

mice/group.

(**H**) SFB loads (y axis), as measured by qPCR relative to host DNA, in jejunal mucosal scrapings of

GF WT mice fed a control or vitamin A deficient diet for 4 weeks and subsequently colonized

with SFB. * *P*<0.05 Mann Whitney test N= 5 mice/group.

(**I**) Total IgA (y axis) in the jejunal contents of WT, GATA4$^{\Delta IEC}$ vehicle-treated, and GATA4$^{\Delta IEC}$ RA-

treated mice (x axis, color) after 14 days. *** *P*<0.001, ** *P*<0.01, * *P*<0.05, ANOVA with Tukey

multiple comparison test. *N*= 4 mice/group.

(**J**) SFB loads (y axis), as measured by qPCR relative to host DNA, in jejunal mucosal scrapings in

PBS-treated WT or GATA4$^{\Delta IEC}$ mice, and IgA-supplemented GATA4$^{\Delta IEC}$ mice (x axis, color). ***

P<0.001, ANOVA with Tukey multiple comparison test. *N*= 5-7 mice/group.

Figure 3 Supplemental. GATA4 regulates regionalization of retinol metabolism and B cell

responses

(**A**) FISH staining of SFB in jejunal (left) and ileal (right) tissue of monocolonized B-cell deficient

(Jh-/-) and littermate control (Jh+/-) mice. Red indicates SFB stained with Cy5-conjugated SFB-

specific 16s probes. Blue is DAPI counterstain.

(**B**) SFB load (y axis), as measured by qPCR relative to host DNA, in mucosal scrapings (dots)

from the jejunum (light blue) of control (Jh+/-) mice and the jejunum (violet) and ileum (dark

purple) of B-cell deficient (Jh-/-) mice. *** $P<0.001$, ** $P<0.01$, Kruskal-Wallis with Dunn

multiple comparison test. N= 7-8 mice/group.

(**C**) Concentration of IgA (y axis)), as measured by ELISA, in culture supernatant (dots) of tissue

explants from the jejunum (dark colors) and ileum (light colors) of WT (blues) and GATA4$^{\Delta IEC}$

(reds) mice after 24 hours in culture. **** $P<0.0001$, ANOVA with Tukey multiple comparison

test. N= 6 mice/group.

(**D**)Concentration of IgA (y axis), as measured by ELISA, in the contents (dots) of the jejunum

(left) or ileum (right) of GF WT (blues) and GATA4$^{\Delta IEC}$ (reds) mice. ** $p<0.01$, t-test.

(**E**)Intensity (y axis), of IgA bacterial coating as described in Fig. 3E. MFI, mean fluorescence

intensity. * $P<0.05$, t-test. N= 4-5 mice/group.

(**F**) Pathways (y axis) in the KEGG database that are significantly enriched (enrichment score, x

axis) in epithelial, region-specific, GATA4-regulated genes.

(**G**) Schema (left) for isolation of sIgA from luminal contents of WT mice, and FACS plots (right)

showing the frequency of IgA-coated bacteria after staining feces of RAG-/- mice with isolated

sIgA.

(**H**) Experimental schema of IgA gavage and SFB colonization experiment in GF WT and

GATA4$^{\Delta IEC}$ mice.

(I)SFB loads (y axis), as measured by qPCR, in ileal mucosal scrapings (dots) in PBS-treated WT

or GATA4$^{\Delta IEC}$ mice, and in IgA-supplemented GATA4$^{\Delta IEC}$ mice (x axis, color). * $P<0.05$, ANOVA

with Tukey multiple comparison test. N= 5-7 mice/group.

Figure 4. Loss of intestinal regionalization impacts host susceptibility to enteropathogenic

infection by *Citrobacter rodentium*.

(**A**) *C. rodentium* load (y axis), measured by qPCR relative to host DNA, in distinct intestinal

segments (x axis) in WT (blue) and GATA4$^{\Delta IEC}$ (red) mice. **** $P<0.0001$, ** $P<0.01$, Mann-

Whitney test. N= 13 mice/group.

(**B**) Bacterial loads (y axis) of wild-type *C. rodentium* (purple, orange) and the ΔEAE mutant

(black), measured by qPCR relative to host DNA, in distinct intestinal segments (x axis) of GF WT

(purple) or GATA4$^{\Delta IEC}$ (orange, black) mice. * $P<0.05$, Kruskal-Wallis with Dunn multiple

comparison test, GF WT + WT *C. rodentium* vs. GF GATA4$^{\Delta IEC}$ + WT *C. rodentium*. N= 7-9

mice/group.

(**C**) Representative FACS plots of IFN-γ (x axis) vs. IL-17a (y axis) in PMA ionomycin stimulated

CD4+ T cells that were isolated from the LP of infected or uninfected mice (left) or mice 10 days

after infection by *C. rodentium* (right).

(**D**) Representative immunohistochemistry (IHC) staining of each intestinal region (panel title)

from WT (top) and GATA4$^{\Delta IEC}$ (bottom) mice 10 days after *C. rodentium* infection.

(**E**) CFUs of *C. rodentium* translocation (y axis) to MLN, liver, and spleen (x axis) 10 days after

infection of WT (blue) and GATA4$^{\Delta IEC}$ (red) mice . **** $P<0.0001$, *** $P<0.001$, * $P<0.05$, Mann-

Whitney test. N= 8-10 mice/group.

(**F**) Percent survival (y axis) of WT (blue) and GATA4$^{\Delta IEC}$ (red) mice 0–15 days (x axis) post *C.*

rodentium infection. ** *P*<0.01 Mantel-Cox test. *N*= 8-9 mice/ group.

Figure 5. Loss of GATA4 expression is associated with lipid metabolic dysfunction and

increased IL-17 signaling in celiac disease.

(**A**) Normalized and batch corrected levels of GATA4 expression in duodenal biopsies from

healthy controls (blue), active celiac disease patients (ACeD, in red), and gluten free diet celiac

patients (GFD, in yellow). *** *P*<0.001, Wilcoxon test.

(**B**) Representative IHC staining for GATA4 in healthy control, active celiac disease patients, and

gluten free diet celiac patients.

(**C**) Bar plot shows the percentages (y-axis) of human-mouse homologous genes specific to

GATA4-hi or GATA4-lo (x-axis) individuals, that are also either jejunum (purple) or ileum-like

(yellow) specific and GATA4-regulated, or neither (gray). *** *P* <10^{-51}, ** *P* <10-6, NS not

significant, Fisher's exact test.

(**D**) Bar plots show the pathways (y axis) and significance ($-\log_{10}$ of adjusted *P*, x axis) of the

most significantly enriched gene ontologies (biological processes) in the 250 genes that are

upregulated in both GATA-hi individuals and WT mouse jejunum.

(**E**) Single-sample gene set enrichment analysis (ssGSEA) scores (y axis) for the retinol

metabolism (left) and IL-17 downstream signaling (right) pathways in GATA4-hi and GATA4-lo (x

axis) individuals (dots) from control (blue), ACeD (red), and GFD (yellow) patient groups. **** *P*

<0.0001 Wilcoxon rank test.

(**F**) Left, heatmap displays the scaled effect size (color) of the absence or presence of relevant 5

bacteria (see Fig. S4I) (rows) on GATA4 expression and on the ssGSEA scores of metabolic and

immune pathways (columns), while accounting for sex, age and technical covariates.

Significance annotated in red, · P <0.1, * P <0.05, ** P <0.01 t-test. Right, bar plot showing the

numbers (x axis) of ACeD samples in which the bacteria (y axis) were detected (orange) or not

(gray).

(**G**) Box plots show GATA4 expression (log_2 CPM, y axis, left panel) and ssGSEA scores (y axis,

other panels) for the retinol metabolism (center) and IL17 downstream signaling (right)

pathways in ACeD patients (dots), grouped (x axis) by the absence (blue) or presence (orange)

of bacteria from the Actinobacillus genus. * P <0.05, Wilcoxon test.

Figure 4 supplemental. Loss of GATA4 is associated with lipid metabolic defects, mucosal

associated bacteria, and increased IL-17 signaling in celiac disease patients.

(**A**) *Gata4* expression (y axis) in GATA4-hi and GATA4-lo groups (x axis) in individuals (dots) from

control (blue), ACeD (red), and GFD (orange) patients.

(**B**) Euler diagram represents the overlaps of 4 sets in a common universe of 11,657 mouse-

human transcriptional homologs: GATA4-regulated jejunum-specific (purple) and ileum-like–

specific (yellow) genes, and GATA4-hi specific (right, gray) and GATA4-lo ACeD specific genes

(left, gray). Enrichment significance as annotated, Fisher's exact test.

(**C**) Retinol pathway ssGSEA scores (y axis) in individuals (dots) from control, ACeD, or GFD

patient groups (x axis, color, as in A). **** P < 0.0001, NS not significant, Wilcoxon rank test.

(**D**) Scatter plot of GATA4 normalized expression (y axis) and ssGSEA scores (x axis) for the

retinol metabolism pathway in individual samples (dots) from control (blue) and ACeD (red)

patient groups. Annotated with Pearson correlation coefficient and p-value.

(**E**) IL-17 downstream signaling pathway ssGSEA scores, analogous to C.

(**F**) Scatter plot of GATA4 normalized expression (y axis) and ssGSEA scores (x axis) for the IL-17

downstream signaling pathway, analogous to D.

(**G** and **H**) Box plots of absolute abundances (y axis) of mucosal 16S (**G**) and Neisseria (**H**) in

biopsies (dots) from control (blue) or ACeD (red) (x axis) patients.

(**I**) Principal component analysis (PCA) biplot shows PC2 (x axis) and PC3 (y axis) of each

individual (dot) from control (blue) or ACeD (red) patient groups; arrows indicate the loadings

of the 5 bacteria most associated with PC2 and PC3.

(**J**) Numbers (x axis) of control or ACeD patients (y axis) with detectable (orange) or

undetectable (gray) levels of the indicated bacteria.

(**K**) Heatmap shows the scaled effect size (color) of each of the bacteria from J (x axis) on GATA4

expression and ssGSEA scores of the indicated pathways in controls, compared to ACeD

patients. $\cdot$ P <0.1, * P <0.05, t-test.

Materials and Methods

Mice

7-12 week old mice were used for experiments, co-housed in specific pathogen-free conditions, and kept *Helicobacter hepaticus*, murine norovirus free at the University of Chicago. Some mice were also housed in gnotobiotic isolators and routinely checked for sterility by culture and 16S PCR or kept SFB monocolonized at the University of Chicago Gnotobiotic Research Animal Facility. GATA4fl/fl villin-cre SPF mice were previously generated in the CD1 background and obtained from the Matzinger laboratory (*19*).This line was rederived GF for this study and backcrossed for 10 generations to C57BL/6J background for T cell transfers. C57BL/6J, B6-Tg(Tcra, Tcrb)2Litt/J SFB TCRtg, B6.SJL-*Ptprc^a Pepc^b*/BoyJ, B.6129S7-Rag1^{tm1mom}/J were obtained from the Jackson Laboratory. CD-1 IGS mice were obtained from Charles River

Laboratories. B cell deficient mice deficient for IgH J segment locus (Jh) were obtained from Dr.

Bendelac at the University of Chicago and generated on a C57BL/6 background using Cas9 with

the protospacers GCTACTGGTACTTCGATGTC and GCCATTCTTACCTGAGGAGA. IgA deficient mice

where the Sα (IgA switch region) and C1α (first exon) were deleted were obtained from Dr.

Bendelac and generated on a C57BL/6 background using Cas9 with the protospacers

AAGCGGCCACAACGTGGAGG and TCAAGTGACCCAGTGATAAT. Jh and IgA deficient mice were

rederived GF at Taconic Biosciences. Littermate controls of GATA4, Jh, and IgA were used for all

experiments in this study. Mice were fed a standard chow diet, vitamin A control diet (Harlan

TD.91280), or vitamin A deficient diet (Harlan TD. 86143). Animal husbandry and experimental

procedures were performed in accordance with Public Health Service policy and approved by

the University of Chicago Institutional Animal Care and Use Committees.

Patients

A duodenal biopsy was obtained from 166 individuals undergoing upper gastrointestinal

endoscopy at the University of Chicago and at Mayo Clinic as previously reported (*38*). There

were 64 control patients, 56 untreated patients with active celiac disease, and 46 patients

treated with a gluten free diet. All control patients underwent endoscopies for issues unrelated

to celiac disease and had normal intestinal histology, no family history of celiac disease, and no

significant levels of anti-TG2 antibodies in the serum. Patients with active celiac disease

contained positive anti-TG2 antibodies and small intestinal enteropathies with increased IEL

infiltration, crypt hyperplasia, and villous atrophy according to the accepted diagnostic

guidelines (*39*). The subjects signed an informed consent as provided by the Institutional

Review Board of each institution (IRB-12623B for the University of Chicago, and IRB-1491-03 for

the Mayo Clinic). DNA and RNA were isolated from each biopsy as described previously (*38*)

using the AllPrep DNA/RNA mini kit (Qiagen).

Isolation of intestinal epithelial cells (IEC), intraepithelial lymphocytes (IEL), and lamina

propria (LP) cells

The segments of the intestine were excised as follows to isolate cells for flow cytometry:

duodenum was taken 12 cm from the stomach, jejunum 12 cm from the middle, and ileum 12

cm from the cecum. Any leftover segments were discarded. The entire colon was taken after

the cecum to the rectum. To isolate IEL and IECs, Peyer's patches were first removed from the

small intestine, and then the segments were opened longitudinally and washed briefly in

phosphate-buffered saline (PBS). Epithelial cells, including IELs and LP cells, were isolated as

previously described (*40*) using ethylenediaminetetraacetic acid (EDTA) containing calcium-free

media and collagenase VIII (Sigma-Aldrich, C2139), respectively. The IEL and LP compartments

were then subjected to a 40% percoll density gradient centrifugation step to remove dead cells

and debris as previously described (*38*). The IEL and LP cells were then counted on a

hemocytometer.

Cytokine stimulation

Up to $2x10^6$ cells were collected and resuspended in RPMI 1640 media with 10% fetal

bovine serum (FBS) and cultured in 48-well plates in the presence of 750 ng/ml of ionomycin,

50 ng/ml of Phorbol 12-myristate 13-acetate (Sigma-Aldrich), and golgi-stop (BD). The cells

were incubated for 2 hours at 37C with 5% CO_2. After stimulation, the reaction was quenched

with ice cold FACS buffer, and the cells were subsequently stained with antibodies for flow

cytometry.

Flow cytometry

The cells were first stained with FC block (CD16/32) to block nonspecific binding and

then were stained with dead dye to exclude dead cells (Aqua, ThermoFisher or Zombie NIR,

Biolegend) for 15 min at 4 °C, followed by staining with cell surface markers for 20 min at 4 °C.

For intracellular cytokine staining, the BD cytofix/cytoperm kit was used, and cells were

incubated with the antibodies for 40 min at 4 °C. For intracellular transcription factors, the

Foxp3 eBioscience kit was used according to the manufacturer's instructions. The antibodies

used are indicated in Table (1). For ALDH staining of IECs, the ALDEFLUOR kit was used

(StemCell Technologies), following the manufacturer's protocol. All cells were gated FSC, SSC,

singlets, and live cells. IECs were gated CD45- EpCAM+. 100,000 IECs from the jejunum or ileum

were sorted with Aria Fusion (BD Biosciences) into RLT buffer (Qiagen) with β-mercaptoethanol

for downstream sequencing analysis. IgA plasma cells were gated as described previously (*41*),

i.e. EpCAM-, CD45+/dim, lineage negative (Ter119, F4/80, CD3, Ly6G, NK1.1, CD19), IgA+, B220-

. CD8αβ IELs were gated TCRβ+, CD4-, CD8α+, CD8β+. CD4 LP T cells were gated TCRβ+, CD4+,

CD8α-. The following antibodies and clones were purchased from Biolegend: CD45 Pacific Blue

(30-F11), CD4 BV785 (GK1.5), CD4 BV605 (GK1.5), IL10 PE-Cy7 (JES5-16E3), CD45.1 Pacific Blue

(A20), Tbet PE (4B10), CD44 PE-Cy7 (IM7), CD62L PE (MEL-14), Epcam PerCP-Cy5.5 (G8.8), CD19

FITC (1D3/CD19), NK1.1 BV605 (PK136), CD11C BV605 (N418), TER119 BV605 (TER-119), F4/80

BV605 (BM8), CD3ε BV605 (145-2C11), Ly6G BV605 (1A8), B220 PE-Cy7 (RA3-6B2). The

following antibodies and clones were purchased from BD: CD8β BUV395 (H35-17.2), CD8α

PerCP-Cy5.5 (53-6.7), NK1.1 PE-CF594 (PK136), TCRβ BUV737 (H57-597), TCRβ BV711 (H57-

597), CD3ε BUV737 (145-2C11), IFN-g APC (XMG1.2), CD45.2 BUV395 (104), vβ14 TCR FITC (14-

2), RORγt BV786 (Q31-37). The following antibodies and clones were purchased from Thermo

Fisher: TCRgd FITC (eBioGL3), IL17a PE (ebio17B7), FOXP3 eFluor450 (FJK-16s), FOXP3 FITC (FJK-

16s), FOXP3 PE-Cy7 (FJK-16s), IgA PE (mA-6E1). The cells were run on the LSRFortessa X-20 Flow

Cytometer (BD Biosciences) or the Cytek Aurora and data were analyzed using FlowJo software

(Treestar).

DNA isolation

For mucosal scrapings for DNA isolation, 5 cm of tissue proximal to the middle of the

intestine was taken for the jejunum, and 5 cm from the ileocecal valve was taken for the ileum.

The entire colon was used for mucosal scrapings. The tissue was excised, opened longitudinally,

scraped with a glass slide, transferred to 2 ml screw cap tube containing 0.1 mm glass beads

(Bio-spec), and snap frozen on dry ice. For luminal content, 50-100 mg of content was taken

from as close to the middle of the jejunum as possible and from the last 5–7 cm of the ileum.

Homogenization was performed after adding 1 ml of inhibitex buffer (Qiagen) using the Bead

Ruptor Elite bead mill homogenizer (Omni, 19040E) on speed 6 for 3 min. DNA was then

extracted using the QIAmp Fast DNA stool mini kit (Qiagen) following the manufacturer's

protocol with the optional high temp (95 °C) lysis step. DNA concentration was determined

using the nanodrop UV spectrophotometer (ThermoFisher).

RNA isolation

For eventual RNA purification, 1 cm of tissue was excised from the beginning of the

duodenum, the middle of the jejunum, the end of the ileum, and the center of the colon and

preserved in RNAprotect (Qiagen) overnight at 4 °C and then transferred to −80 °C for long-

term storage. The tissue was transferred to 600 µl of RLT buffer containing β-mercaptoethanol

(Qiagen) and homogenized for 30 sec with a hand held rotor (Tissue-Tearor, BioSpec). RNA was

purified using RNAeasy plus mini kit (Qiagen) following the manufacturer's protocol with the

optional on column DNase digest (Qiagen).

qPCR

RNA was first reverse-transcribed to cDNA using GoScript Reverse Transcriptase kit

(Promega) following the manufacturers protocol. For qPCR, 10 ng of cDNA or 20 ng of DNA

from mucosal scrapings and content was used. TB green Advantage qPCR Premix (Takara) was

used, and the target gene was quantified and normalized to the housekeeping gene as

described previously (*38*) using $1000*2^{-(Ct\ target-\ Ct\ housekeeping)}$ formula. For host gene expression,

the target gene was normalized to GAPDH. For bacterial load, the target gene was normalized

to either host DNA as described previously (*42*) with primers specific for host argininosuccinate

lyase (ASL) gene or universal 16S primers. The qPCR was performed on the LightCycler 480

System (Roche). The primer pairs and DNA sequences are included in Table (1).

Histology

The tissue was collected in the same manner as for RNA, placed in cassettes and fixed in

10% formalin for H&E staining or Carnoy solution (ThermoFisher) for fluorescent *in situ*

hybridization (FISH) staining overnight at room temperature. Cassettes were transferred to 70%

ethanol for formalin or 100% ethanol after Carnoy fixation to wash out the fixative. The tissue

was embedded in paraffin, and slides were cut at 5 µm thickness. The H&E staining was

performed by the Human Tissue Resource Center at the University of Chicago. For FISH staining,

the paraffin was first removed by running the slides through four 3-min incubations in xylene

and four 3-min incubations in 100% ethanol. The slides were then moved to a polypropylene

slide container and filled with hybridization solution containing the diluted 16S probe (0.9M

NaCl, 20mM Tris-HCL pH 7.5, 0.1% SDS with 0.2ng of probe specific for SFB 16S or universal 16S)

(*43*). The 16S probes used are included in Table (1). The slides were incubated overnight at 50

°C in the dark. The slides were washed three times with the hybridization buffer, briefly rinsed

in H_2O, and then mounted with Prolong diamond antifade with DAPI (ThermoFisher). The slides

were scanned with the CRi Pannoramic SCAN 40x Whole Slide Scanner at the University of

Chicago Integrated Light Microscopy core.

ELISA

To quantify luminal IgA levels, content was collected from the jejunum and ileum and

weighed in 2-ml bead beating tubes containing 0.1mm glass beads. After adding 1 ml of 1X cell

lysis buffer with protease inhibitors (Cell Signaling Technologies), the content was homogenized

on a vortex for 5 min. The debris were pelleted at 13000 rpm for 10 min, and the supernatant

was collected for ELISA. For tissue explants, 1 cm of tissue was excised and opened

longitudinally, washed in PBS, and placed in complete RPMI at 37 °C for 24 h. The culture

supernatant was collected and used for ELISA. The supernatant was diluted in 1X assay diluent

A (ThermoFisher), and the dilution in the middle of the standard curve was used to quantify IgA

levels. IgA mouse uncoated ELISA kit (ThermoFisher) was used following the manufacturer's

protocol, and absorbance was read at 450 nm. The amounts of IgA were back calculated to the

original sample and normalized relative to the weight of the content or to ml of culture

supernatant.

Luminal IgA isolation and *in vivo* treatment

To isolate luminal polyclonal sIgA from the intestine, luminal content was pooled from

the small intestine, large intestine, and cecum from SFB+ 8–12 week old WT CD1 mice (Charles

river). Content was transferred to falcons containing 1X Tris-Buffered Saline, 0.1% Tween 20

(TBST) buffer with proteinase inhibitor (Roche). Falcons were then vortexed for 5 min on max

speed and centrifuged for 10 min at 5000 rpm. The supernatant was collected and spun again

two times to further remove bacteria and debris. Pierce Protein L Magnetic Beads (Thermo

Scientific) were added and incubated for 1 h at room temperature while shaking. After 1 h,

beads selectively bound to IgA through kappa light chain were separated from the supernatant

with EasySep magnetic stand (StemCell technologies). Supernatant was discarded and beads

washed 3 times. IgA was separated from the beads with Pierce IgG Elution Buffer pH 2.0

(ThermoFisher). Elution buffer was incubated with the beads for 10 min at room temperature

on a shaker. Tris-HCl 1M pH 8.5 was added to neutralize the solution. IgA protein concentration

was measured using NanoDrop. The isolated IgA preparation was then filtered with 0.22 µm

sterile syringe filter unit and protease inhibitors were added (ThermoFisher). The IgA

preparation was kept up to one week at 4 °C. When IgA were administered to the mice by

gavage, the isolated IgA preparation was further concentrated with Amicon Ultra-4 Centrifugal

Filter Units (MilliporeSigma) until 250–350µg/0.1ml final concentration was achieved. To

optimize the treatment protocol to restore luminal IgA levels *in vivo* to GATA4$^{\Delta IEC}$ mice, we first

treated RAG-/- mice with 250 µg of the IgA preparation. After 1 h, we assessed the frequency of

IgA+ bacteria in small intestinal contents by flow cytometry as described below, and noted 10–

20% of bacteria were IgA+ after gavage. The bacterial coating was transient due to intestinal

flow and undetectable after two hours. Therefore, continuous IgA gavages were necessary to

sustain luminal IgA and bacterial coating in the small intestine. To administer the IgA

preparation and analyze SFB colonization, GF, WT and GATA4$^{\Delta IEC}$ mice were gavaged with 100 µl

of IgA or PBS. After 1 h, the mice were colonized with SFB and gavaged again with IgA or PBS.

Three more gavages were performed at 2-h intervals. The mice were euthanized 24 h after the

gavage of SFB, and the regionalization of SFB load was assessed in jejunal and ileal mucosal

scrapings by qPCR.

Bacterial staining with luminal IgA

Luminal content was taken from WT, GATA4$^{\Delta IEC}$, or RAG-/- mice and resuspended in 1X

PBS with protease inhibitors at a concentration of 0.1mg/µl, vortexed for 5 min, and spun at

8000 rpm for 5 min. Three fecal pellets from RAG-/- mice were homogenized and pelleted. The

bacterial pellet was resuspended in 50µl PBS and combined with 50µl of luminal supernatant

containing IgA. The IgA was incubated with the bacteria for 1 h at 4 °C. The bacteria were then

washed, pelleted, and stained with SYTO BC (ThermoFisher) diluted 1:5000 and anti-IgA APC

(Southern Biotech) diluted 1:200 for 30 min. Bacteria were gated on FSC, SSC, SYTOBC+, and

IgA+.

***Citrobacter rodentium* infections**

C. rodentium strains DBS100, DBS120 *pler-lux*, or DBS100 ΔEAE were grown at 37 °C in

Luria broth under agitation (*12, 44*). The cultures were diluted 100X and grew to log phase until

the OD^{600nm} reached 0.75. For gavage, 200 µl of bacteria were used, which gave a dose of

2.5X10^9 CFU/mouse. Mice were separated into cages based on genotype for infections, and

male mice were used for survival studies. DBS100 or DBS100 ΔEAE strains were given to GF

mice and DBS120 *pler-lux* was given to SPF mice. The DBS120 strain has a genomic kanamycin

resistance cassette inserted through Tn5. To determine CFUs of DBS120, 2 fecal pellets/mouse

were resuspended in 1 ml of PBS and plated on MacConkey agar containing 50 µg/ml of

kanamycin. The CFU/mg feces concentration was determined as: (#CFU counted*Dilution

factor/(vol plated in ml))/mg feces. To determine the amount of bacterial translocation, the

MLN, liver, and spleen were aseptically dissected, weighed, and homogenized with the Tissue-

Tearor rotor (BioSpec) in 500 µl of PBS. Then 200 µl of homogenate was plated on MacConkey

agar containing 50µg/ml of kanamycin.

Microbial transfers

To colonize mice with SFB (*44*) or rat SFB (*12*), 3–4 fresh fecal pellets from SFB

monocolonized mice were homogenized in 1 ml of PBS, vortexed for 3 min, and spun at 300 g

to remove large debris. Then 200 µl of the homogenate were gavaged to recipient mice. When

possible, SFB donor pellets were taken from monocolonized Jh or IgA deficient mice, which

harbor 10-fold higher levels of SFB. To colonize mice with SFB-free microbiota, C57BL6 mice

from Jackson Labs, which lack SFB in the microbiota, were used as donor mice. Small intestinal

and cecal contents were pooled for one donor mouse homogenized in PBS, and gavaged to

recipient mice with or without SFB supplemented. For WT and GATA4$^{\Delta IEC}$ microbiota transfer to

GF WT and GATA4$^{\Delta IEC}$ hosts, jejunal content was pooled from two donor SPF GATA4$^{\Delta IEC}$ mice or

littermate WT mice. Colonization of ASF strains (Taconic) was performed as described

previously (*45*) and gavaged to recipient WT and GATA4$^{\Delta IEC}$ mice. For all microbial transfers,

mice were colonized at 4 weeks of age and analyzed at 8 weeks.

Vitamin A deficient diet

GF C57BL6 mice were placed on control (Harlan TD.91280) or vitamin A deficient (Harlan

TD. 86143) diets from 4 to 8 weeks of age. At 8 weeks, mice were monocolonized with SFB for

one week as described above, and the amount of SFB in jejunal mucosal scrapings was

quantified by qPCR.

SFB TCRtg adoptive transfer

Naïve SFB TCRtg Vβ8 CD4 T cells were isolated from LNs and spleen of congenically

marked CD45.1 Vβ8+/- female mice using the naïve CD4 T cell isolation kit (Miltenyi), and $2x10^5$

cells/100μl mouse were injected retroorbitally into CD45.2 WT and GATA4$^{\Delta IEC}$. Three days after

transfer, the mice were euthanized to assess T cell priming and activation in the jejunal and ileal

draining MLN as described previously (Esterházy et al., 2019). To assess T cell expansion in the

LP of the jejunum nine days after transfer, 50,000 cells were injected/mouse.

16S sequencing libraries and analysis

The following 16S methods that were used were adapted from a paper currently in

press (*46*). Extracted DNA was amplified, barcoded, and sequenced as described previously (*36,*

47, 48). Briefly, amplification of the variable 4 (V4) region of the 16S rRNA gene was performed

in 20 uL duplicate reactions: 8 uL of 2.5X 5Prime Hotstart Mastermix (VWR, Radnor, PA, USA), 1

uL of 20X Evagreen (VWR), 2 uL each of 5 uM forward and reverse primers (519F, barcoded

806R, IDT, CoralVille, IA, USA), variable input volumes of extracted DNA template and nuclease

free water. Total DNA input (determined by NanoDrop) was limited to 400 ng to prevent

inefficient amplification. Amplification reactions were monitored on a CFX96 RT-PCR machine

(Bio-Rad Laboratories, Hercules, CA, USA), and samples were removed in the late exponential

phase to minimize chimera formation and non-specific amplification (1,4,5). Amplification

cycling conditions were as follows: 94 °C for 3 min, up to 50 cycles of 94 °C for 45 s, 54 °C for 60

s, and 72 °C for 90 s. Successfully amplified duplicates were pooled together and quantified

with KAPA library quantification kit (Roche, Basel, Switzerland), and then all samples were

combined at equimolar concentrations with up to 96 samples per library. AMPureXP beads

(Beckman Coulter, Brea, CA, USA) were used to clean up and concentrate libraries before final

library quantification with a High Sensitivity D1000 Tapestation Chip (Agilent, Santa Clara, CA,

USA). Illumina MiSeq sequencing was performed with a 2x300bp reagent kit by Fulgent

Genetics (Temple City, CA, USA). Raw reads were demultiplexed by Fulgent Genetics.

Demultiplexed forward and reverse reads were processed with QIIME 2 2020.2 (*49*). Loading of

sequence data was performed with the demux plugin followed by quality filtering and denoising

with the dada2 plugin (*50*). Dada2 trimming parameters were set to the base pair where the

average quality score dropped below thirty. All samples were rarefied to the lowest read depth

present in all samples (48,305 reads) to decrease biases from variable sequencing depth across

samples (*51*). The q2-feature-classifier was then used to assign taxonomy to amplicon sequence

variants (ASV) with the Silva 132 99% OTUs references (*52, 53*). Resulting read count tables

were used for downstream analyses in IPython notebooks.

Absolute abundance

The total microbial load (bacteria and archaea) of each sample and the absolute

abundance of each taxon in individual samples was determined as described previously (*36,*

47). Briefly, the Bio-Rad QX200 droplet dPCR system (Bio-Rad Laboratories) was utilized to

measure the 16S concentration in each sample with the following reaction components: 1X

QX200 EvaGreen Supermix (Bio-Rad), 500 nM forward primer, and 500 nM reverse primer (Uni

16S 519F, Uni 16S 806R) and thermocycling conditions: 95 °C for 5 min, 40 cycles of 95 °C for 30

s, 52 °C for 30 s, and 68 °C for 30 s, followed by a dye stabilization step of 4 °C for 5 min and 90

°C for 5 min. The final concentration of 16S rRNA gene copies in each sample was corrected for

dilutions and normalized to the extracted sample total DNA measurement from NanoDrop. We

used total DNA levels as a proxy for tissue mass in biopsy samples. For each sample, the input-

DNA-normalized total microbial load from dPCR was multiplied by each amplicon sequence

variant's (ASV) relative abundance to determine the absolute abundance of each ASV.

Poisson quality filtering

Two separate quality-filtering steps based on Poisson statistics were used to determine

the statistical confidence in the measured values. First, a 95% confidence interval was

calculated from the repeated measures of water blanks. Samples with a total microbial load

below the upper bound of this confidence interval were removed from further analysis.

Second, the limit of detection (LOD) in terms of relative abundance was determined for

each sample. Sequencing can be divided into two separate Poisson sampling steps. First, an

aliquot of sample is taken from the extracted sample and input into the library amplification

reaction. The LOD of the library amplification step (LOD_{amp}) was determined by multiplying the

total microbial load from dPCR by the input volume into the library amplification reaction and

then finding the relative abundance corresponding to an input of three copies. Poisson statistics

tells us that the likelihood of sampling one or more copies with an average input of three copies

is 95%. The second Poisson sampling step in sequencing arises from the number of reads

generated from the amplified library. Here, the relative abundance at which 95% confidence of

detection is observed was previously shown to follow a negative exponential curve with respect

to the total read depth: LOD_{seq} = 7.115 * read_depth$^{-0.115}$ (*36*) The minimum   LOD =

min(LOD_{amp}, LOD_{seq}) of the two described LODs was then determined for each sample. For each

sample, the abundance of any ASV with a relative abundance below the LOD was set to zero.

After this filtering step, data tables for each taxonomic level were generated.

Statistical analysis and correlations

Group comparisons were analyzed using the non-parametric Kruskal-Wallis rank sum

test with Benjamini–Hochberg multiple hypothesis testing correction using *SciPy.stats Kruskal*

function and *statsmodels.stats.multitest multipletests* function with the *fdr_bh* option.

Purified IECs RNA-seq

To perform RNAseq on IECs, 100,000 EPCAM+ CD45- cells were cell sorted from the

jejunum and ileum of WT and GATA4$^{\Delta IEC}$ mice. Three independent cell sorting experiments were

performed and the libraries and sequencing were done on the same batch with 8 mice per

group. The SMART-Seq v4 Ultra Low Input RNA Kit (TaKaRa) was used to generated amplified

cDNA, using 7500 pg of RNA input. The cDNA was generated and purified according to the

manufacturer's specifications, and cDNA was amplified 12 cycles. The Nextera XT DNA Library

Preparation Kit (Illumina) was used to generate the RNA-seq libraries, with an input of 125 pg

cDNA, according to the manufacturer's specifications. Subsequently, the libraries were

multiplexed and sequenced at a depth of 20 million reads per sample (50bp SR) on a

HiSeq4000.

Whole tissue RNA-seq

Whole tissue biopsies stored at −80 °C were thawed on ice and transferred to Starstedt

tubes containing 350uL RLT Plus supplemented with 1% 2-mercaptoethanol and equal

quantities of 1.0 mm and 0.5 mm zirconium oxide beads (Next Advance). Biopsies were bead

beat 3 times for 1 min at a setting of 9 on a Bullet Blender 24, with 1 min of cooling on ice

between each beating. Lysates were processed using the AllPrep DNA/RNA/miRNA Universal Kit

(Qiagen). To generate sample libraries, 500 ng of purified RNA was used as input in the TruSeq

Stranded mRNA Library Prep kit (Illumina) according to manufacturer's specifications. Libraries

were multiplexed and sequenced at a depth of 20 million reads per sample (50 bp SR) on a

HiSeq4000.

Mouse RNA-seq data processing to obtain raw counts

Mouse RNA-seq raw data were processed using a standard workflow based on the

GENPIPES framework (*54*). Specifically, the "stringtie" type "rnaseq" pipeline was used. Reads

were first trimmed using Trimmomatic software (*55*). Trimmed reads were aligned to the

Mus_musculus.GRCm38 mouse reference genome using the STAR aligner (*56*) following a two-

pass mapping protocol. Alignments were then sorted and filtered for duplicates using

Picard(sort, markduplicates) ("Picard Toolkit" Broad Institute

http://broadinstitute.github.io/picard/; Broad Institute). Gene-level read counts for

downstream processing were calculated from spliced alignments using HTseq count (*57*).

RNA-seq data analysis: quality control filtering and normalization

All statistical analyses of the mouse RNA-seq data were performed using R(v4.0.3).The

raw count matrices contained the transcript counts, with columns representing samples and

rows indexed by genes. Genes expressed (*i.e.*, having at least two counts) in fewer than two

samples were removed. The resulting matrices will be referred to as the count matrices. The

variance stabilizing transformation from DESeq2 R package (*58*) was applied to normalize the

count matrices. Batch effects were removed using the *removeBatchEffect()* function from the

limma R package (*59*).

RNA-seq data analysis: differential expression and gene set enrichment

A negative binomial generalized linear model in the DESeq2 package was used to test

whether genes were differentially expressed (DE) between sample groups. The analysis takes

the count matrix as an input and estimates size factors and dispersions prior to fitting the

model. Wald statistics were used to determine the significance and the log2-fold change (LFC)

of the fit for each gene. We used the Benjamini-Hochberg method for reducing the false

discovery rate (FDR), with a cutoff of <0.05 for identifying differentially expressed genes for

further analysis. *P* values reported below are FDR adjusted. The LFCs and FDR-adjusted *P* values

were given as input to the *fgsea()* function from the fgsea R package (*60*), which implements a

preranked gene set enrichment analysis. The rankings of the genes were based on the FDR-

adjusted *P* values. The mouse KEGG pathway database (mmuKegg) (*61*) and/or the Gene

Ontology, Biological Processes (GO-BP) database (*62, 63*) were the gene sets used in the

enrichment analysis. Enriched pathways (*i.e.*, *P* <0.05) were collapsed to independent pathways

to avoid repetition, using the fgsea *collapsePathways()* function (*60*).

(i) Identifying region-specific GATA4-regulated genes

We determined this set of DE genes by grouping together "ileum-like" samples, *i.e.*, WT

ileum and GATA$^{\Delta IEC}$ jejunum samples, and comparing them with WT Jejunum samples. This was

done separately for the tissue and EC RNA-seq analyses. A threshold of >0.25 for the absolute

value of the LFC was used to filter the very high number of DE genes in the tissue RNA-seq data,

whereas no LFC threshold was used for the EC data.

(ii) Microbiota dependent and independent genes

We determined the influence of microbiota on the region-specific GATA4-regulated

genes by first systematically comparing jejunum tissue samples to identify: (1) DE genes

between GATA$^{\Delta IEC}$ SPF in comparison to WT SPF, (2) DE genes between GATA$^{\Delta IEC}$ GF in

comparison to WT GF, and (3) genes whose expression indicated a significant interaction

between genotype (*i.e.*, KO vs WT) and microbiota (*i.e.*, GF vs SPF).

(a) Microbiota dependent genes: Genes that were strongly DE (*i.e.*, $P < 0.01$, $|LFC| > 0.6$)

in Group 1 but not DE (*i.e.*, $P > 0.2$) in Group 2, or vice versa, were deemed microbiota

dependent. Further, genes in group 3 that had a strong, significant interaction term (*i.e.*, P

< 0.01, $|LFC| > 0.9$) were also included.

(b) Microbiota independent genes: Genes that were strongly DE (i.e., $P < 0.01$, $|LFC| >$

0.6) in both groups were deemed microbiota independent. Further, genes in group 3 that also

had a weak or insignificant interaction term (i.e., $P > 0.2$ or $|LFC| < 0.9$) were also added to the

list of microbiota independent genes.

Both these sets of genes were intersected with the region-specific, GATA4-regulated genes,

resulting in region-specific, GATA4-regulated genes that are either microbiota dependent or

independent.

(iii) Genes regulated by retinoic acid

We determined this set of DE genes by grouping together RA-treated, GATA$^{\Delta IEC}$ jejunum

and vehicle-treated WT jejunum samples, and comparing them to vehicle-treated GATA$^{\Delta IEC}$

jejunum samples. Due to the small number (4) of samples per group, many genes were dropped

due to the FDR threshold. To get a better view of the overall trends, DE genes were instead

determined in this step by thresholding on the unadjusted p-value (P <0.05). As a sanity check,

we observed that using the same P value threshold to compare just vehicle-treated WT jejunum

samples with vehicle-treated GATA$^{\Delta IEC}$ jejunum samples resulted in a set of DE genes similar to

the region-specific GATA4-regulated DE genes identified in the first analysis.

RNA-seq data analysis: data visualization

(i) Principal components plots

The principal components analysis (PCA) was done using the *pca()* function from the

PCAtools R package (Blighe K, Lun A, https://github.com/kevinblighe/PCAtools). The top 500

genes selected by highest row variance in the centered (across rows) normalized count matrix

were used to calculate the principal components. Using the full set of genes to calculate the

PCA shows the same patterns but with a larger spread amongst sample groups.

(ii) Heatmaps

Heatmaps were plotted using the *Heatmap()* function from the ComplexHeatmap R

package (*64*) . The z-scored (i.e., across rows) normalized expression values were used to plot

the heatmaps.

Annotation of GATA4-targeted genes

A previously published and publicly available table of annotated GATA4 ChIP-Seq peaks

(*11*) was used to annotate GATA4 targets among the DE genes reported.

Generation of IL-17 and IFNγ gene modules

We curated an IL-17 and IFNg modules, which consisted of known gene pathways from

established databases and publications. The IL-17 module consisted of genes encompassing the

following pathways from mmuKEGG: 04657, 04659 and msigDB: m6335, m19422, m298, m300,

m461, m460, m39560, m8578, m8581, m8579, m8927, m8928, and the following papers (*12,*

18).

The IFNg module consisted of genes encompassing the following pathways from

msigDB: m22085, m5972, m5970, m4551, m9583 , m39363, m161, m6305, m6313, m6696,

m6695, m6689, m6688, m6523, m6522, m6513, m6512, m1898, m2913, m8662, m8657,

m5913, from the Gene Ontology database: GO:0034341, from the Reactome database: R-

913531, and the following papers (*65, 66*).

Human celiac RNA-seq analysis

Adaptors and low-quality bases were trimmed using Trim Galore (v 0.4.4). Resulting

reads were then aligned to the human reference sequence Ensembl GRCh38 release 87 using

Kallisto (*67*). Next, the derived pseudo counts were normalized into log2 counts per million

reads (CPM) using the voom function from the limma package (v3.46.0) (*59*). To evaluate the

transcriptomic changes associated to GATA4 de-regulation in the small intestine and in the

context of celiac disease we defined two contrast groups: GATA4 low, to reflect loss of

regionalization, and GATA4 hi, as a normal jejunum tissue. To define these two groups, we used

the normalized expression of GATA4 were then used to rank all control, ACeD and GFD samples

to define two contrast groups of samples. First, by grouping the top 30% GATA4 expressing

samples into the GATA4 hi group (ACeD n =6, Control n= 18, GFD n= 18) and, second by

considering the bottom 30% of GATA4 expression only from ACeD samples, (n=15) which we

defined as the GATA4 low group. Next, to define a universe set for the enrichment analysis we

generated a set of 11,657 homologous genes from the human and mouse cohort. We then

tested transcriptome wide for significant differences in expression across the GATA4 hi and

GATA4 low groups of samples, by using a linear model while accounting for sex, age, batch and

technical covariates and correcting for multiple testing. Over-enrichment analysis of gene

ontologies biological processes were performed using the enrichGO function from the

clusterProfiler (v3.0.4) (*68*). All statistical analyses within the human cohort and the overlaps

with the mouse GATA4$^{\Delta IEC}$ dataset were performed using R (v4.0.3).

Using on the log2(CPM) expression values, we calculated single sample Gene Set

Enrichment scores (ssGSEA) for the Retinol pathway, IL17 downstream genes and the hallmark

(*69*), using the R Bioconductor Package GSVA (*70*). These ssGSEA scores where then used to test

association between GATA hi vs GATA lo contrasts groups, as well as its association with

presence or absence of bacteria. Associations where evaluated using a linear model while

accounting for sex, age and technical covariates.

1000

1001

1002

1003

1004

1005

1006

1007   **Table 1**

Reagent or Resource	Source	Identifier
Antibodies		
CD45 Pac Blue (30-F11)	Biolegend	Cat#103126
TCRgd FITC (eBioGL3)	Thermo Fisher	Cat#11-5711-82

CD4 BV785 (GK1.5)	Biolegend	Cat#100453
CD4 BV605 (GK1.5)	Biolegend	Cat#100451
CD8β BUV395 (H35-17.2)	BD	Cat#740278
CD8α PerCp/Cy5.5 (53-6.7)	BD	Cat#551162
NK1.1 PE-CF594 (PK136)	BD	Cat#562864
TCRβ BUV737 (H57-597)	BD	Cat#612821
TCRβ BV711 (H57-597)	BD	Cat#563135
CD3ε BUV737 (145-2C11)	BD	Cat#612771
IFNγ APC (XMG1.2)	BD	Cat#554413
IL10 PEcy7 (JES5-16E3)	Biolegend	Cat#505026
IL17a PE (ebio17B7)	Thermo Fisher	Cat#12-7177-81
CD45.1 Pac Blue (A20)	Biolegend	Cat#110722
CD45.2 BUV395 (104)	BD	Cat#553772
vβ14 TCR FITC (14-2)	BD	Cat#553258
RORγt BV786 (Q31-37)	BD	Cat#564723
FOXP3 eflour450 (FJK-16s)	Thermo Fisher	Cat#48-5773-82
FOXP3 FITC (FJK-16s)	Thermo Fisher	Cat#11-5773-82
FOXP3 PE-cy7 (FJK-16s)	Thermo Fisher	Cat#25-5773-82
Tbet PE (4B10)	Biolegend	Cat#644810
CD44 PE-CY7 (IM7)	Biolegend	Cat#103030
CD62L PE (MEL-14)	Biolegend	Cat#104408
Epcam PerCp/Cy5.5 (G8.8)	Biolegend	Cat#118220
CD19 FITC (1D3/CD19)	Biolegend	Cat#152404
NK1.1 BV605 (PK136)	Biolegend	Cat#108753
CD11C BV605 (N418)	Biolegend	Cat#117334
TER119 BV605 (TER-119)	Biolegend	Cat#116239
F4/80 BV605 (BM8)	Biolegend	Cat#123133
CD3ε BV605 (145-2C11)	Biolegend	Cat#100351
Ly6g BV605 (1A8)	Biolegend	Cat#127639
B220 PE-cy7 (RA3-6B2)	Biolegend	Cat#103222
IgA PE (mA-6E1)	Thermo Fisher	Cat#12-4204-81
IgA AF647 goat polyclonal	Southern Biotech	Cat#1040-31
Biological Samples		
Fetal Bovine Serum	Biowest	Cat#S01520; Lot#A11504E
Normal Goat Serum	JacksonImmunoResearch	Cat#005-000-121
Chemicals, Peptides, and		

Recombinant Proteins		
EDTA, 0.5M, pH8.0	Corning	Cat#46-034-CI
1M MgCl2	Thermo Fisher	Cat#AM9530G
Cytiva Percoll™ Centrifugation Media	GE Healthcare	Cat#45-001-747
RNAprotect Tissue Reagent	Qiagen	Cat#76106
2-Mercaptoethanol (BME)	Sigma-Aldrich	Cat#M7154
Phorbol Myristate Acetate	Sigma-Aldrich	Cat#P1585
Ionomycin Calcium Salt from Streptomyces conglobatus	Sigma-Aldrich	Cat#10634
BD GolgiStop Protein Transport Inhibitor	BD	Cat#554724
Ethanol 200 Proof	Decon Labs Inc	Cat#DSP-MD 43
Inhibitex Buffer	Qiagen	Cat#19593
Nuclease-free Water	Ambion	Cat#AM9932
Carnoy Solution	Ricca Chemical	Cat#R18510004C
10% Formalin Solution	Thermo Fisher	Cat#SF98-4
RPMI 1640 with L-Glutamine	Corning	Cat#MT-10043CV
Collagenase from Clostridium histolyticum	Sigma-Aldrich	Cat#C2139-500MG
1M TRIS-HCL pH 7.5	Thermo Fisher	Cat#15567027
10% SDS solution	Thermo Fisher	Cat#15553027
Sodium chloride	Sigma-Aldrich	Cat#S9888
Critical Commercial Assays		
SytoBC	Thermo Fisher	Cat#S34855
LIVE/DEAD® Fixable Aqua Dead Cell Stain Kit	Thermo Fisher	Cat#L34966
Zombie NIR™ Fixable Viability Kit	Biolegend	Cat#423106
BD Cytofix/Cytoperm Plus Fixation/Permeabilization Solution Kit	BD	Cat#554714
eBioscience™ Foxp3 / Transcription Factor Staining Buffer Set	Thermo Fisher	Cat#00-5523-00
QIAamp Fast DNA Stool Mini Kit	Qiagen	Cat#51604
RNeasy Plus Mini Kit	Qiagen	Cat#74136
RNeasy Micro Kit	Qiagen	Cat#74004
GoScript Reverse Transcriptase Kit	Promega	Cat#A5001
SYBR Advantage qPCR Premix	Clontech	Cat#639676

ProLong™ Diamond Antifade Mountant with DAPI	Thermo Fisher	Cat#P36962
Pierce™ Protein L Magnetic Beads	Thermo Fisher	Cat#88850
Pierce™ IgG Elution Buffer, pH 2.0	Thermo Fisher	Cat#21028
1.0 M Tris HCl pH 8.5	VWR	Cat#76236-402
ALDEFLUOR™ Kit	Stemcell technologies	Cat#01700
IgA mouse ELISA	Thermo Fisher	Cat#88-50450-86
QX200 droplet digital PCR system	BioRad	Cat#1864001
QX200 ddPCR EvaGreen Supermix	BioRad	Cat#1864034
5Prime Hotstart Mastermix	QuantaBio	Cat#2200410
Evagreen	Biotium	Cat#31000
KAPA library quantification kit	Roche	Cat#07960140001
AmpureXP beads	Beckman Coulter	Cat#A63880
Oligonucleotides		
GAPDH Forward 5ʹ-AGGTCGGTGTGAACGGATTTG-3ʹ	(*38*)	N/A
GAPDH Reverse 5ʹ-TGTAGACCATGTAGTTGAGGTCA-3ʹ	(*38*)	N/A
IFNγ Forward 5ʹ-ATGAACGCTACACACTGCATC-3ʹ	(*38*)	N/A
IFNγ Reverse 3ʹ-TCTAGGCTTTCAATGACTGTGC-5ʹ	(*38*)	N/A
ASL Forward 5'-TCTTCGTTAGCTGGCAACTCACCT-3'	(*71*)	N/A
ASL Reverse 5'-ATGACCCAGCAGCTAAGCAGATCA-3'	(*71*)	N/A
Uni 16S 340F 5'-ACTCCTACGGGAGGCAGCAGT-3'	(*42*)	N/A
Uni 16S 514R 5'-ATTACCGCGGCTGCTGGC-3'	(*42*)	N/A
SFB 736F 5'-GACGCTGAGGCATGAGAGCAT-3'	(*42*)	N/A
SFB 844R 5'-GACGGCACGGATTGTTATTCA-3'	(*42*)	N/A
ASF457F 5'-TGCAAGAATGAAACTCAAAGGAAT-3'	(*72*)	N/A

ASF457R 5'-TAAGGTTCTTCGGTTAGCATCGA-3'	(72)	N/A
C. rodentium espBF 5'-ATGCCGCAGATGAGACAGTTG-3'	(73)	N/A
C. rodentium espBR 5'-CGTCAGCAGCCTTTTCAGCTA-3'	(73)	N/A
EUB338 5'-Alexa546-GCTGCCTCCCGTAGGAGT-3'	(74)	N/A
EUB338 5'-Cy3-GCTGCCTCCCGTAGGAGT-3'	(74)	N/A
SFB1008 5'-Alexa488-GCGAGCTTCCCTCATTACAAGG-3'	(42)	N/A
SFB1008 5'-Cy5-GCGAGCTTCCCTCATTACAAGG-3'	(42)	N/A
Uni 16S 519F 5'-CAGCMGCCGCGGTAA	(36)	N/A
Uni 16S 806R 5'-GGACTACHVGGGTWTCTAAT	(36)	N/A
Other		
Bead ruptor elite bead mill homogenizer	Omni International	Cat#19-040E
Tissue-Tearor	Biospec Products	Cat#985370-XL
Glass Beads	Biospec Products	Cat#11079101
LightCycler® 480 System	Roche	N/A
LSRFortessa™ X-20 Flow Cytometer	BD	N/A
Cytek® Aurora	Cytek	N/A
CRi Pannoramic SCAN 40x Whole Slide Scanner	3DHistech	N/A
EasyEights™ EasySep™ Magnet	Stemcell technologies	Cat#18103
CFX96 Touch RT-PCR detection system	BioRad	Cat#1855196
Software and Algorithms		
QIIME2: 2020.2.0	(49)	
Silva 132 SSURef NR99	(53)	
Python 3.7.6	https://www.python.org/	
Scipy 1.4.1	https://www.scipy.org/	
Statsmodels 0.10.1	https://www.statsmodels.org/stable/index.html	

Numpy 1.18.1	https://numpy.org/	
Pandas 1.0.3	https://pandas.pydata.org/	
GraphPad Prism 8	https://www.graphpad.com	
FlowJo	https://www.flowjo.com	

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

Acknowledgements:

We would like to thank the patients and their family members, as well as the University of

Chicago Celiac Disease Center, for supporting our research. We would like to thank Betty

Theriault, Kristin Kolar, and the entire Gnotobiotic Research Animal Facility at the University of

Chicago for their help. We also thank the Human Tissue Resource Center, the Integrated Light

Microscopy Core Facility, Flow Cytometry Facility, and the Genomics Facility at the University of

Chicago. We would like to thank Kenya Honda for providing rat SFB, and Gabriel Nuñez for ΔEAE

C. rodentium. We would like to thank Ivaylo Ivanov for providing SFB TCR transgenic mice and

for helpful discussions. We would like to thank Toufic Mayassi, and Cezary Ciszewski for their

thoughtful discussions and insightful comments. Finally, we thank Valerie Abadie for help

editing the manuscript. This work was supported by the National Institutes of Health via T32

AI007090 to Z.M.E., T32 GM007281 to D.G.S., U01AI109695 for A.M., U01 AI125250 to A.B.,

R01 AI144094 to A.B., R01DK103761 to N.S., R01 DK067180 to B.J., and the Digestive Diseases

Research Core Center P30 DK42086 at the University of Chicago to B.J.

1034

Author Contributions:

Z.M.E. and B.J. conceived the study and designed the experiments. N.S., A.M. and P.M, based

on their previously published work and studies performed in gnotobiotic mice, put forward the

concept that absence of GATA4 in epithelial cells was driving inflammatory immune responses

in the jejunum in a microbiota-dependent manner. Z.M.E., W.L., A.K., D.G.S., J.D.E., and J.T.B.

performed the experiments. Z.M.E., W.L., J.J.S, R.A.G., J.T.B., analyzed and interpreted the data.

J.J.S., R.A.G., performed the RNAseq analysis, J.T.B., performed the 16S sequencing and

analysis. D.G.S. and V.D. created the RNAseq libraries, and S.G. performed the DNA alignments.

P.M. supervised the initial GATA4 transcriptomic analysis. I.L.T contributed to and L.B.B

supervised RNA seq analysis of human celiac disease patients, and S.J.R. supervised the overall

RNAseq analysis. R.F.I. supervised the 16S analysis. A.B. supervised the IgA experiments. Z.M.E,

S.J.R, B.J. wrote the manuscript. W.L., J.J.S, J.T.B., R.F.I., A.M., N.S., and A.B. edited the

manuscript. R.F.I., A.M., A.B., N.S., and B.J. acquired funds to support the work; and B.J.

directed the study.

1049

1050   **Declaration of Interests:**

1051   The authors declare no competing interests

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063
1064
References

1.   A. M. Mowat, W. W. Agace, Regional specialization within the intestinal immune system.
Nature reviews. Immunology **14**, 667-685 (2014).
2.   Ivanov, II *et al.*, Induction of intestinal Th17 cells by segmented filamentous bacteria.
Cell **139**, 485-498 (2009).
3.   Z. Shi *et al.*, Segmented Filamentous Bacteria Prevent and Cure Rotavirus Infection. *Cell*
179, 644-658.e613 (2019).
4.   D. Esterházy *et al.*, Compartmentalized gut lymph node drainage dictates adaptive
immune responses. *Nature* **569**, 126-130 (2019).
5.   P. M. Smith *et al.*, The microbial metabolites, short-chain fatty acids, regulate colonic
Treg cell homeostasis. *Science (New York, N.Y.)* **341**, 569-573 (2013).

6. M. E. Johansson *et al.*, The inner of the two Muc2 mucin-dependent mucus layers in
colon is devoid of bacteria. *Proceedings of the National Academy of Sciences of the*
United States of America **105**, 15064-15069 (2008).
7. T. Bosse *et al.*, Gata4 is essential for the maintenance of jejunal-ileal identities in the
adult mouse small intestine. *Mol Cell Biol* **26**, 9060-9070 (2006).
8. M. A. Battle *et al.*, GATA4 is essential for jejunal function in mice. *Gastroenterology* **135**,
1676-1686.e1671 (2008).
9. K. Breuer *et al.*, InnateDB: systems biology of innate immunity and beyond--recent
updates and continuing curation. *Nucleic Acids Res* **41**, D1228-1233 (2013).
10. B. E. Aronson *et al.*, GATA4 represses an ileal program of gene expression in the
proximal small intestine by inhibiting the acetylation of histone H3, lysine 27. *Biochimica*
et biophysica acta **1839**, 1273-1282 (2014).
11. C. A. Thompson *et al.*, GATA4 Is Sufficient to Establish Jejunal Versus Ileal Identity in the
Small Intestine. *Cellular and molecular gastroenterology and hepatology* **3**, 422-446
(2017).
12. K. Atarashi *et al.*, Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial
Cells. *Cell* **163**, 367-380 (2015).
13. Y. Umesaki, H. Setoyama, S. Matsumoto, A. Imaoka, K. Itoh, Differential roles of
segmented filamentous bacteria and clostridia in development of the intestinal immune
system. *Infect Immun* **67**, 3504-3511 (1999).
14. V. Gaboriau-Routhiau *et al.*, The key role of segmented filamentous bacteria in the
coordinated maturation of gut helper T cell responses. *Immunity* **31**, 677-689 (2009).
15. Y. Yang *et al.*, Focused specificity of intestinal TH17 cells towards commensal bacterial
antigens. *Nature* **510**, 152-156 (2014).
16. T. Prakash *et al.*, Complete genome sequences of rat and mouse segmented filamentous
bacteria, a potent inducer of th17 cell differentiation. *Cell host & microbe* **10**, 273-284
(2011).
17. G. P. Donaldson *et al.*, Gut microbiota utilize immunoglobulin A for mucosal
colonization. *Science (New York, N.Y.)* **360**, 795-800 (2018).
18. P. Kumar *et al.*, Intestinal Interleukin-17 Receptor Signaling Mediates Reciprocal Control
of the Gut Microbiota and Autoimmune Inflammation. *Immunity* **44**, 659-671 (2016).
19. N. Shulzhenko *et al.*, Crosstalk between B lymphocytes, microbiota and the intestinal
epithelium governs immunity versus metabolism in the gut. *Nature medicine* **17**, 1585-
1593 (2011).
20. A. Goncalves *et al.*, Fat-soluble vitamin intestinal absorption: absorption sites in the
intestine and interactions for absorption. *Food Chem* **172**, 155-160 (2015).
21. F. M. Herr *et al.*, Intestinal vitamin A metabolism: coordinate distribution of enzymes
and CRBP(II). *Journal of lipid research* **34**, 1545-1554 (1993).
22. J. R. Mora *et al.*, Generation of gut-homing IgA-secreting B cells by intestinal dendritic
cells. *Science (New York, N.Y.)* **314**, 1157-1160 (2006).
23. H. B. Jijon *et al.*, Intestinal epithelial cell-specific RARα depletion results in aberrant
epithelial cell homeostasis and underdeveloped immune system. *Mucosal Immunol* **11**,
703-715 (2018).

24.   S. Gattu *et al.*, Epithelial retinoic acid receptor β regulates serum amyloid A expression
and vitamin A-dependent intestinal immunity. *Proceedings of the National Academy of*
Sciences **116**, 10911 (2019).
25.   B. Jabri, L. M. Sollid, Tissue-mediated control of immunopathology in coeliac disease.
Nature reviews. Immunology **9**, 858-870 (2009).
26.   I. Monteleone *et al.*, Characterization of IL-17A-producing cells in celiac disease mucosa.
Journal of immunology (Baltimore, Md. : 1950) **184**, 2211-2218 (2010).
27.   A. Castellanos-Rubio *et al.*, TH17 (and TH1) signatures of intestinal biopsies of CD
patients in response to gliadin. *Autoimmunity* **42**, 69-73 (2009).
28.   M. Bodd *et al.*, HLA-DQ2-restricted gluten-reactive T cells produce IL-21 but not IL-17 or
IL-22. *Mucosal Immunol* **3**, 594-601 (2010).
29.   T. Mayassi *et al.*, Chronic Inflammation Permanently Reshapes Tissue-Resident
Immunity in Celiac Disease. *Cell* **176**, 967-981.e919 (2019).
30.   P. Brar *et al.*, Change in lipid profile in celiac disease: beneficial effect of gluten-free diet.
Am J Med **119**, 786-790 (2006).
31.   A. C. Bledsoe *et al.*, Micronutrient Deficiencies Are Common in Contemporary Celiac
Disease Despite Lack of Overt Malabsorption Symptoms. *Mayo Clin Proc* **94**, 1253-1260
(2019).
32.   A. Caminero *et al.*, Duodenal bacterial proteolytic activity determines sensitivity to
dietary antigen through protease-activated receptor-2. *Nat Commun* **10**, 1198 (2019).
33.   M. C. Cenit, M. Olivares, P. Codoñer-Franch, Y. Sanz, Intestinal Microbiota and Celiac
Disease: Cause, Consequence or Co-Evolution? *Nutrients* **7**, 6900-6923 (2015).
34.   M. Vuoristo, Y. A. Kesäniemi, H. Gylling, T. A. Miettinen, Metabolism of cholesterol and
apolipoprotein B in celiac disease. *Metabolism* **42**, 1386-1391 (1993).
35.   M. Xu *et al.*, c-MAF-dependent regulatory T cells mediate immunological tolerance to a
gut pathobiont. *Nature* **554**, 373-377 (2018).
36.   J. T. Barlow, S. R. Bogatyrev, R. F. Ismagilov, A quantitative sequencing framework for
absolute abundance measurements of mucosal and lumenal microbial communities. *Nat*
Commun **11**, 2590 (2020).
37.   V. D'Argenio *et al.*, Metagenomics Reveals Dysbiosis and a Potentially Pathogenic N.
flavescens Strain in Duodenum of Adult Celiac Patients. *Am J Gastroenterol* **111**, 879-890
(2016).
38.   V. Abadie *et al.*, IL-15, gluten and HLA-DQ8 drive tissue destruction in coeliac disease.
Nature **578**, 600-604 (2020).
39.   S. Husby *et al.*, European Society for Pediatric Gastroenterology, Hepatology, and
Nutrition Guidelines for the Diagnosis of Coeliac Disease. *Journal of Pediatric*
Gastroenterology and Nutrition **54**, (2012).
40.   L. Lefrançois, N. Lycke, Isolation of mouse small intestinal intraepithelial lymphocytes,
Peyer's patch, and lamina propria cells. *Curr Protoc Immunol* **Chapter 3**, Unit 3.19
(2001).
41.   J. J. Bunker *et al.*, Innate and Adaptive Humoral Responses Coat Distinct Commensal
Bacteria with Immunoglobulin A. *Immunity* **43**, 541-553 (2015).
42.   T. Sano *et al.*, An IL-23R/IL-22 Circuit Regulates Epithelial Serum Amyloid A to Promote
Local Effector Th17 Responses. *Cell* **163**, 381-393 (2015).

43. Z. M. Earley *et al.*, Burn Injury Alters the Intestinal Microbiome and Increases Gut
Permeability and Bacterial Translocation. *PloS one* **10**, e0129996 (2015).
44. J. M. Pickard *et al.*, Rapid fucosylation of intestinal epithelium sustains host-commensal
symbiosis in sickness. *Nature* **514**, 638-641 (2014).
45. M. J. Molloy *et al.*, Intraluminal containment of commensal outgrowth in the gut during
infection-induced dysbiosis. *Cell host & microbe* **14**, 318-328 (2013).
46. J.T. Barlow *et. al*, Quantitative sequencing clarifies the role of disruptor taxa, oral
microbiota, and strict anaerobes in the human small-intestine microbiome. *Microbiome*,
(2021).
47. S. R. Bogatyrev, R. F. Ismagilov, Quantitative microbiome profiling in lumenal and tissue
samples with broad coverage and dynamic range via a single-step 16S rRNA gene DNA
copy quantification and amplicon barcoding. *bioRxiv*, 2020.2001.2022.914705 (2020).
48. S. R. Bogatyrev, J. C. Rolando, R. F. Ismagilov, Self-reinoculation with fecal flora changes
microbiota density and composition leading to an altered bile-acid profile in the mouse
small intestine. *Microbiome* **8**, 19 (2020).
49. E. Bolyen *et al.*, Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).
50. B. J. Callahan *et al.*, DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* **13**, 581-583 (2016).
51. S. Weiss *et al.*, Normalization and microbial differential abundance strategies depend
upon data characteristics. *Microbiome* **5**, 27 (2017).
52. N. A. Bokulich *et al.*, Optimizing taxonomic classification of marker-gene amplicon
sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90 (2018).
53. C. Quast *et al.*, The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590-596 (2013).
54. M. Bourgey *et al.*, GenPipes: an open-source framework for distributed and scalable
genomic analyses. *Gigascience* **8**, (2019).
55. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
56. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
57. S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).
58. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
59. M. E. Ritchie *et al.*, limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).
60. G. Korotkevich *et al.*, Fast gene set enrichment analysis. *bioRxiv*, 060012 (2021).
61. R. H. Waterston *et al.*, Initial sequencing and comparative analysis of the mouse
genome. *Nature* **420**, 520-562 (2002).
62. M. Ashburner *et al.*, Gene ontology: tool for the unification of biology. The Gene
Ontology Consortium. *Nat Genet* **25**, 25-29 (2000).
63. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Res* **49**, D325-d334
(2021).

64.   Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).
65.   S. Y. Liu, D. J. Sanchez, R. Aliyari, S. Lu, G. Cheng, Systematic identification of type I and
type II interferon-induced antiviral factors. *Proceedings of the National Academy of*
Sciences of the United States of America **109**, 4239-4244 (2012).
66.   J. L. Benci *et al.*, Tumor Interferon Signaling Regulates a Multigenic Resistance Program
to Immune Checkpoint Blockade. *Cell* **167**, 1540-1554.e1512 (2016).
67.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525-527 (2016).
68.   G. Yu, L. G. Wang, Y. Han, Q. Y. He, clusterProfiler: an R package for comparing biological
themes among gene clusters. *Omics* **16**, 284-287 (2012).
69.   A. Liberzon *et al.*, The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425 (2015).
70.   S. Hänzelmann, R. Castelo, J. Guinney, GSVA: gene set variation analysis for microarray
and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).
71.   C. S. N. Klose *et al.*, Differentiation of type 1 ILCs from a common progenitor to all
helper-like innate lymphoid cell lineages. *Cell* **157**, 340-356 (2014).
72.   A. Deloris Alexander *et al.*, Quantitative PCR assays for mouse enteric flora reveal strain-
dependent differences in composition that are influenced by the microenvironment.
Mamm Genome **17**, 1093-1104 (2006).
73.   S. Sagaidak, A. Taibi, B. Wen, E. M. Comelli, Development of a real-time PCR assay for
quantification of Citrobacter rodentium. *J Microbiol Methods* **126**, 76-77 (2016).
74.   R. I. Amann *et al.*, Combination of 16S rRNA-targeted oligonucleotide probes with flow
cytometry for analyzing mixed microbial populations. *Appl Environ Microbiol* **56**, 1919-
1925 (1990).
1232

Figure 1. Intestinal epithelial cells control microbiota-dependent and independent regionalization of tissue immunity through GATA4

Figure 2. GATA4 controls the colonization of segmented filamentous bacteria to shape tissue immunity

Figure 3. GATA4 regulates regionalization of retinol metabolism and IgA to limit SFB colonization in the proximal intestine

Figure 4. Loss of intestinal regionalization impacts host susceptibility to enteropathogenic infection by *Citrobacter rodentium*

Figure 5. Loss of GATA4 expression is associated with lipid metabolic dysfunction and increased IL-17 signaling in celiac disease

Figure S1.GATA4 controls regionalization of tissue metabolism and immunity

Figure S2. GATA4 controls the colonization of regional mucosal associated bacteria to shape tissue immunity

A

B

IEL CD8αβ IFNγ+

C

LP CD4 IL-17+

D

Mucispirillum schaedleri

Figure S3. GATA4 regulates regionalization of retinol metabolism and B cell responses

Figure S4. Loss of GATA4 is associated with lipid metabolic defects, mucosal associated bacteria, and increased IL-17 signaling in celiac disease patients