1 **Tissue-resident memory T cells mediate mucosal immunity to recurrent urinary tract infection**
2
3

4 Matthieu Rousseau[1,2*], Livia Lacerda Mariano[1#], Tracy Canton[1], Molly A Ingersoll[1,2]
5
6 [1]Mucosal Inflammation and Immunity, Department of Immunology, Institut Pasteur, 75015 Paris,
7 France and Inserm U1223, Paris France
8
9 [2]Université *Paris Cité*, Institut Cochin, INSERM U1016, CNRS UMR 8104, Paris, 75014, France
10
11 [#]current address: Institut Curie, PSL Research University, INSERM U932, 75005 Paris, France
12
13 *Corresponding author
14 matthieu.rousseau@inserm.fr
15
16

Abstract

Urinary tract infection (UTI) is one of the most prevalent human bacterial infections. Due to the rapid

rise of multidrug-resistant uropathogens in global circulation, new therapeutic approaches, including

vaccination and immunotherapy, are urgently needed. However, the development of these therapies is

impeded by an incomplete understanding of how memory develops during UTI. Here, we found that

reducing bacteria load early in infection, by reducing the quantity of bacteria used for infection or with

antibiotics, completely abrogated the protective memory response. As we observed that a mixed T helper

cell polarization bias, comprised of Th1, Th2, and Th17 T cells, develops during a primary infection,

we hypothesized that reducing antigen load altered T helper cell polarization. Surprisingly, T helper cell

polarization was unchanged in these scenarios. However, we uncovered a population of tissue resident

memory (T_{RM}) T cells that was significantly reduced in the absence of sufficient antigen, demonstrating

that this population is necessary for protection. Supporting that T_{RM} T cells are sufficient to protect

against recurrent infection, depletion of systemic T cells, or treatment with FTY720 to block migration

of memory lymphocytes from lymph nodes to infected tissue, did not compromise the adaptive immune

response to a second infection. Thus, our findings point to an unappreciated mechanism of adaptive

immunity to UTI and key role for T_{RM} cells in the memory response to bacterial infection in this mucosa,

providing a target for non-antibiotic-based immunotherapy and/or new vaccine strategies to prevent

recurrent UTI.

Introduction

Development of immune memory, naturally after infection or by immunomodulatory approaches such as vaccination, protect an organism from experiencing the same infection more than once. Failure to develop adequate immune memory results in repeated or recurrent infection. Mucosal surfaces, which are contiguous with the environment, are particularly at risk for infection. Sterilizing immune memory, however, often fails to develop at these sites, for reasons that are incompletely understood.

The bladder is one such mucosal surface that, in a large proportion of the population, will be infected repeatedly, often with the same pathogen (Flores-Mireles et al., 2015). Indeed, urinary tract infection (UTI), caused predominantly by uropathogenic *E. coli* (UPEC), is one of the most prevalent human bacterial infections (Litwin et al., 2007). Women experience UTI at a significantly higher rate than men; and children, the elderly, pregnant women, and immuno-compromised individuals are at even greater risk than the general population for complications such as chronic infection, pyelonephritis, and kidney damage (Chromek and Brauner, 2007; De Nisco et al., 2019; Dielubanza and Schaeffer, 2011; Hoepelman et al., 1992; Mazor-Dray et al., 2009). Approximately one in two women will have a UTI in her lifetime, with the highest risk between the ages of 16 and 35, when women are 35 times more likely than men to be infected (Foxman, 2002). Additionally, nearly half of all infected women will have a recurrent infection, defined by similar or identical symptoms, within 6 months of their first UTI, suggesting that immune memory fails to develop following infection of the bladder (Ikaheimo et al., 1996). Although the incidence of UTI is much lower in adult men (~1%), in this population, it is always considered a complicated infection, with considerable risk of becoming a recurrent or chronic UTI (Lipsky, 1989). Finally, the prevalence of UTI increases substantially in older individuals, bringing with it increased morbidity and mortality in an aging population (Ruben et al., 1995).

Given the diversity of at-risk populations, the frequency of recurrence, and the considerable increase in multi-drug resistant uropathogen isolates in global circulation, new approaches, such as immune modulation to augment mucosal immunity, are needed to defend against chronic or recurrent UTI (Klein and Hultgren, 2020; Lacerda Mariano and Ingersoll, 2020). First line antibiotic therapy only reduces acute bacterial burden, and no treatment or therapy that protects against initial or recurrent UTI exists (Lacerda Mariano and Ingersoll, 2020). Experimental vaccination in humans and mice was reported to induce antibody production correlating with lower bacterial colonization more than 30 years ago, however, these attempts have not lead to a successful clinically available vaccine (Alteri et al., 2009; Billips et al., 2009; Klein and Hultgren, 2020; Langermann et al., 1997; Uehling et al., 1997; Uehling et al., 2001; Uehling et al., 1994; Uehling et al., 1991). Recurrent UTI likely arises due to several factors, including inadequate development of memory responses after a primary infection, but also due to development of reservoirs in the bladder and gut, which potentially escape elimination by antibiotics (Mora-Bau et al., 2015; Mysorekar and Hultgren, 2006; Spaulding et al., 2017). As the associated costs

of UTI, including treatment and lost productivity, are more than $3 billion in the US (Sivick and Mobley,

2009), with similar expenditures in Europe, a better understanding of how immune memory arises and

how it can be targeted to improve the adaptive response to UTI would have a profoundly positive societal

and economic impact, as well as improve the quality of life of those suffering from recurrent infection.

Notably, how adaptive immunity develops in response to bladder infection is still incompletely

understood (Ingersoll and Albert, 2013; Lacerda Mariano and Ingersoll, 2020). An early key

observation, that mice infected with an ovalbumin-expressing UPEC strain have ovalbumin-specific

antibodies in the serum and reduced bacterial burden following a second or challenge infection,

demonstrates that protective memory can develop during UTI (Thumbikat et al., 2006). We reported

that acute bacterial burden in the bladder following a challenge infection is lower than that observed

during a first or primary UTI (Mora-Bau et al., 2015). By contrast, bacterial burden is not reduced after

challenge infection in RAG2$^{-/-}$ mice, dendritic cell-depleted mice, and CD4$^+$ and CD8$^+$ T cell-depleted

mice, supporting that the lower bacterial burden observed after a challenge infection in wildtype mice

is mediated by an adaptive immune response (Mora-Bau et al., 2015). Together, these data support that

immune memory to UTI does arise, however, similar to the clinical picture in humans, immunity is not

sterilizing in the bladder.

Whether T cells or antibodies are the key players in bacterial clearance in the bladder is unclear. A recent

study suggested that a Th2-biased T cell response develops during UTI to promote bladder epithelial

repair, at the expense of bacterial clearance (Wu et al., 2020). While this is seemingly a potential

explanation for why the memory response to UTI is nonsterilizing, this work fails to consider that

although a type 2 biased response develops in female mice, Th2 polarization is not observed in male

mice (Zychlinsky Scharff et al., 2019). Importantly, despite the absence of a Th2-bias in male mice,

female and male mice exhibit the same degree of nonsterilizing protection against a challenge UTI

(Zychlinsky Scharff et al., 2019). Additionally, the early cytokine response to UTI, which is robust in

female mice and largely absent in male mice, suggests that a Th2 T cell bias is not the determinant factor

in memory development after bladder infection (Mora-Bau et al., 2015; Zychlinsky Scharff et al., 2019).

Thus, to understand how a memory response arises following UTI, we investigated the T cell response,

finding that a mixed Th1, Th2, and Th17 bias arose during a first and second infection. Antigen

persistence was necessary for immunity, as memory was abrogated after early antibiotic treatment or

when mice were infected with a smaller inoculum. Surprisingly, limiting antigen persistence did not

change T cell polarization, but did lead to a significant reduction in a population of T cells, identified as

tissue resident memory (T$_{RM}$) T cells. Depleting T cells from circulation or blocking lymphocyte egress

from lymph nodes did not impair memory, supporting that local mucosal memory responses, and not

systemic immunity, determine protection against UTI, revealing a specific target for new therapeutics.

Results

CD4 or CD8 T cells are sufficient for immune memory to UTI

C57BL/6 mice have significantly lower bacterial burdens in their bladders after a second or challenge infection compared to a primary infection with UPEC (Lacerda Mariano et al., 2020; Mora-Bau et al., 2015; Zychlinsky Scharff et al., 2019). Supporting that this is due to immune memory, bacterial burdens are not significantly reduced after a challenge infection in the absence of B or T cells, such as in RAG2$^{-/-}$ mice or in CD4$^+$ and CD8$^+$ T cell-depleted C57BL/6 mice (Mora-Bau et al., 2015). Given that urine from women with a history of UTI inhibits UPEC binding to human urothelial cells and that antibody depletion of this urine abrogates this inhibitory capacity (Svanborg-Eden and Svennerholm, 1978), we hypothesized that immune memory to UTI is mediated by a specific B cell response.

To test this hypothesis, we intravesically instilled 6 to 8-week-old wildtype female C57BL/6 mice or µMT$^{-/-}$ mice, which lack mature B cells (Kitamura et al., 1991), with 10^7 colony forming units (CFU) of 1 of 2 isogenic UPEC strains with resistance to ampicillin (UTI89-GFP-amp^R) or kanamycin (UTI89-RFP-kana^R). Half of the mice from each group were sacrificed 24 hours post-infection (PI) to assess the bacterial burden in the bladder after the primary infection. Resolution of infection was monitored in the remaining mice by urine sampling twice per week for 4 weeks. The sustained absence of bacteria in the urine was considered to be a resolved infection. At 4 weeks PI, all resolved mice were challenged with 10^7 CFU to model a recurrent UTI, using the isogenic UPEC strain not used for the primary infection. The distinct antibiotic resistance markers between the first and second infection allow us to distinguish UPEC that establish tissue reservoirs following primary infection from those causing the second infection (Mora-Bau et al., 2015). We sacrificed mice 24 hours post-challenge infection to assess bacterial CFU in the bladder. As expected, bacterial burden was significantly reduced 24 hours post challenge (2°) compared to the burden following primary infection (1°) in wildtype mice. Surprisingly, a comparable, statistically significant reduction in CFU was also observed in µMT$^{-/-}$ mice after challenge infection, suggesting that B cells are dispensable for memory to UTI (**Figure 1A**).

Having ruled out a role for B cells, we tested which T cells (CD4$^+$ or CD8$^+$) were necessary for immune memory to UTI. Six- to eight-week-old female C57BL/6 mice were treated with anti-CD4 or anti-CD8 depleting antibodies, or relevant isotype controls, and then infected three days later with 10^7 CFU UPEC. As above, bacterial burden was assessed in half of the animals at 24 hours PI and the remaining mice were challenged with 10^7 CFU of an isogenic UPEC strain following resolution of infection at 4 weeks. In the isotype control groups, bacterial burdens 24 hours after challenge infection were statistically significantly reduced compared to primary UTI, as expected. Unexpectedly, bacterial burdens were also significantly reduced in mice treated with anti-CD4 or anti-CD8 depleting antibodies (**Figure 1B, C**).

Together, our findings support that B cells are dispensable and either CD4$^+$ or CD8$^+$ T cells are sufficient

for immune memory to UTI.

Figure 1. CD4 or CD8 T cells mediate protection against a second UTI. Six- to eight-week-

old female C57BL/6 or μMT$^{-/-}$ mice were instilled intravesically with 10^7 CFU of UPEC strain

UTI89 and bladder bacterial burden was assessed 24 hours post-primary infection (1°, filled

circles). Urine was monitored twice per week for 4 weeks and all resolved mice with sterile urine

were challenged with an isogenic UTI89 strain and bacterial burden was assessed 24 hours after

challenge infection (2°, open circles). Graphs depict 1° and 2° CFU/bladder in (**A**) C57BL/6 or

μMT$^{-/-}$ mice, (**B**) isotype or anti-CD4 antibody-treated mice, or (**C**) isotype or anti-CD8 antibody-

treated mice. Data are pooled from 2-5 experiments, n=2 to 7 mice/group in each experiment.

Each circle is a mouse, lines are medians. Dotted red lines depict the limit of detection of the

assay, 20 CFU/bladder. Significance (adjusted p-value) was determined using a Kruskal-Wallis

test comparing bacterial burden 24 hours post 1° to 2° within a treatment group, with a Dunn's

post hoc test to correct for multiple comparisons. p-values <0.05 are in red.

T cell polarization is mixed during primary and challenge UTI

Having determined that either CD4$^+$ or CD8$^+$ T cells were sufficient for immune protection, we assessed

the T cell bias arising during infection. It was proposed that a Th2-dominated, type 2 immune response

mediates nonsterilizing memory to UTI in female mice (Wu et al., 2020). However, female mice express

a wide variety of cytokines, including those associated with type 2 immunity, but also type 1 and 3

polarization (Mora-Bau et al., 2015; Zychlinsky Scharff et al., 2019). Thus, we hypothesized that

polarization was potentially more complicated than previously appreciated.

To test this idea, we determined bladder-infiltrating CD4$^+$ T cell polarization over time after primary

and challenge UTI. We focused on CD4$^+$ T cells because CD4$^+$ T cell infiltration was more than 100

times higher than CD8$^+$ T cell infiltration following infection of naïve mice (**Supplementary Figure**

1). Following infection with 10^7 CFU of UPEC, six-week-old female C57BL/6 mice were sacrificed at

day 1, 3, and 7 post primary and challenge infection. Using intracellular staining and flow cytometry,

we determined T cell polarization by transcription factor and cytokine expression (T-bet$^+$, IFNγ$^+$: Th1;

Gata-3$^+$, IL-4$^+$: Th2; RORγT$^+$, IL-17$^+$: Th17). Compared to naïve animals (day 0), total CD4$^+$ T cells

increased approximately 10-fold in the bladder over time after primary infection (**Figure 2A**). CD4$^+$ T

cell numbers also increased between mice 7 days post-challenge UTI compared to resolved (R) mice

before a challenge infection, although the baseline in the resolved animals was significantly higher than

in naïve mice (**Figure 2A**). Instead of a distinct polarization, a mixed bias was clearly apparent among

infiltrating CD4$^+$ T cells following primary and challenge UTI (**Figure 2A**). Of the global CD4$^+$ T cell

population, T-bet$^+$ Th1 T cells made up ~10-20% of cells, Gata-3$^+$ Th2-polarized T cells represented

~5-15% of cells, and RORγt$^+$ Th17 T cells made up ~3-30% of cells over time in the primary and

challenge infections (**Figure 2B**). In addition to Th1, Th2, and Th17 T cells, a large number of FoxP3$^+$

regulatory T cells (Tregs) infiltrated the bladder after primary UTI, remained elevated compared to naïve

levels after challenge infection, and made up 10-30% of total CD4$^+$ T cells over time (**Figure 2C**). We

observed a similar mixed bias among T cells in the draining lymph nodes after primary and challenge

UTI with increased numbers of Th2, Th17, IFNγ$^+$ Th1, and IL-17$^+$ Th17 T cells in resolved mice

compared to naïve animals (**Supplementary Figure 2A, B,** and **C**). Strikingly, in the bladder no T

helper cell subset dominated the response, and indeed, only one mouse in all experiments performed

had more than 50% Th2-biased T cells in a single timepoint. Furthermore, while the numbers of IFNγ$^+$

Th1 T cells, IL-17$^+$ Th17 T cells, and Tregs were significantly increased after primary infection, we

detected no IL-4-expressing Th2-biased T cells (**Figure 2D**), in line with our previous finding that IL-

4 levels are 10-100 times lower than any other cytokine measured (Zychlinsky Scharff et al., 2019).

We were struck by the large number of infiltrating Tregs, and given that a balanced Treg/Th T cell ratio

is important for immune tolerance, and also an efficient immune response against pathogens (Falivene

et al., 2015; Yordanova et al., 2019), we calculated the ratio of Tregs to all CD4 Th cells. In the bladder,

the ratio of Tregs to all Th cells was less than 1 for all timepoints measured (**Figure 2E**). The ratio of

Tregs to Th1, Th2, or Th17 T cells individually was ≤1 in naïve mice (Day 0) and mice infected for 24

hours. However, the ratios increased over 1 at day 3 and 7 post-primary infection, and remained around

1 during the second infection (**Figure 2F**). In the lymph nodes, the ratio was close to 2 during primary

UTI and ≥ 1 during the second infection (**Supplementary Figure 2D**). The ratio of Treg to each Th

subset was greatly increased over 1 (medians from 3.5 to 32) over time, however this was more

pronounced during the primary infection compared to the second UTI (**Supplementary Figure 2E**).

Thus, Treg T cells are highly prevalent during UTI, and given their abundance, likely influence the

development of memory T cells. Together, these data support that a dominant T helper bias does not

develop during UTI. Interestingly, all CD4$^+$ Th cell subsets that increased during the first UTI remained

significantly elevated over naïve levels in resolved mice in the bladder, suggesting that these T cells may have undergone tissue-specific imprinting after a primary UTI.

Figure 2. No distinct CD4$^+$ Th cell bias is apparent during primary and recurrent UTI. Six-week-old infected female C57BL/6 mice were sacrificed at the indicated timepoints post-primary (filled circles), or post-challenge infection (open circles). Day 0 are naïve mice and 'R' denotes animals that resolved their primary UTI but were not challenged with a second infection. The graphs depict (**A**) total CD4$^+$, Th1, Th2, and Th17 T cells, (**B**) percentage of the global CD4 T cell population for the indicated Th cell subset, (**C**) percentage of the global CD4 T cell population for Treg, (**D**) total IFNγ$^+$ Th1, IL-17$^+$ Th17 T, and Treg cells in the bladder at the indicated day post primary or challenge infection. Treg/Th cell ratios for the indicated days post-primary (filled circles), or post-challenge infection (open circles) were calculated using the cell numbers presented in **A, D**. (**E**) Ratio of Treg/total CD4$^+$ Th cells and (**F**) ratio of Treg/indicated CD4$^+$ Th cell subset in the bladder. Data are pooled from 2 experiments, n=5 to 7 mice/group in each

experiment. Each circle is a mouse and lines are medians. Nonparametric Mann-Whitney tests

comparing cell numbers between '0' and 'R' were performed. All *p*-values were corrected for

multiple comparisons across all populations shown using the false discovery rate (FDR) method

and *p*-values <0.05 are in red.

Antigen persistence determines the development of adaptive immunity during UTI

We observed that Tregs outnumbered Th cell subsets, potentially inducing a suppressive or tolerant

environment in response to UTI. As tolerance can be induced following suboptimal (too much or too

little) antigen stimulation (Xiao and Link, 1997; Yoshida et al., 1997), we hypothesized that the

abundant Treg infiltration reflected antigen quantity (*i.e.*, bacterial inoculum). We reported that

increasing the inoculum from 10^7 CFU/mouse to 10^8 or 10^9 CFU/mouse does not promote a stronger

memory response to a second UTI, supporting that 10^7 CFU/mouse is sufficient antigen to induce an

adaptive immune response, but may be too high leading to some degree of tolerance (Mora-Bau et al.,

2015). Thus, we tested whether decreasing the inoculum, to reduce antigen load, would induce a stronger

memory response. We intravesically infected female C57BL/6 mice with 10^7, 10^5, or 10^3 CFU of UPEC.

One cohort of mice from each group was sacrificed 24 hours post-primary infection to assess bladder

bacterial burden and a second cohort was monitored for resolution over 4 weeks. Resolved animals were

challenged 28 days later with 10^7 CFU of the isogenic UPEC strain and we assessed CFU in the bladder

24 hours post-challenge infection. Mice infected with 10^5 or 10^3 CFU resolved their primary infection

faster than those infected with 10^7 CFU (**Figure 3A**). Notably, compared to the control infection with

an inoculum of 10^7 CFU, mice infected with 10^5 or 10^3 CFU for the primary infection were no longer

protected against a challenge infection and had higher bladder-associated bacterial burdens after their

second UTI compared to the control group (**Figure 3B**). Indeed, mice infected with 10^3 CFU during

their primary infection had markedly reduced primary bacterial burdens near or at the limit of detection

of the assay and, after a challenge infection, had bacterial burdens closely resembling the level observed

after infection of naïve mice. Together, this supports that immune memory requires persistent and robust

bacterial colonization in the bladder, and low-level exposure to UPEC cannot induce immune memory.

To directly test whether antigen persistence was necessary for immune memory to UTI, we infected

mice with the same inoculum and treated them with antibiotics at specific days PI. In this scenario, the

initial antigen exposure of 10^7 CFU is the same, but antigen persistence is disrupted at different

timepoints post-infection. The advantage of this approach is that antibiotics are first line therapy to treat

UTI, and thus, we modeled the impact of a clinical scenario on immune memory. UPEC-infected female

C57BL/6 mice were given trimethoprim/sulfamethoxazole, an antibiotic combination commonly

prescribed for UTI, in the drinking water for 5 days starting at 1, 2, 3, or 5 days PI. All resolved animals

were challenged with an isogenic UPEC strain 28 days PI and bacterial burdens at 24 hours post-

challenge infection were compared to CFU at 24 hours post-primary infection. When antibiotics were

given at day 1 post-primary infection, immune memory was abrogated, whereas when antibiotics were delayed until day 2, 3, or 5 post-primary infection, bacterial burdens were significantly reduced in challenge infection, similar to that observed in the control untreated group (**Figure 3C**). These findings support that antigen persistence is necessary for the development of memory in UTI. Additionally, they raise the important question as to whether a previous UTI is actually a risk factor for recurrent infection (Hooton et al., 1996), or rather, antibiotic treatment contributes to an individual's risk for recurrent UTI due to poor immune memory.

Figure 3. Antigen persistence is necessary for immune memory to UTI. 6 to 8-week-old female C57BL/6 mice were infected with (**A-B**) 10^7 (solid line), 10^5 (dashed line), or 10^3 CFU (dotted line) of UPEC isolate UTI89-GFP-amp^R or UTI89-RFP-kan^R and challenged with 10^7 CFU of an isogenic strain after resolution, or (**C**) infected with 10^7 CFU UPEC isolate UTI89-GFP-amp^R or UTI89-RFP-kan^R, treated with trimethoprim/sulfamethoxazole 1, 2, 3, or 5 days PI, and challenged with 10^7 CFU of an isogenic strain after resolution. (**A**) Resolution after primary infection was monitored twice per week by assessing the presence of UPEC in the urine. The graph depicts the number of infected mice over time and an increase in the number of infected mice in the course of the experiment is due to a spontaneous recurrent UTI. (**B-C**) Mice were sacrificed 24 hours post-primary (1°, filled circles) or post-challenge infection (2°, open circles) for CFU assessment. The graphs show bladder bacterial burdens following (**B**) primary infection with the indicated inocula and challenge infection with 10^7 CFU UPEC and (**C**) primary infection and challenge infection in mice treated with trimethoprim/sulfamethoxazole at the indicated days post-primary UTI. The control (Ctrl) group did not receive antibiotics. **A** depicts representative data from 1 experiment of 2 (n=7 mice/group). Data in **B** are pooled from 2 experiments, n=5 to 7 mice/group in each experiment. Data in **C** are pooled from 2-4 experiments, n=6 to 7 mice/group in each experiment. Each circle is a mouse, lines are medians, dotted red lines depict the limit of detection of the assay, 20 CFU/bladder. In **B,** significance was determined by Kruskal-Wallis test comparing bacterial burden 24 hours post 1° to 2° within each inoculum group and in **C**, each 2° group was compared to the 1° CFU, with Dunn's post hoc test to correct for multiple comparisons and adjusted p-values for each comparison are presented, p-values <0.05 are in red.

Limiting antigen persistence does not impact Th cell polarization

When antigen load was reduced, by decreasing the inoculum or giving antibiotics on day 1 PI, protection against a second infection was lost. Previously, we reported that depletion of tissue resident macrophages, prior to challenge infection, results in increased infiltration of Th1 T cells and improved bacterial clearance in the bladder (Lacerda Mariano et al., 2020). To test whether the loss of immune memory observed in the two infection scenarios was due to a change in the proportions of polarized Th cell populations, we either infected mice with 10^7 or 10^3 CFU of UTI89, or we infected a cohort of mice with 10^7 CFU of UTI89 and treated half of these mice with antibiotics at day 1 PI. 28 days later, we challenged all resolved mice with 10^7 CFU of UTI89 and assessed CD4$^+$ T cell polarization 24 hours post-challenge. In mice infected with 10^3 CFU for their first UTI, we observed a significant reduction in the total number of Th1, Th2, and Th17 T cells, as well as decreases in IFNγ$^+$ Th1 T cells, IL-17$^+$ Th17 T cells, and Tregs compared to the control group (**Figure 4A**). Disproving our hypothesis, however, all Th cell subsets and Tregs decreased similarly, so proportion of each subset and therefore, the global Th cell polarization was the same between mice infected with 10^7 CFU or 10^3 CFU (**Figure 4B**). Further supporting our conclusion that the loss of immune memory was not due to a change in T cell polarization, we observed no differences in the numbers of Th1, Th2, or Th17 T cell subsets; IFNγ$^+$ Th1 T cells or IL-17$^+$ Th17 T cells; or Tregs between groups of mice that were antibiotic-treated or not (**Figure 4C**). As the number of T cells in each subset did not change in antibiotic-treated mice, the proportions of each Th cell subset were unchanged compared to the control condition (**Figure 4D**). Supporting that memory was not lost due changes in the systemic T cell compartment, the draining lymph nodes also had no differences among any of the parameters measured in the experimental groups compared to the control conditions (**Supplementary Figure 3**). Notably, although both experimental groups had abrogated immune memory (**Figure 3**), in one scenario, T cell infiltration was globally reduced (**Figure 4A**), whereas in the second, T cell infiltration was comparable to the control group (**Figure 4C**). That there was no change in proportions of polarized Th cells or, in the case of antibiotic treatment, no change in the number of Th cell subsets present, supports that the polarization and quantity of infiltrating Th cells in response to UTI are not major determinants of memory development in UTI.

Figure 4. Th T cell polarization is unchanged when antigen persistence is limited. (A-B) Six-

week-old female C57BL/6 mice were intravesically infected with either 10^7 or 10^3 CFU of UTI89,

and at day 28, resolved mice were challenged with 10^7 CFU of UTI89 and bladders analyzed by

flow cytometry 24 hours post-challenge infection. Graphs depict (**A**) the total number of the

specified Th cell populations per bladder and (**B**) the frequency among CD4 T cells of the Th cell

subsets in **A**. (**C-D**) Six-week-old female C57BL/6 mice were infected with 10^7 CFU of UTI89.

One cohort of mice was treated 24 hours PI with antibiotics and all resolved mice were challenged

with 10^7 CFU of UTI89 28 days later. Bladders were analyzed by flow cytometry 24 hours post-

challenge infection. Graphs show (**C**) the total number of the specified Th cell populations per

bladder and (**D**) the frequency of each Th cell subset among CD4$^+$ T cells. Data are pooled from

2 experiments, n=4 to 6 mice/group in each experiment. Each circle represents a mouse and lines

are medians. In **A** and **C,** nonparametric Mann-Whitney tests comparing cell numbers between

each condition were performed and all *p*-values were corrected for multiple comparisons among

the same experiment using the false discovery rate (FDR) method. In **B** and **D,** significance was

determined by Kruskal-Wallis tests comparing the frequency of indicated T cell subsets for each

condition with Dunn's post hoc test to correct for multiple comparisons were performed. *p*-values

<0.05 are in red.

 Tissue resident memory T cells are necessary to mediate immune memory to recurrent UTI

We were surprised that Th cell polarization was not changed when immune memory to a recurrent UTI

was abrogated following approaches that limited antigen persistence. To understand this unexpected

result, we considered that immunological memory can be mediated through B cells, circulating central

(T_{CM}) and (T_{EM}) effector memory T cells, or a more recently described subset of long-lived stationary T

cells called tissue resident memory T cells (T_{RM}) (Akkaya et al., 2020; Mueller et al., 2013). Our prior

work, as well as our findings in **Figure 1**, ruled out a role for B cells, and demonstrated T cells are

necessary for an adaptive response (Mora-Bau et al., 2015). This T cell-mediated memory response to

UTI is rapid as a 2-log reduction in bacterial burden is observed in the bladder at 24 hours post challenge

infection (Lacerda Mariano et al., 2020; Mora-Bau et al., 2015; Zychlinsky Scharff et al., 2019). Given

the speed of this response, we hypothesized that T_{RM} cells mediate memory to recurrent UTI.

To test this hypothesis, we infected mice with 10^7 or 10^3 CFU of UTI89, or we treated mice infected

with 10^7 CFU of UTI89 with antibiotics 1 day PI. 28 days post-primary infection, we challenged all

resolved mice with 10^7 CFU of UTI89 and analyzed bladder resident T cell populations by flow

cytometry. Defining these cells as $CD3^+$, $CD4^+$, $CD8^+$, $CD44^{high}$, $CD62L^-$, and expressing CD69 and/or

CD103, we observed a significant reduction in the number of $CD4^+$ and $CD8^+$ T_{RM} cells in mice infected

with 10^3 CFU compared to mice infected with 10^7 CFU (**Figure 5A**) or in mice treated with antibiotics

at 1 day PI compared to control-treated mice (**Figure 5B**). Supporting that it is specifically bladder-

associated T_{RM} cells that mediate memory, there were no differences in the number of T cells with a T_{RM}

phenotype in the draining lymph nodes in any of the tested conditions (**Supplementary Figure 4**). Thus,

reducing antigen load or persistence negatively impacted only local bladder T_{RM} cell populations,

correlating with abrogation of protection following a recurrent UTI.

One defining feature of T_{RM} cells is that their tissue resident status renders them inaccessible to

intravascular staining strategies. Therefore, to provide supporting evidence that the

$CD3^+CD4^+CD44^{high}CD62L^-$, or $CD3^+CD8^+CD44^{high}CD62L^-$ cells expressing CD69 and/or CD103 were

tissue resident, we injected fluorescently-labelled anti-CD3 antibody intravenously (IV) after resolution

of a primary infection and collected blood and bladders 3 minutes after injection. Circulating $CD4^+$ and

$CD8^+$ T cells were nearly universally labeled by the injected antibody (**Figure 5C**). By contrast, bladder-

associated $CD4^+$ and $CD8^+$ T cells, including $CD4^+$ and $CD8^+$ T_{RM} cells, were almost entirely negative

for the injected antibody IV, but were identified by using an anti-CD3 antibody with a different

fluorophore after bladder digestion for flow cytometry (**Figure 5D**). As a complementary approach to

test whether these cells were tissue resident while avoiding the caveats of tissue disruption, we used

confocal microscopy following IV-labeling of T cells. Again, blood $CD4^+$ and $CD8^+$ T were uniformly

positive for an IV-injected fluorophore labeled anti-CD3 antibody (**Figure 5E**). However, using an anti-

hamster secondary antibody to amplify the IV-injected primary anti-CD3 antibody signal in the bladder,

we did not detect any T cells labeled by IV antibody administration (**Figure 5F**). The absence of signal

with anti-hamster labeling was not due to a lack of T cells as when we used an anti-CD3 antibody to

immunostain bladder tissue sections, we observed an abundance of T cells in the urothelium and lamina

propria of post-infected tissue (**Figure 5F**). Altogether, these data support that $CD3^+CD4^+$, $CD8^+$,

$CD44^{high}$, $CD62L^-$ cells expressing CD69 and/or CD103 are T_{RM} cells.

Figure 5: Tissue resident memory T cell accumulation requires antigen persistence. Graphs
depict the total number of the specified T_{RM} cell populations per bladder, identified by flow
cytometry, after (**A**) six-week-old female C57BL/6 mice were instilled intravesically with 10^7 or
10^3 CFU of UTI89, and following resolution, challenged with 10^7 CFU of UTI89 and sacrificed
24 hours post challenge infection or (**B**) mice were infected with10^7 CFU of UTI89 and half of
the animals were treated with antibiotics 24 hours PI. All resolved mice were challenged with 10^7
CFU of UTI89 28 days later and sacrificed 24 hours post-challenge infection. (**C-F**) Six-week-
old female C57BL/6 mice were instilled intravesically with 10^7 CFU of UTI89 and following
resolution 28 days later, received 3μg of anti-CD3 antibody intravenously. Blood and bladders
were collected and analyzed by flow cytometry or confocal microscopy. (**C**) Representative dot
plots and graph show the proportion of IV-injected Pacific blue-labeled anti-CD3 antibody
staining in circulating T cells. (**D**) Representative dot plots and graph show the proportion of IV-
injected Pacific blue-labeled anti-CD3 antibody staining and post-tissue digestion PE-CF594 anti-
CD3 antibody staining in bladder-associated T cells. (**E**) Graph shows the proportion of IV-
injected PE-labeled hamster anti-CD3 antibody staining in circulating T cells. (**F**) Representative
confocal images of 3 different bladders from C57BL6/J mice at 20x and 40x magnification.
Merged images and single channels with the target of interest are shown. DAPI, 4',6'-diamidino-
2-phenymlindole. Data in **A** and **B** are pooled from 2 experiments, n=3 to 6 mice/group in each
experiment. Data in (**C-D**) are pooled from 2 experiments n=5 mice/group in each experiment.
(**E** and **F**) are from 1 experiment with 5 mice. Each circle is a mouse, lines are medians.
Significance was determined in **A** and **B** by nonparametric Mann-Whitney tests comparing cell
numbers between each condition and calculated *p*-values were corrected for multiple comparisons
using the false discovery rate (FDR) method. *p*-values <0.05 are in red.

Tissue resident memory T cells are sufficient to mediate immune memory to recurrent UTI
To formally demonstrate that T_{RM} cells mediate protection against a recurrent UTI, we hypothesized
that their elimination should abrogate immune memory. However, there are currently no specific tools
that target T_{RM} cells without also impacting circulating T cell populations. As an alternative approach,
we chose to eliminate all T cells subsets except T_{RM} cells. Similar to their inaccessibility to intravascular
labeling, T_{RM} cells are protected from depleting antibodies when administered using certain doses or
schedules (Davies et al., 2017; Galvez-Cancino et al., 2018; Gebhardt et al., 2011; Turner et al., 2018).
Thus, we created a scenario in which the only memory T cells present during a challenge infection would
be T_{RM} cells in the bladder. Mice were infected with UPEC and a cohort of animals was sacrificed 24
hours PI infection. After resolution of infection, at day 28, mice received anti-CD4 and CD8 depleting
antibodies or their respective isotype controls. Mice were challenged 24 hours later with 10^7 CFU of
UPEC (**Figure 6A**). In T cell-depleted but uninfected mice, we measured T cell depletion in the blood,

draining lymph nodes, and bladder. $CD4^+$ and $CD8^+$ T cells were significantly reduced to nearly zero in

the circulation of antibody-treated mice (depicted in red) compared to control isotype-treated animals

(**Figure 6B**). In bladder-draining lymph nodes of antibody-depleted mice, $CD4^+$ and $CD8^+$ T cells with

a T_{RM} cell phenotype (**Figure 6C**), as well as $CD4^+$ and $CD8^+$ T_{CM} and T_{EM} cells (**Figure 6D**) were

significantly reduced to nearly zero, indicating depleting antibodies accessed the lymph nodes and any

memory T cells resident in the lymph nodes were not available to infiltrate the bladder upon challenge

infection to mediate protection. Importantly, analysis of the T_{RM} populations in the bladder revealed that

T_{RM} cells were protected from antibody depletion and there was no reduction of $CD4^+$ or $CD8^+$ T_{RM} cells

in the bladders of mice that received CD4 and CD8 depleting antibodies compared to mice that were

treated with isotype controls (**Figure 6E**). Thus, only bladder T_{RM} cells were present in the mice when

we challenged them with a second UPEC infection.

Following challenge UTI, isotype-treated mice had bacterial burdens that were significantly reduced

compared to bacterial CFU measured after a primary infection, as expected. Remarkably, anti-CD4 and

CD8 antibody-treated mice were also protected to the same extent from challenge infection (depicted

by red open circles), with significantly reduced bacterial CFU in the bladder despite a near global

circulating T cell depletion, demonstrating that immune memory was intact in these animals (**Figure**

6F). As an alternative approach, we used FTY720, an inhibitor of the sphingosine-1-phosphate receptor

pathway (Chiba et al., 1998), which prevents lymphocyte migration from the lymph nodes. Mice were

infected with UPEC, and after resolution of infection at day 28, mice received 3 doses of either control

diluent or FTY720 prior to challenge infection (**Figure 6A**). Control or FTY720-treated mice (depicted

by blue open circles) were equally protected 24 hours post-challenge with no significant difference in

their bacterial burdens after a second UTI (**Figure 6G**). Together, these results support that T_{RM} cells,

specifically in the bladder, and not peripheral memory T cells, mediate protection against recurrent UTI.

Figure 6: Tissue resident memory T cells mediate protection against recurrent UTI (A) Schematic of experiments in which six-week-old female C57BL/6 mice were infected with 10^7 CFU of UTI89 and at 28 days post-infection treated with either (**B-F**) anti-CD4 and anti-CD8 antibodies or their respective isotype controls or (**G**) FTY720 or a control diluent. Resolved mice were sacrificed (**B-E**) to assess depletion or (**F-G**) challenged with 10^7 CFU of UTI89 and

analyzed 24 hours post-challenge infection. Graphs depict the total number of specified cell
populations in (**B**) blood, (**C** and **D**) bladder-draining lymph nodes, and (**E**) bladder in resolved
but unchallenged mice. Graphs show bacterial burden in (**F**) control and antibody-depleted mice
and (**G**) control and FTY720-treated mice. Data in (**B-F**) are pooled from 3 experiments, n=3 to
6 mice/group in each experiment. Data in **G** are pooled from 4 experiments, n=3 to 5 mice/group
in each experiment. Each circle is a mouse, lines are medians. In **F** and **G**, dotted red lines depict
the limit of detection of the assay. Significance in **B-E** was determined by nonparametric Mann-
Whitney tests comparing cell numbers between each condition and all calculated p-values were
corrected for multiple comparisons using the false discovery rate (FDR) method, in **F** by Kruskal-
Wallis test comparing bladder bacterial burden 24 hours post primary infection (filled circles) to
bacterial CFU 24 hours post challenge infection (open circles) in isotype (black open circles) or
depleting antibody-treated mice (red open circles), with Dunn's post hoc test to correct for
multiple comparisons, and in **G** by nonparametric Mann-Whitney test comparing bacterial burden
24 hours post challenge between diluent (black open circles) and FTY720-treated mice (blue open
circles). p-values <0.05 are in red.

468
469
Discussion

Overwhelmingly, studies of immunity to UTI focus on innate immunity, and despite how common
recurrent UTI is, we have a poor understanding of how memory develops after an acute infection
(Lacerda Mariano and Ingersoll, 2020). Supporting that humoral responses are important, UPEC-
specific antibody production is observed in children with pyelonephritis (Jodal et al., 1974) and, many
years ago, antibodies were proposed as a biomarker for human pyelonephritis (Percival et al., 1964).
Thus, it was surprising to rule out a role for B cells in our model of recurrent UTI. Notably, however, in
a similar mouse UTI model, antibodies to UPEC arise only when bacteria colonized both the bladder
and kidneys (Chan et al., 2013). We used C57BL/6 mice in our studies, and given that this background
has a very low incidence of vesicoureteral reflux and kidney colonization (Murawski et al., 2010), and
that antibodies are present specifically in cases of human pyelonephritis, our findings support that B
cells are likely dispensable for memory to cystitis. This has consequences for future vaccines or other
immunomodulatory strategies for recurrent UTI. Indeed, it may be that targeting the bladder to promote
antibody production will not provide efficacious protection against cystitis, and may explain why,
despite more than 25 years of research, vaccination strategies for UTI have so far not been successful.

Immune memory to UTI develops, but it is not sterilizing (Mora-Bau et al., 2015). Inadequate or
inefficient memory responses likely play a role in the frequency of recurrent UTI. Recently, a study
proposed that a type 2 immune biased response during UTI in mice is the reason for why memory is not
sterilizing (Wu et al., 2020). While a Th2 biased response would be expected to favor tissue repair over

strong cellular immunity, our data show that immune memory can be ablated without changing the
proportion, or even the number, of Th1, Th2, or Th17 subsets and that there is no prevailing T helper
cell bias in the first 7 days of a primary or recurrent infection. In addition, female and male mice develop
comparable protective memory responses after an initial UTI, although male mice do not express type
2 cytokines (Zychlinsky Scharff et al., 2019). In trying to understand this paradox, we found through
immunophenotyping that the immune response is not specifically biased towards any single Th subset,
but rather that a mixed Th1, Th2, and Th17 response develops, with a robust infiltration of regulatory T
cells. While this mixed response was not previously appreciated, it is not entirely surprising, as in the
first 24 hours of infection, UPEC induces extensive tissue damage, including robust inflammation and
urothelial cell exfoliation, and will be both intracellular and extracellular (Anderson et al., 2003;
Ingersoll et al., 2008; Justice et al., 2004; Mora-Bau et al., 2015; Mulvey et al., 1998; Zychlinsky Scharff
et al., 2019), necessitating a Th2 response to mediate repair, regulatory T cells to control effector cell
activation, Th1-polarized T cells to eliminate intracellular pathogens and a Th17 response to target
mucosal and extracellular pathogens concomitantly (Gieseck et al., 2018; Kaiko et al., 2008; Shevyrev
and Tereshchenko, 2019; Vignali et al., 2008; Zhu et al., 2010). Indeed, this mixed response reflects the
complexity of the host-pathogen interaction and immune response to UTI, as well as the fine equilibrium
that must be established between resolution of infection and maintenance of bladder integrity.
507
Adding to our understanding of memory to UTI, the quantity and persistence of bacteria is a key
parameter in the development of an adaptive immune response, and a minimum exposure of greater than
24 hours is needed for the development of adaptive immunity after a primary UTI. While additional
studies are necessary to determine the mechanisms underlying this result, alarmingly, it may be that the
rapid availability of antibiotics to individuals with a UTI actually places them at risk for recurrent
infections by impeding the development of immune memory. Determining whether delaying antibiotic
treatment improves the memory response is feasible and may ultimately decrease the incidence of
recurrence. Treatments that ameliorate symptoms but do not eliminate bacteria themselves, followed by
delayed antibiotic therapy in those not at risk for pyelonephritis, may be a better therapeutic approach
for lasting memory. Supporting this idea, while nonsteroidal anti-inflammatory drugs (NSAIDs)
administered in place of antibiotics are not superior to antibiotic treatment (Gagyor et al., 2015;
Kronenberg et al., 2017; Vik et al., 2018), a meta-analysis of 962 patients comparing antibiotic treatment
to placebo for uncomplicated cystitis did not find a difference in the development of pyelonephritis
(Falagas et al., 2009). This suggests that delaying antibiotics may not put individuals at greater risk for
progression to pyelonephritis, and may support the development of a better memory response. Further
supporting this alternative treatment approach, NSAID-treated individuals had a significant reduction
in the number of recurrent UTI in their first month post-treatment compared to those treated with
antibiotics (Bleidorn et al., 2016). Thus, it would be of great value to test whether briefly delaying

antibiotic treatment promotes a better memory response and a reduction in recurrent UTI in humans,
which would, in turn, reduce reliance on antibiotics for this infection.

We previously reported that macrophage depletion before a primary or a challenge infection improves
bacterial clearance during the challenge infection (Lacerda Mariano et al., 2020; Mora-Bau et al., 2015).
This phenotype is T cell-dependent, and macrophage depletion specifically before a second UTI leads
to an increase in type 1-biased effector immune cell infiltration (Lacerda Mariano et al., 2020; Mora-
Bau et al., 2015). Thus, when we observed that memory was negatively impacted by decreasing the
inoculum or administering antibiotics, we expected a change in T cell polarization, potentially towards
a Th2 bias. We were surprised to observe changes only in the numbers of T_{RM} cell populations, and not
in the proportions of Th subsets, which suggests to us that resident macrophages directly influence
bladder resident T cell development or maintenance, potentially as a mechanism to maintain bladder
integrity. Bladder resident macrophages likely modulate the microenvironment, which shapes the local
T_{RM} cell response. In other mucosal tissues, lung resident macrophages limit the development of
influenza-specific CD8$^+$ T_{RM} cells (Goplen et al., 2019). In *Yersinia pseudotuberculosis* infection, type
I IFN and IL-12 in the intestinal microenvironment is key for differentiation and persistence of T_{RM}
cells, and depletion of CCR2$^+$ IL-12-producing cells, including macrophages or monocytes, impedes
T_{RM} cell differentiation (Bergsbaken et al., 2017). Deciphering the mechanisms by which macrophages
control local T cell responses in the bladder will be key to design immunomodulatory approaches.

The discovery of T_{RM} cells localized in non-lymphoid tissues, which are anatomically well positioned
to detect and respond to a second infection faster than T_{CM} or T_{EM} cells, has driven development of
vaccine strategies to promote their differentiation. UTI needs new therapeutic approaches to combat the
rapid global spread of multidrug resistant bacteria. Vaccination strategies promoting the differentiation
of bladder T_{RM} cells would likely be of great benefit for patients suffering from recurrent UTI and would
have a positive economic impact by reducing the cost of medical care. The failure of previous vaccine
strategies for UTI may be because the correct vaccination strategy to promote the differentiation of T_{RM}
cells has not been identified. For example, a BCG mucosal vaccination better protects mice against
tuberculosis compared to a subcutaneous vaccination, and adoptive transfer of T_{RM} cells from vaccinated
mice protects naïve animals from disease (Perdomo et al., 2016). Similarly, a mucosal vaccine approach
provides superior protection against coronavirus infections compared to a subcutaneous route (Zhao et
al., 2016). In addition to the route of vaccination, the type of vaccine used is also a key parameter.
Indeed, an attenuated influenza vaccine promotes T_{RM} cell generation in the lung, inducing long term
protection, whereas an inactivated vaccine does not, despite intranasal delivery (Zens et al., 2016).
Intranasal immunization with *Chlamydia trachomatis* promotes uterine mucosa protection and oral or
gastrointestinal vaccination with *Chlamydia muridarum* protects the genital tract from infection (Stary

et al., 2015; Wang et al., 2018). These observations will likely inform vaccine strategies for UTI and
suggest that the bladder mucosa does not need to be directly targeted.

In sum, we demonstrated that bladder T_{RM} cells mediate memory to recurrent UTI, and do so in the
absence of circulating or lymph node-resident T cells. Thus, we uncovered a specific, targetable
population for development of new therapeutic approaches for this common infection plagued by
antibiotic resistance. These findings greatly improve our understanding of adaptive immunity to UTI
and provide valuable information for the design and development of much needed new non-antibiotic-
based therapies.

Materials and methods
Study design
This study was conducted using a preclinical mouse model in controlled laboratory experiments to
investigate adaptive immunity UTI. Our objective was to determine how an adaptive immune response
develops after a UTI. Mice were assigned to groups by random partition into cages. In all experiments,
a minimum of 2 and a maximum of 7 mice made up an experimental group and all experiments were
repeated 2 to 5 times, with the exception of confocal imaging, in which five mice were assessed in total.
Data from all repetitions were pooled before any statistical analysis to reach a predetermined *n* of at
least 10 mice per group. As determined *a priori*, all animals with abnormal kidneys (atrophied, enlarged,
and/or white in color) at the time of sacrifice were excluded from all analyses, as we have observed that
abnormal kidneys correlate with the inability to resolve infection. Animals with a recurrent or
unresolved UTI, defined as having persistent bacteria in the urine up to 1 day prior to challenge infection,
were also excluded. End points were determined before the start of experiments and researchers were
not blinded to experimental groups.

Ethics Statement
Animal experiments were conducted in accordance with approval of protocol number 2012-0024 and
2016-0010 by the *Comité d'éthique en expérimentation animale Paris Centre et Sud* and the *Comités*
d'Ethique pour l'Expérimentation Animale Institut Pasteur (the ethics committee for animal
experimentation), and APAFIS #34290 by SC3 - CEEA34 – Université de Paris Cité, at Institut Cochin,
in application of the European Directive 2010/63 EU. In all experiments, mice were anesthetized by
intraperitoneal injection of 100 mg/kg ketamine and 5 mg/kg xylazine and sacrificed by carbon dioxide
inhalation or cervical dislocation after isoflurane inhalation.

Mice
Female mice between the ages of 6 and 8 weeks were used in this study. Female C57BL/6J mice were

obtained from Charles River Laboratories France. μMT$^{-/-}$ mice were bred in house at Institut Pasteur,

Paris, and were a kind gift from Claude Leclerc.

UTI and determination of bacterial burden

Female mice were anesthetized as described above, catheterized transurethrally, and infected with

$1×10^7$, $1×10^5$, or $1×10^3$ colony forming units (CFU) of UTI89-GFP-amp^R or UTI89-RFP-kan^R in 50 μL

PBS as previously described (Mora-Bau et al., 2015). UTI89-GFP-amp^R and UTI89-RFP-kan^R infect

with equal efficiency and were used interchangeably (Mora-Bau et al., 2015). For flow cytometry, only

the nonfluorescent parental UTI89 strain was used. Infection was monitored by detection of bacterial

growth from urine samples collected twice per week. 2 μL of urine were diluted directly into 8 μL PBS

spotted on agar plates containing antibiotics as appropriate (kanamycin (50 μg/ml) or ampicillin (100

μg/ml)). The presence of any bacterial growth was counted as positive for infection. The limit of

detection (LOD) for this assay is 500 bacteria per mL of urine. Mice were sacrificed at indicated time

points, bladders homogenized in sterile PBS, serially diluted, and plated on LB agar plates with

antibiotics, as appropriate, to determine CFU. The LOD for CFU in the bladder is 20 CFU per bladder

and is indicated by a dotted line in graphs. All sterile organs are reported at the LOD.

Flow cytometry of bladder tissue, draining lymph nodes and blood

Samples were acquired on a BD LSRFortessa using DIVA software (v8.0.1), and data were analyzed

by FlowJo (Treestar) software. Bladder and blood analyses were performed as described previously

(Lacerda Mariano et al., 2020; Mora-Bau et al., 2015; Zychlinsky Scharff et al., 2017). Briefly, bladders

were dissected, cut into small pieces, and digested using Liberase (0.34 U/ml) in phosphate-buffered

saline (PBS) at 37°C for 1 hour with robust manual agitation every 15 min. Digestion was stopped by

adding FACS buffer (PBS supplemented with 2% fetal bovine serum and 0.2 μM EDTA). Single cell

suspensions were washed and resuspended in brilliant stain buffer (BD) with anti-mouse CD16/CD32

to block Fc receptors. Antibody mixes (**Table 1**) in brilliant stain buffer were added directly to the

samples after 10 minutes. Total cell counts were determined by addition of AccuCheck counting beads

to a known volume of sample after staining, just before cytometer acquisition. To determine cell

populations in the circulation, whole blood was incubated with BD PharmLyse and stained with

antibodies (**Table 1**). Total cell counts per μL of blood were determined by the addition of AccuCheck

counting beads to 10 μl of whole blood in 1-step BD Fix/Lyse Solution. Draining lymph nodes were

disrupted with 27G needles and passed through 70 μm filters (Miltenyi). Single cell suspensions were

washed and resuspended in brilliant stain buffer with anti-mouse CD16/CD32 to block Fc receptors and

subsequently stained as above for bladders. Total cell counts were determined as they were for the

bladder. Memory T cell staining was performed at 37°C for 30 minutes to enhance CCR7 staining

(Berhanu et al., 2003).

For intracellular staining, single cell suspensions were resuspended in 1 mL of Golgi Stop protein transport inhibitor diluted 1:1500 in RPMI with 10% FBS, 1% sodium pyruvate, 1X HEPES, 1X nonessential amino acid, 1% penicillin-streptomycin, phorbol 12-myristate 13-acetate (50 ng/ml), and ionomycin (1 μg/ml), and incubated for 4 hours at 37°C. Samples were washed once with FACS buffer and Fc receptors blocked with anti-mouse CD16/CD32. Samples were stained with antibodies listed in **Table 1** against surface markers and fixed and permeabilized with 1X fixation and permeabilization buffer and incubated at 4°C for 40 to 50 min protected from light. After incubation, samples were washed two times with 1X permeabilization and wash buffer from the transcription factor buffer kit (BD Biosciences) and stained with antibodies against IFN-γ, IL-17, IL-4, and the transcription factors RORγT, Gata-3, T-bet, and FoxP3 (**Table 1**), diluted in 1X permeabilization and wash buffer at 4°C for 40 to 50 min protected from light. Finally, samples were washed two times with 1X permeabilization and wash buffer and resuspended in FACS buffer. Total cell counts were determined by addition of counting beads to a known volume of sample after staining, just before cytometer acquisition.

Table 1: Antibodies used for flow cytometry in this study

Antibodies	Clone	Source	Identifier
Anti-mouse NK1.1 APC	PK136	BD Biosciences	cat# 550627
Anti-mouse B220 AF700	RA3-6B2	BD Biosciences	cat# 557957
Anti-mouse I-A/I-E (MHCII) APC-efluor780	MS/114.15.2	Invitrogen	cat# 47-5321-82
Anti-mouse CD8 FITC	53-6.7	BD Biosciences	cat# 553031
Anti-mouse GR1 PerCP-Cy5.5	RB6-8C5	BD Biosciences	cat# 552093
Anti-mouse CD3 Pacific Blue	500A2	BD Biosciences	cat# 558214
Anti-mouse CD45 BV605	30-F11	BD Biosciences	cat# 563053
Anti-mouse CD115 PE	AFS98	Invitrogen	cat# 12-1152-82
Anti-mouse SiglecF PE-CF594	E50-2440	BD Biosciences	cat# 562757
Anti-mouse CD4 PE-Cy7	GK1.5	ebioscience	cat# 25-0041-82
Anti-mouse NK1.1 AF700	PK136	BD Biosciences	cat# 560515
Anti-mouse CD127 APC-eFluor780	A7R34	Invitrogen	cat# 47-1271-82
Anti-mouse CD69 FITC	H1.2F3	BD Biosciences	cat# 553236
Anti-mouse CD8a PerCP-Cy5.5	53-6.7	BD Biosciences	cat# 551162
Anti-mouse Ly6C BV421	HK1.4	Biolegend	cat# 128032

Anti-mouse CD103 BV510	M290	BD Biosciences	cat# 563087
Anti-mouse CD4 BV605	RM4-5	Biolegend	cat# 100548
Anti-mouse CCR7 BV650	4B12	BD Biosciences	cat# 564356
Anti-mouse KLRG1 BV711	2F1	BD Biosciences	cat# 564014
Anti-mouse CD45 BUV395	30-F11	BD Biosciences	cat# 564279
Anti-mouse CD44 BUV737	IM7	BD Biosciences	cat# 612799
Anti-mouse $\gamma\delta$ TCR PE	eBioGL3 (GL3)	Invitrogen	cat# 12-5711-82
Anti-mouse CD3 PE-CF594	145-2C11	Biolegend	cat# 100348
Anti-mouse CD62L PE-Cy7	MEL-14	Invitrogen	cat# 25-0621-82
Anti-mouse CD25 APC	PC61	BD Bioscience	cat# 557192
Anti-mouse CD11b AF700	M1/70	BD Biosciences	cat# 557960
Anti-mouse FoxP3 AF488	MF-14	Biolegend	cat# 126406
Anti-mouse RORγT BV421	Q31-378	BD Bioscience	cat# 562894
Anti-mouse $\gamma\delta$ TCR BV510	GL3	BD Biosciences	cat# 563218
Anti-mouse IL-17A BV650	TC11-18H10	BD Biosciences	cat# 564170
Anti-mouse NK1.1 BV711	PK136	BD Biosciences	cat# 740663
Anti-mouse IFNγ BV785	XMG1.2	Biolegend	cat# 505837
Anti-human/mouse T-bet PE	eBio4B10 (4B10)	Invitrogen	cat# 12-5825-80
Anti-mouse CD3 PE-CF594	145-2C11	Biolegend	cat# 100348
Anti-mouse Gata-3 PE-Cy7	L50-823	BD Biosciences	cat# 560405
Anti-mouse IL-4 BV711	11B11	BD Biosciences	cat# 564005

Immune cell depletion

To deplete T cells, 100 μg of CD4 and/or 100 μg of CD8 (clone GK1.5, clone YTS 169.4, respectively, Bio X Cell) depleting antibodies were injected intraperitoneally in 100 μl sterile PBS per mouse 3 days before primary infection, or 24 hours before challenge infection. 100 μg or 200 μg of the isotype control (clone LTF-2, Bio X Cell) were injected intraperitoneally in 100 μl of sterile PBS per mouse at the same time in control groups. Antibody injections given before the primary infection were repeated 7 days after the first injection, *i.e.*, day 4 PI.

Intravascular staining

Female mice were anesthetized as describe above, and 3μg of Pacific Blue anti-CD3 (clone 500A2, BD bioscience), or 3μg of PE anti-CD3 (clone 145-2C11, BD bioscience) were injected intravenously in the retroorbital sinus. 3 minutes later mice are sacrificed as described above, blood and bladder are collected and analyzed by flow cytometry or by confocal microscopy.

Histological and immunostaining for confocal microscopy

Whole bladders were fixed with 4% paraformaldehyde (PFA) in PBS for 1 hour and subsequently washed with PBS. Samples were then dehydrated in 30% sucrose in PBS for 24 hours. Samples were cut transversally and embedded in optimal cutting temperature compound, frozen, and sectioned at 30 μm. Sections were blocked for 1 hour with blocking buffer [3% bovine serum albumin (BSA) + 0.1% Triton X-100 + goat serum (1:20) in PBS] and washed three times. Immunostaining was performed using rat anti-mouse CD3 (1:100) in staining buffer (0.5% BSA + 0.1% Triton X-100 in PBS) overnight. Sections were washed and stained with phalloidin (1:350), goat anti-hamster (1:350) and goat anti-rat (1:1000) antibodies in staining buffer for 4 hours. Last, sections were washed and stained with 4′,6-diamidino-2-phenylindole. Confocal images were acquired on a Leica SP8 confocal microscope. Final image processing was done using Fiji (version 2.0.0-rc-69/1.52p).

FTY720 treatment

Female mice received by an intraperitoneal injection 50μl of aqueous solution, or FTY720 at 0,4mg/ml diluted in aqueous solution (0,02mg/mouse ≈ 1mg/kg) at 3 consecutive days (day -2, -1, and at the time of challenge infection).

Statistical analysis

Statistical analysis was performed in GraphPad Prism 9 (GraphPad, USA) for Mac OS X applying the nonparametric Mann-Whitney test for unpaired data in the case of two group comparisons. To correct for comparisons made within an entire analysis or experiment, calculated *p*-values were corrected for multiple testing with the false discovery rate (FDR) method to determine the FDR-adjusted *p*-value. In the case that more than two groups were being compared, Kruskal-Wallis tests were performed, with Dunn's post hoc test to correct for multiple comparisons. All adjusted *p*-values are shown in the figures.

Acknowledgments

We are thankful for insightful discussion with Dr. Elizabeth Wohlfort. Funding: LLM was part of the Pasteur-Paris University (PPU) International PhD Program, which received funding from the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant

695 agreement no. 665807 and from the Labex Milieu Intérieur (ANR-10-LABX-69-01). MAI was
696 supported by funding from the *Agence Nationale de la Recherché* (French National Research Agency)
697 ANR-17-CE17-0014 and ANR-19-CE15-0015 and *Chaires d'excellence de l'IdEx Université de Paris*.
698
699 **Author contributions**
700 Conceptualization: MR and MAI; Methodology: MR, LLM, TC, MAI; Investigation and data analysis:
701 MR, LLM, TC, MAI; Writing - Original Draft: MR and MAI; Writing - Review & Editing: MR, LLM,
702 TC, MAI; Funding Acquisition: MAI; Supervision: MAI.
703
704 **Competing interests**
705 The authors declare no competing interests.
706
707 **Data and materials availability**
708 Further information and requests for resources and reagents should be directed to and will be fulfilled
709 by Matthieu Rousseau (matthieu.rousseau@inserm.fr) and Molly A. Ingersoll
710 (molly.ingersoll@pasteur.fr).

References

Akkaya, M., Kwak, K., and Pierce, S.K. (2020). B cell memory: building two walls of protection against
pathogens. Nat Rev Immunol *20*, 229-238.

Alteri, C.J., Hagan, E.C., Sivick, K.E., Smith, S.N., and Mobley, H.L. (2009). Mucosal immunization
with iron receptor antigens protects against urinary tract infection. PLoS Pathog *5*, e1000586.

Anderson, G.G., Palermo, J.J., Schilling, J.D., Roth, R., Heuser, J., and Hultgren, S.J. (2003).
Intracellular bacterial biofilm-like pods in urinary tract infections. Science *301*, 105-107.

Bergsbaken, T., Bevan, M.J., and Fink, P.J. (2017). Local Inflammatory Cues Regulate Differentiation
and Persistence of CD8(+) Tissue-Resident Memory T Cells. Cell Rep *19*, 114-124.

Berhanu, D., Mortari, F., De Rosa, S.C., and Roederer, M. (2003). Optimized lymphocyte isolation
methods for analysis of chemokine receptor expression. J Immunol Methods *279*, 199-207.

Billips, B.K., Yaggie, R.E., Cashy, J.P., Schaeffer, A.J., and Klumpp, D.J. (2009). A live-attenuated
vaccine for the treatment of urinary tract infection by uropathogenic Escherichia coli. J Infect Dis *200*,
263-272.

Bleidorn, J., Hummers-Pradier, E., Schmiemann, G., Wiese, B., and Gagyor, I. (2016). Recurrent
urinary tract infections and complications after symptomatic versus antibiotic treatment: follow-up of a
randomised controlled trial. Ger Med Sci *14*, Doc01.

Chan, C.Y., St John, A.L., and Abraham, S.N. (2013). Mast cell interleukin-10 drives localized tolerance
in chronic bladder infection. Immunity *38*, 349-359.

Chiba, K., Yanagawa, Y., Masubuchi, Y., Kataoka, H., Kawaguchi, T., Ohtsuki, M., and Hoshino, Y.
(1998). FTY720, a novel immunosuppressant, induces sequestration of circulating mature lymphocytes
by acceleration of lymphocyte homing in rats. I. FTY720 selectively decreases the number of circulating
mature lymphocytes by acceleration of lymphocyte homing. J Immunol *160*, 5037-5044.

Chromek, M., and Brauner, A. (2007). Urinary Tract Infection. Why Do Some Children Get
Complications, While Others Dont? Current Pediatric Reviews *3*, 35-44.

Davies, B., Prier, J.E., Jones, C.M., Gebhardt, T., Carbone, F.R., and Mackay, L.K. (2017). Cutting
Edge: Tissue-Resident Memory T Cells Generated by Multiple Immunizations or Localized Deposition
Provide Enhanced Immunity. J Immunol *198*, 2233-2237.

De Nisco, N.J., Neugent, M., Mull, J., Chen, L., Kuprasertkul, A., de Souza Santos, M., Palmer, K.L.,
Zimmern, P., and Orth, K. (2019). Direct Detection of Tissue-Resident Bacteria and Chronic
Inflammation in the Bladder Wall of Postmenopausal Women with Recurrent Urinary Tract Infection.
J Mol Biol *431*, 4368-4379.

Dielubanza, E.J., and Schaeffer, A.J. (2011). Urinary tract infections in women. Med Clin North Am
95, 27-41.

Falagas, M.E., Kotsantis, I.K., Vouloumanou, E.K., and Rafailidis, P.I. (2009). Antibiotics versus
placebo in the treatment of women with uncomplicated cystitis: a meta-analysis of randomized
controlled trials. J Infect *58*, 91-102.

Falivene, J., Ghiglione, Y., Laufer, N., Socias, M.E., Holgado, M.P., Ruiz, M.J., Maeto, C., Figueroa,
M.I., Giavedoni, L.D., Cahn, P., *et al.* (2015). Th17 and Th17/Treg ratio at early HIV infection associate
with protective HIV-specific CD8(+) T-cell responses and disease progression. Sci Rep *5*, 11511.

Flores-Mireles, A.L., Walker, J.N., Caparon, M., and Hultgren, S.J. (2015). Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. Nat Rev Microbiol *13*, 269-284.

Foxman, B. (2002). Epidemiology of urinary tract infections: incidence, morbidity, and economic costs.
Am J Med *113 Suppl 1A*, 5S-13S.

Gagyor, I., Bleidorn, J., Kochen, M.M., Schmiemann, G., Wegscheider, K., and Hummers-Pradier, E.
(2015). Ibuprofen versus fosfomycin for uncomplicated urinary tract infection in women: randomised
controlled trial. BMJ *351*, h6544.

Galvez-Cancino, F., Lopez, E., Menares, E., Diaz, X., Flores, C., Caceres, P., Hidalgo, S., Chovar, O.,
Alcantara-Hernandez, M., Borgna, V*., et al.* (2018). Vaccination-induced skin-resident memory CD8(+)
T cells mediate strong protection against cutaneous melanoma. Oncoimmunology *7*, e1442163.

Gebhardt, T., Whitney, P.G., Zaid, A., Mackay, L.K., Brooks, A.G., Heath, W.R., Carbone, F.R., and
Mueller, S.N. (2011). Different patterns of peripheral migration by memory CD4+ and CD8+ T cells.
Nature *477*, 216-219.

Gieseck, R.L., 3rd, Wilson, M.S., and Wynn, T.A. (2018). Type 2 immunity in tissue repair and fibrosis.
Nat Rev Immunol *18*, 62-76.

Goplen, N.P., Huang, S., Zhu, B., Cheon, I.S., Son, Y.M., Wang, Z., Li, C., Dai, Q., Jiang, L., and Sun,
J. (2019). Tissue-Resident Macrophages Limit Pulmonary CD8 Resident Memory T Cell Establishment.
Front Immunol *10*, 2332.

Hoepelman, A.I., van Buren, M., van den Broek, J., and Borleffs, J.C. (1992). Bacteriuria in men
infected with HIV-1 is related to their immune status (CD4+ cell count). AIDS *6*, 179-184.

Hooton, T.M., Scholes, D., Hughes, J.P., Winter, C., Roberts, P.L., Stapleton, A.E., Stergachis, A., and
Stamm, W.E. (1996). A prospective study of risk factors for symptomatic urinary tract infection in
young women. N Engl J Med *335*, 468-474.

Ikaheimo, R., Siitonen, A., Heiskanen, T., Karkkainen, U., Kuosmanen, P., Lipponen, P., and Makela,
P.H. (1996). Recurrence of urinary tract infection in a primary care setting: analysis of a 1-year follow-
up of 179 women. Clin Infect Dis *22*, 91-99.

Ingersoll, M.A., and Albert, M.L. (2013). From infection to immunotherapy: host immune responses to
bacteria at the bladder mucosa. Mucosal immunology *6*, 1041-1053.

Ingersoll, M.A., Kline, K.A., Nielsen, H.V., and Hultgren, S.J. (2008). G-CSF induction early in
uropathogenic Escherichia coli infection of the urinary tract modulates host immunity. Cell Microbiol
10, 2568-2578.

Jodal, U., Ahlstedt, S., Carlsson, B., Hanson, L.A., Lindberg, U., and Sohl, A. (1974). Local antibodies
in childhood urinary tract infection: a preliminary study. Int Arch Allergy Appl Immunol *47*, 537-546.

Justice, S.S., Hung, C., Theriot, J.A., Fletcher, D.A., Anderson, G.G., Footer, M.J., and Hultgren, S.J.
(2004). Differentiation and developmental pathways of uropathogenic Escherichia coli in urinary tract
pathogenesis. Proc Natl Acad Sci U S A *101*, 1333-1338.

Kaiko, G.E., Horvat, J.C., Beagley, K.W., and Hansbro, P.M. (2008). Immunological decision-making:
how does the immune system decide to mount a helper T-cell response? Immunology *123*, 326-338.

Kitamura, D., Roes, J., Kuhn, R., and Rajewsky, K. (1991). A B cell-deficient mouse by targeted
disruption of the membrane exon of the immunoglobulin mu chain gene. Nature *350*, 423-426.

Klein, R.D., and Hultgren, S.J. (2020). Urinary tract infections: microbial pathogenesis, host-pathogen
interactions and new treatment strategies. Nat Rev Microbiol *18*, 211-226.

Kronenberg, A., Butikofer, L., Odutayo, A., Muhlemann, K., da Costa, B.R., Battaglia, M., Meli, D.N.,
Frey, P., Limacher, A., Reichenbach, S., and Juni, P. (2017). Symptomatic treatment of uncomplicated
lower urinary tract infections in the ambulatory setting: randomised, double blind trial. BMJ *359*, j4784.

Lacerda Mariano, L., and Ingersoll, M.A. (2020). The immune response to infection in the bladder. Nat
Rev Urol *17*, 439-458.

Lacerda Mariano, L., Rousseau, M., Varet, H., Legendre, R., Gentek, R., Saenz Coronilla, J., Bajenoff,
M., Gomez Perdiguero, E., and Ingersoll, M.A. (2020). Functionally distinct resident macrophage
subsets differentially shape responses to infection in the bladder. Sci Adv *6*.

Langermann, S., Palaszynski, S., Barnhart, M., Auguste, G., Pinkner, J.S., Burlein, J., Barren, P.,
Koenig, S., Leath, S., Jones, C.H., and Hultgren, S.J. (1997). Prevention of mucosal Escherichia coli
infection by FimH-adhesin-based systemic vaccination. Science *276*, 607-611.

Lipsky, B.A. (1989). Urinary tract infections in men. Epidemiology, pathophysiology, diagnosis, and
treatment. Ann Intern Med *110*, 138-150.

Litwin, M.S., Saigal, C.S., and editors. (2007). Urologic Diseases in America. Washington. P.H.S. U.S.
Department of Health and Human Services, National Institutes of Health, National Institute of Diabetes
and Digestive and Kidney Diseases, ed. (D.C, U.S. Government Publishing Office).

Mazor-Dray, E., Levy, A., Schlaeffer, F., and Sheiner, E. (2009). Maternal urinary tract infection: is it
independently associated with adverse pregnancy outcome? J Matern Fetal Neonatal Med *22*, 124-128.

Mora-Bau, G., Platt, A.M., van Rooijen, N., Randolph, G.J., Albert, M.L., and Ingersoll, M.A. (2015).
Macrophages Subvert Adaptive Immunity to Urinary Tract Infection. PLoS Pathog *11*, e1005044.

Mueller, S.N., Gebhardt, T., Carbone, F.R., and Heath, W.R. (2013). Memory T cell subsets, migration
patterns, and tissue residence. Annu Rev Immunol *31*, 137-161.

Mulvey, M.A., Lopez-Boado, Y.S., Wilson, C.L., Roth, R., Parks, W.C., Heuser, J., and Hultgren, S.J.
(1998). Induction and evasion of host defenses by type 1-piliated uropathogenic Escherichia coli.
Science *282*, 1494-1497.

Murawski, I.J., Maina, R.W., Malo, D., Guay-Woodford, L.M., Gros, P., Fujiwara, M., Morgan, K., and
Gupta, I.R. (2010). The C3H/HeJ inbred mouse is a model of vesico-ureteric reflux with a susceptibility
locus on chromosome 12. Kidney Int *78*, 269-278.

Mysorekar, I.U., and Hultgren, S.J. (2006). Mechanisms of uropathogenic Escherichia coli persistence
and eradication from the urinary tract. Proc Natl Acad Sci U S A *103*, 14170-14175.

Percival, A., Birumfitt, W., and Delouvois, J. (1964). Serum-Antibody Levels as an Indication of
Clinically Inapparent Pyelonephritis. Lancet *2*, 1027-1033.

Perdomo, C., Zedler, U., Kuhl, A.A., Lozza, L., Saikali, P., Sander, L.E., Vogelzang, A., Kaufmann,
S.H., and Kupz, A. (2016). Mucosal BCG Vaccination Induces Protective Lung-Resident Memory T
Cell Populations against Tuberculosis. mBio *7*.

Ruben, F.L., Dearwater, S.R., Norden, C.W., Kuller, L.H., Gartner, K., Shalley, A., Warshafsky, G.,
Kelsey, S.F., O'Donnell, C., Means, E., and et al. (1995). Clinical infections in the noninstitutionalized
geriatric age group: methods utilized and incidence of infections. The Pittsburgh Good Health Study.
Am J Epidemiol *141*, 145-157.

Shevyrev, D., and Tereshchenko, V. (2019). Treg Heterogeneity, Function, and Homeostasis. Front
Immunol *10*, 3100.

Sivick, K.E., and Mobley, H.L. (2009). Waging War against Uropathogenic Escherichia coli: Winning
Back the Urinary Tract. Infect Immun.

Spaulding, C.N., Klein, R.D., Ruer, S., Kau, A.L., Schreiber, H.L., Cusumano, Z.T., Dodson, K.W.,
Pinkner, J.S., Fremont, D.H., Janetka, J.W.*, et al.* (2017). Selective depletion of uropathogenic E. coli
from the gut by a FimH antagonist. Nature *546*, 528-532.

Stary, G., Olive, A., Radovic-Moreno, A.F., Gondek, D., Alvarez, D., Basto, P.A., Perro, M., Vrbanac,
V.D., Tager, A.M., Shi, J.*, et al.* (2015). VACCINES. A mucosal vaccine against Chlamydia trachomatis
generates two waves of protective memory T cells. Science *348*, aaa8205.

Svanborg-Eden, C., and Svennerholm, A.M. (1978). Secretory immunoglobulin A and G antibodies
prevent adhesion of Escherichia coli to human urinary tract epithelial cells. Infect Immun *22*, 790-797.

Thumbikat, P., Waltenbaugh, C., Schaeffer, A.J., and Klumpp, D.J. (2006). Antigen-specific responses
accelerate bacterial clearance in the bladder. J Immunol *176*, 3080-3086.

Turner, D.L., Goldklang, M., Cvetkovski, F., Paik, D., Trischler, J., Barahona, J., Cao, M., Dave, R.,
Tanna, N., D'Armiento, J.M., and Farber, D.L. (2018). Biased Generation and In Situ Activation of Lung
Tissue-Resident Memory CD4 T Cells in the Pathogenesis of Allergic Asthma. J Immunol *200*, 1561-
1569.

Uehling, D.T., Hopkins, W.J., Balish, E., Xing, Y., and Heisey, D.M. (1997). Vaginal mucosal
immunization for recurrent urinary tract infection: phase II clinical trial. J Urol *157*, 2049-2052.

Uehling, D.T., Hopkins, W.J., Beierle, L.M., Kryger, J.V., and Heisey, D.M. (2001). Vaginal mucosal
immunization for recurrent urinary tract infection: extended phase II clinical trial. J Infect Dis *183 Suppl*
1, S81-83.

Uehling, D.T., Hopkins, W.J., Dahmer, L.A., and Balish, E. (1994). Phase I clinical trial of vaginal
mucosal immunization for recurrent urinary tract infection. J Urol *152*, 2308-2311.

Uehling, D.T., James, L.J., Hopkins, W.J., and Balish, E. (1991). Immunization against urinary tract
infection with a multi-valent vaginal vaccine. J Urol *146*, 223-226.

Vignali, D.A., Collison, L.W., and Workman, C.J. (2008). How regulatory T cells work. Nat Rev
Immunol *8*, 523-532.

Vik, I., Bollestad, M., Grude, N., Baerheim, A., Damsgaard, E., Neumark, T., Bjerrum, L., Cordoba, G.,
Olsen, I.C., and Lindbaek, M. (2018). Ibuprofen versus pivmecillinam for uncomplicated urinary tract
infection in women-A double-blind, randomized non-inferiority trial. PLoS Med *15*, e1002569.

Wang, L., Zhu, C., Zhang, T., Tian, Q., Zhang, N., Morrison, S., Morrison, R., Xue, M., and Zhong, G.
(2018). Nonpathogenic Colonization with Chlamydia in the Gastrointestinal Tract as Oral Vaccination
for Inducing Transmucosal Protection. Infect Immun *86*.

Wu, J., Hayes, B.W., Phoenix, C., Macias, G.S., Miao, Y., Choi, H.W., Hughes, F.M., Jr., Todd Purves,
J., Lee Reinhardt, R., and Abraham, S.N. (2020). A highly polarized TH2 bladder response to infection
promotes epithelial repair at the expense of preventing new infections. Nat Immunol *21*, 671-683.

Xiao, B.G., and Link, H. (1997). Mucosal tolerance: a two-edged sword to prevent and treat autoimmune
diseases. Clin Immunol Immunopathol *85*, 119-128.

Yordanova, I.A., Cortes, A., Klotz, C., Kuhl, A.A., Heimesaat, M.M., Cantacessi, C., Hartmann, S., and
Rausch, S. (2019). RORgammat(+) Treg to Th17 ratios correlate with susceptibility to Giardia infection.
Sci Rep *9*, 20328.

Yoshida, T., Hachimura, S., and Kaminogawa, S. (1997). The oral administration of low-dose antigen
induces activation followed by tolerization, while high-dose antigen induces tolerance without
activation. Clin Immunol Immunopathol *82*, 207-215.

Zens, K.D., Chen, J.K., and Farber, D.L. (2016). Vaccine-generated lung tissue-resident memory T cells
provide heterosubtypic protection to influenza infection. JCI Insight *1*.

Zhao, J., Zhao, J., Mangalam, A.K., Channappanavar, R., Fett, C., Meyerholz, D.K., Agnihothram, S.,
Baric, R.S., David, C.S., and Perlman, S. (2016). Airway Memory CD4(+) T Cells Mediate Protective
Immunity against Emerging Respiratory Coronaviruses. Immunity *44*, 1379-1391.

Zhu, J., Yamane, H., and Paul, W.E. (2010). Differentiation of effector CD4 T cell populations (*).
Annu Rev Immunol *28*, 445-489.

Zychlinsky Scharff, A., Albert, M.L., and Ingersoll, M.A. (2017). Urinary Tract Infection in a Small
Animal Model: Transurethral Catheterization of Male and Female Mice. J Vis Exp.

Zychlinsky Scharff, A., Rousseau, M., Lacerda Mariano, L., Canton, T., Consiglio, C.R., Albert, M.L.,
Fontes, M., Duffy, D., and Ingersoll, M.A. (2019). Sex differences in IL-17 contribute to chronicity in
male versus female urinary tract infection. JCI Insight *5*.
891
892

893
894   Supplementary Information for
895
896
897 **Tissue-resident memory T cells mediate mucosal immunity to recurrent urinary tract infection**
898
899

900   Matthieu Rousseau[1,2][*], Livia Lacerda Mariano[1][#], Tracy Canton[1], Molly A Ingersoll[1,2]
901

902 [1]Mucosal Inflammation and Immunity, Department of Immunology, Institut Pasteur, 75015 Paris,
903 France and Inserm U1223, Paris France
904 [2]Université *Paris Cité*, Institut Cochin, INSERM U1016, CNRS UMR 8104, Paris, 75014, France
905 [#]current address: Institut Curie, PSL Research University, INSERM U932, 75005 Paris, France

906
907

909
910

911 **Supplementary Figure 1: CD8 T cells poorly infiltrate the bladder.** Graph shows the total number

912 of the specified T cell populations per bladder 24 hours post primary infection. A representative

913 experiment from 3 experiments is presented. Each circle is a mouse and lines are medians.

Supplementary Figure 2: No distinct CD4⁺ Th cell bias arises during primary and recurrent UTI in the lymph nodes. Six-week-old infected female C57BL/6 mice were sacrificed at the indicated timepoints post-primary (filled circles), or post-challenge infection (open circles). Day 0 are naïve mice and 'R' denotes animals that resolved their primary UTI but were not challenged with a second infection. The graphs depict total (**A**) CD4$^+$, Th1, Th2, or Th17 T cells, (**B**) IFNγ$^+$ Th1 or IL-17$^+$ Th17 T cells, and (**C**) Tregs, in the draining lymph nodes at the indicated day post primary or challenge infection. Treg/Th cell ratios for the indicated days post-primary (filled circles), or post-challenge infection (open circles) were calculated using the cell numbers in **Supplementary Figure 2A and C**. (**D**) Ratio of Treg/total CD4$^+$ Th cells and (**E**) ratio of Treg/indicated CD4$^+$ Th cell subset in the draining lymph nodes. Data are pooled from 2 experiments, n=5 to 7 mice/group in each experiment. Results from the bladders of these mice are shown in **Figure 2**. Each circle is a mouse and lines are medians. Nonparametric Mann-Whitney tests comparing cell numbers between '0' and 'R' were performed. All p-values were corrected for multiple comparisons across all populations shown using the false discovery rate (FDR) method and p-values <0.05 are in red.

932
933
Supplementary Figure 3. Limiting antigen persistence does not change Th cell polarization in
bladder-draining lymph nodes. Graphs show the total number of specified cell populations per
bladder-draining lymph nodes from mice presented in **Figure 4A-D** in which (**A**) shows draining lymph
nodes of mice presented in **Figure 4A** and **B, (B)** shows draining lymph nodes of mice presented in
Figure 4C and **D,** analyzed by flow cytometry. Data are pooled from 2 experiments, n=4 to 6 mice/group
in each experiment (**A**), or n=5 to 6 mice/group in each experiment (**B**). Each circle is a mouse and lines
are medians. Nonparametric Mann-Whitney tests comparing cell numbers between each condition were
performed. *p*-values were corrected for multiple comparisons using the false discovery rate (FDR)
method. Corrected *p*-values are shown.
943

945
946
Supplementary Figure 4. Limiting antigen persistence does not impact bladder-draining lymph
node T cells with a tissue resident memory phenotype. (**A** and **B**) Graphs show the total number of
specified cell populations per bladder draining lymph nodes from mice (**A**) presented in **Figure 5A** and
(**B**) presented in **Figure 5B**. Data are pooled from 2 experiments, n=3 to 6 mice/group in each
experiment. Nonparametric Mann-Whitney tests comparing cell numbers between each condition were
performed. *p*-values were corrected for multiple comparisons using the false discovery rate (FDR)
method. Corrected *p*-values for each comparison are presented.