Coevolution of relative brain size and life expectancy in parrots

Simeon Q. Smeele[1,2,3,4,*], Dalia A. Conde[6,4,5], Annette Baudisch[4], Simon Bruslund[7,8], Andrew Iwaniuk[9],

Johanna Staerk[4,5,6], Timothy F. Wright[10], Anna M. Young[11], Mary Brooke McElreath[1,2,^], Lucy

Aplin[1,12,^]

ORCIDs: SQS 0000-0003-1001-6615, DAC 0000-0002-7923-8163, AB 0000-0002-4202-089X, SB

0000-0003-4701-1754, AI 0000-0001-9273-3655, JS 0000-0001-6965-019X, TFW 0000-0003-2859-

5360, AMY 0000-0003-3321-1878, MBM 0000-0003-3206-5485, LA 0000-0001-5367-826X

9
[1] Cognitive & Cultural Ecology Research Group, Max Planck Institute of Animal Behavior, Radolfzell,

Germany

[2] Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary

Anthropology, Leipzig, Germany

[3] Department of Biology, University of Konstanz, Konstanz, Germany

[4] Interdisciplinary Centre on Population Dynamics, University of Southern Denmark, Odense,

Denmark

[5] Department of Biology, University of Southern Denmark, Odense, Denmark

[6] Species360 Conservation Science Alliance, Bloomington, USA

[7] Vogelpark Marlow gGmbH, Marlow, Germany

[8] European Association of Zoos and Aquaria, Parrot Taxon Advisory Group, Amsterdam, The

Netherlands

[9] Department of Neuroscience, University of Lethbridge, Lethbridge, Canada

[10] Biology Department, New Mexico State University, Las Cruces, USA

[11] Department of Biology and Earth Science, Otterbein University, Westerville, USA

[12] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany

^ Joint senior authors

* Corresponding author: ssmeele@ab.mpg.de

Abstract

Previous studies have demonstrated a correlation between longevity and brain size in a variety of taxa. Little research has been devoted to understanding this link in parrots; yet parrots are well-known for both their exceptionally long lives and cognitive complexity. We employed a large-scale comparative analysis that investigated the influence of brain size and life history variables on longevity in parrots. Specifically, we addressed two hypotheses for evolutionary drivers of longevity: the *Cognitive Buffer Hypothesis*, which proposes that increased cognitive abilities enable longer life spans, and the *Expensive Brain Hypothesis*, which holds that increases in life span are caused by prolonged developmental time of, and increased parental investment in, large brained offspring. We estimated life expectancy from detailed zoo records for 133,818 individuals across 244 parrot species. Using a principled Bayesian approach that addresses data uncertainty and imputation of missing values, we found a consistent correlation between relative brain size and life expectancy in parrots. This correlation was best explained by a direct effect of relative brain size. Notably, we found no effects of developmental time, clutch size, or age at first reproduction. Our results suggest that selection for enhanced cognitive abilities in parrots have in turn promoted longer lifespans.

Keywords

Psittaciformes; longevity; cognitive evolution; Bayesian structural equation model; cognitive buffer hypothesis; expensive brain hypothesis

Introduction

Evolutionary theories of ageing predict the inevitability of senescence in most iteroparous multicellular organisms (1–4). However, recent studies have highlighted the diversity of patterns and timing in which different taxa experience senescence, revealing species-specific patterns of longevity linked with allometry and life history variables (5,6). Generally, larger bodied species tend to live longer (7), but longevity is also associated with other variables such as diet, latitude and sociality (8,9). Perhaps of most recent interest, brain size has been correlated with longevity across diverse taxa ranging from amphibians (10) to primates (11). While some studies have proposed a negative relationship between brain size and longevity, suggesting a trade-off between the energetic costs into larger brains and investments in defences against ageing (e.g., (12)), the large majority of studies have suggested a positive effect of larger brain sizes on longevity (10,11,13–17). However, the causal pathways for this relationship between brain size and longevity are not yet well established.

Three non-mutually exclusive hypotheses have been proposed to explain the correlated evolution of larger brains and longer lifespans. First, the *Cognitive Buffer Hypothesis* posits that increased cognitive flexibility enabled by a relatively larger brain allows species to solve problems that would otherwise increase their extrinsic mortality, hence allowing for increased longevity (15). Second, the *Expensive Brain Hypothesis* argues that there is an indirect association between brains and longevity, with an investment in expensive brain tissue slowing down the pace of life through increased developmental time and increased parental investment per offspring (18). Third, the *Delayed Benefits Hypothesis* extends the *Expensive Brain Hypothesis* and reverses the directionality of its argument, positing that a shift to a higher quality, skill-based diet lowered adult mortality rates and supported a longer juvenile period that facilitated inter-generational skill transmission. This extended development in turn allows for investment in brain growth that further promotes skill-based foraging niches. In other words, long-lived, extractive foraging, species evolve larger brains because they can benefit most from learning (17). Previous work in mammals, amphibian and birds has found mixed support for all three hypotheses (13,16). For example, Isler et al. (18) showed that larger brained, monotokous (single offspring per reproduction), precocial mammals had longer developmental periods. This longer developmental period led to a prolonged life span; in other words, the effect of brain size on longevity

was indirect. In contrast, Jiménez-Ortega et al. (14) showed both a direct and an indirect effect of

absolute brain size on lifespan in birds, with larger brained species also living longer independently

from their developmental period.

Parrots (Psittaciformes) are famous for both their long lives and complex cognition (19,20), with

lifespans and relative brain size on par with primates (21). Indeed, recent studies on the genetics of

longevity and cognition in parrots have revealed positive selection on lifespan-prolonging genes, as

well as genes related to increased cognitive abilities and cell repair (22–24). Parrots are also

morphologically and ecologically diverse, with an extensive global distribution of almost 400 species,

ranging in size from adult yellow-capped pygmy parrots (*Micropsitta keiensis*, 12 g) to kakapo

(*Strigops habroptilus*, 3000 g) (25). In the first comparative study to examine longevity in parrots,

Munshi-South et al. (19) used maximum longevity records from 162 species, and found that both diet

and communal roosting were correlated with longevity, with granivorous and communal roosting

species living the longest on average. While not considering longevity, the potential drivers of the

evolution of brain size in Neotropical parrots were explored in Schuck-Paim et al. (26), finding that

brain size is associated with environmental and seasonal variability. Finally, highlighting the

importance of life history variation, Young et al. (27) found that longer lived parrots were more likely to

be threatened. To date, however, little research effort has been invested in understanding the link

between longevity and brain size in parrots.

One of the greatest challenges for comparative life history studies is sourcing good quality data (28).

For instance, the above studies all depended on maximum (or median) recorded lifespan, many used

regressions on residuals (see e.g., Gonzáles-Lagos et al. (29)) and some only included absolute brain

size (see e.g., Jiménez-Ortega et al. (14)). Maximum recorded lifespan can be a problematic measure

because it represents the longest-lived known individual and is therefore highly sensitive to sample

size. Making matters worse, how much sample size influences results depends on the pattern of age-

related mortality itself (30). For species where most individuals die around the same age, smaller

samples are more likely to approximate maximum longevity than in species with many extreme ages

of death. Therefore, a measure that accounts for all information available is preferable to a single-

point measure. Life expectancy is one such a measure and has been found to be the most

appropriate measure of pace of life (31). It calculates the average age at death based on information

across the full age range and therefore takes into account all available information. While life

expectancy can be sensitive to both intrinsic and extrinsic sources of mortality, the use of captive

records allows the removal of extrinsic sources of mortality as much as possible, thereby focusing on

senescence. Yet even when using captive data, other variables and shared evolutionary history

create confounds that need to be addressed within a multivariate framework. A principled way to

decide which covariates to include is the use of Directed Acyclical Graphs (DAG) (32,33). Based on a

specific hypothesis, a DAG represents all potential causal paths in the system by arrows. Conditional

on the DAG being true, the back-door criterion informs which variables should be included and which

should not be included (34). We additionally controlled for variables that only influence life expectancy

to improve accuracy of the model estimates.

Here, we present a phylogenetic comparative analysis focused on brain size and its effects on

longevity in parrots. First, we estimate life expectancy from Species360's Zoological Information

Management System (ZIMS) with records of 133,818 individuals across 244 parrot species. We then

test for a correlation between life expectancy and relative brain size after removing the effect of

covariates. Third, we used a DAG to distinguish between two possible pathways for this correlation.

The *Cognitive Buffer Hypothesis* predicts a direct effect of relative brain size on life expectancy, with

larger brained species living longer (15), while the *Expensive Brain Hypothesis* predicts that the effect

of brain size on life expectancy is indirect, emerging from increased developmental time and parental

investment per offspring (18). In this case, we expect that any relationship between brain size and life

expectancy will be reduced when also including parental investment (clutch size) and developmental

time in the model. While the *Delayed Benefits Hypothesis* would also predict a direct relationship

between relative brain size and longevity (17), it would argue for strong effect of diet, as well as

reversed directionality (extended longevity leads to larger brain sizes). While we included diet in our

models, our analysis focused explicitly on how brain size could affect longevity, and so we did not fully

explore this hypothesis. Overall, our study demonstrates a robust methodology for comparative life

history analysis using a comprehensive measure of life expectancy in a Bayesian statistical

framework. Moreover, it provides the most comprehensive analysis of longevity in Psittaciformes to

date, and contributes to a broader understanding of this understudied group.

Materials and Methods

Estimating life expectancy

We obtained data on birth and death dates from Species360's ZIMS. After cleaning (see

Supplementary Methods) we included records for 133,818 individuals across 244 species. To

estimate life expectancy, we implemented Bayesian Survival Trajectory Analysis (BaSTA, (35)), which

allowed us to make inferences on age-specific survival based on census data when ages of some

individuals are unknown. The model, implemented in R (36), uses a Markov Chain Monte Carlo

(MCMC) algorithm with Metropolis-Hastings sampling of mortality parameters and latent times of birth.

Here, we used a Siler hazard model (37) for each species, given by

149
$$\mu(x) = \exp[a_0 - a_1 x] + c + \exp[b_0 + b_1 x],$$

150   where $a_1, c, b_1 > 0$ and $a_0, b_0 \in (-\infty, \infty)$. These five parameters can fit infant and juvenile mortality

151   (controlled by a_0 and a_1), age independent (adult) mortality (c) as well as senescent mortality

152   (controlled by b_0 for initial mortality and b_1 for the rate of aging). Cumulative survival can be calculated

153   as

154
$$S(x) = \exp[-\int_0^x \mu(t)dt].$$

155   Life expectancy at birth is calculated as

156
$$e_0 = \int_0^\infty S(x)dx.$$

157   We used the Gelman-Rubin statistic (Rhat, (38)) to determine if models converged and visually

158   assessed the traces and model goodness of fit. When models did not converge, they were rerun with

159   longer burn-in and more iterations. If models clearly did not fit the data, the results were excluded.

160   This was the case for 27 out of 244 species. In most cases this was due to issues with data quality

161   (e.g., when the number of individuals without a recorded date of death was too high).

Life-history covariates

We collected body mass data from ZIMS. Additional body mass measurements were included from
the literature if no captive records were available for a species (28). We then used a Bayesian multi-
level model to extract species-level averages and standard errors (see Supplemental Methods more
details). Brain mass was collected by AI, from Iwaniuk et al. (39), from Schuck-Paim et al. (26) and
from Ksepka et al. (40), and similarly to body size, we fitted a Bayesian multi-level model to extract
species-level averages and standard errors. We also collected data for six additional potential
explanatory variables, based on previously proposed causal relationships with life expectancy: diet
(estimated protein content of main food items) (19), insularity (whether a species includes a
continental range or not) (19), maximum latitudinal range (as a proxy for environmental variability)
(41), clutch size (42), developmental time (from the start of incubation until fledging) and age of first
possible reproduction (AFR) (18). Diet, insularity, maximum latitude range, clutch size and
developmental time were collected from the literature. When data were not freely available, we
collected estimates directly from experts (see Supplemental Methods for the details). Finally, AFR is
unknown for the large majority of parrot species. We therefore estimated it directly from the
distribution of first breeding records in ZIMS, using the 5% percentile. To control for possible issues
arising from low sample sizes, we restricted this analysis to species with at least 30 breeding
individuals.

We used a DAG (see Figure 1) to decide how to incorporate variables in the statistical models,
accounting for their influences on each other in proposed causal pathways. It is important to note that
evolutionary time is not included explicitly in the DAG, thus arrows can potentially go in both
directions, representing evolutionary feedbacks. However, in our view, it represents the most
principled representation of the potential causal relationships for evolution of longevity in parrots,
based on available data and current knowledge. Although not depicted in the DAG, phylogenetic co-
variance was assumed to influence all variables and was included in all analyses using the L2-norm

(which calculates the covariance between two species based on a maximum possible covariance and

the squared distance between the two species) and the phylogenetic tree from Burgio et al. (43).

Figure 1 *Directed Acyclic Graph of the potential causal pathways that could drive parrot life expectancy. Colours*
represent different covariate groups and are kept consistent throughout the manuscript. Solid lines represent
assumed causal effects in all models (see Statistical analysis for model definitions). Dashed lines represent
additional causal relationships in model 2 and 3.

Statistical analysis

To test for a correlation between life expectancy and relative brain size, we first constructed a

Bayesian structural equation model (model 1) with life expectancy as the main variable to be

explained by relative brain size and four other potential covariates. We included a total of 360 species

for which at least one variable was known. The structure of this first model was as follows: LE ~ I +

BO + RB + LA + D, where LE = standardised log life expectancy, I = insularity (binary), BO =

standardised log body mass, RB = relative brain size, LA = standardised maximum latitude range and

D = protein content diet (ordinal). Relative brain size was calculated as: BR – pBR, where BR =

standardised log brain mass and pBR = predicted brain mass from a second model that ran

simultaneously: pBR ~ BO. Relative brain size has been shown to correlate with innovation rates in

birds (44) and we therefore use it as a proxy for cognitive flexibility. Our implementation is similar to

residual brain size in multiple regressions, but since both models are evaluated at each step of the

sampling, information flows in both directions and measurement error is modelled correctly (45). We

included standard error around the mean for life expectancy, body mass and brain mass. We also

included a phylogenetic variance-covariance matrix based on the phylogenetic distances calculated

from Burgio et al. (43), using the L2-norm. For each variable with missing data, missing values were

imputed using a multinormal distribution with mean and standard deviation based on the observed

data, variance-covariance based on the phylogenetic signal and means further informed by the causal

relationships outlined in Figure 1. For life expectancy we had data for 244 species, but the models

only converged for 217 species. Life expectancy for the remaining 143 parrot species was therefore

imputed (see Supplemental Methods for details).

To test whether any correlation between relative brain size and longevity could be indirectly caused

by developmental time, delayed juvenile periods, and/or parental investment, we ran a second model

(model 2) where developmental time (incubation period plus fledging period in model 2) and clutch

size were included as additional covariates. Both variables were log transformed and standardised.

Since data on AFR (a third measure of developmental time) was only available for 89 species and the

available data was biased towards later AFR (see Supplemental Methods for more detail), we did not

attempt to impute this variable, but tested its effect in a third model (model 3) limited to cases where

AFR was known.

To assess which hypothesis was best supported by the data, we compared the effect of relative brain

size in the three models. If an increase in relative brain size directly causes an increase in life

expectancy (*Cognitive Buffer Hypothesis*), we would expect the coefficient of the brain size effect to

be positive and similar in all three models. If an increase in relative brain size only causes an increase

in developmental time (*Expensive Brain Hypothesis*), we would expect the coefficient of the brain size

effect to be positive in model 1 and much reduced or zero in model 2 and 3.

Results

Overall, we were able to estimate life expectancy for 217 species out of 244 species for which we had

data. This included representatives of all eight major genera (i.e., those with at least ten species) and

over half of the extant parrot species. The shortest-lived genera were the small-bodied

Psittaculirostris and *Charmosyna*, e.g., with a life expectancy of less than 2 years for *Psittaculirostris*

desmarestii. The longest-lived genera were the large-bodied *Ara* and *Cacatua*, e.g., with a life

expectancy of more than 35 years for *Ara macao* (full distribution of values across the phylogenetic

tree is shown in Figure 2). Similarly, there was large variability in other covariates, e.g., with brain size

ranging from 1 to 22 grams, and age of first reproduction ranging from 7 months to 6 years. There

was a strong phylogenetic signal in life expectancy (Figure 2b), however, covariance was generally

low between species that belonged to different genera (Figure 3c).

Figure 2. *Phylogenetic tree of the 217 parrot species included in the study. Branches are coloured according to life expectancy (see density plot in bottom right), and phylogeny is based on Burgio et al. (43). Genera are named if they contain at least two species. For a version with all species named see Supplemental Figure S1.*

Model 1 (without developmental time and parental investment) as well as model 2 and 3 (including these potential indirect paths) had similar estimates for the direct effect of relative brain size. As expected, body size was strongly and positively correlated with life expectancy (see Figure 3c for model 2, Supplementary Results for model 1 and 3). Relative brain size also had a small, but consistently positive, effect on life expectancy ($\beta = 0.22$ in model 1, $\beta = 0.18$ in model 2 and $\beta = 0.16$ in model 3; overlap with zero < 0.03 for all models; Figures 3a, 4). Of the other life history variables included, none appeared to have a large effect on life expectancy (see Figure 3d-h). In particular,

model 2 showed no effect of developmental time (β = 0.01, overlap with zero > 0.22) or clutch size (β

= -0.08, overlap with zero > 0.88) on longevity, and there was no clear effect of AFR on longevity in

model 3 (β = -0.11, overlap with zero > 0.88). However, it should be noted that these models were

designed to test the effect of relative brain size, so other parameter estimates should be interpreted

with caution (46).

Figure 3. Parameter estimates for model 2. For results of model 1 and 3 see Figure S2 and S4. Grey density
plots and lines are the regularising priors. Coloured areas are the posterior densities for the parameter estimates
controlling the effect of the covariates on life expectancy. Black lines are 20 samples of the posterior for the
phylogenetic covariance. For insularity the difference between islandic and continental species is shown.

273
274 **Figure 4.** *Standardised relative log brain size vs life expectancy for model 2. Black points represent 217 species*
275 *where life expectancy was available, vertical black lines represent the SE for life expectancy, horizontal black*
276 *lines represent the 89% percentile intervals for standardised relative log brain size. Purple lines represent 20*
277 *samples from the posterior for the slope (beta) of the effect of standardised relative log brain size on life*
278 *expectancy.*

279
Discussion

Using an extensive database from captive parrots, our study showed a clear and positive correlation

between relative brain size and life expectancy in parrots. We further tested two hypotheses to

explain this observed correlation between relative brain size and life expectancy: the *Cognitive Buffer*

Hypothesis (15) and the *Expensive Brain Hypothesis* (18). Our results best supported a direct

relationship between larger brains and longer life expectancy, as predicted under the *Cognitive Buffer*

Hypothesis. It should be noted that this result is also consistent with the *Delayed Benefits Hypothesis*

(17). We would, however, also expect a strong effect of diet on life expectancy, since this hypothesis

argues that long life spans allow species to invest more time in learning foraging skill which requires

larger brains, and only pays off with an extended juvenile period. To fully explore this hypothesis we

would need data on post-fledging parental care and future studies could additionally try to use

process-based approaches (where evolution is modelled explicitly), such as generative inference (47)

or Bayesian ancestral state reconstruction (48) to disentangle the direction of causality. We found no

evidence that the relationship between relative brain size and life expectancy was explained by the

need for longer development times (here measured by incubation to fledging time, and by age of first

reproduction), or by increased parental investment (here represented by clutch size), as predicted by

the *Expensive Brain Hypothesis*. Interestingly, our results differ from a previous study in parrots by

Munshi-South et al. (19). This study found that the protein content of diets and communal roosting

best explained variation in maximum longevity. Data on sociality is largely lacking for parrots, so we

did not test for an effect of sociality, but we found no effect of diet. However, Munshi-South et al. did

not consider brain size in their analysis. Since diet potentially determines whether and how quickly

brains can grow (49), protein intake could still have an indirect effect on longevity via its potential link

with brain size.

The lack of support for the *Expensive Brain Hypothesis* is contrary to previous studies in primates

(11,50), other mammals (29,51), and amphibians (10), all of which show a positive correlation

between developmental time or AFR and life expectancy. However, it is in line with previous work

examining the evolution of longevity in birds (14). To explain this discrepancy between birds and

mammals, Isler et al. (16) suggested that bird species with allomaternal care (care provided for

mother or offspring by either the father or helpers) can provide enough nutrition for relatively larger

brained offspring without the need to prolong developmental periods or reduce clutch size to an extent

that would lead to the co-evolution of increased lifespans. All parrots have relatively large brain sizes

compared to most other birds, and all parrot species exhibit biparental care. Almost all parrots are

also cavity nesters. Cavity nests are less vulnerable to predation, and often have extensive nest

defence strategies, and so can have relatively relaxed selective pressure on fledging times as

compared to open-cup nesters (52). Perhaps the combination of these factors provides enough

flexibility to deal with heightened nutritional demands of rearing large-brained offspring without

selection on developmental times. This does not, however, diminish the importance of cognitive

development in parrots. The extended juvenile periods observed in many parrot species of up to six

years may provide enhanced opportunities for social learning, as proposed for another large-brained

bird taxon, the corvids (53). This hypothesis remains to be tested in parrots.

To our knowledge this is the first study of life expectancy and/or brain size that uses a bespoke

Bayesian model to include: 1) uncertainty about variable estimates; 2) imputation of missing values;

3) a principled representation of relative brain size; and 4) phylogenetic signal. We believe this

method has some major advantages. Most notably, we could estimate both life expectancy and its

uncertainty in each species. This allowed us to fully exploit the fact that we have a hundred-fold more

data for some species, instead of relying on a single point estimate of maximum longevity as in

previous studies of longevity in parrots (19,27). We also imputed life expectancy for species which

have no data. This is likely to be important in most datasets to account for biased data collection, but

it is especially important when using data from captivity, because zoos do not randomly pick species

to be included in their population, but have a general bias toward larger and longer-lived species (54).

Complete case analysis will introduce bias in this case (55) and we therefore chose to impute missing

values. The use of DAGs and structural equation models is very similar to path analysis. The main

advantage of our implementation is that it allows for a statistically robust definition of relative brain

size and can handle uncertainty and missing data. Our model structure can be easily adapted to

impute any continuous variable.

Our study also departs from most previous studies of longevity by using data from captivity on life

expectancy (41,56–58). This provided several important advantages. First, it provided a large sample

size, both improving the estimation of life expectancy per species and allowing us to have a fuller

representation of species. Second, captivity reduces external sources of mortality as much as

possible (little predation, starvation, etc.). However, captive data poses different challenges. First, as

with data from the wild, birth and death dates can be missing (e.g., for individuals born in the wild or

transferred from institutions that are not part of ZIMS). The BaSTA implementation that we used

imputed these missing values, and we believe that our thorough cleaning procedure, coupled with the

sheer magnitude of the dataset, means that any gaps, data entry errors or biases should have

minimal effect on the life expectancies presented here. Second, there may be differences in causes of

death in captivity and the wild, for example if some species are difficult to keep or prone to negative

behavioural responses to captivity which is also true for some of the shortest-lived genera included in

the study such as *Psittaculirostris* and *Charmosyna* which have been historically difficult to manage in

captivity. We dealt with this by excluding potentially problematic species from the initial life

expectancy estimations, and instead imputed values in the final model (see Supplemental Methods

for details). We can still not be completely sure that the patterns observed in the data are all

representative of the evolutionary processes that shaped them, but it is highly unlikely that the clear

positive correlation between relative brain size and life expectancy is due to captivity. It could even be

expected that large brained species live shorter in captivity, because of the higher metabolic rates

required to keep the large brain supplied with glucose. This has been shown to be the case within

species in captive guppies (12). Since such an effect would be opposite to the one observed in our

study, its presence would not change the conclusions drawn from our results.

Conclusions

Overall, our results are consistent with the *Cognitive Buffer Hypothesis*, suggesting that relatively

large brains may have buffered parrots against environmental variability and/or predation threats

reducing sources of extrinsic mortality and allowing longer lifespans. This result is consistent with

previous studies in other birds, suggesting that common processes may explain longevity in altricial

birds. In addition to their longevity, parrots are famous for their complex cognition. It remains largely

unknown what evolutionary processes have driven cognitive evolution in parrots, but given the results

of our study, in addition to those of Munshi-South et al. (19), future work should further investigate the

potentially complex feedbacks between these two factors and sociality and diet. Unfortunately, longer

lived species are also more likely to be threatened (27), showing the vulnerability of this order. Having

life expectancy and other life history variables for hundreds of species will hopefully aid in future

conservation efforts for this globally threatened order.

Data, code and materials

Data, code and materials are publicly available at https://doi.org/10.5061/dryad.sbcc2fr7x. A

repository is also publicly available at

https://github.com/simeonqs/Coevolution_of_relative_brain_size_and_life_expectancy_in_parrots.

Competing interests

The authors have no competing interests with this study.

Author Contributions

AMY, DAC, LMA, MBM and SQS conceived the idea. AI, AMY, LA, TFW, SB and SQS collected the data. SQS analysed the data under supervision from LA and MBM. AB, DAC, LA, MBM and SQS drafted the initial manuscript, and all authors contributed to writing and editing the final article. All contributors are listed in alphabetical order.

Acknowledgments

We would like to thank Dr. Dieter Lukas, Dr. Cody Ross, Prof. Fernando Colchero, Prof. Richard McElreath and Dr. Rita da Silva for their advice on the analysis. We would further like to thank the more than 1,300 Species360 members for registering their animals in the ZIMS database. We thank Prof. Alejandro Salinas, Geddes Hislop and Johann C. Carstens for contributing data on diet.

Funding

This project was possible thanks to the financial support of the sponsor members of the Species360 Conservation Science Alliance (Copenhagen Zoo, the World Association of Zoos and Aquariums, and the Wildlife Reserves of Singapore), the Interdisciplinary Centre on Population Dynamics and the Biology Department at the University of Southern Denmark. We would like to thank Species360 for granting access to the data under permission number #86892. This work was supported by the Max Planck Society. LMA was funded by a Max Planck Independent Group Leader Fellowship. SQS received additional funding from the International Max Planck Research School for Organismal Biology.

References

1. Medawar PB, University College London. An Unsolved problem of biology; an inaugural lecture delivered at University College, London, 6 December, 1951. London: H.K. Lewis and Co.; 1952.

2. Williams GC. Pleiotropy, natural selection, and the evolution of senescence. Evolution. 1957;398–411.

3. Hamilton WD. The moulding of senescence by natural selection. J Theor Biol. 1966;12(1):12–45.

4. Kirkwood TB. Evolution of ageing. Nature. 1977;270(5635):301–4.

5. Jones OR, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB, et al. Diversity of ageing across the tree of life. Nature. 2014;505(7482):169–73.

6. Colchero F, Jones OR, Conde DA, Hodgson D, Zajitschek F, Schmidt BR, et al. The diversity of population responses to environmental change. Ecol Lett. 2019;22(2):342–53.

7. Lindstedt SL, Calder III WA. Body size, physiological time, and longevity of homeothermic animals. Q Rev Biol. 1981;56(1):1–16.

8. Wasser D, Sherman P. Avian longevities and their interpretation under evolutionary theories of senescence. J Zool. 2010;280(2):103–55.

9. Scholer MN, Strimas-Mackey M, Jankowski JE. A meta-analysis of global avian survival across species and latitude. Ecol Lett. 2020;23(10):1537–49.

10. Yu X, Zhong MJ, Li DY, Jin L, Liao WB, Kotrschal A. Large-brained frogs mature later and live longer. Evolution. 2018;72(5):1174–83.

11. Street SE, Navarrete AF, Reader SM, Laland KN. Coevolution of cultural intelligence, extended life history, sociality, and brain size in primates. Proc Natl Acad Sci. 2017;114(30):7908–14.

12. Kotrschal A, Corral-Lopez A, Kolm N. Large brains, short life: selection on brain size impacts intrinsic lifespan. Biol Lett. 2019;15(5):20190137.

13. DeCasien AR, Thompson NA, Williams SA, Shattuck MR. Encephalization and longevity evolved in a correlated fashion in Euarchontoglires but not in other mammals. Evolution. 2018;72(12):2617–31.

14. Jiménez-Ortega D, Kolm N, Immler S, Maklakov AA, Gonzalez-Voyer A. Long life evolves in large-brained bird lineages. Evolution. 2020;74(12):2617–28.

15. Allman J, McLaughlin T, Hakeem A. Brain weight and life-span in primate species. Proc Natl Acad Sci. 1993;90(1):118–22.

16. Isler K, Van Schaik CP. Why are there so few smart mammals (but so many smart birds)? Biol Lett. 2009;5(1):125–9.

17. Kaplan H, Hill K, Lancaster J, Hurtado AM. A theory of human life history evolution: Diet, intelligence, and longevity. Evol Anthropol Issues News Rev Issues News Rev. 2000;9(4):156–85.

18. Isler K, van Schaik CP. The expensive brain: a framework for explaining evolutionary changes in brain size. J Hum Evol. 2009;57(4):392–400.

19. Munshi-South J, Wilkinson GS. Diet influences life span in parrots (Psittaciformes). The Auk. 2006;123(1):108–18.

20. Auersperg AM, von Bayern AM. Who'sa clever bird—now? A brief history of parrot cognition.
Behaviour. 2019;156(5–8):391–407.

21. Olkowicz S, Kocourek M, Lučan RK, Porteš M, Fitch WT, Herculano-Houzel S, et al. Birds have
primate-like numbers of neurons in the forebrain. Proc Natl Acad Sci. 2016;113(26):7255–60.

22. Costantini D, Racheli L, Cavallo D, Dell'Omo G. Genome size variation in parrots: longevity and
flying ability. J Avian Biol. 2008;39(4):453–9.

23. Andrews CB, Gregory TR. Genome size is inversely correlated with relative brain size in parrots
and cockatoos. Genome. 2009;52(3):261–7.

24. Wirthlin M, Lima NC, Guedes RLM, Soares AE, Almeida LGP, Cavaleiro NP, et al. Parrot
genomes and the evolution of heightened longevity and cognition. Curr Biol. 2018;28(24):4001-
4008. e7.

25. Dunning Jr JB. CRC handbook of avian body masses. CRC press; 2007.

26. Schuck-Paim C, Alonso WJ, Ottoni EB. Cognition in an ever-changing world: climatic variability is
associated with brain size in neotropical parrots. Brain Behav Evol. 2008;71(3):200–15.

27. Young AM, Hobson EA, Lackey LB, Wright TF. Survival on the ark: life-history trends in captive
parrots. Anim Conserv. 2012;15(1):28–43.

28. Conde DA, Staerk J, Colchero F, da Silva R, Schöley J, Baden HM, et al. Data gaps and
opportunities for comparative and conservation biology. Proc Natl Acad Sci. 2019;116(19):9658–
64.

29. González-Lagos C, Sol D, Reader S. Large-brained mammals live longer. J Evol Biol.
2010;23(5):1064–74.

30. Vaupel JW. Post-darwinian longevity. Popul Dev Rev. 2003;29:258–69.

31. Wrycza TF, Baudisch A. The pace of aging: Intrinsic time scales in demography. Demogr Res.
2014;30:1571–90.

32. Laubach ZM, Murray EJ, Hoke KL, Safran RJ, Perng W. A biologist's guide to model selection
and causal inference. Proc R Soc B. 2021;288(1943):20202815.

33. Pearl J, Glymour M, Jewell NP. Causal inference in statistics: A primer. John Wiley & Sons; 2016.

34. Pearl, Judea M Dana. The book of why. 2018.

35. Colchero F, Jones OR, Rebke M. BaSTA: an R package for Bayesian estimation of age-specific
survival from incomplete mark–recapture/recovery data with covariates. Methods Ecol Evol.
2012;3(3):466–70.

36. R Core Team. R: A language and environment for statistical computing. 2013; Available from:
https://www.R-project.org/

37. Siler W. A competing-risk model for animal mortality. Ecology. 1979;60(4):750–7.

38. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian data analysis. CRC
press; 2013.

39. Iwaniuk AN, Dean KM, Nelson JE. Interspecific allometry of the brain and brain regions in parrots
(Psittaciformes): comparisons with other birds and primates. Brain Behav Evol. 2005;65(1):40–
59.

40. Ksepka DT, Balanoff AM, Smith NA, Bever GS, Bhullar B-AS, Bourdon E, et al. Tempo and
pattern of avian brain size evolution. Curr Biol. 2020;30(11):2026-2036. e3.

41. Scholer MN, Strimas-Mackey M, Jankowski JE. A meta-analysis of global avian survival across
species and latitude. Ecol Lett. 2020;23(10):1537–49.

42. Ghalambor CK, Martin TE. Fecundity-survival trade-offs and parental risk-taking in birds. Science.
2001;292(5516):494–7.

43. Burgio KR, Davis KE, Dreiss LM, Cisneros LM, Klingbeil BT, Presley SJ, et al. Phylogenetic
supertree and functional trait database for all extant parrots. Data Brief. 2019;24:103882.

44. Overington SE, Morand-Ferron J, Boogert NJ, Lefebvre L. Technical innovations drive the
relationship between innovativeness and residual brain size in birds. Anim Behav.
2009;78(4):1001–10.

45. Freckleton RP. On the misuse of residuals in ecology: regression of residuals vs. multiple
regression. J Anim Ecol. 2002;542–5.

46. Westreich D, Greenland S. The table 2 fallacy: presenting and interpreting confounder and
modifier coefficients. Am J Epidemiol. 2013;177(4):292–8.

47. Kandler A, Powell A. Generative inference for cultural evolution. Philos Trans R Soc B Biol Sci.
2018;373(1743):20170056.

48. Ringen E, Martin JS, Jaeggi A. Novel phylogenetic methods reveal that resource-use
intensification drives the evolution of "complex" societies. 2021;

49. Robson S. Breast milk, diet, and large human brains. Curr Anthropol. 2004;45(3):419–25.

50. Powell LE, Barton RA, Street SE. Maternal investment, life histories and the evolution of brain
structure in primates. Proc R Soc B. 2019;286(1911):20191608.

51. Barton RA, Capellini I. Maternal investment, life histories, and the costs of brain growth in
mammals. Proc Natl Acad Sci. 2011;108(15):6169–74.

52. Martin TE. Nest predation and nest sites. BioScience. 1993;43(8):523–32.

53. Uomini N, Fairlie J, Gray RD, Griesser M. Extended parenting and the evolution of cognition.
Philos Trans R Soc B. 2020;375(1803):20190495.

54. Frynta D, Lišková S, Bültmann S, Burda H. Being attractive brings advantages: the case of parrot
species in captivity. PloS One. 2010;5(9):e12568.

55. Little RJ, Rubin DB. Statistical analysis with missing data. Stat Anal Missing Data. 2002;

56. Blumstein DT, Møller AP. Is sociality associated with high longevity in North American birds? Biol
Lett. 2008;4(2):146–8.

57. Beauchamp G. Group-foraging is not associated with longevity in North American birds. Biol Lett.
2010;6(1):42–4.

58. Arnold KE, Owens IP. Cooperative breeding in birds: a comparative test of the life history
hypothesis. Proc R Soc Lond B Biol Sci. 1998;265(1398):739–45.

519