1
9   **Vagus nerve stimulation accelerates motor learning through cholinergic modulation.**
10

11   Jordan Hickman*[1], Spencer Bowles*[1,2], Xiaoyu Peng*[1,2], W. Ryan Williamson[3], Rongchen Huang[1,2],
12   Kayden Washington[1], Dane Donegan[1,2], Cristin G Welle[1,2,4]

31   [1]Department of Physiology and Biophysics, [2]Department of Neurosurgery, [3]IDEA Core, University of
32   Colorado School of Medicine. * These authors contributed equally to this work. [4]Corresponding author.
33   Email: cristin.welle@cuanschutz.edu.

Abstract

Vagus nerve stimulation (VNS) is a novel therapeutic option to treat a broad and rapidly expanding set of neurologic conditions. While classically used to treat epilepsy and depression, more recently VNS has received FDA approval as a therapeutic option for stroke rehabilitation and is under preclinical and clinical investigation for other disorders. Despite benefits across a diverse range of neurological disorders, the mechanism for how VNS influences central nervous system circuitry is not well described. A deeper understanding of the influence of VNS on neural circuits and activity is needed to optimize the use of VNS therapy.

To define the complex dynamics between VNS, neuronal activity, plasticity and behavior, we performed chronic VNS in mice during the learning of a dexterous motor task, and leveraged genetic tools to perform optogenetic circuit dissection and longitudinal *in vivo* imaging calcium activity in cortical neurons. We find VNS has the most robust effect on motor learning when paired with successful movement outcome, while randomized stimulation impairs learning, consistent with VNS serving as a reinforcement cue. In motor cortex, VNS paired with movement outcome selectively modulates neurons that are representing outcome, but not other movement-related neurons. Finally, cholinergic signaling from basal forebrain is required both for VNS-driven improvements in motor learning and the effects on neural activity in M1. This suggests that the effect of VNS on motor learning is mediated by cholinergic signaling, and also presents a novel role for cholinergic signaling in endogenous motor learning. These data imply that VNS therapy may be mediated by augmenting reinforcement cues with precisely-timed cholinergic neuronal activity, presenting strategies for optimizing the use of VNS to treat neurologic conditions.

Introduction

Vagus Nerve Stimulation (VNS) is currently used in clinical care to treat epilepsy[1,2] and depression[3], but novel responsive stimulation paradigms are being explored to treat a broad and growing range of neurologic injuries [4–6]. Recently, VNS paired with motor rehabilitation was approved by the FDA for the treatment of motor deficits associated with stroke [6]. Preclinical and early clinical studies suggest that paired VNS speed functional recovery from other neurologic conditions, including spinal cord injury, peripheral nerve injury and traumatic brain injury [7–9]. Despite the wide-ranging etiology of these conditions, the therapeutic model is similar: VNS is paired with a relevant rehabilitation protocol. It is hypothesized that this precise timing of stimulation drives targeted circuit plasticity for recovery from injury[5,10]. Yet, the lack of a clear circuit mechanism limits optimization of VNS therapy to treat neurologic injury.

The circuitry that mediates VNS effects on central nervous system plasticity remains poorly understood. Vagus nerve afferents terminate in the brainstem nucleus tractus solitarius (NTS) [11], which in turn activates many subcortical and cortical brain regions, including major subcortical neuromodulatory nuclei [12–14]. Lesions of major neuromodulatory centers, including the cholinergic basal forebrain (BF), limit VNS-driven cortical map plasticity [15,16] and functional rehabilitation after nerve damage [9]. In addition, the cholinergic basal forebrain has been indicated as necessary for motor learning[17], and phasic cholinergic signals are thought to play critical roles in reinforcement learning and outcome representation [18,19]. Together, these suggest a possible role for phasic cholinergic signaling in mediating the effects on VNS-driven learning, although this remains to be substantiated. In addition to activation of neuromodulatory nuclei, VNS has a well-document effect of decorrelating neural activity in cortex and across other brain areas[20–22]. Recent in vivo imaging studies have confirmed a broad, excitatory effect of VNS across multiple cortical regions[14]. Yet, the effects of VNS are specific to a paired sensory or motor stimuli, suggesting VNS may confer more specific modulation of key neural circuits.

To understand the mechanisms by which VNS can modulate neural circuits to optimally enhance motor behavior, we compared the effect VNS timing on skilled reach learning and then used optogenetic cholinergic circuit manipulation, kinematic analysis, and *in vivo* calcium imaging in the motor cortex of freely behaving animals to dissect the circuit that underlies VNS action. Paired-VNS accelerated the learning of a dexterous reach task, but only when applied as a reinforcement signal for a successful reach (Success VNS). Cholinergic neural activity in BF was required for the effects of VNS on motor learning. Improved reach performance by Success VNS was achieved by increased reach consolidation over learning, which also depended on cholinergic signaling. Finally, VNS altered specific neural populations relevant to reach representation in primary motor cortex, and the effects of VNS in M1 were mitigated by cholinergic antagonists. Our results reveal that VNS serves as a reinforcement signal during early learning and acts

through cholinergic signaling to modulate the activity of behaviorally-relevant neural populations in motor

cortex, resulting in increased reach consolidation and improved motor learning.

Results

VNS enhances motor skill learning when paired with successful task outcome

To induce motor rehabilitation and cortical plasticity, VNS must be paired with motor output[23,24] at a

relatively short latency[8], yet a single optimal pairing protocol has not been identified. To explore the optimal

timing of VNS during skilled motor learning, we applied multiple VNS pairing protocols as freely moving

mice learned a dexterous forelimb reaching task. Using a newly developed chronic VNS approach for mice

[25], we implanted a microfabricated stimulation cuff on the left cervical vagus nerve (**Fig. 1a**) connected to

a skull-mounted headcap. Mice were trained daily for 14 days (**Fig. 1b,c,**).

102   To identify an optimal VNS pairing strategy for enhancing dexterous motor learning, we explored

four pairing protocols (**Fig. 1b**). First, to test if VNS drives plasticity by increasing widespread cortical

excitation and inducing general arousal during a session [14], VNS was applied at pseudo-random intervals

(Random VNS) during the 20-minute training session. Alternatively, VNS may modulate short-term

attention, or influence spike timing dependent plasticity [26]. To test this, VNS was applied at the initiation

of a subset of reach movements (Reach VNS). A third option is that VNS may augment reward or

reinforcement pathways, and is better paired with movement outcome [27,28]. To explore this option, a third

cohort received VNS after successful reach completion (Success VNS). The surgical control cohort was

implanted with stimulation cuffs and connected to a stimulation isolation unit that was turned 'off' (Sham

VNS). Current amplitude was the same across groups and the number of stimulation trains delivered during

training sessions did not correlate with reach success (**Supplementary Fig.1a-c**).

113   Animals in all cohorts (Sham VNS, Random VNS, Reach VNS, and Success VNS) learned the

114   dexterous reach task (Sham VNS: p=0.0001, Random VNS p=0.0001, Reach VNS p=0.0002, Success VNS

115   p=0.001; Supplementary **Fig. 2b-e**). Neither Random nor Reach VNS altered the success rate of the animals

116   relative to Sham VNS (Random, 47.4% ± 3.9%; Reach 53.6% ± 3.5%; Sham, 46.3% ± 3.2%; **Fig. 1g**).

117   However, Success VNS improved the success rate over Sham VNS (59.2% ± 3.1% vs 46.3% ± 3.2% **Fig.**

118   **1g**), demonstrating for the first time, that paired VNS can accelerate motor learning in healthy animals.

Figure 1 | VNS modulates forelimb reach learning and requires temporally specific stimulation. a-c, Illustration of VNS surgery, behavior timeline, and stimulation protocol. **d-f,** Random VNS, Reach VNS, and Success VNS success rate across 14 sessions of training. **g,** Comparison of mean performance across all days between control and stimulated groups (Success VNS: p=0.0065, f=9.24, Random VNS: p>0.05, Reach VNS: p>0.05, Student T test). **h,** Comparison of mean success rate for control and Random VNS mice during early (p=0.028, f=7.07, Student T test) and late learning (p>0.05). **i,** Comparison of mean success rate for control and Reach VNS mice during early and late learning (p>0.05). **j,** Comparison of mean success rate for control and Success VNS mice during early (p=0.0031) and late learning (p=0.0126). **k,l,** in-session learning trajectories for each group. **m,** Comparison of within session learning between all groups across 4 days of learning. **n, o,** VNS mice performed a conditioned place preference test after 3 days being stimulated in one of two distinct rooms. *p < 0.05, **p < 0.01, ***p< 0.001 bars and error bars represent the mean ± s.e.m.

In addition, these results indicate the importance of pairing VNS as a reinforcement cue during
skilled motor learning.

VNS serves as a reinforcement cue during early learning but is not inherently rewarding.
Prior work on learning of a dexterous reach has suggested a multiphasic approach to learning, with distinct
early and late learning phases [29,30], yet the time of early to late transition has not been empirically
demonstrated. Using a Weibull growth curve nonlinear model ($55.49\% \pm 6.81$) we identified early learning
as days 1-4 and late learning as days 5-14 (**Supplementary Fig. 2a)**. We next examined if VNS exerted
distinct effects during different learning stages. Interestingly, despite having no effect on the 14-day average
results, Random VNS impaired early learning ($27.3\% \pm 7.3\%$ versus $37.4\% \pm 8.9\%$), but performance
recovered during late learning (**Fig. 1h)**. Reach VNS had no influence on success rates at any phase of
learning (**Fig. 1i)**. Success VNS increased success rates during the early phase and late phase (early: 50.6%
$\pm 9.4\%$ versus $37.4\% \pm 8.9\%$; late: $63.6\% \pm 6.9$ versus $49.6\% \pm 13.2\%$ **Fig. 1j)**. These data suggest that
VNS paired with a successful outcome accelerates learning and increases the final proficiency of a forelimb
task, while random application of stimulation temporarily impairs learning during early learning.

To determine if VNS alters within-session learning or between-session learning [31], training session
data was grouped into 4 blocks of 5 trials each (**Fig. 1k,l)**, and then the within-session learning slope was
quantified over the first 4 days (**Fig.1m)**. Within-session learning slope was not significantly different
between conditions, suggesting that Success VNS likely operates by promoting between-session learning.
One means by which Success VNS might confer learning benefit is through activation of a classical reward
system [32]. To determine if VNS serves as a rewarding or aversive stimulus, we used a well-documented
behavioral assessment of reward, the conditioned place preference test (CPP) (**Fig. 1n)**. Implanted mice
were introduced to two rooms with distinct visual and olfactory cues, with VNS applied in one room. On
the final day probe session, mice spent equal time in the conditioned (stimulated) room on the initial and
final test days ($44.8\% \pm 2.0\%$ and $41.4\% \pm 2.3\%$, **Fig.1o**), indicating that VNS is not inherently rewarding
or aversive.

VNS confers both short-term performance benefits during the execution of known tasks.
Despite a lack of evidence for within-session learning, we wanted to confirm the behavioral results were
due to learning, and not short-term modulation of attention. Since Success VNS is applied after reach
outcome, the success rate of trials that immediately follow a stimulation (post-success) were compared to

Figure 2 | VNS improves success rate within sessions and in learned mouse during rehearsal of forelimb reach task. a, trials before and after a stimulated success are investigated in success VNS **b,** Comparison of success rate for reaches preceding or following stimulated reach. **c,** For Success VNS, day 7 and 14 trials are divided into equal blocks of unstimulated and stimulated trials. **d,** Comparison of unstimulated (light orange) and sham (grey) trials on day 14 (p=0.0059, t=3.34, Student's T test). **e&f,** Comparison of stimulated (dark orange) and unstimulated (light orange) trials on days 7 & 14 (Day 7: p>0.05; Day 14: p=0.0499, t=2.57, Ratio paired t test). **g,** Reach VNS schematic **h,** Success rates across learning training sessions for Sham (grey), Stimulated Reach VNS (dark green), and Unstimulated Reach VNS (light green). **i,** Comparison of stimulated Reach VNS and Sham VNS trials in early (p>0,05) and late learning (p=0.024, t=2.47, Student's T test). **j,** Comparison between stimulated Reach VNS and unstimulated Reach VNS trails in early (p=0.004, t=3.98, paired t test) and late learning (p<0.0001, t=10.08, paired t test). **k,** Success VNS and Reach VNS applied during rehearsal of reach task in naively trained mice. **l,** Stimulated Reach VNS trials improve success rate during rehearsal (p=0.015, t=4.058, paired t test). **m,** Stimulated Success VNS trials improve success rate during rehearsal (p=0.005, t=5.62, paired t test). **n,** Normalized improvement of stimulated Reach VNS (p=0.047, t=3.56) and stimulated Success VNS trials (p=0.028, t=4.16) compared to unstimulated trials (Sidak's multiple comparison's test in RM one-way ANOVA). *p < 0.05, **p < 0.01, ***p < 0.001; bars and error bars represent the mean ± s.e.m.

those immediately prior (pre-success), and there was no trial-level difference (**Fig. 2a,b**). We next

compared the response to Success VNS to an unstimulated probe trial block included on days 7 and 14(**Fig.**

2c). The success rate for unstimulated trials on Day 14 was greater than Sham VNS (61.9% ± 6.3% versus

46.6% ± 9.2% **Fig. 2d)**, suggesting that Success VNS led to stimulation-independent, lasting learning. Yet
also, performance during stimulated blocks was greater than unstimulated blocks on day 14 (69.7% ± 10.2%
and 61.9% ± 6.3; **Fig. 2f**), but not day 7 (**Fig. 2e**), implying a short-term performance benefit in mice that
have learned the task, in addition to the improved learning.

To further explore potential short-term benefits of VNS, we took advantage of the Reach VNS trial
design, which allowed subgroup analysis of only stimulated or unstimulated reaches (**Fig. 1g**). The success
rate of only stimulated trials is greater than sham during late learning (64.0% ± 9.4% versus 50.8% ± 13.7)
(**Fig. 2i),** while unstimulated trials are not different from sham. Paired analysis for individual animals shows
a higher success rate for stimulated trials compared to unstimulated trials in both the early and late phase
(early: 41.9% ± 10.2% vs 35.3% ± 9.7; late: 64.0% ± 9.4% vs 53.7% ± 9.5%) (**Fig. 2j).** Taken together,
Reach VNS provides a short-term performance boost for stimulated over unstimulated trials throughout
learning. Yet, like Success VNS, Reach VNS most effectively modulates short-term performance for motor
skills that are already learned.

Finally, to confirm that VNS augments the performance of a known task, paired VNS was applied
to animals already proficient in the skilled reaching task (**Fig. 2k**). Both Success VNS (**Fig. 2m**) and Reach
VNS (**Fig. 2l**), delivered on alternate days for 10 days, improved performance over trial blocks without
VNS (Success VNS 46.0% ± 8.5% vs 40.8% ± 8.9%; Reach VNS 49.0% ± 10.2% versus 38.8% ± 9.5%),
confirming that either pairing protocol is sufficient to modulate the short-term performance of a known
task.

VNS drives neural activity in the basal forebrain (BF)
Cholinergic neuromodulation is associated with reinforcement-driven plasticity[18,33] and is required for
motor learning[34–36]. The basal forebrain (BF) is the source of cortically-projecting cholinergic neurons[37],
however the response of BF neurons to VNS *in vivo* remains unknown. To address this question, tetrodes
were implanted into the BF of mice with VNS cuffs (**Fig. 3a**). Extracellular activity was recorded during
VNS in awake animals. (30 Hz, 0.6 mA, 100us pulse, 500ms train). Neurons in the BF displayed an altered
firing rate following VNS onset, as compared to baseline (**Fig. 3b,c**). VNS elicited activation or suppression
in 43% of recorded units, with 61% of activated units showing increased activity (**Fig. 3d**) which began
during the stimulation train and persisted after stimulation ended (**Fig. 3e**).

The BF contains multiple neuronal cell types [38], so opto-tagging was used to identify cholinergic
BF neurons[39] and interrogate their VNS response. Acute recordings were performed in anesthetized ChAT-
ChR2 transgenic mice (B6.Cg-Tg(Chat-COP4*H134R/EYFP,Slc18a3)6Gfng/J) (**Fig. 3f**). Optrodes were

Figure 3 | VNS drives BF neural activity in anesthetized and awake mice.
a, Tetrodes were implanted in the left BF of mice and recordings were obtained during awake behavior. **b,** Example raster (top) and average firing rate from a response to VNS. Grey box denotes stimulus delivery. **c,** Average responses of all recorded neurons to VNS (grey box). **d,** Percent of neurons that respond to VNS (N = 5 mice, 53 neurons). **e,** Average activity of all 'activated' neurons in response to VNS. Dashed lines mark significance, shading represents SE. **f,** Extracellular recordings were obtained using optrodes while under light anesthesia. **g,** Optrodes were targeted at the HDB in *ChAT-ChR2* transgenic mice. Green fluorescence denotes the presence of ChR2. **h,** Example cholinergic neuron responding consistently to pulses of 488 nm light. **i,** Average activity of all cholinergic neurons during opto-tagging. Each row represents a neuron. **j,** Stimulus-associated latency tests (SALT) separate light responsive neurons from non-light responsive neurons. **k,** Mean baseline FR of cholinergic and non-cholinergic neurons (p=0.013, N = 5 mice, 53 neurons). **l,** Percent of neurons categorized as cholinergic (light green) and non-cholinergic (dark green) (N = 6 mice, 76 neurons). **m,** Percent of units that are VNS-responsive in cholinergic (left) and noncholinergic (right) populations. **n,** Average response to VNS for all 'activated' neurons. **o,** Mean peak activation during VNS. **p,** Average delay of peak activation from VNS onset. **q,** Mean duration of significantly elevated activity after VNS (cholinergic vs awake p=0.0087, non-cholinergic vs awake p=0.0003).

were identified by their rapid response to light (**Fig. 3h,i**) and confirmed using SALT analysis[18] (latency 5.2±1.3ms) (**Fig. 3j**). Cholinergic neurons exhibited a lower baseline firing rate (3.4±2.1hz) than the non-cholinergic population (6.8±6.2hz) (**Fig. 3k**). Of 76 units, roughly ⅓ were cholinergic (**Fig. 3l**) and half of both neuron populations were VNS responsive (52% of cholinergic, 49% of non-cholinergic; **Fig. 3m**). Of the VNS-responsive units, most cholinergic units and non-cholinergic units showed increased activity (**Fig. 3n**). This data suggests that VNS increases activity in both cholinergic and non-cholinergic populations of the BF.

A comparison of VNS-driven activation in anesthetized recordings to awake recordings suggest that the response to VNS depends on arousal state (**Fig. 3e&n**). While the peak response magnitude and timing to VNS do not change between awake and anesthetized animals (**Fig. 3o&p**), awake animals have a longer response than anesthetized animals (awake 652.7±439 ms; cholinergic: 293.7±94 ms; non-cholinergic: 235.8±127 ms; **Fig. 3q**). These results demonstrate that VNS alters BF activity during awake behavior, making it a strong candidate for mediating VNS-driven behavioral effects.

Optogenetic cholinergic inhibition prevents VNS-enhanced motor learning

Cholinergic projections are required for VNS-driven cortical map plasticity[15], but the role of cholinergic signaling in VNS-driven motor learning remains unknown. To address this gap, an inhibitory opsin (AAV-EF1a-DIO-eArch3.0-EYFP) was injected into the BF of ChAT-Cre transgenic mice, followed by implanted optical fibers and VNS cuffs (see methods) (**Fig. 4a&b**). Mice then learned to perform the skilled reach task (**Fig. 4c**). One cohort received Success VNS, a second received Success VNS paired with optical inhibition of the BF (Arch+VNS), while the control cohort was unstimulated (**Fig. 4d**). VNS animals (40.55±7.1) performed significantly better than controls (31.23±10.4), while Arch+VNS animals performed at control levels (25.59±9.3) (**Fig. 4e&f**). While all cohorts learned the task (**Fig. 4g**), cholinergic inhibition prevented VNS-driven performance increases in both the early and late (19.02±9.5) (28.66±4.1) learning phases compared to VNS mice (early: 32.81±17.1 L; late: 48.28±11.6) (**Fig. 4g**), demonstrating that phasic cholinergic signaling is necessary for VNS-enhanced motor learning.

VNS reduces off-target failures through increased reach consistency

To understand how VNS may improve reach success rates, we used a behavior apparatus (CLARA) to automate the reaching task, track forelimb kinematics in real-time, and apply VNS following automated classification of reach outcome [40] (**Fig. 5a-c**). Using video data acquired by CLARA, trials were categorized into one of four categories: success; reach failures; grasp failures; and retrieval failures (**Fig. 5d**). Success-VNS mice made fewer reach failure errors than control and Arch+VNS mice (VNS: 54.14±10.4%; control: 72.16±7.0%; Arch+VNS: 71.66±4.5%), and more on-target grasp errors (VNS: 38.19±9.2%, Arch+VNS:

Figure 4 | Outcome-paired VNS motor learning enhancement requires cholinergic neuromodulation.
a, Schematic of VNS implant surgery. **b,** A subset of VNS-implanted *ChAT-Cre* transgenic mice also received injections of viral constructs containing Archaerhodpsin3.0 bilaterally in the BF (right). Mice were chronically implanted with bilateral fiberoptic cannulas for light delivery. **c,** Timeline of experimental set up and training. Each training session lasts for 20 minutes. **d,** Depending on cohort, mice receive VNS, or continuous 532nm light and VNS simultaneously, after successful reach attempts. **e,** Average success rate for all mice over the course of learning(VNS N=11, Arch+VNS N=9, Control N=12). **f,** Mean performance across all days between VNS and control (p=0.0409 (top)), and Arch VNS and control (p>0.05 (bottom)). **g,** Mean success rate of all groups between early and late phases (Control p=0.0001, VNS p=0.0009, Arch VNS p=0.0379). **h,** Mean success rate for control and VNS mice during early (p=0.0458) and late (p=0.0001) learning phases and control and Arch VNS mice (p>0.05).

24.00±4.8%, Control: 22.14±6.0%) (**Fig. 5e,f**). Given the reduction in 'off-target' reach failures, which implies an improved accuracy in reach trajectory, we explored if VNS drives a speed/accuracy trade-off[41]. We measured reach endpoint accuracy and outward reach velocity (see methods) (**Supplementary Fig. 3a&b**), and found that Success VNS does not alter endpoint accuracy or speed of reach attempts (**Supplementary Fig. 3c**), suggesting that it does not influence performance through modulation of either speed or endpoint accuracy.

Motor learning leads to increases in reach consistency towards an expert reach[29,42]. To determine if VNS influences reach consistency, an expert trajectory was defined for each mouse (see methods) (**Fig. 5g**) and 'expert reaches' were identified by having a >0.95 correlation with the expert trajectory. While stimulation does not alter initial reach variance (**Supplementary Fig. 3d**), VNS mice made significantly more 'expert reaches' during late learning compared to control mice (VNS: 45.22±4.4%; control: 34.22±6.6%) while Arch+VNS did not differ from controls (23.58±6.2%) (**Fig. 5i**). Expert reach attempts correlate strongly with behavioral performance (p=0.0001, r^2=0.621), demonstrating that they are a kinematic proxy to task performance (**Fig. 5h**). Likewise, in reach failures, which are not captured in the expert reach population (**Supplementary Fig. 5e**), VNS also increased the similarity of the failures to the expert trajectory during late learning as compared to control mice, while Arch+VNS mice did not differ from controls (VNS: 50.18±16.6; Control: 29.65±19.9; Arch VNS: 33.19±11.6) (**Fig. 5j**). Additional

kinematic features also show a VNS-driven improvement of the feature consistency between early and late

phases (indicated by a narrower histogram with a higher peak) (**Fig. 5k,l**). This data suggests that VNS

decreases off-target reach attempts through improved selection and execution of an expert trajectory late in

learning. Further, cholinergic inhibition prevents VNS-driven changes of kinematic features.

VNS drives acute neural suppression and activation in motor cortex

VNS paired with forelimb movement alters motor cortical map plasticity [24], but the effect of VNS on

neuronal function in motor cortex is unknown. Given that neural activity in mice motor cortex (M1) is

required for both motor skill learning and execution [42,43](**Supp. Fig. 4a**), we hypothesize that VNS will

modulate the neural activity and movement representation in M1. To investigate the effects of VNS in M1

neural activity, we first imaged neural responses to VNS in freely-moving animals expressing calcium

indicator GCaMP6m in M1 using a head-mounted miniature microscope in the homecage (UCLA

miniscope V3, http://miniscope.org, **Fig. 6a**).

In response to VNS, neurons demonstrated either activation (**Fig. 6b**, red, cell 25) or suppression (**Fig. 6b**,

blue, cell 36), without a change in the overall firing rate of the neuron population (**Supp. Fig. 5a&b**).

Approximately 30% of all neurons showed acute response to a VNS delivery, with roughly similar

percentages of neurons showing activation and suppression (activation: $13.8 \pm 5.8\%$; suppression: $18.1 \pm$

15.8%; **Fig. 6d**), and only a small fraction ($0.7 \pm 1.4\%$) show bidirectional modulation (**Fig. 6d**). The

number of neurons modulated by VNS showed a dose-dependence on stimulation intensity (**Fig. 6e**).

We next investigated the temporal dynamics of VNS activation and suppression (**Fig. 6c,f**). The

mean peak timing of the suppression precedes the activation by 1.2s (**Fig. 6g**; suppression: 1.6 s from VNS

onset; activation: 2.8 s), a relationship consistent across a range of behaviorally relevant stimulation

Figure 5 | VNS improves performance through improved consolidation of reach trajectory.
a, The Closed-loop automated reaching apparatus (CLARA) provides 3D tracking of the paw and pellet. **b**, A custom CLARA data pipeline automatically identifies reaching events (one mouse one control session). **c**, CLARA rapidly delivers stimulation after reach end in a closed-loop fashion (180±5ms). Top: duration of all stimulated control trials, yellow dot denotes stimulus delivery. Bottom: a histogram of reach timepoints normalized to reach end.

262 intensities (**Fig. 6h**). Similarly, the onset of suppression preceding activation by 0.6 seconds (**Supp. Fig.**
263 **5c**). These together suggest that in the forelimb region of the mouse primary motor cortex, VNS first drives
264 acute neural suppression, followed by activation in a subpopulation of neurons without altering the mean
265 population firing rate.
266
Success VNS modifies the task-related neural representation of the skilled reach during learning
We examined the influence of Success VNS on movement representation during early learning. The neural
activity in M1 of freely-moving mice implanted with VNS cuff and trained to reach in CLARA was
measured by imaging GCaMP6m in neurons with a head-mounted miniscope. Each reach was subdivided
into a reach and an outcome phase using post hoc analysis (**Fig 7a**). The reach phase includes the outward
paw movement from reach initiation to reach max (~100 ms) and the return movement from reach max to
reach end (to 200 ms). Reach outcome (success or failure) is typically detected to 350 ms after reach max
and ~200ms after reach end. As anticipated [44–47], the average population activity is significantly modulated
during movement (either the reach or outcome phase; **Fig. 7b**). Nearly half of all neurons were movement
modulated (44% of all neurons), with 13.3% of all neurons modulated during reach and 31.7% modulated
during success outcome. For success-outcome-modulated neurons, roughly half are activated and half
suppressed (**Fig. 7c**, success-activated: 16.3 ± 6.4%; success-suppressed: 15.4 ±10.2%). The outcome
representation of success differs from failure, both at the level of the population average response (**Supp.**
Fig. 6 a&b) and individual neural responses (**Supp. Fig. 6 c&d**).
During Success VNS, stimulation is delivered at reach outcome, and so the acute neural response
to VNS is likely to overlap with the intrinsic response to success outcome. To accurately detect VNS-related
neural activity, mice participated in two sessions of training, one with VNS and one without, administered
on a pseudo-randomized schedule (**Fig. 7d**), and the average neural response was compared between VNS
and no-stimulation sessions. During VNS sessions, the success-activated neurons' average response was
first attenuated, then slightly enhanced (**Fig. 7e**). In contrast, success-suppressed (**Fig. 7f**), nonmodulated

Figure 5 | VNS improves performance through improved consolidation of reach trajectory (con't).
d, Example images of the three subcategories of failed reaches (see METHODS). **e**, Breakdown of failure outcomes for each group over 8 days of learning. Light colors: reach failures; intermediate : grasp failures; dark: retrieval failures. **f**, A comparison of types of failed attempts between control and VNS (reach errors: p=0.0005; grasp errors: p=0.0035) and between control and Arch+VNS mice (p>0.05) (VNS N=8, Arch+VNS N=8, Control N=8). **g**, Examples of all outward trajectories (reach initiation – reach maximum) during a session on day 8. Black lines represent each mouse's 'expert reach'. **h**, Correlation of 'expert' reaches and task performance for all mice, r^2=0.62. i, Percent of reaches that are 'expert'. Comparisons were made for the mean 'expert' reaches in the late learning phase (grey box) between control and VNS mice (p=0.0142) and control and Arch+VNS mice (p>0.05) (VNS N=8, Arch+VNS N=6, Control N=8). **j**, Percent of improvement in 'reach failure' toward expert trajectory as measured by increase in correlation coefficient. Comparisons were made during late learning between control and VNS mice (p=0.0455) and control and Arch+VNS mice (p>0.05). **k**, Distribution of trajectory lengths from all failure attempts during early (grey) and late (purple) learning phases. **l**, Normalized improvement in reach features from early to late learning phases.

Figure 6 | VNS drives acute neural suppression and activation in forelimb motor cortex
a, Placement of the grin lens and miniscope above contralateral M1, VNS cuff in the neck and its wire connectors on top back of skull. **b,** Left, representative neural ROIs from one mouse M1 (n = 156 neurons): VNS-activated (red) and suppressed (blue), scale bar = 100 μm. Middle and right: two representative neurons' responses aligned by VNS onset (gray: individual trials; red: VNS-activated; blue: VNS-suppressed). **c,** Top: individual neurons' average responses z-scored to inactive phases of all neurons from the representative mouse in **b**; bottom: average neural responses of all neurons from the same mouse. **d,** Average % of total neurons response after 0.6 to 0.8mA VNS delivery (N = 7 mice, 767 neurons). **e,** % of total neurons that are activated or suppressed by VNS across different current amplitudes. (N = 7 mice, n= 747 to 807 total neurons from each stimulation group, One-way ANOVA and multiple group comparison to 0 to 0.1 mA group, * p < 0.05, ** p < 0.01, *** p < 0.001). **f,** Neural response of all activated and suppressed neurons. **g,** Average neural activity of all activated and suppressed neurons aligned at VNS onset (N=7 mice, 82 activated neurons, 125 suppressed neurons, 0.6mA VNS). **h,** Cumulative distribution of neural response peak time of activated and suppressed neurons measured as the peak value of the average trace 0 to 5 seconds after VNS onset (82 to 151 neurons from each group, Kruskal-Wallis test followed by Dunn's multiple comparisons test, p < 0.0001).

population response (**Fig. 7e**), nearly half of individual neuronal responses were enhanced by VNS (43.3%; **Fig. 7i,k**) and a similar portion were attenuated (40.9%; **Fig. 7j, k**). We next considered if attenuated neural responses preceded the enhanced responses, similar to the temporal dynamics of activation and suppression in the homecage (**Fig. 6g, h**). Indeed, the peak attenuation occurs with similar latency to homecage VNS suppression (1.85 ± 1.20s vs. 1.78 ± 1.10s; **Supp. Fig. 6g**), followed by peak enhancement at a similar latency to peak activation in the homecage context (2.65±1.50s vs.2.74±1.20s, **Supp. Fig. 6g**), demonstrating a similar temporal structure to the VNS response both during and outside of the reach behavior.

Having established that many success-activated neurons are acutely modulated by VNS during success outcome, we next wanted to explore if this neuronal population displays additional activity differences during other phases of the reach. To measure activity across the reach, neural activity was compared to a pre-reach baseline epoch, and differences between the reach representation were compared between the Success VNS and no-stimulation sessions (**Fig. 7l**). Approximately 60% of success-activated VNS-modulated neurons show activity differences onset prior to VNS (**Fig. 7m**). More broadly, during the

Figure 7 | Success VNS selectively modulates activity of task-activated M1 neurons in the reach task.
a, Peri-event histogram of the task related events aligned at reach max (n = 6 control mice, day 4, n = 278 single-reach success trials). Dashed lines indicate median reach initiation (-93 ms), reach max (0 s), reach end (180 ms) and outcome recognition (360 ms). **b**, Average neural activity of success trials (black) and random control trials (grey, n = 488 neurons). Gray dashed line indicates 2 s.d. from the baseline mean. **c**, Left: representative neural responses: red success-activated, blue success-suppressed, gray individual trials. Right: % neurons modulated in the task (903 neurons). **d**, Assignment of VNS or no-stimulation sessions. **e-g**, Top: Average responses of success-activated (d,115 and 101 neurons), success-suppressed (e, 122 and 92 neurons) and success-nonmodulated neurons (f, 383 and 394 neurons) in VNS (orange) and in no-stimulation session (gray). Bottom: the difference trace. **h**, Registered neurons (white) in VNS (green) and no-stimulation session (Magenta). **i&j**, Left: single trial responses in no-stimulation session and in VNS session (orange ticks: VNS onset). Right: the average and difference responses of the same neurons aligned by VNS onset. **k**, Percentage of success-activated neurons (n = 77 registered) that are enhanced (40.9 ± 13.7%) or attenuated (43.3 ± 21.2%) in VNS compared to no-stimulation session. **l**, One success-activated neuron modulated by VNS. Arrowhead: onset of increased activity. **m**, Onset of VNS-driven modulation. **n,** % neurons modulated in reach in VNS-nonmodulated versus VNS-enhanced neurons.

VNS session, all VNS-enhanced neurons are more likely to be active during the reach phase than non-

VNS modulated neurons (38.1 ± 4.7% vs. 25.6 ± 4.9%; **Fig. 7n**). Since VNS alters neural response during

the entire reaching movement, this suggests that VNS effects persist beyond the initial acute modulation

during outcome.

VNS-driven acute neural modulation is mediated through acetylcholine receptors (AChR)

To determine if the acute effects of VNS on neural activity in M1 are mediated through cholinergic

signaling, we injected a systemic acetylcholine receptor antagonist cocktail and measured the acute neural

response to VNS in M1 in the awake, freely-moving animal. The baseline acute VNS response in M1 was

first measured in homecage, then 15 minutes following administration of AChR antagonist cocktail, and

finally in a washout session ~24 hours later (**Fig. 8a**). The percentage of the VNS-modulated neural

population was not changed by AChR antagonism (supp. **Fig. 7**). However, the response amplitudes of

VNS modulated neurons, both activated and suppressed, were reduced by cholinergic antagonism (**Fig. 8b-**

e), demonstrating that AChR mediated signaling is required for VNS-driven acute neural activation and

suppression.

Figure 8 | VNS driven acute neural modulation is mediated through AChRs.
a, Diagram of experimental design: three sessions including a control VNS session in home cage, VNS session with AChR blocker, recovery VNS session on the 2nd day. **b&d,** Average neural activity of VNS-activated neurons (**b,** n = 104 ~116 neurons) or VNS-suppressed neurons (**d,** n = 124~143 neurons) in control VNS session, VNS session with AchR blocker and the 2nd day recovery VNS session. **c,** Average neural activity comparison of VNS-activated neurons quantified from 0.8 to 2.8 seconds after VNS onset, defined based onset and peak delay of VNS-driven neural activation in Figure. 6 (N = 7 mice x two repeats each mouse, Repeated measures ANOVA, p < 0.001). **e,** Average neural activities comparison of VNS suppressed-neurons quantified from 0.2 to 1.6 seconds from VNS onset defined based onset and peak delay of VNS-driven neural suppression in Figure. 6 (N = 7 mice x two repeats each mouse, Repeated measures ANOVA, p = 0.02,).

Discussion

Vagus nerve stimulation paired with rehabilitation is proposed as therapeutic treatment for a wide range of neurologic conditions, yet the mechanism by which VNS may drive neuronal activity to produce changes in behavior remains relatively unexplored. In this study, we establish that VNS optimally enhances motor learning when paired with reach success. VNS drives activity of cholinergic neurons in BF, and optogenetic inhibition of cholinergic neurons is sufficient to eliminate the effects of paired VNS on motor learning, and to reverse VNS-driven kinematic improvements of the reach. Longitudinal in vivo imaging of neurons of the primary motor cortex shows that VNS selectively modulates neurons that represent reach outcome, suggesting a mechanism by which VNS influences reinforcement cues. The effects of VNS on M1 neural activity depend on cholinergic signaling. Together, these results demonstrate that paired-VNS accelerates motor learning through reinforcement, mediated by cholinergic-dependent changes in neural representation and reach kinematics.

VNS paired with reach success optimally enhances motor learning

To our knowledge, we are the first to demonstrate the importance of pairing VNS to movement outcome to enhance motor learning. VNS induces cortical map plasticity in healthy animals when paired with specific movements[24] or auditory tones[48] but has not been shown to improve task performance [24,49]. In contrast, we find that VNS paired with a successful reach, but not reach initiation, enhanced motor learning and ultimate performance. The enhanced motor learning may be due to the pairing of VNS stimulation with a successful task outcome, perhaps to augment task reward or reinforcement. Endogenous activity of the vagal nerve has been linked to reward and motivation[50], and VNS in human subjects drives motivation towards reward [51] and improves reinforcement learning[52]. However, VNS does not seem to activate classical dopaminergic reward pathway [32] as conditioned place preference test results indicate that VNS is not inherently rewarding of aversive. Instead, VNS may augment reinforcement cues, leading to improved selection of successful trajectory[53,54] and the associated neural ensembles that underlie those movements[55–57]. By augmenting reinforcement cues, VNS may help to select the appropriate neural circuits, strengthening those connections for lasting improvements in functional outcome.

The importance of VNS timing appears to differ across the learning process. During early learning, Success VNS has a unique ability to improve reach learning. Moreover, Random VNS delivered during early learning impairs reach learning, suggesting that poorly timed VNS can disrupt normal learning processes. During late learning, Success VNS maintains improved performance over control animals, while the subset of stimulated trials in Reach VNS also show performance improvements over controls (but this improvement is abolished when all trials are included). In expert animals, both Success and Reach VNS provide short-term improvements over unstimulated blocks of trials. Thus, it appears critical that VNS be

paired with reach outcome during early learning, but there is more flexibility in the timing of VNS pairing during the rehearsal of a known task.

How might reinforcement-paired VNS contribute to skilled motor learning? Motor learning travels along an exploration-exploitation axis, with early exploration, expressed as motor variability, reducing as the motor behavior consolidates onto an expert solution [58–60]. Moreover, early learning motor variability predicts improved later performance of the expert motor solution [54,61,62], pointing to a reinforcement learning process by which increased exploration allows for faster identification of the expert solution. Reward is also known to influence the exploration-exploitation relationship, with conditions of increased reward frequency leading to reduced variability and increased consolidation onto a motor solution [53,60,63]. Our results show that paired-VNS does not increase variability in early learning, but instead improves kinematic consolidation onto an expert reach. Perhaps, in this context, VNS acts to reinforce the reward salience, increasing the exploitation of the expert solution. Future experiments are needed to explore if VNS can reinforce movements in the without association to a food reward.

VNS-driven motor learning is mediated by cholinergic signaling

VNS activates multiple neuromodulatory systems in the central nervous system[16,64,65], and the effects of VNS on cortical plasticity are mitigated with lesions of neuromodulatory nuclei including the locus coeruleus, raphe nucleus and the cholinergic basal forebrain[15,16]. While each of these neuromodulators could play a role in motor learning, cholinergic neuromodulatory systems are critical for use-dependent plasticity[34,66–69], are most closely associated with reinforcement signaling[18,33], and encode task outcome[19,70]. Lesions[17,71] or pharmacological suppression of cholinergic neurons[72] are detrimental to motor learning and VNS-enhanced motor rehabilitation[9]. Leveraging the temporal precision of optogenetic suppression of cholinergic neurons, we find that a brief cholinergic inhibition is sufficient to prevent VNS-driven enhancement in motor learning. This suggests that the effects of Success VNS are mediated through cholinergic signaling in the basal forebrain, and is consistent with the role of the cholinergic BF to encode outcome salience[18,19,70].

Only limited evidence exists demonstrating an anatomical or functional connection between the vagus nerve and the cholinergic basal forebrain[73]. Using optotagging approaches, we were able to demonstrate robust functional connectivity between the BF and the vagus nerve. Stimulation of the vagus nerve elicited robust responses in nearly half of the cholinergic and non-cholinergic units recorded under anesthesia and more than 40% of the units recorded in awake animals. The variable timing of BF neuronal responses to VNS suggests the involvement of a multi-synaptic pathway, possibly through the locus coeruleus (LC), which is known to send direct projections to the BF[74–76] and is activated by VNS[13,77].

394   Additionally, VNS leads to the sequential activation of noradrenergic and cholinergic axons in cortex, with
395   noradrenergic activity preceding cholinergic activity[14].
396
Motor cortical neurons are modulated by VNS via cholinergic activity
Neurons in the primary motor cortex represent movement preparation and execution for dexterous
movements [43,78,79]. During motor learning, these neural representations are updated to improve motor output
[46,80,81] by incorporating feedback from error and reinforcement signals generated throughout multiple
regions of the central nervous system, including cortical regions [55,82–84]. Recent work demonstrated that, in
addition to movement preparation and execution, primary motor cortex (M1) pyramidal neurons also report
movement outcome. Neurons in superficial L2/3 of M1 represent success and failure, independent of
kinematics and the food reward consumption [47]. Our data demonstrate that Success VNS attenuates the
population representation of a success outcome by selectively modulating success-outcome responsive
neurons. This same neural population is also more likely to have different representation of movement
preparation and reach execution as well, suggesting that VNS changes neural activity beyond the acute
response to stimulation. The specificity of the population of neurons that are modulated by Success VNS
may indicate that VNS adds selectivity to outcome representation, to optimize outcome signals for enhanced
learning.
The relative selectivity of the effects of Success VNS on movement representation are somewhat
in contrast to recent observations of widespread, long-lasting excitatory responses to VNS [14]. However, in
the previous study, in head-fixed mice VNS elicited locomotion and whisking, both of which correlate to
increased general arousal and widespread cortical activation [85–87], making it difficult to disentangle direct
VNS effects from changes in arousal. In contrast, another recent work demonstrated a VNS-driven
suppression of neural response to an auditory tone that persisted even after arousal state was regressed from
the neural response [88]. Our trial design controls for movement state, as the comparison between the VNS
trial and no-stimulation trial occurs at the exact same reach position, and VNS did not produce any
noticeable acute motor response. By eliminating the confound of behavioral states such as locomotion or
quiet resting, we can detect that VNS produces an initial suppression of the outcome response followed by
excitation. This effect is seen only in neurons that respond to outcome, indicating that when applied during
a reach, VNS acts on a specific population of neurons that are already engaged in the representation of
reach outcome.
424
Mechanisms of other neurotechnology during motor learning and performance
Motor learning and performance have been modulated utilizing non-invasive stimulation devices including
transcranial stimulation (ie. tCDS, TRMS) and transcranial magnetic stimulation (TMS). While both tCDS

and TMS have relatively unknown mechanisms, they are postulated to operate by inducing polarity changes

in cortical neurons altering their excitability[89–91] and promote BDNF-mediated synaptic plasticity[92]. Still,

the underlying mechanisms and optimal stimulation protocols are a current area of exploration[93]. For

example, some tCDS studies have shown improved reaction times in motor sequence tasks[89,94], yet others

have reported no effect on reaction time or performance [95,96]. As with VNS, further exploration of the

mechanisms of action of non-invasive stimulation modalities, along with optimized stimulation protocols,

is needed to address specific therapeutic needs across a wide range of neurologic disorders.

Conclusion

In conclusion, we demonstrated that VNS augments reinforcement cues to accelerate skilled motor learning

and enhance kinematic consolidation on an optimal motor plan in healthy animals. VNS alters neural coding

of outcome in a select neural population, and changes neuronal activity of this population across the entire

reach. The behavioral, kinematic and motor representation effects of VNS are mediated by phasic

cholinergic activity. Understanding the behavioral and circuit mechanisms of VNS allows for future

optimization of rehabilitation protocols and new avenues for the use of cholinergic manipulation to treat

neurologic conditions.

Figure 1 | VNS modulates forelimb reach learning and requires temporally specific stimulation. a-c,

Illustration of VNS surgery, behavior timeline, and stimulation protocol. **d-f,** Random VNS, Reach VNS,

and Success VNS success rate across 14 sessions of training. **g,** Comparison of mean performance across
all days between control and stimulated groups (Success VNS: $p=0.0065$, $f=9.24$, Random VNS: $p>0.05$,
Reach VNS: $p>0.05$, Student T test). **h,** Comparison of mean success rate for control and Random VNS
mice during early ($p=0.028$, $f=7.07$, Student T test) and late learning ($p>0.05$). **i,** Comparison of mean
success rate for control and Reach VNS mice during early and late learning ($p>0.05$). **j,** Comparison of
mean success rate for control and Success VNS mice during early ($p=0.0031$) and late learning ($p=0.0126$).
k,l, in-session learning trajectories for each group. **m,** Comparison of within session learning between all
groups across 4 days of learning. **n, o,** VNS mice performed a conditioned place preference test after 3 days
being stimulated in one of two distinct rooms. $*p<0.05$, $**p<0.01$, $***p<0.001$ bars and error bars
represent the mean $\pm$ s.e.m.
460
461

Figure 2 | VNS improves success rate within sessions and in learned mouse during rehearsal of
forelimb reach task. a, trials before and after a stimulated success are investigated in success VNS **b,**
Comparison of success rate for reaches preceding or following stimulated reach. **c,** For Success VNS, day
7 and 14 trials are divided into equal blocks of unstimulated and stimulated trials. **d,** Comparison of
unstimulated (light orange) and sham (grey) trials on day 14 (p=0.0059, t=3.34, Student's T test). **e&f,**
Comparison of stimulated (dark orange) and unstimulated (light orange) trials on days 7 & 14 (Day 7:
p>0.05; Day 14: p=0.0499, t=2.57, Ratio paired t test). **g,** Reach VNS schematic **h,** Success rates across
learning training sessions for Sham (grey), Stimulated Reach VNS (dark green), and Unstimulated Reach
VNS (light green). **i,** Comparison of stimulated Reach VNS and Sham VNS trials in early (p>0,05) and late
learning (p=0.024, t=2.47, Student's T test). **j,** Comparison between stimulated Reach VNS and
unstimulated Reach VNS trails in early (p=0.004, t=3.98, paired t test) and late learning (p<0.0001, t=10.08,

paired t test). **k,** Success VNS and Reach VNS applied during rehearsal of reach task in naively trained
mice. **l,** Stimulated Reach VNS trials improve success rate during rehearsal (p=0.015, t=4.058, paired t
test). **m,** Stimulated Success VNS trials improve success rate during rehearsal (p=0.005, t=5.62, paired t
test). **n,** Normalized improvement of stimulated Reach VNS (p=0.047, t=3.56) and stimulated Success VNS
trials (p=0.028, t=4.16) compared to unstimulated trials (Sidak's multiple comparison's test in RM one-
way ANOVA). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; bars and error bars represent the mean ± s.e.m.
481
482

Figure 3 | VNS drives BF neural activity in anesthetized and awake mice.

a, Tetrodes were implanted in the left BF of mice and recordings were obtained during awake behavior. **b,**

Example raster (top) and average firing rate (bottom from an 'activated' neuron) in response to VNS. Grey

box denotes stimulus delivery. **c,** Heatmap showing the average responses of all recorded neurons to VNS

(grey box). White represents increased activity; black represents decreased activity. **d,** Percent of neurons

that respond to VNS (N = 5 mice, 53 neurons). **e,** Average activity of all 'activated' neurons in response to

VNS. Dashed lines mark significantly altered activity, shading represents SE. **f,** Extracellular recordings

were obtained using optrodes while under light anesthesia. **g,** Optrodes were targeted at the HDB in *ChAT-*

ChR2 transgenic mice. Green fluorescence denotes the presence of ChR2. **h,** Example cholinergic neuron

responding consistently to pulses of 488 nm light. **i,** Average activity of all cholinergic neurons during opto-

tagging. Each row represents a neuron. **j,** Stimulus-associated latency tests (SALT) separate light

responsive neurons from non-light responsive neurons. **k,** Mean baseline FR of cholinergic and non-

cholinergic neurons (p=0.013, N = 5 mice, 53 neurons). **l,** Percent of neurons categorized as cholinergic

(light green) and non-cholinergic (dark green) (N = 6 mice, 76 neurons). **m,** Percent of units that are VNS-

responsive in cholinergic (left) and noncholinergic (right) populations. **n,** Average response to VNS for all

'activated' neurons. **o,** Mean peak activation during VNS in cholinergic, non-cholinergic and awake-

recorded(red) neurons. **p,** Average delay of peak activation from VNS onset cholinergic, non-cholinergic

and awake-recorded(red) neurons. **q,** Mean duration of significantly elevated activity after VNS

(cholinergic vs awake p=0.0087, non-cholinergic vs awake p=0.0003).

Figure 4 | Outcome-paired VNS motor learning enhancement requires cholinergic neuromodulation.
a, Schematic of VNS implant surgery. **b,** A subset of VNS-implanted *ChAT-Cre* transgenic mice also
received injections of viral constructs containing Archaerhodpsin3.0 bilaterally in the BF (right). Mice were
chronically implanted with bilateral fiberoptic cannulas for light delivery (left). **c,** Timeline of experimental
set up and training. Each training session lasts for 20 minutes. **d,** Depending on cohort, mice receive VNS,
or continuous 532nm light and VNS simultaneously, after successful reach attempts. **e,** Average success
rate for all mice over the course of learning (VNS N=11, Arch+VNS N=9, Control N=12). **f,** Comparison
of mean performance across all days between VNS and control (p=0.0409 (top)), and Arch VNS and control

515   (p>0.05 (bottom)). **g,** Comparison of mean success rate of all groups between early and late phases (Control

516   p=0.0001, VNS p=0.0009, Arch VNS p=0.0379). H). Comparison of mean success rate for control and

517   VNS mice during early (p=0.0458) and late (p=0.0001) learning phases and control and Arch VNS mice

518   (p>0.05).

Figure 5 | VNS improves performance through improved consolidation of reach trajectory.

a, The Closed-loop automated reaching apparatus (CLARA) provides 3D tracking of the paw and pellet. **b,** A custom CLARA data pipeline automatically identifies reaching events (one mouse one control session). **c**, CLARA rapidly delivers stimulation after reach end in a closed-loop fashion (180±5ms). Top: duration of all stimulated control trials, yellow dot denotes stimulus delivery. Bottom: a histogram of reach timepoints normalized to reach end. **d**, Example images of the three subcategories of failed reaches (see METHODS). **e**, Breakdown of failure outcomes for each group over 8 days of learning. Light colors: reach failures; intermediate: grasp failures; dark: retrieval failures. **f**, A comparison of types of failed attempts between control and VNS (reach errors: p=0.0005; grasp errors: p=0.0035) and between control and Arch+VNS mice (p>0.05) (VNS N=8, Arch+VNS N=8, Control N=8). **g**, Examples of all outward trajectories (reach initiation – reach maximum) during a session on day 8. Black lines represent each mouse's 'expert reach'. **h**, Correlation of 'expert' reaches and task performance for all mice, r^2=0.62. i, Percent of reaches that are 'expert'. Comparisons were made for the mean 'expert' reaches in the late learning phase (grey box) between control and VNS mice (p=0.0142) and control and Arch+VNS mice (p>0.05) (VNS N=8, Arch+VNS N=6, Control N=8). **j**, Percent of improvement in 'reach failure' toward expert trajectory as measured by increase in correlation coefficient. Comparisons were made during late learning between control and VNS mice (p=0.0455) and control and Arch+VNS mice (p>0.05). **k**, Distribution of trajectory lengths from all failure attempts during early (grey) and late (purple) learning phases. **l**, Normalized improvement in reach features from early to late learning phases.

546 Figure 6 | VNS drives acute neural suppression and activation in forelimb motor cortex

a, Placement of the grin lens and miniscope above contralateral M1, VNS curf in the neck and its wire

connectors on top back of skull. **b,** Left, representative neural ROIs from the field of view of one mouse

M1 (n = 156 neurons): neurons are pseudo color labeled as VNS-activated (red) and suppressed (blue),

scale bar = 100 μm. Middle and right: two representative neurons' Ca responses aligned by VNS onset

(gray: individual trials; red: VNS-activated; blue: VNS-suppressed). **c,** Top: individual neurons' average

response z-scored to inactive phases of all neurons from the representative mouse in **b**; bottom: average

neural responses of all neurons from the same mouse. **d,** Average % of total neurons activated, suppressed,

bidirectionally modulated and nonmodulated after 0.6 to 0.8mA VNS delivery (N = 7 mice, 767 neurons).

e, % of total neurons that are activated or suppressed by VNS across different current amplitudes. (N = 7

mice, n= 747~807 total neurons from each stimulation group, One-way ANOVA and multiple group

comparison to 0~0.1 mA group, * p < 0.05, ** p < 0.01, *** p < 0.001). **f,** Neural response heatmap of all

activated neurons and all suppressed neurons aligned at VNS onset. **g,** Average neural activity of all

activated neurons and all suppressed neurons aligned at VNS onset (N=7 mice, 82 activated neurons, 125

suppressed neurons, 0.6mA VNS). **h,** Cumulative distribution of neural response peak time of activated

and suppressed neurons measured as the peak value of the average trace 0 to 5 seconds after VNS onset (82

to 151 neurons from each group, Kruskal-Wallis test followed by Dunn's multiple comparisons test, p <

0.0001).

Figure 7 | Success VNS selectively modulates activities of a subpopulation of task-activated M1 neurons in the reach task.

a, Peri-event histogram of the task related events aligned at reach max (N = 6 control mice, day 4, n = 278 single-reach success trials). Dashed lines indicate median reach initiation -93 ms, reach max 0 s, reach end 180 ms and CLARA success recognition 360 ms. Magenta indicates full reaches, green success recognition. **b**, Average neural activity of success trials (black) and random control trials (grey, n = 488 neurons). Gray dashed line indicates 2 s.d. from the baseline mean. c, Left: representative neural responses: red success-activated, blue success-suppressed, gray individual trials. Right: % neurons modulated in the task (903 neurons; 16.3 ± 6.4% success-activated, 15.4 ± 10.2% success-suppressed, 13.3 ± 5.5% preparation/reach modulated). **d**, Assignment of VNS or no-stimulation sessions. **e-g**, Top: Average responses of success-activated (d,115 and 101 neurons), success-suppressed (e, 122 and 92 neurons) and success-nonmodulated neurons (f, 383 and 394 neurons) in VNS (orange) and in no-stimulation session (gray). Bottom: the difference trace. **h**, Registered neurons (white) in VNS (green) and no-stimulation session (Magenta). **i&j**, Left: trial responses of success-activated neurons in no-stimulation session and in VNS session (orange ticks: trials' VNS onset). Right: the average and difference responses of the same neurons aligned by VNS onset. **k**, Percentage of success-activated neurons (n = 77 registered) that are enhanced (40.9 ± 13.7%) or attenuated (43.3 ± 21.2%) in VNS compared to no-stimulation session. **l**, One success-activated neuron modulated by VNS (supp. Fig. 6h) also have higher neural activity before reach end in VNS session. Arrowhead: onset of increased activity. **m**, Onset timing histogram of VNS-driven modulation of success-activated neurons. n, % neurons modulated in reach in VNS-nonmodulated versus VNS-enhanced neurons.

Figure 8 | VNS driven acute neural modulation is mediated through AChRs.

a, Diagram of experimental design: three sessions including a control VNS session in home cage, VNS session with AChR blocker, recovery VNS session on the 2nd day. **b&d,** Average neural activity of VNS-activated neurons (**b,** n = 104 ~116 neurons) or VNS-suppressed neurons (**d,** n = 124~143 neurons) in control VNS session, VNS session with AchR blocker and the 2nd day recovery VNS session. **c,** Average neural activity comparison of VNS-activated neurons quantified from 0.8 to 2.8 seconds after VNS onset, defined based onset and peak delay of VNS-driven neural activation in Figure. 6 (N = 7 mice x two repeats each mouse, Repeated measures ANOVA, p < 0.001). **e,** Average neural activities comparison of VNS suppressed-neurons quantified from 0.2 to 1.6 seconds from VNS onset defined based onset and peak delay of VNS-driven neural suppression in Figure. 6 (N = 7 mice x two repeats each mouse, Repeated measures ANOVA, p = 0.02,).

Supplementary Figure 1 | Stimulation Amount between Random VNS, Reach VNS, and Success VNS does not predict success.

a-c, a simple linear regression was calculated to predict success rate based on the amount of stimulation trains given across groups in a particular session. **a,** For Random VNS a regression equation was found $(F(1.03)\ p = 0.3117)$, with an R^2 of 0.009. **b,** For Reach VNS a regression equation was found $(F(2.017)\ p = 0.22)$, with an R^2 of 0.011 **c,** For Success VNS a regression equation was found $(F(0.11)\ p = 0.7359.)$, with an R^2 of 0.001.

Supplementary Figure 2 | All experiment groups learn the behavior task.

a, nonlinear model of sham animal learning trajectories used to identify early and late learning phases (Methods). **b-e,** Comparison of performance during early and late learning phases in Sham VNS (paired T test, n=12, p=0.0001), Random VNS (paired T test, n=8, p=0.0001), Reach VNS (paired T test, n=9, p=0.0002), and Success VNS (paired T test, n=10, p=0.001).

Supplemental Figure 3 | VNS does not change speed, accuracy or initial exploration.

a, Endpoint targeting of all control reaches on day 1 (grey) and day 8 (purple) of learning. Orange dot depicts pellet center. **b**, Mean absolute velocity between reach initiation and reach maximum on day 1 and day 8 of learning. Reaches are time warped to be an equal arbitrary length. **c,** Comparison of endpoint accuracy (top) and absolute velocity (bottom) between control, VNS and Arch+VNS stimulation (p>0.05, RM ANOVA, VNS N=8, Arch+VNS N=6, Control N=8). **d,** Reach variability on day 1 for all groups, as measured by average correlation to expert trajectory (p>0.05, One-way ANOVA, VNS N=8, Arch+VNS N=6, Control N=8). **e,** Percent of 'Reach Failures' that qualify as expert reaches over learning.

Supplementary Figure 4 | The M1 activity is necessary for successful execution of the skilled reach task.

a, Comparison of success rate of the skilled reach task before and after muscimol infusion into M1 (n = 3 mice, one-way ANOVA, p<0.05). **b,** Comparison of reach attempts per second in the same mice before and after muscimol infusion (n = 3 mice, one-way ANOVA, control ~0.3 reaches per minute, muscimol 1.1 reaches, p<0.05).

Supplementary Figure 5 | The M1 activity is constant on minutes' time scale and the VNS response onset timing analysis.

a, Heatmap plot of 40 neurons' GCaMP6m calcium response in L2/3 M1 from a representative mouse in homecage two minutes before VNS delivery and two minutes with VNS trains delivery. Each orange arrow indicates one VNS pulse train delivery (15 pulses at 0.1 ms duration, 30 Hz, 0.4 mA). **b,** Ca transients quantified as area under the curve (AUC) of the fluorescence trace per minute in the 2nd minute after VNS application starts in VNS mice (n = 7 mice, 676 to 732 neurons, plot as median with interquantile range. Kruskal -Wallis test followed by Dunn's multiple comparison, p > 0.05 for all VNS stimulation groups compared to control. Controls are the same mice without VNS delivery). **c,** Cumulative distribution of neural response onset (measured as the first value goes above 2SD of the baseline mean in the average trace 0~5 seconds after VNS onset) of VNS-activated or suppressed neurons (82 to 151 neurons from each group, 7 mice, Kruskal-Wallis test followed by Dunn's multiple comparisons test, p < 0.0001).

a

Success trials   Failure trials   Swapped trials   ΔZ
Time from reachMax (s) Time from reachMax (s) Time from reachMax (s)

b
success
failure
ReachMax
0.1 ΔF
1s

c

Success-activated neurons   Failure-activated neurons   Universal outcome-activated neurons   ΔZ

Success trials

Failure trials

2 ΔZ
2 s

ReachMax   ReachMax   ReachMax

d

Total neurons

11.6%   4.6%   5.2%

Success-activated neurons
Failure-activated neurons
Universal outcome-activated neurons

e

Success-activated   Success-suppressed   Success-nonmodulated

% of all neurons   % of all neurons   % of all neurons

No VNS   VNS   No VNS   VNS   No VNS   VNS

f
Failure-activated

0 -

0.2 ΔZ
2 s

Failure outcome

g

VNS

2 ΔZ
1 s

h

Peak timing of VNS driven modulation
in success activated neurons

% of neurons

**

VNS enhanced success activated neurons
VNS enhanced in homecage
VNS attenuated success activated neurons
VNS attenuated in homecage

Time from VNS onset (s)

Supplementary Figure 6 | The M1 neural dynamics in the reach task.

a, Heatmap plot of individual neurons' response sorted by the peak timing: reach max is 0 second (N = 6 mice, n = 488 neurons). **b,** Average neural activity of all neurons in single-reach success trials (black line), and single-reach failure trials (dotted line). Gray block indicates 2 SD from the baseline mean during -6 to -3 s before reach max. **c,** Heatmap plot of trials and average response of representative success-activated neuron, failure-activated neuron and universal outome-activated neuron aligned by reach max. **d,** % of success activated-neurons, failure-activated neurons and universal outome-activated neurons (1 mice, 172 neurons). **e,** The percentage of success-activated, -suppressed and non-modulated neurons in VNS sessions and in no-stimulation sessions (N = 6 mice, total 488 neurons). **f,** Average neural activity of failure-activated neurons in VNS session (orange, 77 neurons) and in no-stimulation session (gray, 73 neurons), aligned at outcome recognition by CLARA. **g,** Cumulative distribution of neural response peak time of VNS enhancement or attenuation of outcome-activated neurons' outcome response. The VNS driven enhancement or attenuation are measured as the peak value of the difference trace (individual neuron's VNS session average – no-stimulation session average) 0~5 seconds after VNS onset (51~54 neurons from each group, 6 mice, Kruskal-Wallis test, $p < 0.0001$). **h,** The response to VNS of representative success-activated neuron in fig. 7l aligned at VNS and normalized to the -3 ~0 seconds before VNS.

Supplementary Figure 7 | The neural populations analysis with AChR blocker.

a, Percentage of VNS-activated neurons and VNS-suppressed neurons in the total neuron populations after 0.4~0.6 mA VNS delivery in control VNS session, VNS session with AChR blocker, recovery VNS session 2nd day (N = 7 mice. total 627 neurons, Brown-Forsythe ANOVA test, $p > 0.5$)

Methods

Animal care

All animal procedures and experiments were performed in accordance with protocols approved by the Institutional Animal Care and Use Committee at the University of Colorado Anschutz Medical Campus. Male and female adult C57BL/6 wild-type mice between the age of 2 and 10 months old were used for all experiments unless otherwise noted. Mice were group-housed before surgery and single-housed following surgery and throughout behavior training. Mice were kept on 14h light/10h dark schedule with ad libitum access to food and water with exception from behavior-related food restriction (Forelimb Reach training).

Surgery

Vagus nerve stimulating cuff implantations

Commercial cuffs from Micro-leads (150um cuffs) and Cortec (100 um microsling cuffs) soldered to gold pins or Plastics1 connectors were implanted on the cervical vagus nerve. Mice were anesthetized with 4.5% isoflurane anesthesia for induction and maintained with 1.5%. 1% injectable lidocaine was used locally at incision sites. Eye ointment was applied to the eye to prevent corneal drying. Temperature was regulated at 37ºC with a thermostat-controlled heating pad. The vagus nerve was accessed with an incision in the ventral cervical region, and the nerve was bluntly dissected from the carotid sheath. The cuff was tunneled subcutaneously to the ventral cervical incision from an incision at the base of the dorsal skull. The vagus nerve was placed in the cuff. Kwik-Sil (WPI)The ventral cervical incision was sutured using 6-0 absorbable sutures. The dorsal skull was cleaned using saline and ethanol, and electrical connectors were fixed to the skull using dental cement (C&B Metabond). GLUture (WPI) was used to seal the skin around the dental cemented headcap. Stimulation efficacy was measured using peripheral biomarkers such as breathing rate changes and heart rate reduction[25] on the day of surgery and weekly until the end of experiments with paw sensor (Mouse Stat Jr., Kent Scientific). Mice received sub-cutaneous lactated ringers (~100μL as necessary), intramuscular gentamicin (3mg/kg), and intraperitoneal meloxicam (5mg/kg) following surgery and as needed in cases of dehydration, infection, or pain. Mice were monitored for 7-14 days to ensure proper recovery from surgery before any subsequent experiments were conducted.

Viral injections and optical fiber implantation

All surgeries were performed on mice that express Cre recombinase driven by the ChAT promoter (ChAT-IRES-Cre, Jackson Labs stock #0064100 [97]). Animals were prepared for surgery as described above, and the hair was removed prior to an incision over the dorsal skull. 200um diameter fiber optics with 1cm ceramic cannulas were fabricated in-house using ONECore facilities. The skull at the dorsal incision was cleaned using sterile saline and ethanol. Two craniotomies were opened above the basal forebrain in each

hemisphere (0.35mm posterior, ±1.6mm lateral of bregma) using a dental drill. Glass pipettes containing a

floxed inhibitory archaerhodopsin (AAV-EF1a-DIO-eArch3.0-EYFP, UNC) or YFP control (AAV-EF1a-

DIO-EYFP-WPRE-pA, UNC) were then inserted bilaterally into the basal forebrain using a stereotaxic

device (-4.75mm from dorsal surface of the brain). Approx. 210nL of viral construct was injected over 5

minutes. The pipettes were removed and 200um fiber optic cannulas were inserted above each injection site

(-4.65mm from the dorsal surface of the brain). The craniotomies were then sealed using a surgical silicone

(Kwik-Sil). The cannulas were fixed to the skull using dental cement (C&B Metabond). Gluture was used

to seal the skin around the headcap. Mice received intramuscular gentamicin (3mg/kg), and intraperitoneal

meloxicam (5mg/kg) following surgery and as needed in cases of infection, or pain. Mice were monitored

for 5-7 days to ensure proper recovery. After 14 days, mice underwent VNS implantation described above

(see *Vagus nerve stimulating cuff implantations*).

Electrode implantation

<u>Chronic tetrodes:</u> Custom twisted wire tetrodes, built in-house, were implanted into the left basal forebrain

of mice. Surgical preparation was as described above. Then, two craniotomies were opened using a dental

drill; one above the left basal forebrain (0.35mm posterior, 1.6mm lateral of bregma) and one above the

cerebellum (~1mm posterior of lambda, midline). A tetrode was then inserted into the basal forebrain

craniotomy using a stereotaxic device (-4.75mm from the dorsal surface of the brain). A gold ground pin

was inserted into the cerebellum. Both craniotomies were sealed with surgical silicone (Kwik-Sil) and the

tetrode was fixed to the skull using dental cement C&B Metabond). Gluture was used to seal the skin around

the dental cemented headcap. Mice received intramuscular gentamicin (3mg/kg), and intraperitoneal

meloxicam (5mg/kg) following surgery and as needed in cases of infection, or pain. Mice were monitored

for 7 days to ensure proper recovery after which they underwent VNS implantation described above (see

Vagus nerve stimulating cuff implantations).

<u>Acute Optrodes:</u> Vagus nerve cuffs were implanted in transgenic mice expressing Channelrhodopsin2 in

cholinergic neurons (ChAT-ChR2 [98]) using the protocol described above (see *Vagus nerve stimulating cuff*

implantations). After cuff implantation, while mice were still under anesthesia (1.5% isoflurane), mice were

moved to a stereotaxic apparatus and a cranial window (2.5x2.5mm) was opened above the left BF (0.35mm

posterior, 1.6mm lateral of bregma). The stereotaxic apparatus was placed in a Faraday cage and single

shank Optrodes from Neuronexus (A1x32-Edge-10mm-20-177-OA32LP) were inserted into the BF (-

4.75mm from the dorsal surface of the brain). Extracellular recordings were then performed while mice

then underwent VNS (see *BF response to VNS*) and Opto-tagging (see *Opto-tagging*) protocols. Mice were

sacrificed at the end of the experiment and the cuff and optrode were recovered.

Cranial window surgery for miniscope objective lens and baseplate installation

Mice were anesthetized with isoflurane and maintained similarly as described above until skull is exposed. A round cranial window (~1.8mm diameter, ML 1.5mm, AP 0.3 mm for center) was made on contralateral M1 using a dental drill. A viral vector (AAV1.Syn.GCaMP6m.WPRE.SV40) was infused at $2x10^{12}$ titer to 200 ~300 um beneath brain surface at 3~4 sites in the cranial window around the center, with ~200 nl each site. Objective grin lenses (Edmund Optics, #64520, 1.8mm, 0.23mm WD) were lowered through the cranial window and pressed against the brain surface. The lens' side was sealed by surgical silicone (Kwik-Sil) and secured by dental cement. The exposed part of the lens above the skull was further coated with black nail polish. 3~4 weeks later, a metal baseplate was mounted to the skull over the lens with loctite glue (loctite 454 prism), guided by miniscope for optimal field of view while the mouse were anesthetized with isoflurane (1.5%). After the baseplate were securely mounted, the miniscope was taken off, a cap was attached to the baseplate and the mouse was returned to homecage.

Muscimol cannula surgery and infusion

With the same craniotomy surgery procedure described above, a round cranial window (~1.8 mm diameter, ML 1.5mm, AP 0.3 mm for center) was made on contralateral M1 and a plastic cannula (2~3 mm long pipet tip) was inserted into the window right above the brain surface. Then the pipet tip was secured by kwik-sil and dental cement. The top of the cannula will be sealed with kwik-sil if the mouse is not going to the behavior test in a couple of hours. Before behavior tests, the mice were briefly anesthetized with isofluorane the seal will be removed and 1~2 μL of muscimol at a concentration of 1 mg/mL will be infused into the pipet tip.

Behavior

Manual training of skilled forelimb reach task

Mice were trained and scored on a skilled reach task [99]. Mice were food restricted and maintained at 85-90% of their weight throughout training. Following food restriction, mice were habituated to the training box for 20 minutes where mice were given 20mg food pellets (BIO-serv) near the window of the box where reaching occurs. The training box is a custom-built plexiglass box with a 1cm wide opening that provides access to a platform or post with a divot to hold a pellet located approximately 1cm away with a left offset from the center of the opening to force right forepaw reaching. Learning sessions then occurred for 14 consecutive days where mice perform a reach to grasp task with the right forepaw for food pellets. Rehearsal sessions occurred 7-10 days after training and featured stimulated and unstimulated trials. For both training and rehearsal sessions, mice were scored on a per trial basis until 20 successful attempts or 20 minutes

passed. A trial terminates in a success or the pellet being knocked off the pellet holder by the mouse. Trial
outcomes were recorded by the trainer in real time. A success was defined by when the mouse grabbed the
pellet and returned it into the cage. Errors were subcategorized into: "reach error" (failure to correctly
target), "grasp error" (failure to grasp the pellet), and "retrieval error" (successful grasp of pellet, but failure
to return it into the box).

VNS experiment groups

Experiment groups for training were based on stimulation protocols. Sham VNS cohort were implanted
with a cuff but were not stimulated at any point during learning. Success VNS were manually stimulated
following every successful trial -- days 7 and 14 featured designated blocks without stimulation to track
baseline learning levels versus trial-to-trial performance. Reach VNS received manual stimulation prior to
movement onset on a pseudo-random 50% subset of trials to normalize VNS trains delivered. Random VNS
received stimulation at random intervals as generated by an Arduino board to achieve between 15-20
stimulations per day, matching stimulation rates to other groups. All groups were trained on the forelimb
reach task for 14 days, Reach VNS and Success VNS were tested during rehearsal in naively trained mice.

CLARA training and behavior data acquisition

Mice were food restricted and habituated identically to manually trained animals. Dimensions of the
behavior box were also identical in manual and CLARA cohorts. Behavior box used for miniscope
recording was modified so that the front panel had an alcove above the height of mouse head to
accommodate the miniscope when the mouse was close to the slit to reach. On day 1, the mice were primed
to have one success before CLARA training session started. Learning sessions then occurred for 14
consecutive days (or specified otherwise in results) where mice perform a reach to grasp task with the right
forepaw for food pellets. Each trial started as the automated dispenser placing a food pellet on the post, as
the mouse reached to successfully retrieve it or knocked it off, the CLARA would mark success or failure
as the trial outcome as the end of this trial. Each session lasted for 20 minutes and mice were scored on a
per attempt basis. Using the CLARA behavior system, high speed (150hz) video data was recorded from
three FLIR Blackfly® S (model BFS-U3-16S2M-CS, Edmund Optics) cameras placed in front of the box,
lateral to the box, and at 45° angle above the box from the opposite side from the lateral camera (**Fig. 6a**).
A neural network was trained prior to experiment sessions using manually annotated frames of the skilled
reach behavior labeling the hand center and the pellet. Video frames from all cameras we sent through this
network in real time to identify the location and state of the hand and pellet. This information was used
initiate trials via pellet placement and to categorize attempt outcomes as either success or failure so that
stimulation could be delivered in a closed-loop manner (for additional details, see [40]). The timing of pellet

placement, success or failure outcome, VNS delivery, and optogenetic light delivery were recorded through

CLARA. In miniscope cohort, the timing of miniscope neural recording is cross registered with behavior

video frames through CLARA controlled arduino board.

Optogenetic+VNS experiment groups

Experiment groups for training were based on stimulation type. The control cohort was injected with a

floxed YFP construct that did not contain an opsin and received light stimulation (see *Light stimulation*

parameters), the Success VNS cohort was electrically stimulated (see *VNS stimulation parameters*), and

the Arch w VNS cohort received both light and electrical stimulation. All groups automatically received

stimulation following every successful trial through the CLARA system.

Miniscope groups

Mice wore miniscopes for 5~10 minutes for a few times in their home cages or CLARA training box to

habituate the weight. When recording VNS response in homecage, mice wearing miniscopes and VNS

wires were put in homecage. About 4 minutes spontaneous neural activity were recorded as mice freely

moving in homecage, then 30~40 VNS were delivered every 20~30 seconds. Afterward, another ~4

minutes spontaneous activity were recorded. In sessions with AChR antagonists, scopolamine (1 mg/kg

body weight) and mecamylamine (10 mg/kg body weight) were dissolved in saline and delivered to mice

through Intraperitoneal (IP) injection. The concentration of scopolamine and mecamylamine cocktail was

chosen to have effects in brain circuits of memory and learning without debilitating effects according to

previous studies [100,101]. After 15 minutes, the neural response was recorded in homecage session.

For reach training recording, food restricted mice were mounted with miniscope and VNS wires and put in

training box to start CLARA training session. The minisope acquisition was turned on immediately as

CLARA training session started and each frame of the video was cross registered with the CLARA video

frames. All mice were primed without VNS to have one success reach before the first session started. On

the first day, VNS mice participated in one 20 minutes' Success VNS session. From day 2 to day 4, each

VNS mouse participated in two sessions of training, with one of them being a Success VNS session, the

other a no-stimulation session in which VNS was not delivered, given on a pseudorandomized schedule

(Fig. 7d). Control mice also received two training sessions without VNS each day. On days when mice

receive two sessions of training, the two sessions are 1~3 hours apart.

Place preference test

We used a standard conditioned place preference test to examine if VNS is rewarding or aversive. The

behavior apparatus contained two compartments separated by a gate. Mice were tested for baseline

preference in an initial 20-minute session where mice can freely navigate between compartments. Mice then were trained for three days with two 20 minute sessions each day where they received stimulation in only one compartment. Stimulation was delivered pseudo-random approximately once per minute. On the day of testing, no stimulation was given, and mice were tested to see which compartment they spent more time in [102]. The amount of time spent in each compartment was compared between the baseline and testing day. Experiments conducted with assistance from CU Anschutz Behavior Core.

Stimulation parameters

VNS stimulation parameters

For all VNS experiment groups, VNS was delivered as a 500ms train of 15 pulses with 100µs phase duration at 30Hz. Current amplitudes were 0.4-0.6mA. Stimulation parameters were controlled and delivered using Master8, PulsePal, or CLARA+Arduino connected to a stimulation isolation unit (A-M Systems, Model 2200 Analog Stimulus Isolator) to control amperage.

Light inhibition parameters

For all light stimulated groups, 561nm light was delivered continuously for 500ms. Light was delivered through a 200nm fiber-optic cable from a Class IIIb diode pumped solid state laser (Cobalt) at 0.5mW (calculated based on output efficiency from the bottom of the optical fiber).

Behavior and kinematic analysis

Manual behavior analysis

A success percent was generated for each session of each animal by determining the number of trials that resulted in a successful retrieval out of all trials initiated. Success percent was compared between stimulation groups across all days of training, as well as by early and late learning phases. Early learning phase refers to days 1-4 of training, Late phase refers to days 5-14 and were defined using a Weibull growth curve (See Quantification and Statistical Analysis). On days where animals received blocks of stimulation, such as rehearsal groups, stimulated and unstimulated trials were compared on a per mouse basis within days.

Behavior curator analysis

Videos acquired during CLARA training sessions were processed by custom python scripts overnight to extract key reach timepoints: reach initiation (ReachInit), when the hand leaves the box; reach max (ReachMax), the outward point of maximum distance from reach initiation; reach end (ReachEnd), when the hand returns to the cage; and stimulation onset (stim), when a trial received a trigger pulse for VNS or

light stimulation. The stamps of reachInit, reachMax and reachEnd were further manually screened for

consistency. The accuracy of CLARA trial outcome classifications were verified, and failures were

subcategorized *post-hoc* into reach, grasp and retrieval failures.

Kinematic analysis

3D Tracked data of paw center and pellet were tracked during reach attempts (between reach initiation and

reach end) were extracted using custom MATLABMATLAB scripts (MATLAB Simulink) and

documented as 3D data arrays for kinematic analysis. and documented as 3D data arrays.

Positional data for gross targeting experiments was determined by selecting the 3D location of the hand and

pellet at the reach max timepoint. Points were normalized such that the pellet center was 0,0,0. Euclidean

distance between the hand center and pellet center was then calculated using the norm function from

MATLAB. The mean distance from pellet was compared in early and late phases and between stimulation

groups. Reach velocity was obtained by measuring the absolute velocity between reach initiation and reach

end and averaging the velocity over that period. The mean velocity was compared in early and late phases

of training and between stimulation groups.

Reach consistency and expert reach

Positional data between these timepoints were then normalized such that the pellet center was 0,0,0. Reach

trajectory was defined as the time between reach initiation and reach end. Each trajectory was then

temporally warped to be the same arbitrary 'length' of time using dynamic time warping[103]. An expert

expert trajectory was constructed for each mouse by averaging the trajectories of all successful reach

attempts made on each mouse's last two days of training. Reach consistency was determined through

comparison of reach trajectories to each mouse's expert trajectory. Reach trajectories were compared to the

expert trajectory through a correlation coefficient to obtain the mean correlation coefficient for each mouse

in each training session. Additionally, any individual reach that had a correlation of 0.95 or higher with the

expert trajectory was defined as an 'expert reach', and the percent of expert reaches were also recorded for

each day. Number of expert reaches and mean correlation coefficients were then compared between VNS,

Arch+VNS and control groups in early and late phases.

Feature consistency

Several reach features were extracted from each reach attempt: Start location (X, Y, Z), End location (X,

Y, Z), Mean absolute velocity, Max absolute velocity, Pathlength (length of full trajectory, and Reach

consistency (defined above). The distribution of each feature was calculated for the early and late learning

phases based on the stimulation group. The distribution of each early-late pair was normalized using the

interquartile range of the early phase distribution. The normalized late interquartile range was subtracted

from the normalized early range and the difference was defined as 'delta feature consistency'. A positive

delta means that the distribution was tighter during the late phase compared to the early phase.

Electrophysiology recording and analysis

VNS electrophysiological recording in BF

While mice were either under maintained anesthesia (1.5% isoflurane), or awake in a homecage, VNS was

repeatedly delivered while recording from the left basal forebrain (see Optrode implantation). No

behavioral task was performed during recording. Mice received several (10-20) trains of VNS (0.5s, 30hz,

100us pulse-width, 0.6mA), delivered approximately 90 seconds apart. Data was recorded with Cheetah

acquisition software at 30kHz using a Digital Lynx SX (Neuralynx). TTL pulses were sent from the Master-

8 (A.M.P.I.) to the Digital Lynx SX for each pulse of a light or electric stimulation train. In acute

experiments, mice were opto-tagged after all VNS trains were delivered to identify cholinergic units.

Opto-tagging protocol

While mice were under maintained anesthesia (1.5% isoflurane), recordings were performed in the left

basal forebrain (see Optrode implantation). Light was delivered using a class 3b diode pumped solid state

laser (Cobalt) attached to the optrode through a ceramic ferule. Opto-tagging stimulus consisted of several

(10-20) trains of 5mW 488nm light delivered just above the basal forebrain through a 105um diameter fiber

optic spaced ~30 seconds apart. Trains consisted of 10 pulses of light at 20hz with a 10ms pulse duration.

Neural classification of BF response

After recording, units were clustered manually using clustering software SpikeSort 3d (Neuralynx) and

imported into MATLAB. Isolation distance and L-ratio were used to quantify cluster quality and noise

contamination[104]. The start of each train was identified post hoc using custom scripts (Mathworks) and

defined as a trial. The trial window, referred to as the 'VNS stimulation window' was defined as a 1 second

baseline before stimulation (-1 to 0s), VNS delivery (0 to 0.5s) and 1 second after the end of VNS (0.5 to

1.5s). Firing rate during the trial window was calculated using a 100ms moving average, shifted by 1ms

from the start of the trial window to 100ms after the end of the trial window. Baseline firing rate was defined

as the mean firing rate during the baseline period (-1 to 0s). Units were screened, and any unit with a mean

firing rate below 0.5Hz in anesthetized recordings or below 1Hz in awake recordings were removed from

the pool. +/-1ms around each stimulation pulse was removed to account for electrical noise. Firing rate was

converted into a Z score normalized to the mean firing rate and standard deviation baseline activity. If a

unit's normalized firing rate was ±2.56 s.d. from baseline firing rate for >100ms during the VNS delivery

window (0 to 0.5s), the unit was defined as VNS responsive. If the change in firing rate was 2.56 s.d above

baseline it was further subclassified as Activated, and if it was 2.56 s.d. below baseline, it was subclassified

as Suppressed. A unit that met both criteria was classified based on which occurred first. Peak response,

peak delay and response duration were compared between cholinergic and non cholinergic units in

anesthetized mice, and units recorded in awake mice. Peak response refers to the maximum normalized

firing rate of VNS activated units after stimulation onset (0 to 1.5s). Peak response delay refers to the

amount of time in ms from train onset (0s) to peak response. Duration refers to the total amount of time that

a VNS activated unit had normalized activity <2.56 s.d above baseline after stimulation onset (0 to 1.5s).

Neural classification of opto-tagging in BF

After recording, units were clustered manually using clustering software SpikeSort 3d (Neuralynx). Each

pulse was identified post hoc using custom MATLAB scripts (Mathworks) and defined as a trial. Tagged

units were identified using a stimulus-associated latency test (SALT,[18]). This test compares the distribution

of onset time of the first spike recorded during light delivery trials (10ms) to the onset time of spikes during

control windows of similar length (10ms). Units whose p-value from the SALT test was <0.05 were defined

as cholinergic, all other units were defined as non-cholinergic. Firing rates of cholinergic and non-

cholinergic units were calculated using the baseline firing rate of the unit from the VNS session.

Calcium imaging and analysis

Data aqcuisition

Miniscopes components an DAQ board were purchased and assembled by Optogenetics and Neural

Engineering (ONE) core according to UCLA miniscope (http://miniscope.org/) V3 guidelines. The

objective GRIN lens were Edmund optics # 64-520 and achromatic lens were #45-407. Images were

acquired at 30 Hz with MiniscopeControl data acquisition package. affiliated with UCLA miniscope. Each

imaging session was 15~20 minutes homecage or CLARA reach training session. The calcium signal

images were saved as TIFF stacks through USB3.0 port to SSD hard drive to reduce frame drops. For some

homecage sessions, the mice behavior was recorded simultaneously by a LG webcam controlled by

MiniscopeControl. For CLARA reach training session, behaviors were recorded by the CLARA system

and the timing of miniscope frames and behavior camera frames were coordinated.

Imaging analysis

Neural signal preprocessing

Image stacks acquired through miniscope were motion corrected using NoRMCorre packages

[105](https://github.com/flatironinstitute/NoRMCorre). For most sessions, rigid motion correction module

was sufficient to yield good results; in sessions with non-even shifting of the field of view, non-rigid motion

correction module was used. After motion correction, the field of view was cropped and spatially down

sampled 4x to reduce the file size for subsequent neural signal extraction. The neural signal were extracted

from images with CNMF-E package [106](https://github.com/zhoupc/CNMF_E). The results were manually

curated to discard non-neuronal ROIs. The result C-raw matrix which was a scaled dF x ROIs was used for

further analysis. For individual ROI, the time series dF of the whole session was further organized as 2D

matrix dF per trial x trials.

1006

Neural classification

To identify VNS responsive neurons, each trial's dF were organized 10 seconds before VNS onset and 10

seconds after, z-scale normalized to the dF values 3 seconds prior to the onset of VNS for each trial. We

call this trial-based z-score process which normalizes baseline period's average to 0 and disregards

spontaneous activity's effects on baseline dF level. Average z-scored dF of individual neurons was the

average of all trials of dF of the neuron. The response of individual neurons to VNS was measure as

maximum and minimum value of the average z-scored dF in the 3 seconds after VNS onset. Noise response

was estimated by calculating average z-scored dF for individual neurons using randomize VNS onset time

in the same session (excluding the 3 seconds window after each VNS onset in the whole session) and by

pooling 1000 repeats of such randomized procedures. For individual neuron, if the real maximum z-scored

dF in the 3 seconds window after VNS is higher than 95% value of the 1000 randomized maximum z-

scored dF for that neuron, the neuron was defined as activated by VNS; if the real minimum z-scored dF in

the 3 seconds window after VNS is lower than 95% value of randomized z-scored dF, the neuron was

defined as suppressed by VNS. The bi-directional modulated neurons were rare.

To measure the onset or peak timing for the VNS response, VNS responsive neurons' dF were z-scaled

with the whole session baseline mean calculated from time points lacking Ca2+ activity [107](defined as time

points with fluorescence values less than the 0.50 quantile of all fluorescence values). This general-based

z-score process allows dF level above or below 0 before VNS onset each trial and keeps the average

dynamics of VNS response better. For individual VNS activated neurons, the onset of VNS activation was

the first time point when the z-scored dF value in the 5 seconds after VNS onset reached above 2 s.d. of the

mean of 3 seconds baseline before VNS onset. For individual VNS suppressed neurons, the onset of VNS

suppression was the first time point when the z-scored dF value in the 5 seconds after VNS onset reached

below 2 s.d. of the mean of 3 seconds baseline before VNS onset.

To identify reach task responsive neurons, similar trial-based z-score process where employed. Trial dFs

were aligned by reach max. The baseline was taken as -6 to -3 seconds before reach max time. The response

of individual neurons to the task was measure as maximum and minimun value of the average z-scored dF

in the -800 to -300 ms (reach preparation), -300 to 200 ms (reach), 200 to 1700 ms (outcome) in reference

to reach max as time 0. The cut off value for maximum or minimum dF for these time windows were

estimated by similar randomization procedure as above in which that the same number of reach trials were

randomized across the whole session 1000 times. Success trials and failure trials were analyzed separately

unless noted otherwise. Several sub-groups of neurons were categorized as preparation-activated,

preparation-suppressed, reach-activated, reach-suppressed, success-activated, success-suppressed, failure-

activated and failure- suppressed. Preparation and reach modulated neurons were grouped together as reach

modulated neurons in figure 7. After neural classification, general-based z-score procedure were used to

obtain average success modulated neurons response VNS and no-stimulation sessions.

1042

Cross-registration of multiple sessions and cross session VNS modulated neurons definition

Due to computation power limit, we chose to process neural activity data from each session and register

neurons across sessions. CellReg package was used [108] https://github.com/zivlab/CellReg) to identify the

same neurons from multiple sessions based on spatial footprints of cellular activity (ROIs). Pairs of

neurons with correlation > 0.65 were regarded as the same neurons.

For individual neurons, if it was categorized as success-activated neurons in one of the two training sessions,

the neuron is regarded as a success-activated neuron. To look for significant modulation after VNS onset

in VNS sessions of these success activated neurons, the success outcome response of individual neuron was

aligned to VNS onset and measured in no-stimulation session and VNS session; the difference response

was obtained by no-stimulation response - VNS session response. The difference response in the 0~3s after

VNS onset that was higher or lower than 2.5 standard deviation of the mean of baseline (-3 to 0 before

VNS onset) for at least 0.15 s was regarded significant enhancement or attenuation in the VNS session. To

look for significant modulation before VNS onset, the neural response was aligned to reach end. The

baseline window was set from -6 to -3 seconds before reach end. This Reach end alignment allows us to

evaluate differences in neural representation of reach preparation, execution, and outcome. Differences

between the reach representation were compared between the Success VNS and no-stimulation sessions.

The difference response in the -3 to 3s around reach end that was higher or lower than 2.5 standard deviation

of the mean of baseline (-6 to -3 in reference to reach end) for at least 0.15 s was regarded significant

enhancement or attenuation. The onset time of the modulation was defined as the first time point of this

enhancement or attenuation.

The control sessions, AchR antagonists and recovery sessions were processed similarly as VNS stimulation

in homecage sessions described above. In addition, these sessions were temporally downsample to 15Hz

frame rate to reduce file size so that each mouse's three sessions imaging data were motion corrected and

concatenated together to save the post hoc cross registration step. For VNS-modulated neurons, the VNS

activation window was empirically measured and defined as rise onset to peak timing in **Fig. 6g&h** (0.8 to

2.8 seconds after VNS onset, suppression window as 0.2 to 1.6 seconds after VNS onset).

1069

Quantitation and Statistical Analysis

Statistical analyses were conducted using Graphpad, JMP (SAS), or MATLAB (MathWorks). No normality

tests were performed but individual data points are plotted to visualize distribution. We used parametric

statistics including paired and unpaired Student's t-tests, or one-way ANOVA with Tukey's HSD post-hoc

tests. Two-tailed tests and a < 0.05 were employed unless otherwise stated. We employed a mixed model

(Restricted Maximum Likelihood model (REML)) for all learning experiments. REML enables us to test

how fixed effects (dependent variables) and known random effects (individual mouse, sex, age) correlate

to an outcome variable.

1078   *Outcome = fixed effect + known random effect + error*

Models were constructed with one or two fixed effects. In cases where there were two fixed effects, we ran

full factorial models.

To determine early and late phases of learning a Weibull growth curve was applied to Sham VNS learning

data. The formula: $a*(1-Exp(-(Day/b)^c))$ a = asymptote, b = inflection point, c = growth rate. The AICc =

1295.90 and the R^2 = 0.21. We used the asymptote (55.50% +/- 6.8%) to represent the plateau in success

rate. The late learning phase was selected based on which training day exceeded the lower 95th percentile

of the asymptote (42.15%) which was day 4. We thus called late learning days 5-14 and early learning days

1-4.

1087

1088

1089

1090

1091

1092

1.   Ben-Menachem, E. *et al.* Vagus Nerve Stimulation for Treatment of Partial Seizures: 1. A

Controlled Study of Effect on Seizures. *Epilepsia* **35**, 616–626 (1994).

2.   The Vagus Nerve Stimulation Study Group. A randomized controlled trial of chronic vagus

nerve stimulation for treatment of medically intractable seizures. The Vagus Nerve

Stimulation Study Group. *Neurology* **45**, 224–230 (1995).

3.   Rush, A. J. *et al.* Vagus Nerve Stimulation for Treatment-Resistant Depression: A

Randomized, Controlled Acute Phase Trial. *Biological Psychiatry* **58**, 347–354 (2005).

4.   Shi, C., Flanagan, S. R. & Samadani, U. Vagus Nerve Stimulation to Augment Recovery

from Severe Traumatic Brain Injury Impeding Consciousness: A Prospective Pilot Clinical

Trial. *Neurol Res* **35**, 263–276 (2013).

5.   Tyler, R. *et al.* Vagus Nerve Stimulation Paired with Tones for the Treatment of Tinnitus: A

Prospective Randomized Double-blind Controlled Pilot Study in Humans. *Sci Rep* **7**, 11960

(2017).

6.   Dawson, J. *et al.* Vagus nerve stimulation paired with rehabilitation for upper limb motor

function after ischaemic stroke (VNS-REHAB): a randomised, blinded, pivotal, device trial.

The Lancet **397**, 1545–1553 (2021).

7.   Pruitt, D. T. *et al.* Vagus Nerve Stimulation Delivered with Motor Training Enhances

Recovery of Function after Traumatic Brain Injury. *J. Neurotrauma* **33**, 871–879 (2016).

8.   Ganzer, P. D. *et al.* Closed-loop neuromodulation restores network connectivity and motor

control after spinal cord injury. *Elife* **7**, (2018).

9.   Meyers, E. C. *et al.* Enhancing plasticity in central networks improves motor and sensory

recovery after nerve damage. *Nat Commun* **10**, 5782 (2019).

10. Kimberley, T. J. *et al.* Vagus Nerve Stimulation Paired With Upper Limb Rehabilitation

After Chronic Stroke. *Stroke* **49**, 2789–2792 (2018).

11. Krahl, S. E., Clark, K. B., Smith, D. C. & Browning, R. A. Locus coeruleus lesions suppress

the seizure-attenuating effects of vagus nerve stimulation. *Epilepsia* **39**, 709–714 (1998).

12. Beaumont, E. *et al.* Cervical vagus nerve stimulation augments spontaneous discharge in

second- and higher-order sensory neurons in the rat nucleus of the solitary tract. *Am J*

Physiol Heart Circ Physiol **313**, H354–H367 (2017).

13. Hulsey, D. R. *et al.* Parametric characterization of neural activity in the locus coeruleus in

response to vagus nerve stimulation. *Exp. Neurol.* **289**, 21–30 (2017).

14. Collins, L., Boddington, L., Steffan, P. J. & McCormick, D. Vagus nerve stimulation induces

widespread cortical and behavioral activation. *Current Biology* **31**, 2088-2098.e3 (2021).

15. Hulsey, D. R. *et al.* Reorganization of Motor Cortex by Vagus Nerve Stimulation Requires

Cholinergic Innervation. *Brain Stimul* **9**, 174–181 (2016).

16. Hulsey, D. R., Shedd, C. M., Sarker, S. F., Kilgard, M. P. & Hays, S. A. Norepinephrine

and serotonin are required for vagus nerve stimulation directed cortical plasticity. *Exp.*

Neurol. **320**, 112975 (2019).

17. Conner, J. M., Kulczycki, M. & Tuszynski, M. H. Unique Contributions of Distinct

Cholinergic Projections to Motor Cortical Plasticity and Learning. *Cereb Cortex* **20**, 2739–

2748 (2010).

18. Hangya, B., Ranade, S. P., Lorenc, M. & Kepecs, A. Central Cholinergic Neurons Are

Rapidly Recruited by Reinforcement Feedback. *Cell* **162**, 1155–1168 (2015).

19. Zhang, K., Chen, C. D. & Monosov, I. E. Novelty, salience, and surprise-timing are

signaled by neurons in the basal forebrain. *Curr Biol* **29**, 134-142.e3 (2019).

20. Chase, M. H., Sterman, M. B. & Clemente, C. D. Cortical and subcortical patterns of

response to afferent vagal stimulation. *Exp. Neurol.* **16**, 36–49 (1966).

21. Fraschini, M. *et al.* VNS induced desynchronization in gamma bands correlates with

positive clinical outcome in temporal lobe pharmacoresistant epilepsy. *Neurosci. Lett.* **536**,

14–18 (2013).

22. Usami, K. *et al.* Modulation of cortical synchrony by vagus nerve stimulation in adult rats.

Conf Proc IEEE Eng Med Biol Soc **2013**, 5348–5351 (2013).

23. Engineer, N. D. *et al.* Reversing pathological neural activity using targeted plasticity.

Nature **470**, 101–104 (2011).

24. Porter, B. A. *et al.* Repeatedly pairing vagus nerve stimulation with a movement

reorganizes primary motor cortex. *Cereb. Cortex* **22**, 2365–2374 (2012).

25. Mughrabi, I. T. *et al.* Development and characterization of a chronic implant mouse model

for vagus nerve stimulation. *eLife* **10**, e61270 (2021).

26. Feldman, D. E. The spike timing dependence of plasticity. *Neuron* **75**, 556–571 (2012).

27. Dayan, P. & Balleine, B. W. Reward, Motivation, and Reinforcement Learning. *Neuron* **36**,

285–298 (2002).

28. Leong, Y. C., Radulescu, A., Daniel, R., DeWoskin, V. & Niv, Y. Dynamic Interaction

between Reinforcement Learning and Attention in Multidimensional Environments. *Neuron*

93, 451–463 (2017).

29. Peters, A. J., Lee, J., Hedrick, N. G., O'Neil, K. & Komiyama, T. Reorganization of

corticospinal output during motor learning. *Nat. Neurosci.* **20**, 1133–1141 (2017).

30. Padmashri, R. & Dunaevsky, A. Modulation of excitatory but not inhibitory synaptic inputs

in the mouse primary motor cortex in the late phase of motor learning. *Neurosci Lett* **709**,

134371 (2019).

31. Censor, N., Sagi, D. & Cohen, L. G. Common mechanisms of human perceptual and motor

learning. *Nat Rev Neurosci* **13**, 658–664 (2012).

32. Wickens, J. R., Reynolds, J. N. & Hyland, B. I. Neural mechanisms of reward-related motor

learning. *Current Opinion in Neurobiology* **13**, 685–690 (2003).

33. Guo, W., Robert, B. & Polley, D. B. The Cholinergic Basal Forebrain Links Auditory Stimuli

with Delayed Reinforcement to Support Learning. *Neuron* **103**, 1164-1177.e6 (2019).

34. Ramanathan, D., Tuszynski, M. H. & Conner, J. M. The Basal Forebrain Cholinergic

System Is Required Specifically for Behaviorally Mediated Cortical Map Plasticity. *J.*

Neurosci. **29**, 5992–6000 (2009).

35.   Ramanathan, D. S., Conner, J. M., Anilkumar, A. A. & Tuszynski, M. H. Cholinergic

systems are essential for late-stage maturation and refinement of motor cortical circuits. *J*

Neurophysiol **113**, 1585–1597 (2015).

36.   Kucinski, A., Kim, Y. & Sarter, M. Basal forebrain chemogenetic inhibition disrupts the

superior complex movement control of goal-tracking rats. *Behav. Neurosci.* **133**, 121–134

(2019).

37.   Gielow, M. R. & Zaborszky, L. The Input-Output Relationship of the Cholinergic Basal

Forebrain. *Cell Reports* **18**, 1817–1830 (2017).

38.   Do, J. P. *et al.* Cell type-specific long-range connections of basal forebrain circuit. *eLife* **5**,

e13214 (2016).

39.   Lima, S. Q., Hromádka, T., Znamenskiy, P. & Zador, A. M. PINP: A New Method of

Tagging Neuronal Populations for Identification during In Vivo Electrophysiological

Recording. *PLoS One* **4**, e6099 (2009).

40.   Bowles, S., Williamson, W. R., Nettles, D., Hickman, J. & Welle, C. G. Closed-loop

automated reaching apparatus (CLARA) for interrogating complex motor behaviors. *J.*

Neural Eng. (2021) doi:10.1088/1741-2552/ac1ed1.

41.   Shmuelof, L., Krakauer, J. W. & Mazzoni, P. How is a motor skill learned? Change and

invariance at the levels of task success and trajectory control. *Journal of Neurophysiology*

108, 578–594 (2012).

42.   Kawai, R. *et al.* Motor cortex is required for learning but not executing a motor skill. *Neuron*

86, 800–812 (2015).

43.   Guo, J.-Z. *et al.* Cortex commands the performance of skilled movement. *eLife* **4**, e10774

(2015).

44.   Donoghue, J. P. & Sanes, J. N. Motor areas of the cerebral cortex. *J Clin Neurophysiol* **11**,

382–396 (1994).

1196 45. Sanes, J. N. & Donoghue, J. P. Plasticity and primary motor cortex. *Annu Rev Neurosci*

1197   **23**, 393–415 (2000).

1198 46. Peters, A. J., Chen, S. X. & Komiyama, T. Emergence of reproducible spatiotemporal

1199   activity during motor learning. *Nature* **510**, 263–267 (2014).

1200 47. Levy, S. *et al.* Cell-Type-Specific Outcome Representation in the Primary Motor Cortex.

1201   *Neuron* **107**, 954-971.e9 (2020).

1202 48. Borland, M. S. *et al.* Cortical Map Plasticity as a Function of Vagus Nerve Stimulation

1203   Intensity. *Brain Stimulation* **9**, 117–123 (2016).

1204 49. Reed, A. *et al.* Cortical map plasticity improves learning but is not necessary for improved

1205   performance. *Neuron* **70**, 121–131 (2011).

1206 50. Han, W. *et al.* A Neural Circuit for Gut-Induced Reward. *Cell* **175**, 887–888 (2018).

1207 51. Neuser, M. P. *et al.* Vagus nerve stimulation boosts the drive to work for rewards. *Nat*

1208   *Commun* **11**, 3555 (2020).

1209 52. Weber, I. *et al.* Trust your gut: vagal nerve stimulation in humans improves reinforcement

1210   learning. *Brain Communications* **3**, (2021).

1211 53. Pekny, S. E., Izawa, J. & Shadmehr, R. Reward-Dependent Modulation of Movement

1212   Variability. *J. Neurosci.* **35**, 4015–4024 (2015).

1213 54. Uehara, S., Mawase, F., Therrien, A. S., Cherry-Allen, K. M. & Celnik, P. Interactions

1214   between motor exploration and reinforcement learning. *Journal of Neurophysiology* **122**,

1215   797–808 (2019).

1216 55. Wolpert, D. M., Diedrichsen, J. & Flanagan, J. R. Principles of sensorimotor learning. *Nat*

1217   *Rev Neurosci* **12**, 739–751 (2011).

1218 56. Athalye, V. R., Santos, F. J., Carmena, J. M. & Costa, R. M. Evidence for a neural law of

1219   effect. *Science* **359**, 1024–1029 (2018).

1220 57. Oby, E. R. *et al.* New neural activity patterns emerge with long-term learning. *Proc Natl*

1221   *Acad Sci U S A* **116**, 15210–15215 (2019).

1222   58.   Aronov, D., Andalman, A. S. & Fee, M. S. A Specialized Forebrain Circuit for Vocal

1223   Babbling in the Juvenile Songbird. *Science* **320**, 630–634 (2008).

1224   59.   Garst-Orozco, J., Babadi, B. & Ölveczky, B. P. A neural circuit mechanism for regulating

1225   vocal variability during song learning in zebra finches. *eLife* **3**, e03697 (2014).

1226   60.   Dhawale, A. K., Smith, M. A. & Ölveczky, B. P. The Role of Variability in Motor Learning.

1227   *Annu Rev Neurosci* **40**, 479–498 (2017).

1228   61.   Wu, H. G., Miyamoto, Y. R., Castro, L. N. G., Ölveczky, B. P. & Smith, M. A. Temporal

1229   structure of motor variability is dynamically regulated and predicts motor learning ability.

1230   *Nat Neurosci* **17**, 312–321 (2014).

1231   62.   Athalye, V. R., Ganguly, K., Costa, R. M. & Carmena, J. M. Emergence of Coordinated

1232   Neural Dynamics Underlies Neuroprosthetic Learning and Skillful Control. *Neuron* **93**, 955-

1233   970.e5 (2017).

1234   63.   Dhawale, A. K., Miyamoto, Y. R., Smith, M. A. & Ölveczky, B. P. Adaptive Regulation of

1235   Motor Variability. *Current Biology* **29**, 3551-3562.e7 (2019).

1236   64.   Perez, S. M., Carreno, F. R., Frazer, A. & Lodge, D. J. Vagal Nerve Stimulation Reverses

1237   Aberrant Dopamine System Function in the Methylazoxymethanol Acetate Rodent Model

1238   of Schizophrenia. *J Neurosci* **34**, 9261–9267 (2014).

1239   65.   Farrand, A. Q. *et al.* Vagus nerve stimulation improves locomotion and neuronal

1240   populations in a model of Parkinson's disease. *Brain Stimul* **10**, 1045–1054 (2017).

1241   66.   Sawaki, L. *et al.* Cholinergic Influences on Use-Dependent Plasticity. *Journal of

1242   Neurophysiology* **87**, 166–171 (2002).

1243   67.   Shinoe, T., Matsui, M., Taketo, M. M. & Manabe, T. Modulation of Synaptic Plasticity by

1244   Physiological Activation of M1 Muscarinic Acetylcholine Receptors in the Mouse

1245   Hippocampus. *J Neurosci* **25**, 11194–11200 (2005).

1246   68.   Kang, J. I., Huppé-Gourgues, F. & Vaucher, E. Boosting visual cortex function and

1247   plasticity with acetylcholine to enhance visual perception. *Front. Syst. Neurosci.* **8**, (2014).

69. Jiang, L. *et al.* Cholinergic signaling controls conditioned-fear behaviors and enhances

plasticity of cortical-amygdala circuits. *Neuron* **90**, 1057–1070 (2016).

70. Lin, S.-C. & Nicolelis, M. A. L. Neuronal ensemble bursting in the basal forebrain encodes

salience irrespective of valence. *Neuron* **59**, 138–149 (2008).

71. Conner, J. M., Culberson, A., Packowski, C., Chiba, A. A. & Tuszynski, M. H. Lesions of

the Basal Forebrain Cholinergic System Impair Task Acquisition and Abolish Cortical

Plasticity Associated with Motor Skill Learning. *Neuron* **38**, 819–829 (2003).

72. Puzerey, P. A., Maher, K., Prasad, N. & Goldberg, J. H. Vocal learning in songbirds

requires cholinergic signaling in a motor cortex-like nucleus. *Journal of Neurophysiology*

120, 1796–1806 (2018).

73. Détári, L., Juhász, G. & Kukorelli, T. Effect of stimulation of vagal and radial nerves on

neuronal activity in the basal forebrain area of anaesthetized cats. *Acta Physiol Hung* **61**,

147–154 (1983).

74. Smiley, J. F., Subramanian, M. & Mesulam, M.-M. Monoaminergic–cholinergic interactions

in the primate basal forebrain. *Neuroscience* **93**, 817–829 (1999).

75. Zaborszky, L., Rosin, D. L. & Kiss, J. Alpha-adrenergic receptor (alpha(2 A)) is colocalized

in basal forebrain cholinergic neurons: a light and electron microscopic double

immunolabeling study. *J Neurocytol* **33**, 265–276 (2004).

76. Schwarz, L. A. & Luo, L. Organization of the Locus Coeruleus-Norepinephrine System.

Current Biology **25**, R1051–R1056 (2015).

77. Groves, D. A., Bowman, E. M. & Brown, V. J. Recordings from the rat locus coeruleus

during acute vagal nerve stimulation in the anaesthetised rat. *Neuroscience Letters* **379**,

174–179 (2005).

78. Whishaw, I. Q., O'Connor, W. T. & Dunnett, S. B. THE CONTRIBUTIONS OF MOTOR

CORTEX, NIGROSTRIATAL DOPAMINE AND CAUDATE-PUTAMEN TO SKILLED

FORELIMB USE IN THE RAT. *Brain* **109**, 805–843 (1986).

79. Lemon, R. N. Descending Pathways in Motor Control. *Annual Review of Neuroscience* **31**,

195–218 (2008).

80. Adler, A., Zhao, R., Shin, M. E., Yasuda, R. & Gan, W.-B. Somatostatin-Expressing

Interneurons Enable and Maintain Learning-Dependent Sequential Activation of Pyramidal

Neurons. *Neuron* **102**, 202-216.e7 (2019).

81. Biane, J. S., Takashima, Y., Scanziani, M., Conner, J. M. & Tuszynski, M. H.

Reorganization of Recurrent Layer 5 Corticospinal Networks Following Adult Motor

Training. *J Neurosci* **39**, 4684–4693 (2019).

82. Luft, A. R. & Schwarz, S. Dopaminergic signals in primary motor cortex. *International*

Journal of Developmental Neuroscience **27**, 415–421 (2009).

83. Hosp, J. A., Pekanovic, A., Rioult-Pedotti, M. S. & Luft, A. R. Dopaminergic Projections

from Midbrain to Primary Motor Cortex Mediate Motor Skill Learning. *J. Neurosci.* **31**,

2481–2487 (2011).

84. Heffley, W. *et al.* Coordinated cerebellar climbing fiber activity signals learned

sensorimotor predictions. *Nat Neurosci* **21**, 1431–1441 (2018).

85. Eggermann, E., Kremer, Y., Crochet, S. & Petersen, C. C. H. Cholinergic signals in mouse

barrel cortex during active whisker sensing. *Cell Rep* **9**, 1654–1660 (2014).

86. Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity

in cortex. *Nat Commun* **7**, 13289 (2016).

87. Musall, S., Urai, A. E., Sussillo, D. & Churchland, A. K. Harnessing behavioral diversity to

understand neural computations for cognition. *Current Opinion in Neurobiology* **58**, 229–

238 (2019).

88. Lai, J. & David, S. V. Short-Term Effects of Vagus Nerve Stimulation on Learning and

Evoked Activity in Auditory Cortex. *eNeuro* **8**, (2021).

89. Nitsche, M. A. *et al.* Facilitation of implicit motor learning by weak transcranial direct

current stimulation of the primary motor cortex in the human. *J Cogn Neurosci* **15**, 619–626

(2003).

90. Khanna, P. *et al.* Low-frequency stimulation enhances ensemble co-firing and dexterity

after stroke. *Cell* **184**, 912-930.e20 (2021).

91. Potok, W., Bächinger, M., Groen, O. van der, Cretu, A. L. & Wenderoth, N. Transcranial

Random Noise Stimulation Acutely Lowers the Response Threshold of Human Motor

Circuits. *J. Neurosci.* **41**, 3842–3853 (2021).

92. Fritsch, B. *et al.* Direct current stimulation promotes BDNF-dependent synaptic plasticity:

Potential implications for motor learning. *Neuron* **66**, 198–204 (2010).

93. Ammann, C., Spampinato, D. & Márquez-Ruiz, J. Modulating Motor Learning through

Transcranial Direct-Current Stimulation: An Integrative View. *Front Psychol* **7**, 1981 (2016).

94. Kang, E. K., Kim, Y. K., Sohn, H. M., Cohen, L. G. & Paik, N.-J. Improved picture naming

in aphasia patients treated with cathodal tDCS to inhibit the right Broca's homologue area.

Restor Neurol Neurosci **29**, 141–152 (2011).

95. Stagg, C. J. *et al.* Polarity and timing-dependent effects of transcranial direct current

stimulation in explicit motor learning. *Neuropsychologia* **49**, 800–804 (2011).

96. Albuquerque, L. L. de *et al.* An acute application of transcranial random noise stimulation

does not enhance motor skill acquisition or retention in a golf putting task. *Human*

Movement Science **66**, 241–248 (2019).

97. Rossi, J. *et al.* Melanocortin-4-receptors Expressed by Cholinergic Neurons Regulate

Energy Balance and Glucose Homeostasis. *Cell Metab* **13**, 195–204 (2011).

98. Zhao, S. *et al.* Cell-type Specific Optogenetic Mice for Dissecting Neural Circuitry Function.

Nat Methods **8**, 745–752 (2011).

99. Whishaw, I. Q., Whishaw, P. & Gorny, B. The Structure of Skilled Forelimb Reaching in the

Rat: A Movement Rating Scale. *J Vis Exp* 816 (2008) doi:10.3791/816.

100. Riekkinen, P., Sirviö, J., Aaltonen, M. & Riekkinen, P. Effects of concurrent manipulations

of nicotinic and muscarinic receptors on spatial and passive avoidance learning.

Pharmacol Biochem Behav **37**, 405–410 (1990).

101. Riekkinen, P., Riekkinen, M. & Sirviö, J. Cholinergic drugs regulate passive avoidance

performance via the amygdala. *J Pharmacol Exp Ther* **267**, 1484–1492 (1993).

102. Prus, A. J., James, J. R. & Rosecrans, J. A. Conditioned Place Preference. in *Methods of*

Behavior Analysis in Neuroscience (ed. Buccafusco, J. J.) (CRC Press/Taylor & Francis,

2009).

103. Li, Q. *et al.* Refinement of learned skilled movement representation in motor cortex deep

output layer. *Nature Communications* **8**, 15834 (2017).

104. Schmitzer-Torbert, N., Jackson, J., Henze, D., Harris, K. & Redish, A. D. Quantitative

measures of cluster quality for use in extracellular recordings. *Neuroscience* **131**, 1–11

(2005).

105. Pnevmatikakis, E. A. & Giovannucci, A. NoRMCorre: An online algorithm for piecewise

rigid motion correction of calcium imaging data. *J Neurosci Methods* **291**, 83–94 (2017).

106. Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from

microendoscopic video data. *eLife* **7**, e28728 (2018).

107. Jimenez, J. C. *et al.* Anxiety Cells in a Hippocampal-Hypothalamic Circuit. *Neuron* **97**, 670-

683.e6 (2018).

108. Sheintuch, L. *et al.* Tracking the Same Neurons across Multiple Days in Ca2+ Imaging

Data. *Cell Rep* **21**, 1102–1115 (2017).

1345