Systems-based approach for optimization of a scalable bacterial ST mapping

assembly-free algorithm

Natasha Pavlovikj[1¶], Joao Carlos Gomes-Neto[2,3¶], Jitender S. Deogun[1], Andrew

K. Benson[2,3*]

[1] Department of Computer Science and Engineering, University of Nebraska-

Lincoln, Lincoln, Nebraska, United States of America

[2] Department of Food Science and Technology, University of Nebraska-Lincoln,

Lincoln, Nebraska, United States of America

[3] Nebraska Food for Health Center, University of Nebraska-Lincoln, Lincoln,

Nebraska, United States of America

* Corresponding Author:

E-mail: abenson1@unl.edu (AKB)

¶ These authors contributed equally to this work.

Abstract

Epidemiological surveillance of bacterial pathogens requires real-time data analysis with a fast turn-around, while aiming at generating two main outcomes: 1) Species level identification; and 2) Variant mapping at different levels of genotypic resolution for population-based tracking, in addition to predicting traits such as antimicrobial resistance (AMR). With the recent advances and continual dissemination of whole-genome sequencing technologies, large-scale population-based genotyping of bacterial pathogens has become possible. Since bacterial populations often present a high degree of clonality in the genomic backbone (i.e., low genetic diversity), the choice of genotyping scheme can even facilitate the understanding of ancestral relationships and can be used for prediction of co-inherited traits such as AMR. Multi-locus sequence typing (MLST) fits that purpose and can identify sequence types (ST) based on seven ubiquitous genome-scattered loci that aid in genotyping isolates beneath the species level. ST-based mapping also standardizes genotyping across laboratories and can be consistently used worldwide. However, ST-based algorithms, when using Illumina paired-end sequences, often rely on genome assembly prior to classification. That hinders rapid genotyping and scalability which are essential aspects of genomic epidemiology. stringMLST is a kmer-based ST method with the capacity to solve both hurdles. Yet, a comprehensive scalable comparison of its use in contrast to a standard MLST program for a wide array of phylogenetically divergent Public Health-relevant bacterial pathogens is lacking. Herein, we first demonstrated that stringMLST is a fast tool that can be deployed for ST-based epidemiological

inquiries of bacterial populations. Additionally, we systematically evaluated and

showed the impact of genome-intrinsic and -extrinsic features, as well as the

optimal kmer length in maximizing the performance of stringMLST on species-

by-species basis, and highlighted a few instances where this program may not be

applicable in its current format. Furthermore, we integrated stringMLST as part of

our freely available and scalable hierarchical-based population genomics platform

called ProkEvo. Besides facilitating automatable and reproducible bacterial

population guided analysis, ProkEvo now offers a rapidly deployable genomic

epidemiology tool for ST mapping, with specific guidance on how to optimize its

performance, that can be widely applicable by microbiological laboratories and

epidemiological agencies.

Introduction

Modern epidemiological investigation of bacterial pathogens is primarily focused

on real-time, fast turn-around characterization of many thousands of isolates,

routinely received by Public Health laboratories and regulatory agencies [1][2].

Additionally, due to the large-scale availability of whole-genome sequencing

(WGS) and an emerging emphasis on retrieving accurate metadata, three major

goals can be achieved with population-based inquiries: 1) Species identification;

2) Genotyping at different levels of hierarchical resolution; and 3) Prediction of

co-inherited traits such as antimicrobial resistance (AMR) via loci mapping, or

based on the assessment of population-inherited linkage between the core- and

accessory-genomes [3]. In general, bacterial populations contain a genomic

backbone (i.e., clonal-frame), for which the degree of clonality is predicted to be a

heritable trait, leading to a measurable degree and pattern of co-inheritance

between core- and accessory-genes (loci) – high *linkage disequilibrium* (LD)

[4][5][6][7]. Core loci are those found in 99% of the genomes or more; whereas

accessory loci represent a sparse ensemble present in less than 99% of genomes,

which combined form the species-specific pan-genomic content [66]. In theory,

one should be able to identify a genotyping scheme that not only facilitates

sufficient isolate differentiation beneath the species level, but can also reveal the

pattern of population diversification and structuring, while being used in

phenotypic prediction such as for AMR traits. This optimal level of genotypic

resolution can be considered an informative genotypic unit that facilitates both

ecological and epidemiological inquiries.

Multi-locus sequence typing (MLST) is a well-established and widely used

genotyping technique that classifies bacterial genomes into sequence types (ST)

[8]. ST classification is achieved by mapping seven ubiquitous genome-scattered

loci using highly curated, and species-specific allelic databases. Essentially,

sequences for those seven loci can be generated by polymerase chain reaction

(PCR)-based assays, or WGS [8][9][44]. Regardless of the methodology, partial

sequences for each locus are mapped against multiple ever-expanding public

allelic databases [27][30][67]. The combination of all seven allelic numbers

defines which ST number the isolate is classified into. ST-based classification

provides useful genotyping approach below the species level, while revealing the

population structure and retrieving ancestral relationships, since when five or

more loci are identical to one another between two genomes, both belong to the

same clonal complex (i.e., group of STs that have shared a common ancestor very

recently) [8][9][28][44]. Moreover, ST-based genotyping standardizes the

nomenclature for intra- and inter-laboratorial diagnostics and epidemiological

inquiries worldwide [9][10][11]. Lastly, genes that are part of the MLST scheme

can co-vary with other accessory loci important for phenotypic predictions such

serotyping (e.g., *Salmonella enterica*) and inter-species AMR predictions

[12][13][14].

Harnessing this intrinsic LD property of bacterial pan-genomes has been the

basis for recent innovation in epidemiological investigations, whereby heuristic

mapping of STs led to accurate prediction of AMR profiles [14][15]. However,

ST-based classification, using the most widely distributed Illumina sequencing

technology, is often dependent on genome assemblies [27][30][67]. That results in

slower turn-around for data analysis and hinders ST-based surveillance efforts for

enhancing scalability when working with many thousands of genomes [16][21].

One approach to overcome those hurdles is to use kmer-based ST classification

directly from Illumina paired-end raw reads (assembly-free). stringMLST is an

approach that successfully accomplishes that goal [17], but has not yet been

systematically tested for its analytical performance while classifying thousands of

genomes from phylogenetically divergent bacterial pathogens.

Therefore, the purpose of this work was to test whether stringMLST can be

used as a rapid and accurate replacement of the standard MLST programs for

scalable genotyping of phylogenetic divergent bacterial pathogens, with direct

Public Health implications, using Illumina raw sequences. In contrast to the

original work [17], our systematic approach demonstrated a comprehensive

comparison between a standard MLST program vs. stringMLST, while

classifying many thousands of genomes across 15 pathogens, including:

*Acinetobacter baumannii (*Phylum: Proteobacteria)*, Clostridioides difficile*

*(*Phylum: Firmicutes)*, Enterococcus faecium (*Phylum: Firmicutes)*, Escherichia*

*coli (*Phylum: Proteobacteria)*, Haemophilus influenzae (*Phylum: Proteobacteria)*,*

*Helicobacter pylori (*Phylum: Proteobacteria)*, Klebsiella pneumoniae (*Phylum:

Proteobacteria)*, Mycobacterium tuberculosis (*Phylum: Actinobacteria)*, Neisseria*

*gonorrhoeae (*Phylum: Proteobacteria)*, Pseudomonas aeruginosa (*Phylum:

Proteobacteria)*, Streptococcus pneumoniae (*Phylum: Firmicutes)*, Campylobacter*

jejuni (Phylum: Proteobacteria)*, Listeria monocytogenes (*Phylum: Firmicutes),

Salmonella enterica (Phylum: Proteobacteria), and *Staphylococcus aureus*

(Phylum: Firmicutes). Additionally, we have also analyzed the optimal kmer

length for 23 most relevant zoonotic serovars of *Salmonella enterica* subsp.

enterica lineage I (*S. enterica*) individually given its cryptic population structure

and diverse ecology [12]. For comparison between programs, we use the

following analytical outcomes as proxies for algorithmic performance-based

assessment: 1) Computational runtime and memory used for genome

classification; 2) ST richness and diversity metrics; and 3) Proportion of non-

classified STs and concordance between programs. [17]Importantly, we measured

the statistical contribution of genome-intrinsic (genome size and composition) and

-extrinsic (ST database size) factors on classification accuracy at species level of

resolution, including the identification of the optimal kmer length per species.

This comprehensive approach revealed how stringMLST is a deployable ready-to-

use program that can be further optimized (parameter fine-tunning) based on the

species dataset, while attempting to scale its application for practical

implementation in microbiological laboratories and epidemiological agencies.

Lastly, we added stringMLST to our computational platform called ProkEvo,

aiming at facilitating its automated, reproducible, and scalable use, in

combination with other standard assembly-based hierarchical genotypic and pan-

genomic mapping approaches for bacterial population genomic analyses.

Materials and methods

This systems-based comparison between mlst and stringMLST was centered at

capturing their differences in computational and statistical performances, and was

accomplished through the following steps: 1) Narrow-scope comparative analysis

across four phylogenetic distinct pathogens species; 2) Further examination of

algorithmic performance within a single ecologically diverse bacterial species;

and 3) Wide-scope comparison between phylogenetic divergent pathogenic

species with Public Health relevance and with databases available on pubMLST

(https://pubmlst.org/) for direct contrast between stringMLST and mlst.

Datasets used for narrow-scope analysis

WGS data from four major bacterial pathogens, including *Campylobacter jejuni*, *Listeria monocytogenes*, *Salmonella enterica* subps. *enterica* lineage I (*S. enterica*) and *Staphylococcus aureus,* were selected to be used in this first part of the study. Our basis for that choice was due to three *a priori* defined criteria: 1) Select bacterial species from two main phylogenetic divergent Phyla: Firmicutes (*L. monocytogenes* and *S. aureus*) and Proteobacteria (*C. jejuni* and *S. enterica*); 2) Select zoonotic pathogens that continually cause human illnesses worldwide [18]; and 3) Consider their epidemiological relevance according to the Centers for Disease Control and Prevention (CDC) [19]. Specifically for *S. enterica*, 20 of the CDC most investigated serovars were represented in the dataset, which includes: *S.* Agona, *S.* Anatum, *S.* Braenderup, *S.* Derby, *S.* Dublin, *S.* Enteritidis, *S.* Hadar, *S.* Heidelberg, *S.* Infantis, *S.* Javiana, *S.* Johannesburg, *S.* Kentucky, *S.* Mbandaka, *S.* Montevideo, *S.* Muenchen, *S.* Newport, *S.* Schwarzengrund, *S.* Senftenberg, *S.* Thompson, and *S.* Typhimurium [20]. All publicly available raw paired-end Illumina reads for these organisms were downloaded from NCBI using parallel-fastq-dump [58]. Genomes used for all analyses were randomly selected from a previously downloaded samples of isolates containing *C. jejuni* (n = 21,919 genomes), *L. monocytogenes* (n = 19,633 genomes), *S. enterica* (n = 25,284 genomes), and *S. aureus* (n = 11,990 genomes) that were processed through the computational platform ProkEvo [21]. Specifically, our study design was comprised of random sampling of 600 genomes from each species, except for *S. enterica* for which 600 genomes were randomly drawn per serovar (list of all 20

serovars is shown in S1 Table). For each species and all *S. enterica* serovars, all

~600 genomes were randomly split into three independent batches, with ~200

genomes each. The batches were created to measure the degree of variation in

classification accuracy when comparing the two ST-based genotyping programs.

While for the majority of *S. enterica* serovars there were a total of 600 genomes

available, the total number of raw reads publicly available on NCBI and

ultimately used for the analyses for *S.* Agona, *S.* Derby, *S.* Johannesburg, *S.*

Mbandaka and *S.* Senftenberg was 565, 590, 534, 535 and 563 respectively. The

final total number of genomes used per species was: *C. jejuni* (n = 600), *L.*

monocytogenes (n = 600), *S. enterica* (n = 11,787), and *S. aureus* (n = 600). Text

file containing all genome NCBI SRA identifications is available here,

https://figshare.com/articles/dataset/_/16735411.

Software tools

mlst

mlst is a standard approach for scanning genome assemblies against traditional

PubMLST typing schemes [22]. The genome assemblies can be in

FASTA/GenBank/EMBL formats [22]. mlst (version 2.16.2) was installed using

Anaconda, a package and environment manager that supports maintaining and

installing various open-source conda packages [26]. mlst uses genome assemblies

as an input. In order to generate assemblies from the raw Illumina paired-end

reads, multiple pre-processing steps were performed. Quality trimming and

adapter clipping were performed using Trimmomatic [50], while FastQC was

used to check and verify the quality of the trimmed reads [51]. The paired-end

reads were assembled *de novo* into contigs using SPAdes with the default

parameters [52]. The quality of the assemblies was evaluated using QUAST [53].

The information obtained from QUAST was used to discard assemblies with 0 or

more than 300 contigs, or assemblies with N50 value of less than 25,000 [21].

Finally, the assemblies that passed the quality control were used with mlst, where

they are categorized into specific variants based on the allele combinations from

seven ubiquitous, house-keeping genes [22]. A list of the exact versions of the

bioinformatics tools used for generating assemblies for mlst are shown on S2

Table. We used mlst with the default options (e.g., *mlst --legacy --scheme*

<scheme> --csv <assembly.fasta> > <output>) and the following schemes:

"senterica" (for *S. enterica*), "campylobacter" (for *C. jejuni)*, "lmonocytogenes"

(for *L. monocytogenes*), "saureus" (for *S. aureus*). The distribution of mlst comes

with set of pre-downloaded ST schemes. More details about these MLST

schemes, such as the number of alleles in the seven genes and the number of ST

classifications available are shown on S3 Table. To obtain the ST classifications

of all datasets, mlst was run as part of the computational platform ProkEvo [21].

Additionally, a separate run of the mlst program was used to conduct a pairwise

comparison between the computational performance (runtime and memory usage)

of mlst and stringMLST. The used mlst script can be found here,

https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/scripts/ml

st.submit.

stringMLST

stringMLST is an assembly- and alignment-free rapid tool for ST-based

classification of Illumina paired-end raw reads based on kmers [17]. For the

analyses performed in this paper, we used stringMLST version 0.6.3. stringMLST

was installed using Anaconda [26]. The first step of using stringMLST was to

download the respective MLST scheme from PubMLST. In order to do this, a

species name and a kmer length were needed. The default kmer length used and

suggested by the developers of stringMLST for reads with lengths between 55 and

150 base pairs or nucleotides is 35 (common read length for Illumina paired-end

reads) [17]. We used stringMLST with the default options (e.g., *stringMLST.py --*

getMLST --species=<species_name> -P <output_prefix> -k <kmer>) and the

following species names, "Salmonella enterica", "Campylobacter jejuni",

"Listeria monocytogenes", "Staphylococcus aureus", and kmer lengths of 10, 20,

30, 35, 45, 55, 65, 70, 80, 90 independently

(https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/scripts/str

ingMLST_dbs.submit). More details about the downloaded MLST schemes, such

as the number of alleles in the seven genes and the number of ST classifications

available are shown on S3 Table. After the MLST scheme was downloaded and

prepared, the final step was to run "stringMLT.py –predict" for the ST

classification. For this, we ran stringMLST with the databases *a priori* created and

the respective paired-end raw reads and kmer lengths of 10, 20, 30, 35, 45, 55, 65,

70, 80, 90 independently (e.g., *stringMLST.py --predict -d*

<directory_raw_reads> -p -r -t -x -P <database_prefix> -k <kmer> -o

<output>)

(https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/scripts/str

ingMLST.submit). Our choice of using an increasing gradient of kmer lengths

was to evaluate whether the kmer length parameter could be optimized to enhance

ST-based classification accuracy across species. Lastly, stringMLST was also

integrated as part of the computational platform ProkEvo for a rapid ST-based

genotyping as part of a hierarchical genotypic scheme [21][57]. This

implementation can be found here,

https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/Prokevo_st

ringMLST.

ProkEvo-based MLST classifications

In order to compare the ST-based classification accuracy and conduct other

statistical analysis (e.g., identifying major contributing factors influencing ST-

based classifications) between mlst version 2.16.2 (assembly-dependent) and

stringMLST version 0.6.3 (assembly-independent), all initial ST calls for all

selected genomes, across all four species, were done using mlst [22] through the

computational platform ProkEvo [21]. In brief, ProkEvo uses bacterial Illumina

raw paired-end sequences as an input, and the following steps are sequentially

done prior to ST-based genotyping using mlst: Trimmomatic for sequence

trimming [50], FastQC for quality control of the trimmed reads [51], SPAdes for

de novo genome assembly [52], and QUAST for quality assessment of the

genome assemblies [53]. More information on how to install and use ProkEvo for

hierarchical bacterial population genomic analyses can be found here,

https://github.com/npavlovikj/prokevo.

277   Genome-intrinsic and –extrinsic factors that can

278   influence algorithmic performance

Both genome-intrinsic and –extrinsic factors were considered to determine their

contribution on the accuracy of ST classifications when comparing mlst vs.

stringMLST.

The genome-intrinsic variables considered in these analyses were: number of

contigs per genome, total number of nucleotides per genome (genome length),

GC% content per genome, and dinucleotide composition of genomes. The number

of contigs per genome, as well as the genome length, were calculated using the

assembled contigs from SPAdes [52]. The number of contigs was calculated for

each genome using the Linux "grep" utility (e.g., *grep ">" assembly.fasta | wc -*

l). The total number of nucleotides per genome was calculated using the

"getlengths" function from the AMOS package [54]. For this analysis, we used

AMOS v3.1. "getlengths" provides the length for each contig, and a custom Bash

script was used to summarize these values per genome. The GC% content was

calculated using the program FastQC [51]. FastQC is used to check and verify the

quality of the raw Illumina paired-end raw reads. With each pair of raw reads

from all datasets, FastQC v0.11 was used. One of the statistics checked for read

quality is GC% and this value was extracted with custom Bash script from the file

"fastqc_data.txt" once the FastQC output was generated. Since FastQC outputs

the GC% per read, the average of both reads was calculated as the final read

GC%. The dinucleotide composition of the genomes was calculated with the

function "compseq" from the EMBOSS package [55]. "compseq" calculates the

frequency of words of a specific length (e.g., length is 2 in the case of

dinucleotides) from given input genome sequences. For these analyses we used

EMBOSS v6.6 with the command "*compseq -word 2 -outfile <output>*

assembly.fasta" for all datasets and genomes *a priori* assembled with SPAdes

[52]. Next, customized Bash script was used to count the total number of

occurrences of each dinucleotide for each genome across all bacterial species.

Finally, all these outputs were merged per genome using custom Python script to

facilitate statistical analyses and data visualization. The used scripts can be found

here,

https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/scripts.

The genome-extrinsic variables used in the analyses presented here were: the

total count of unique STs per database and the total count of unique alleles across

all seven loci used for ST classification across all bacterial species. These

genome-extrinsic variables were extracted from the PubMLST databases for both

stringMLST and mlst using custom Bash scripts. While the first step of

stringMLST is to download the newest available MLST scheme from PubMLST,

the distributed version of mlst comes with set of pre-downloaded ST and allelic

schemes. For each MLST scheme, the mlst distribution has a separate directory

with 8 files - seven are ".tfa" files with the fasta sequences of the alleles for each

locus, and one file (e.g., senterica.txt) contains the ST information (i.e., the total

number of STs mapped including their specific allelic composition across all

seven loci for that given species). To calculate the total number of unique STs, we

used the Linux utility "wc" with the text file with ST information (e.g., *wc -l*

senterica.txt). To calculate the total count of unique alleles across the seven loci,

the "grep" Linux utility was used with the seven ".tfa" files (e.g., *grep ">" *.tfa |*

wc -l). All calculations were done per bacterial species. The downloaded MLST

scheme with stringMLST is in a separate directory for each organism and used

kmer length. This directory had 12 files - seven are ".tfa" files with fasta

sequences for all alleles across all seven loci, and one file has the ST profiles

(e.g., Salmonella_enterica_profile.txt), while the remaining files contained

information about the extracted kmers and additional config and log information.

Similarly, the total number of unique STs for stringMLST was counted using the

Linux utility "wc" with the text file with ST profile information (e.g., *wc -l*

Salmonella_enterica_profile.txt) and the total count of unique alleles per loci was

extracted using the "grep" Linux utility with the seven ".tfa" files (e.g., *grep ">"*

.tfa | wc -l). Similarly, all ST and allelic counts were carried out per bacterial

species. With stringMLST, the MLST schemes are downloaded and prepared

separately for each different kmer length used. However, the kmer length did not

affect the number of STs and unique alleles per organism. Thus, these values are

the same across organisms and kmer lengths for stringMLST.

Kmer-based distribution across ST programs

In order to assess the potential impact of random mapping or occurrence of kmers of different lengths across different bacterial species we randomly chose 100 raw Illumina paired-end reads from the initial *C. jejuni*, *L. monocytogenes*, *S. aureus* and *S.* Typhimurium (major representative zoonotic serovar of *S. enterica*) isolates. For each read, we extracted all unique kmers of length 10, 20, 30, 35, 45, 55, 65, 70, 80 and 90 respectively, and counted their occurrence in the corresponding raw reads. This was done using DSK v2.2.0 [56] (https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/scripts/ds k.submit). Next, the total number of kmer frequency was summarized per organism and kmer length, and the mean value was calculated to examine the distribution of different kmers across the raw reads. For each database created with stringMLST, a file with the kmer frequency for the used ST scheme was generated. Using the kmers generated from the raw reads and the stringMLST database, a relative frequency of the common kmers was calculated (calculated as a ratio between the common kmers and the unique kmers from all the kmers generated between the raw reads and the stringMLST database, e.g., *(common_kmers/unique_total_observations)*100*). The code used for this can be found in our GitHub repository (https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/figures_c ode).

Agreement in ST classification between programs

In order to assess the overall accuracy of stringMLST compared to the standard mlst approach for ST calls, a percentage of agreement in ST classification was calculated. For this, the initial dataset composed of 600 genomes from either *C. jejuni*, or *L. monocytogenes*, or *S. aureus* was selected, in addition to a total of 11,787 genomes across twenty zoonotic serovars of *S. enterica* (~600 genomes per serovar, S1 Table). The program stringMLST was run with increasing kmer lengths ranging from 10 to 90 nucleotides. If both stringMLST and mlst produced identical ST calls, either "good" or "bad" ones, the call was a match. A "good" and "bad" call represent ST with a number or a missing/blank value, respectively. The remaining combinations were classified as a mismatch. Next, the percentage of agreement (concordance) was calculated with custom R base script (https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/figures_c ode).

Computational platforms

All computational analyses performed for this paper were done on Crane - one of the high-performance computing clusters at the University of Nebraska-Lincoln Holland Computing Center [23]. Crane is Linux cluster, having 548 Intel Xeon nodes with RAM ranging from 64 GB to 1.5 TB. The scalability of ProkEvo with stringMLST was tested on the Open Science Grid (OSG), a distributed, high-throughput computational platform for large-scale scientific research [24][25]. OSG is a national consortium of more than 100 academic institutions and

386   laboratories that provide storage and tens of thousands of resources to OSG users.

387   These sites share their idle resources via OSG for opportunistic usage. The OSG

388   resources are Linux-based, and due to the different sites involved, the hardware

389   specifications of the resources are different and vary.

391   Computational performance

To evaluate the computational performance of stringMLST in comparison to the

mlst program, we assessed the runtime and memory usage of both programs. For

this, we chose four different datasets, *C. jejuni*, *L. monocytogenes*, *S. aureus* and

S. Typhimurium (major representative zoonotic serovar of *S. enterica*), with three

different batches of 200 genomes each, with a total of 600 genomes each. We ran

mlst with all required steps, such as quality trimming and adapter clipping, *de*

novo assembly and assembly discarding on each dataset (see Section Software

tools: mlst for more detailed description). Separately, we ran stringMLST with a

range of 10 different kmer lengths (10, 20, 30, 35, 45, 55, 65, 70, 80, 90) on each

dataset. For each organism, the runtime was calculated as an average of all 200

genomes per batch. In general, the runtime depends on multiple factors, such as

the specification and capabilities of the used computational platform. Since the

runtime can vary depending on these various factors, the average statistics was

used to show the central tendency of the runtime when comparing stringMLST vs.

mlst. The runtime was calculated using the "date" command integrated in the

Unix operating systems (e.g., *t=`date +%s`; mlst --legacy --scheme senterica --*

csv assembly.fasta > <output>; tt=`date +%s`; total_time=$((tt-t))). For each

organism, the memory was calculated as the maximum memory recorded from all

200 genomes per batch, since all genomes were analyzed separately and

concurrently. In the case of mlst, the recorded memory was the maximum

memory of all the steps ran prior to mlst, such as trimming, *de novo* assembly,

quality checking, filtering and ST typing. The memory used for these steps

considerably varies from a few MBs to a few GBs (e.g., filtering vs. *de novo*

assembly), and since the memory is a physical limitation of the computational

platform, the maximum used memory was calculated for each organism and

batch. The memory used was calculated using the "cgget" command that tracks

various parameters from the Linux Control Groups (cgroups) per running job

(e.g., *mlst --legacy --scheme senterica --csv assembly.fasta > <output>; r=`cgget*

-r memory.usage_in_bytes /slurm/uid_${UID}/job_${SLURM_JOBID}/`;

mem=`echo $r | awk -F: '{print $3}'`).

Incorporating stringMLST in ProkEvo

ProkEvo is a freely available and scalable computational platform capable of

facilitating bacterial population genomics analyses while combining various

independent algorithms in a portable pipeline [21]. One of the advantages of

ProkEvo is its ability to facilitate the addition and removal of new steps and

programs. For instance, more details about adding new program to ProkEvo are

given here https://github.com/npavlovikj/ProkEvo/wiki/4.1.-Add-new-

bioinformatics-tool-to-ProkEvo. By following these instructions, we were able to

successfully add stringMLST to the current ProkEvo platform. The ultimate

description of how stringMLST was integrated into ProkEvo can be found here,

https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/Prokevo_st

ringMLST.

Comparison between mlst and stringMLST performance using ProkEvo

In order to compare the performance/accuracy of MLST and stringMLST as part

of the ProkEvo platform, two subsets of the *C. jejuni*, *L. monocytogenes*, *S.*

Typhimurium and *S. aureus* datasets used in this paper were selected. One subset

was composed of 100 randomly selected genomes, while the second one

contained 1,000. The subsets were randomly selected from the original isolates

used in this paper. As part of ProkEvo, stringMLST was run with the default kmer

length of 35. The ProkEvo workflows with mlst and stringMLST and the two

datasets were individually run on Crane - one of the high-performance computing

clusters at the Holland Computing Center. Once the four workflows finished, the

performance of ProkEvo with mlst and stringMLST and the datasets with 100 and

1,000 genomes, respectively, was compared using: i) the total running time; ii) the

percentage of non-classified STs; and iii) the percentage of agreement between

programs. Since ProkEvo is an automated platform, a list of NCBI-SRA

identifications was provided with the ProkEvo implementations with both mlst

and stringMLST. In brief, ProkEvo manages all the dependencies and

intermediate steps, and produces the final ST classification as an output.

stringMLST-based kmer length optimization across phylogenetic divergent bacterial pathogens

In order to identify the optimal species-specific kmer length that minimizes the frequency of ST miscalls, we ran stringMLST with a range of different kmer lengths across phylogenetic divergent pathogenic species. First, we chose twenty-three *S. enterica* serovars (*S.* Agona, *S.* Anatum, *S.* Braenderup, *S.* Derby, *S.* Dublin, *S.* Enteritidis, *S.* Hadar, *S.* Heidelberg, *S.* Infantis, *S.* Javiana, *S.* Johannesburg, *S.* Kentucky, *S.* Mbandaka, *S.* Montevideo, *S.* Muenchen, *S.* Newport, *S.* Oranienburg, *S.* Poona, *S.* Saintpaul, *S.* Schwarzengrund, *S.* Senftenberg, *S.* Thompson, *S.* Typhimurium), and for each dataset we randomly selected 100 paired-end Illumina reads from NCBI-SRA. Second, for each dataset we ran mlst and stringMLST with kmer lengths ranging from 20, 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 90. The kmer length of 10 was excluded due to its poor performance in previous analyses. Additionally, we use data from fourteen pathogens with Public Health relevance to widen the scope of the analysis and assess the necessity of fine-tunning the kmer length on a more broadly selected collection of species. In particular, we chose the following pathogens:

Acinetobacter baumannii, Clostridioides difficile, Enterococcus faecium, Escherichia coli, Haemophilus influenzae, Helicobacter pylori, Klebsiella pneumoniae, Mycobacterium tuberculosis, Neisseria gonorrhoeae, Pseudomonas aeruginosa, Streptococcus pneumoniae, Campylobacter jejuni, Listeria monocytogenes, Staphylococcus aureus. For each pathogen, we randomly selected and downloaded 1,000 paired-end reads from NCBI-SRA and processed these

reads separately with mlst and stringMLST. stringMLST was run with kmer

lengths ranging from 20, 30, 35, 45, 55, 65, 70, 80, 90 and different schemes for

the different pathogens. Similarly to the *S. enterica* datasets, the kmer length of 10

was excluded from the analysis.

Across all datasets, the percentage of ST miscalls was calculated for

stringMLST for each kmer length, whereby miscalls were defined as "bad" ST

calls - calls with a missing or blank value. Next, for each dataset, the kmer length

that equated with the lowest percentage of ST miscalls was recorded. For some

datasets, multiple kmer lengths generated an identical lowest percentage for ST

miscalls. In this case, we applied a two-folded approach to select the most optimal

kmer length: 1) if kmer of length 35 was part of the kmer lengths that showed the

most optimal results, we recorded kmer 35 as the optimal kmer length since that is

the default and recommended value for stringMLST; or 2) if kmer of length 35

was not part of the kmer lengths that showed the most optimal results, we

recorded the kmer with the highest value as the most optimal one, since in general

our analysis showed that longer kmers consumed less computational resources

and speed up the entire analysis. Ultimately, the optimal kmer length and the

percentage of ST miscalls were visualized onto a core-genome phylogeny

generated for all twenty-three *S. enterica* serovars, as well as for all fourteen

pathogens including all twenty-three *S. enterica* serovars which jointly totaled

fifteen pathogens (total of 37 genomes, one per species including one per serovar

of *S. enterica*, were used to construct the core-genome phylogeny for

visualization purposes). The core-genome alignment was generated using Roary

with this set of parameters, "*roary -s -e --mafft -p 8 -cd 70 -i 70*

./prokka_output/*.gff -f roary_output"

(https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/scripts/ro

ary.submit).The phylogenetic tree was produced using FastTree [76] and

visualized using iTOL [77], and the recorded statistics were extract with custom R

scripts

(https://github.com/npavlovikj/MLST_stringMLST_analyses/blob/main/figures_c

ode/figures_code.Rmd).

In addition to calculating the percentage of ST miscalls for different kmer

lengths with stringMLST, for each dataset we calculated the percentage of

agreement (concordance) between mlst and stringMLST on ST calls ("good" or

"bad"), as previously described here. Of note, when the stringMLST and mlst

results were combined, the number of returned ST calls wasn't always 1,000 (the

original size of the used datasets). If 1,000 reads are used with stringMLST,

stringMLST generates ST calls for all 1,000 reads. On the other hand, when using

mlst, a set of steps are used before mlst, including filtering, and a fraction of

assemblies were disregarded due to poor quality. Thus, only genome sequences

that passed through the mlst program and yielded a "good" or "bad" call were

ultimately used to compare with stringMLST. The number of raw reads for each

dataset, as well as the number of final reads from mlst used for these analyses are

shown on S4 Table.

Statistical analyses

In order to compare the overall performance and accuracy of mlst vs. stringMLST on ST-based classifications, the following statistics were used across all bacterial species datasets: 1) ST richness; 2) Simpson's D index ($1 - D$) of diversity using ST counts as input data; 3) Proportion of non-classified STs (missing values or blank calls); and 4) Standard deviation of the proportion of non-classified STs. These statistics were calculated to evaluate the algorithmic performance on ST-based classification accuracy within and between bacterial species selected to be used in the narrow scope analysis (*C. jejuni*, *S. aureus*, *L. monocytogenes*, and *S. enterica*). ST richness was calculated by identifying the number of distinct STs present in each species. The Simpson's D index of diversity ($1 - D$) was used to calculate the degree of genotypic diversity across species, using the diversity() function available in the vegan (version 2.5-6) R library [29]. The proportion of non-classified STs was calculated using the counts of isolates or genomes that were not assigned a ST number after each run of either mlst or stringMLST. The standard deviation of the proportion of non-classified STs was calculated using the sd() function which is derived from an unbiased estimate of the sample variance corrected by $n - 1$ (n for number of observations). The frequency of genomes used for all analyses was calculated per batch and program across all species, including across serovars for *S. enterica*. The relative frequency of the most dominant ST lineages was also assessed across bacterial species.

PERMANOVA univariate or multivariate models were used to assess the degree of association between the genome-intrinsic and –extrinsic factors with the

following dependent variables: ST richness, Simpson's D index of diversity, or

proportion of non-classified STs. Statistical models were built for each of the

dependent variables separately. Multivariate models included either the

combination of bacterial species and program, or serovars in the case of *S.*

enterica and program. These multivariate models were stated to calculate the

main and synergistic effects of the explanatory variables (e.g., species*program or

serovar*program). Univariate models were also assessed for each of the

dependent variables, using one of the following independent/explanatory

variables: 1) Genome-intrinsic variables: median number of contigs, mean of the

total count of nucleotides per genome, mean of the average GC% content per

genome, standard deviation of the number of contigs, standard deviation of the

total count of nucleotides per genome, and standard deviation of the average

GC% content per genome; 2) Genome-extrinsic variables: species, serovar of *S.*

enterica, program (mlst vs. stringMLST with kmer lengths of 10, 20, 30, 35, 45,

55, 65, 70, 80, 90), mean of the total count of unique STs per program, mean of

the total count of unique alleles across all genes per program, and the Simpson's

D index of diversity per species. Statistical significance and strength of

association between the dependent and independent variables were measured with

p-values ($p < 0.05$) and *R*-squared, respectively. In the case of contig size

(median), total number of nucleotides per genome (mean), and GC% content per

genome, summary statistic values (median or mean) were calculated grouped by

species and batch (there was a total of three batches per bacterial species or

serovar). For the total count of STs and total number of alleles in the database,

summary statistic values (mean) were calculated grouped by species, batch, and

program. Lastly, the standard deviation of number of contigs, total count of

nucleotides per genome, or GC% content per genome were calculated grouped by

species. PERMANOVA models were run using the adonis() function with 1,000

permutations using the vegan (version 2.5-6) R library [29]. Principal component

analysis (PCA) was used to analyze the dinucleotide distribution across species

and across serovars for *S. enterica* with two dimensions using the prcomp()

function. The PCA calculations and the selection of the number of PCs were done

using the factoextra (version 1.0.7) library. Bar-plots, box-and-whiskers plots, and

bivariate/trivariate scatter plots were used to assess the distribution and

associations within and between dependent and independent/explanatory

variables. The R software (version 4.0.3) and R libraries such as Tidyverse

(version 1.3.0) were used to conduct all statistical analyses, and all R scripts are

available here

(https://github.com/npavlovikj/MLST_stringMLST_analyses/tree/main/figures_c

ode). Data quality control was achieved with R base functions, in addition to the

following packages: skimr (version 2.1.3) and visdat (version 0.5.3). Graphical

visualizations were achieved using ggplot2 (version 3.3.2), GGally (version

2.1.2), and plotly (version 4.9.4.1). R code integrity was checked using the

assertive (version 0.3-6) package.

Results

The computational and analytical approaches used in this paper are shown on Fig

1. Our analytical approach was sub-divided into a narrow- and wide-scope

analysis aiming at accomplishing two goals: 1) Comparing the computational and

statistical performance of mlst vs. stringMLST; and 2) Optimizing the use of

stringMLST on a bacterial species basis and ultimately implementing it as part of

the ProkEvo computational genomics platform. First, we used freely available raw

Illumina paired-end sequence data from *C. jejuni*, *L. monocytogenes*, *S. enterica*

and *S. aureus*, to run stringMLST and mlst independently in order to compare the

accuracy in ST-based classifications and assess the computational needs and

performance in the overall analysis (narrow-scope step). In particular, for this

narrow-scope stage we performed a detailed comparative analysis between these

two programs including: i) analyses of computational performance and resources

needed (e.g., average runtime per genome and maximum memory needed to

analyze all genomes), and ii) statistical analyses to determine the accuracy of

classifications (e.g., ST richness, Simpson's D index of ST-based diversity,

proportion of miscalls, and percentage of agreement or concordance between

programs). For the wide-scope step, we systematically analyzed the accuracy and

concordance between mlst and stringMLST across a broad array of phylogenetic

divergent pathogens with direct implication for Public Health (*Acinetobacter*

baumannii, Clostridioides difficile, Enterococcus faecium, Escherichia coli,

Haemophilus influenzae, Helicobacter pylori, Klebsiella pneumoniae,

Mycobacterium tuberculosis, Neisseria gonorrhoeae, Pseudomonas aeruginosa,

Streptococcus pneumoniae, Campylobacter jejuni, Listeria monocytogenes,

Salmonella enterica and *Staphylococcus aureus*). Combined with the intra-species

analysis done across 23 serovars of *S. enterica*, our assessment aimed at revealing

the optimized kmer length to be used with stringMLST in order to: i) minimize

the percentage of ST miscalls, and ii) maximize the use of computational

resources by speeding up the analysis. Lastly, we provided an implementation of

stringMLST within ProkEvo - a freely available and scalable computational

platform that facilitates hierarchical genotyping of bacterial populations including

pan-genomic mapping [21].

Computational performance

The computational performance between stringMLST and mlst was measured

using two metrics: 1) The average computational runtime per genome; and 2) The

maximum memory used per dataset. The average runtime in minutes per genome

per batch between mlst and stringMLST with different kmer lengths, for *C. jejuni*,

L. monocytogenes, *S. aureus*, and *S.* Typhimurium (major representative of *S.*

enterica), is shown on Fig 2. While the runtime of mlst varies between 20 and 80

minutes per genome depending on the dataset used, all stringMLST runs with

different kmers finished within a few minutes (ranging from ~1-16 minutes when

kmer 10 was included and ~1-5 minutes when kmer 10 was excluded). Apart from

stringMLST with kmer 10, all other kmer lengths showed a uniform runtime. The

longer runtime observed with kmer 10 can be partially explained by the higher

number of kmers that were generated and used for mapping (S1 Fig). The

obtained results show that ST-based classifications are accomplished considerably

more rapidly when carried out using stringMLST compared to the standard MLST

program.

Additionally, a comparison of maximum memory used when both stringMLST

and mlst were run for *C. jejuni*, *L. monocytogenes*, *S. aureus*, and *S.* Typhimurium

(major representative of *S. enterica*) is shown on S2 Fig. Across all species, the

range of maximum memory usage for mlst and stringMLST (across all kmers)

was ~2-16GBs and ~3-30GBs respectively. Although the memory used across

datasets is variable, none of the analyses we ran exceeded 30GBs of RAM. Since

most high-performance computers can consistently provide resources from 32GBs

to a few TBs of RAM, the memory available should not be considered a

bottleneck for running either program.

Factors that can influence ST-based classification

First, we describe the characteristics and composition of the data utilized for

comparison between programs regarding ST-based classification in the narrow-

scope approach (utilization of fewer phylogenetic diverse pathogen datasets). The

frequency of genomes utilized per species and across programs is shown on S3A-

D Fig. The frequency of *S. enterica* genomes was higher than other species

because an equal sample of ~600 genomes was taken from 20 representative

zoonotic serovars (S4A-D Fig). An assessment of the proportion of the most

dominant STs across species (proportion $\geq$ 2% - S5A-D Fig) or serovars of *S.*

enterica (proportion $\geq$ 15% - S4N Fig) initially revealed a similar ST-based

distribution across programs. Furthermore, genome-intrinsic and -extrinsic factors

that could potentially impact the mlst vs. stringMLST algorithmic comparison and

performance were *a priori* determined and considered in the analysis. Among the

genome-intrinsic factors considered across species were the number of contigs per

genome (Fig 3A), the total number of nucleotides per genome (Fig 3B), GC%

content per genome (Fig 3C), and the distribution and composition of

dinucleotides per species (Fig 3D and S3E-F Fig). Similarly, the distribution of

the genome-intrinsic factors was analyzed across all twenty serovars of *S. enterica*

(S4G-L Fig). A correlogram (pairwise correlation analysis) was also used to

assess the bivariate correlation (Pearson's correlation coefficient) across genome-

intrinsic variables, for either all four bacterial species (S3G Fig) or serovars

across *S. enterica* (S4M Fig). At large, the differences observed in the distribution

of genomic-intrinsic variables were species driven, with a strong uniformity found

across serovars of *S. enterica*.

As for the genome-extrinsic variables, the total count of unique STs (for

species - Fig 3E) and unique number of alleles across all seven loci (for species -

Fig 3F), across all batches, were selected as factors that could influence the

comparative analysis between mlst and stringMLST. Similarly, the genome-

extrinsic variables were analyzed across all twenty serovars of *S. enterica* (S4E-F

Fig). Of note, database size differences (number of STs and alleles) may directly

influence the number of miscalls since it is expected that the larger the database

is, the more likely STs are to be classified, or to find a match, and not be

miscalled [30]. Considering the differences in genome-intrinsic and -extrinsic

variable distribution across species, such factors were further utilized for

assessing their statistical contribution in the accuracy of ST-based classification

between mlst vs. stringMLST.

Assessing the contribution of genome-intrinsic and –

extrinsic variables

In order to assess the statistical association and contribution of each genomic-

intrinsic and -extrinsic variable onto the accuracy of mlst vs. stringMLST on ST

calls (narrow-scope analysis since it only included four bacterial species, *C.*

jejuni, *S. aureus*, *L. monocytogenes*, and *S. enterica*), the following dependent

variables (outcomes) were used in the PERMANOVA models: 1) ST richness

(Fig 4A); 2) Simpson's D index of ST diversity (Fig 4B); and 3) Proportion of

non-classified STs (Fig 4C). Additionally, the standard deviation of the proportion

of non-classified STs was measured as an auxiliary metric for accuracy (Fig 4D).

At the species level, a multivariate model was used to examine the interaction of

species and program (mlst vs. stringMLST); whereas, the remaining analyses

were done using univariate models containing each genome-intrinsic and -

extrinsic variable for all three outcomes (S6A-L Fig, S7A-K Fig, S8A-L Fig).

For each variable, the significance and strength of association were assessed by

jointly examining the p-value ($p < 0.05$) and R-squared, respectively. For both ST

richness (Fig 4A) and the Simpson's D index of diversity (Fig 4B), the difference

between species explained the majority of the variation with ~98.3% and ~99%,

respectively. As expected, based on the phylogenetic divergence of the four

chosen pathogens, differences across species could largely be explained by

genome-intrinsic variables associated with genome composition, such as: GC%

content (p ~0.0009, R-squared ~44%) for ST richness, and the number of contigs

per genome (p ~0.0009, R-squared ~39.5%) for the Simpson's D index of

diversity (Fig 4A-B). Notably, for both ST richness and the Simpson's D index of

diversity most of the differences between species could be explained by variation

in genome composition (Fig 4A-B). Not surprisingly, co-linearity was observed

between ST richness and the Simpson's D index of diversity across species (Fig

4A). In the case of the proportion of non-classified STs (ST miscalls) (Fig 4C),

most of the variation was explained by inter-species differences (p ~0.0009, R-

squared ~33%), with the number of contigs per genome being the most important

genome-intrinsic contributing factor (p ~0.0009, R- squared ~27%). As for the

kmer length parameter used by stringMLST, results for ST richness and the

Simpson's D index of diversity were uniform across all lengths (Fig 4A-B).

However, when examining the proportion of miscalls (Fig 4C) and the standard

deviation of that proportion (Fig 4D), the data pointed toward the optimal kmer

length being between 35 and 65 across all four species (narrow-scope analysis).

Specifically, this kmer length range was defined based on two criteria: i)

minimization of the proportion of miscalls; and ii) less variation (standard

deviation) around the average of ST-based miscalls. Of note, mlst has the highest

proportion of miscalls and standard deviation of that proportion for both *L.*

monocytogenes and *C. jejuni* (Fig 4C-D), and the kmer length 10 for stringMLST

yielded very low accuracy and null results for ST richness and Simpson's D index

of diversity (Fig 4A-D). Differences between species across ST richness,

Simpson's D index of diversity, and proportion of ST miscalls along with all

genome-intrinsic and -extrinsic variables across programs (mlst vs. stringMLST)

were further examined here (Fig 5A-D, S9A-O Fig). Nonetheless, differences in

ST-based calls across programs were largely influenced by the bacterial species

dataset.

Given the complexity and diversity of the *S. enterica* population structure [12],

the stringMLST performance was analyzed across twenty zoonotic serovars

(S4O-R Fig), and resulted in a significant and predominant contribution of the

"serovar groupings" across all outcomes and PERMANOVA models (S10A-L

Fig, S11A-K Fig, S12A-L Fig): ST richness (p ~0.0009, R-squared ~75.4%),

Simpson's D index of diversity (p ~0.0009, R-squared ~88%), and proportion of

ST miscalls (p ~0.0009, R- squared ~35.4%). By assessing the distribution of the

model outcomes, along with PERMANOVA model results and bivariate

association between dependent and explanatory variables (S13A-R Fig), the

results recapitulated the species-level results with the optimal kmer length for

stringMLST being around 35 and 65, but also revealed the need to consider

difference across *S. enterica* serovars prior to implementation. Combined, these

accuracy-based results suggest that: i) stringMLST minimizes the ST miscalls

compared to mlst in a species-specific fashion, and by consequence the optimal

kmer length for stringMLST ranged from 35 to 65 overall; ii) the performance

and accuracy of stringMLST can vary across species and serovars of *S. enterica*

allowing for data-driven fine-tunning of the kmer length; and iii) the use of

sequence platform with longer reads which would maximize the number of

contigs per genome could directly alter both mlst and stringMLST accuracy in ST

calls across species.

Concordance between programs

Concordance between programs was calculated as the percentage of cases in

which outputs from both mlst vs. stringmlst agreed in the call ("good" or "bad").

Results demonstrating the percentage agreement in ST calls between mlst and

stringMLST with different kmer lengths are shown on Fig 6. With the exception

of kmer 10, across all species, the percentage of agreement between mlst and

stringMLST varies between 81.50% and 97.50%. In the case of *L.*

monocytogenes, *C. jejuni*, and *S. aureus*, the kmer length of 35 appears to be the

optimal value to reach the same accuracy as mlst, which matches the original

default and recommended parameter value for stringMLST [17]. However, for *S.*

enterica a higher percentage of agreement with MLST was achieved for kmer

lengths of 55 and 65 (Fig 6). This *S. enterica*-related observation recapitulated the

initial findings of decreased proportion of ST miscalls with higher kmer lengths

(Fig 4C). Of note, our finding collectively showed that the kmer length of 10

yielded low accuracy when compared to mlst and other stringMLST kmer lengths.

The most likely explanation for lower accuracy generated by kmer 10 is that

shorter kmers are more likely to map unambiguously onto a genome when

compared to other lengths. That high frequency of kmer length 10 on a given

dataset reflects their higher likelihood of mapping to multiple regions of a genome

(S1 Fig, S14A-B Fig). Overall, stringMLST is a rapidly deployable and

optimizable ST-based genotyping algorithm that in this narrow-scope analysis

proved to be applicable to four phylogenetic distinct pathogens.

778   Optimization of stringMLST kmer length across

779   phylogenetic divergent species

As previously proposed [17], the default kmer length for Illumina paired-end

reads for stringMLST is 35. However, our narrow-scope analysis across four

distinct pathogens (see above) suggested a species-specific variation in the

optimal kmer length capable of minimizing the proportion of ST miscalls.

Therefore, we systematically investigated what kmer length would give the fewest

ST miscalls (optimized length) with stringMLST across a diverse array of

phylogenetic divergent pathogens. Given our previous results, we first deepened

our investigation into the *S. enterica* population given the genetic and ecological

diversity across serovars. For that, we selected data from twenty-three *S. enterica*

zoonotic serovars and ran stringMLST with wide range of kmer lengths (20, 30,

35, 40, 45, 50, 55, 60, 65, 70, 80, 90). Fig 7A shows the core-genome phylogeny

mapping of the optimized kmer length across all twenty-three serovars along with

their corresponding percentage of ST miscalls. More detailed information on the

distribution of the percentage of ST miscalls for all used kmer lengths (20, 30, 35,

40, 45, 50, 55, 60, 65, 70, 80, 90) is shown on S15A Fig. As it can be seen on Fig

7A, many serovars (*S.* Anatum, *S.* Braenderup, *S.* Javiana, *S.* Mbandaka, *S.*

Montevideo, *S.* Oranienburg, *S.* Poona, *S.* Schwarzengrund, *S.* Senftenberg, *S.*

Typhimurium) have 0% of miscalls when the default kmer length 35 was used. *S.*

Infantis and *S.* Derby show the lowest percentage of ST miscalls (3% and 2%

respectively) with higher value of kmer, e.g., 90. Interestingly, *S.* Saintpaul

showed the highest percentage of ST miscalls when only considering the range of

kmer lengths used for the initial analyses (10-90). To investigate this further, we

ran stringMLST for *S.* Saintpaul with kmer lengths up to 240 (240 was chosen

because the maximum read length for the *S.* Saintpaul dataset is 250 base pairs or

nucleotides) (S15C-D Fig). As it can be seen on S15C Fig, the fewest ST miscalls

for *S.* Saintpaul were produced when kmer of length 140 was used (22%). When

comparing the percentage of ST miscalls between mlst and stringMLST, mlst

outperformed stringMLST for the used datasets and range of kmer lengths. In

addition to the percentage of ST miscalls, we calculated the percentage of ST

agreement between mlst and stringMLST with the range of kmer lengths (S15B

Fig). While for some serovars this percentage is the highest when kmer with

length 35 is used (e.g., *S.* Anatum, *S.* Braenderup, *S.* Javiana, *S.* Mbandaka, *S.*

Montevideo, *S.* Oranienburg, *S.* Poona, *S.* Schwarzengrund, *S.* Senftenberg, *S.*

Typhimurium), for other serovars (e.g., *S.* Derby, *S.* Dublin, *S.* Enteritidis, *S.*

Hadar, *S.* Heidelberg, *S.* Infantis, *S.* Kentucky, *S.* Saintpaul) the percentage of ST

agreement between the two programs was higher with higher kmer lengths.

In order to widen the scope of our phylogenetic-based analysis, we assessed the

percentage of ST miscalls across varying kmer lengths for divergent bacterial

pathogens with Public Health relevance. We selected 14 distinct organisms and

ran stringMLST with wide range of kmer lengths (20, 30, 35, 45, 55, 65, 70, 80,

90). Fig 7B depicts the core-genome phylogeny mapped results including the

optimal kmer length that minimized the percentage of ST miscalls. Of note, the

phylogeny contained fourteen distinct pathogens and twenty-three genomes across

each serovar of *S. enterica*. The distribution of the percentage of ST miscalls for

all used kmer lengths (20, 30, 35, 40, 45, 50, 55, 60, 65, 70, 80, 90) is shown on

S15E Fig. While the percentage of ST miscalls varied between 0% and 22%

across the *S. enterica* serovars as shown in Fig 7A, the percentage of miscalls is

more variable for the fourteen bacterial pathogens, ranging from 1.2% to 74.9%.

The datasets for *A. baumannii, C. jejuni, H. influenzae, K. pneumoniae, L.*

monocytogenes, N. gonorrhoeae, S. aureus and *S. pneumoniae* showed the lowest

percentage of ST calls with the default kmer length of 35. *C. difficile* and *M.*

tuberculosis had minimized ST miscalls with kmer lengths of 20 and 30

respectively, while *P. aeruginosa* with kmer length of 65. Interestingly, for *E.*

faecium and *H. pylori*, the optimal kmer lengths were 35 and 20, even though the

percentage of miscalls was high (74.9% and 67.6%). To further investigate this,

we ran stringMLST for *E. faecium* and *H. pylori* with kmer lengths up to 140 (140

was chosen because the maximum read length for the two datasets is 150 base

pairs or nucleotides) (S15G-H Fig, S15K-L Fig). As it can be seen on the Figures,

the percentage of miscalls was higher with higher kmer lengths, and the lower

kmer lengths yielded fewer miscalls, even though this number was still high.

Additionally, we ran stringMLST on another set of randomly selected 100 paired-

end reads for *E. faecium* (S15M-N Fig), *H. pylori* (S15I-J Fig) and *Enterococcus*

faecalis (S15O-P Fig). These 100 reads were not part of the initial datasets and

were chosen to validate that the initial random data selection was not completely

biased. We also added *E. faecalis* here due to its close phylogenetic association

with *E. faecium*. For *E. faecium* and *H. pylori* we observed the same pattern with

100 reads as with 1,000 reads. On the other hand, the pattern for *E. faecalis* was

quite opposite and as expected, with lowest percentage of ST miscalls of 5.43%

for kmer 35. When comparing the percentage of ST miscalls between mlst and

stringMLST, for some datasets, such as *H. pylori*, *C. jejuni*, *L. monocytogenes*, *M.*

tuberculosis, *N. gonorrhoeae*, *S. aureus*, mlst performed worse than stringMLST.

In addition to the percentage of ST miscalls, we calculated the percentage of ST

agreement between mlst and stringMLST with the range of kmer lengths (S15F

Fig). Of note, in the case of stringMLST, when the optimal kmer length was

above the default parameter of 35, the ultimately selected kmer length was picked

based on our empirical evidence for longer kmers being capable of speeding up

the computational analysis.

In summary, while the default kmer length of 35 used by stringMLST performs

accurately across many organisms, our systems-based approach that encompassed

the analysis of a variety of phylogenetic divergent organisms revealed: i) intra-

and inter-species variation in the percentage of ST miscalls requires fine-tunning

of the kmer length parameter; ii) lack of association between taxonomy or

phylogenetic placement of organisms and the optimal kmer length; and iii) unique

species behave as outliers for which stringMLST cannot be directly applied with

the default settings.

Incorporating stringMLST in ProkEvo

ProkEvo was recently developed as an automated and scalable computational

platform for bacterial population genomics analyses that uses the Pegasus

Workflow Management System (WMS) [31] that allows for distributed use on

different computational platforms and rapid integration of novel programs [21]. In

particular, ProkEvo facilitates the use of a hierarchical approach for population

stratification with different layers of genotypic resolution. MLST-based

classification of genomes into STs is part of this hierarchical approach that has

been proven to be predictive of ecological traits such as AMR in *S. enterica*

lineages [57]. However, ProkEvo currently only uses the standard mlst algorithm

for ST calls [21]. As part of this paper, the stringMLST program was incorporated

into ProkEvo without any disruption in its workflow. The workflow design of

ProkEvo with both mlst and stringMLST is shown on S16 Fig.

In order to compare the performance of ProkEvo with mlst and stringMLST,

randomly shuffled subsets derived from the original datasets used for *C. jejuni*, *L.*

monocytogenes, *S.* Typhimurium, and *S. aureus* were used. One random subset

contained 100 genomes, while the second one had 1,000 genomes. ProkEvo was

run using either mlst or stringMLST on Crane, one of the high-performance

computing clusters at the Holland Computing Center [23]. For mlst, the pipeline

used was previously established and included a few required steps, such as quality

trimming and adapter clipping, *de novo* assembly and assembly discarding prior

to the ST mapping [21]. Based on the inter-species results shown here (Fig 6), the

default kmer length of 35 was used with stringMLST for this comparison. The

outcomes measured for this analysis were: i) total running time (Fig 8A); ii) the

percentage of non-classified STs (Fig 8B); and iii) the percentage of agreement

between programs (Fig 8C).

While the runtime of using ProkEvo with mlst varied from ~8 to 34 hours for

the subset containing 100 genomes, the runtime of ProkEvo with stringMLST

varied from ~25 minutes to 3 hours (Fig 8A). Similarly, for the larger datasets

containing 1,000 genomes, the runtime of ProkEvo with mlst varied from ~17 to

39 hours, while the runtime of ProkEvo with stringMLST varied from ~4 to 8

hours. Regardless of the pathogen species tested, stringMLST speeded up the

analyses ~4 times when utilizing 1,000 genomes across species.

In terms of accuracy in ST classifications, the use of stringMLST considerably

decreased the number of non-classified STs, regardless of the dataset size (100 or

1,000 genomes) and bacterial species (Fig 8B). In accordance, stringMLST

resulted in a higher frequency of genomes classified as novel STs (ST numbers

that were not classified by mlst) (S17 Fig). Additionally, the overall concordance

between mlst and stringMLST varies from 82% to 100% across all datasets. The

percentage of agreement is the lowest for *S.* Typhimurium, while it is the highest

for *S. aureus* (Fig 8C). The lower proportion of ST miscalls and high percentage

of agreement between programs for *S. aureus*, compared to other species, is

associated with its higher degree of genetic homogeneity (fewer dominant STs)

(S5 Fig). This difference in miscalls and concordance between programs may be

further explained by the variation in database sizes, since the PubMLST schemes

used for mlst have fewer alleles across all seven loci which results in fewer STs

compared to stringMLST as shown on S3 Table.

Previously, the scalability of ProkEvo was assessed by a comparative analysis

of its computational performance on Crane and OSG, using two datasets with

2,392 and 23,045 genomes each (10 X difference), and the standard mlst approach

for ST calling [21]. To further demonstrate the gain in computational runtime

obtained with the use of stringMLST within ProkEvo, the complete $S.$

Typhimurium dataset containing 23,045 genomes was run on OSG. While

ProkEvo with mlst finished all ST calls in 26 days and 6 hours when OSG was

used as a computational platform [21], ProkEvo with stringMLST completed the

task in 3 days and 6 hours. Altogether, stringMLST provides an accurate and

rapid alternative to mlst for scalable ST genotyping that is portable to be

implemented in any high-performance and high-throughput platform, with its use

being further facilitated by its implementation in ProkEvo.

Discussion

The incorporation of WGS technology has advanced the study of bacterial

populations, since it has facilitated genotyping at different levels of resolution,

which in turn has proven to be predictive of, or associated with, inferable

ecological traits or epidemiological patterns [21] [32][33][34][35][36][37][38]. In

particular, the use of a hierarchical population structure analysis allows for

ancestral relationships and patterns of diversification to be inferred, while

determining the most important informative genotypic unit to be tracked over
time [21][39][40][41][12][42][43]. ST-based classification is an integral part of
the hierarchical genotyping approach [21][27]. ST lineages are formed based on
the utilization of allelic mapping across seven genome scattered loci that are
ubiquitously present across phylogenetic divergent bacterial species [8][9][44].
Such ST lineages can be further combined in clonal complexes, when sharing five
or more of the seven loci combinations - also called eBURST groups (eBG)
[10][45]. Thus, ST-based genotyping is widely used for a variety of reasons,
including: i) classification of genomes below the species level [8][9][44]; ii)
providing stable informative genotypic unit that can be used intra- and inter-
laboratory for mapping and tracking of populations [8][9][44]; iii) predictability
of ecological traits such as serovar in the case of *S. enterica*, and AMR across
bacterial species due to the linkage disequilibrium between MLST and accessory
loci [13][14][15][57]; and iv) inferring ancestral relationships through eBG
profiles [10][45]. ST-based genotyping is typically dependent on genome
assembly, which is efficient and accurate but not scalable and of rapid turn-
around [30][22]. However, stringMLST is a program capable of rapidly
classifying genomes into STs independently of genome assemblies [17]. Yet, a
systematic and scalable comparison between the standard MLST and stringMLST
programs is lacking [16]. Therefore, this study sought to comprehensively assess
the computational performance and accuracy of mlst vs. stringMLST across
phylogenetic divergent bacterial pathogens with direct implication for Public
Health. Additionally, this algorithmic comparison was designed to consider the

intra- and inter-species variation, in addition to the statistical contribution of

genome-intrinsic and -extrinsic factors on classification accuracy, aiming at

identifying actionable approaches that may be used to further optimize the

implementation of stringMLST.

Characterization of bacterial pathogens and performing molecular typing

provides valuable epidemiological information important for Public Health

agencies. There are multiple tools available for MLST classification, such as mlst

[22], ARIBA **Error! Reference source not found.**, stringMLST [17],

MentaLiST **Error! Reference source not found.**, STing **Error! Reference**

source not found.. In general, the available tools can be categorized based on the

input data they use - some tools use raw Illumina paired-end sequence data, while

others use *de novo* assemblies [16]. In order to generate the *de novo* assemblies, a

few pre-processing steps need to be performed, such as quality control, trimming,

assembly and filtering, that can be costly and require lots of computational

resources, such as memory and time. Using raw sequence data for ST-based

classification has a tremendous advantage especially in pathogen surveillance,

since all the costly steps prior to the *de novo* assembly are bypassed and the STs

calls are made as the sequence reads are generated. mlst uses *de novo* genome

assemblies as an input and performs mapping in order to align sequences to pre-

downloaded allelic files across all target loci. ARIBA identifies AMR-associated

genes, single nucleotide polymorphisms and ST calls using Illumina paired-end

raw sequencing reads. ARIBA clusters the raw reads by mapping them to genes,

and then performs local assembly within clusters to identify AMR genes and ST

calls. On the other hand, stringMLST and MentaLiST rely on kmer matching

between raw sequence reads and available ST schemes that allows for fast

mapping and ST-based typing. Both tools are shown to be accurate and fast for

standard MLST classification, while providing comparable accuracy with

MentaLiST albeit using less computational resources **Error! Reference source**

not found.. STing is the successor of stringMLST - it uses the same algorithmic

approach with additional computational applications for large MLST schemes

such as ribosomal MLST (rMLST) and core-genome MLST (cgMLST) **Error!**

Reference source not found.. All these tools have integrated ST schemes and/or

provide utilities for downloading the available PubMLST databases. There are a

few available comparisons of such tools for ST classification, mostly focusing on

the computational resources used and the percentage of correctly classified STs

[16]**Error! Reference source not found.Error! Reference source not found.**.

When tools were tested with real outbreak datasets (*L. monocytogenes*, *E. coli*, *C.*

jejuni, *S. enterica*) comprising 85 samples, stringMLST showed the fastest

running time of 80.8 minutes and high accuracy in ST calls (100%) [16]. While

MentaLiST does not scale well when reads with high coverage are used, it

performs well on MLST schemes with up to a few thousand genes and alleles,

such as cgMLST (~3,000 genes) **Error! Reference source not found.**. While

most ST tools perform satisfactorily, there are some relevant bottlenecks to be

considered. For example, some tools use out-of-date MLST databases that require

manual curation, and can directly affect the accuracy of ST calls, especially when

mixed and low coverage samples are used [16]. ST tools that are assembly and

alignment free, such as stringMLST, STing and MentaLiST, show quite a few

advantages in term of accuracy and efficiency that make them applicable for real-

time molecular epidemiology and surveillance. Thus, we chose stringMLST as a

representative of the kmer-based ST tools to perform a systems-based

comparative analysis that assess the computational and statistical efficacy of ST

calls across divergent pathogens in contrast to the legacy MLST approach.

As shown here, the stringMLST accuracy can be affected by the species being

tested without any specific phylogenetic patterns. In particular, the choice of kmer

length used directly impacts the proportion of ST miscalls across species, and in

certain cases it may not be applied as designed even after parameter tunning,

which is likely a reflect of their varying population structure and pattern of

genome diversification and architecture (e.g., horizontal gene transfer (HGT), and

acquisition of mobile elements such as prophages and insertion sequences, etc.)

[12][41][68][69][70][71][72][73]. A clear example is *S. enterica*, for which the

accuracy of stringMLST varied across ecologically distinct serovars that are

known to have unique pan-genomic composition as exemplified by their

predictive prophage distribution [12][71][74][75]. To the best of our knowledge,

the currently available comparisons between ST tools have not considered any

systematic approach for parameter tunning across phylogenetic divergent

species known to vary in population structure [21][27][67].

In evaluating genome-intrinsic and -extrinsic variables that could contribute to

differences in accuracy between mlst and stringMLST, it was found that the

species level variation was mostly explained by the uniqueness of their genomic

composition and number of contigs per genome. As genomic composition is an

inheritable property of the bacterial species and reflects their evolutionary history

and speciation patterns, this association with algorithmic performance was

somewhat expected [46][47]. However, the contribution of the number of contigs

making the overall difference between programs poses forth the hypothesis that

by using sequencing platforms that generate longer reads, such as PacBio and

Oxford Nanopore Technologies (ONT), both mlst and stringMLST accuracy in

ST calls may be considerably altered in species-specific fashion, whereby

accuracy would be expected to improve if HGT occurs at high rates. However,

these sequencing technologies produce reads with lower accuracy (~80-90%) that

may inflate the number of false allelic calls and consequently alter the distribution

of STs - likely this would split major STs into sub-populations [48]**Error!**

Reference source not found.Error! Reference source not found.. Therefore,

while more work is needed in this field, current studies using hybrid assembly

approaches of both Illumina short reads and ONT long reads **Error! Reference**

source not found., as well as only polished ONT reads **Error! Reference source**

not found. for performing ST-based classification showed promising cost-

effective results for this kind of molecular typing. Hence, we expect that

stringMLST, or its successor STing, will be optimized for their implementation

with longer read sequencing platform such as PacBio and Oxford Nanopore

Technologies [14][49], which will in turn facilitate real-time surveillance of

pathogens using hierarchical genotypes such as ST calls.

While the kmer length of 35 is currently recommended as a default value of

stringMLST, our systems-based approach demonstrated that for specific bacterial

species it will result in increasing the frequency of ST miscalls which in turn may

hinder epidemiological investigations. Across phylogenetic divergent pathogenic

bacterial species, the optimal kmer length ranged from 20 to 140, regardless of

their ancestral relationship or speciation pattern. The varying population structure,

the pattern of genome diversification and architecture (e.g., impact of HGT), as

well as sequence coverage may be some of the reasons underlying the observed

statistics [12][41][68][69][70][71][72][73]. Although we hypothesize that longer

sequence reads will help overcome this limitation, there is still a context-

dependent consideration for parameter tunning and overall algorithmic

implementation. Therefore, in the case of stringMLST, we suggest the following

actionables to maximize its utilization, including: i) developers to consider

implementing a pre-step that heuristically searches for the optimal kmer length

(minimizes ST miscalls) in dataset-dependent fashion (sampling from the testing

data), perhaps even by comparing with the standard MLST as positive controls;

and/or ii) researchers to run wide range of kmer lengths on a subset of the dataset

in order to select the optimal kmer length that minimizes the percentage of ST

miscalls. Given the speed and scalability of stringMLST, using multiple kmer

lengths is not likely to add much overhead to the analyses, and this provides an

empirical statistical approach for kmer selection and optimization of ST

classifications. With this data-driven fine-tunning of the kmer length, stringMLST

is a powerful program that can be efficiently and effectively used in

microbiological and epidemiological laboratories.

We recently developed ProkEvo, a freely available scalable platform for

performing hierarchical-based bacterial population genomics analyses [21].

ProkEvo: 1) uses the Pegasus Workflow Management System to ensure

reproducibility, scalability, and modularity; 2) uses high-performance and high-

throughput computational platforms; 3) automates and scales multitude of

computational analyses of a few to tens of thousands of bacterial genomes; 4) can

run many thousands of analyses concurrently if the computational resources are

available; 5) is easily modifiable and expandable platform that can incorporate

additional algorithmic steps and custom scripts. The initial implementation of ST-

based classifications through ProkEvo, as part of a hierarchical genotyping

strategy to map and track populations, was done using the assembly-dependent

MLST program [21][22]. Running mlst inside ProkEvo allows for parallelization

of the genome assemblies (run per isolate or genome) which enhances scalability

and facilitates the optimal use of computation resources. Theoretically, if there

are n isolates and n cores available on the computational platform, ProkEvo can

linearly utilize all resources and run all n independent tasks simultaneously.

Typically, ST-based classifications are time consuming because the mapping

process is run sequentially in a set of genomes instead of running them

independently. Thus, using modular and distributed platforms such

as ProkEvo for performing ST-based genotyping provides great benefit, especially

if additional features such as other hierarchical genotypes and pan-genomic

mapping tools are part of the same platform [21]. As part of this work, we

modified ProkEvo to not only offer the standard assembly-dependent MLST

mapping approach, but it now contains stringMLST, and our tests showed a

significant speed-up in runtime for datasets ranging from a few hundreds to tens

of thousands of genomes. To use ProkEvo with stringMLST, the researcher only

needs to provide a list of SRA identifications and run the submit script without

any advanced experience in high-performance or high-throughput computing.

Depending on the configuration set, ProkEvo can use locally downloaded

sequence data or download the data from NCBI directly. The Pegasus Workflow

Managements System that is used by ProkEvo automatically handles the

dependencies, as well as all the intermediate and final files. Thus, using platforms

such as ProkEvo with fast tool for hierarchical genotyping, such as stringMLST,

allows for robust and efficient population-based genomics analyses that facilitate:

i) mapping and tracking of variants or lineages for epidemiological inquiries; ii)

population structure analysis; and iii) ecological trait prediction using pan-

genomic mapping to specific genotypes.

1108

Conclusion

In conclusion, stringMLST largely proved to be an accurate, rapid, and scalable

tool for ST-based classifications that could be readily implemented in

microbiological laboratories and epidemiological agencies. Notably, this

comprehensive analysis of stringMLST across phylogenetic divergent bacterial

pathogens, with varying degrees of clonality, revealed the potential for enhancing

its accuracy by parameter tunning (kmer length) in a dataset-dependent fashion.

Specifically, we propose that the kmer length can be optimized in two ways on a

case-by-case basis: 1) intrinsically by implementing a pre-step inside the

algorithm to sample from the target data and select the optimal kmer length; or 2)

by the user through a heuristic data mining approach to select the optimal kmer

length prior to finalizing the ST calls. Also, by assessing genome-intrinsic and -

extrinsic factors that could affect the stringMLST performance, our work suggests

that longer sequence reads have the potential to improve its accuracy for specific

bacterial species. Furthermore, the integration of stringMLST into ProkEvo

allows users to take advantage of other hierarchical genotyping strategies,

including pan-genomic mapping, which reproducibly facilitates ecological and

epidemiological inquiries at scale. Ultimately, this work emphasizes the

importance of developing robust algorithmic tools for mining WGS data that can

have direct implications for mapping and tracking of bacterial populations.

1129

Acknowledgements

This work was completed by utilizing the Holland Computing Center of the

University of Nebraska, which receives support from the Nebraska Research

Initiative, and using resources provided by the Open Science Grid, which is

supported by the National Science Foundation and the U.S. Department of

Energy's Office of Science. This research used the Pegasus Workflow

Management Software funded by the National Science Foundation under grant

#1664162. This publication made use of the PubMLST website

(https://pubmlst.org/) developed by Keith Jolley (Jolley & Maiden 2010, BMC

Bioinformatics, 11:595) and sited at the University of Oxford. The development

of that website was funded by the Wellcome Trust. We would like to greatly

thank Mats Rynge for his extensive assistance and valuable suggestions while

setting up and running ProkEvo on the Open Science Grid. We also thank Dr.

Derek Weitzel and Karan Vahi for their technical support.

1144

References

[1]   Bedford J, Farrar J, Ihekweazu C, Kang G, Koopmans M, Nkengasong J.

A new twenty-first century science for effective epidemic response.

Nature. 2019 Nov;575(7781):130-6.

[2]   Lewnard JA, Reingold AL. Emerging challenges and opportunities in

infectious disease epidemiology. American journal of epidemiology. 2019

May 1;188(5):873-82.

[3]   Armstrong GL, MacCannell DR, Taylor J, Carleton HA, Neuhaus EB,

Bradbury RS, Posey JE, Gwinn M. Pathogen genomics in public health.

New England Journal of Medicine. 2019 Dec 26;381(26):2569-80.

[4]   Achtman M. How old are bacterial pathogens?. Proceedings of the Royal

Society B: Biological Sciences. 2016 Aug 17;283(1836):20160990.

[5]   Selander RK, Musser JM, Caugant DA, Gilmour MN, Whittam TS.

Population genetics of pathogenic bacteria. Microbial pathogenesis. 1987

Jul 1;3(1):1-7.

1160 [6] Shapiro BJ. How clonal are bacteria over time?. Current opinion in

1161   microbiology. 2016 Jun 1;31:116-23.

1162 [7] Smith JM, Smith NH, O'Rourke M, Spratt BG. How clonal are bacteria?.

1163   Proceedings of the National Academy of Sciences. 1993 May

1164   15;90(10):4384-8.

1165 [8] Maiden MC, Bygraves JA, Feil E, Morelli G, Russell JE, Urwin R, Zhang

1166   Q, Zhou J, Zurth K, Caugant DA, Feavers IM. Multilocus sequence

1167   typing: a portable approach to the identification of clones within

1168   populations of pathogenic microorganisms. Proceedings of the National

1169   Academy of Sciences. 1998 Mar 17;95(6):3140-5.

1170 [9] Maiden MC. Multilocus sequence typing of bacteria. Annu. Rev.

1171   Microbiol.. 2006 Oct 13;60:561-88.

1172 [10] Feil EJ, Li BC, Aanensen DM, Hanage WP, Spratt BG. eBURST:

1173   inferring patterns of evolutionary descent among clusters of related

1174   bacterial genotypes from multilocus sequence typing data. Journal of

1175   bacteriology. 2004 Mar 1;186(5):1518-30.

1176 [11] Croucher NJ, Harris SR, Grad YH, Hanage WP. Bacterial genomes in

1177   epidemiology—present and future. Philosophical Transactions of the

1178   Royal Society B: Biological Sciences. 2013 Mar 19;368(1614):20120202.

1179 [12] Alikhan NF, Zhou Z, Sergeant MJ, Achtman M. A genomic overview of

1180   the population structure of Salmonella. PLoS genetics. 2018 Apr

1181   5;14(4):e1007261.

[13]   Achtman M, Wain J, Weill FX, Nair S, Zhou Z, Sangal V, Krauland MG,

Hale JL, Harbottle H, Uesbeck A, Dougan G. Multilocus sequence typing

as a replacement for serotyping in Salmonella enterica. PLoS Pathog. 2012

Jun 21;8(6):e1002776.

[14]   Břinda K, Callendrello A, Ma KC, MacFadden DR, Charalampous T, Lee

RS, Cowley L, Wadsworth CB, Grad YH, Kucherov G, O'Grady J. Rapid

inference of antibiotic resistance and susceptibility by genomic neighbour

typing. Nature microbiology. 2020 Mar;5(3):455-64.

[15]   MacFadden DR, Coburn B, Břinda K, Corbeil A, Daneman N, Fisman D,

Lee RS, Lipsitch M, McGeer A, Melano RG, Mubareka S. Using genetic

distance from archived samples for the prediction of antibiotic resistance

in Escherichia coli. Antimicrobial agents and chemotherapy. 2020 Mar

9;64(5):e02417-19.

[16]   Page AJ, Alikhan NF, Carleton HA, Seemann T, Keane JA, Katz LS.

Comparison of classical multi-locus sequence typing software for next-

generation sequencing data. Microbial genomics. 2017 Aug;3(8).

[17]   Gupta A, Jordan IK, Rishishwar L. stringMLST: a fast kmer based tool for

multilocus sequence typing. Bioinformatics. 2017 Jan 1;33(1):119-21.

[18]   Abebe E, Gugsa G, Ahmed M. Review on major food-borne zoonotic

bacterial pathogens. Journal of tropical medicine. 2020 Jun 29;2020.

[19]   Centers for Disease Control and Prevention. Foodborne germs and

illnesses. Centers for Disease Control and Prevention. https://www. cdc.

gov/foodsafety/foodborne-germs. html. 2016.

[20]   Centers for Disease Control and Prevention. Individual Salmonella

serotypes reports. Centers for Disease Control and Prevention.

https://www.cdc.gov/salmonella/reportspubs/salmonella-atlas/serotype-

reports.html. 2020.

[21]   Pavlovikj N, Gomes-Neto JC, Deogun JS, Benson AK. ProkEvo: an

automated, reproducible, and scalable framework for high-throughput

bacterial population genomics analyses. PeerJ. 2021 May 21;9:e11376.

[22]   Seemann T, mlst Github https://github.com/tseemann/mlst.

[23]   HCC. 2008. Holland computing center | Nebraska.

[24]   Pordes R, Petravick D, Kramer B, Olson D, Livny M, Roy A, Avery P,

Blackburn K, Wenaus T, Würthwein F+6 more. 2007. The open science

grid. Journal of Physics: Conference Series 78:12057

[25]   Sfiligoi I, Bradley DC, Holzman B, Mhashilkar P, Padhi S, Wurthwein F.

2009. The pilot way to grid resources using glideinWMS.

[26]   Anaconda. 2012. Anaconda | The World's Most Popular Data Science

Platform.

[27]   Zhou Z, Alikhan NF, Mohamed K, Achtman M, Agama Study Group. The

user's guide to comparative genomics with EnteroBase. Three case

studies: micro-clades within Salmonella enterica serovar Agama, ancient

and modern populations of Yersinia pestis, and core genomic diversity of

all Escherichia. Biorxiv. 2019 Jan 1:613554.

[28]   Feil EJ, Cooper JE, Grundmann H, Robinson DA, Enright MC, Berendt T,

Peacock SJ, Smith JM, Murphy M, Spratt BG, Moore CE. How clonal is

Staphylococcus aureus?. Journal of bacteriology. 2003 Jun

1;185(11):3307-16.

[29] Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB,

Simpson GL, Solymos P, Stevens MH, Wagner H, Oksanen MJ. Package

'vegan'. Community ecology package, version. 2013 Dec 12;2(9):1-295.

[30] Jolley KA, Bray JE, Maiden MC. Open-access bacterial population

genomics: BIGSdb software, the PubMLST. org website and their

applications. Wellcome open research. 2018;3.

[31] Deelman E, Singh G, Su MH, Blythe J, Gil Y, Kesselman C, Mehta G,

Vahi K, Berriman GB, Good J, Laity A. Pegasus: A framework for

mapping complex scientific workflows onto distributed systems. Scientific

Programming. 2005 Jan 1;13(3):219-37.

[32] Burnett E, Ishida M, De Janon S, Naushad S, Duceppe MO, Gao R,

Jardim A, Chen JC, Tagg KA, Ogunremi D, Vinueza-Burgos C. Whole-

genome sequencing reveals the presence of the blactx-m-65 gene in

extended-spectrum β-lactamase-producing and multi-drug-resistant clones

of salmonella serovar infantis isolated from broiler chicken environments

in the Galapagos Islands. Antibiotics. 2021 Mar;10(3):267.

[33] Cooper AL, Low AJ, Koziol AG, Thomas MC, Leclair D, Tamber S,

Wong A, Blais BW, Carrillo CD. Systematic evaluation of whole genome

sequence-based predictions of Salmonella serotype and antimicrobial

resistance. Frontiers in microbiology. 2020 Apr 3;11:549.

[34]   Dallman TJ, Byrne L, Ashton PM, Cowley LA, Perry NT, Adak G,

1251   Petrovska L, Ellis RJ, Elson R, Underwood A, Green J. Whole-genome

1252   sequencing for national surveillance of Shiga toxin–producing Escherichia

1253   coli O157. Clinical Infectious Diseases. 2015 Aug 1;61(3):305-12.

1254   [35]   Pightling AW, Pettengill JB, Luo Y, Baugher JD, Rand H, Strain E.

1255   Interpreting whole-genome sequence analyses of foodborne bacteria for

1256   regulatory applications and outbreak investigations. Frontiers in

1257   microbiology. 2018 Jul 10;9:1482.

1258   [36]   Sheppard SK, Jolley KA, Maiden MC. A gene-by-gene approach to

1259   bacterial population genomics: whole genome MLST of Campylobacter.

1260   Genes. 2012 Jun;3(2):261-77.

1261   [37]   Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ,

1262   Bentley SD, Maiden MC, Parkhill J, Falush D. Genome-wide association

1263   study identifies vitamin B5 biosynthesis as a host specificity factor in

1264   Campylobacter. Proceedings of the national academy of sciences. 2013 Jul

1265   16;110(29):11923-7.

1266   [38]   Alba P, Leekitcharoenphon P, Carfora V, Amoruso R, Cordaro G, Di

1267   Matteo P, Ianzano A, Iurescia M, Diaconu EL, Pedersen SK, Guerra B.

1268   Molecular epidemiology of Salmonella Infantis in Europe: insights into

1269   the success of the bacterial host and its parasitic pESI-like megaplasmid.

1270   Microbial genomics. 2020 May;6(5).

[39]   Cheng L, Connor TR, Sirén J, Aanensen DM, Corander J. Hierarchical

and spatially explicit clustering of DNA sequences with BAPS software.

Molecular biology and evolution. 2013 Feb 13;30(5):1224-8.

[40]   Tonkin-Hill G, Lees JA, Bentley SD, Frost SD, Corander J. Fast

hierarchical Bayesian analysis of population structure. Nucleic Acids

Research. 2019 Jun 20;47(11):5539-49.

[41]   Didelot X, Bowden R, Street T, Golubchik T, Spencer C, McVean G,

Sangal V, Anjum MF, Achtman M, Falush D, Donnelly P. Recombination

and population structure in Salmonella enterica. PLoS genetics. 2011 Jul

28;7(7):e1002191.

[42]   Gymoese P, Kiil K, Torpdahl M, Østerlund MT, Sørensen G, Olsen JE,

Nielsen EM, Litrup E. WGS based study of the population structure of

Salmonella enterica serovar Infantis. BMC genomics. 2019 Dec;20(1):1-1.

[43]   Liao J, Orsi RH, Carroll LM, Wiedmann M. Comparative genomics

reveals different population structures associated with host and geographic

origin in antimicrobial-resistant Salmonella enterica. Environmental

microbiology. 2020 Jul;22(7):2811-28.

[44]   Maiden MC, Van Rensburg MJ, Bray JE, Earle SG, Ford SA, Jolley KA,

McCarthy ND. MLST revisited: the gene-by-gene approach to bacterial

genomics. Nature Reviews Microbiology. 2013 Oct;11(10):728-36.

[45]   Spratt BG, Hanage WP, Li B, Aanensen DM, Feil EJ. Displaying the

relatedness among isolates of bacterial species–the eBURST approach.

FEMS microbiology letters. 2004 Dec 1;241(2):129-34.

[46] Bobay LM, Ochman H. Impact of recombination on the base composition

of bacteria and archaea. Molecular biology and evolution. 2017 Oct

1;34(10):2627-36.

[47] Raghavan R, Kelkar YD, Ochman H. A selective force favoring increased

G+ C content in bacterial genes. Proceedings of the National Academy of

Sciences. 2012 Sep 4;109(36):14504-7.

[48] De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J,

Wick R, AbuOun M, Stubberfield E, Hoosdally SJ, Crook DW.

Comparison of long-read sequencing technologies in the hybrid assembly

of complex bacterial genomes. Microbial genomics. 2019 Sep;5(9).

[49] Page AJ, Keane JA. Rapid multi-locus sequence typing direct from

uncorrected long reads using Krocus. PeerJ. 2018 Jul 31;6:e5233.

[50] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for

Illumina sequence data. Bioinformatics. 2014 Aug 1;30(15):2114-20.

[51] Andrews S. FASTQC: a quality control tool for high throughput sequence

data. Available at

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. 2010.

[52] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,

Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV. SPAdes: a

new genome assembly algorithm and its applications to single-cell

sequencing. Journal of computational biology. 2012 May 1;19(5):455-77.

[53] Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment

tool for genome assemblies. Bioinformatics. 2013 Apr 15;29(8):1072-5.

[54]   Treangen TJ, Sommer DD, Angly FE, Koren S, Pop M. Next generation

sequence assembly with AMOS. Current Protocols in Bioinformatics.

2011 Mar;33(1):11-8.

[55]   Rice P, Longden I, Bleasby A. EMBOSS: the European molecular biology

open software suite. Trends in genetics. 2000 Jun 1;16(6):276-7.

[56]   Rizk G, Lavenier D, Chikhi R. DSK: k-mer counting with very low

memory usage. Bioinformatics. 2013 Mar 1;29(5):652-3.

[57]   Gomes-Neto JC, Pavlovikj N, Cano C, Abdalhamid B, Al-Ghalith GA,

Loy JD, Knights D, Iwen PC, Chaves BD, Benson AK. Heuristic and

hierarchical-based population mining of Salmonella enterica lineage I pan-

genomes as a platform to enhance food safety. Front. Sustain. Food Syst.

5:725791. doi: 10.3389/fsufs.2021.725791.

[58]   Valieris R. Parallel-fastq-dump. GitHub. Available at

https://github.com/rvalieris/parallelfastq-dump. 2020.

[59]   Rang FJ, Kloosterman WP, de Ridder J. From squiggle to basepair:

computational approaches for improving nanopore sequencing read

accuracy. Genome biology. 2018 Dec;19(1):1-1.

[60]   Rhoads A, Au KF. PacBio sequencing and its applications. Genomics,

proteomics & bioinformatics. 2015 Oct 1;13(5):278-89.

[61]   Chen Z, Erickson DL, Meng J. Benchmarking hybrid assembly

approaches for genomic analyses of bacterial pathogens using Illumina

and Oxford Nanopore sequencing. BMC genomics. 2020 Dec;21(1):1-21.

[62] Liou CH, Wu HC, Liao YC, Lauderdale TL, Huang IW, Chen FJ.

nanoMLST: accurate multilocus sequence typing using Oxford Nanopore

Technologies MinION with a dual-barcode approach to multiplex large

numbers of samples. Microbial genomics. 2020 Mar;6(3).

[63] Feijao P, Yao HT, Fornika D, Gardy J, Hsiao W, Chauve C, Chindelevitch

L. MentaLiST–A fast MLST caller for large MLST schemes. Microbial

genomics. 2018 Feb;4(2).

[64] Espitia-Navarro HF, Chande AT, Nagar SD, Smith H, Jordan IK,

Rishishwar L. STing: accurate and ultrafast genomic profiling with exact

sequence matches. Nucleic acids research. 2020 Aug 20;48(14):7681-9.

[65] Hunt M, Mather AE, Sánchez-Busó L, Page AJ, Parkhill J, Keane JA,

Harris SR. ARIBA: rapid antimicrobial resistance genotyping directly

from sequencing reads. Microbial genomics. 2017 Oct;3(10).

[66] Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT,

Fookes M, Falush D, Keane JA, Parkhill J. Roary: rapid large-scale

prokaryote pan genome analysis. Bioinformatics. 2015 Nov

15;31(22):3691-3.

[67] Zhou Z, Alikhan NF, Mohamed K, Fan Y, Achtman M, Brown D,

Chattaway M, Dallman T, Delahay R, Kornschober C, Pietzka A. The

EnteroBase user's guide, with case studies on Salmonella transmissions,

Yersinia pestis phylogeny, and Escherichia core genomic diversity.

Genome research. 2020 Jan 1;30(1):138-52.

[68]   Sheppard SK, Colles FM, McCARTHY ND, Strachan NJ, Ogden ID,

1362   Forbes KJ, Dallas JF, Maiden MC. Niche segregation and genetic

1363   structure of Campylobacter jejuni populations from wild and agricultural

1364   host species. Molecular ecology. 2011 Aug;20(16):3484-90.

1365   [69]   Sheppard SK, Cheng L, Méric G, De Haan CP, Llarena AK, Marttinen P,

1366   Vidal A, Ridley A, Clifton-Hadley F, Connor TR, Strachan NJ. Cryptic

1367   ecology among host generalist Campylobacter jejuni in domestic animals.

1368   Molecular ecology. 2014 May;23(10):2442-51.

1369   [70]   Yahara K, Didelot X, Ansari MA, Sheppard SK, Falush D. Efficient

1370   inference of recombination hot regions in bacterial genomes. Molecular

1371   biology and evolution. 2014 Jun 1;31(6):1593-605.

1372   [71]   Mottawea W, Duceppe MO, Dupras AA, Usongo V, Jeukens J, Freschi L,

1373   Emond-Rheault JG, Hamel J, Kukavica-Ibrulj I, Boyle B, Gill A.

1374   Salmonella enterica prophage sequence profiles reflect genome diversity

1375   and can be used for high discrimination subtyping. Frontiers in

1376   microbiology. 2018 May 4;9:836.

1377   [72]   den Bakker HC, Desjardins CA, Griggs AD, Peters JE, Zeng Q, Young

1378   SK, Kodira CD, Yandava C, Hepburn TA, Haas BJ, Birren BW.

1379   Evolutionary dynamics of the accessory genome of Listeria

1380   monocytogenes. PLoS One. 2013 Jun 25;8(6):e67511.

1381   [73]   Castillo-Ramírez S, Corander J, Marttinen P, Aldeljawi M, Hanage WP,

1382   Westh H, Boye K, Gulay Z, Bentley SD, Parkhill J, Holden MT.

1383   Phylogeographic variation in recombination rates within a global clone of

methicillin-resistant Staphylococcus aureus. Genome biology. 2012

Dec;13(12):1-3.

[74]   Laing CR, Whiteside MD, Gannon VP. Pan-genome analyses of the

species Salmonella enterica, and identification of genomic markers

predictive for species, subspecies, and serovar. Frontiers in microbiology.

2017 Jul 31;8:1345.

[75]   Ferrari RG, Rosario DK, Cunha-Neto A, Mano SB, Figueiredo EE, Conte-

Junior CA. Worldwide epidemiology of Salmonella serovars in animal-

based foods: a meta-analysis. Applied and environmental microbiology.

2019 Jul 1;85(14):e00591-19.

[76]   Price MN, Dehal PS, Arkin AP. FastTree: computing large minimum

evolution trees with profiles instead of a distance matrix. Molecular

biology and evolution. 2009 Jul 1;26(7):1641-50.

[77]   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. Nucleic acids research. 2021 Jul

2;49(W1):W293-6.

Figure 1. Computational workflow describing the analytical steps for a comparative analysis of two algorithms used for ST-based classification. From top-down, the first step (narrow-scope) of the analytical approach entailed the acquisition and processing of Illumina paired-end raw reads from four distinct pathogens (*C. jejuni*, *L. monocytogenes*, *S. enterica* and *S. aureus*), through an assembly-dependent (mlst) or assembly-free (stringMLST) approach for ST-based classification. Next, a set of comparative analyses encompassing measuring the computational performance, statistical metrics, and modeling were used to assess the accuracy and efficiency of mlst vs. stringMLST. Additionally, the contribution of genome-intrinsic and –extrinsic variables were used to identify explanatory factors that could impact the algorithmic efficiency across phylogenetic divergent species. Upon identification of inter-species differences in the performance of stringMLST, a wide-scope analysis was done to assess its accuracy across an array of other fourteen phylogenetic divergent pathogenic species of bacteria with Public Health relevance. Ultimately, stringMLST was added to the computational platform ProkEvo to facilitate ST-based classification at scale, as part of a hierarchical-based approach for population genomic analyses.

Figure 2. Box-and-whiskers plot showing the comparison of the average runtime per genome per batch (in minutes) needed by mlst and stringMLST for ST classification of genomes across four distinct bacterial species.

In order to compare the average runtime used by mlst and stringMLST with different kmer values, we chose four different datasets, including four phylogenetic divergent bacterial pathogenic species: *C. jejuni*, *L. monocytogenes*, one major serovar of *S. enterica* (*S.* Typhimurium) and *S. aureus* - using 600 randomly selected genomes for each species. These 600 genomes were randomly split into three batches with 200 genomes each. We then ran mlst with all required steps, such as quality trimming and adapter clipping, *de novo* assembly and assembly discarding, on each batch and dataset. Separately, we ran stringMLST with a range of 10 different kmer values (10, 20, 30, 35, 45, 55, 65, 70, 80, 90) on each dataset, including the default length of 35 (y-axis). For each organism, the runtime was calculated as an average of 200 genomes per batch - since there were three batches, three datapoints were used to depict the distribution of runtime in minutes (x-axis).

Figure 3. Genome-intrinsic and –extrinsic variables that can impact the accuracy of ST-based classification using either mlst (MLST-based genotyping) or stringMLST algorithmic approaches.

Box-and-whiskers plot showing genome-intrinsic variables, varying in distribution according to the bacterial species (A-C as y-axis), that may affect ST-based classification, include: (A) Number of contigs per genome (x-axis); (B) Total number of nucleotides per genome (x-axis); (C) GC% content per genome (x-axis); and (D) Dinucleotide composition of genomes. (D) Inter-species PCA using the relative frequency of all pairs of dinucleotides (16 pairs) present in the genome as input data. Only two PCs are shown, and the percentage of variance explained by either PC is depicted in parenthesis. Bar plots showing genome-extrinsic variables that may influence the performance of mlst vs. stringMLST across species include but are not limited to: (E) Total count of unique STs per database (ST richness in the database used for mapping of raw reads or assemblies) (x-axis); and (F) Total count of unique alleles across all seven loci used for ST classification (x-axis). Specifically, the differences in ST richness and allelic composition in the databases reflect difference between mlst vs. stringMLST, and were not impacted by the kmer length (E-F, y-axis).

Figure 4. Statistical analysis of ST-based classification outcomes for comparison between mlst and stringMLST performance across bacterial species.
(A-C) Box-and-whiskers plots A-C demonstrate the relationship between ST richness (x-axis), Simpson's index of diversity $(1 - D)$ based on ST composition (x-axis), or the proportion of non-classified STs (x-axis) across bacterial species (color-coded differently) and programs (y-axis), respectively. Along with plots A-C are depicted all PERMANOVA results including p-values ($p < 0.05$) and the univariate or synergistic contribution of factors measured by R-squared. PERMANOVA modeling was done in two specific ways: 1) A model including species, program, and their interaction, considering that those were the main variables of interest; and 2) All other results were calculated using univariate models and included modeling using genome-intrinsic (number of contigs per genome, total number of nucleotides per genome, and average GC% content) and – extrinsic (Simpson's D index of diversity, ST and allelic counts per database) variables. (D) Bar plot depicting the distribution of the standard deviation (SD, y-axis) for the proportion of non-classified STs based on species (color-coded differently) and programs (y-axis).

Figure 5. Multi-dimensional analysis of ST-based classification outcomes across different species using mlst vs. stringMLST.

(A) Tri-dimensional scatter plot demonstrating species grouping based on the outcomes calculated using the ST classification across programs, including: 1) Simpson's index of diversity ($1 - D$, Simpson's index); 2) ST richness; and 3) proportion of non-classified STs. (B-D) Biplots demonstrating groupings formed across species and programs based on the same outcomes. Scatter plot B depicts groupings produced based on the relationship between Simpson's index of diversity vs. ST richness (Richness); whereas scatter plot C shows the relationship between the Simpson's index of diversity and the proportion of non-classified STs; and lastly, scatter plot D depicts the relationship between ST richness (Richness) and the proportion of non-classified STs.

Figure 6. Box-and-whiskers plot depicting the concordance between mlst and stringMLST in ST calls.

Four different datasets belonging to four phylogenetic distinct bacterial pathogens, including *C. jejuni* (600 genomes), *L. monocytogenes* (600 genomes), *S. enterica* (11,787 genomes from 20 different serovars) and *S. aureus* (600 genomes) were run with mlst and stringMLST for ST-based classification. In the case of stringMLST, kmer lengths varied from 10 to 90 to identify the optimal value (highest percentage of agreement with the standard MLST approach), across all four species (y-axis). If both programs outputted identical ST calls (either number of missing/blank value), the call was defined as a match; otherwise, it was identified as a mismatch, and the percentage of agreement (x-axis, concordance) was calculated accordingly. The dashed line on the x-axis represents the percentage agreement for the kmer value of 35 which is used as a default parameter by stringMLST.

Figure 7. Phylogeny-guided display of optimal kmer length and algorithmic performance when using stringMLST for ST mapping across bacterial species. (A) Phylogeny-based display of stringMLST results across the twenty-three zoonotic serovars of *Salmonella enterica* subsp. *enterica* lineage I (*S. enterica*). The branches are colored based on the optimal kmer length which gives the lowest percentage of ST miscalls (ST calls that returned missing/blank values for stringMLST). The outer ring present in the phylogeny is colored based on the corresponding ST miscall percentages associated with each optimal kmer length. The dataset used to identify the optimal kmer length and percentage of ST miscalls was composed of 2,300 genomes (100 genomes per serovar) and the phylogenetic tree was generated using twenty-three genomes (one of each serovar to facilitate data visualization); (B) Phylogeny-based display of stringMLST results across fourteen phylogenetic divergent bacterial pathogens, including twenty-three representative genomes across each zoonotic serovar of the *S. enterica* species. The tree branches are colored based on the optimal kmer length which minimizes the percentage of ST miscalls (ST calls that returned missing/blank values for stringMLST). The outer ring present in the phylogeny corresponds to ST miscall percentage associated with each optimal kmer length. The dataset used to identify the optimal kmer length and percentage of ST miscalls was composed of 14,000 genomes (1,000 genomes for each bacterial pathogen) and 2,300 *Salmonella* genomes (100 genomes per serovar). The phylogeny was ultimately generated using 37 genomes (one of each dataset used to facilitate visualization). All phylogeny-based visualization were generated using iTOL version 6.4.

Figure 8. Comparison between the computational and statistical performance of mlst and stringMLST when using ProkEvo to run both programs.

Two subsets, one with 100 and the second one with 1,000 randomly chosen genomes, were selected from *C. jejuni*, *L. monocytogenes*, one major serovar of *S. enterica* (*S.* Typhimurium) and *S. aureus* to compare the performance of running mlst or stringMLST through ProkEvo. The performance and statistical metrics used for comparison were: (A) Total runtime of individual workflow in minutes; (B) Percentage of non-classified STs (ST calls that returned missing/blank values); and (C) Percentage of agreement (concordance) between programs ("good" or "bad" ST calls that matched between mlst and stringMLST).