The non-specific Lipid Transfer Protein (nsLTP) is involved at early and late

stages of symbiosis between *Alnus glutinosa* and *Frankia alni*.

Mélanie GASSER[1], Nicole ALLOISIO[1], Pascale FOURNIER[1], Severine BALMAND[2], Ons

KHARRAT[1], Joris TULUMELLO[1,3], Abdelaziz HEDDI[2], Pedro Da SILVA[2], Philippe

NORMAND[1], Hasna BOUBAKRI[1#], Petar PUJIC[1]

1-Université de Lyon, F-69361, Lyon, France ; Université Claude Bernard Lyon 1, CNRS,

UMR 5557, INRAE UMR1418, Ecologie Microbienne, F-69622, Villeurbanne, France

2-INSA-Lyon, INRAE, UMR203 BF2i, Biologie Fonctionnelle Insectes et Interactions,

Villeurbanne, France

3-present address: Aix Marseille Univ, CEA, CNRS, BIAM, Lab Microbial Ecology of the

Rhizosphere (LEMiRE), F-13108, Saint-Paul-Lez-Durance, France; BioIntrant 139, Rue

Philippe de Girard, Pertuis F-84120, France.

#corresponding author: hasna.boubakri@univ-lyon1.fr, tel: 33-472 44 82 00

Summary

Alnus glutinosa response to *Frankia alni* is driven by several sequential physiological

modifications that include calcium spiking, root hair deformation, penetration, induction

of primordium, formation and growth of nodule. Here, we have conducted a

transcriptomic study to analyse plant responses to *Frankia alni* at early stages of

symbiosis establishment.

Forty-two genes were significantly activated by either with a *Frankia* culture

supernatant or with living cells separated from the roots by a dialysis membrane

permitted to identify plant genes which expression changes upon early contact with

Frankia. Most of these genes encode biological processes, including oxidative stress and

response to stimuli. The most upregulated gene is the non-specific lipid transfer protein

(nsLTP) encoding gene with a fold change of 141. Physiological experiments showed

that nsLTP increases *Frankia* nitrogen fixation at sub-lethal concentration.

Immunohistochemistry experiments conducted at an early infection stage indicated that

nsLTP protein is localized at the deformed root hair region after *Frankia* inoculation and

later in nodules, precisely around bacterial vesicles. Taken together, these results

suggest that nsLTP acts at early and late stages of symbiosis, probably by increasing

nitrogen uptake by *Frankia*.

Keywords: nitrogen fixation, antimicrobial peptide, immunity, root hairs, oxidative

stress, lipid, infection threads, nodule

Introduction

Symbiosis between *Alnus glutinosa* and the actinobacterium *Frankia alni* permits the

establishment of root nodules in which nitrogen fixation takes place. This helps alder

and other plants collectively called "actinorhizal" to grow on nitrogen-poor soils and

initiate ecological successions. The determinants of this interaction are poorly known,

due among other reasons to the lack of a transformation system in alder. *Alnus glutinosa*

is the type species of the genus (Navarro *et al.*, 2003), it grows in pioneer biotopes such

as glacial moraines, volcanic ashes and mine spoils (Normand & Fernandez, 2019) and

its genome has been deciphered recently (Griesmann *et al.*, 2018).

The presence of a symbiotic bacterium like *Frankia* or simply that of its exudates,

triggers in the host plant tissues a series a events leading to root hair deformation and

calcium spiking (Granqvist *et al.*, 2015). Later, penetration, primordium formation and

growth of nodules follow with subsequent exchange of nutrients. Omic analyses have

permitted to gain a global view of the physiological changes taking place upon specific

challenges. In particular, transcriptomics of 21 day post-inoculation (dpi) nodules

permitted to see on the plant side the presence of homologs of the whole common

symbiotic signalling cascade or CSSP (Hocher, V. *et al.*, 2011; Hocher, Valérie *et al.*,

2011), some of which have been lost in non-symbiotic neighbours (Griesmann *et al.*,

2018).

This important conservation on the plant side of the common symbiotic pathway

(Hocher, V. *et al.*, 2011) contrasts with the absence in most *Frankia* genomes (Normand

et al., 2007) of the canonical *nod* genes which are involved in the biosynthesis of Nod

Factor (NF) in majority of Rhizobia (D'Haeze & Holsters, 2002). In mature nodules, the

bacterial genes *nif, hup, shc* and *suf* are upregulated but no trace of a symbiotic island

was found (Alloisio *et al.*, 2010). A study of the *Frankia alni* early response (2.5 dpi)

showed that several determinants were upregulated among which a K-transporter, lipid

modifying enzymes and a conserved cellulose synthase cluster (Pujic *et al.*, 2019).

Proteomics was performed on a related species, *Frankia coriariae* that unravelled an

upregulation of cell wall remodelling enzymes, signal transduction and host signal

processing proteins (Ktari *et al.*, 2017). Metabolomics on an *Alnus*-infective strain

showed the presence of various compounds absent from roots (Hay *et al.*, 2017), high

levels of TCA intermediates citrate, fumarate and malate (Carro *et al.*, 2015a) and high

levels of citrulline, glutamate and pyruvate (Brooks & Benson, 2016; Hay *et al.*, 2020).

Signalling between alder and *Frankia alni* involves on the part of the bacterium the

synthesis of an uncharacterized root hair deforming factor (Ghelue *et al.*, 1997;

Cérémonie *et al.*, 1999) and that of the auxin PAA (Hammad *et al.*, 2003). On the plant

side, less is known besides the upregulation of defensins that modify the porosity of

Frankia membranes (Carro *et al.*, 2015b; Carro *et al.*, 2016).

We undertook the present study to better understand the initial steps of the actinorhizal

symbiosis on the plant side where the symbiont must be recognized to pave the way for

its internalization. To analyse this early plant responses to *Frankia* symbiont we have

conducted a cell-free contact through two conditions: indirect contact with *Frankia*

trapped in dialysis tube or its supernatant and by targeting 2.5 dpi response, by which

time extensive root hair deformation has occurred (Berry & Torrey, 1983) and before

primordium formation at 7dpi (Lalonde, 1979). We further focussed on the most

upregulated gene coding a non-specific Lipid Transfer Protein (nsLTP), which is also one

of the most overexpressed genes in the nodule. We showed that nsLTP increases *Frankia*

nitrogen fixation at sub-lethal concentration. Taken into account that this protein shown

to localize at the deformed root hairs and in *Frankia's* vesicles inside nodules, we

hypothesized that nsLTP acts at early and late stages of symbiosis.

Materials and methods

Strains and growth condition before total RNA extraction

Plant samples at early stages of infection were obtained as described previously(Pujic *et*

al., 2019). Briefly, *Frankia alni* ACN14a (Normand & Lalonde, 1982) was grown in BAP-

PCM media until log-phase, collected by centrifugation, washed twice with sterile ultra-

pure water and suspended in Farhaeus medium without KNO_3. *Frankia* cells were

homogenised by forced passage through a series of needles (21G, 23G, 25G, 27G) before

inoculation.

Alnus glutinosa seeds obtained from a single individual growing on the banks of the river

Rhone in Lyon were surface sterilised and grown as described earlier (Pujic *et al.*, 2019).

Seedlings were transferred to Fåhraeus' solution (Fahraeus, 1957) in opaque plastic

pots (8 seedlings/ pot) and grown for four weeks with 0.5 g.L^{-1} KNO$_3$, followed by one

week without KNO$_3$ before inoculation.

Two independent experiments were performed. In experiment #1 (Exp 1- *Frankia*

indirect contact (*F*IC)), 8ml of *Frankia* cells were transferred to dialysis tubing

(MWCO=100kDa) and arranged into a plastic pot containing 8 seedlings and filled with

Fåhraeus' solution without KNO$_3$. Five biological replicates were performed with 4

plastic pots per replicate. In experiment #2 (Exp2-*Frankia* supernatant direct contact

(SupC)), *Frankia* supernatants were applied directly on *Alnus* roots. Supernatant extract

was prepared as follow: 250 ml log-phase culture cells were removed by centrifugation

and the supernatant filtered through 0.22 μm membranes (Millipore, Billerica, MA).

Solid phase extractions of the supernatants were done using benzene sulphonic acid

cation exchanger on silica (Macherey Nagel, Hoerdt, France). Elution was done using

50:50 v/v methanol-water followed by 100 % methanol. Two fractions were lyophilised

under vacuum, dissolved in water and tested for root hair deformation activity on

independent plant roots. In parallel, as a control condition without supernatant, an

equivalent volume of culture medium was treated with the same extraction protocol.

Three biological replicates were performed with 6 seedlings per replicate.

The root hair deformation process took place similarly in both direct and indirect

contact conditions, confirmed with stereomicroscope observations (Leica MZ8, Wetzlar,

Germany). After 2.5 days (64 hours) with root hairs highly deformed, a 2 cm long

segment representing the central part of the root was cut at 2 cm from the distal end,

washed with sterile water before freezing in liquid nitrogen and storing at -80° C. In

addition, a T0 control condition without *Frankia* was performed and plant roots were

washed and frozen as described above.

Transcriptomics of alder at 2.5dpi

Transcriptomics was done as described earlier (Hocher, V. *et al.*, 2011). Total RNA was purified from roots using RNeasy plant mini kits (Qiagen, Courtaboeuf, France) and treated with DNases as before. Residual DNA was removed using the Turbo DNA free kit (Ambion, Thermo Fisher Scientific, Wilmington, DE), quantified using a NanoDrop spectrophotometer (Thermo Fischer Scientific) and qualitatively assessed using a Bioanalyzer 2100 (Agilent, Waldbronn, Germany).

Microarrays were designed (13909 probes; 1 probe/*A. glutinosa* unigene), manufactured and hybridised by Imaxio (http://www.imaxio.com/index.php), using Agilent Technologies (http://www.home.agilent.com/agilent/home.jspx) as previously described (Hocher, V. *et al.*, 2011). The microarrays were scanned with an Agilent G2505C Scanner. The Feature Extraction software (Agilent, version 11.5.1.1) was used to quantify the intensity of fluorescence signals and microarray raw data were analysed using GeneSpring GX 12.0 software (Agilent technologies). Normalisation per chip (to the 75th percentile) and per probe (to the median) were performed to allow comparison of samples. The two experiments Exp 1 and Exp 2 were analysed separately. In order to limit false positive results, microarray data were filtered according to the flag parameter "detected". Thus, probes taken into account are uniform, non-outlying, non-saturated and displayed an intensity level above the background in at least one of the two biological conditions.

In Exp1, 1590 probes yielding an intensity level significantly below the background in the 10 samples were discarded. In addition, 1146 probes did not yield the flag "detected" in at least one of the two biological conditions and were also discarded. Thus, 11 173 probes were taken into account for subsequent analyses of Exp 1.

In Exp 2, 1765 probes yielded an intensity level significantly below the background in

the 6 samples and were discarded. In addition, 1113 probes did not yield the flag

"detected" in at least one of the two biological conditions and were also discarded. Thus,

11031 probes were taken into account for subsequent analyses of Exp 2. T-tests

comparing *F*IC roots *vs.* control roots in Exp 1 and SupC roots *vs.* control roots in Exp 2

were applied and only those genes with an average fold change (FC) above 2 (up-

regulated) or below 0.5 (down-regulated) with a *p*-value<0.05 were considered

significant. The normalized and raw microarray data values have been deposited in the

Gene Expression Omnibus database (www.ncbi.nlm.nih.gov/geo; accession nos. E-

MTAB-8936 and E-MTAB-8937).

Quantitative real-time RT-PCR

Reverse transcription (RT) and real time quantitative PCR (qRT-PCR) were performed

with the same biological replicates used for microarray experiments. For *A. glutinosa*

analyses, RT was performed with 5 µg of total mRNA using Transcriptor Reverse

Transcriptase and oligo (dT)$_{15}$ primer (Roche, Mannheim, Germany). QRT-PCR was run

on a LightCycler 480 (Roche) using LightCycler 480 SYBR Green I Master (Roche) under

the following conditions: 95 °C for 5 min; 45 cycles of 95 °C for 20 s, 60 °C for 20 s and

72 °C for 15 s. Primer sets were designed using ProbeFinder (Roche) and Primer 3

softwares and can be seen as Supporting Information (Table S1). Two qRT-PCR

reactions were run for each biological replicate and each primer set. Expression values

were normalised using the expression level of the *Ag-ubi* gene that encodes ubiquitin

(Hocher, V. *et al.*, 2011).

Cloning of *agLTP24* gene

The *agltp24* gene was PCR amplified from cDNA prepared above using designed

primers (Table S1) under the following conditions: 98 °C for 30 s; 30 cycles of 98 °C for 7

s, 63 °C for 20 s and 72 °C for 8 s. The reaction mixture of 50 µl PCR contained 1X

Phusion HF buffer, 200 µM dNTPs, 0.5 µM Forward primer, 0.5 µM reverse primer, 87 ng

template DNA and 1-units Phusion DNA Polymerase (NEB, Evry, France). The PCR

product was cloned using the hot fusion method (Fu *et al.*, 2014) in pET30a+ vector

(Merck, Molsheim France) pre-digested with *Eco*RI and *Bgl*II in order to fusion AgLTP24

with a N- terminal 6XHis flag. Ligation mixture was transformed in *Escherichia coli*

DH5α chemocompetent cells (Chung *et al.*, 1989) for plasmid DNA propagation and

sequencing (pET30a-HIS-AgLTP24, supplemental Fig. S1). Plasmids DNA was prepared

using (Nucleospin Plasmid DNA kit Macherey Nagel) from 4 independent clones, 2 ml of

each *E. coli* culture grown overnight in 5ml LB medium at 37°C with shaking. Cloned

insert DNA was checked by Sanger sequencing (Biofidal, Lyon, France) and used for

electrotransformation into iSHuffle T7 *lysY E. coli* cells (NEB, Evry, France), a strain used

to obtain proteins with disulphide bonds (Lobstein *et al.*, 2012).

AgLTP24 production and purification

Cultures were grown in LB medium supplemented with kanamycin (40µg.ml^{-1}) at 30°C

at 130 rpm until log phase was reached (OD$_{600}$ ~0,5) then the HIS-AgLTP24 expression

was induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG). Induced cultures

were incubated at 30°C for 3h at 130 rpm and harvested by centrifugation (5100 g for

10 min at 4°C). Expression was analysed by 4-20% Tricine SDS-PAGE (Schagger, 2006).

Pellets were lysed by freeze thaw and resuspended in a binding buffer (100 mM Tris,

500 mM NaCl, 5mM Imidazole, pH 7.4). A second lysis was performed using silica beads

and Fastprep 24™ 5G (MP Biomedicals). The cell lysate was harvested by centrifugation

at 5100 g, 4°C for 30min and used for the purification of HIS-AgLTP24 using His

GravityTrap columns (GE Healthcare) with modified binding (100mM Tris, 500mM NaCl,

5mM Imidazole, pH 7.4) and elution buffers (100 mM Tris, 500 Nacl, 500 mM Imidazole,

pH 7.4). Purification of HIS-AgLTP24 was verified by 4-20% Tricine SDS-PAGE. Elution

fractions was applied to a 3K MWCO Amicon Ultra-15 centrifugal filter (Merck Millipore)

to buffer exchange with the enterokinase buffer (20 mM Tris, 50 mM NaCl, 2 mM $CaCl_2$,

pH 6.8) and concentrate. The HIS tag was removed from the mature sequence of

AgLTP24 with enterokinase (NEB, Evry, France) for 16 h at room temperature. Cleavage

was confirmed by 4-20% Tricine SDS-PAGE. The enterokinase was removed from the

mixture using the Enterokinase Removal Kit (Sigma). His tag removal was done with a

His GravityTrap column (GE-Healthcare) using an imidazole free binding buffer (100

mM Tris, 500 mM NaCl, pH 7.4). Wash fractions containing AgLTP24 purified were

applied to a 3K MWCO Amicon Ultra-15 centrifugal filter (Merck Millipore) for buffer

exchange with NH_4^+-free FBM medium (FBM-). Final concentration was determined with

Qubit Protein Assay Kit (ThermoFisher). The purified AgLTP24 protein were further

analysed by UPLC-ESI-HRMS. The system used for UPLC-ESI-HRMS was an ultra-

performance liquid chromatography Ultimate 3000 (Thermo Fisher Scientific, Villebon-

sur-Yvette, France) coupled to a high resolution hybrid quadrupole-time of flight mass

spectrometer (Impact II, Bruker, Brême, Germany) equipped with electrospray

ionization source (ESI) (Bruker, Brême Germany). Instrument control and data

collection were performed using Data Analysis 5.0 software.

Immunolocalization of AgLTP24

Immunostaining and fluorescence microscopy was done as described earlier (Login *et*

al., 2011) with the synthesised epitope (DKHSTADFEKLAPCGKAAQD) injected to a

rabbit (Covalab). Plant immunolocalization was performed on *A. glutinosa* roots (2.5

dpi) as described above in Exp 1 (*F*indC). In addition, nodules (21dpi) obtained as a

previous study (Carro *et al.*, 2015b) were also used here to localise agLTP24 at the

mature stage of symbiosis. T0 non-infected seedlings were used as control.

Physiological assays with AgLTP24 on *Frankia alni*.

The antimicrobial activity of AgLTP24 (using 1μM, 2μM, 5μM concentrations), was done

using the resazurin assay in 24-well microplates (Greiner bio-one, Les Ulis, France). A 2-

week-old *Frankia* ACN14a culture grown in FBM medium supplemented by 5mM of

NH$_4$Cl (FBM+) was centrifuged at 5100 g for 15 minutes, the supernatant was removed,

and the pellet homogenized in FBM- by repeated passage through syringes. *Frankia*

culture was done in 1ml of FBM- per well with a final optical density of 0.02 and with an

appropriate concentration of AgLTP24 peptide. Four replicates were made for each

condition. Negative (*Frankia* ACN14 in FBM-) and positive controls (*Frankia* ACN14a in

FBM- supplemented by kanamycin at 40μg.ml^{-1} final concentration) were performed.

Microplates were incubated at 28°C, 80 rpm in a humid incubator. After 6 days of

incubation, resazurin (Fisher Scientific SAS, Illkirch, France) was added to a final

concentration of 6.25μg.ml^{-1} in each well and the microplates were incubated at 28°C, 80

rpm for 16 hours in the dark. Fluorescence measurements were performed with a

microplate reader (infinite M200PRO, TECAN) with an excitation wavelength of 530nm

and an emission wavelength of 590nm. The results obtained were normalised to the

mean fluorescence of the negative control. Since data were normally distributed, mean

comparisons with the negative control were performed with a Student's *t*-test.

A second bioassay was made by growing *F. alni* in FBM- and incubating from 1 to 14

days at 28°C (5 replicates by kinetic point and by condition) as described previously. At

each kinetic point, the effects nitrogen fixation (or ARA), respiration (IRA) and on

morphology were monitored (Prin *et al.*, 1990; Carro *et al.*, 2015b). Since data were

normally distributed or not, different statistical tests were made using GraphPad Prism

9.2.0 (GraphPad software Inc; San Diego, CA, USA).

Bioinformatics

Blast analyses and GO assignation were performed on Blast2GO v5.2 using BLASTx

against NR, Gene Ontology (GOs) and Inter-ProScan (Conesa & Gotz, 2008). Fishers'

exact test implemented in Blast2GO was used to identify significantly enriched GO

categories. A GO category was considered significantly enriched only when the *p*-value

for that category was < 0.05 after applying FDR correction. Molecular mass of proteins

and their isolectric point were calculated through the Expasy software

(https://web.expasy.org/protparam/)

Results

Transcriptomics response of alder at 2dpi

In the first early response transcriptomic experiment (Exp 1), 300 and 225 genes were

found to be significantly up- or down- regulated in *A. glutinosa* after 2.5 days post

inoculation (dpi) of *Frankia* in a separated dialysis bag (cut off 100 KDa -*Frankia*

indirect contact (*F*indC) roots/ control roots) by having a fold change (FC) ≥ 2 or ≤ 0.5 (p

≤ 0.05), respectively (Table S2). In Exp2, 248 *A. glutinosa* genes were found to be

significantly regulated 2.5 dpi with *Frankia* supernatant (*Frankia* supernatant contact

(*F*supC) roots/control roots). Of these, 176 and 76 genes were also up- and down-

regulated in Exp2; respectively (Table S3). By comparing ESTs present in both

experiments, 84 genes were found significantly up- or down-regulated (See SI Table S4).

Biological processes implicated in early steps of symbiosis

A GO-based analysis of the biological processes enriched in up- and down-regulated

genes within each experiment yielded 12 to 23 % of genes with no Blast hit (Fig. S2).

Among the sequences with a Blast hit, 8 to 14% correspond to unknown proteins and

could not be associated to any biological process. The annotated sequences retrieved

were distributed in 21 different processes (Fig. **1a**). The presence of *Frankia* at early

stages strongly activated genes involved in response to stimuli and catabolic processes

among which oxidative stress (oxidase, peroxidase) and defence (chitinase, defensin,

thaumatin…) were upregulated in both conditions, as well as a large portion of genes

involved in nitrogen compound metabolic processes with kinases, transferases, lyases or

glucosidases. Globally, the number of sequences retrieved was higher in Exp1 than Exp2

(Fig. **1a**), suggesting a stronger signal detected in indirect interaction through dialysis

tubing.

A statistical analysis permitted to identify certain enzyme families by comparing each

pool of genes. Even this Fisher test made in both experiment, only Exp1 gave differences

in GO assignation between up and down-regulated genes (Fig. **1b**).

In Exp1, an overrepresentation of genes coding enzymes involved in oxidation-reduction

biological processes and oxidative stress responses was seen with numerous

peroxidases, catalases, oxidases or lipoxygenases overexpressed (Table S1). Also,

chemical reactions involving alpha and aromatic amino acid metabolism and fatty acid

beta-oxidation, selenocysteine methyltransferase, phenylalanine ammonia lyase or

caffeic acid O-methyltransferase were seen following indirect contact with *Frankia* cells.

Nucleotide binding proteins such as a receptor kinase (AG-N01f_037_C05), a uridine

kinase (AG-N01f_030_B06) or a calcium transporting membrane (AGCL1701Contig1)

were also more abundant in the set of up-regulated proteins of Exp1. Finally, several

genes coding defence response were overrepresented such as endochitinase, disease

resistance protein, defensin, thaumatin, allergen or nsLTP. Conversely, we observed an

enrichment in down-regulated genes of one molecular function of GO assignation:

isomerase such as inositol-3-phosphate synthase, ribose-5-phosphate isomerase or

beta-amyrin synthase.

The most upregulated genes in both experimentations

By compiling differentially regulated genes found in both experiments (Table 1 and

Table S4), we compared them through GO-based analysis. An overrepresentation of only

one biological process (lipid metabolism) in down-regulated genes was found with the

same statistical restriction (FDR<0.05). Indeed, several genes, including synthase,

reductase and oxidase and involved in triterpenoid (AG-N01f_010_H21) isopentenyl

diphosphate (IPP) and dimethylallyl diphosphate (AG-N01f_014_J09); geranyl or

geranylgeranyl diphosphate (GPP and GGPP) (AG-N01f_005_E08; AG-J07f_004_F15) or

terpene (AG-J07f_004_M14; AGCL3086Contig1) biosynthesis were detected as more

abundant in down-regulated genes and already observed by our first GO assignation

(Fig. **1a**).

Amongst the 42 most upregulated genes found in both conditions (Table 1), the most

upregulated gene encodes a non-specific lipid transfer protein or nsLTP (fold changes:

140.7 (*F*indC) and 74.8 (*F*supC)), followed by a betabutilin and a pectin methylesterase

inhibitor. Remarkably, 20 out these 42 up-regulated genes encoded proteins with a

peptide signal (Table 1). Among them, numerous upregulated genes encode secreted

peptides (47%), some of them are classified as antimicrobial: three defensins (Ag5, Ag3

and Ag11), a nsLTP and a thaumatin like-protein.

To confirm the level of expression at this early step, we focussed on two genes, one

coding the nsLTP and the second coding the basic blue copper protein (fold changes:

(fold changes: 140.7 (FindC) and 74.8 (FsupC), 4.7 (FindC) and 11.7 (FsupC),

respectively) as candidate genes for qRT-PCR with two primer pairs per gene (Table S1).

Up-regulation was confirmed by qRT-PCR with the same biological replicates previously

used (Table S5).

In order to examine the temporal distribution of expression, we extracted from our

previous work (Hocher, V. *et al.*, 2011) the fold change obtained with the same

microarray but at 21 dpi when the nodule is formed and compared it to non-infected

roots (Table 1, Fig. 2).

Almost down-regulated genes at early stage were either repressed or not modulated at

nodule stage. Among them, we observed an enrichment of proteins associated to lipid

metabolic process and lyase activity. Conversely, genes upregulated at 2.5 dpi are

classified in three profiles. Sixteen are still induced when 17 are not modulated and 8

are repressed in nodule. For instance, genes encoding proteins involved in stress

response are found in the three profiles (Fig. 2) but we found mainly AMPs (defensin,

thaumatin) in the first group when chitinase and peroxidase enriched the two other

groups (Table 1). The nsLTP found as the most upregulated gene at early stage

presented also a very high expression at 21 dpi (FC=2145 in nodule, Table 1) suggesting

a requirement all along the symbiosis process. NsLTPs are classified as AMPs and

annotated as systematic acquired resistance according GO analysis suggest its potential

role in defense response. For all these reasons, we decided to focus our functional

analysis on this protein.

Structure and classification of AgLTP24

The nsLTP from *A. glutinosa* overexpressed during its interaction with *Frankia* is named AgLTP24. The total nucleotide sequence is 552 base pairs and contains one exon. This sequence encodes a putative processed protein of 114 amino acids including a signal peptide with a sequence of 22 amino acids that allows its secretion [29]. The mature peptide has a molecular weight of 9700.28 Dalton and a cationic isoelectric point of 8.46, this nsLTPs is classified according to Edstam et al. (Edstam *et al.*, 2011) as Type D. It contains the characteristic motif of nsLTPs composed of 8 cysteines: C-X13-C-X14CC-X9-C-X1-C-X24-C-X10-C which stabilises the 3D structure by folding the α-helices domains with disulphide bridges forming a hydrophobic cavity.

Immunolocalization of AgLTP24 in planta

In order to determine where this nsLTP is secreted during symbiosis establishment, we performed an immunohistolocalization within plant tissues. Antibodies anti-AgLTP24 were applied to different alder tissues inoculated or not by *Frankia*. First, we produced deformed root hairs after an indirect contact of *Frankia* (F indC) and observed signals in extracellular nooks (Fig. **3b** and Fig. **3c**). These nooks were the specific site for *Frankia* binding. No signal was detected with control rabbit serum (Fig. **3a**) or with anti-AgLTP24 against non-infected roots (Fig. S3). Microarrays and qRT-PCR showed overexpression of this protein in mature 21 dpi nodules. We observed a specific immunolocalization in plant cells infected by *Frankia* situated in the fixation zone of the nodule (Fig. **3e**), specifically on *Frankia*'s vesicles (Fig. **3f**).

Biological production of AgLTP24

The nsLTPs are low molecular weight peptides rich in cysteines. It has a signal sequence

that allows the plant to address it to a target compartment. Those nsLTPs are

characterized by a conserved motif of 8 cysteines in their mature sequence forming 4

disulfide bonds important for the formation of the hydrophobic cavity, which makes

their production difficult. Synthetic production is limited to small peptides with few

disulfide bonds, so we developed a biological production of mature AgLTP24, i.e.

without its signal peptide, in *E. coli* shuffle T7 *lys*Y (Fig. S1) which are capable of forming

disulfide bonds (Lobstein *et al.*, 2012). We obtained a yield of 0.1mg of pure protein per

litre of culture. After peptide purification, UPLC-ESI-HRMS analysis of the purified

AgLTP24 protein revealed a monoisotopic $[M+H]^+$ m/z 9686.8008 (theoretical

monoisotopic m/z calculated for $C_{415}H_{676}N_{123}O_{126}S_9$: 9686.7760) consistent with the

formation of four disulfide bridges (Fig. S4). This analysis was made after each run of

purification and before bioassays to check the purity and the good conformation of the

protein with disulphide bridges.

Physiology of *Frankia alni* in contact with AgLTP24

A miniaturised, rapid and fluorescent bioassay using resazurin dye was successfully

developed to determine whether AgLTP24 had antimicrobial activity against *Frankia*

alni at 7 dpi. For this, we applied a range of concentrations of AgLTP24 from 1 to 5μM

(Fig. 4) and used kanamycin as positive control of inhibition of *Frankia*. The cell viability

of *Frankia* is reduced from a concentration of 5μM AgLTP24 (54 ± 16% of cell viability)

while no effect was detected below this concentration. AgLTP24 can thus be considered

an antibacterial peptide against the symbiotic partner at this concentration. In addition,

microscopy observation showed a negative effect of AgLTP24 on vesicles' production

(Fig. **4b**). Indeed, at 5μM of AgLTP24, no vesicle was observed whereas they are present

at sub-inhibitory concentrations. However, *Frankia* must be viable and efficient in the

nodule to ensure nitrogen fixation for trophic exchange with its plant partner. This

conducted us to perform a second physiological assay by using sub-inhibitory

concentrations of AgLTP24. The second physiological assay was conducted with a range

of AgLTP24 from 1nM to 1µM (Fig. 5). In this test, nitrogen fixation, respiratory activity

(OD_{490nm}) and growth effects (OD_{600nm}) were monitored over a 10 days' time course (Fig.

S5).

That range of AgLTP24 had little effect on *Frankia* growth. We noted a negative effect at

10nM of AgLTP24 on respiratory activity at early time of the time course (Fig. **S5b**) but

it was not maintained over time. Regarding respiratory activity, we observed a positive

but not significant effect with AgLTP24 above 100nM and 5 days of growth. The most

striking effect was observed on nitrogen fixation after 7 days of culture (Fig. 5) with a

concentration above 100nM leading to inhibitory effect whereas no effect was observed

on growth or respiratory activity. Conversely, at 1nM of this peptide, the AgLTP24

improved this activity. Microscopy did show any effect on vesicle morphology (data not

shown).

Discussion

Early steps of the interaction between alder and *Frankia* start with the molecular

dialogue through the secretion of flavonoids by the plant (Benoit & Berry, 1997; Hughes

et al., 1999) and the bacterial "Nod-like" factors by *Frankia* (Cérémonie *et al.*, 1999;

Perrine-Walker *et al.*, 2011). This factor also called root hair deforming factor (RHDF)

because it triggers plant response by reorientating root-hair tip growth followed by the

formation of an infection thread (IT) and cell division induction in inner cortical cells.

This process leads to a nodule where *Frankia* will be housed and exchange with its plant

partner. Even though RHDF structure remains unknown, previous studies have shown

that this factor of around 3KDa is present in *Frankia* cell free supernatant and acts at

nanomolar dose (Cissoko *et al.*, 2018). In addition, this factor is able to induce a high

frequency nuclear Ca^{2+} spiking (Granqvist *et al.*, 2015). Due to the lack of a genetic

transformation system for alder, other approaches besides genetic inactivation have

been used to identify the host symbiotic determinants.

Focusing at nodule step (Hocher, V. *et al.*, 2011), transcriptomic analysis showed that

the common symbiotic cascade known in Legumes (interacting with rhizobia) and in

most land plants (interacting with VAM fungi) (Genre & Russo, 2016) was also present

in actinorhizal plants (Gherbi *et al.*, 2008; Hocher, V. *et al.*, 2011; Hocher, Valérie *et al.*,

2011). Previously, differential screening of cDNA libraries from root and nodules of

Alnus glutinosa with nodule and root cDNAs permitted to identify several symbiotic

genes among which a subtilisin-like protease (Ribeiro *et al.*, 1995), a sucrose synthase,

an enolase (van Ghelue *et al.*, 1996), and a dicarboxylate transporter (Jeong *et al.*, 2004).

These genes are evocative of tissue reorganization and trophic exchanges but shed no

light on the triggering of organogenesis. The present study aimed at investigating the

plant molecular response at early symbiotic stages by using same transcriptomic

microarray as in our previous work at nodule step (Hocher, V. *et al.*, 2011).

Global plant response at early step of symbiosis

Firstly, the two experiments induced different level of response in the plant partners

with stronger signal in the Exp1, suggesting that even though the supernatant is

sufficient to trigger root-hair deformation, a sustained dynamic of interaction with

Frankia is more efficient. These results suggest that *Frankia* secretome produced during

Exp1 could be more diverse triggering a strong perception of its presence by plant.

Indeed, *Frankia* secretes a variety of proteolytic enzymes such as glycosidases,

esterases, or proteases presumably involved in root infection (Mastronunzio *et al.*,

2008).

Plant global response reveals a common pattern with a strong modulation of genes

involved in stress response. This biological process, specific to early response (Hocher,

V. *et al.*, 2011) is constituted by genes encoding putative proteins involved in oxidative

stress (oxidase, peroxidase); to defense against pathogens such as pathogenesis related

(PR) proteins (chitinase, Defensin, thaumatin or nsLTP). Beside these biological

processes, we can propose different steps in the plant response after *Frankia* contact.

Oxidative stress in hairy roots repressed in nodule

Focusing on oxidative stress, both the overall GO analysis showed a strong induction of

genes involved in oxido-reduction metabolism (oxidase, peroxidase). These enzymes

involved in oxidative stress to reduce reactive oxygen species (ROS) suggesting that

Frankia supernatant induces a ROS stress such as previously observed for rhizobia at

early steps of interaction (Peleg-Grossman *et al.*, 2009; Peleg-Grossman *et al.*, 2012). In

legume roots, the enzymatic activities of catalase, ascorbate peroxidase, glutathione

reductase or NADPH oxidase significantly increase upon inoculation with bacterial

symbiont (Bueno *et al.*, 2001; Den Herder *et al.*, 2007; Peleg-Grossman *et al.*, 2009;

Peleg-Grossman *et al.*, 2012). After this important release, ROS production reduced

drastically to prevent root hair curling and IT formation (Shaw & Long, 2003; Lohar *et*

al., 2007).

To follow this dynamics of expression over the course of symbiosis establishment, we

extracted transcriptional data from our previous work made at nodule step (Hocher, V.

et al., 2011). Few genes are not modulated or repressed in the nodule steps (Table S4,

Fig. **2**), suggesting a similar profile. Consequently, plant after *Frankia* internalization

would repress this stress response but the measure of ROS abundance over the time

course of the symbiosis is necessary to support this tendency.

Perception of Frankia factor, signaling and hormonal function

Little is known about the symbiotic signals produced by *Frankia* but their perception

requires plant receptor. Here, a statistical analysis revealed a moderate

overrepresentation of gene involved in signal transduction (Table 1; AG-N01f_037_C05)

encoding a putative receptor like kinase (RLK) somewhat repressed in the nodule

suggesting its potential implication in *Frankia* recognition. However, other RLK similar

to Lys6/Lys7/Nfr1 of legume NF receptors (AG-R01f_025_F02) already found in *A.*

glutinosa (Hocher, V. *et al.*, 2011) are not modulated after 2.5 days, which raises the

question of *Frankia* recognition. Plant RLKs have been shown to control the initiation,

development, and maintenance of symbioses with beneficial mycorrhizal fungi and

rhizobia (Buendia *et al.*, 2018; Chiu & Paszkowski, 2020) but the mechanism in

actinorhizal symbiosis remain unknown (Svistoonoff *et al.*, 2014).

The CSSP transduction will trigger primordium organogenesis where hormonal balance

plays a crucial function and particularly auxin. Indeed, exogenous auxins treatments

lead to the formation of thick lateral roots resembling nodules in actinorhizal Fagales

(Hammad *et al.*, 2003; Svistoonoff *et al.*, 2003) and an auxin influx inhibitor perturbs the

formation of nodules in another actinorhizal model, *Casuarina glauca* (Peret *et al.*,

2008). Transcriptional data at early steps sustain the pivotal function of auxin by

demonstrating the overexpression of several genes encoding auxin binding proteins

(Table 1: AGCL1169Contig1) or auxin inducing proteins (Table S2: AGCL1376Contig1;

AG-N01f_043_C20). We did not detected genes involved in salicylic acid (SA) or jasmonic

acid (JA) biosynthesis in early steps of symbiosis however a fine bioinformatics study of

metabolic routes using the genome alder is required to facilitate putative assignation

and to validate or not their absence in those omic data.

In any case, perception of pathogen by plant triggers SA and JA accumulation and leads

to the accumulation of PR proteins to minimize pathogen load or disease onset. In

legume symbiosis, those hormones inhibit bacterial infection and nodule development

(Liu *et al.*, 2018) suggesting that NF recognition represses this accumulation.

Defense mechanisms activation

After oxidative stress, following *Frankia* contact, alder activates numerous genes

encoding PR proteins which are key components of plant innate immune system against

both biotic and abiotic stresses (Ali *et al.*, 2018). More precisely, PR proteins have

diverse functions such as glucanase, chitinases, thaumatin like, peroxidases, defensins,

nsLTPs or thionins.

We found numerous upregulated chitinases as observed in legumes (Staehelin *et al.*,

1994; Goormachtig *et al.*, 1998; Xie *et al.*, 1999) and described as important for rhizobia

infection and IT formation (Malolepszy *et al.*, 2018). Also, genes for oxidative stress and

chitinases are not modulated or repressed in late step of actinorhizal symbiosis (Table

S4, Fig. **2**) suggesting they are important in the early steps of *Frankia* infection.

Furthermore, genes encoding one uncharacterized PR proteins is upregulated after

Frankia contact (AG-R01f_016_N10) and rapidly repressed at nodule step (Table 1). A

similar profile was observed for PR10 in *Medicago trunculata* when associated to

symbiont or pathogen infection, it is activated but in symbiotic pathway, its expression

is transitory suggesting that defense signaling pathways are suppressed during the

establishment of symbiosis (Chen *et al.*, 2017). Here, the transitory activation of the

Alder PR protein could be also induced by *Frankia* factors even though a specific

response must be demonstrated by comparing to other biotic and abiotic stresses.

Some of the PR proteins are classified as AMPs based on their small size, the

conservation of cysteine rich motif and their potential action as antimicrobial compound

(Tam *et al.*, 2015). In alder transcriptome, we distinguished the overexpression of gene

encoding plant defensins (Ag3, Ag5 and Ag11), thaumatins (AG-N01f_038_P13) and

nsLTPs (AGCL115Contig1). Here, all AMPs induced at early steps are still overexpressed

in nodule (Table S4, Fig. **2**).

Like defensin, thaumatin like proteins considered as a member of defense protein but

also in plant development (de Jesús-Pires *et al.*, 2020) is overinduced in alder during

actinorhizal symbiosis. This expression profile is quite different to the well-documented

thaumatin like gene in soybean. Indeed, the *rj4* thaumatin like gene was described as

constitutively and similarly transcribed in roots and nodules and intervene at a very

early stage of IT formation to inhibit nodule formation with an incompatible strain

(Hayashi *et al.*, 2014). This suggests that thaumatin-like proteins in alder could

intervene in incompatibility function but its overexpression in nodule step also raise

question about another function in nodule organogenesis.

AgLTP24: from *Frankia* infection to nodule functioning

Function at early step

Global omic approach made at early steps of alder symbiosis pointed to a strong signal

from one protein belonging to nsLTPs family named here AgLTP24. The

immunolocalization on curled hairy roots induced after *Frankia* contact showed the

specific binding of AgLTP24 in nooks (Fig. **3**) where *Frankia* embedding before

internalization through IT formation. This suggests that AgLTP24 acts at the early phase

of infection. This is supported the induction of an nsLTP (MtN5) in *M. trunculata* at early

stage of symbiosis and in the nodule (Pii *et al.*, 2009; Pii *et al.*, 2012). Because its

silencing resulted in an increased number of curling events with a reduced number of

invading primordia, whereas its overexpression resulted in an increased number of

nodules, authors concluded that MtN5 is important at very early step of infection for the

successful establishment of the symbiotic interaction but not in nodule formation (Pii *et*

al., 2013). Similar to MtN5 (Pii *et al.*, 2009), AgLTP24 possesses a slight antimicrobial

activity *in vitro* against its symbiont at 5μM. As *Frankia* cell integrity must be preserved

for infection, we hypothesize that AgLTP24 concentration in contact with *Frankia* must

be below to 5μM. Further investigation to assess protein concentration in plant tissues is

a great challenge but requires the development of high-performance mass spectrometry

technology coupled to imagery (Gemperline *et al.*, 2015; Gemperline *et al.*, 2016).

Finally, AgLTP24 could be a gene player at early step of symbiosis to transduce signal

within plant roots to permit IT formation as well as induce stress response in *Frankia*

cells.

Function at late step

AgLTP24 is also present in nodule (Table 1) and targets specifically *Frankia's* vesicle in

nodule fixation zone (Fig. **3f**) but this localization is different from MtN5 binding in the

distal zone of the nodule (Pii *et al.*, 2012). However, a nsLTP (AsE246) found in

Astragalus sinicus specifically symbiosome membranes through binding a lipid

component: digalactosyldiacylglycerol (DGG) (Lei *et al.*, 2014). In this model, the *asE246*

silencing impacts nodule development with fewer matured infected plant cells.

Thus, AgLTP24 could fix vesicle wall composed by a multilamellate hopanoid lipid

envelopes (Berry *et al.*, 1993; Nalin *et al.*, 2000) or plant perisymbiontic membrane

coating *Frankia* cells in nodule. The perisymbiosome membrane composition is less

documented but non-inoculated *Alnus rubra* roots are mainly composed of glycerol,

phosphatidylethanolamine, phosphatidylserine and phosphatidylcholine (Berry *et al.*,

1991). A lipidomic study in *A. glutinosa* roots using recent technology followed by a

bioassay binding is a great perspective to decipher AgLTP24 mechanisms involved.

As observed *in vitro*, AgLTP24 could act as antimicrobial compound by reducing cell

viability and vesicles formation of *Frankia* (Fig. 4) suggesting a potential role by

controlling *Frankia* proliferation in nodule. However, because expression of nitrogen

fixation gene cluster is more active in symbiotic conditions compared to N-free culture

condition (Alloisio *et al.*, 2010; Lurthy *et al.*, 2018), the concentration probably present

in the nodule fixation zone would be lower to this lethal concentration. It is worth to

note that at 1nM concentration of AgLTP24, the nitrogen fixation activity is induced a

higher but not statistically so significant without perturbing *Frankia* growth (Fig. 5). The

improvement of nitrogen fixation was also observed for the AgDef5 defensin

translocated by *A. glutinosa* to *Frankia*'s vesicle in nodule (Carro *et al.*, 2015b).

In addition, this defensin is also upregulated at 2.5 dpi (Table 1) suggesting the plant

could deliver a cocktail of molecules including AgLTP24 to improve bacterial infection

and nitrogen fixation in nodule. A depth investigation of this synergic effect *in vitro*

opens a new perspective in addition to their genetic silencing in plant as an exciting

challenge to complete the gap of knowledge regarding their role in actinorhizal

symbiosis.

Acknowledgements

Thanks are expressed to Elise Lacroix and the Greenhouse facility (FR 41), to Danis

Abrouk in the ibio platform for help in submission of omic data to international

database, Jonathan Gervaix in the AME platform to access ARA measurement and

database, Jonathan Gervaix in the AME platform to access ARA measurement and

Philippe Bulet for his technical advice on peptide purification. We thank the PGE

platform for other measurements. This project has been funded by the FR BioEnviS

(Biodiversité, Environnement et Santé, Université Lyon1, France) and the EC2CO

(Ecosphère Continentale et Côtière) grant (reference: 10459).

Author contribution

M.G., N.A., A.H., P.N., P.P. and H.B. conceived and designed the study. M.G., N.A., P.F., S.B.,

O.K, J.T., P.D.S., P.P and H.B carried out the experiments. M.G., P.N., P.P. and H.B.

performed the data analysis. M.G., P.D.S., and H.B. performed the figure drawing. M.G.,

P.N., P.P., A.H. and H.B. provided critical biological interpretations of the data. P.N. and

H.B. wrote the manuscript.

Competing Interests

The authors declare no competing interests.

Bibliography

Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA, Tyagi A, Islam ST, Mushtaq M,
Yadav P, et al. 2018. Pathogenesis-related proteins and peptides as promising
tools for engineering plants with multiple stress tolerance. *Microbiological*
Research **212-213**: 29-37.

Alloisio N, Queiroux C, Fournier P, Pujic P, Normand P, Vallenet D, Medigue C,
Yamaura M, Kakoi K, Kucho K. 2010. The *Frankia alni* symbiotic transcriptome.
Mol Plant Microbe Interact **23**(5): 593-607.

Benoit LF, Berry AM. 1997. Flavonoid-like compounds from seeds of red alder (Alnus
rubra) influence host nodulation by Frankia (Actinomycetales). *Physiologia*
Plantarum **99**(4): 588-593.

Berry AM, Harriott OT, Moreau RA, Osman SF, Benson DR, Jones AD. 1993.
Hopanoid lipids compose the *Frankia* vesicle envelope, presumptive barrier of

oxygen diffusion to nitrogenase. *Proceedings of the National Academy of Sciences*
90(13): 6091-6094.

Berry AM, Moreau RA, Jones AD. 1991. Bacteriohopanetetrol: abundant lipid in
frankia cells and in nitrogen-fixing nodule tissue. *Plant Physiol* **95**(1): 111-115.

Berry AM, Torrey JG. 1983. Root hair deformation in the infection process of Alnus
rubra. *Botany* **61**: 2863-2876.

Brooks JM, Benson DRJS. 2016. Comparative metabolomics of root nodules infected
with Frankia sp. strains and uninfected roots from Alnus glutinosa and Casuarina
cunninghamiana reflects physiological integration. *Symbiosis* **70**(1): 87-96.

Buendia L, Girardin A, Wang T, Cottret L, Lefebvre B. 2018. LysM Receptor-Like
Kinase and LysM Receptor-Like Protein Families: An Update on Phylogeny and
Functional Characterization. *Frontiers in Plant Science* **9**(1531).

Bueno P, Soto MJ, Rodríguez-Rosales MP, Sanjuan J, Olivares J, Donaire JP. 2001.
Time-course of lipoxygenase, antioxidant enzyme activities and H2O2
accumulation during the early stages of Rhizobium–legume symbiosis. *New*
Phytologist **152**(1): 91-96.

Carro L, Persson T, Pujic P, Alloisio N, Fournier P, Boubakri H, Pawlowski K,
Normand P. 2015a. Organic acids metabolism in *Frankia alni. Symbiosis* **70**(1):
37-48.

Carro L, Pujic P, Alloisio N, Fournier P, Boubakri H, Hay AE, Poly F, Francois P,
Hocher V, Mergaert P, et al. 2015b. *Alnus* peptides modify membrane porosity
and induce the release of nitrogen-rich metabolites from nitrogen-fixing *Frankia*.
ISME J **9**(8): 1723-1733.

Carro L, Pujic P, Alloisio N, Fournier P, Boubakri H, Poly F, Rey M, Heddi A,
Normand P. 2016. Physiological effects of major up-regulated Alnus glutinosa
peptides on Frankia sp. ACN14a. *Microbiology* **162**(7): 1173-1184.

Cérémonie H, Debellé F, Fernandez MP. 1999. Structural and functional comparison
of Frankia root hair deforming factor and rhizobia Nod factor. *Canadian Journal of*
Botany **77**(9): 1293-1301.

Chen T, Duan L, Zhou B, Yu H, Zhu H, Cao Y, Zhang Z. 2017. Interplay of Pathogen-
Induced Defense Responses and Symbiotic Establishment in Medicago truncatula.
Frontiers in Microbiology **8**: 973-973.

Chiu CH, Paszkowski U. 2020. Receptor-Like Kinases Sustain Symbiotic
Scrutiny1 [OPEN]. *Plant Physiology* **182**(4): 1597-1612.

Chung CT, Niemela SL, Miller RH. 1989. One-step preparation of competent
Escherichia coli: transformation and storage of bacterial cells in the same
solution. *Proc Natl Acad Sci U S A* **86**(7): 2172-2175.

Cissoko M, Hocher V, Gherbi H, Gully D, Carré-Mlouka A, Sane S, Pignoly S,
Champion A, Ngom M, Pujic P, et al. 2018. Actinorhizal Signaling Molecules:
Frankia Root Hair Deforming Factor Shares Properties With NIN Inducing Factor.
Frontiers in Plant Science **9**(1494).

Conesa A, Gotz S. 2008. Blast2GO: A comprehensive suite for functional analysis in
plant genomics. *Int J Plant Genomics* **2008**: 619832.

D'Haeze W, Holsters M. 2002. Nod factor structures, responses, and perception during
initiation of nodule development. *Glycobiology* **12**(6): 79R-105R.

de Jesús-Pires C, Ferreira-Neto JRC, Pacifico Bezerra-Neto J, Kido EA, de Oliveira
Silva RL, Pandolfi V, Wanderley-Nogueira AC, Binneck E, da Costa AF, Pio-
Ribeiro G, et al. 2020. Plant Thaumatin-like Proteins: Function, Evolution and
Biotechnological Applications. *Curr Protein Pept Sci* **21**(1): 36-51.

Den Herder J, Lievens S, Rombauts S, Holsters M, Goormachtig S. 2007. A Symbiotic
Plant Peroxidase Involved in Bacterial Invasion of the Tropical Legume Sesbania
rostrata *Plant Physiology* **144**(2): 717-727.

Edstam MM, Viitanen L, Salminen TA, Edqvist J. 2011. Evolutionary history of the
non-specific lipid transfer proteins. *Mol Plant* **4**(6): 947-964.

Fahraeus G. 1957. The infection of clover root hairs by nodule bacteria studied by a
simple glass slide technique. *J Gen Microbiol* **16**(2): 374-381.

Fu C, Donovan WP, Shikapwashya-Hasser O, Ye X, Cole RH. 2014. Hot Fusion: an
efficient method to clone multiple DNA fragments as well as inverted repeats
without ligase. *PLoS One* **9**(12): e115318.

Gemperline E, Jayaraman D, Maeda J, Ane JM, Li L. 2015. Multifaceted investigation
of metabolites during nitrogen fixation in Medicago via high resolution MALDI-
MS imaging and ESI-MS. *J Am Soc Mass Spectrom* **26**(1): 149-158.

Gemperline E, Keller C, Jayaraman D, Maeda J, Sussman MR, Ané JM, Li L. 2016.
Examination of Endogenous Peptides in Medicago truncatula Using Mass
Spectrometry Imaging. *J Proteome Res* **15**(12): 4403-4411.

Genre A, Russo G. 2016. Does a common pathway transduce symbiotic signals in plant–
microbe interactions? *Front Plant Sci.* **7**: 96.

Ghelue MV, Løvaas E, Ringø E, Solheim B. 1997. Early interactions between Alnus
glutinosa and Frankia strain ArI3. Production and specificity of root hair
deformation factor(s). *Physiologia Plantarum* **99**(4): 579-587.

Gherbi H, Markmann K, Svistoonoff S, Estevan J, Autran D, Giczey G, Auguy F, Peret
B, Laplaze L, Franche C, et al. 2008. SymRK defines a common genetic basis for
plant root endosymbioses with arbuscular mycorrhiza fungi, rhizobia, and
Frankiabacteria. *Proc Natl Acad Sci U S A* **105**(12): 4928-4932.

Goormachtig S, Lievens S, Van de Velde W, Van Montagu M, Holsters M. 1998.
Srchi13, a Novel Early Nodulin from Sesbania rostrata, Is Related to Acidic Class
III Chitinases. *The Plant Cell* **10**(6): 905-915.

Granqvist E, Sun J, Op den Camp R, Pujic P, Hill L, Normand P, Morris RJ, Downie JA,
Geurts R, Oldroyd GE. 2015. Bacterial-induced calcium oscillations are common
to nitrogen-fixing associations of nodulating legumes and nonlegumes. *New*
Phytol **207**(3): 551-558.

Griesmann M, Chang Y, Liu X, Song Y, Haberer G, Crook MB, Billault-Penneteau B,
Lauressergues D, Keller J, Imanishi L, et al. 2018. Phylogenomics reveals
multiple losses of nitrogen-fixing root nodule symbiosis. *Science* **361**(6398):
eaat1743.

Hammad Y, Nalin R, Marechal J, Fiasson K, Pepin Rg, Berry AM, Normand P,
Domenach A-M. 2003. A possible role for phenyl acetic acid (PAA) on Alnus
glutinosa nodulation by Frankia. *Plant Soil* **254**(1): 193-205.

Hay A-E, Herrera-Belaroussi A, Rey M, Fournier P, Normand P, Boubakri H. 2020.
Feedback Regulation of N Fixation in Frankia-Alnus Symbiosis Through Amino
Acids Profiling in Field and Greenhouse Nodules. *Molecular Plant-Microbe*
Interactions® **33**(3): 499-508.

Hay AE, Boubakri H, Buonomo A, Rey M, Meiffren G, Cotin-Galvan L, Comte G,
Herrera-Belaroussi A. 2017. Control of endophytic *Frankia* sporulation by
Alnus nodule metabolites. *Mol Plant Microbe Interact* **30**: 205-214.

Hayashi M, Shiro S, Kanamori H, Mori-Hosokawa S, Sasaki-Yamagata H, Sayama T,
Nishioka M, Takahashi M, Ishimoto M, Katayose Y, et al. 2014. A Thaumatin-
Like Protein, Rj4, Controls Nodule Symbiotic Specificity in Soybean. *Plant and Cell*
Physiology **55**(9): 1679-1689.

Hocher V, Alloisio N, Auguy F, Fournier P, Doumas P, Pujic P, Gherbi H, Queiroux C,
Da Silva C, Wincker P, et al. 2011. Transcriptomics of actinorhizal symbioses
reveals homologs of the whole common symbiotic signaling cascade. *Plant*
Physiol **156**(2): 700-711.

Hocher V, Alloisio N, Bogusz D, Normand P. 2011. Early signaling in actinorhizal
symbioses. *Plant signaling & behavior* **6**(9): 1377-1379.

Hughes M, Donnelly C, Crozier A, Wheeler CT. 1999. Effects of the exposure of roots
of Alnus glutinosa to light on flavonoids and nodulation. *Canadian Journal of*
Botany **77**(9): 1311-1315.

Jeong J, Suh S, Guan C, Tsay YF, Moran N, Oh CJ, An CS, Demchenko KN, Pawlowski
K, Lee Y. 2004. A nodule-specific dicarboxylate transporter from alder is a
member of the peptide transporter family. *Plant Physiol* **134**(3): 969-978.

Ktari A, Gueddou A, Nouioui I, Miotello G, Sarkar I, Ghodhbane-Gtari F, Sen A,
Armengaud J, Gtari M. 2017. Host Plant Compatibility Shapes the
Proteogenome of *Frankia coriariae*. *Front Microbiol* **8**: 720.

Lalonde M. 1979. Immunological and Ultrastructural Demonstration of Nodulation of
the European Alnus glutinosa (L.) Gaertn. Host Plant by an Actinomycetal Isolate
from the North American Comptonia peregrina (L.) Coult. Root Nodule. *Botanical*
Gazette **140**: S35-S43.

Lei L, Chen L, Shi X, Li Y, Wang J, Chen D, Xie F, Li Y. 2014. A Nodule-Specific Lipid
Transfer Protein AsE246 Participates in Transport of Plant-Synthesized Lipids to
Symbiosome Membrane and Is Essential for Nodule Organogenesis in Chinese
Milk Vetch. *Plant Physiology* **164**(2): 1045-1058.

Liu H, Zhang C, Yang J, Yu N, Wang E. 2018. Hormone modulation of legume-rhizobial
symbiosis. *Journal of Integrative Plant Biology* **60**(8): 632-648.

Lobstein J, Emrich CA, Jeans C, Faulkner M, Riggs P, Berkmen M. 2012. SHuffle, a
novel Escherichia coli protein expression strain capable of correctly folding
disulfide bonded proteins in its cytoplasm. *Microbial cell factories* **11**: 56-56.

Login FH, Balmand S, Vallier A, Vincent-Monégat C, Vigneron A, Weiss-Gayet M,
Rochat D, Heddi A. 2011. Antimicrobial peptides keep insect endosymbionts
under control. *Science* **334**(6054): 362-365.

Lohar DP, Haridas S, Gantt JS, VandenBosch KA. 2007. A transient decrease in
reactive oxygen species in roots leads to root hair deformation in the legume-
rhizobia symbiosis. *New Phytol* **173**(1): 39-49.

Lurthy T, Alloisio N, Fournier P, Anchisi S, Normand P, Pujic P, Boubakri H. 2018.
Molecular response to nitrogen starvation by *Frankia alni* ACN14a revealed by
transcriptomics and functional analysis with a fosmid library in Escherichia coli.
Research in Microbiology

Malolepszy A, Kelly S, Sørensen KK, James EK, Kalisch C, Bozsoki Z, Panting M,
Andersen SU, Sato S, Tao K, et al. 2018. A plant chitinase controls cortical
infection thread progression and nitrogen-fixing symbiosis. *eLife* **7**: e38874.

Mastronunzio JE, Tisa LS, Normand P, Benson DR. 2008. Comparative secretome
analysis suggests low plant cell wall degrading capacity in *Frankia* symbionts.
BMC Genomics **9**: 47.

Nalin R, Putra S, Domenach AM, Rohmer M, Gourbière F, Berry A. 2000. High
hopanoid/total lipids ratio in Frankia mycelia is not related to the nitrogen
status. *Microbiology* **146 (Pt 11)**: 3013-3019.

Navarro E, Bousquet J, Moiroud A, Munive A, Piou D, Normand P. 2003. Molecular
phylogeny of Alnus (Betulaceae), inferred from nuclear ribosomal DNA ITS
sequences. *Plant and Soil* **254**(1): 207-217.

Normand P, Fernandez MP 2019. *Frankia. Bergey's Manual of Systematics of Archaea*
and Bacteria: (eds W.B. Whitman, F.A. Rainey, P. Kämpfer, M.E. Trujillo, P. DeVos,
B. Hedlund and S. Dedysh), 1-19.

Normand P, Lalonde M. 1982. Evaluation of *Frankia* strains isolated from provenances
782   of two Alnus species. *Canadian Journal of Microbiology* **28**(10): 1133-1142.

783   **Normand P, Lapierre P, Tisa LS, Gogarten JP, Alloisio N, Bagnarol E, Bassi CA, Berry**
784   **AM, Bickhart DM, Choisne N, et al. 2007.** Genome characteristics of
785   facultatively symbiotic *Frankia* sp. strains reflect host range and host plant
786   biogeography. *Genome Res* **17**(1): 7-15.

787   **Peleg-Grossman S, Golani Y, Kaye Y, Melamed-Book N, Levine A. 2009.** NPR1
788   protein regulates pathogenic and symbiotic interactions between Rhizobium and
789   legumes and non-legumes. *PLoS One* **4**(12): e8399.

790   **Peleg-Grossman S, Melamed-Book N, Levine A. 2012.** ROS production during
791   symbiotic infection suppresses pathogenesis-related gene expression. *Plant*
792   *signaling & behavior* **7**(3): 409-415.

793   **Peret B, Svistoonoff S, Lahouze B, Auguy F, Santi C, Doumas P, Laplaze L. 2008.** A
794   Role for auxin during actinorhizal symbioses formation? *Plant Signal Behav* **3**(1):
795   34-35.

796   **Perrine-Walker F, Gherbi H, Imanishi L, Hocher V, Ghodhbane-Gtari F, Lavenus J,**
797   **Benabdoun FM, Nambiar-Veeti M, Svistoonoff S, Laplaze L. 2011.** Symbiotic
798   signaling in actinorhizal symbioses. *Curr Protein Pept Sci* **12**(2): 156-164.

799   **Pii Y, Astegno A, Peroni E, Zaccardelli M, Pandolfini T, Crimi M. 2009.** The Medicago
800   truncatula N5 gene encoding a root-specific lipid transfer protein is required for
801   the symbiotic interaction with Sinorhizobium meliloti. *Mol Plant Microbe Interact*
802   **22**(12): 1577-1587.

803   **Pii Y, Molesini B, Masiero S, Pandolfini T. 2012.** The non-specific lipid transfer
804   protein N5 of *Medicago truncatula* is implicated in epidermal stages of *rhizobium*-
805   host interaction. *BMC Plant Biol* **12**: 233.

806   **Pii Y, Molesini B, Pandolfini T. 2013.** The involvement of Medicago truncatula non-
807   specific lipid transfer protein N5 in the control of rhizobial infection. *Plant*
808   *signaling & behavior* **8**(7): e24836-e24836.

809   **Prin Y, Neyra M, Diem HG. 1990.** Estimation of *Frankia* growth using Bradford protein
810   and INT reduction activity estimations: application to inoculum standardization.
811   *FEMS Microbiology Letters* **69**(1-2): 91-95.

812   **Pujic P, Alloisio N, Fournier P, Roche D, Sghaier H, Miotello G, Armengaud J, Berry**
813   **AM, Normand P. 2019.** Omics of the early molecular dialogue between *Frankia*
814   *alni* and *Alnus glutinosa* and the cellulase synton. *Environmental Microbiology*
815   **0**(0).

816   **Ribeiro A, Akkermans AD, van Kammen A, Bisseling T, Pawlowski K. 1995.** A
817   nodule-specific gene encoding a subtilisin-like protease is expressed in early
818   stages of actinorhizal nodule development. *Plant Cell* **7**(6): 785-794.

Schagger H. 2006. Tricine-SDS-PAGE. *Nat Protoc* **1**(1): 16-22.

Shaw SL, Long SR. 2003. Nod factor inhibition of reactive oxygen efflux in a host
legume. *Plant Physiology* **132**(4): 2196-2204.

Staehelin C, Granado J, Müller J, Wiemken A, Mellor RB, Felix G, Regenass M,
Broughton WJ, Boller T. 1994. Perception of Rhizobium nodulation factors by
tomato cells and inactivation by root chitinases. *Proceedings of the National*
Academy of Sciences **91**(6): 2196-2200.

Svistoonoff S, Hocher V, Gherbi H. 2014. Actinorhizal root nodule symbioses: what is
signalling telling on the origins of nodulation? *Curr Opin Plant Biol* **20C**: 11-18.

Svistoonoff S, Laplaze L, Auguy F, Runions J, Duponnois R, Haseloff J, Franche C,
Bogusz D. 2003. cg12 expression is specifically linked to infection of root hairs
and cortical cells during *Casuarina glauca* and *Allocasuarina verticillata*
actinorhizal nodule development. *Mol Plant Microbe Interact* **16**(7): 600-607.

Tam JP, Wang S, Wong KH, Tan WL. 2015. Antimicrobial Peptides from Plants.
Pharmaceuticals **8**(4): 711-757.

van Ghelue M, Ribeiro A, Solheim B, Akkermans AD, Bisseling T, Pawlowski K.
1996. Sucrose synthase and enolase expression in actinorhizal nodules of Alnus
glutinosa: comparison with legume nodules. *Mol Gen Genet* **250**(4): 437-446.

Xie Z-P, Staehelin C, Wiemken A, Broughton WJ, Müller J, Boller T. 1999. Symbiosis-
stimulated chitinase isoenzymes of soybean (Glycine max (L.) Merr.). *Journal of*
Experimental Botany **50**(332): 327-333.
840

Figure legends

Table 1: The up-regulated genes during early root hair deformation of *Alnus glutinosa* in

both experiments, with *Frankia* indirect contact (FIndC) and with *Frankia* supernatant

contact (FSupC).

Figure 1: Gene Ontology (GO) functional classification analysis. A. Distribution of

annotated genes into functional categories according to GO biological process. B. GO

functional classification analysis was made by comparing (A) up and down-regulated

genes found in Exp1 (Table S2) according to biological process, molecular function and

cellular component through Fisher test. Number of sequences (Nr) were indicated for

each category. BLAST2GO pipeline was used to get this GO annotation.

Figure 2: Expression levels of the genes which were commonly upregulated and

downregulated in FindC and FsupC conditions. The expression level values in nodule

were extracted from our previous work ((5); Table S4). The expression values were

expressed as Fold change (FC) with a color gradient from red (down-regulated; ≤0.5)

and orange-yellow (no-regulated) to green (up-regulated; ≥2) in infected roots or

nodules compared to control roots without Frankia. The GO annotations were assigned

by noting preferentially biological process and if not present, the molecular function.

Certain genes have no GO assignation after blast2Go analysis. The heatmap was

generated by GraphPad Prism 9.2.0.

Figure 3: AgLTP24 localization in roots after 2.5 days (A, B,C) or nodule of 21 dpi (D, E,

F). Hair magnification of a specific zone of the infected root (C) or nodule (F).

Immunofluorescence localization of AgLTP24 in hairy roots (B, C) and around Frankia

vesicles in infected cells (E and F). Negative control with rabbit serum before

intramuscular injection of AgLTP24 epitope (A, D) showed no green fluorescence in

hairy roots (A) or infected cells (D).

Blue: DAPI; red: autofluorescence; green: Alexa Fluor 488 dye anti-rabbit antibody (Life

Technologies, Saint Aubin, France). c, infected cells; nc, noninfected cells; v, *Frankia*

vesicles; x, xylem.

Figure 4: physiological bioassays on *Frankia* growth and cellular development. **A.**

Resazurin bioassay on *Frankia* ACN14a growth after supplementation of 1 to 5µM of

AgLTP24. Cell viability was calculated by normalizing the fluorescence of the assay with

the mean of the negative control (*Frankia* without AgLTP24). Kanamycin ($40\mu g.ml^{-1}$)

was used as positive control. Data are expressed as mean values ± SD. Differences

between normalized data were assessed by the Mann Whitney test (bilateral and

unpaired) compared to control. Graphic representation and statistical analysis of results

was conducted with GraphPad Prism version 9.2.0. *p value <0.05, **p value <0.01.

B. *Frankia* cells observation without or with gradual concentration of AgLTP24 from 0

(Control) to $5\mu M$ and obtained during the different physiological bioassays. v, *Frankia*

vesicles

Figure 5: effect of AgLTP24 in physiology of *Frankia* after 7 days of growth on **A**.

Nitrogen fixation; **B**. respiratory activity and **C**. growth by measuring OD_{600nm}. Data were

extracted from the kinetic assay (Fig. S5).

Data are expressed as mean values ± SD. Differences between means were assessed by

the Mann Whitney test (bilateral and unpaired) compared to control. Graphic

representation and statistical analysis of results was conducted with GraphPad Prism

9.2.0. *p value <0.05, **p value < 0.01.

Fig. 1: Gene Ontology (GO) functional classification analysis. **a.** Distribution of annotated genes into functional categories according to GO biological process. **b.** GO functional classification analysis was made by comparing up and down-regulated genes found in Exp1 (Table S2) according to biological process, molecular function and cellular component through Fisher test. Number of sequences (Nr) were indicated for each category. BLAST2GO pipeline was used to get this GO annotation.

Fig. 2: Expression levels of the genes which were commonly upregulated and downregulated in *F*indC and *F*supC conditions. The expression level values in nodule were extracted from our previous work ((5); Table S4). The expression values were expressed as Fold change (FC) with a color gradient from red (down-regulated; ≤0.5) and orange-yellow (no-regulated) to green (up-regulated; ≥2) in infected roots or nodules compared to control roots without *Frankia*. The GO annotations were assigned by noting preferentially biological process and if not present, the molecular function. Certain genes have no GO assignation after blast2Go analysis. The heatmap was generated by GraphPad Prism 9.2.0.

Fig. 3: **AgLTP24 localization in roots after 2.5 days (a, b, c) or nodule of 21 dpi (d, e, f).** Hair magnification of a specific zone of the infected root **(c)** or nodule **(f)**. Immunofluorescence localization of AgLTP24 in hairy roots **(b, c)** and around *Frankia* vesicles in infected cells **(e, f)**. Negative control with rabbit serum before intramuscular injection of AgLTP24 epitope showed no green fluorescence in hairy roots **(a)** or infected cells **(d).** Blue: DAPI; red: autofluorescence; green: Alexa Fluor 488 dye anti-rabbit antibody (Life Technologies, Saint Aubin, France). c, infected cells; nc, noninfected cells; v, *Frankia* vesicles; x, xylem; rh, root hairs.

Fig. 4: physiological bioassays on *Frankia* growth and cellular development. **a.** Resazurin bioassay on *Frankia* ACN14a growth after supplementation of 1 to 5μM of AgLTP24. Cell viability was calculated by normalizing the fluorescence of the assay with the mean of the negative control (*Frankia* without AgLTP24). Kanamycin (40μg.ml^{-1}) was used as positive control. Data are expressed as mean values ± SD. Differences between normalized data were assessed by the Mann Whitney test (bilateral and unpaired) compared to control. Graphic representation and statistical analysis of results was conducted with GraphPad Prism version 9.2.0. *p value <0.05, **p value <0.01.
b. *Frankia* cells observation without or with gradual concentration of AgLTP24 from 0 (Control) to 5μM and obtained during the different physiological bioassays. v, *Frankia* vesicles

Fig. 5: effect of AgLTP24 in physiology of *Frankia* after 7 days of growth on **a.** Nitrogen fixation; **b.** respiratory activity and **c.** growth by measuring OD_{600nm}. Data were extracted from the kinetic assay (Fig. S5).

Data are expressed as mean values ± SD. Differences between means were assessed by the Mann Whitney test (bilateral and unpaired) compared to control. Graphic representation and statistical analysis of results was conducted with GraphPad Prism 9.2.0. *p value <0.05, **p value < 0.01.

Table 1: The up-regulated genes during early root hair deformation of *Alnus glutinosa* in both experiments, with *Frankia* indirect contact (*F*IndC) and with *Frankia* supernatant contact (*F*SupC)

For contigs, **EMBL-Accession** numbers of all the contigated EST sequences are indicated.

Singleton or contig **nucleotide sequences** are indicated when they were not of high-enough quality for EMBL submission acceptance.

Fold Change (**FC**) is the ratio between *F*IndC roots and control roots and between *F*SupC roots and control roots, respectively.

The significant difference of gene expression between *F*IndC roots and control roots and between *F*SupC roots and control roots (FC ≥ 2 or ≤ 0.05) was assessed by a T- test at p<0.05

*:Results of the previously reported microarray study with *Alnus glutinosa* nodules are indicated: **FC*** is the ratio between nodules and non inoculated root and **p*** is the p value of a T-student test comparing these two conditions (Hocher et al, 2011). ns indicated non significant difference between nodules and non inoculated root (p<0.05)

For EMBL-accession number, see Suppl Table 3.

** AGCL3038Contig1 sequence is the 3' non coding region of basic blue copper gene (AG-N01f_023_I15).

Clone Name	*Frankia* indirect contact (*F*IndC)		*Frankia* supernatant contact (*F*SupC)		Nodule*		Sequence homology	GO annotation	Peptide signal	Theoretical pI/Mw
	FC	p < 0.05	FC	p < 0.05	FC*	p*				
AGCL115Contig1	140.65	1.78E-06	74.8	1.12E-05	2145.65	1.38E-02	lipid transfer protein	systemic acquired resistance	Yes	8.46 / 9700.28
AG-N01f_017_D15	24.72	1.72E-06	11.7	2.09E-03	173.71	5.66E-03	betatubulin 14	microtubule-based process	No	ND
AG-R01f_023_D12	10.02	3.30E-04	8.4	3.79E-04	0.55	NS	pectin methylesterase inhibitor	pectinesterase inhibitor activity	Yes	4.41 / 16267.16
AGCL581Contig1	8.78	5.70E-04	7.1	4.17E-03	2941.04	7.08E-04	NA		0 Yes	10.07 / 11702.98*
AGCL3639Contig1	8.57	1.35E-05	3.1	1.66E-02	55.49	9.27E-03	NA		0 ?	ND
AG-R01f_043_B03	7.89	4.73E-04	12.0	6.70E-03	0.90	NS	Major pollen allergen Aln g 1	cellular response to stimulus	No	5.57 / 15622.87
AG-R01f_030_G24	7.07	4.26E-08	17.4	5.13E-04	288.39	4.27E-03	plant basic secretory protein (BSP), peptidase M		0 Yes	ND
AG-N01f_038_P13	7.06	3.27E-04	2.9	1.54E-02	9.13	1.52E-02	thaumatin like protein precursor	cellular response to stimulus	0 Yes	ND
AGCL3713Contig1	6.77	5.79E-03	4.1	1.27E-02	155.02	1.24E-02	defensin cysteine rich antimicrobial protein, Ag5	cellular response to stimulus	Yes	9.22 / 8145.27
AGCL1289Contig1	6.04	4.08E-08	2.9	2.15E-03	1.56	6.22E-03	proline-rich extensin-related protein-like		0 Yes	9.60 / 16039.35
AG-R01f_030_E08	6.04	4.68E-04	3.4	7.87E-03	46.10	1.19E-02	defensin cysteine rich antimicrobial protein, Ag11	cellular response to stimulus	Yes	7.01 / 6276.13
AGCL3038Contig1	5.67	4.96E-06	11.3	2.83E-04	31.43	4.86E-03	3' non coding region of basic blue copper gene		0 Non coding region	ND
AGCL649Contig1	5.21	1.80E-05	2.3	8.49E-03	0.43	3.64E-03	class III peroxidase	oxidation-reduction process	Yes	ND
AGCL1440Contig1	5.20	9.33E-06	2.2	2.81E-02	1.01	NS	chitinase / Hevein / PR-4 / Wheatwin2	cellular response to stimulus	Yes	5.01 / 18391.98
AGCL1113Contig1	4.71	1.92E-03	2.3	4.94E-02	0.14	2.92E-03	NA		0 Very small peptide	ND
AG-R01f_023_I15	4.71	5.29E-07	11.7	3.03E-05	28.29	4.31E-03	basic blue copper protein	oxidation-reduction process	Yes	9.82 / 10324.85
AG-R01f_032_F18	4.56	3.81E-06	2.5	1.20E-02	0.42	5.29E-03	cytochrome P450	oxidation-reduction process	Yes	ND
AG-R01f_020_G10	4.41	3.52E-07	5.9	9.86E-05	0.11	1.52E-02	germin like protein 5		0 Yes	ND
AG-N01f_006_G15	4.37	6.67E-07	2.7	4.29E-04	0.91	7.86E-02	cationic peroxidase	oxidation-reduction process	Yes	ND
AG-N01f_002_C06	4.34	6.70E-03	3.6	6.67E-03	125.38	7.39E-04	defensin cysteine rich antimicrobial protein, Ag3		0 Yes	9.36 / 6178.00
AGCL1514Contig1	4.19	1.79E-04	4.5	3.88E-03	0.36	4.66E-03	class IV chitinase	chitin catabolic process	Partial sequence	ND
AG-J02f_002_N06	3.88	9.15E-06	4.0	2.54E-02	0.06	1.41E-02	class V chitinase	carbohydrate metabolic process	Partial sequence	ND
AG-R01f_019_G06	3.64	2.46E-05	2.1	3.41E-02	16.08	1.32E-02	NA		0 ?	ND
AG-R01f_001_A18	3.59	3.12E-07	4.5	2.70E-03	1.03	NS	PRp27-like protein		0 Yes	6.10 / 23102.67
AGCL1169Contig1	3.29	4.34E-03	2.9	7.86E-03	12.42	2.83E-02	auxin-binding protein ABP20	cellular response to stimulus	Yes	ND
AG-R01f_024_N21	3.14	4.19E-06	3.5	4.05E-02	1.11	NS	PRp27 like protein		0 Partial sequence	ND
AG-N01f_036_L04	3.03	2.01E-05	3.6	1.09E-03	1.10	NS	PRp27 like protein		0 Partial sequence	ND
AGCL194Contig1	2.99	5.21E-05	3.9	9.30E-05	0.92	NS	germin like protein 5	cellular response to stimulus	Yes	6.07 / 21323.34
AGCL1654Contig1	2.81	2.39E-05	2.5	2.47E-02	2.70	1.03E-02	glutathione S-transferase		0 No	ND
AG-N01f_024_P14	2.56	1.89E-04	2.4	3.49E-02	0.45	1.20E-02	cationic peroxidase	oxidation-reduction process	Partial sequence	ND
AGCL91Contig1	2.50	4.18E-05	2.5	1.21E-02	0.83	NS	PRp27 like protein		0 Yes	5.79 / 23032.54
AGCL1559Contig1	2.45	8.83E-05	2.9	1.87E-02	7.18	2.70E-03	mannitol dehydrogenase	oxidation-reduction process	Partial sequence	ND
AG-J02f_002_F23	2.37	8.29E-06	2.4	4.34E-05	1.77	NS	NA		0 Small sequence	ND
AG-N01f_037_C05	2.26	9.10E-05	38.9	2.25E-05	1.97	6.01E-03	receptor protein kinase perk1 like protein	cellular response to stimulus	No	ND
AG-R01f_012_L02	2.12	1.18E-02	2.3	3.32E-03	0.22	3.61E-05	phenylalanine ammonia lyase		0 Partial sequence	ND
gi_38453640_emb_AJ420771.1	2.18	9.83E-03	2.31	3.66E-03	ND	ND	phenylalanine ammonia-lyase		No	ND