GATA2 haploinsufficiency causes an epigenetic feedback mechanism resulting in myeloid
and erythroid dysplasia

Emanuele Gioacchino[1], Wei Zhang[1, *], Cansu Koyunlar[1, *], Joke Zink[1], Hans de Looper[1,2],

Kirsten J. Gussinklo[1], Remco Hoogenboezem[1], Dennis Bosch[1], Eric Bindels[1], Ivo P. Touw[1] and

Emma de Pater[1,2].

Affiliations

[1] Department of Hematology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

[2] Cancer Genome Editing Center, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

[*] equally contributed

Corresponding author: e.depater@erasmusmc.nl

Abstract

The transcription factor GATA2 has pivotal roles in hematopoiesis. Germline *GATA2*

mutations in patients result in GATA2 haploinsufficiency syndrome characterized by

immunodeficiency, bone marrow failure, and predispositions to myelodysplastic syndrome

(MDS) and acute myeloid leukemia (AML). Clinical symptoms in GATA2 patients are diverse

and mechanisms driving GATA2 related phenotypes largely unknown. To explore the impact

of GATA2 haploinsufficiency on hematopoiesis, we generated a zebrafish model carrying a

heterozygous mutation of *gata2b* (*gata2b*[+/-]), an orthologue of *GATA2*. Morphological

analysis revealed myeloid and erythroid dysplasia in *gata2b*[+/-] kidney marrow (KM). single

nucleus (sn)-ATAC-seq showed that the co-accessibility between the transcription start site

(TSS) and a +3.5-4.1kb enhancer was more robust in *gata2b*[+/-] zebrafish HSPCs compared to

wild type, increasing *gata2b* expression. This is suggestive of an auto-regulatory feedback

mechanism, where *gata2b* expression remains at sufficient levels after the loss of a single

allele to maintain the HSPC pool. As a result, *gata2b*[+/-] chromatin is also more accessible in

the erythroid and myeloid lineage, causing several defects. scRNA-seq data revealed a

differentiation delay in erythroid progenitors, hallmarked by downregulation of intrinsic

signals like cytoskeletal transcripts, aberrant proliferative signatures, and downregulation of

Gata1a, a master regulator of erythropoiesis, likely preceding erythroid dysplasia. This
shows that the cell intrinsic compensatory mechanisms for the maintenance of normal
levels of Gata2b to maintain HSPC integrity result in aberrant lineage differentiation and a
preleukemia syndrome.

Keywords

Gata2b, Zebrafish, MDS, Haploinsufficiency, HSCs

Introduction

The transcription factor GATA2 plays a major role in the generation and maintenance of
the hematopoietic system[1-3]. In humans, heterozygous germline mutations in GATA2 often
lead to a loss of function of one allele, causing GATA2 haploinsufficiency. The clinical
manifestations of GATA2 haploinsufficiency are broad and include immunodeficiency,
pulmonary-, vascular- and/or lymphatic dysfunctions and a strong propensity to develop
myelodysplastic syndromes (MDS) or acute myeloid leukemia (AML)[4,5]. These conditions,
now collectively referred to as GATA2 deficiency syndromes, were previously known as
Emberger syndrome[6], Monocytopenia and Mycobacterium Avium Complex (MonoMAC)
syndrome[7], dendritic cell, monocyte, B- and natural killer (NK) cell deficiency (DCML)[8], and
familial forms of AML[9]. In addition to the various disease phenotypes, the risk of developing
MDS/AML in GATA2 deficient patients is approximately 80% before the age of 40[4,5]. No clear
correlation is found between the occurrence of GATA2 mutations and the severity of
hematopoietic deficiencies, even among family members who share the same mutation[10,11].
Therefore, it is essential to gain insight into the mechanism of underlying GATA2 deficiencies
in well-defined uniform experimental models.

In mice, Gata2 has an essential regulatory function in hematopoietic stem cell (HSC)
generation and maintenance[1-3]. However, whereas Gata2-null mice are lethal at embryonic
day (E) 10.5[3], *Gata2* heterozygous (*Gata2$^{+/-}$*) mice survive to adulthood with normal blood
values. Notwithstanding this apparently normal blood phenotype, *Gata2$^{+/-}$* mice have a
diminished HSC compartment in the bone marrow (BM) showing a reduced repopulation
capacity in competitive transplantation studies[12,13]. Whereas mouse models thus emerged
as a useful source to identify the function of *GATA2* in HSC generation and fitness, they leave
the mechanisms causing the different aspects of GATA2 deficiency syndromes largely

undiscovered.

To better understand the biology of GATA2 haploinsufficiency syndromes, zebrafish
serves as an attractive alternative model. Zebrafish have the advantage of having two *GATA2*
orthologues; *Gata2a* and *Gata2b*. *Gata2a* is expressed predominantly in the vasculature[14]
and is required for programming of the hemogenic endothelium[15,16]. *Gata2b* is expressed in
hematopoietic stem/progenitor cells (HSPCs)[14] and homozygous deletion (*gata2b*[-/-])
redirects HSPC differentiation to the lymphoid lineage in expense of myeloid differentiation
causing a lymphoid bias with an incomplete B-cell differentiation in the kidney marrow (KM),
thus mimicking one of the GATA2 haploinsufficiency phenotypes found in patients.
Additionally, the most primitive HSC compartment was lost in *gata2b*[-/-] KM, but none of the
animals displayed signs of dysplasia[16,17]. Because patients carry heterozygous rather than
homozygous *GATA2* mutations, we specifically focused on how Gata2b haploinsufficiency
could be mechanistically linked to erythro-myelodysplasia, the major clinical hallmark of
GATA2 patients.

To validate the model, we first assessed hematopoietic cell differentiation in *gata2b*
heterozygous zebrafish (*gata2b*[+/-]) KM and observed erythroid dysplasia in all *gata2b*[+/-] but
not in wild-type (WT) zebrafish and myeloid dysplasia in 25% of the *gata2b*[+/-] zebrafish.
Subsequent single-cell RNA and ATAC sequencing analysis revealed an auto-regulatory
feedback mechanism of Gata2b where *gata2b* chromatin was over-accessible, underlying
the hematopoietic lineage differentiation defects in adults. Further characterization of
gata2b[+/-] zebrafish showed reduction of HSC proliferation, failure to initiate the "GATA2 to
GATA1 switch" and depletion of *Gata1a* and its co-factor *FOG1* in erythroid progenitor cells.
Taken together, these results reveal a dosage-dependent function of *Gata2b* and provide a
plausible explanation for the hematopoietic defects observed in GATA2-deficient patients.

Material and methods

Generation and genotyping of Gata2b heterozygous zebrafish

Gata2b[+/-] and wild type (WT) clutch mates were used for all analyses[16] and animals
were maintained under standard conditions. A knockout allele was generated by
introducing a 28bp out-of-frame insertion in exon 3 as previously described[16].

Zebrafish embryos were kept at 28.5°C on a 14h/10h light-dark cycle in HEPES-buffered
E3 medium. Zebrafish were anesthetized using tricaine and euthanized by ice-water. Animal

studies were approved by the animal Welfare/Ethics Committee in accordance to Dutch
legislation.

Kidney marrow isolation and analysis
Kidney marrow was dispersed mechanically using tweezers and dissociated by
pipetting in phosphate buffered saline (PBS)/10% fetal calf serum (FCS) to obtain single-cell
suspensions as previously described[16]. The KM cells were sorted in non-stick cooled micro
tubes (Ambion) containing 10% FCS in PBS. Proliferation was assessed by anti-Ki67 staining
in fixed (4% paraformaldehyde) KM cells. 7-AAD (7-amino-actinomycin D) (Stem-Kit
Reagents) 0.5mg/L or DAPI 1mg/L was used for live/dead discrimination. FACS sorting and
analysis were performed using FACS AriaIII (BD Biosciences).

May–Grünwald-Giemsa stain of KM smears
Kidney marrow smears were fixed in 100% MeOH before staining in May-Grünwald
solution (diluted 1:1 in phosphate buffer) and Giemsa solution (diluted 1:20 in phosphate
buffer) followed by a last rinsing step in tap water.
Morphological analysis was performed by pathologist by counting 200-500
hematopoietic cells of each kidney marrow smear; excluding mature erythrocytes and
thrombocytes. Cells were categorized as: blast, myelocyte, neutrophil, eosinophil,
lymphocyte or erythroblast. Furthermore, if dysplasia was observed within a specific
lineage, the percentage of dysplastic cells within that lineage was determined by additional
counting of at least 50 cells within that specific lineage.

Single cell omics sequencing
For single cell transcriptomics sequencing (scRNA-seq), kidney marrow cells were
isolated and 7×10^4 viable cells were sorted from 2 pooled *Tg(CD41:GFP[18]; runx1:DsRed[19])*
WT or *gata2b$^{+/-}$* male zebrafish at 1 year of age. For additional replicates, 7×10^4 single viable
cells were sorted from kidney marrows of one WT and one *gata2b$^{+/-}$* female zebrafish
between 18-20 months of age. cDNA was prepared using the manufacturers protocol
(Chromium Single Cell 3' reagent kits kit v3, 10x Genomics) and sequenced on a Novaseq
6000 instrument (Illumina). After sample processing and quality control analysis, 18,147

cells for WT and 10,849 cells for gata2b[+/-] were processed for further analysis. The read
depth was over 20K reads for all replicates.
We collected the same KM cell composition as scRNA-seq for single nucleus chromatin
accessibility sequencing (snATAC-seq, Chromium Single Cell ATAC reagent kits, 10x
Genomics). After low quality data filtering, 4,190 nuclei for WT and 7,572 nuclei for
gata2b[+/-] were processed for downstream analysis.
Data was analyzed using the Seurat[20] and Signac[21] pipeline. Trajectory inference was
performed using Monocle3[22] R packages.
133

Statistics

135   Statistical analysis was carried out in GraphPad Prism 8 (GraphPad Software). Unless
136   otherwise specified, data were analyzed using unpaired, 2-tailed Student's t-test. Statistical
137   significance was defined as $p<0.05$. Graphs are means ± standard error of mean (SEM) and
138   the number of replicates is indicated in the figure legend.
139

Results

Gata2b[+/-] zebrafish have erythroid and myeloid dysplasia in the kidney marrow

We first assessed hematopoietic cell morphology in KM smears of WT and *gata2b[+/-]*
zebrafish with ages ranging from 9 months post fertilization (mpf) to 18 mpf. Morphological
analysis showed that, while WT zebrafish had normal KM cell morphology, all *gata2b[+/-]* KM
samples had a considerable fraction of dysplastic cells in the erythroid lineage (Figure 1A,
panel 1-6). On average 0.5% of WT erythroid cells showed dysplastic features, compared to
9.9% of *gata2b[+/-]* erythroid cells (Figure 1A and B), the latter representing 4.5% of the total
kidney marrow population of *gata2b[+/-]* zebrafish. Myeloid lineage dysplasia was seen in the
gata2b[+/-] KM in 25% of the fish (Figure 1A, panel 7 and 8, and B). In these samples, 30% of
myeloid cells were dysplastic compared to 0.3% in WT. Myeloid dysplasia was mostly
represented by multi-lobulated nuclei, whereas the erythroid abnormalities ranged from
nuclear deformities and double nuclei to irregular cytoplasm or an almost complete lack of
cytoplasm (Figure 1A). The remaining cell types were not affected morphologically by
Gata2b haploinsufficiency. These results indicate that *gata2b* heterozygosity induces
dysplasia, predominantly in erythroid and myeloid progenitors.
156

To investigate if hematopoietic lineage differentiation ratios were still intact, WT and

gata2b[+/-] KM was assessed by flow cytometry (supplemental Figure 1A). *Gata2b*[+/-] zebrafish

KM showed no significant difference in the distribution of either mature myeloid-, erythroid-,

lymphoid- or progenitor- and HSPC populations compared to WT (supplemental Figure 1B-E).

Since GATA2 haploinsufficiency manifestations might require longer periods of time to

become evident[4,5], we tested the effect of *gata2b* heterozygosity in zebrafish up to 18

months of age. However, no significant differences were found in the ratios of different

lineages in *gata2b*[+/-] kidney marrow compared to WT during this period (supplemental

Figure 1B-E). Interestingly, the erythroid population significantly increased, indiscriminately

of genotype, after 12 months of age (P<0.001) (supplemental Figure 1D), while the

remaining myeloid, lymphoid and HSPC and progenitor populations did not vary as

dramatically (supplemental Figure 1B, C and E), suggesting aging results in an erythroid

biased hematopoietic system in zebrafish. To confirm the presence of a normal

hematopoietic lineage distribution in *gata2b*[+/-] kidney marrow we bred Gata2b-zebrafish

with transgenic GFP reporter zebrafish. Lineage analysis in transgenic lines specifically

marking neutrophils by *Tg(mpx:GFP)* (supplemental Figure 2A-B), T-cells by *Tg(lck:GFP)*

(supplemental Figure 2C-D), B-cells by *Tg(Igm:GFP)* (supplemental Figure 2E-F) and

Tg(mpeg:GFP) (supplemental Figure 2G-H) or monocytes by *Tg(mpeg:GFP)*[23,24]

(supplemental Figure 2I-J) showed no significant alterations in lineage distribution between

WT and *gata2b*[+/-] KM [25-28]. KM smear quantification by May–Grünwald-Giemsa (MGG)

staining revealed that *gata2b*[+/-] KM had a significant decrease in eosinophils compared to

WT (supplemental Figure 2K), representing less than 5% of the total KM cells. In summary,

based on flow cytometry analysis and transgenic reporters we conclude that the

differentiation in major hematopoietic lineages is not altered in *gata2b*[+/-] KM and that the

erythroid and myeloid dysplasia does not affect this.

Single cell RNA- and ATAC sequencing analysis reveals a cell maturation delay in gata2b[+/-]

KM

To investigate the molecular mechanisms underlying dysplasia in *gata2b*[+/-] KM, we

assessed chromatin accessibility, a measure for gene activation, and transcriptional

differences in dysplastic cell populations. We flow-sorted 4 cell populations based on light

scatter and observed dysplastic cells in the progenitor and lymphoid + HSPCs population of

gata2b[+/-] KM, indicating that dysplastic cells could be viably sorted (Figure 1A, panel 6). We

did not identify a uniquely separated population of dysplastic cells, possibly caused by a
heterogeneity in their shape. Therefore, we sorted progenitor and lymphoid + HSPC
populations from WT and *gata2b*[+/-] KMs and performed single-cell (sc) RNA- and single
nucleus (sn) ATAC-sequencing to identify transcriptome and epigenetic regulation signatures
defining dysplastic cells (Figure 2A). After filtering out low quality cells, we obtained 28,996
cells (WT = 18,147, *gata2b*[+/-] = 10,849) for scRNA-seq analysis and 11,762 cells (WT = 4,190,
gata2b[+/-] = 7,572) for snATAC-seq analysis (Figure 2B).

scRNA-sequencing resulted in 19 clusters based on a nearest neighbor algorithm using
the R Seurat package[20] (Figure 2B). Each cluster was classified based on differentially
expressed genes (supplemental Figure 3A-D) and known differentiation markers[29-34] (Figure
2C). We then used this labeled scRNA-seq dataset as a reference to interpret and identify 16
clusters in the snATAC-seq dataset based on the R Signac package[21] (Figure 2B, supplemental
Figure 4A-C). Cluster proportion analysis indicated an overall similar distribution of cells
within clusters between *gata2b*[+/-] and WT (Figure 2C). However, a clear difference in
differentiation trajectory was observed between WT and *gata2b*[+/-] KM in scRNA-seq and
snATAC-seq data based on the R package Monocle3[22], and particularly note the difference in
pseudo-time scores between WT and *gata2b*[+/-] erythroid progenitors 1 and myeloid
progenitors 1, suggesting a delayed differentiation in *gata2b*[+/-] KM (Figure 2D and E,
supplemental figure 4D). Whereas the pseudo-time score for HSPCs was similar between WT
and *gata2b*[+/-] zebrafish, the pseudo-time scores of differentiation of erythroid and myeloid
lineages were reduced in *gata2b*[+/-] KM (Figure 2D and E). Moreover, the expression of
gata2b was reduced in *gata2b*[+/-] cells compared to WT in the terminal trajectory of
hematopoietic development, where erythroid and myeloid progenitors are located (Figure
2F and G). Based on these findings, we conclude that *gata2b*[+/-] KM showed an overall
lineage maturation delay. As expected, a clear distinction in open chromatin patterns of
lineage specific transcription factors (TF) between immature and mature cells was seen in
WT KM (Supplemental figure 5A, C and E, left panels). In contrast, this distinction was less
pronounced in *gata2b*[+/-] KM (Supplemental figure 5B, D and F, left panels). Specifically, the
ETV family of transcription factors and the GATA family transcription factors were affected in
erythroid differentiation, SPI1, SPIB, and CEBPA family of transcription factors were affected
in myeloid differentiation, and TCF4 was most affected in lymphoid differentiation and
showed significantly different patterns of activation (Supplemental figure 5A-F). Surprisingly,
this did not always lead to a reduction in gene expression (Supplemental   figure5A-F, right

panels), particularly note the increased expression of *gata2b* in erythroid lineage
differentiation (Supplemental   figure 5B, right panel).
225

gata2b heterozygosity results in over-accessible gata2b chromatin leading to self-renewal
defects in HSPCs.
To further investigate the mechanism underlying the effects of haploinsufficient *gata2b*
expression, we assessed the accessible peak distribution of *gata2b* and its upstream region.
First of all, the chromatin accessibility signals of *gata2b* were strongest in the HSPCs cluster
(Figure 3A, top line), then gradually disappeared with cell maturation (Figure 3A, all other
lines), which is consistent with the GATA2 function in HSC maintenance[1-3]. However, when
we compared the signals between WT on the left and *gata2b*[+/-] on the right, we observed
that the chromatin of the *gata2b* gene body and upstream regions was more accessible in
gata2b[+/-] HSPCs. Chromatin co-accessibility analysis, which was used to predict
cis-regulatory interaction in the genome, suggested that there were two regions, +3.5-4.1kb
termed enhancer 1(E1) and +5.4-6.2kb, termed enhancer 2 (E2), which could be strongly
interacting with each other (Figure 3A, black arrows in Links). Moreover, E1 had stronger
interaction with *gata2b* transcription start site (TSS) in *gata2b*[+/-] KM (Figure 3A,   red arrows
in Links). These two upstream regions might work as enhancers for *gata2b* transcription.
When we assessed the cell distribution of the two enhancers and *gata2b* TSS (Figure 3B-G),
we observed that the accessibility signals of +5.4-6.2kb region-E2, (Figure 3B, C), was
accessible in HSPCs and *gata2b*[+/-] erythroid progenitors, while a +3.5-4.1kb region-E1,
(Figure 3D, E) was mainly accessible in HSPCs and *gata2b*[+/-] myeloid progenitors. This result
suggests that E1 and E2 might be responsible for different functions in hematopoiesis, and
particularly in *gata2b*[+/-] KM, E1 affects myeloid differentiation and E2 affects erythroid
differentiation. The percentage of cells with *gata2b* mRNA expression did not show a
significant difference as a result from the up-regulated chromatin accessibility and stronger
enhancer-TSS co-accessibility (Supplementary figure 6A and B), but we did see an increase
in the level of *gata2b* expression in the HSPC cluster as a result of the increase in
enhancer-TSS co-accessibility (Figure 3A, H, and I and Supplementary Figure 5B, right panel).
252

The functional consequence of epigenetic changes of the gata2b locus in HSPCs is a
reduction in G2-M phase in cell cycle
Because the most significant signal differences occurred in HSPC clusters and the cell

differentiation velocity differences appeared at "progenitors" clusters (Figure 2D), we

compared the different TF activity in progenitor-like clusters, including "HSPCs",

"progenitors", and "HSPCs and thrombocytes". *Gata2b*[+/-] progenitor-like clusters showed

significantly up-regulated chromatin accessibility of motifs of the BATF family of

transcription factors (Figure 4A). The BATF family of TFs are known to function in limiting the

self-renewal of HSCs in response to the accumulation of DNA damage[35], suggesting a defect

in self-renewal of *gata2b*[+/-] HSCs. To investigate this, proliferation was assessed in

CD41:GFP[int] purified HSCs (Figure 4B). Although numerically unaffected by *gata2b*

heterozygosity (Figure 4C), cell cycle phase distribution was markedly different in *gata2b*[+/-]

HSCs compared to WT (Figure 4D and E), with increased G1 phase and decreased G2-M

phase. As stem cells are known to have a short G1 phase to reduce cellular differentiation

potential, this may already be an indication that these HSCs lose self-renewal capacity and

differentiate, which is halted in the next differentiation stages[36].

Epigenetic dysregulation results in delayed erythroid maturation in gata2b[+/-] *kidney marrow*

Next, we sought to reveal the molecular mechanism behind the erythroid dysplasia in

Gata2b[+/-] KM. Compared to WT, *Gata2b*[+/-] progenitor-like clusters had significantly reduced

accessibility of GATA2 and GATA1::TAL1 motifs (Figure 4A). Despite the increase in *gata2b*

expression, this indicated an overall reduction of Gata2b function in activation of

downstream targets in the *gata2b*[+/-] zebrafish genome. Activation of GATA2 motifs were

mainly enriched in the erythroid progenitors (Figure 5A). An important downstream target

of Gata2 is Gata1, which is the master regulator of erythropoiesis[37,38]. We therefore

investigated *Gata1a* accessibility and expression in our dataset (Figure 5B-E). The *gata1a* TSS

was generally more accessible in *gata2b*[+/-] KM (Figure 5B), but not in erythroid progenitors 2

(Figure 5C). Furthermore, we found downregulation of *gata1a* mRNA expression (Figure 5D)

and the GATA1::TAL1 motif was less accessible (Figure 5E), further supporting the reduced

functionality of Gata1a in erythroid progenitors. This could be due to reduced expression of

zfpm1 (also known as *Friend of GATA (FOG1)*), which can facilitate *GATA1* expression[39](Figure

5F). The increased expression of *gata2b* in HSPCs (Figure 3I) and reduced expression of

gata1a in erythroid progenitors (Figure 5D) suggested that the "GATA factor switch", which is

a shift from GATA2 to GATA1 occupation during erythroid lineage differentiation[40-42], might

be silenced, causing deficiencies in erythroid lineage differentiation.

Considering that the dysplastic cells in KM smears were of myeloid or erythroid origin,

we hypothesized that dysplastic cells occupy a proportion of erythroid or myeloid clusters.
For a more precise examination of small populations along the myeloid and erythroid
differentiation lineage, we sub-clustered "HSPCs", "proliferative progenitors", "progenitors",
"myeloid progenitors 1", "myeloid progenitors 2", "erythroid progenitors 1", and "erythroid
progenitors 2" clusters and identified 5 significantly overrepresented sub-clusters in
gata2b[+/-] KM (Figure 5G-I, supplemental Figure 7 A-H). These sub-clusters represented 5.6%
of the total sequenced *gata2b*[+/-] cells, comparable to the proportion of dysplastic cells
observed in *gata2b*[+/-] KM. Differentially expressed genes (DEGs) analysis of overrepresented
sub-clusters in *gata2b*[+/-] KM compared to the rest of that cluster showed downregulation of
tubulin transcripts (not shown), suggesting a loss of cytoskeletal structure, a characteristic of
dysplasia. Besides, TF motif score analysis with snATAC-seq indicated a general reduction for
lineage-specific determination and differentiation motifs in *gata2b*[+/-] KM (supplemental
Figure 5A-F), suggesting that *gata2b*[+/-] cell maturation could be impaired resulting in a
differentiation block, as this is often the case in dysplasia[43].

To identify the mechanisms behind the erythroid differentiation delay in *gata2b*[+/-] KM,
we compared the DEGs and their function in the "erythroid progenitors 1" cluster. The result
showed that *gata2b*[+/-] "erythroid progenitors 1" cluster cells had upregulated signatures of
DNA replication together with the downregulation of transcripts necessary for mitosis
(Figure 5J), which can explain the origin of multi-lobulated nuclei and other nuclear
abnormalities observed in *gata2b*[+/-] dysplastic cells[44,45]. Pseudo-time trajectory inference
showed a diminished differentiation in *gata2b*[+/-] "erythroid progenitors 1" cluster with a
lower pseudo-time score (Figure 5K, L, and T). Pseudo-time dynamic gene expression level
also indicated the erythropoiesis differentiation delay in the *gata2b*[+/-] "erythroid progenitors
1" cluster with persistent high open chromatin, but lower expression of *gata1a*, which
should gradually decrease toward terminal erythroid maturation, and the decreased initial
level of *mki67*, indicative of incomplete differentiation (comparing Figure 5M-O to Figure
5P-R). This could be due to a diminished open chromatin structure at the locus leading to a
reduction in Gata2b binding sites to drive proper *Gata1a* expression, to support the
erythroid differentiation process. Furthermore, TF motif score analysis of the Erythroid
progenitors 1 cluster revealed a significant depletion for GATA family motifs, but an
enrichment for ETS and ETV family of transcription factor motifs in *gata2b*[+/-] "Erythroid
progenitors 1" cells (Figure 5S), suggesting an epigenetic dysregulation related to decreased
maturation of erythroblasts resulting from Gata2b haploinsufficiency.

Discussion

In humans, a balanced *GATA2* expression is essential for proper hematopoiesis.

Consequently, more than 80% of *GATA2* mutation carriers progress to hematological

malignancy by 40 years of age[4]. While the clinical consequences of *GATA2* mutations became

obvious over the last decades, the regulation of *GATA2* activity and its contribution to

human bone marrow failure syndromes and progression to hematological malignancy

remain incompletely understood.

Here, we used transgenic reporters, morphological phenotyping, and single-cell omics

sequencing to analyze *gata2b* heterozygous zebrafish. We showed that, while major

differentiation lineages remain intact, *gata2b* heterozygosity causes dysplasia in erythroid

and myeloid progenitors of zebrafish KM. snATAC-seq revealed an auto-regulatory feedback

mechanism for Gata2b haploinsufficiency, characterized by *gata2b* chromatin

over-accessibility and stronger co-accessibility of *gata2b* enhancers and TSS, indicating

epigenetic dysregulation in hematopoiesis in the zebrafish model. scRNA-seq analysis did

not identify a single population of dysplastic cells. Interestingly, a single erythroid progenitor

cluster was identified using snATAC-seq, and could indicate a specific dysplastic erythroid

population. The difference in recognition of the dysplastic cluster between scRNA-seq and

snATAC-seq may point to the underlying mechanism of the dysplasia where we have

evidence that the epigenetic compensatory mechanism of maintaining normal levels of

Gata2b may cause the dysplastic features during lineage differentiation, particularly in the

GATA2 to GATA1 switch and therefore affects erythroid lineage differentiation most. This also

underscores the importance of epigenetic analysis in other types of MDS and the fact that

many epigenetic regulators are mutated in MDS and AML[46]. In *gata2b*[+/-] erythroid

progenitors we found increased proliferative signatures together with decreased expression

of genes related to mitosis indicating an impaired cell cycle progression. Furthermore, we

found the depletion of GATA family factors indicating a failure of the "GATA factor switch"

that is indispensable for erythroid lineage differentiation[40-42]. We propose that these

alterations could play a role in the onset of dysplasia found in *gata2b*[+/-] zebrafish.

Interestingly, whereas *gata2b*[-/-] zebrafish display abrogated myeloid lineage

differentiation and a bias toward lymphoid differentiation [16,17], *gata2b*[+/-] did not simply

result in an intermediate phenotype between WT and *gata2b*[-/-]. Instead, Gata2b

haploinsufficiency uniquely caused dysplasia, not observed in $gata2b^{-/-}$, potentially caused
by compensatory mechanisms in haploinsufficiency that are otherwise overcome in
$gata2b^{-/-}$ zebrafish[47]. This may also be the reason that missense mutations have a different
molecular mechanism, because these alleles are not prone to mRNA degradation and may
thus not be compensated by epigenetic deregulation. The differences in the homozygous
and heterozygous Gata2b knockout phenotype support a role for gene dosage underlying
the GATA2 deficiency phenotype, possibly explaining the phenotypic heterogeneity between
patients. Since both erythroid and myeloid dysplasia can be observed in GATA2 patients[4], we
propose that the presence of dysplastic cells in $gata2b^{+/-}$ resembles the clinical phenotypes
associated with GATA2 heterozygosity. In the future, the isolation of single dysplastic cells
could help us to further explore the effect of Gata2 haploinsufficiency in malignant
transformation. Nevertheless, it remains to be established how $gata2b^{+/-}$ HSPCs would
respond to secondary insults such as infections or severe bleeding.

In conclusion, while the major lineage differentiation remains intact, $gata2b^{+/-}$ zebrafish
possess a stressed proliferative HSPC compartment which leads to the generation of
erythroid and myeloid dysplastic cells. Taken together, our model provides insights into the
consequences of Gata2b dosage, illustrated the alteration of the microenvironment, and
reveal how changes in epigenetics affect the outcome in lineage differentiation after Gata2b
haploinsufficiency in zebrafish.

Acknowledgements
We thank Dr Monteiro (University of Birmingham) for careful reading of the manuscript. We
thank the Experimental Animal Facility of Erasmus MC for animal husbandry and the
Erasmus Optical Imaging Center for confocal microscopy services. This research is supported
by the European Hematology Association (junior and senior non clinical research fellowship)
(EdP), the Dutch Cancer Foundation KWF/Alpe d'HuZes (SK10321)(EdP), the Daniel den Hoed
Foundation for support of the Cancer Genome Editing Center (IT) and the Josephine Nefkens
Foundation for purchase of the Chromium 10x (IT).

Authorship contributions
EdP and EG conceived the study; EG, CK, HdL, JZ, DB, and EB performed experiments; EG, WZ,
CK, RH, KG and EdP analysed results; IPT provided resources and EG, WZ, CK and EdP wrote
the manuscript and IPT revised the manuscript.

Disclosures

The authors declare no conflicts of interests

References

1.   Gao X, Johnson KD, Chang YI, et al. Gata2 cis-element is required for hematopoietic stem cell generation in the mammalian embryo. *J Exp Med*. Dec 16 2013;210(13):2833-42. doi:jem.20130733 [pii]

20130733 [pii]

10.1084/jem.20130733

2.   de Pater E, Kaimakis P, Vink CS, et al. Gata2 is required for HSC generation and survival. *J Exp Med*. Dec 16 2013;210(13):2843-50. doi:jem.20130751 [pii]

20130751 [pii]

10.1084/jem.20130751

3.   Tsai FY, Keller G, Kuo FC, et al. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature*. Sep 15 1994;371(6494):221-6. doi:10.1038/371221a0

4.   Donadieu J, Lamant M, Fieschi C, et al. Natural history of GATA2 deficiency in a survey of 79 French and Belgian patients. *Haematologica*. Aug 2018;103(8):1278-1287. doi:haematol.2017.181909 [pii]

1031278 [pii]

10.3324/haematol.2017.181909

5.   Spinner MA, Sanchez LA, Hsu AP, et al. GATA2 deficiency: a protean disorder of hematopoiesis, lymphatics, and immunity. *Blood*. Feb 6 2014;123(6):809-21. doi:S0006-4971(20)36040-7 [pii]

2013/515528 [pii]

10.1182/blood-2013-07-515528

6.   Ostergaard P, Simpson MA, Connell FC, et al. Mutations in GATA2 cause primary lymphedema associated with a predisposition to acute myeloid leukemia (Emberger syndrome). *Nat Genet*. Sep 4 2011;43(10):929-31. doi:ng.923 [pii]

10.1038/ng.923

7. Hsu AP, Sampaio EP, Khan J, et al. Mutations in GATA2 are associated with the

autosomal dominant and sporadic monocytopenia and mycobacterial infection

(MonoMAC) syndrome. *Blood.* Sep 8 2011;118(10):2653-5.

doi:S0006-4971(20)40686-X [pii]

2011/356352 [pii]

10.1182/blood-2011-05-356352

8. Dickinson RE, Griffin H, Bigley V, et al. Exome sequencing identifies GATA-2

mutation as the cause of dendritic cell, monocyte, B and NK lymphoid deficiency.

Blood. Sep 8 2011;118(10):2656-8. doi:S0006-4971(20)40687-1 [pii]

10.1182/blood-2011-06-360313

9. Hahn CN, Chong CE, Carmichael CL, et al. Heritable GATA2 mutations

associated with familial myelodysplastic syndrome and acute myeloid leukemia. *Nat*

Genet. Sep 4 2011;43(10):1012-7. doi:ng.913 [pii]

10.1038/ng.913

10. Mutsaers PG, van de Loosdrecht AA, Tawana K, Bodor C, Fitzgibbon J, Menko

FH. Highly variable clinical manifestations in a large family with a novel GATA2

mutation. *Leukemia.* Nov 2013;27(11):2247-8. doi:leu2013105 [pii]

10.1038/leu.2013.105

11. Wang X, Muramatsu H, Okuno Y, et al. GATA2 and secondary mutations in

familial myelodysplastic syndromes and pediatric myeloid malignancies.

Haematologica. Oct 2015;100(10):e398-401. doi:haematol.2015.127092 [pii]

100e398 [pii]

10.3324/haematol.2015.127092

12. Rodrigues NP, Janzen V, Forkert R, et al. Haploinsufficiency of GATA-2 perturbs

adult hematopoietic stem-cell homeostasis. *Blood.* Jul 15 2005;106(2):477-84.

doi:S0006-4971(20)53269-2 [pii]

10.1182/blood-2004-08-2989

13. Ling KW, Ottersbach K, van Hamburg JP, et al. GATA-2 plays two functionally

distinct roles during the ontogeny of hematopoietic stem cells. *J Exp Med.* Oct 4

2004;200(7):871-82. doi:jem.20031556 [pii]

20031556 [pii]

10.1084/jem.20031556

14. Butko E, Distel M, Pouget C, et al. Gata2b is a restricted early regulator of

hemogenic endothelium in the zebrafish embryo. *Development*. Mar 15

2015;142(6):1050-61. doi:142/6/1050 [pii]

DEV119180 [pii]

10.1242/dev.119180

15. Dobrzycki T, Mahony CB, Krecsmarik M, et al. Deletion of a conserved Gata2

enhancer impairs haemogenic endothelium programming and adult Zebrafish

haematopoiesis. *Commun Biol*. Feb 13 2020;3(1):71.

doi:10.1038/s42003-020-0798-3 [pii]

798 [pii]

10.1038/s42003-020-0798-3

16. Gioacchino E, Koyunlar C, Zink J, et al. Essential role for Gata2 in modulating

lineage output from hematopoietic stem cells in zebrafish. *Blood Adv*. Jul 13

2021;5(13):2687-2700. doi:S2473-9529(21)00348-7 [pii]

10.1182/bloodadvances.2020002993

17. Avagyan S, Weber MC, Ma S, et al. Single-cell ATAC-seq reveals

GATA2-dependent priming defect in myeloid and a maturation bottleneck in lymphoid

lineages. *Blood Adv*. Jul 13 2021;5(13):2673-2686. doi:S2473-9529(21)00347-5 [pii]

2020/ADV2020002992 [pii]

10.1182/bloodadvances.2020002992

18. Ma D, Zhang J, Lin HF, Italiano J, Handin RI. The identification and

characterization of zebrafish hematopoietic stem cells. *Blood*. Jul 14

2011;118(2):289-97. doi:S0006-4971(20)44778-0 [pii]

2010/327403 [pii]

10.1182/blood-2010-12-327403

19. Tamplin OJ, Durand EM, Carr LA, et al. Hematopoietic stem cell arrival triggers

dynamic remodeling of the perivascular niche. *Cell*. Jan 15 2015;160(1-2):241-52.

doi:S0092-8674(14)01638-9 [pii]

10.1016/j.cell.2014.12.032

20. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal

single-cell data. *Cell*. Jun 24 2021;184(13):3573-3587 e29.

doi:S0092-8674(21)00583-3 [pii]

10.1016/j.cell.2021.04.048

21. Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state

analysis with Signac. *Nat Methods*. Nov 2021;18(11):1333-1341.
doi:10.1038/s41592-021-01282-5 [pii]

10.1038/s41592-021-01282-5

22. Cao J, Spielmann M, Qiu X, et al. The single-cell transcriptional landscape of
mammalian organogenesis. *Nature*. Feb 2019;566(7745):496-502.
doi:10.1038/s41586-019-0969-x [pii]

10.1038/s41586-019-0969-x

23. Boatman S, Barrett F, Satishchandran S, Jing L, Shestopalov I, Zon LI. Assaying
hematopoiesis using zebrafish. *Blood Cells Mol Dis*. Dec 2013;51(4):271-6.
doi:S1079-9796(13)00160-5 [pii]

10.1016/j.bcmd.2013.07.009

24. Traver D, Paw BH, Poss KD, Penberthy WT, Lin S, Zon LI. Transplantation and in
vivo imaging of multilineage engraftment in zebrafish bloodless mutants. *Nat Immunol*.
Dec 2003;4(12):1238-46. doi:ni1007 [pii]

10.1038/ni1007

25. Langenau DM, Ferrando AA, Traver D, et al. In vivo tracking of T cell
development, ablation, and engraftment in transgenic zebrafish. *Proc Natl Acad Sci U*
S A. May 11 2004;101(19):7369-74. doi:0402248101 [pii]

1017369 [pii]

10.1073/pnas.0402248101

26. Renshaw SA, Loynes CA, Trushell DM, Elworthy S, Ingham PW, Whyte MK. A
transgenic zebrafish model of neutrophilic inflammation. *Blood*. Dec 15
2006;108(13):3976-8. doi:S0006-4971(20)52189-7 [pii]

10.1182/blood-2006-05-024075

27. Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. mpeg1 promoter
transgenes direct macrophage-lineage expression in zebrafish. *Blood*. Jan 27
2011;117(4):e49-56. doi:S0006-4971(20)58625-4 [pii]

2010/314120 [pii]

10.1182/blood-2010-10-314120

28. Page DM, Wittamer V, Bertrand JY, et al. An evolutionarily conserved program of
B-cell development and activation in zebrafish. *Blood*. Aug 22 2013;122(8):e1-11.
doi:S0006-4971(20)54254-7 [pii]

2012/471029 [pii]

10.1182/blood-2012-12-471029

29. Carmona SJ, Teichmann SA, Ferreira L, et al. Single-cell transcriptome analysis

of fish immune cells provides insight into the evolution of vertebrate immune cell

types. *Genome Res*. Mar 2017;27(3):451-461. doi:gr.207704.116 [pii]

10.1101/gr.207704.116

30. Danilova N, Bussmann J, Jekosch K, Steiner LA. The immunoglobulin

heavy-chain locus in zebrafish: identification and expression of a previously unknown

isotype, immunoglobulin Z. *Nat Immunol*. Mar 2005;6(3):295-302. doi:ni1166 [pii]

10.1038/ni1166

31. Kortum AN, Rodriguez-Nunez I, Yang J, et al. Differential expression and ligand

binding indicate alternative functions for zebrafish polymeric immunoglobulin receptor

(pIgR) and a family of pIgR-like (PIGRL) proteins. *Immunogenetics*. Apr

2014;66(4):267-79. doi:10.1007/s00251-014-0759-4

32. Macaulay IC, Svensson V, Labalette C, et al. Single-Cell RNA-Sequencing

Reveals a Continuous Spectrum of Differentiation in Hematopoietic Cells. *Cell Rep*.

Feb 2 2016;14(4):966-977. doi:S2211-1247(15)01538-7 [pii]

10.1016/j.celrep.2015.12.082

33. Tang Q, Iyer S, Lobbardi R, et al. Dissecting hematopoietic and renal cell

heterogeneity in adult zebrafish at single-cell resolution using RNA sequencing. *J Exp*

Med. Oct 2 2017;214(10):2875-2887. doi:jem.20170976 [pii]

20170976 [pii]

10.1084/jem.20170976

34. Athanasiadis EI, Botthof JG, Andres H, Ferreira L, Lio P, Cvejic A. Single-cell

RNA-sequencing uncovers transcriptional states and fate decisions in

haematopoiesis. *Nat Commun*. Dec 11 2017;8(1):2045.

doi:10.1038/s41467-017-02305-6 [pii]

2305 [pii]

10.1038/s41467-017-02305-6

35. Wang J, Sun Q, Morita Y, et al. A differentiation checkpoint limits hematopoietic

stem cell self-renewal in response to DNA damage. *Cell*. Mar 2 2012;148(5):1001-14.

doi:S0092-8674(12)00145-6 [pii]

10.1016/j.cell.2012.01.040

36. Pauklin S, Vallier L. The cell-cycle state of stem cells determines cell fate

propensity. *Cell*. Sep 26 2013;155(1):135-47. doi:S0092-8674(13)01025-8 [pii]

10.1016/j.cell.2013.08.031

37. Welch JJ, Watts JA, Vakoc CR, et al. Global regulation of erythroid gene

expression by transcription factor GATA-1. *Blood*. Nov 15 2004;104(10):3136-47.

doi:S0006-4971(20)55869-2 [pii]

10.1182/blood-2004-04-1603

38. Wang F, Zhu Y, Guo L, et al. A regulatory circuit comprising GATA1/2 switch and

microRNA-27a/24 promotes erythropoiesis. *Nucleic Acids Res*. Jan

2014;42(1):442-57. doi:gkt848 [pii]

10.1093/nar/gkt848

39. Cantor AB, Orkin SH. Coregulation of GATA factors by the Friend of GATA (FOG)

family of multitype zinc finger proteins. *Semin Cell Dev Biol*. Feb 2005;16(1):117-28.

doi:S1084-9521(04)00102-8 [pii]

10.1016/j.semcdb.2004.10.006

40. Grass JA, Boyer ME, Pal S, Wu J, Weiss MJ, Bresnick EH. GATA-1-dependent

transcriptional repression of GATA-2 via disruption of positive autoregulation and

domain-wide chromatin remodeling. *Proc Natl Acad Sci U S A*. Jul 22

2003;100(15):8811-6. doi:1432147100 [pii]

1008811 [pii]

10.1073/pnas.1432147100

41. Pal S, Cantor AB, Johnson KD, et al. Coregulator-dependent facilitation of

chromatin occupancy by GATA-1. *Proc Natl Acad Sci U S A*. Jan 27

2004;101(4):980-5. doi:0307612100 [pii]

1010980 [pii]

10.1073/pnas.0307612100

42. Bresnick EH, Lee HY, Fujiwara T, Johnson KD, Keles S. GATA switches as

developmental drivers. *J Biol Chem*. Oct 8 2010;285(41):31087-93.

doi:S0021-9258(19)88840-3 [pii]

R110.159079 [pii]

10.1074/jbc.R110.159079

43. Monika Belickova M, Merkerova MD, Votavova H, et al. Up-regulation of

ribosomal genes is associated with a poor response to azacitidine in myelodysplasia

and related neoplasms. *Int J Hematol*. Nov 2016;104(5):566-573.

doi:10.1007/s12185-016-2058-3 [pii]

10.1007/s12185-016-2058-3

44. Nakayama Y, Yamaguchi N. Multi-lobulation of the nucleus in prolonged S phase

by nuclear expression of Chk tyrosine kinase. *Exp Cell Res*. Apr 1

2005;304(2):570-81. doi:S0014-4827(04)00703-7 [pii]

10.1016/j.yexcr.2004.11.027

45. Ullah Z, Lee CY, Depamphilis ML. Cip/Kip cyclin-dependent protein kinase

inhibitors and the road to polyploidy. *Cell Div*. Jun 2 2009;4:10. doi:1747-1028-4-10

[pii]

10.1186/1747-1028-4-10

46. Haferlach T, Nagata Y, Grossmann V, et al. Landscape of genetic lesions in 944

patients with myelodysplastic syndromes. *Leukemia*. Feb 2014;28(2):241-7.

doi:leu2013336 [pii]

10.1038/leu.2013.336

47. Rossi A, Kontarakis Z, Gerri C, et al. Genetic compensation induced by

deleterious mutations but not gene knockdowns. *Nature*. Aug 13

2015;524(7564):230-3. doi:nature14580 [pii]

10.1038/nature14580

Figure legends

Figure. 1: Gata2b$^{+/-}$ kidney marrow shows erythroid and myeloid dysplasia

(A) Representative pictures of kidney marrow smears after May–Grünwald-Giemsa staining

of WT KM smears (panel 1 and 2) and *gata2b$^{+/-}$* KM smears (panel 3-8). 3) Blebbing in

cytoplasm of proerythroblast; 4) Irregular cytoplasm in erythroid precursor; 5) Lobed

nucleus and micronucleus in erythroblast; 6) Blebbing in cytoplasm of blast of sorted cell

after cytospin; 7) Binucleated promyelocyte; 8) Multinucleated promyelocyte.

(B) Frequency of dysplastic cells of the erythroid, myeloid and lymphoid lineage in KM

smears of WT (n=8) and *gata2b$^{+/-}$* (n=8) zebrafish. ***: P < 0.001

Figure. 2: Single cell transcriptome and chromatin accessibility analysis reveals a

differentiation block in gata2b$^{+/-}$ KM

(A) The flow cytometry-based sorting strategy for kidney marrow cells for scRNA-seq and

snATAC-seq analysis.

(B) UMAP analysis of scRNA-seq (top panels) and snATAC-seq (bottom panels) showing all

cell types, which was 19 cell types in scRNA-seq and 16 cell types in snATAC-seq. UMAP atlas

was separated between WT and *gata2b$^{+/-}$* cells.

(C) Cluster distribution quantification of scRNA-seq (left panel) and snATAC-seq (right panel)

between WT and *gata2b$^{+/-}$* cells, with significantly differentially distributed clusters

represented in pink. A statistical significance threshold of FDR < 0.05 and log2 fold change >

1 were applied to determine the significance of the observed differences.

(D-E) Box plots representing the comparison of pseudo-time scores across progenitor,

myeloid, and erythroid clusters in D) scRNA-seq data and E) snATAC-seq data. *: P < 0.05,

**: P < 0.01, ***: P < 0.0001, NS: not significant.

(F) WT (left panel) and $^{+/-}$ (right panel) expression of *gata2b* in scRNA-seq data in

pseudo-time. Dots are depicted if *gata2b* expression is detected and colored according to

cell type. The curved line shows the dynamic expression of *gata2b* throughout pseudo-time

trajectory.

(G) WT (left panel) and *gata2b$^{+/-}$* (right panel) chromatin accessibility of the *gata2b* locus in

snATAC-seq data in pseudo-time. Dots are depicted if open chromatin was detected and are

colored according to the cell types. The curved line shows the open chromatin at the *gata2b*

locus throughout pseudo-time trajectory.

Figure. 3: Differences in chromatin accessibility of Gata2b result in upregulation of Gata2b

expression in gata2b$^{+/-}$ HSPCs

(A) Coverage plots showing significant differences of *gata2b* chromatin accessibility

between WT and *gata2b$^{+/-}$* cells with the location of the *gata2b* gene block in blue. The

normalized peak signal range is 0 to 40. Highlighted genomic regions within grey blocks

indicate the location of two enhancers, enhancer 1 (E1) and enhancer 2 (E2). The link

between the blocks illustrate predicted *cis*-regulatory interactions between the *gata2b* gene

body and its upstream region, based on examination of genome co-accessibility. Link colors

correspond to predicted co-accessibility scores, with yellow indicating stronger predicted

interactions. Black arrows indicated the co-accessibility between E2 and E1. Red arrows

indicated the co-accessibility between E1 and TSS which is stronger in $gata2b^{+/-}$ KM.

(B-G) UMAP feature plots depict the open chromatin of Enhancer 2 (E2; B and C), Enhancer

1 (E1; D and E), and transcription start site (TSS; F and G) of $gata2b$. UMAP atlas were

separated by WT (B, D and F) and $gata2b^{+/-}$ cells (C, E and G).

(H) Open chromatin of the $gata2b$ locus per HSPCs cluster cell comparing WT and $gata2b^{+/-}$

HSPCs.

(I) Violin plot displaying the expression level of $gata2b$ in HSPCs in WT (pink) and $gata2b^{+/-}$

(turquois).

Figure 4. Functional consequences in HSPCs upon Gata2b overexpression

(A) The dot plot displays the log2 fold change values of differentially activated transcription

factor (TF) motifs in progenitor clusters, including "HSPCs", "progenitors", and "HSPCs and

thrombocytes", between WT and $gata2b^{+/-}$ cells. The TF motifs are ranked in increasing

order based on log2 fold change values. Dotted lines indicate the threshold for significance,

with absolute log2 fold change values > 0.58. The blue and red colors represent TF motifs

that are statistically significant (P value < 0.05) with absolute log2 fold change values > 0.58.

(B) Representative figure of identification of CD41:GFP^{int} population (in green) and

distribution in the FSC-A SSC-A kidney marrow population.

(C) Quantification of the percentage of CD41:GFP^{int} cells in WT (n=12) and $gata2b^{+/-}$ (n=12)

KM single live cells.

(D) Representative flow cytometry plots of cell cycle analysis by anti-Ki67 and DAPI staining

in WT and *gata2b*$^{+/-}$ CD41:GFP^{int} KM.

(E) Quantification of the percentages of WT (n=3) and *gata2b*$^{+/-}$ (n=3) CD41:GFP^{int} cells in

different cell cycle stages (* P value<0.05). Data represents as mean ± standard error of the

mean.

Figure. 5: Epigenetic changes resulting in the erythroid differentiation block in gata2*$^{+/-}$ *KM

(A) UMAP feature plots depicting open chromatin of GATA2 motif sequence indicating

activity of GATA2. UMAP atlas were separated by WT (left panel) and *gata2b*$^{+/-}$ cells (right

panel). The top left corner of UMAP plots display the TF sequence logo corresponding to the

analyzed motif, generated by the function *MotifPlot*.

(B) Coverage plots showing differences of *gata1a* chromatin accessibility between WT and

gata2b$^{+/-}$ cells.

(C) Percentage of cells derived from WT and *gata2b*$^{+/-}$ KM with accessibility of the *gata1a*

locus across clusters in snATAC-seq data.

(D) Percentage of cells derived from WT and *gata2b*$^{+/-}$ KM expressing *gata1a* across clusters

in scRNA-seq data.

(E) UMAP plots depict the activity of combined motif GATA1::TAL1. UMAP atlas were

separated by WT (left panel) and *gata2b*$^{+/-}$ cells (right panel). The top left corner of the

UMAP plots display the TF sequence logo corresponding to the analyzed motif, generated by

the function *MotifPlot*.

(F) Feature plot showing the *zfpm1* (FOG1) expression in WT and *gata2b*$^{+/-}$ KM.

(G-H) UMAP plots of the subclustering of the erythroid progenitors 1 cluster in scRNA-seq

data. UMAP atlas were separated by WT(G) and $gata2b^{+/-}$ cells (H)

(I) Quantification of distribution between WT and $gata2b^{+/-}$ erythroid progenitor 1

subclusters. Significantly differentially distributed clusters in pink. Dotted box indicates

overrepresented subcluster in $gata2b^{+/-}$ cells.

(J) STRING network of upregulated and downregulated transcripts in $gata2b^{+/-}$ "erythroid

progenitors 1". Only networks with more than 2 interactions were represented. Highlighted

in red DNA replication genes from KEGG pathways and highlighted in blue mitotic

cytokinesis genes from biological processes (Gene Ontology).

(K-L) Lineage trajectory analysis of WT (K) and $gata2b^{+/-}$ (L) erythroid progenitor 1

subclusters.

(M-R) WT (M-O) and $gata2b^{+/-}$ (P-R) pseudo-time expression of individual genes *hemoglobin*

beta adult 1 (*hbba1*) (M and P), *GATA binding protein 1 alpha (gata1a)* (N and Q), and *mki67*

(O and R). Dotted red boxes indicate the different initial expression of *mki67* (O and R). Fold

change > 0.05 & adjusted P value < 0.05. FDR=False discovery rate. FD=Fold difference.

(S) Dot plot displaying log2 fold change values of differentially activated TF motifs in the

erythroid progenitor 1 cluster between WT and $gata2b^{+/-}$ cells in snATAC-seq data. The TF

motifs are ranked in descending order based on log2 fold change values. Dotted lines

indicate the threshold (absolute log2 fold change < -0.58 and >0.58) for significance, The

blue (downregulated) and red (upregulated) colors represent TFs that are statistically

significant (P value < 0.05).

(T) Box plots representing the comparison of pseudo-time scores between WT and

gata2b$^{+/-}$across all subclusters in erythroid progenitors 1. *: P < 0.05, **: P < 0.01, NS: no

significance.

Supplementary figure. 1: Lineage gating of total KM in WT and gata2b$^{+/-}$ zebrafish

(A) Gating strategy of flowcytometry analysis of whole kidney marrow of WT and gata2b$^{+/-}$

zebrafish. Percentages represent the average of all zebrafish analyzed per genotype.

(B-E) Quantitation as percentages of the different cell populations in single viable cells of

myeloid (B), lymphoid and HSCs (C), erythrocytes (D) and progenitors (E) in WT and

gata2b$^{+/-}$ zebrafish kidney marrow over time. Statistical analysis showing as mean ±

standard error of the mean. ***: P < 0.001.

Supplementary figure. 2: Transgenic zebrafish reporter lines show no overall

differentiation difference in gata2b$^{+/-}$

(A) *mpx* positive cells, depicted in green, with distribution in FSC-A SSC-A graph.

(B) Quantification of the percentage of mpx:GFP+ cells in WT (n = 8) and gata2b$^{+/-}$ (n = 5)

kidney marrow single live cells.

(C) *Lck* positive cells, depicted in green, with distribution in FSC-A SSC-A graph.

(D) Quantification of the percentage of lck:GFP+ cells in WT (n = 4) and gata2b$^{+/-}$ (n = 7)

kidney marrow single live cells.

(E) *Igm* positive cells, depicted in green, with distribution in FSC-A SSC-A graph.

(F) Quantification of the percentage of IgM:GFP+ cells in WT (n = 3) and *gata2b*[+/-] (n = 3)

kidney marrow single live cells.

(G-J) *mpeg* positive cells, depicted in green, mark monocytes (G) and phagocytic B-cells (I).

Quantification of the percentage of mpeg:GFP+ cells (H and J) in WT (n = 7) and *gata2b*[+/-] (n

= 7) kidney marrow single live cells.

(K) Frequency of differentiated cell types in KM cells in smears of WT (n=8) and *gata2b*[+/-]

(n=8) zebrafish. Differentiated erythrocytes were excluded from quantification. *P

value<0.5. Data represents mean ± Standard error of the mean.

Supplementary Figure. 3: Transcriptomic heatmap and the expression of marker genes

(A) Unbiased heatmap representing the expression level of the top 5 expressed transcripts

per cluster. Transcripts highly expressed in multiple clusters were not repeated in the list

(like hemoglobin in erythroid clusters). Transcripts identified only by their chromosome

location were not included.

(B-D) FeaturePlot gene expression analysis of *CD41* (B) in HSPCs and thrombocytes, *mki67*

(C) in Proliferative progenitors, and *pcna* (D) predominantly in undifferentiated clusters.

Supplementary Figure. 4: Expression and accessibility of marker genes used for cluster

annotation and the pseudo-time trajectories inference

(A) Feature plot showing the scRNA-seq expression of marker gene *hemoglobin beta adult 1*

(*hbba1*) in the erythroid lineage, *granulin1* (*grn1*) in the myeloid lineage, *cluster of*

differentiation 37 (*CD37*) in the B cell lineage, and *GATA binding protein 2b* (*gata2b*) in

HSPCs to show cluster annotation.

(B) Feature plot showing the snATAC-seq gene activity of *hbba1* in the erythroid lineage,

grn1 in the myeloid lineage, *CD37* in the B lineage and *gata2b* in HSPCs to show cluster

annotation.

(C) Coverage plot illustrating the chromatin accessibility peaks of *hbba1* in the erythroid

lineage, *grn1* in the myeloid lineage, *CD37* in the B lineage, and *gata2b* in hematopoietic

stem and progenitor cells (HSPCs) to show cluster annotation.

(D) UMAP plot of scRNA-seq data (upper panels) and snATAC-seq data (bottom panels)

colored by pseudo-time scores representing cell developmental trajectories split by WT (left

panels) and *gata2b*$^{+/-}$ (right panels) cells. Lower pseudo-time scores correspond to the root

of trajectories, while higher pseudo-time scores represent the terminal differentiation

trajectories. The lines plotted on the UMAP atlas illustrate the differentiation paths of cells.

Supplementary Figure. 5: Motif regulation of lineages differentiation along pseudo-time

trajectories show reduced differentiation in gata2b$^{+/-}$ KM compared to WT.

(A) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) of erythroid cells differentiation along pseudo-time trajectory in WT

KM. Left panel depicting clusters in snATAC-seq data, including "HSPCs", "Progenitors",

"HPSCs and thrombocytes", "Erythroid progenitors 1", and "Erythroid progenitors 2", while

right panel depicting clusters in scRNA-seq data, including "HSPCs", "Progenitors",

"Proliferative progenitors", "HPSCs and thrombocytes", "Erythroid progenitors 1",

"Erythroid progenitors 2", and "Erythroid progenitors 3".

(B) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) during erythroid cell differentiation along pseudo-time trajectory in

gata2b$^{+/-}$ KM. The same clusters as those depicted in Figure S5A were used to generate the

heatmaps.

(C) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) during myeloid cells differentiation along pseudo-time trajectory in

WT KM. Left panel depicting clusters in snATAC-seq data, including "HSPCs", "Progenitors",

"HPSCs and thrombocytes", "Myeloid progenitors 1", "Myeloid progenitors 2",

"Neutrophils", and "Monocytes", while right panel depicting clusters in scRNA-seq data,

including "HSPCs", "Progenitors", "Proliferative progenitors", "HPSCs and thrombocytes",

"Myeloid progenitors 1", "Myeloid progenitors 2", "Neutrophils", and "Monocytes".

(D) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) during myeloid cell differentiation along pseudo-time trajectory in

gata2b$^{+/-}$ KM. The same clusters as those depicted in Figure S5C were used to generate the

heatmaps.

(E) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) during lymphoid cell differentiation along pseudo-time trajectory in

WT KM. Left panel depicting clusters in snATAC-seq data, including "HSPCs", "Progenitors",

"HPSCs and thrombocytes", "B cells and progenitors", "B cells", "T cells", and "NK", while

right panel depicting clusters in scRNA-seq data, including "HSPCs", "Progenitors",

"Proliferative progenitors", "HPSCs and thrombocytes", "B cells", "T cells", "NK", and "NK2".

(F) Heatmap plots showing the dynamic motif activity (left panel) and relative gene

expression (right panel) during lymphoid cells differentiation along pseudo-time trajectory

in *gata2b*$^{+/-}$ KM. The same clusters as those depicted in Figure S5E were used to generate

the heatmaps.

Supplementary Figure 6. Cell percentages with snATAC activity of the gata2b locus and

gata2b expression per cluster.

A) Cell percentages with detectable open chromatin of the *gata2b* locus per cluster. B) Cell

percentages with detectable *gata2b* expression per cluster.

Supplementary Figure 6. gata2b$^{+/-}$ ***cells have a different distribution of subclusters***

compared to WT

Split UMAP representing the cell distribution in the various subclusters (distinguishable by

different colors) in WT and *gata2b*$^{+/-}$ for HSPCs (A), Proliferative progenitors (B), Progenitors

(C), Myeloid progenitors 1 (D), Myeloid progenitors 2 (E), Erythroid progenitors 2 (F), HSPCs

and thrombocytes (G) , and Erythroid progenitors 3 (H), with respective quantification of

distribution between genotypes. Significantly differentially distributed subclusters in pink.

FDR<0.05 & Log2 fold change >1.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5