Compensatory mechanisms upon Gata2b haploinsufficiency causes aberrant
transcriptional regulatory signatures in HSPCs resulting in myeloid and erythroid dysplasia
in zebrafish

Emanuele Gioacchino[1], Wei Zhang[1, *], Cansu Koyunlar[1, *], Joke Zink[1], Hans de Looper[1,2],
Kirsten J. Gussinklo[1], Remco Hoogenboezem[1], Dennis Bosch[1], Eric Bindels[1], Ivo P. Touw[1] and
Emma de Pater[1,2].

Affiliations

[1] Department of Hematology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

[2] Cancer Genome Editing Center, Erasmus MC Cancer Institute, Rotterdam, The Netherlands

[*] equal contribution

Corresponding author: e.depater@erasmusmc.nl

13
Abstract

The transcription factor GATA2 has pivotal roles in hematopoiesis. Germline *GATA2*
mutations in patients result in GATA2 haploinsufficiency syndrome characterized by
immunodeficiency, bone marrow failure, and predispositions to myelodysplastic syndrome
(MDS) and acute myeloid leukemia (AML). Clinical symptoms in GATA2 patients are diverse
and mechanisms driving GATA2 related phenotypes largely unknown. To explore the impact
of GATA2 haploinsufficiency on hematopoiesis, we generated a zebrafish model carrying a
heterozygous mutation of *gata2b* (*gata2b*[+/-]), an orthologue of *GATA2*. Morphological
analysis revealed myeloid and erythroid dysplasia in *gata2b*[+/-] kidney marrow (KM). single
nucleus (sn)-ATAC-seq showed that the co-accessibility between the transcription start site
(TSS) and a +3.5-4.1kb enhancer was more robust in *gata2b*[+/-] zebrafish HSPCs compared to
wild type, increasing *gata2b* expression. This is suggestive of an auto-regulatory feedback
mechanism, where *gata2b* expression remains at sufficient levels after the loss of a single
allele to maintain the HSPC pool. As a result, *gata2b*[+/-] chromatin is also more accessible in
the erythroid and myeloid lineage, causing several defects. scRNA-seq data revealed a
differentiation block in erythroid progenitors, hallmarked by downregulation of intrinsic

signals like cytoskeletal transcripts, aberrant proliferative signatures, and downregulation of
Gata1a, a master regulator of erythropoiesis, likely preceding erythroid dysplasia. This
shows that the cell intrinsic compensatory mechanisms for the maintenance of normal
levels of Gata2b to maintain HSPC integrity result in aberrant lineage differentiation and a
preleukemia syndrome.

Keywords

Gata2b, Zebrafish, MDS, Haploinsufficiency, HSCs

Introduction

The transcription factor GATA2 plays a major role in the generation and maintenance of
the hematopoietic system[1-3]. In humans, heterozygous germline mutations in GATA2 often
lead to a loss of function of one allele, causing GATA2 haploinsufficiency. The clinical
manifestations of GATA2 haploinsufficiency are broad and include immunodeficiency,
pulmonary-, vascular- and/or lymphatic dysfunctions and a strong propensity to develop
myelodysplastic syndromes (MDS) or acute myeloid leukemia (AML)[4,5]. These conditions,
now collectively referred to as GATA2 deficiency syndromes, were previously known as
Emberger syndrome[6], Monocytopenia and Mycobacterium Avium Complex (MonoMAC)
syndrome[7], dendritic cell, monocyte, B- and natural killer (NK) cell deficiency (DCML)[8], and
familial forms of AML[9]. In addition to the various disease phenotypes, the risk of developing
MDS/AML in GATA2 deficient patients is approximately 80% before the age of 40[4,5]. No clear
correlation is found between the occurrence of GATA2 mutations and the severity of
hematopoietic deficiencies, even among family members who share the same mutation[10,11].
Therefore, it is essential to gain insight into the mechanism of underlying GATA2 deficiencies
in well-defined uniform experimental models.

In mice, Gata2 has an essential regulatory function in hematopoietic stem cell (HSC)
generation and maintenance[1-3]. However, whereas Gata2-null mice are lethal at embryonic
day (E) 10.5[3], *Gata2* heterozygous (*Gata2^{+/-}*) mice survive to adulthood with normal blood
values. Notwithstanding this apparently normal blood phenotype, *Gata2^{+/-}* mice have a
diminished HSC compartment in the bone marrow (BM) which show a reduced repopulation
capacity in competitive transplantation studies[12,13]. Whereas mouse models thus emerged
as a useful source to identify the function of *GATA2* in HSC generation and fitness, they leave

the mechanisms causing the different aspects of GATA2 deficiency syndromes largely
undiscovered.

To better understand the biology of GATA2 haploinsufficiency syndromes, zebrafish
serves as an attractive alternative model. Zebrafish have the advantage of having 2 *GATA2*
orthologues; *Gata2a* and *Gata2b*. *Gata2a* is expressed predominantly in the vasculature[14]
and is required for programming of the hemogenic endothelium[15,16]. *Gata2b* is expressed in
hematopoietic stem/progenitor cells (HSPCs)[14] and homozygous deletion (*gata2b$^{-/-}$*)
redirects HSPC differentiation to the lymphoid lineage in expense of myeloid differentiation
causing a lymphoid bias with an incomplete B-cell differentiation in the kidney marrow (KM),
thus mimicking one of the GATA2 haploinsufficiency phenotypes found in patients.
Additionally, the most primitive HSC compartment was lost in *gata2b$^{-/-}$* KM, but none of the
animals displayed signs of dysplasia[16,17]. Because patients carry heterozygous rather than
homozygous *GATA2* mutations, we hypothesize that Gata2b haploinsufficiency could be a
driver for leukemia onset. To investigate this, we assessed hematopoietic cell differentiation
in *gata2b* heterozygous zebrafish (*gata2b$^{+/-}$*) KM. We observed erythroid dysplasia in all
gata2b$^{+/-}$ but not in wild-type (WT) zebrafish and myeloid dysplasia in 25% of the *gata2b$^{+/-}$*
zebrafish. Single-cell RNA and ATAC sequencing analysis revealed an auto-regulatory
feedback mechanism of Gata2b where *gata2b* chromatin was over-accessible, underlying
the hematopoietic lineage differentiation defects in adults. Further characterization of
gata2b$^{+/-}$ zebrafish showed reduction of HSC proliferation, failure to initiate the "GATA2 to
GATA1 switch" and depletion of *Gata1a* and its co-factor *FOG1* in erythroid progenitor cells.
Taken together, our results reveal a concentration-dependent function of *Gata2b* and
hematopoietic defects in zebrafish hematopoiesis that reflect its dysfunction in
GATA2-deficient patients.

Material and methods

Generation and genotyping of Gata2b heterozygous zebrafish

Gata2b$^{+/-}$ and wild type (WT) clutch mates were used for all analyses[16] and animals
were maintained under standard conditions. A knockout allele was generated by
introducing a 28bp out-of-frame insertion in exon 3 as previously described[16].

Zebrafish embryos were kept at 28.5°C on a 14h/10h light-dark cycle in HEPES-buffered
E3 medium. Zebrafish were anesthetized using tricaine and euthanized by ice-water. Animal

studies were approved by the animal Welfare/Ethics Committee in accordance to Dutch
legislation.

Kidney marrow isolation and analysis

Kidney marrow was dispersed mechanically using tweezers and dissociated by
pipetting in phosphate buffered saline (PBS)/10% fetal calf serum (FCS) to obtain single-cell
suspensions as previously described[16]. The KM cells were sorted in non-stick cooled micro
tubes (Ambion) containing 10% FCS in PBS. Proliferation was assessed by anti-Ki67 staining
in fixed (4% paraformaldehyde) KM cells. 7-AAD (7-amino-actinomycin D) (Stem-Kit
Reagents) 0.5mg/L or DAPI 1mg/L was used for live/dead discrimination. FACS sorting and
analysis were performed using FACS AriaIII (BD Biosciences).

May–Grünwald-Giemsa stain of KM smears

Kidney marrow smears were fixed in 100% MeOH before staining in May-Grünwald
solution (diluted 1:1 in phosphate buffer) and Giemsa solution (diluted 1:20 in phosphate
buffer) followed by a last rinsing step in tap water.

Morphological analysis was performed by pathologist by counting 200-500
hematopoietic cells of each kidney marrow smear; excluding mature erythrocytes and
thrombocytes. Cells were categorized as: blast, myelocyte, neutrophil, eosinophil,
lymphocyte or erythroblast. Furthermore, if dysplasia was observed within a specific
lineage, the percentage of dysplastic cells within that lineage was determined by additional
counting of at least 50 cells within that specific lineage.

Single cell omics sequencing

For single cell transcriptomics sequencing (scRNA-seq), kidney marrow cells were
isolated and $7x10^4$ viable cells were sorted from 2 pooled *Tg(CD41:GFP[18]; runx1:DsRed[19])*
WT or *gata2b[+/-]* male zebrafish at 1 year of age. For additional replicates, $7x10^4$ single viable
cells were sorted from kidney marrows of one WT and one *gata2b[+/-]* female zebrafish
between 18-20 months of age. cDNA was prepared using the manufacturers protocol
(Chromium Single Cell 3' reagent kits kit v3, 10x Genomics) and sequenced on a Novaseq
6000 instrument (Illumina). After sample processing and quality control analysis, 18,147

cells for WT and 10,849 cells for *gata2b[+/-]* were processed for further analysis. The read
depth was over 20K reads for all replicates.

We collected the same KM cell composition as scRNA-seq for single nucleus chromatin
accessibility sequencing (snATAC-seq, Chromium Single Cell ATAC reagent kits, 10x
Genomics). After low quality data filtering, 4,190 nuclei for WT and 7,572 nuclei for
gata2b[+/-] were processed for downstream analysis.

Data was analyzed using the Seurat[20] and Signac[21] pipeline. Trajectory inference was
performed using Monocle3[22] R packages.

Statistics

Statistical analysis was carried out in GraphPad Prism 8 (GraphPad Software). Unless
otherwise specified, data were analyzed using unpaired, 2-tailed Student's t-test. Statistical
significance was defined as $p<0.05$. Graphs are means ± standard error of mean (SEM) and
the number of replicates is indicated in the figure legend.

Results

Gata2b[+/-] zebrafish have erythroid and myeloid dysplasia in the kidney marrow

We first assessed hematopoietic cell morphology in KM smears of WT and *gata2b[+/-]*
zebrafish with ages ranging from 9 months post fertilization (mpf) to 18 mpf. Morphological
analysis showed that, while WT zebrafish had normal KM cell morphology, all *gata2b[+/-]* KM
samples had a considerable fraction of dysplastic cells in the erythroid lineage (Figure 1A,
panel 1, 2, 4, 6, and 8-10). On average 0.5% of WT erythroid cells showed dysplastic features,
compared to 9.9% of *gata2b[+/-]* erythroid cells (Figure 1A and B), the latter representing 4.5%
of the total kidney marrow population of *gata2b[+/-]* zebrafish. Furthermore, we found
evidence of myeloid lineage dysplasia in the *gata2b[+/-]* KM in 25% of the fish (Figure 1A,
panel 3, 5 and 7, and B). In these samples, 30% of myeloid cells were dysplastic compared to
0.3% in WT. Myeloid dysplasia was mostly represented by multi-lobulated nuclei, whereas
the erythroid abnormalities ranged from nuclear deformities and double nuclei to irregular
cytoplasm or an almost complete lack of cytoplasm (Figure 1A). The remaining cell types
were not affected morphologically by Gata2b haploinsufficiency. These results indicate that
gata2b heterozygosity induces dysplasia, predominantly in erythroid progenitors.

To investigate if hematopoietic lineage differentiation ratios were still intact, WT and
gata2b[+/-] KM was assessed by flow cytometry (supplemental Figure 1A). *Gata2b*[+/-] zebrafish
KM showed no significant difference in the distribution of either mature myeloid-, erythroid-,
lymphoid- or progenitor- and HSPC populations compared to WT (supplemental Figure 1B-E).
Since GATA2 haploinsufficiency manifestations might require longer periods of time to
become evident[4,5], we tested the effect of *gata2b* heterozygosity in zebrafish up to 18
months of age. However, we found no significant differences in the ratios of different
lineages in *gata2b*[+/-] kidney marrow compared to WT during this period (supplemental
Figure 1B-E). Interestingly, the erythroid population significantly increased, indiscriminately
of genotype, after 12 months of age (P<0.001) (supplemental Figure 1D), while the
remaining myeloid, lymphoid and HSPC and progenitor populations did not vary as
dramatically (supplemental Figure 1B, C and E). To confirm the presence of a normal
hematopoietic lineage distribution in *gata2b*[+/-] kidney marrow we bred our zebrafish with
transgenic reporter zebrafish. Lineage analysis in transgenic lines specifically marking
neutrophils by *Tg(mpx:GFP)* (supplemental Figure 2A-B), T-cells by *Tg(lck:GFP)* (supplemental
Figure 2C-D), B-cells by *Tg(Igm:GFP)* and *Tg(mpeg:GFP)* (supplemental Figure 2E-F and I-J) or
monocytes by *Tg(mpeg:GFP)*[23,24] (supplemental Figure 2G-H) showed no significant
alterations in lineage distribution between WT and *gata2b*[+/-] KM [25-28]. KM smear
quantification by May–Grünwald-Giemsa (MGG) staining revealed that *gata2b*[+/-] KM had a
significant decrease in eosinophils compared to WT (supplemental Figure 2I), representing
less than 5% of the total KM cells. In summary, based on flow cytometry analysis and
transgenic reporters we conclude that the differentiation in major hematopoietic lineages is
not altered in *gata2b*[+/-] KM.
180
Single cell RNA- and ATAC sequencing analysis reveals a cell maturation delay in gata2b[+/-]
KM
To investigate the molecular mechanisms underlying dysplasia in *gata2b*[+/-] KM, we
assessed chromatin accessibility, a measure for gene activation, and transcriptional
differences in dysplastic cell populations. We flow-sorted 4 cell populations based on light
scatter and observed dysplastic cells in the progenitor and lymphoid + HSPCs population of
gata2b[+/-] KM, indicating that dysplastic cells could be viably sorted (Figure 1A, panel 9 and
10). We did not identify a uniquely separated population of dysplastic cells, possibly caused
by a heterogeneity in their shape. Therefore, we sorted progenitor and lymphoid + HSPC

populations from WT and *gata2b*[+/-] KMs and performed single-cell (sc) RNA- and
ATAC-sequencing to identify transcriptome and epigenetic regulation signatures defining
dysplastic cells (Figure 2A). After filtering out low quality cells, we obtained 28,996 cells (WT
= 18,147, *gata2b*[+/-] = 10,849) for scRNA-seq analysis and 11,762 cells (WT = 4,190, *gata2b*[+/-]
= 7,572) for snATAC-seq analysis (Figure 2B).

scRNA-sequencing resulted in 19 clusters based on a nearest neighbor algorithm using
the R Seurat package[20] (Figure 2B). Each cluster was classified based on differentially
expressed genes (supplemental Figure 3A-D) and known differentiation markers[29-34] (Figure
2C). We then used this labeled scRNA-seq dataset as a reference to interpret and identify 16
clusters in the snATAC-seq dataset based on the R Signac package[21] (Figure 2B-E). Cluster
proportion analysis indicated an overall similar distribution of cells within clusters between
gata2b[+/-] and WT (Figure 2F). However, a clear difference in differentiation trajectory was
observed between WT and *gata2b*[+/-] KM in snATAC-seq data based on the R package
Monocle3[22], and particularly note the difference in pseudo-time scores between WT and
gata2b[+/-] erythroid progenitors 1, suggesting a delayed differentiation in *gata2b*[+/-] KM
(Figure 2G). Whereas the initial pseudo-time score for cell differentiation at the HSPCs stage
was similar (Supplemental figure 4A), the pseudo-time scores of differentiation of erythroid,
myeloid, and lymphoid lineages were reduced in *gata2b*[+/-] KM (Figure 2G and supplemental
Figure 4A). Based on these findings, we conclude that *gata2b*[+/-] KM showed an overall
lineage maturation delay. As expected, a clear distinction in open chromatin patterns of
lineage specific transcription factors (TF) between immature and mature cells was seen in
WT KM (Supplemental Figure 4B, D and F, left panels). In contrast, this distinction was less
pronounced in *gata2b*[+/-] KM (Supplemental Figure 4C, E and G, left panels). Specifically, the
ETV family of transcription factors and the GATA family transcription factors were affected in
erythroid differentiation, SPI1, SPIB, and CEBPA family of transcription factors were affected
in myeloid differentiation, and TCF4 was most affected in lymphoid differentiation and
showed significantly different patterns of activation (supplemental Figure 4B-G). Surprisingly,
this did not always lead to a reduction in gene expression (Supplemental Figure 4 B-G, right
panels), particularly note the increased expression of *gata2b* in erythroid lineage
differentiation (Supplemental Figure 4C, right panel).

gata2b heterozygosity results in over-accessible gata2b chromatin leading to self-renewal
defects in HSPCs.

To further investigate the mechanism underlying the effects of haploinsufficient *gata2b*
expression, we assessed the accessible peak distribution of *gata2b* and its upstream region.
First of all, the chromatin accessibility signals of *gata2b* were strongest in the HSPCs cluster
(Figure 3A, top line), then gradually disappeared with cell maturation (Figure 3A, all other
lines), which is consistent with the GATA2 function in HSC maintenance[1-3]. However, when
we compared the signals between WT on the left and *gata2b*[+/-] on the right, we observed
that the chromatin of the *gata2b* gene body and upstream regions was more accessible in
gata2b[+/-] HSPCs. Chromatin co-accessibility analysis, which was used to predict
cis-regulatory interaction in the genome, suggested that there were two regions, +3.5-4.1kb
termed enhancer 1(E1) and +5.4-6.2kb, termed enhancer 2 (E2), which could be strongly
interacting with *gata2b* transcription start site (TSS) (Figure 3A, yellow lines in Links). These
two upstream regions might work as enhancers for *gata2b* transcription. When we assessed
the cell distribution of the two enhancers and *gata2b* TSS (Figure 3B-G), we observed that
the accessibility signals of +5.4-6.2kb region-E2, (Figure 3B, C), was accessible in HSPCs and
gata2b[+/-] erythroid progenitors, while a +3.5-4.1kb region-E1, (Figure 3D, E) was mainly
accessible in HSPCs and *gata2b*[+/-] myeloid progenitors. This result suggests that E1 and E2
might be responsible for different functions in hematopoiesis, and particularly in *gata2b*[+/-]
KM, E1 affects myeloid differentiation and E2 affects erythroid differentiation. The
percentage of cells with *gata2b* mRNA expression did not show a significant difference as a
result from the up-regulated chromatin accessibility and stronger enhancer-TSS
co-accessibility (Supplementary Figure 5A, B), but we did see an increase in the level of
gata2b expression in the HSPC cluster as a result of the increase in enhancer-TSS
co-accessibility (Figure 3A, H, and I and Supplementary Figure 4C, right panel).
246
The functional consequence of epigenetic changes of the gata2b locus in HSPCs is a
reduction in G2-M phase in cell cycle
Because the most significant signal differences occurred in HSPCs clusters and the cell
differentiation velocity differences appeared at "progenitors" clusters, we compared the
different TF activity in progenitor-like clusters, including "HSPCs", "progenitors", and "HSPCs
and thrombocytes". *Gata2b*[+/-] progenitor-like clusters showed significantly up-regulated
chromatin accessibility of motifs of the BATF family of transcription factors. The BATF family
of TFs are known to function in limiting the self-renewal of HSCs in response to the
accumulation of DNA damage[35] (Figure 4A), suggesting a defect in self-renewal of *gata2b*[+/-]

HSCs. To investigate this, proliferation was assessed in CD41:GFP^{low} purified HSCs (Figure 4B).
Although numerically unaffected by *gata2b* heterozygosity (Figure 4C), cell cycle phase
distribution was markedly different in *gata2b*$^{+/-}$ HSCs compared to WT (Figure 4D and E),
with increased G1 phase and decreased G2-M phase. As stem cells are known to have a
short G1 phase to reduce cellular differentiation potential, this may already be an indication
that these HSCs lose self-renewal capacity and differentiate, which is halted in the next
differentiation stages[36].
263
Epigenetic dysregulation results in an erythroid differentiation block in gata2b$^{+/-}$ kidney
marrow
Next, we sought to reveal the molecular mechanism behind the erythroid dysplasia in
Gata2$^{+/-}$ KM. Compared to WT, *Gata2b*$^{+/-}$ progenitor-like clusters had significantly reduced
accessibility of GATA2 and GATA1::TAL1 motifs (Figure 4A). Despite the increase in *gata2b*
expression, this indicated an overall reduction of Gata2b function in activation of
downstream targets in the *gata2b*$^{+/-}$ zebrafish genome. Activation of GATA2 motifs were
mainly enriched in the erythroid progenitors (Figure 5A). An important downstream target
of Gata2 is Gata1, which is the master regulator of erythropoiesis. We therefore investigated
Gata1a accessibility and expression in our dataset (Figure 5B-F). The *gata1a* TSS was
generally more accessible in *gata2b*$^{+/-}$ KM (Figure 5B), but not in erythroid progenitors 1
(Figure 5C). Furthermore, we found downregulation of *gata1a* mRNA expression (Figure 5D)
and the GATA1::TAL1 motif was less accessible (Figure 5E), further supporting the reduced
functionality of Gata1a in erythroid progenitors. This could be due to reduced expression of
zfpm1 (also known as *Friend of GATA* (*FOG1*)), which can facilitate *GATA1* expression[37](Figure
5F). The different pattern of the GATA family (Increased expression of *gata2b* in HSPCs
(Figure 3I) and reduced expression of *gata1a* in erythroid progenitors (Figure 5D) suggested
that the "GATA factor switch", which is a shift from GATA2 to GATA1 occupation during
erythroid lineage differentiation[38-40], might be silenced, causing deficiencies in erythroid
lineage differentiation.
Considering that the origin of dysplastic cells were readily recognized as myeloid or
erythroid in KM smears, we hypothesized that dysplastic cells occupy a proportion of
erythroid or myeloid clusters. For a more precise examination of small populations along the
myeloid and erythroid differentiation lineage, we sub-clustered "HSPCs", "proliferative
progenitors", "progenitors", "myeloid progenitors 1" and "erythroid progenitors 2" clusters

and identified 5 significantly overrepresented sub-clusters in *gata2b*[+/-] KM (Figure 5G-I,
supplemental Figure 6 A-H). These clusters are only present in *gata2b*[+/-] KM and could
therefore be the dysplastic cells. These sub-clusters represented 5.6% of the total sequenced
gata2b[+/-] cells, comparable to the proportion of dysplastic cells observed in *gata2b*[+/-] KM.
Differentially expressed genes (DEGs) analysis of overrepresented sub-clusters in *gata2b*[+/-]
KM compared to the rest of that cluster showed downregulation of tubulin transcripts (not
shown), suggesting a loss of cytoskeletal structure, a characteristic of dysplasia. Besides, TF
motif score analysis with snATAC-seq indicated a general reduction for lineage-specific
determination and differentiation motifs in *gata2b*[+/-] KM (supplemental Figure 5B-G),
suggesting that *gata2b*[+/-] cell maturation could be impaired resulting in a differentiation
block, as this is often the case in dysplasia[41].

To identify the differentiation block mechanisms in *gata2b*[+/-] KM, we compared the
DEGs and their function in the "erythroid progenitor 2" cluster. The result showed that
gata2b[+/-] "erythroid progenitor 2" cluster cells had upregulated signatures of DNA
replication together with the downregulation of transcripts necessary for mitosis (Figure 5J),
which can explain the origin of multi-lobulated nuclei and other nuclear abnormalities
observed in *gata2b*[+/-] dysplastic cells[42,43]. Pseudo-time trajectory inference showed a clearly
diminished differentiation in *gata2b*[+/-] "erythroid progenitor 2" cluster with a lower
pseudo-time score (Figure 5K and L). Pseudo-time dynamic gene expression level also
indicated the erythropoiesis block in *gata2b*[+/-] "erythroid progenitor 2" cluster with
persistent high open chromatin, but lower expression of *gata1a*, which should gradually
decrease toward terminal erythroid maturation, and the increasing level of *mki67*, which
should be decreasing during maturation, indicative of incomplete differentiation (comparing
Figure 5M-O to Figure 5P-R). This could be due to a diminished open chromatin structure at
the locus leading to a reduction in Gata2b binding sites to drive proper *Gata1a* expression,
to support the erythroid differentiation process. Furthermore, TF motif score analysis of the
Erythroid progenitor 2 cluster revealed a significant depletion for GATA family motifs, but an
enrichment for Ets and Etv family of transcription factor motifs in *gata2b*[+/-] Erythroid
progenitors 2 cells (Figure 5S), suggesting an epigenetic dysregulation related to decreased
maturation of erythroblasts resulting from Gata2b haploinsufficiency.

319
320

Discussion

In humans, a balanced *GATA2* expression is essential for proper hematopoiesis.
Consequently, more than 80% of *GATA2* mutation carriers progress to hematological
malignancy by 40 years of age[4]. While the clinical consequences of *GATA2* mutations became
obvious over the last decades, the regulation of *GATA2* activity and its contribution to
human bone marrow failure syndromes and progression to hematological malignancy
remain incompletely understood.

Here, we used transgenic reporters, morphological phenotyping, and single-cell omics
sequencing to analyze *gata2b* heterozygous zebrafish. We showed that, while major
differentiation lineages remain intact, *gata2b* heterozygosity causes dysplasia in erythroid
and myeloid progenitors of zebrafish KM. snATAC-seq revealed an auto-regulatory feedback
mechanism for Gata2b haploinsufficiency, characterized by *gata2b* chromatin
over-accessibility and stronger co-accessibility of *gata2b* enhancers and TSS, indicating
epigenetic dysregulation in hematopoiesis in the zebrafish model. scRNA-seq analysis did
not identify a single population of dysplastic cells. Interestingly, a single erythroid progenitor
cluster was identified using snATAC-seq, and could indicate a specific dysplastic erythroid
population. The difference in recognition of the dysplastic cluster between scRNAseq and
snATACseq may point to the underlying mechanism of the dysplasia where we have evidence
that the epigenetic compensatory mechanism of maintaining normal levels of Gata2b may
cause the dysplastic features during lineage differentiation, particularly in the GATA2 to
GATA1 switch and therefore affects erythroid lineage differentiation most. This also
underscores the importance of epigenetic analysis in other types of MDS and the fact that
many epigenetic regulators are mutated in MDS and AML[46]. In *gata2b*[+/-] erythroid
progenitors we found increased proliferative signatures together with decreased expression
of genes related to mitosis indicating an impaired cell cycle progression. Furthermore, we
found the depletion of GATA family factors indicating a failure of the "GATA factor switch"
that is indispensable for erythroid lineage differentiation[38-40]. We propose that these
alterations could play a role in the onset of dysplasia found in *gata2b*[+/-] zebrafish.

Interestingly, whereas *gata2b*[-/-] zebrafish display abrogated myeloid lineage
differentiation and a bias toward lymphoid differentiation [16,17], *gata2b*[+/-] did not simply
result in an intermediate phenotype between WT and *gata2b*[-/-]. Instead, Gata2b
haploinsufficiency uniquely caused dysplasia, not observed in *gata2b*[-/-], potentially caused
by compensatory mechanisms in haploinsufficiency that are otherwise overcome in
gata2b[-/-] zebrafish[47]. This may also be the reason that missense mutations have a different

molecular mechanism, because these alleles are not prone to mRNA degradation and may thus not be compensated by epigenetic deregulation. The differences in the homozygous and heterozygous Gata2b knockout phenotype support a role for gene dosage underlying the GATA2 deficiency phenotype, possibly explaining the phenotypic heterogeneity between patients. Since both erythroid and myeloid dysplasia can be observed in GATA2 patients[4], we propose that the presence of dysplastic cells in *gata2b*[+/-] resembles the clinical phenotypes associated with GATA2 heterozygosity. In the future, the isolation of single dysplastic cells could help us to further explore the effect of Gata2 haploinsufficiency in malignant transformation. Nevertheless, it remains to be established how *gata2b*[+/-] HSPCs would respond to secondary insults such as infections or severe bleeding.

In conclusion, while the major lineage differentiation remains intact, *gata2b*[+/-] zebrafish possess a stressed proliferative HSPC compartment which leads to the generation of erythroid and myeloid dysplastic cells. Taken together, our model provides insights into the consequences of Gata2b dosage, illustrated the alteration of the microenvironment, and reveal how changes in epigenetics affect the outcome in lineage differentiation after Gata2b haploinsufficiency in zebrafish.

Acknowledgements

We thank Dr Monteiro (University of Birmingham) for careful reading of the manuscript. We thank the Experimental Animal Facility of Erasmus MC for animal husbandry and the Erasmus Optical Imaging Center for confocal microscopy services. This research is supported by the European Hematology Association (junior and senior non clinical research fellowship) (EdP), the Dutch Cancer Foundation KWF/Alpe d'HuZes (SK10321)(EdP), the Daniel den Hoed Foundation for support of the Cancer Genome Editing Center (IT) and the Josephine Nefkens Foundation for purchase of the Chromium 10x (IT).

Authorship contributions

EdP and EG conceived the study; EG, CK, HdL, JZ, DB, and EB performed experiments; EG, WZ, CK, RH, KG and EdP analysed results; IT provided resources and EG, WZ, CK and EdP wrote the manuscript and IT revised the manuscript.

Disclosures

The authors declare no conflicts of interests

References

1. Gao X, Johnson KD, Chang YI, et al. Gata2 cis-element is required for
hematopoietic stem cell generation in the mammalian embryo. *J Exp Med*. Dec 16
2013;210(13):2833-42. doi:jem.20130733 [pii]
10.1084/jem.20130733

2. de Pater E, Kaimakis P, Vink CS, et al. Gata2 is required for HSC generation and
survival. *J Exp Med*. Dec 16 2013;210(13):2843-50. doi:jem.20130751 [pii]
10.1084/jem.20130751

3. Tsai FY, Keller G, Kuo FC, et al. An early haematopoietic defect in mice lacking
the transcription factor GATA-2. *Nature*. Sep 15 1994;371(6494):221-6.
doi:10.1038/371221a0

4. Donadieu J, Lamant M, Fieschi C, et al. Natural history of GATA2 deficiency in a
survey of 79 French and Belgian patients. *Haematologica*. Aug
2018;103(8):1278-1287. doi:haematol.2017.181909 [pii]
10.3324/haematol.2017.181909

5. Spinner MA, Sanchez LA, Hsu AP, et al. GATA2 deficiency: a protean disorder of
hematopoiesis, lymphatics, and immunity. *Blood*. Feb 6 2014;123(6):809-21.
doi:S0006-4971(20)36040-7 [pii]
10.1182/blood-2013-07-515528

6. Ostergaard P, Simpson MA, Connell FC, et al. Mutations in GATA2 cause
primary lymphedema associated with a predisposition to acute myeloid leukemia
(Emberger syndrome). *Nat Genet*. Sep 4 2011;43(10):929-31. doi:ng.923 [pii]
10.1038/ng.923

7. Hsu AP, Sampaio EP, Khan J, et al. Mutations in GATA2 are associated with the
autosomal dominant and sporadic monocytopenia and mycobacterial infection
(MonoMAC) syndrome. *Blood*. Sep 8 2011;118(10):2653-5.
doi:blood-2011-05-356352 [pii]
10.1182/blood-2011-05-356352

8. Dickinson RE, Griffin H, Bigley V, et al. Exome sequencing identifies GATA-2
mutation as the cause of dendritic cell, monocyte, B and NK lymphoid deficiency.

Blood. Sep 8 2011;118(10):2656-8. doi:S0006-4971(20)40687-1 [pii]

10.1182/blood-2011-06-360313

9. Hahn CN, Chong CE, Carmichael CL, et al. Heritable GATA2 mutations

associated with familial myelodysplastic syndrome and acute myeloid leukemia. *Nat*

Genet. Sep 4 2011;43(10):1012-7. doi:ng.913 [pii]

10.1038/ng.913

10. Mutsaers PG, van de Loosdrecht AA, Tawana K, Bodor C, Fitzgibbon J, Menko

FH. Highly variable clinical manifestations in a large family with a novel GATA2

mutation. *Leukemia*. Nov 2013;27(11):2247-8. doi:leu2013105 [pii]

10.1038/leu.2013.105

11. Wang X, Muramatsu H, Okuno Y, et al. GATA2 and secondary mutations in

familial myelodysplastic syndromes and pediatric myeloid malignancies.

Haematologica. Oct 2015;100(10):e398-401. doi:haematol.2015.127092 [pii]

10.3324/haematol.2015.127092

12. Rodrigues NP, Janzen V, Forkert R, et al. Haploinsufficiency of GATA-2 perturbs

adult hematopoietic stem-cell homeostasis. *Blood*. Jul 15 2005;106(2):477-84.

doi:2004-08-2989 [pii]

10.1182/blood-2004-08-2989

13. Ling KW, Ottersbach K, van Hamburg JP, et al. GATA-2 plays two functionally

distinct roles during the ontogeny of hematopoietic stem cells. *J Exp Med*. Oct 4

2004;200(7):871-82. doi:jem.20031556 [pii]

10.1084/jem.20031556

14. Butko E, Distel M, Pouget C, et al. Gata2b is a restricted early regulator of

hemogenic endothelium in the zebrafish embryo. *Development*. Mar 15

2015;142(6):1050-61. doi:10.1242/dev.119180

15. Dobrzycki T, Mahony CB, Krecsmarik M, et al. Deletion of a conserved Gata2

enhancer impairs haemogenic endothelium programming and adult Zebrafish

haematopoiesis. *Commun Biol*. Feb 13 2020;3(1):71.

doi:10.1038/s42003-020-0798-3

10.1038/s42003-020-0798-3 [pii]

16. Gioacchino E, Koyunlar C, Zink J, et al. Essential role for Gata2 in modulating

lineage output from hematopoietic stem cells in zebrafish. *Blood Adv*. Jul 13

2021;5(13):2687-2700. doi:S2473-9529(21)00348-7 [pii]

10.1182/bloodadvances.2020002993

17. Avagyan S, Weber MC, Ma S, et al. Single-cell ATAC-seq reveals
GATA2-dependent priming defect in myeloid and a maturation bottleneck in lymphoid
lineages. *Blood Adv.* Jul 13 2021;5(13):2673-2686. doi:S2473-9529(21)00347-5 [pii]
10.1182/bloodadvances.2020002992

18. Ma D, Zhang J, Lin HF, Italiano J, Handin RI. The identification and
characterization of zebrafish hematopoietic stem cells. *Blood.* Jul 14
2011;118(2):289-97. doi:blood-2010-12-327403 [pii]
10.1182/blood-2010-12-327403

19. Tamplin OJ, Durand EM, Carr LA, et al. Hematopoietic stem cell arrival triggers
dynamic remodeling of the perivascular niche. *Cell.* Jan 15 2015;160(1-2):241-52.
doi:10.1016/j.cell.2014.12.032

20. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal
single-cell data. *Cell.* Jun 24 2021;184(13):3573-3587 e29.
doi:S0092-8674(21)00583-3 [pii]
10.1016/j.cell.2021.04.048

21. Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state
analysis with Signac. *Nat Methods.* Nov 2021;18(11):1333-1341.
doi:10.1038/s41592-021-01282-5 [pii]
10.1038/s41592-021-01282-5

22. Cao J, Spielmann M, Qiu X, et al. The single-cell transcriptional landscape of
mammalian organogenesis. *Nature.* Feb 2019;566(7745):496-502.
doi:10.1038/s41586-019-0969-x [pii]
10.1038/s41586-019-0969-x

23. Boatman S, Barrett F, Satishchandran S, Jing L, Shestopalov I, Zon LI. Assaying
hematopoiesis using zebrafish. *Blood Cells Mol Dis.* Dec 2013;51(4):271-6.
doi:S1079-9796(13)00160-5 [pii]
10.1016/j.bcmd.2013.07.009

24. Traver D, Paw BH, Poss KD, Penberthy WT, Lin S, Zon LI. Transplantation and in
vivo imaging of multilineage engraftment in zebrafish bloodless mutants. *Nat Immunol.*
Dec 2003;4(12):1238-46. doi:10.1038/ni1007
ni1007 [pii]

25. Langenau DM, Ferrando AA, Traver D, et al. In vivo tracking of T cell

development, ablation, and engraftment in transgenic zebrafish. *Proc Natl Acad Sci U*

S A. May 11 2004;101(19):7369-74. doi:10.1073/pnas.0402248101

0402248101 [pii]

26. Renshaw SA, Loynes CA, Trushell DM, Elworthy S, Ingham PW, Whyte MK. A

transgenic zebrafish model of neutrophilic inflammation. *Blood*. Dec 15

2006;108(13):3976-8. doi:blood-2006-05-024075 [pii]

10.1182/blood-2006-05-024075

27. Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. mpeg1 promoter

transgenes direct macrophage-lineage expression in zebrafish. *Blood*. Jan 27

2011;117(4):e49-56. doi:blood-2010-10-314120 [pii]

10.1182/blood-2010-10-314120

28. Page DM, Wittamer V, Bertrand JY, et al. An evolutionarily conserved program of

B-cell development and activation in zebrafish. *Blood*. Aug 22 2013;122(8):e1-11.

doi:blood-2012-12-471029 [pii]

10.1182/blood-2012-12-471029

29. Carmona SJ, Teichmann SA, Ferreira L, et al. Single-cell transcriptome analysis

of fish immune cells provides insight into the evolution of vertebrate immune cell

types. *Genome Res*. Mar 2017;27(3):451-461. doi:10.1101/gr.207704.116

30. Danilova N, Bussmann J, Jekosch K, Steiner LA. The immunoglobulin

heavy-chain locus in zebrafish: identification and expression of a previously unknown

isotype, immunoglobulin Z. *Nat Immunol*. Mar 2005;6(3):295-302. doi:ni1166 [pii]

10.1038/ni1166

31. Kortum AN, Rodriguez-Nunez I, Yang J, et al. Differential expression and ligand

binding indicate alternative functions for zebrafish polymeric immunoglobulin receptor

(pIgR) and a family of pIgR-like (PIGRL) proteins. *Immunogenetics*. Apr

2014;66(4):267-79. doi:10.1007/s00251-014-0759-4

32. Macaulay IC, Svensson V, Labalette C, et al. Single-Cell RNA-Sequencing

Reveals a Continuous Spectrum of Differentiation in Hematopoietic Cells. *Cell Rep*.

Feb 2 2016;14(4):966-977. doi:S2211-1247(15)01538-7 [pii]

10.1016/j.celrep.2015.12.082

33. Tang Q, Iyer S, Lobbardi R, et al. Dissecting hematopoietic and renal cell

heterogeneity in adult zebrafish at single-cell resolution using RNA sequencing. *J Exp*

Med. Oct 2 2017;214(10):2875-2887. doi:10.1084/jem.20170976

34. Athanasiadis EI, Botthof JG, Andres H, Ferreira L, Lio P, Cvejic A. Single-cell

RNA-sequencing uncovers transcriptional states and fate decisions in

haematopoiesis. *Nat Commun.* Dec 11 2017;8(1):2045.

doi:10.1038/s41467-017-02305-6 [pii]

2305 [pii]

10.1038/s41467-017-02305-6

35. Wang J, Sun Q, Morita Y, et al. A differentiation checkpoint limits hematopoietic

stem cell self-renewal in response to DNA damage. *Cell.* Mar 2 2012;148(5):1001-14.

doi:S0092-8674(12)00145-6 [pii]

10.1016/j.cell.2012.01.040

36. Pauklin S, Vallier L. The cell-cycle state of stem cells determines cell fate

propensity. *Cell.* Sep 26 2013;155(1):135-47. doi:S0092-8674(13)01025-8 [pii]

10.1016/j.cell.2013.08.031

37. Cantor AB, Orkin SH. Coregulation of GATA factors by the Friend of GATA (FOG)

family of multitype zinc finger proteins. *Semin Cell Dev Biol.* Feb 2005;16(1):117-28.

doi:S1084-9521(04)00102-8 [pii]

10.1016/j.semcdb.2004.10.006

38. Grass JA, Boyer ME, Pal S, Wu J, Weiss MJ, Bresnick EH. GATA-1-dependent

transcriptional repression of GATA-2 via disruption of positive autoregulation and

domain-wide chromatin remodeling. *Proc Natl Acad Sci U S A.* Jul 22

2003;100(15):8811-6. doi:10.1073/pnas.1432147100

1432147100 [pii]

39. Pal S, Cantor AB, Johnson KD, et al. Coregulator-dependent facilitation of

chromatin occupancy by GATA-1. *Proc Natl Acad Sci U S A.* Jan 27

2004;101(4):980-5. doi:10.1073/pnas.0307612100

0307612100 [pii]

40. Bresnick EH, Lee HY, Fujiwara T, Johnson KD, Keles S. GATA switches as

developmental drivers. *J Biol Chem.* Oct 8 2010;285(41):31087-93.

doi:S0021-9258(19)88840-3 [pii]

10.1074/jbc.R110.159079

41. Monika Belickova M, Merkerova MD, Votavova H, et al. Up-regulation of

ribosomal genes is associated with a poor response to azacitidine in myelodysplasia

and related neoplasms. *Int J Hematol.* Nov 2016;104(5):566-573.

doi:10.1007/s12185-016-2058-3

10.1007/s12185-016-2058-3 [pii]

42. Nakayama Y, Yamaguchi N. Multi-lobulation of the nucleus in prolonged S phase

by nuclear expression of Chk tyrosine kinase. *Exp Cell Res*. Apr 1

2005;304(2):570-81. doi:S0014-4827(04)00703-7 [pii]

10.1016/j.yexcr.2004.11.027

43. Ullah Z, Lee CY, Depamphilis ML. Cip/Kip cyclin-dependent protein kinase

inhibitors and the road to polyploidy. *Cell Div*. Jun 2 2009;4:10. doi:1747-1028-4-10

[pii]

10.1186/1747-1028-4-10

44. Do C, Xing Z, Yu YE, Tycko B. Trans-acting epigenetic effects of chromosomal

aneuploidies: lessons from Down syndrome and mouse models. *Epigenomics*. Feb

2017;9(2):189-207. doi:10.2217/epi-2016-0138

45. Brault V, Duchon A, Romestaing C, et al. Opposite phenotypes of muscle

strength and locomotor function in mouse models of partial trisomy and monosomy

21 for the proximal Hspa13-App region. *PLoS Genet*. Mar 2015;11(3):e1005062.

doi:PGENETICS-D-14-01836 [pii]

10.1371/journal.pgen.1005062

46. Haferlach T, Nagata Y, Grossmann V, et al. Landscape of genetic lesions in 944

patients with myelodysplastic syndromes. *Leukemia*. Feb 2014;28(2):241-7.

doi:leu2013336 [pii]

10.1038/leu.2013.336

47. Rossi A, Kontarakis Z, Gerri C, et al. Genetic compensation induced by

deleterious mutations but not gene knockdowns. *Nature*. Aug 13

2015;524(7564):230-3. doi:nature14580 [pii]

10.1038/nature14580

Figure legends

Figure. 1: Gata2b^{+/-} kidney marrow shows erythroid and myeloid dysplasia

(A) Representative pictures of kidney marrow smears after May–Grünwald-Giemsa staining

of WT KM smears (panel 1 and 2) and *gata2b*$^{+/-}$ KM smears (panel 3-10). 3) Binucleated

erythroblast; 4) Blebbing in cytoplasm of proerythroblast; 5) Binucleated promyelocyte; 6)

Lobed nucleus and micronucleus in erythroblast; 7) Multinucleated promyelocyte; 8)

Irregular cytoplasm in erythroid precursor; 9) Lobed nucleus in erythrocytes of sorted cell

after cytospin; 10) Blebbing in cytoplasm of blast of sorted cell after cytospin.

(B) Frequency of dysplastic cells of the erythroid, myeloid and lymphoid lineage in KM

smears of WT (n=8) and *gata2b*$^{+/-}$ (n=8) zebrafish.

Figure. 2: Single cell transcriptome and chromatin accessibility analysis reveals a

differentiation block in gata2b$^{+/-}$ *KM*

(A) The flow cytometry-based sorting strategy for kidney marrow cells for scRNA-seq and

snATAC-seq analysis.

(B) UMAP analysis of scRNA-seq (top panels) and snATAC-seq (bottom panels) showing all

cell types, which was 19 cell types in scRNA-seq and 16 cell types in snATAC-seq. UMAP atlas

was separated between WT and *gata2b*$^{+/-}$ cells.

(C) Feature plot showing the scRNA-seq expression of marker gene *hemoglobin beta adult 1*

(*hbba1*) in the erythroid lineage, *granulin1* (*grn1*) in the myeloid lineage, *cluster of*

differentiation 37 (*CD37*) in the B cell lineage, and *GATA binding protein 2b* (*gata2b*) in

HSPCs to show cluster annotation.

(D) Feature plot showing the snATAC-seq gene activity of *hbba1* in the erythroid lineage,

grn1 in the myeloid lineage, *CD37* in the B lineage and *gata2b* in HSPCs to show cluster

annotation.

(E) Coverage plot illustrating the chromatin accessibility peaks of *hbba1* in the erythroid

lineage, *grn1* in the myeloid lineage, *CD37* in the B lineage, and *gata2b* in hematopoietic

stem and progenitor cells (HSPCs) to show cluster annotation.

(F) Cluster distribution quantification of scRNA-seq (left panel) and snATAC-seq (right panel)

between WT and *gata2b*[+/-] cells, with significantly differentially distributed clusters

represented in pink. A statistical significance threshold of FDR < 0.05 and log2 fold change >

1 were applied to determine the significance of the observed differences.

(G) UMAP plot colored by pseudotime scores representing cell developmental trajectories

split by WT and *gata2b*[+/-] cells. Lower pseudotime scores correspond to the root of

trajectories, while higher pseudotime scores represent the terminal differentiation

trajectories. The lines plotted on the UMAP atlas illustrate the differentiation paths of cells,

while the circled numbers indicate the location and order of the branch points along the cell

differentiation trajectories.

Figure. 3: Differences in chromatin accessibility of Gata2b result in upregulation of Gata2b

expression in gata2b*[+/-] *HSPCs

(A) Coverage plots showing significant differences of *gata2b* chromatin accessibility

between WT and *gata2b*[+/-] cells. The normalized peak signal range is 0 to 40. Highlighted

genomic regions within grey blocks indicate the location of two enhancers, enhancer 1 (E1)

and enhancer 2 (E2). The gene block depicts gene bodies contained within the plotted

genomic regions, *gata2b* is depicted in blue. The link between the blocks illustrate predicted

cis-regulatory interactions between the *gata2b* gene body and its upstream region, based

on examination of genome co-accessibility. Link colors correspond to predicted

co-accessibility scores, with yellow indicating stronger predicted interactions.

(B-G) UMAP feature plots depict the open chromatin of Enhancer 2 (E2; B and C), Enhancer

1 (E1; D and E), and transcription start site (TSS; F and G) of *gata2b*. UMAP atlas were

separated by WT (left panel) and *gata2b*$^{+/-}$ cells (right panel).

(H) Open chromatin of the *gata2b* locus per HSPC cluster cell comparing WT and *gata2b*$^{+/-}$

HSPCs.

(I) Violin plot displaying the expression level of *gata2b* in HSPCs in WT (pink) and *gata2b*$^{+/-}$

(turqois).

Figure 4. Functional consequences in HSPCs upon Gata2b overexpression

(A) The dot plot displays the log2 fold change values of differentially activated transcription

factor (TF) motifs in progenitor clusters, including "HSPCs", "progenitors", and "HSPCs and

thrombocytes", between WT and *gata2b*$^{+/-}$ cells. The TF motifs are ranked in descending

order based on log2 fold change values. Dotted lines indicate the threshold for significance,

with absolute log2 fold change values > 0.58. The blue and red colors represent TF motifs

that are statistically significant (P value < 0.05) with absolute log2 fold change values > 0.58.

(B) Representative figure of identification of CD41:GFP^{int} population (in green) and

distribution in the FSC-A SSC-A kidney marrow population.

(C) Quantification of the percentage of CD41:GFP^{int} cells in WT (n=12) and *gata2b$^{+/-}$* (n=12)

KM single live cells.

(D) Representative flow cytometry plots of cycle analysis by anti-Ki67 and DAPI staining in

WT and *gata2b$^{+/-}$* CD41:GFP^{int} KM.

(E) Quantification of the percentages of WT (n=3) and *gata2b$^{+/-}$* (n=3) CD41:GFP^{int} cells in

different cell cycle stages (* P value<0.5). Data represents as mean ± standard error of the

mean.

Figure. 5: Epigenetic changes resulting in the erythroid differentiation block in gata2$^{+/-}$ KM

(A) UMAP feature plots depicting open chromatin of GATA2 motif sequence indicating

activity of GATA2. UMAP atlas were separated by WT (left panel) and *gata2b$^{+/-}$* cells (right

panel). The top left corner of UMAP plots display the TF sequence logo corresponding to the

analyzed motif, generated by the function *MotifPlot*.

(B) Coverage plots showing differences of *gata1a* chromatin accessibility between WT and

gata2b$^{+/-}$ cells.

(C) Percentage of cells derived from WT and *gata2b$^{+/-}$* KM with accessibility of the *gata1a*

locus across clusters.

(D) Percentage of cells derived from WT and *gata2b$^{+/-}$* KM expressing *gata1a* across clusters.

(E) UMAP plots depict the activity of combined motif GATA1::TAL1. UMAP atlas were

separated by WT (left panel) and *gata2b$^{+/-}$* cells (right panel). The top left corner of the

UMAP plots display the TF sequence logo corresponding to the analyzed motif, generated by

the function *MotifPlot*.

(F) Feature plot showing the *zfpm1* (FOG1) expression in WT and *gata2b*$^{+/-}$ KM.

(G-H) UMAP plots of the subclustering of the erythroid progenitors 2 cluster in scRNA-seq

data. UMAP atlas were separated by WT(G) and *gata2b*$^{+/-}$ cells (H)

(I) Quantification of distribution between WT and *gata2b*$^{+/-}$ erythroid progenitor 2

subclusters. Significantly differentially distributed clusters in pink. Dotted box indicates

overrepresented subcluster in *gata2b*$^{+/-}$ cells.

(J) STRING network of upregulated and downregulated transcripts in *gata2b*$^{+/-}$ "erythroid

progenitors 2". Only networks with more than 2 interactions were represented. Highlighted

in red DNA replication genes from KEGG pathways and highlighted in blue mitotic

cytokinesis genes from biological processes (Gene Ontology).

(K-L) Lineage trajectory analysis of WT (K) and *gata2b*$^{+/-}$ (L) erythroid progenitor 2

subclusters. Note the difference in pseudo-time scale between the WT and *gata2b*$^{+/-}$

trajectories.

(M-R) WT (M-O) and *gata2b*$^{+/-}$ (P-R) pseudo-time expression of individual genes *hemoglobin*

beta adult 1 (*hbba1*) (M and P), *GATA binding protein 1 alpha (gata1a)* (N and Q)*,* and *mki67*

(O and R). Fold change > 0.05 & adjusted P value < 0.05. FDR=False discovery rate. FD=Fold

difference.

(S) Dot plot displaying log2 fold change values of differentially activated TF motifs in the

erythroid progenitor 2 cluster between WT and *gata2b*$^{+/-}$ cells in snATAC-seq data. The TF

motifs are ranked in descending order based on log2 fold change values. Dotted lines

indicate the threshold (absolute log2 fold change < -0.58 and >0.58) for significance, The

blue (downregulated) and red (upregulated) colors represent TFs that are statistically

significant (P value < 0.05).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

Supplementary figure 4

Supplementary Figure 5

Supplementary figure 6