VOCAL TRACT ALLOMETRY IN A MAMMALIAN VOCAL LEARNER

Koen de Reus[1,2,3*#], Daryll Carlson[3,4*], Alice Lowry[3,5], Stephanie Gross[6], Maxime Garcia[7,8], Ana Rubio-Garcia[3], Anna Salazar-Casals[3], Andrea Ravignani[1,3,9#]

[1]Comparative Bioacoustics Group, Max Planck Institute for Psycholinguistics, Nijmegen, the Netherlands
[2]Artificial Intelligence Lab, Vrije Universiteit Brussel, Brussels, Belgium
[3]Research Department, Sealcentre Pieterburen, Pieterburen, the Netherlands
[4]Department of Earth System Science, Stanford University, Stanford, USA
[5]School of Environmental Sciences, University of Liverpool, Liverpool, United Kingdom
[6]Institute for Terrestrial and Aquatic Wildlife Research (ITAW), University of Veterinary Medicine Hannover, Foundation, Büsum, Germany
[7]Animal Behaviour, Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zürich, Switzerland
[8]Center for the Interdisciplinary Study of Language Evolution, University of Zurich, Zürich, Switzerland
[9]Center for Music in the Brain, Department of Clinical Medicine, Aarhus University & The Royal Academy of Music Aarhus/Aalborg, Denmark

* These authors share joint first authorship.
# Corresponding Authors: koen.de.reus@mpi.nl; andrea.ravignani@mpi.nl

Abstract

Acoustic allometry occurs when features of animal vocalisations can be predicted from body size measurements. Despite this being considered the norm, allometry sometimes breaks, resulting in species sounding smaller or larger than expected for their size. A recent hypothesis suggests that allometry-breaking mammals cluster into two groups: those with anatomical adaptations to their vocal tracts and those capable of learning new sounds (vocal learners). Here we test which mechanism is used to escape from acoustic allometry by probing vocal tract allometry in a proven mammalian vocal learner, the harbour seal (*Phoca vitulina*). We test whether vocal tract structures and body size scale allometrically in 68 young individuals. We find that both body length and body mass accurately predict vocal tract length and one tracheal dimension. Independently, body length predicts vocal fold length while body mass predicts a second tracheal dimension. All vocal tract measures are larger in weaners than in pups and some structures are sexually dimorphic within age classes. We conclude that harbour seals do comply with anatomical allometric constraints. However, allometry between body size and vocal fold length seems to emerge after puppyhood, suggesting that ontogeny may modulate the anatomy-learning distinction previously hypothesised as clear-cut. We suggest that seals, like other species producing signals that deviate from those expected from their vocal tract dimensions, may break allometry without morphological adaptations. In seals, and potentially other vocal learning

40 mammals, advanced neural control over vocal organs may be the main mechanism for breaking

41 acoustic allometry.

43 **Key words:** pinniped, harbour seal, vocal anatomy, acoustic allometry, trachea, larynx, vocal

44 tract

Introduction

In many species, acoustic signals help mediate social interactions such as competition for mates and territory, and parent-offspring recognition (Bradbury and Vehrencamp, 1998; Martin et al., 2017). Signals can encode information about the caller's biology which can be readily deciphered by the receiver, including age (Reby and McComb, 2003; Charlton et al., 2009), sex (Vignal and Kelley, 2007; Charlton et al., 2009), body size (Fitch, 1997; Charlton et al., 2009; Charlton et al., 2011; Garcia et al., 2016), hormone levels (Koren and Geffen, 2009), and physical condition (Wyman et al., 2008; Koren and Geffen, 2009).

In particular, body size often shapes mammalian sounds by constraining the geometry of the vocal tract (Fitch, 2000; Reby and McComb, 2003). Acoustic cues relating to the body size of the caller can inform the receiver about the caller's competitive ability and reproductive success (Poole, 1999; Reby and McComb, 2003; Kuester et al., 1995; Pfefferle and Fischer, 2006). For example, in primates and carnivores, there is an inverse relationship between body size and call frequency parameters, where larger animals produce calls with lower frequencies, i.e., have a 'deeper' voice (Bowling et al., 2017). This relationship between acoustical call features and body size, where one accurately reflects the other, is known as 'acoustic allometry' (Taylor and Reby, 2010; Fitch, 1997). Here, signalling is considered *honest* when the acoustic parameters of observed vocalisations accurately reflect an individual's body size (Zahavi, 1997; Fitch and Hauser, 2003). Deviations from allometry can generate *dishonest* signals, with animals sounding unexpectedly small or large for their body size (Garcia and Ravignani, 2020). Dishonest signals may be produced when an animal shows 1) a lack of allometric scaling between their vocal tract and their body size, or 2) shows enhanced control over their vocal organs which allows them to learn new vocalisations or modify existing vocalisations: an ability known as 'vocal learning' (Janik and Slater, 1997; Lattenkamp and Vernes, 2018). Recent work indeed showed that, given a cross-species regression between sounds produced and body size, outlier species seem to cluster either well below the regression line—those with anatomical adaptations—or markedly above—the vocal learners. This led to a morphology vs. learning hypothesis (Garcia and Ravignani, 2020; Ravignani and Garcia, 2021): dishonest signals in mammals may arise either from anatomical adaptations or vocal learning capacities. This prediction has the potential to identify new vocal learners or species with unexpected vocal tract morphology. Vocal learners

should therefore be able to violate acoustic allometry while possessing a vocal tract that scales

allometrically with the rest of their body. For the first time, we test this prediction, asking

whether vocal tract allometry is present in a vocal learning species which is known to violate

acoustic allometry.

Harbour seals (*Phoca vitulina*) are vocal learners that escape acoustic allometry by producing

sounds with different frequencies than expected from their body size, allowing them to transmit

dishonest body size information. Indeed, they stand out as outliers in cross-species allometric

regressions between body mass and frequency parameters (Ravignani and Garcia, 2021; see

Figure 1). Moreover, previous studies have shown that harbour seals can actively modulate the

call frequencies they produce based on auditory experience. In one special case, a human-raised

harbour seal, named Hoover, was found capable of mimicking human speech sounds (Ralls et al.,

1985). In a more recent study on harbour seal pups, young animals were found capable of

lowering their fundamental frequency (f_0) in the presence of background noise (Torres Borda et

al., 2021). Do the environmental noise conditions in which vocalisations are produced have a

stronger influence on the f_0 values than body size? To address this, we complemented acoustic

data from Torres Borda and colleagues (2021) with body mass information and reanalysed it to

show that acoustic allometric relationships do indeed break down in this species due to the large

vocal plasticity observed within individuals (see Figure 2, its caption and detailed explanations in

the Supplement). These re-analyses indicate that, also within-species, individual harbour seals

may sound bigger or smaller than predicted by body size. Seals can therefore escape the

constraints of acoustic allometry, both across and within species.

Harbour seals are particularly vocal during the first few weeks following birth (Perry and

Renouf, 1988). Pups produce individually distinctive mother attraction calls (Renouf, 1984)

which vary with age, sex, and body length (Khan et al., 2006; Sauvé et al., 2015). After weaning,

however, these calls disappear entirely from their vocal repertoire, with most vocalisations

ceasing aside from occasional clicks and growls (Renouf, 1984). During adulthood, female

harbour seals remain almost entirely vocally inactive (Van Parijs and Kovacs, 2002), but males

start vocalising again, producing underwater calls during the mating season (Hanggi and

Schusterman, 1994). The large variation in vocal repertoire observed across individuals, sexes,

and age classes makes harbour seals ideal candidates to test the morphology vs. learning

hypothesis, i.e., whether a vocal learning mammal does indeed escape acoustic allometry via

learning instead of via anatomical adaptations.

Most mammalian vocalizations are described using the source-filter theory of vocal production.

Within this framework, vocal signals are initially produced by a *source* and are then *filtered* by

the vocal tract before being released into the environment (Fant, 1970). In mammals, the source

of sound production consists of the vocal folds in the larynx, and the filter is composed of the

cavities making up the upper vocal tract (Fant, 1970) (see Figure 3). The vocal folds are shelves

of tissue lying across the airway that attach ventrally and laterally to the thyroid cartilage and

dorsally to the arytenoid cartilage (see Figure 4A). When vocalising, the air expelled from the

lungs rushes between the vocal folds, causing them to vibrate and produce sound (Elemans et al.,

2015). The sound then continues to propagate along the upper vocal tract and is modified by its

geometry (i.e., filtered) before being emitted as vocalisation. The source-filter framework

highlights which vocal tract structures determine specific features present in acoustic signals.

The rate of vibration of the vocal folds determines the f_0 and the cavities of the vocal tract

determine formant frequencies (Taylor and Reby, 2010). Measurements of these vocal tract

structures can thus be used to estimate certain acoustic features of vocalisations.

Bioacoustics studies often investigate allometric relationships between acoustic signal features

and body size, without consideration of the underlying allometric scaling between body size and

vocal anatomy. Most mammals show allometry between body size and upper vocal tract length

because the upper vocal tract is constrained by bony structures (Fitch, 1997; Fitch and Giedd,

1999; Fitch, 2000; Plotsky et al., 2013; Garcia et al., 2016). However, allometry between body

size and the size of the vocal folds is less common: the larynx is surrounded by cartilaginous

structures and is thus less constrained, suggesting that vocal fold length can be decoupled from

overall body size, as found in nonhuman primates (Fitch and Hauser, 1995; Fitch, 1997; Garcia

et al., 2017). In mammals, formants, the acoustic proxy of vocal tract length, are thus often a

stronger body size predictor than f_0, the acoustic proxy of vocal fold length (Fitch, 1997; Garcia

et al., 2016).

Within the larger framework of the hypothesis above, this study tests for allometric relationships

between body size and vocal anatomy measurements in young harbour seals and tests how these

relationships vary with sex and age. Preliminary work found that harbour seals' body length

correlates with upper vocal tract length and tracheal diameter, but not with vocal fold length

(Ravignani et al., 2017). Here, we aim to expand on these findings by using a larger sample size

(353% increase), adding refined anatomical measurements, and comparing different age classes

(to test for developmental effects). Based on previous literature, we expect to find allometry

between body size and vocal tract structures that are surrounded—and hence constrained—by

bony structures, such as vocal tract length. However, based on harbour seals' vocal learning

abilities (Ralls et al., 1985; Torres Borda et al., 2021; Janik and Slater, 1997), we expect their

vocal flexibility to offer favourable grounds to find deviations from body size allometry for vocal

tract components surrounded by cartilage, such as the trachea and vocal fold length.

Materials and methods

Sample collection

Larynges were collected during necropsies on 68 young harbour seals (35 males). Fifty-two

samples came from seals that stranded on the Dutch coastline, the rest from animals found on the

German coastline (Schleswig-Holstein). Forty-two animals died in captivity at Sealcentre

Pieterburen, Pieterburen, the Netherlands, either naturally during rehabilitation despite intensive

care or by means of euthanasia due to the presence of severe clinical signs without any indication

for recovery. Euthanasia was performed by trained veterinarians, after sedation, with

pentobarbital sodium (100 mg/kg) using the method described in Greer and colleagues (2001).

The other 26 animals died in the wild, either naturally or were mercy killed by trained hunters

due to severe signs of illness (see Table 1 of the Supplement). No animals were euthanised or

mercy killed for the purpose of this study.

At the time of death, the seals studied were aged between 9 days and 12 months (median 6

months). The age of new-born individuals was estimated in number of days by expert seal

veterinarians based on the condition of the umbilical cord or the umbilicus. Older individuals

with a closed umbilicus were assigned June as their birth month, which is consistent with the

majority of harbour seal births in the Wadden Sea (Osinga et al., 2012; Reijnders et al., 2010).

Animals aged 1 month or younger were classified as pups, while those between 1 and 12 months

in age were classified as weaners, making age a binary variable. Of the 68 individuals included

in this study, 14 (8 males) were classified as pups and 54 (26 males) were classified as weaners.

A Fisher's exact test showed no significant association between age and sex ($\chi^2 = 0.765$, $p >$

.05), suggesting our sample is balanced between sexes and ages.

Sample treatment and measurements

Post-mortem examinations were performed by veterinarians who all trained at Sealcentre

Pieterburen and thereby used the same necropsy protocol (Pugliares et al., 2007). Dutch seals

were examined at Sealcentre Pieterburen and German seals were necropsied at the Institute for

Terrestrial and Aquatic Wildlife Research (ITAW), Büsum, Germany. Necropsies were

performed on either cooled or defrosted carcasses. Body mass, body length and axillary girth

were all measured prior to the start of the necropsy. Body length was measured from the tip of

the nose to the end of the tail in a non-curvilinear fashion, while the animal was in supine

position, and axillary girth was measured as the body circumference directly caudal to the front

flippers. The vocal apparatus including the upper vocal tract, the larynx, and part of the trachea

was then removed and immediately frozen at -20°C. All samples were in a similar condition (i.e.,

none presented signs of decomposition), comparable to pinniped vocal tracts in Schneider (1962)

and Ravignani and colleagues (2017).

Prior to measurement, samples were thawed in a refrigerator at 8°C and each larynx was cut

medially to produce two hemi-vocal tracts. The measurements taken on these hemi-vocal tracts

(see Figures 4B and 4C) include vocal tract length (VTL), vocal fold length (VFL), vocal fold

thickness (VFT), and tracheal measurements in the form of subglottic-tracheal dorsoventral

distances (STDVs) (called subglottic-tracheal anterior-posterior distance, STAP, in Roers et al.,

2009) using a calliper to an accuracy of ±0.01 mm. Although the vocal tract can be divided into

lower (below larynx) and upper (above larynx) sections, formants (the resonant frequencies

which often encode information about body size) are only determined by the upper vocal tract

(Lester and LaGasse, 2008). VTL will henceforth refer to the length of the upper vocal tract.

VTL was measured as the linear distance from the caudal end of the epiglottis to the rostral end

of the tongue muscle while the tongue was kept straight. VFL was measured as the distance from

the ventral attachment of the vocal fold on the thyroid cartilage to the dorsal attachment of the

vocal fold on the arytenoid cartilage. VFT was measured as the distance between the anterior and

posterior sides of the vocal folds. The first STDV was measured as the distance between the

cricothyroid ligament and the caudal end of the arytenoid. The second STDV was measured as

the diameter of the first tracheal ring. All measurements were performed independently by two

raters (KdR and AR), different from the veterinarians who performed the dissections. For both

raters, VTL, VFL, and VFT were measured 4 times, twice for each hemi-vocal tract, and STDVs

were taken twice, once for each hemi-larynx, because the start and end measuring points were

composed of cartilage (as opposed to soft tissue) and hence, we assumed that the inter-rater

reliability for STDVs would be higher than for other measurements.

Statistical analysis

Statistical analyses were performed in RStudio version 1.1.463 (R version 4.0.4). First, for both

raters, the medians for VTL, VFL and VFT were computed from all values reported for every

right and left hemi-larynx. Second, using the medians from the first step, the median values for

all measurements including STDV1 and STDV2 were computed for each larynx. This provided,

for each larynx and rater, five measurements: VTL, VFL, VFT, STDV1 and STDV2. The inter-

rater reliability for VTL, VFL, VFT, STDV1 and STDV2 was evaluated using Pearson's

correlations. Finally, the overall median values between raters were computed for all

measurements. Using these new values, Spearman's correlations between body size and vocal

anatomy measurements were then calculated (see Table 1). For each measurement, normality

was assessed using the Shapiro-Wilk test and homogeneity of variance was assessed using an F-

test. If both assumptions were met, a two-tailed independent samples t-test was computed to

check for age and sex differences. When variables were not normally distributed, but samples

had equal variance, a Mann Whitney U-test was performed to assess group differences instead.

Predictive modelling was done using generalised linear models (GLMs) with the *stats* package

(R Core Team, 2013). A series of models were produced for all anatomical measurements with

high inter-rater reliability ($r > 0.70$; Salkind, 2010, p. 627). For every response variable, the full

model included the fixed effects body length, body mass, girth, sex, age and the interaction

effects of sex with all body size predictors, age with all body size predictors and the interaction

of age and sex. The reduced model was then obtained through stepwise regression based on

Akaike Information Criterion (AIC) values. An analysis of variance (ANOVA) test was

performed to ensure that the reduced model was not performing significantly worse than the full

one. Variance inflation factors (VIF) scores were calculated for all predictors included in the

reduced models using the *car* package (Fox and Weisberg, 2019). Multicollinearity was

considered problematic for subsequent model selection if VIF scores were greater than 5

(Akinwande et al., 2015). For all selected models, deviance explained was calculated from the

model output (1 – residual deviance / null deviance) and expressed as a percentage. Plots

displaying the predicted effects of every predictor retained in the final models were produced to

assess their relationship with the response variable. Diagnostic residual plots were used to verify

the model assumptions. Independence of residuals was tested using a Durbin Watson test (Fox

and Weisberg, 2019). Normality of residuals was assessed visually by plotting model fit against

the observed data. Homoscedasticity (i.e., constant variance) of residuals was also assessed

visually using quantile-quantile plots. Finally, influential data points were assessed by

calculating Cook's distance.

248
<u>Results</u>
Inter-rater reliability for VTL, VFL, VFT, and both STDVs was evaluated using Pearson

correlations. VTL ($r = 0.94$), VFL ($r = 0.88$), STDV1 ($r = 0.97$) and STDV2 ($r = 0.93$) showed

high inter-rater reliability. VFT ($r = 0.59$) showed lower inter-rater reliability and was

consequently excluded from further analysis. All correlations were significant at $p < 0.001$.

All Spearman correlations between body size and vocal anatomy measurements showed positive

relationships and significance at the 0.05 level (see Table 1). There were high correlations

between body mass and body length ($r_s = 0.70$), and between body mass and girth ($r_s = 0.86$).

Other notable correlations included those between VTL and VFL ($r_s = 0.72$), VTL and STDV1

($r_s = 0.70$), VFL and STDV1 ($r_s = 0.82$), VFL and STDV2 ($r_s = 0.76$). Spearman correlations for

pups and weaners can be found in Table 2 of the Supplement.

All anatomical measurements were non-normally distributed but showed equal variances across

age and sex groups. A Mann Whitney U-test was used to test for group differences as only the

assumption for homogeneity of variance was satisfied. All anatomical measurements were

significantly larger in weaners than in pups ($p < 0.001$; see Table 2 and Figure 5). No significant

sex differences were found when considering pups and weaners together ($p > 0.05$). When

considering pups alone, both the normality and homoscedasticity assumptions were met. A two-

tailed independent samples t-test found significant sex differences for vocal tract length ($t = $ -

3.42, $p < 0.05$; see Figure 6A). Male pups (86.0 mm ± 2.9) had a larger mean VTL than females

(79.8 mm ± 3.7). When considering weaners alone, variables showed non-normal distribution,

but equal variances. A series of Mann Whitney U-tests found that only the first subglottic-

tracheal dorsoventral distance was significantly different across sexes ($U = 218$, $p < 0.05$; see

Figure 6B). Weaned males (25.1 mm ± 1.5) had a wider mean STDV1 compared to weaned

females (24.2 mm ± 1.2).

A reduced GLM, obtained by stepwise regression based on AIC values, was produced for every

vocal tract measurement with high inter-rater reliability, including VTL, VFL, STDV1 and

STDV2. All VIF scores were lower than 5 suggesting that multicollinearity was not problematic

in the selected models. All model assumptions were satisfied. Moreover, ANOVA testing

indicated that the reduced models did not perform significantly worse than the full models ($p > $

0.90). GLM results showed that most vocal tract dimensions were best explained by body length,

body mass, age, and sex (see Table 3). Girth was not retained as a predictor term in any of the

selected models. For each model, the predictor estimates with their confidence intervals can be

found in Table 3 of the Supplement and plots of the predicted effects can be found in Figures 1-5

of the same document. Significant interaction effects are shown in Figure 7.

Discussion

This study reports on the allometric relationships between body size and vocal tract dimensions

in harbour seals. It shows that body length accurately predicts VTL, VFL, and STDV1, and body

mass predicts VTL and both tracheal measurements (STDVs). We also find age and sex to be

important predictors for the size of vocal tract structures. This is evidenced by significant

differences in measurements between age classes and significant sexual differences within age

classes.

Previous work showed that upper vocal tract (i.e., filter) dimensions in mammals are predicted

by body size measurements (Fitch, 1997; Fitch and Giedd, 1999; Fitch, 2000; Plotsky et al.,

2013; Garcia et al., 2016, Ravignani et al., 2017) and our results provide additional evidence to

support such allometry. Although most studies have used body length as a proxy for body size,

we find that body mass can also be used to predict VTL in harbour seals. In the first years of life,

harbour seals show a linear growth rate for both body length (Haukkson, 2006) and mass

(Markussen et al., 1989), suggesting that VTL may develop in a similar fashion during this

period. Acoustic proxies for the filter could thus provide a good estimation of a harbour seal's

size. In mammals, formant frequencies and formant spacing can be predicted from VTL and vice

versa (see Reby and McComb, 2003). Other acoustic proxies include energy quartiles, the

frequency of amplitude peaks, and the ratios between these amplitudes (Sauvé et al., 2015).

These parameters also encode individual signatures, suggesting that acoustic individuality may

partially be an allometric by-product (Ravignani et al., 2017). Harbour seals have the vocal tract

predispositions to produce vocalisations that accurately reflect body size whilst also sharing

individual-specific information, suggesting that learning does not need to be invoked to explain

individuality.

Across mammals, source-related features such as f_0 can sometimes predict body size despite

showing weaker allometric scaling than filter-related features (Reby and McComb, 2003;

Charlton et al., 2011; Pfefferle et al., 2007; Charlton and Reby, 2016); it was unclear whether

this holds for harbour seals (Ravignani et al., 2017; Bowling et al., 2017). Our findings indicate

that VFL, which may be used to approximate f_0, can be predicted by body size in harbour seals.

Moreover, Sauvé and colleagues (2015) reported a decrease in f_0 with an increase in body length

of harbour seal pups. Taken together, this suggests that a harbour seal's f_0 can be predicted from

vocal anatomy. Previous evidence against allometric scaling for VFL could be explained by low

statistical power or lack of testing for age effects on vocal tract measurements (Ravignani et al.,

2017). It is indeed notable that age is included in both interactions which were retained in the

selected VFL model. Our results, including both pups and weaners, show that allometric scaling

between body size and VFL only emerges after weaning, suggesting that VFL may not be

constrained in harbour seal pups (see bottom panel of Figure 7). This begs the following

question: how would escaping acoustic allometry for source-related features be beneficial for

pups? Broadcasting honest body size information may be detrimental for harbour seal pups as

they are significantly more likely to be displaced by larger conspecifics during agonistic

interactions (Neumann, 1999). However, pups may be able to benefit from lowering the f_0

(Torres Borda et al., 2021) of their calls to create an impression of size exaggeration. On the

other hand, pups may also benefit from increasing the f_0 of their calls to create an impression of

distress to the mother (Briefer, 2012). Future playback studies could and should contrast these

hypotheses.

Several phocid species use the trachea for sound production (Bryden and Felts, 1974), but this

could be a by-product of adaptive modifications to the respiratory tract required for diving

(Kooyman and Andersen, 1969; Tyack and Miller, 2002). Our results support the correlation

between tracheal diameter and body length found by Ravignani and colleagues (2017), but also

provide evidence that tracheal dimensions can be predicted by body mass. Previous literature

found that the trachea may potentially convey body size information if its size influences

acoustic call features (Ravignani et al., 2017). In humans, a wider tracheal diameter partially

predicts turbulence (i.e., unsteady air movements) for large airflows (Van den Berg et al., 1957).

Applying the same logic to other mammals, larger seals would have wider tracheal dimensions

which, in turn, would make vocalisations noisier. This could explain, for instance, why the

harmonics-to-noise ratio decreases as harbour seals get older (de Reus, 2017). Future work on

sound production in this species could test this prediction using sound-anatomy correlations and

excised larynx set-ups. Moreover, playback experiments could test whether adding noise to

vocalisations alters interactive behaviour to determine if harmonics-to-noise ratio may encode

body size information. Understanding whether and how the trachea is involved in sound

production will thus require further research.

As expected, all anatomical measurements are larger for weaners than they are for pups. In

Ravignani and colleagues (2017), animals up to 108 days old were classified as pups. However,

in the wild, the lactation period for harbour seals ranges from 23 to 42 days, after which the pups

are weaned (Renouf, 2012). Hence, for the sake of simplicity, we consider animals up to one

month old as pups and animals older than one month as weaners. Through this categorical

classification, we were able to identify how allometric trends develop over the harbour seal's

early life. At the time of data collection, we only had very few larynges from subadults and

adults, leading us to not include these data points in our analysis to avoid potential problems

caused by small sample size. Future research including larynges from subadults and adults will

further extend our knowledge of how vocal allometry develops in harbour seals.

There were no sexual differences when considering the sample size as a whole, but significant

sexual differences existed within age classes. These differences could be attributed to differing

levels of steroid hormones acting on the laryngeal structures in males and females (Aufdemorte

et al., 1983; Sauvé et al., 2015). In some mammals, sex hormones affect the structural

development of the larynx and the viscoelastic properties of the vocal fold tissue (Fitch and

Giedd, 1999; Beckford et al., 1985). At puberty of these animals, the male larynx descends in the

vocal tract causing an elongation of the length of the upper vocal tract, allowing males to convey

an exaggerated impression of size (Fitch and Giedd, 1999; Fitch and Reby, 2001). In harbour

seal pups of similar body size, males have larger VTLs than females, suggesting that laryngeal

descent in males possibly occurs early in life. Once weaned, however, females show a clear

increase in VTL whereas it remains relatively constant in males (see top panel of Figure 7),

suggesting that VTL differences across sexes may become less pronounced over time. In

mammalian males, sex hormone action also causes a rapid increase in cartilage size leading to an

enlarged larynx and an increase in the vibrating portion of the vocal folds (Fitch and Hauser,

2003). This could explain why, in weaners, STDV1 is larger in males than in females.

Nevertheless, these findings are somewhat surprising as young harbour seals normally show little

sexual dimorphism (Le Boeuf, 1991). In particular, there is a lack of evidence for sexual

differences regarding birth mass and growth rates among harbour seal pups (Bowen et al., 1994).

In our sample, there are no significant body size differences between sexes ($p > 0.05$), however,

male pups are slightly larger than female pups in body length (M = 81.6 cm ± 4.4, F = 77.5 cm ±

4.5), which could partially explain the VTL differences observed in this age class. Male (9.8 kg ±

1.5) and female (9.8 kg ± 1.6) pups do not differ in body mass, but it is important to note that the

sampled animals were sick and/or in poor condition; hence body mass values are not

representative of healthy individuals and should be interpreted with caution. In short, based on

these observed differences in vocal anatomy across sexes, formants are expected to differ in pups

and harmonics-to-noise ratio is expected to differ in weaners. The anatomical structures that

determine these acoustic features both show strong allometric scaling, hence these parameters
may provide distinct body size cues across age classes, potentially facilitating the discrimination
of male and female conspecific calls. Future research should investigate how sex hormones
affect the elastic properties of harbour seal laryngeal tissues. Hormone levels can be measured by
taking blood samples from healthy male and female seals at different developmental stages, and
results can be combined with magnetic resonance imaging (MRI) mapping of laryngeal tissue
elasticity.
395
The high inter-rater reliability observed for VTL, VFL and both STDVs demonstrates that these
quantities can be measured and replicated easily, making them reliable landmarks for vocal tract
measurements. However, tissue properties such as the viscoelasticity of certain vocal tract
structures, like the vocal folds, are significant obstacles to getting accurate measurements.
Indeed, raters struggled to produce precise data for VFT. Future research in the field of pinniped
vocal anatomy would benefit from improved measuring techniques using 2D pictures,
radiography, MRI and computed tomography scans as this would enable more accurate
measurements for structures that are difficult to handle. Finally, future similar studies should
include measurements of another vocal tract structure: the corniculate cartilage. Although widely
absent in terrestrial carnivores, harbour seals have rather large corniculate cartilages that help
close the trachea together with the epiglottis (Adams et al., 2020). These cartilages are located
close to the vocal folds and are possibly innervated by the same nerves and controlled by the
same muscles. It may be possible that these cartilages play a role in sound production by, for
example, lowering the f_0 by adding weight to the vocal folds. Taken together, these suggestions
will provide a more precise and detailed picture of the harbour seal's vocal anatomy.
411
Observed species-specific vocalisations are determined by both the species' vocal anatomy and
their capacity for vocal learning (Garcia and Ravignani 2020; Ravignani and Garcia 2021). The
vocal anatomy generates vocal predispositions by imposing biomechanical constraints, whereas
neural processes determine the degree of control species have over their vocal organs (Garcia
and Manser, 2020). Particularly, vocal learners, like the harbour seal, are capable of actively
modulating sounds, suggesting that they are less constrained by anatomy and have a refined
capacity for vocal motor control. Unfortunately, the relative contribution of both sound

production mechanisms is unclear. Here, we test a hypothesis trying to segregate anatomical vs.

learning mechanisms (Garcia and Ravignani, 2020; Ravignani and Garcia 2021). As shown here,

by testing for allometric relationships between body size and vocal tract structures, one can start

to disentangle the respective contributions of vocal anatomy and vocal motor control in shaping

acoustic signals. We find that harbour seals are mechanistically constrained by their vocal

anatomy, and their large vocal flexibility (Ralls et al., 1985; Torres Borda et al., 2021), which

may result in the production of dishonest signals, thus points towards extensive volitional control

over their vocalisations. In brief, we provide support for the morphology vs. learning hypothesis,

showing however that this relation may be mediated by ontogeny.

In sum, we provide evidence of allometry between body size and vocal tract measurements in

harbour seals. Body length is a strong predictor for VTL, VFL, and STDV1, and body mass is a

strong predictor for VTL and both tracheal measurements (STDVs). Age and sex are also

important in predicting the dimensions of these anatomical structures. Taken together, the

combined findings demonstrate that harbour seal vocal tracts do indeed scale with body size,

although allometry between VTL and body size may only emerge after weaning. One could now

make inferences about the vocal predispositions of harbour seals (e.g., f_0, formants), based on

either their body size or the size of their vocal tract. However, to accurately predict f_0, further

studies are needed in harbour seals to determine the range of stress they apply to their vocal folds

while vocalising and to infer the tissue density of their vocal folds (Titze et al., 1989). Once such

predictions are made, comparing them to data obtained from observed natural vocalisations

would shed light on the range of vocal flexibility resulting from their extensive vocal motor

control. Although formant spacing could be predicted from vocal tract length (Titze, 1994),

bioacousticians have not yet been able to consistently extract formants from harbour seal

vocalisations, meaning that predictions cannot currently be compared to observed vocalisations.

Finally, a critical next step to directly relate acoustic features to sound production structures is to

connect harbour seals' vocal anatomy measurements to the vocalisations they produce while

alive. Integrating such results with investigations of call function will eventually inform on

which vocal structures are responsible for generating the individual- and species-specific

information encoded in harbour seals' vocalisations.

List of symbols and abbreviations

f_0 – Fundamental frequency

C – Celsius

VTL – Vocal Tract Length

VFL – Vocal Fold Length

VFT – Vocal Fold Thickness

STDV – Subglottic-Tracheal Dorsoventral distance

STAP – Subglottic-Tracheal Anterior-Posterior distance

GLM – Generalised Linear Model

AIC – Akaike Information Criterion

ANOVA – Analysis Of Variance

VIF – Variance Inflation Factor

M – Male

F – Female

mg - milligram

kg – kilogram

cm – centimetre

MRI – Magnetic Resonance Imaging

Acknowledgments

We are grateful to Letty Stupers, Rebecca Andreini, Maria Jose Robles for help in gathering and preparing samples for data collection. We thank Aline Joustra for her illustrations of seal vocal tracts. We also greatly appreciate the extensive comments provided by Taylor Hersh, Laura Verga and Yannick Jadoul. We would also like to thank all members and volunteers of Sealcentre Pieterburen for their continued support in this research.

Competing interests

The authors declare no competing interests.

Author contributions

KdR, SG, MG, ARG, ASC and AR conceived the research and designed the experiment. SG, ARG, and ASC performed the post-mortem examinations. KdR and AR performed the measurements. KdR, DC, and AL performed the statistical analysis. All authors interpreted the results, drafted the article, and helped to critically revise it. KdR and AR addressed the reviewers' comments and incorporated them in the manuscript with support from SG, MG, ARG, and ASC.

Funding

This work was supported by a Max Planck Research Group (MPRG) and a FWO Pegasus Marie Curie fellowship 12N5517N awarded to AR. KdR was supported by the FWO project 'Interactive vocal rhythms' G034720N awarded to Bart de Boer. The German Ministry of Energy, Agriculture, the Environment, Nature and Digitalization (MELUND) funded the marine mammal stranding scheme and the necropsies in Schleswig-Holstein. The Sealcentre Pieterburen funded the seal stranding scheme and the necropsies at the Sealcentre Pieterburen. MG was supported by a research grant from the Ethologische Gesellschaft e.V.

Reference list

Adams, A., Vogl, W., Dawson, C., Raverty, S., Haulena, M., & Skoretz, S. A. (2020). Laryngeal and soft palate valving in the harbour seal (Phoca vitulina). *Journal of Experimental Biology*, *223*(20).

Akinwande, M. O., Dikko, H. G., & Samson, A. (2015). Variance inflation factor: as a condition for the inclusion of suppressor variable (s) in regression analysis. *Open Journal of Statistics*, *5*(07), 754.

Aufdemorte, T. B., Sheridan, P. J., & Holt, G. R. (1983). Autoradiographic evidence of sex steroid receptors in laryngeal tissues of the baboon (Papio cynocephalus). *The Laryngoscope*, *93*(12), 1607–1611. https://doi.org/10.1288/00005537-198312000-00013

Beckford, N. S., Schaid, D., Rood, S. R., & Schanbacher, B. (1985). Androgen stimulation and laryngeal development. *Annals of Otology, Rhinology and Laryngology*, *94*(6 I), 634–640. https://doi.org/10.1177/000348948509400622

Bowen, W., Oftedal, O., Boness, D., & Iverson, S. (1994). The effect of maternal age and other factors on birth mass in the harbour seal. *Canadian Journal of Zoology*, *72*, 8–14.

Bowling, D. L., Garcia, M., Dunn, J. C., Ruprecht, R., Stewart, A., Frommolt, K. H., & Fitch, W. T. (2017). Body size and vocalization in primates and carnivores. *Scientific reports*, *7*(1), 1-11.

Bradbury, J. W., & Vehrencamp, S. L. (1998). *Principles of animal communication*. Sunderland, MA: Sinauer Associates.

Briefer, E. F. (2012). Vocal expression of emotions in mammals: mechanisms of production and evidence. *Journal of Zoology*, *288*(1), 1-20.

Bryden, M. M., & Felts, W. J. (1974). Quantitative anatomical observations on the skeletal and muscular systems of four species of Antarctic seals. *Journal of Anatomy*, *118*(Pt 3), 589–600.

Charlton, B., Zhihe, Z., & Snyder, R. (2009). The information content of giant panda, Ailuropoda melanoleuca, bleats: acoustic cues to sex, age and size. *Animal Behaviour*, *78*(4), 893–898.

Charlton, B., Ellis, W., McKinnon, A., Cowin, G., Brumm, J., Nilsson, K., & Fitch, W. (2011). Cues to body size in the formant spacing of male koala (Phascolarctos cinereus) bellows: honesty in an exaggerated trait. *Journal of Experimental Biology*, *214*(20), 3414–3422. https://doi.org/10.1242/jeb.061358

Charlton, B. D., & Reby, D. (2016). The evolution of acoustic size exaggeration in terrestrial mammals. *Nature Communications*, *7*(1), 1–8. https://doi.org/10.1038/ncomms12739

Elemans, C. P., Rasmussen, J. H., Herbst, C. T., Düring, D. N., Zollinger, S. A., Brumm, H., … & Švec, J. G. (2015). Universal mechanisms of sound production and control in birds and mammals. *Nature communications*, *6*(1), 1-13.

Fant, G. (1970). *Acoustic theory of speech production* (2nd ed.). The Hague: Mouton.

Fitch, W. T., & Hauser, M. D. (1995). Vocal production in nonhuman primates: Acoustics, physiology, and functional constraints on "honest" advertisement. *American Journal of Primatology*, *37*(3), 191–219. https://doi.org/10.1002/ajp.1350370303

Fitch, W. T. (1997). Vocal tract length and formant frequency dispersion correlate with body size in rhesus macaques. *The Journal of the Acoustical Society of America*, *102*(2), 1213–1222. https://doi.org/10.1121/1.421048

Fitch, W. T., & Giedd, J. (1999). Morphology and development of the human vocal tract: A study using magnetic resonance imaging. *The Journal of the Acoustical Society of America*, *106*(3), 1511–1522. https://doi.org/10.1121/1.427148

Fitch, W.T. (2000). The evolution of speech: a comparative review. *Trends in Cognitive Sciences*, *4*(7), 258–267.

Fitch, W.T., & Reby, D. (2001). The descended larynx is not uniquely human. *Proceedings of the Royal Society of*
London. Series B: Biological Sciences, *268*(1477), 1669–1675. https://doi.org/10.1098/rspb.2001.1704

Fitch, W. T., & Hauser, M. D. (2003). Unpacking "Honesty": Vertebrate Vocal Production and the Evolution of
Acoustic Signals. In *Acoustic Communication* (pp. 65–137). Springer-Verlag. https://doi.org/10.1007/0-
387-22762-8_3

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression*, Third edition. Sage, Thousand Oaks CA.
https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Garcia, M., Wondrak, M., Huber, L., & Fitch, W. T. (2016). Honest signaling in domestic piglets (Sus scrofa
domesticus): vocal allometry and the information content of grunt calls. *The Journal of Experimental*
Biology, *219*, 1913–1921. https://doi.org/10.1242/jeb.138255

Garcia, M., Herbst, C. T., Bowling, D. L., Dunn, J. C., & Fitch, W. T. (2017). Acoustic allometry revisited:
morphological determinants of fundamental frequency in primate vocal production. *Scientific reports*, *7*(1),
1-11.

Garcia, M., & Manser, M. (2020). Bound for Specific Sounds: Vocal Predisposition in Animal Communication.
Trends in Cognitive Sciences, *24*(9), 690-693.

Garcia, M., & Ravignani, A. (2020). Acoustic allometry and vocal learning in mammals. *Biology Letters*, *16*(7),
20200081.

Greer, L. L., Whaley, J., Rowles, T. K. (2001). Euthanasia. In: L. Dierauf & F. M. D. Gulland (Eds.), *CRC*
Handbook of Marine Mammal Medicine: Health, Disease and Rehabilitation (pp. 729-737). Boca Raton,
FL: CRC Press, Inc.

Hanggi, E. B., & Schusterman, R. J. (1994). Underwater acoustic displays and individual variation in male harbour
seals, Phoca vitulina. *Animal Behaviour*, *48*(6), 1275–1283. https://doi.org/10.1006/anbe.1994.1363

Hauksson, E. (2006). Growth and reproduction in the Icelandic common seal (Phoca vitulina L., 1758). *Marine*
Biology Research, *2*(01), 59-73.

Janik, V. M., & Slater, P. J. (1997). Vocal learning in mammals. In: P. Slater, C. Snowdon, J. Rosenblatt, & M.
Milinski (Eds.), *Advances in the Study of Behaviour*, *26*, 59-100.

Khan, C. B., Markowitz, H., & McCowan, B. (2006). Vocal development in captive harbor seal pups, Phoca vitulina
richardii: Age, sex, and individual differences. *The Journal of the Acoustical Society of America*, *120*(3),
1684–1694. https://doi.org/10.1121/1.2226530

Kooyman, G., & Andersen, H. (1969). Deep diving. *The Biology of Marine Mammals*, 65–94.

Koren, L., & Geffen, E. (2009). Androgens and social status in female rock hyraxes. *Animal Behaviour*, *77*(1), 233–
238.

Kuester, J., Paul, A., & Arnemann, J. (1995). Age-related and individual differences of reproductive success in male
and female barbary macaques (Macaca sylvanus). *Primates*, *36*(4), 461–476.

Lattenkamp, E. Z., & Vernes, S. C. (2018). Vocal learning: a language-relevant trait in need of a broad cross-species
approach. *Current opinion in behavioral sciences*, *21*, 209-215.

Le Boeuf, B. J. (1991). Pinniped mating systems on land, ice and in the water: Emphasis on the Phocidae. In *The*
Behaviour of Pinnipeds (pp. 45–65). Springer Netherlands. https://doi.org/10.1007/978-94-011-3100-1_2

Lester, B. M., & LaGasse, L. L. (2008). Crying. In *Encyclopedia of Infant and Early Childhood Development* (Vol.
1–3, pp. 332–343). Elsevier Inc. https://doi.org/10.1016/B978-012370877-9.00045-1

Markussen, N. H., Bjørge, A., & Øritsland, N. A. (1989). Growth in harbour seals (Phoca vitulina) on the
Norwegian coast. *Journal of Zoology*, *219*(3), 433-440.
602
603 Martin, K., Tucker, M. A., & Rogers, T. L. (2017). Does size matter? Examining the drivers of mammalian
604   vocalizations. *Evolution*, *71*(2), 249–260. https://doi.org/10.1111/evo.13128
605
606 Neumann, D. R. (1999). Agonistic behavior in harbor seals (Phoca vitulina) in relation to the availability of haul-out
607   space. *Marine Mammal Science*, *15*(2), 507-525.
608
609 Osinga, N., Nussbaum, S. B., Brakefield, P. M., & de Haes, H. A. U. (2012). Response of common seals (Phoca
610   vitulina) to human disturbances in the Dollard estuary of the Wadden Sea. *Mammalian Biology*, *77*(4),
611   281–287. https://doi.org/10.1016/j.mambio.2012.02.005
612
613 Perry, E. A., & Renouf, D. (1988). Further studies of the role of harbour seal (Phoca vitulina) pup vocalizations in
614   preventing separation of mother–pup pairs. *Canadian Journal of Zoology*, *66*(4), 934–938.
615   https://doi.org/10.1139/z88-138
616
617 Pfefferle, D., & Fischer, J. (2006). Sounds and size: identification of acoustic variables that reflect body size in
618   hamadryas baboons, Papio hamadryas. *Animal Behaviour*, *72*(1), 43–51.
619
620 Pfefferle, D., West, P. M., Grinnell, J., Packer, C., & Fischer, J. (2007). Do acoustic features of lion, Panthera leo,
621   roars reflect sex and male condition? *The Journal of the Acoustical Society of America*, *121*(6), 3947.
622   https://doi.org/10.1121/1.2722507
623
624 Plotsky, K., Rendall, D., Riede, T., & Chase, K. (2013). Radiographic analysis of vocal tract length and its relation
625   to overall body size in two canid species. *Journal of Zoology*, *291*(1), 76–86.
626   https://doi.org/10.1111/jzo.12048
627
628 Poole, J. H. (1999). Signals and assessment in African elephants: Evidence from playback experiments. *Animal
629   Behaviour*, *58*(1), 185–193. https://doi.org/10.1006/anbe.1999.1117
630
631 Pugliares, K. R., Bogomolni, A., Touhey, K. M., Herzig, S. M., Harry, C. T., & Moore, M. J. (2007). Marine
632   mammal necropsy: an introductory guide for stranding responders and field biologists. *Woods Hole
633   Oceanographic Institution Technical Report*, *6*, 1-133.
634
635 R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical
636   Computing, Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org/.
637
638 Ralls, K., Fiorelli, P., & Gish, S. (1985). Vocalizations and vocal mimicry in captive harbor seals, Phoca vitulina.
639   *Canadian Journal of Zoology*, *63*(5), 1050–1056. https://doi.org/10.1139/z85-157
640
641 Ravignani, A., Gross, S., Garcia, M., Rubio-Garcia, A., & De Boer, B. (2017). How small could a pup sound? The
642   physical bases of signaling body size in harbor seals. *Current Zoology*, *63*(4), 457–465.
643
644 Ravignani, A., & Garcia, M. (2021). A cross-species framework to identify vocal learning abilities in mammals.
645   *Philosophical Transactions of the Royal Society B: Biological Sciences*. 376, 20200394.
646   https://doi.org/10.1098/rstb.2020.0394
647
648 Reby, D., & McComb, K. (2003). Anatomical constraints generate honesty: Acoustic cues to age and weight in the
649   roars of red deer stags. *Animal Behaviour*, *65*(3), 519–530. https://doi.org/10.1006/anbe.2003.2078
650

651 Reijnders, P. J. H., Brasseur, S. M. J. M., & Meesters, E. H. W. G. (2010). Earlier pupping in harbour seals, Phoca
652   vitulina. *Biology Letters*, *6*(6), 854–857. https://doi.org/10.1098/rsbl.2010.0468
653
654 Renouf, D. (1984). The vocalization of the Harbour seal pup (Phoca vitulina) and its role in the maintenance of
655   contact with the mother. *Journal of Zoology*, *202*(4), 583–590. https://doi.org/10.1111/j.1469-
656   7998.1984.tb05055.x
657
658 Renouf, D. (2012). *The behaviour of pinnipeds*. Springer Science & Business Media.
659
660 de Reus, K. (2017). Talking seals: vocal development in Eastern Atlantic harbour seal pups (*Phoca vitulina
661   vitulina*). MSc Thesis, Royal Veterinary College, University of London.
662
663 Roers, F., Mürbe, D., & Sundberg, J. (2009). Predicted singers' vocal fold lengths and voice classification—a study
664   of x-ray morphological measures. *Journal of Voice*, *23*(4), 408-413.
665
666 Salkind, N. J. (2010). *Encyclopedia of research design* (Vols. 1-0). Thousands Oaks, CA: SAGE Publications.
667
668 Sauvé, C., Beauplet, G., Hammill, M., & Charrier, I. (2015). Acoustic analysis of airborne, underwater, and
669   amphibious mother attraction calls by wild harbor seal pups (Phoca vitulina). *Journal of Mammalogy*,
670   *96*(3), 591–602.
671
672 Schneider, R. (1962). Vergleichende untersuchungen am kehlkopf der robben (mammalia, carnivora, pinnipedia).
673   *Gegenbaurs Morph Jb*, *103*, 177–262.
674
675 Taylor, A. M., & Reby, D. (2010). The contribution of source-filter theory to mammal vocal communication
676   research. *Journal of Zoology*, *280*(3), 221–236.
677
678 Titze, I. R. (1994). *Principles of Voice Production.* Englewood Cliffs, New Jersey: Prentice Hall.
679
680 Titze, I. R., Luschei, E. S., & Hirano, M. (1989). Role of the thyroarytenoid muscle in regulation of fundamental
681   frequency. *Journal of Voice*, *3*(3), 213-224.
682
683 Torres Borda, L., Jadoul, Y., Rasilo, H., Casals, A. S., & Ravignani, A. (2021). Vocal plasticity in harbour seal
684   pups. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 376, 20200456.
685   https://doi.org/10.1098/rstb.2020.0456
686
687 Tyack, P., & Miller, E. (2002). Vocal anatomy, acoustic communication and echolocation. In *Marine mammal
688   biology: An evolutionary approach* (pp. 142–184). Oxford: Wiley-Blackwell.
689
690 Van den Berg, J., Zantema, J. T., & Doornenbal, P. (1957). On the Air Resistance and the Bernoulli Effect of the
691   Human Larynx. *The Journal of the Acoustical Society of America*, *29*(5), 626–631.
692   https://doi.org/10.1121/1.1908987
693
694 Van Parijs, S. M., & Kovacs, K. M. (2002). In-air and underwater vocalizations of eastern Canadian harbour seals,
695   Phoca vitulina. *Canadian Journal of Zoology*, *80*(7), 1173–1179. https://doi.org/10.1139/z02-088
696
697 Vignal, C., & Kelley, D. (2007). Significance of temporal and spectral acoustic cues for sexual recognition in
698   Xenopus laevis. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1609), 479–488.
699   https://doi.org/10.1098/rspb.2006.3744
700
701 Wyman, M., Mooring, M., McCowan, B., Penedo, M., & Hart, L. (2008). Amplitude of bison bellows reflects male
702   quality, physical condition and motivation. *Animal Behaviour*, *76*(5), 1625–1639.
703
704 Zahavi, A. (1977). The cost of honesty (further remarks on the handicap principle). *Journal of theoretical
705   Biology*, *67*(3), 603-605.
706

 Figure legends

Figure 1. PGLS regressions between frequency parameters (frequency range in the left panel,

maximum frequency in the right panel) and body mass across 164 mammalian species. All

variables are log-transformed and the figure is adapted from Ravignani and Garcia (2021). The

dotted lines represent a threshold at 2,5 standard deviations from the main regression lines used

to define outliers. Non-outlier species (which show acoustic allometry between frequency

parameters and body mass) are represented by smaller sized circles and outlier species (which

escape acoustic allometry) are represented by bigger sized circles. The two red data points,

representing harbour seals, are both outliers.

Figure 2. Lack of acoustic allometry relationships in harbour seals. Panel A shows the

correlations between median f_0 for each noise condition (silence, low and high) and body mass.

The respective correlation coefficients (τ) and associated p-values (p) for each correlation are

reported above the regression line. At first sight, the characteristic inverse relationship between

f_0 and body size may seem present, but there is some overlap in the range of f_0 values (whiskers

on the right side of the plot) produced by individuals of differing body size between noise

conditions. Non-significant p-values suggest that, at least in this sample, there is a lack of

acoustic allometry. In addition, allometry may break if calls are produced in different noise

conditions. In other words, do the environmental conditions in which vocalisations are produced

strongly affect the f_0 values, as much as or even more than body mass? We produced density

distributions (panels B-D) by computing 10,000 different combinations of randomly selected

median f_0 values (1 of the 3 median frequency values per seal) to assess if allometric

relationships hold across noise conditions. The coloured vertical lines in these plots represent the

respective values for each of the noise conditions. The median value of the distribution is

represented by black circle on the density curve. Panel B shows the density distribution of the

Kendall rank correlation coefficients. The median value lies around -0.18, pointing to a weak

negative correlation. Panel C shows the density distribution of the correlation p-values associated

with the correlations from Panel B. The median p-value is 0.38 which means that in most of the

simulated cases we would not reject the null hypothesis (i.e., the correlation is not significantly

different from 0). In fact, in only 2.2% of cases (217 out of 10,000) is the correlation significant;

this is indicated by the red vertical line. In other words, in 10,000 simulated samples of 8 seals,

we generally find no acoustic allometry. Panel D shows the density distribution of the simulated

linear regression coefficients, where the median value is -10.8 Hz. Given a 5.1 kg difference in

body mass between the smallest and the largest seal, we would expect, on average, a frequency

shift of 55.08 Hz. For every individual, we calculated the difference of the median f_0 values

between the silent and high noise condition; the median range across all individuals is 73.6 Hz.

This suggests that the differences caused by individual variability in f_0 in response to noise

conditions are larger than the f_0 differences expected from body mass differences alone. Seals of

differing body sizes (e.g., 7 vs. 12 kg) could thus potentially produce the same f_0 value. This

would mean that, in harbour seal pups, vocal plasticity can outweigh and mask acoustic

allometric relationships.

Figure 3. Illustration of the source-filter theory of sound production using the vocal anatomy of

the harbour seal.

752
Figure 4. Vocal anatomy of the harbour seal. (A) shows the main anatomical structures

composing the vocal tract, (B) depicts the measurements shown on a digital rendering and (C)

depicts the measurements shown on a picture of a hemi-larynx from a harbour seal pup. In panel

C, the black square outlined on the white paper serves as reference and is exactly 1 cm^2. The

vocal tract measurements taken include (1) vocal tract length (VTL), (2) vocal fold length (VFL),

(3) vocal fold thickness (VFT), (4) subglottic-tracheal dorsoventral distance 1 (STDV1), and (5)

subglottic-tracheal dorsoventral distance 2 (STDV2).

760
Figure 5. Boxplots illustrating the significant age differences between pups and weaners for (A)

VTL, (B) VFL, (C) STDV1, and (D) STDV2. The level of significance is denoted by asterisks,

where *** = 0.001.

Figure 6. Boxplots illustrating the significant sex differences for (A) VTL in pups and (B)

STDV1 in weaners. The level of significance is denoted by asterisks, where * = 0.05 and ** =

0.01.

Figure 7. Predicted effects of the body length and sex interaction for VTL including both sexes

(top), and the body mass and age interaction for VFL including both age classes (bottom). The

shading around each line of best fit indicates the 95% confidence interval.

Tables

Table 1

Pairwise Spearman correlations

Variable	Body Length (cm)	Body Mass (kg)	Girth (cm)	VTL (mm)	VFL (mm)	VFT (mm)	STDV1 (mm)
Body Mass (kg)	0.70						
Girth (cm)	0.53	0.86					
VTL (mm)	0.62	0.69	0.63				
VFL (mm)	0.73	0.79	0.65	0.72			
VFT (mm)	0.44	0.67	0.68	0.52	0.60		
STDV1 (mm)	0.63	0.78	0.65	0.70	0.82	0.69	
STDV2 (mm)	0.58	0.72	0.62	0.67	0.76	0.60	0.81

780   *Note.* All correlations were significant at $p < 0.05$ after correcting for multiple comparisons using
781   the Holm-Bonferroni method.

782   Table 2

784   *Means and standard deviations*

Variable	All		Pups		Weaners	
	Mean	SD	Mean	SD	Mean	SD
Body Length (cm)	88.07	8.03	79.86	4.79	90.19	7.32
Body Mass (kg)	14.53	3.92	9.81	1.46	15.75	3.39
Girth (cm)	57.55	9.62	46.54	5.66	60.41	8.30
VTL (mm)	91.43	6.77	83.36	4.48	93.53	5.60
VFL (mm)	10.92	1.16	9.26	0.90	11.35	0.77
VFT (mm)	5.15	0.61	4.50	0.43	5.35	0.51
STDV1 (mm)	23.97	1.95	21.35	1.37	24.65	1.43
STDV2 (mm)	17.43	1.82	15.17	1.04	18.02	1.49

787   Table 3

789   *Selected models for each vocal tract structure*

Measurement	Selected model	Deviance explained (%)
VTL	B<u>L</u> + <u>BM</u> + <u>A</u> + <u>S</u> + B<u>L*S</u>	59.30
VFL	<u>BL</u> + BM + A + <u>S</u> + A*BM + <u>A*S</u>	74.89
STDV1	<u>BL</u> + <u>BM</u> + <u>A</u> + <u>S</u>	69.99
STDV2	BL + <u>BM</u> + <u>A</u> + S	58.38

790   *Note.* Models included the predictors body length (BL), body mass (BM), age class (A) and sex

791   (S). Predictor terms joined by an asterisk denote an interaction effect. Significant predictor terms

792   are shown as underlined.

A.

CORNICULATE
CARTILAGE

ARYTENOID
MUSCLE

ARYTENOID
CARTILAGE

SOFT
PALATE

NASAL
CAVITY

HARD
PALATE

OESOPHAGUS

VOCAL
FOLDS

TRACHEAL
CARTILAGE

CRICOID
CARTILAGE

CRICOTHYROID
LIGAMENT

THYROID
CARTILAGE

HYOID
BONE

EPIGLOTTIS

TONGUE

B.

1.

2.

3.

4.

5.

C.