A crosstalk between hepcidin and IRE/IRP pathways controls ferroportin expression and determines serum iron levels in mice

Edouard Charlebois[1], Carine Fillebeen[1], Angeliki Katsarou[1], Aleksandr Rabinovich[2#],

Kazimierz Wisniewski[2†], Vivek Venkataramani[3,4], Bernhard Michalke[5], Anastasia Velentza[2‡]

and Kostas Pantopoulos[1]*

[1] Lady Davis Institute for Medical Research, Jewish General Hospital and Department of Medicine, McGill University, Montreal, Quebec, Canada

[2] Ferring Research Institute Inc, San Diego, CA

[3] Department of Medicine II, Hematology/Oncology, University Hospital Frankfurt, Frankfurt, Germany

[4] Institute of Pathology, University Medical Center Göttingen (UMG), Göttingen, Germany

[5] Helmholtz Zentrum München GmbH – German Research Center for Environmental Health, Research Unit Analytical BioGeoChemistry, Neuherberg, Germany

[#] Current address: 8234 Caminito Maritimo La Jolla CA 92037

[†] Current address: Peptide Logic, LLC, 6185 Cornerstone Ct E, Suite 112, San Diego, CA 92121

[‡] Current address: Plexium, 11494 Sorrento Valley Rd, San Diego, CA 92121

* Corresponding author; tel.: (514) 340-8260 ext. 25293; email: kostas.pantopoulos@mcgill.ca

Abstract

The iron hormone hepcidin is transcriptionally activated by iron or inflammation via distinct, partially overlapping pathways. We addressed how iron affects inflammatory hepcidin levels and the ensuing hypoferremic response. Dietary iron overload did not mitigate hepcidin induction in LPS-treated wt mice but prevented effective inflammatory hypoferremia. Likewise, LPS modestly decreased serum iron in hepcidin-deficient Hjv-/- mice, model of hemochromatosis. Synthetic hepcidin triggered hypoferremia in control but not iron-loaded wt animals. Furthermore, it dramatically decreased hepatic and splenic ferroportin in Hjv-/- mice on standard or iron-deficient diet, but only triggered hypoferremia in the latter. Mechanistically, iron antagonized hepcidin responsiveness by inactivating IRPs in the liver and spleen, to stimulate ferroportin mRNA translation. Prolonged LPS treatment eliminating ferroportin mRNA permitted hepcidin-mediated hypoferremia in iron-loaded mice. Thus, *de novo* ferroportin synthesis is critical determinant of serum iron and finetunes hepcidin-dependent functional outcomes. Our data uncover a crosstalk between hepcidin and IRE/IRP systems that controls tissue ferroportin expression and determines serum iron levels. Moreover, they suggest that hepcidin supplementation therapy is more efficient combined with iron depletion.

Introduction

Systemic iron balance is controlled by hepcidin a peptide hormone that is produced by hepatocytes in the liver and operates in target cells by binding to the iron exporter ferroportin (1, 2). This results in ferroportin internalization and lysosomal degradation but also directly inhibits ferroportin function by occluding its iron export channel (3, 4). Ferroportin is highly expressed in duodenal enterocytes and tissue macrophages, which are instrumental for dietary iron absorption and iron recycling from senescent erythrocytes, respectively. Ferroportin is also expressed in hepatocytes, where excess iron is stored and can be mobilized on demand. Hepcidin-mediated ferroportin inactivation inhibits iron entry into plasma. This is a critical homeostatic response against iron overload, but also an innate immune response against infection (5). Thus, hepcidin expression is induced when systemic iron levels are high to prevent dietary iron absorption, or under inflammatory conditions to promote iron retention within ferroportin-expressing cells and render the metal unavailable to extracellular pathogens.

The hepcidin-encoding *HAMP* gene is transcriptionally induced by iron or inflammatory stimuli via BMP/SMAD (6) or IL-6/STAT3 (7) signaling, respectively. These pathways crosstalk at different levels. For instance, the BMP co-receptor hemojuvelin (HJV), a potent enhancer of iron-dependent BMP/SMAD signaling, is also essential for inflammatory induction of hepcidin. Thus, Hjv-/- mice, a model of juvenile hemochromatosis characterized by severe iron overload and hepcidin deficiency (8), exhibit blunted inflammatory induction of hepcidin and fail to mount a hypoferremic response following LPS treatment or infection with *E. coli* (9). Excess iron inhibits hepcidin induction via the BMP/SMAD and IL-6/STAT3 signaling pathways in cultured cells (10, 11), but the *in vivo* relevance of these findings is not known.

Hepcidin-dependent inhibition of ferroportin activity and expression is a major but not the sole contributor to inflammatory hypoferremia (12, 13). This is related to the fact that ferroportin expression is regulated by additional transcriptional and post-transcriptional mechanisms (14). Thus, ferroportin transcription is induced by iron (15) and suppressed by inflammatory signals (16), while translation of *Fpn(+IRE)* mRNA, a ferroportin transcript harboring an "iron responsive element" (IRE) within its 5' untranslated regions (5' UTR) is controlled by "iron regulatory proteins" (IRPs), IRP1 and IRP2. The IRE/IRP system accounts for coordinate post-transcriptional regulation of iron metabolism proteins in cells (17, 18). In a homeostatic response to iron deficiency, IRPs bind to the (IRE) within the *Fpn(+IRE)* and ferritin (*Fth1* and *Ftl1*) mRNAs, inhibiting their translation. IRE/IRP interactions do not take place in iron-loaded cells, allowing *de novo* ferroportin and ferritin synthesis to promote iron efflux and storage, respectively. The impact of the IRE/IRP system on regulation of tissue ferroportin and serum iron is not well understood.

The aim of this work was to elucidate mechanisms by which systemic iron overload affects hepcidin expression and downstream responses, especially under inflammatory conditions. Utilizing wild type and Hjv-/- mice, we demonstrate that serum iron levels reflect regulation of ferroportin in the liver and spleen by multiple signals. We further show that effective hepcidin-mediated hypoferremia is antagonized by compensatory mechanisms aiming to prevent cellular iron overload. Our data uncover a crosstalk between hepcidin and the IRE/IRP system that controls ferroportin expression in hepatocytes, as well as liver and splenic macrophages, and thereby determines serum iron levels.

Results

Dietary iron overload does not prevent further inflammatory Hamp mRNA induction in LPS-treated wt mice, but mitigates hepcidin responsiveness. In an exploratory experiment, wt mice were subjected to dietary iron loading by feeding a high-iron diet (HID) for short (1 day), intermediate (1 week) or long (5 weeks) time intervals; control animals remained on standard diet (SD). As expected, mice on HID for 1 day manifested maximal increases in serum iron (Fig. S1A) and transferrin saturation (Fig. S1B). They retained physiological liver iron content (LIC; Fig. S1E) and serum ferritin (Fig. S1C), a reflection of LIC. Serum iron and transferrin saturation plateaued after longer HID intake, while LIC and serum ferritin gradually increased to peak at 5 weeks. The dietary iron loading promoted gradual upregulation of serum hepcidin (Fig. S1D) and liver *Hamp* mRNA (Fig. S1F), with highest values at 5 weeks. This could not prevent chronic dietary iron overload, in agreement with earlier findings (19, 20).

LPS triggered appropriate hepcidin induction and a robust hypoferremic response in control mice. Interestingly, LPS-induced inflammation resulted in further proportional increase in hepcidin and *Hamp* mRNA in dietary iron-loaded mice (Fig. S1D and S1F). This was accompanied by significant drops in serum iron and transferrin saturation. However, values did not reach the nadir of LPS-treated control animals and were increasing in mice on HID for longer periods, despite significant hepcidin accumulation. These data suggest that hepatic iron overload does not prevent inflammatory induction of hepcidin; however, it impairs its capacity to decrease serum iron.

Uncoupling inflammatory hepcidin induction from hypoferremic response in wt and Hjv-/- mice following dietary iron manipulations. To further explore the potential of hepcidin to

promote hypoferremia under iron overload, wt and Hjv-/- mice, a model of hemochromatosis,

were subjected to dietary iron manipulations. Wt mice were fed SD or HID, and Hjv-/- mice

were fed SD or an iron-deficient diet (IDD) for 5 weeks, to achieve a broad spectrum of hepcidin

regulation. Wt mice on HID and Hjv-/- mice on SD or IDD manifested similarly high serum iron

and transferrin saturation (Fig. 1A-B). Serum non-transferrin bound iron (NTBI) levels appeared

modestly elevated in the dietary and genetic iron overload models and seemed to decrease in

Hjv-/- mice on IDD (Fig. 1C). LIC was substantially reduced in Hjv-/- mice in response to IDD,

but also compared to wt mice on HID (Fig. 1E). The quantitative LIC data were corroborated

histologically by Perls staining (Fig. 1F). Dietary iron loading increased splenic iron in wt mice

and confirmed that Hjv-/- mice fail to retain iron in splenic macrophages (Fig. S2). As expected,

serum hepcidin (Fig. 1D) and liver *Hamp* mRNA (Fig. 1G) were maximally induced in HID-fed

wt mice and were low in Hjv-/- mice on SD, and further suppressed to undetectable levels

following IDD intake.

LPS reduced serum iron and transferrin saturation in hyperferremic wt mice on HID and

Hjv-/- mice on SD or IDD, but not below baseline of control wt mice on SD, the only animals

that developed a robust hypoferremic response (Fig. 1A-B). The LPS treatment was associated

with significant accumulation of hepcidin (Fig. 1D) and induction of *Hamp* mRNA (Fig. 1G) in

all experimental groups, while NTBI (Fig. 1C) and LIC (Fig. 1E) were unaffected. Notably,

LPS-treated wt mice on HID and Hjv-/- mice on IDD exhibited dramatic differences in *Hamp*

mRNA but similar blunted hypoferremic response to the acute inflammatory stimulus. Thus, the

profound hepcidin induction in iron-loaded wt mice cannot decrease serum iron below that of

iron-depleted Hjv-/- mice with negligible hepcidin, which indicates reduced hepcidin

responsiveness. In support to this interpretation, *Id1* and *Socs3* mRNAs (Fig. 1H-I), which are

markers of BMP/SMAD and IL-6/STAT3 signaling, respectively, were appropriately induced by

dietary iron loading or LPS treatment in wt mice. Thus, the major hepcidin signaling pathways

were intact under these experimental conditions.

Fpn(+IRE) mRNA levels were significantly increased in the liver of iron-loaded wt mice

on HID and Hjv-/- mice on SD (Fig. 1J), which is consistent with transcriptional induction (15).

The LPS treatment strongly suppressed liver *Fpn(+IRE)* mRNA (Fig. 1J) and induced *Dmt1*,

Zip14 and *Lcn2* mRNAs in all animals (Fig. 1K-M). These encode the divalent metal transporter

DMT1, the NTBI transporter Zip14 and the siderophore-binding protein Lcn2, respectively;

Lcn2 mRNA induction was dramatic. The transferrin receptor 1-encoding *Tfrc* mRNA was

largely unaffected by LPS, except for a reduction in Hjv-/- mice on IDD (Fig. 1N).

To assess the downstream function of hepcidin, we first analyzed ferroportin in the liver,

an organ that contributes to iron sequestration during inflammation. Immunohistochemical

staining of liver sections revealed strong ferroportin expression in Kupffer cells, predominantly

in periportal areas, under all experimental conditions (Figs. 2A and S3). Hepatocellular

ferroportin staining is also evident in the iron overload models, mostly in periportal hepatocytes

(Fig. S3), and in line with recent data (21). LPS triggered redistribution and decreased expression

of ferroportin in Kupffer cells from wt but not Hjv-/- mice (Fig. S3), as reported (9).

We further analyzed ferroportin in liver homogenates by Western blotting. Levels of

biochemically detectable liver ferroportin differed substantially between wt and Hjv-/- mice.

Thus, they were relatively low in the former and highly induced in the latter (Fig. 2B),

independently of iron load. The differences were more dramatic compared to those observed by

immunohistochemistry (Figs. 2A and S3). Conceivably, the strong ferroportin signal in Western

blots from Hjv-/- liver homogenates reflects high ferroportin expression in hepatocytes, which

are the predominant cell population and make up ~80% of the liver cell mass (22). Yet,

hepatocellular ferroportin is less visible by immunohistochemistry because the signal is

substantially weaker compared to that in Kupffer cells (see also Fig. 5E). Interestingly, the LPS

treatment visibly suppressed total liver ferroportin in Hjv-/- mice on SD but not IDD, while it

appeared to modestly affect it in wt mice (Fig. 2B). These data are consistent with negative

regulation of ferroportin by residual LPS-induced hepcidin in Hjv-/- mice on SD, which could

explain the small drop in serum iron and transferrin saturation under these acute inflammatory

conditions, as reported (9). However, liver ferroportin remained detectable and apparently

functional, as it did not allow significant iron sequestration and dramatic drop in serum iron.

Notably, persistence of relatively high serum iron is also evident in LPS-treated wt mice on HID,

despite maximal hepcidin and minimal liver ferroportin levels.

Next, we analyzed ferroportin in the spleen, an organ with erythrophagocytic

macrophages that plays an important role in body iron traffic (23). Immunohistochemical

analysis shows that LPS reduced ferroportin in red pulp splenic macrophages from wt mice on

SD, but this effect was less evident in wt mice on HID and in Hjv-/- mice on SD or IDD (Figs.

2C and S4). Western blot analysis shows a stronger ferroportin signal in splenic extracts from

Hjv-/- animals (Fig. 2D), consistent with immunohistochemistry. However, in this assay LPS

appeared to suppress splenic ferroportin in wt animals (though statistical significance was not

reached) and in Hjv-/- mice on SD, but not IDD. This could be a result of residual hepcidin

upregulation (Figs. 1D and 1G), while the lack of significant splenic ferroportin suppression in

Hjv-/- mice on IDD may indicate hepcidin insufficiency. In any case, the relatively high

circulating iron levels in dietary iron-loaded and LPS-treated wt mice indicates continuous iron

efflux to plasma despite hepcidin excess.

Insufficient hepcidin leads to blunted hypoferremic response in iron overload. We used

human synthetic hepcidin to address whether the failure of mouse models of iron overload to

mount an appropriate hypoferremic response to acute inflammation is caused by endogenous

hepcidin insufficiency or other mechanisms. Wt and Hjv-/- mice subjected to dietary iron

manipulations received 2.5 μg/g synthetic hepcidin every two hours for a total of four

intraperitoneal injections. Each dose corresponds to ~200-fold excess over endogenous

circulating hepcidin in control wt animals. The treatment caused hypoferremia in wt mice on SD

but not HID, where the decrease in serum iron was significant but well above baseline of

untreated wt controls (Fig. 3A-B). Likewise, synthetic hepcidin significantly decreased serum

iron but failed to cause dramatic hypoferremia in hepcidin-deficient Hjv-/- mice on SD.

Strikingly, hepcidin administration was much more effective in relatively iron-depleted Hjv-/-

mice on IDD, and lowered serum iron and transferrin saturation below baseline. The treatments

significantly reduced NTBI in Hjv-/- mice on SD, with a trend in mice on IDD (Fig. 3C) but did

not affect LIC or splenic iron content (SIC) under any experimental conditions (Figs. 3D-E and

S5). Serum iron represents <2% of total tissue iron and therefore its acute fluctuations are not

expected to dramatically alter LIC or SIC.

Notably, synthetic hepcidin led to significant reduction of endogenous *Hamp* mRNA in

wt mice on SD (Fig. 3F), as earlier reported (24). Conceivably, this is related to destabilization of

the Hamp inducer Tfr2 in the liver (Fig. 3G), a known response to hypoferremia (25). Synthetic

hepcidin did not promote inflammation, iron perturbations or alterations in BMP/SMAD

signaling in the liver, as judged by the unaltered expression of hepatic *Fpn, Socs3, Bmp6* and *Id1*

mRNAs (Fig. S6A-D). Moreover, synthetic hepcidin did not affect *Dmt1*, *Zip14*, *Lcn2* or *Tfrc*

mRNAs (Fig. S6E-H), which encode iron transporters; *Zip14* and *Lcn2* are also inflammatory

markers.

Next, we analyzed liver ferroportin by immunohistochemistry. Figs. 4A and S7 show that

exogenous hepcidin decreased ferroportin signal intensity in all animal groups to varying

degrees. The hepcidin effect was particularly noticeable in hepatocytes from Hjv-/- mice (see

low magnification images in Fig. S7). Kupffer cells seemed to retain some ferroportin in all

groups except Hjv-/- mice on IDD. Interestingly, while synthetic hepcidin decreased ferroportin

signal intensity in Kupffer cells, it did not alter intracellular ferroportin distribution as would be

expected based on the data in LPS-treated wt mice (Fig. 4A).

Western blotting confirmed that total liver ferroportin is highly induced in Hjv-/- mice

(Fig. 4B). Again, the signal intensity can be attributed to protein expressed in hepatocytes. The

treatment with synthetic hepcidin did not significantly affect liver ferroportin in wt mice (either

on SD or HID), but substantially reduced it in Hjv-/- mice, to almost control levels. The effect

appeared more pronounced in Hjv-/- mice on IDD; nevertheless, ferroportin remained detectable.

Splenic ferroportin was reduced in all animal groups following hepcidin treatment, with

stronger effects visualized by immunohistochemistry in wt mice on SD and Hjv-/- mice on IDD

(Figs. 4C and S8). At the biochemical level, ferroportin expression was again much stronger in

the spleen of Hjv-/- mice (Fig. 4D). Synthetic hepcidin did not significantly affect splenic

ferroportin in wt mice, but dramatically reduced it in all Hjv-/- mice.

Taken together, our data suggest that synthetic hepcidin overcomes endogenous hepcidin

deficiency in Hjv-/- mice. However, it only triggers hypoferremia in these animals following

relative iron depletion. On the other hand, in iron-loaded wt mice with already high endogenous

hepcidin, excess synthetic hepcidin fails to promote hypoferremia.

Dietary iron manipulations are sensed by IRPs in the liver and spleen of wt and Hjv-/-

mice. The IRE/IRP system orchestrates homeostatic adaptation to cellular iron supply (17, 18).

To evaluate the responses of IRPs in the whole liver and spleen to the above-described dietary

iron manipulations, we analyzed tissue extracts from wt and Hjv-/- mice by an electrophoretic

mobility shift assay (EMSA) using a ^{32}P-labelled IRE probe. The data in Fig. 5A-B show that

HID intake decreases the IRE-binding activities of IRP1 and IRP2 in both liver and spleen of wt

mice. Conversely, IDD intake stimulates the IRE-binding activities of IRP1 and IRP2 in the liver

and spleen of Hjv-/- mice. IRE/IRP2 interactions are better visible in longer exposures (left

panels). EMSAs with tissue extracts previously treated with 2-mercaptoethanol (2-ME) were

performed as loading controls (26) and are shown in the bottom panels.

To clarify which cell types of the liver account for the responses of IRPs to dietary iron,

separate EMSAs were performed using extracts from isolated hepatocytes or non-parenchymal

liver cells. The data in Fig. 5C-D uncover that IRP1 and IRP2 in both liver cell populations from

wt and Hjv-/- mice are sensitive to dietary iron loading or restriction, respectively. The EMSA

analysis of non-parenchymal liver cells, which contains Kupffer cells, showed a high

experimental variability (Fig. 5D). Nevertheless, the overall results are consistent with those

obtained with splenic extracts, which contain red pulp macrophages (Fig. 5B).

Relative expression of ferroportin in hepatocytes and non-parenchymal liver cells from

wt and Hjv-/- mice. We determined the relative abundance of ferroportin in hepatocytes and non-

parenchymal liver cells from wt and Hjv-/- mice on SD by Western blotting. As expected,

ferroportin expression (normalized to β-actin) was ~1.5-2-fold higher in the non-parenchymal

cell fraction as compared to hepatocytes in both wt and Hjv-/- mice (Fig. 5E). In comparison

across genotypes, ferroportin expression was ~2-fold higher in hepatocytes and ~50% higher in

non-parenchymal cells from Hjv-/- vs wt mice.

Iron-dependent regulation of ferroportin mRNA translation in the liver. Having

established that dietary iron manipulations trigger IRP responses in the liver and spleen, we

hypothesized that the functional outcomes of exogenous hepcidin may not merely depend on its

capacity to degrade tissue ferroportin, but also on iron-dependent ferroportin regeneration via *de*

novo synthesis. Ferroportin is encoded by two alternatively spliced transcripts, *Fpn(+IRE)* and

Fpn(-IRE), which exhibit different tissue distribution (27). *Fpn(+IRE)* mRNA is predominant in

the mouse liver and spleen (27) and is considered as a target of IRPs.

Thus, we assessed the effects of dietary iron on whole liver *Fpn(+IRE)* mRNA

translation by polysome profile analysis. We focused on the liver because this organ contains the

highest number of iron-recycling macrophages (28) and can also export iron to plasma from

ferroportin-expressing parenchymal cells. Liver extracts from wt mice on SD or HID, and Hjv-/-

mice on SD or IDD were fractionated on sucrose gradients to separate translationally inactive

light monosomes from translating heavy polysomes. The relative distribution of *Fpn(+IRE)*,

Fth1 (positive control for iron regulation) and *Actb* (negative control) mRNAs within the

different fractions was quantified by qPCR. Fig. 6A shows that dietary iron loading stimulates

Fpn(+IRE) (and *Fth1*) mRNA translation in wt mice (note the shifts from monosomes to

polysomes). Conversely, dietary iron depletion inhibits *Fpn(+IRE)* (and *Fth1*) mRNA translation

in Hjv-/- mice. We also attempted to obtain polysome profiles of *Fpn(-IRE)* mRNA but it was

undetectable after fractionation. These data indicate that in mice subjected to iron overload, iron-

stimulated ferroportin synthesis in the liver antagonizes hepcidin-mediated ferroportin

degradation and prevents an appropriate hypoferremic response. Considering that levels of

Fpn(+IRE) mRNA are elevated in iron-loaded wt and Hjv-/- mice (Figs. 1J and S6A), it is

possible that increased *de novo* ferroportin synthesis is further enhanced by transcriptional

induction.

Quantification of liver iron by ICP-MS (Fig. 6B) validated iron loading of wt mice by

HID, and relative iron depletion of Hjv-/- mice by IDD intake, respectively (see also Fig. 1E).

Iron redox speciation analysis by CE-ICP-MS revealed a profound increase in Fe^{2+}/Fe^{3+} ratios in

livers of Hjv-/- mice on SD, which was corrected by dietary iron depletion (Fig. 6C).

Nevertheless, there was no difference in Fe^{2+}/Fe^{3+} ratios among livers of wt mice on SD or HID,

and Hjv-/- mice on IDD. We conclude that an increase in total iron content, rather than excessive

accumulation of redox active Fe^{2+} drives *Fpn(+IRE)* (and *Fth1*) mRNA translation in the liver.

Restoration of effective hypoferremic response under iron overload following maximal

Fpn mRNA suppression. We reasoned that complete inactivation of ferroportin mRNA would

restore hepcidin-induced hypoferremia despite iron overload. An 8-hour treatment of mice with

LPS suppresses liver *Fpn(+IRE)* mRNA below detection levels (Fig. 7A), as reported (9). The

same holds true for the *Fpn(-IRE)* isoform (Fig. 7B), which is 290 times less abundant in control

mouse livers compared to *Fpn(+IRE)* (ΔCt=8.18), in agreement with published data (27). We

went on to examine the effects of synthetic hepcidin on serum iron under these conditions of

maximal *Fpn* mRNA suppression. Importantly, the prolonged LPS treatment decreased serum

iron in wt mice on HID below the control baseline (Fig. 7C). Furthermore, when combined with

synthetic hepcidin, it promoted an effective hypoferremic response in wt mice on HID and Hjv-/-

mice on SD (or IDD) (Figs. 7C-D) and tended to decrease NTBI (Fig. 7E). These data strongly

suggest that the expression of actively translating *Fpn* mRNA in iron-exporting tissues under

systemic iron overload mitigates hepcidin-induced drop in serum iron.

Discussion

We sought to analyze how iron overload affects hepcidin-mediated inflammatory

responses. We and others reported that excess iron inhibits the major hepcidin signaling

pathways (BMP/SMAD and IL-6/STAT3) in cultured cells (10, 11). To explore the physiological

relevance of these findings, wt mice were subjected to variable degrees of dietary iron loading

and then treated with LPS. All iron-loaded mice could further upregulate hepcidin in response to

LPS-induced acute inflammation (Fig. S1). This is consistent with other relevant findings (29)

and apparently contradicts the *in vitro* data. While experimental iron loading of cultured cells is

rapid, dietary iron loading of mice is gradual (20) and most of excess iron is effectively

detoxified within ferritin, which is highly induced (30). By contrast, the suppression of hepcidin

preceded ferritin induction in cultured cells (10), which may explain the discrepancy with the *in*

vivo data.

The unimpaired inflammatory induction of hepcidin in iron-loaded wt mice correlated

with significant drops in serum iron, but these appeared inversely proportional to the degree of

systemic iron loading (Fig. S1). Thus, LPS-treated mice on 5 weeks HID failed to reduce serum

iron below a baseline of untreated control mice on SD. This indicates a defect in mounting a

hypoferremic response to inflammation, which can be attributed to mechanisms antagonizing

hepcidin action. To explore how iron modulates the capacity of hepcidin to trigger inflammatory

hypoferremia, we established conditions of iron overload using wt and Hjv-/- mice with extreme

differences in hepcidin expression. Figs. 1 and 2 demonstrate that iron overload prevents

effective inflammatory hypoferremia independently of hepcidin and tissue ferroportin levels.

We used a ~200-fold excess of synthetic hepcidin to directly assess its capacity to divert

iron traffic in iron-loaded mice. Hepcidin injection caused hypoferremia in control wt mice on

SD and significantly reduced serum iron in wt mice on HID and Hjv-/- mice on SD, but not

below baseline (Fig. 3). Thus, synthetic hepcidin failed to drastically drop serum iron levels in

iron overload models with either high or low endogenous hepcidin. Importantly, synthetic

hepcidin promoted robust hypoferremia in relatively iron-depleted Hjv-/- mice on IDD, with

undetectable endogenous hepcidin. It should be noted that synthetic hepcidin had similar effects

on tissue ferroportin among wt or Hjv-/- mice, regardless of iron diet (Fig. 4). It reduced

intensity of the ferroportin signal in Kupffer cells and splenic macrophages of wt mice without

significantly affecting biochemically detectable total protein levels. In addition, it dramatically

reduced total ferroportin in the liver and spleen of Hjv-/- mice. However, in all experimental

settings there was residual tissue ferroportin, which appears to be functionally significant.

We reasoned that at steady-state, tissue ferroportin may consist of fractions of newly

synthesized protein, and protein that is *en route* to hepcidin-mediated degradation. Conceivably,

the former may exhibit more robust iron export activity, at least before its iron channel gets

occluded by hepcidin. Increased *de novo* synthesis of active ferroportin could explain why

synthetic hepcidin cannot drastically drop serum iron levels under iron overload. In fact, Fig. 6

demonstrates that dietary iron overload augments *Fpn(+IRE)* mRNA translation in the liver of

wt mice. Conversely, relative dietary iron depletion inhibits *Fpn(+IRE)* mRNA translation in the

liver of Hjv-/- mice, in line with the restoration of hepcidin-mediated hypoferremic response

(Fig. 3).

Our data are consistent with translational control of liver ferroportin expression via the

IRE/IRP system and do not exclude the possibility for an additional contribution of iron-

dependent transcriptional regulation of *Fpn(+IRE)* mRNA. Direct evidence for activation of IRP

responses in the liver and spleen to dietary iron manipulations is provided in Fig. 5. While

translational control of ferritin in tissues is established (31), regulation of ferroportin by the

IRE/IRP system is less well characterized and has hitherto only been documented in cell models

(32, 33), the mouse duodenum (34), and the rat liver (35). Moreover, the physiological relevance

of this mechanism remained speculative. The data in Figs. 5 and 6 show that the IRE/IRP system

is operational and controls *Fpn(+IRE)* mRNA translation in both hepatocytes and non-

parenchymal liver cells. Presumably, this offers a compensatory mechanism to protect these cells

from iron overload and iron-induced toxicity. On the other hand, this mechanism attenuates

hepcidin responsiveness and promotes a state of hepcidin resistance, as higher amounts of

hepcidin are required to achieve effective hypoferremia. Because hepcidin has a short plasma

half-life, it is reasonable to predict that the use of more potent hepcidin analogs (36) will

overcome the antagonistic effects of increased ferroportin mRNA translation under iron

overload.

The critical role of *de novo* ferroportin synthesis in fine-tuning hepcidin-dependent

functional outcomes is also highlighted in Fig. 7. Thus, synthetic hepcidin was highly effective

as promoter of hypoferremia in dietary iron-loaded wt mice when administered together with

LPS. LPS is known to suppress *Fpn* mRNA in cell lines (16) and mouse tissues, with a nadir in

the liver reached at 8 h (9). The recovery of hepcidin effectiveness in mouse models of iron

overload was only possible when *Fpn* mRNA was essentially eliminated. Under these conditions,

LPS treatment alone was sufficient to decrease serum iron in dietary iron-loaded wt mice below

baseline.

Tissue iron uptake may be another important determinant of the hypoferremic response to

inflammation. LPS did not affect *Tfrc* mRNA levels in the liver (Fig. 1N), which argues against

increased uptake of transferrin-bound iron via TfR1. On the other hand, LPS induced *Zip14*,

Dmt1 and *Lcn2* mRNAs (Figs 1K-M). Zip14 is the NTBI transporter accounting for

hepatocellular iron overload in hemochromatosis (37) and is upregulated by inflammatory cues

in hepatocytes (38). DMT1 is dispensable for NTBI uptake by hepatocytes (39), and its

inflammatory induction might promote iron acquisition by macrophages (16, 40). Nevertheless,

considering that the fraction of NTBI represents <2% of total serum iron even in the iron

overload models (Figs. 1A and C), it is implausible that NTBI uptake by Zip14 and/or DMT1

substantially contributes to inflammatory hypoferremia. Lcn2 is an acute phase protein that can

sequester intracellular iron bound to catecholate siderophores (41), and is more likely to transport

iron to tissues during infection. In any case, synthetic hepcidin did not affect expression of iron

transporters (Fig. S6E-H). This excludes the possibility for a synergistic effect on LPS-induced

tissue iron uptake that could promote effective hypoferremia in the iron overload models.

In conclusion, our data reveal a crosstalk between the hepcidin pathway and the IRE/IRP

system in the liver and spleen for the control of tissue ferroportin and serum iron levels.

Furthermore, they suggest that application of hepcidin therapeutics for treatment of iron overload

disorders should be combined with iron depletion strategies to mitigate *Fpn* mRNA translation

and increase hepcidin efficacy. Future work is expected to clarify whether optimizing the

hypoferremic response to inflammation under systemic iron overload decreases susceptibility to

pathogens.

Materials and Methods

Animals. Wild type C57BL/6J and isogenic Hjv-/- mice (42) were housed in macrolone cages (up to 5 mice/cage, 12:12 h light-dark cycle: 7 am - 7 pm; $22 \pm 1°C$, $60 \pm 5\%$ humidity). The mice were fed either a standard diet (200 ppm iron), an iron-deficient diet (2-6 ppm iron) or a high-iron diet (2% carbonyl iron) (43). Where indicated, mice were injected intraperitoneally with 1 µg/g LPS (serotype 055:B5; Sigma-Aldrich) or subcutaneously with 2.5 µg/g synthetic hepcidin; control mice were injected with phosphate-buffered saline. At the endpoints, animals were sacrificed by CO_2 inhalation and cervical dislocation. Experimental procedures were approved by the Animal Care Committee of McGill University (protocol 4966).

Serum biochemistry. Blood was collected via cardiac puncture. Serum was prepared by using micro Z-gel tubes with clotting activator (Sarstedt) and was kept frozen at $-20°C$ until analysis. Serum iron, total iron binding capacity (TIBC) and, where indicated serum ferritin, were determined at the Biochemistry Department of the Montreal Jewish General Hospital using a Roche Hitachi 917 Chemistry Analyzer. Transferrin saturation was calculated from the ratio of serum iron and TIBC. Serum hepcidin was measured by using an ELISA kit (HMC-001; Intrinsic LifeSciences).

Quantification of serum non-transferrin bound iron (NTBI). NTBI was measured by adapting the method developed by Esposito *et al* (44). Briefly, iron samples of known concentration were created by mixing 70 mM nitrilotriacetate (NTA) (pH = 7.0) with 20 mM ferrous ammonium sulfate. Fe^{2+} was allowed to oxidize to Fe^{3+} in ambient air for at least 30 min and then the solution was diluted to 0.2 mM before further serial dilutions to create a ladder. 5 µl

of ladder was loaded in a 96-well plate containing 195 μl plasma-like medium with or without

100 μM deferiprone. The composition of the plasma-like medium was: 40 mg/ml bovine serum

albumin, 1.2 mM sodium phosphate dibasic, 120 μM sodium citrate, 10 mM sodium bicarbonate

in iron-free HEPES-buffered saline (HEPES 20 mM, NaCl 150 mM, treated with Chelex-100

chelating resin [Bio-Rad, Hercules, CA], 0.5 mM NTA, 40 μM ascorbic acid, 50 μM

dihydrorhodamine, pH=7.4). 5 μl of sample was loaded in a 96-well plate containing 195 μl of

iron-free HEPES-buffered saline with or without 100 μM deferiprone. Microplates were read

every 2 minutes at 37°C over 40 min at 485/520 nm (ex/em). Final NTBI was calculated by

comparing the oxidation rate of DHR in the presence or absence of the strong chelator

deferiprone.

Hepcidin synthesis. Human hepcidin (DTHFPICIFCCGCCHRSKCGMCCKT) was

synthesized at Ferring Research Institute, San Diego, CA. The linear peptide was assembled on

Rink amide resin using Tribute peptide synthesizer and the peptide was cleaved from the resin

with the TFA/TIS/EDT/H_2O 91:3:3:3 (v/v/v/v) cocktail. The solvents were evaporated, and the

crude peptide was precipitated with diethyl ether, reconstituted in 50% aqueous acetonitrile and

lyophilized. The lyophilizate was dissolved in 30% aqueous acetonitrile at the concentration of

0.05 mM and the pH of the solution was adjusted to 7.8 with 6 M ammonium hydroxide. Folding

was achieved within 4 hours using the cysteine/cystine redox (peptide/Cys/Cys_2 1:6:6 molar

ratio). The reaction mixture was acidified to pH 3, loaded onto HPLC prep column and purified

in a TFA based gradient. The identity of the peptide was confirmed by mass spectrometry and by

coelution with a commercially available sample (Peptide International, #PLP-3771-PI).

Quantitative real-time PCR (qPCR). RNA was extracted from livers by using the RNeasy

kit (Qiagen). cDNA was synthesized from 1 μg RNA by using the OneScript® Plus cDNA

Synthesis Kit (Applied Biological Materials Inc.). Gene-specific primers pairs (Table S1) were validated by dissociation curve analysis and demonstrated amplification efficiency between 90-110 %. SYBR Green (Bioline) and primers were used to amplify products under following cycling conditions: initial denaturation 95°C 10 min, 40 cycles of 95°C 5 s, 58°C 30 s, 72°C 10 s, and final cycle melt analysis between 58°-95°C. Relative mRNA expression was calculated by the $2^{-\Delta\Delta Ct}$ method (45). Data were normalized to murine ribosomal protein L19 (*Rpl19*). Data are reported as fold increases compared to samples from wild type mice on standard diet (SD).

Polysome fractionation. RNA was freshly prepared from frozen livers. Linear sucrose gradients were prepared the day before the experiment by using 5% (w/v) and 50% (w/v) sucrose solutions with 10x gradient buffer (200 mM HEPES pH=7.6, 1 M KCl, 50 mM $MgCl_2$, 0.1 mg/ml Cycloheximide, 1 tablet cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche), 200 U/mL Recombinant RNasin® Ribonuclease Inhibitor (Promega), 2 mM DTT). Linear gradients were prepared in Polyallomer Centrifuge Tubes (Beckman Coulter). Tubes were marked using a gradient cylinder (BioComp), and 5% sucrose solution was added using a syringe with a layering needle (BioComp) until solution level reached the mark. Then, 50% sucrose solution was layered underneath the 5% solution until the interface between the two solutions reached the mark. Tubes were capped with rate zonal caps (BioComp) and linearized using a Gradient Master 108 (Biocomp). All reagents were nuclease-free and all solutions were kept on ice or at 4°C. Sample preparation was adapted from Liang *et al.* (46). Briefly, livers were flash frozen upon collection. Roughly 30-80 mg of tissue was crushed using a mortar and pestle in the presence of liquid nitrogen to prevent thawing. Tissues were lysed in up to 1 ml of hypotonic lysis buffer (5 mM Tris-Hcl pH=7.5, 1.5 mM KCl, 2.5 mM $MgCl_2$, 2 mM DTT, 1 mg/ml Cycloheximide, 200 U/ml Recombinant RNasin® Ribonuclease Inhibitor (Promega), 1 tablet

cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche), 0.5% (v/v) Triton X-100,

0.5% (v/v) Sodium Deoxycholate) and homogenized using Dounce homogenizers (60

movements with both loose and tight pestles) on ice. Samples were centrifuged at 4°C, 16,060g

for 4 minutes and supernatants were collected. Sample optical density was measured at 260 nM

and samples were normalized to either the lowest value or 30 ODs. 450 µl of sucrose gradient

was removed from the top and replaced with normalized sample. Tube weights were balanced by

weight before centrifugation at 200,000g for 2 h at 4°C in a SW 41 Ti rotor and a Beckman

Optima L-60 Ultracentrifuge. Samples were fractionated using a BR-188 Density Gradient

Fractionation System (Brandel). Immediately upon collection, 800 µl of samples were mixed

with 1 ml of TRIzol™ and kept on ice before storage at -80°C. Polysomal RNA was processed

according to the manufacturer's protocol. mRNA distribution was analyzed as previously

described (47).

Electrophoretic mobility shift assay (EMSA). IRE-binding activities from liver and spleen

were analyzed by EMSA using a radioactive ^{32}P-labelled IRE probe, according to established

procedures (26). EMSAs were also performed in extracts from hepatocytes and non-parenchymal

cells, which were separated by using a 2-step collagenase perfusion technique, as previously

described (9).

Western blotting. Livers were washed with ice-cold PBS and dissected into pieces.

Aliquots were snap frozen at liquid nitrogen and stored at -80^{o}C. Protein lysates were obtained

as described (21). Lysates containing 40 µg of proteins were analyzed by SDS-PAGE on 9-13%

gels and proteins were transferred onto nitrocellulose membranes (BioRad). The blots were

blocked in non-fat milk diluted in tris-buffered saline (TBS) containing 0.1% (v/v) Tween-20

(TBS-T), and probed overnight with antibodies against ferroportin (48) (1:1000 diluted

monoclonal rat anti-mouse 1C7, kindly provided by Amgen Inc), β-actin (1:2000 diluted;

Sigma), or Tfr2 (1:1000 diluted rabbit polyclonal; Alpha Diagnostics). Following a 3x wash with

TBS-T, the membranes were incubated with peroxidase-coupled secondary antibodies for 1 h.

Immunoreactive bands were detected by enhanced chemiluminescence with the Western

Lightning ECL Kit (Perkin Elmer).

Immunohistochemistry. Tissue specimens were fixed in 10% buffered formalin and

embedded in paraffin. Samples from 3 different mice for each experimental condition were cut at

4-μm, placed on SuperFrost/Plus slides (Fisher) and dried overnight at 37°C. The slides were

then loaded onto the Discovery XT Autostainer (Ventana Medical System) for automated

immunohistochemistry. Slides underwent de-paraffinization and heat-induced epitope retrieval.

Immunostaining was performed by using 1:500 diluted rabbit polyclonal antibodies against

ferroportin (49) and an appropriate detection kit (Omnimap rabbit polyclonal HRP, #760-4311

and ChromoMap-DAB #760-159; Roche). Negative controls were performed by the omission of

the primary antibody. Slides were counterstained with hematoxylin for four minutes, blued with

Bluing Reagent for four minutes, removed from the autostainer, washed in warm soapy water,

dehydrated through graded alcohols, cleared in xylene, and mounted with Permount (Fisher).

Sections were analyzed by conventional light microscopy and quantified by using the Aperio

ImageScope software (Leica Biosystems) (9).

Perls Prussian blue staining. To visualize non-heme iron deposits, deparaffinized tissue

sections were stained with Perls' Prussian blue using the Accustain Iron Stain kit (Sigma).

Quantification of liver iron content (LIC). Total liver iron was quantified by using the

ferrozine assay (20) or inductively coupled plasma mass spectrometry (ICP-MS) (50).

Iron speciation analysis. Iron redox speciation analysis in the liver was performed by

capillary electrophoresis (CE) coupled to ICP-MS (CE-ICP-MS). Dynamic reaction cell (DRC)

technology (ICP-DRC-MS) with NH_3 as DRC-gas was applied for non-interfered monitoring of

the iron isotopes. A "PrinCe 706" CE system (PrinCe Technologies B.V., Emmen, Netherlands)

was employed for separation of iron species at +20 kV. Temperature settings for sample/buffer

tray and capillary were set to 20^oC. An uncoated capillary (100 cm x 50 µm ID; CS-

Chromatographie Service GmbH, Langerwehe, Germany) was used for separation and

hyphenation to the ICP–DRC-MS. A CE-ICP-MS interface (50, 51) was installed which

provided the electrical connection between CE capillary end and outlet electrode. The self-

aspiration mode allowed for best flow rate adjustment and avoided suction flow. Electrolytes for

sample stacking and electrophoretic separation were 10% HCl = leading electrolyte, 0.05 mM

HCl= terminating electrolyte and 50 mM HCl = background electrolyte. The Fe^{2+}/Fe^{3+} ratio was

calculated from quantitative determined concentrations of Fe-species.

Statistics. Statistical analysis was performed by using the Prism GraphPad software

(version 9.1.0). Lognormally distributed data including qPCR and ELISA results were first log

transformed before analysis with ordinary two-way ANOVA (Tukey's multiple comparisons

test) for comparisons within same treatment groups (denoted by a or b on figures) or with

multiple unpaired t tests using the Holm-Sidak method to compare effects between treatments.

Normally distributed data was analyzed by two-way ANOVA using either Sidak's method for

comparisons between treatment groups or Tukey's multiple comparisons test within treatments

groups. Probability value $p<0.05$ was considered statistically significant.

Acknowledgments

We thank Dr. Naciba Benlimame and Lilian Canetti for assistance with histology and immunohistochemistry. This work was supported by a grant from the Canadian Institutes of Health Research (CIHR; PJT-159730). EC was funded by a fellowship from the Natural Sciences and Engineering Research Council of Canada (NSERC) and is currently a recipient of a fellowship from the *Fonds de recherche du Québec – Santé* (FRQS). The work of VV and BM was financially supported by the Deutsche Forschungsgemeinschaft (DFG) through the Priority Program "Ferroptosis: from Molecular Basics to Clinical Applications" (SPP 2306).

Competing interests

The authors declare no competing financial interests.

References

1. Camaschella C, Nai A, Silvestri L. Iron metabolism and iron disorders revisited in the hepcidin era. Haematologica. 2020;105(2):260-72.
2. Katsarou A, Pantopoulos K. Basics and principles of cellular and systemic iron homeostasis. Mol Aspects Med. 2020;75:100866.
3. Aschemeyer S, Qiao B, Stefanova D, Valore EV, Sek AC, Ruwe TA, et al. Structure-function analysis of ferroportin defines the binding site and an alternative mechanism of action of hepcidin. Blood. 2018;131(8):899-910.
4. Billesbolle CB, Azumaya CM, Kretsch RC, Powers AS, Gonen S, Schneider S, et al. Structure of hepcidin-bound ferroportin reveals iron homeostatic mechanisms. Nature. 2020;586(7831):807-11.
5. Ganz T, Nemeth E. Iron homeostasis in host defence and inflammation. Nat Rev Immunol. 2015;15(8):500-10.
6. Wang CY, Babitt JL. Liver iron sensing and body iron homeostasis. Blood. 2019;133(1):18-29.
7. Schmidt PJ. Regulation of Iron Metabolism by Hepcidin under Conditions of Inflammation. J Biol Chem. 2015;290(31):18975-83.
8. Huang FW, Pinkus JL, Pinkus GS, Fleming MD, Andrews NC. A mouse model of juvenile hemochromatosis. J Clin Invest. 2005;115(8):2187-91.

9.   Fillebeen C, Wilkinson N, Charlebois E, Katsarou A, Wagner J, Pantopoulos K. Hepcidin-
mediated hypoferremic response to acute inflammation requires a threshold of
Bmp6/Hjv/Smad signaling. Blood. 2018;132(17):1829-41.
10. Charlebois E, Pantopoulos K. Iron overload inhibits BMP/SMAD and IL-6/STAT3 signaling
to hepcidin in cultured hepatocytes. PLoS ONE. 2021;16(6):e0253475.
11. Yu LN, Wang SJ, Chen C, Rausch V, Elshaarawy O, Mueller S. Direct modulation of
hepatocyte hepcidin signaling by iron. World J Hepatol. 2021;13(10):1378-93.
12. Guida C, Altamura S, Klein FA, Galy B, Boutros M, Ulmer AJ, et al. A novel inflammatory
pathway mediating rapid hepcidin-independent hypoferremia. Blood. 2015;125(14):2265-
75.
13. Deschemin JC, Vaulont S. Role of hepcidin in the setting of hypoferremia during acute
inflammation. PLoS ONE. 2013;8(4):e61050.
14. Drakesmith H, Nemeth E, Ganz T. Ironing out Ferroportin. Cell Metab. 2015;22(5):777-87.
15. Aydemir F, Jenkitkasemwong S, Gulec S, Knutson MD. Iron loading increases ferroportin
heterogeneous nuclear RNA and mRNA levels in murine J774 macrophages. J Nutr.
2009;139(3):434-8.
16. Ludwiczek S, Aigner E, Theurl I, Weiss G. Cytokine-mediated regulation of iron transport
in human monocytic cells. Blood. 2003;101:4148-54.
17. Wang J, Pantopoulos K. Regulation of cellular iron metabolism. Biochem J.
2011;434(3):365-81.
18. Muckenthaler MU, Galy B, Hentze MW. Systemic iron homeostasis and the iron-responsive
element/iron-regulatory protein (IRE/IRP) regulatory network. Annu Rev Nutr.
2008;28:197-213.
19. Corradini E, Meynard D, Wu Q, Chen S, Ventura P, Pietrangelo A, et al. Serum and liver
iron differently regulate the bone morphogenetic protein 6 (BMP6)-SMAD signaling
pathway in mice. Hepatology. 2011;54(1):273-84.
20. Daba A, Gkouvatsos K, Sebastiani G, Pantopoulos K. Differences in activation of mouse
hepcidin by dietary iron and parenterally administered iron dextran: compartmentalization is
critical for iron sensing. J Mol Med (Berl). 2013;91:95-102.
21. Katsarou A, Gkouvatsos K, Fillebeen C, Pantopoulos K. Tissue-Specific Regulation of
Ferroportin in Wild-Type and Hjv-/- Mice Following Dietary Iron Manipulations. Hepatol
Commun. 2021;5(12):2139-50.
22. Schulze RJ, Schott MB, Casey CA, Tuma PL, McNiven MA. The cell biology of the
hepatocyte: A membrane trafficking machine. J Cell Biol. 2019;218(7):2096-112.
23. Kurotaki D, Uede T, Tamura T. Functions and development of red pulp macrophages.
Microbiol Immunol. 2015;59(2):55-62.
24. Laftah AH, Ramesh B, Simpson RJ, Solanky N, Bahram S, Schumann K, et al. Effect of
hepcidin on intestinal iron absorption in mice. Blood. 2004;103:3940-4.
25. Johnson MB, Enns CA. Diferric transferrin regulates transferrin receptor 2 protein stability.
Blood. 2004;104:4287-93.
26. Fillebeen C, Wilkinson N, Pantopoulos K. Electrophoretic mobility shift assay (EMSA) for
the study of RNA-protein interactions: the IRE/IRP example. J Vis Exp. 2014(94).
27. Zhang DL, Hughes RM, Ollivierre-Wilson H, Ghosh MC, Rouault TA. A ferroportin
transcript that lacks an iron-responsive element enables duodenal and erythroid precursor
cells to evade translational repression. Cell Metab. 2009;9(5):461-73.

28. Krenkel O, Tacke F. Liver macrophages in tissue homeostasis and disease. Nat Rev
Immunol. 2017;17(5):306-21.
29. Enculescu M, Metzendorf C, Sparla R, Hahnel M, Bode J, Muckenthaler MU, et al.
Modelling Systemic Iron Regulation during Dietary Iron Overload and Acute Inflammation:
Role of Hepcidin-Independent Mechanisms. PLoS Comput Biol. 2017;13(1):e1005322.
30. Kent P, Wilkinson N, Constante M, Fillebeen C, Gkouvatsos K, Wagner J, et al. Hfe and
Hjv exhibit overlapping functions for iron signaling to hepcidin. J Mol Med (Berl).
2015;93(5):489-98.
31. Wilkinson N, Pantopoulos K. The IRP/IRE system in vivo: insights from mouse models.
Front Pharmacol. 2014;5:176.
32. Lymboussaki A, Pignatti E, Montosi G, Garuti C, Haile DJ, Pietrangelo A. The role of the
iron responsive element in the control of ferroportin1/IREG1/MTP1 gene expression. J
Hepatol. 2003;39:710-5.
33. Nairz M, Ferring-Appel D, Casarrubea D, Sonnweber T, Viatte L, Schroll A, et al. Iron
Regulatory Proteins Mediate Host Resistance to Salmonella Infection. Cell Host Microbe.
2015;18(2):254-61.
34. Galy B, Ferring-Appel D, Becker C, Gretz N, Grone HJ, Schumann K, et al. Iron regulatory
proteins control a mucosal block to intestinal iron absorption. Cell Rep. 2013;3(3):844-57.
35. Garza KR, Clarke SL, Ho YH, Bruss MD, Vasanthakumar A, Anderson SA, et al.
Differential translational control of 5' IRE-containing mRNA in response to dietary iron
deficiency and acute iron overload. Metallomics. 2020;12(12):2186-98.
36. Katsarou A, Pantopoulos K. Hepcidin Therapeutics. Pharmaceuticals (Basel). 2018;11(4).
37. Jenkitkasemwong S, Wang CY, Coffey R, Zhang W, Chan A, Biel T, et al. SLC39A14 Is
Required for the Development of Hepatocellular Iron Overload in Murine Models of
Hereditary Hemochromatosis. Cell Metab. 2015;22(1):138-50.
38. Liuzzi JP, Lichten LA, Rivera S, Blanchard RK, Aydemir TB, Knutson MD, et al.
Interleukin-6 regulates the zinc transporter Zip14 in liver and contributes to the
hypozincemia of the acute-phase response. Proc Natl Acad Sci USA. 2005;102:6843-8.
39. Wang CY, Knutson MD. Hepatocyte divalent metal-ion transporter-1 is dispensable for
hepatic iron accumulation and non-transferrin-bound iron uptake in mice. Hepatology.
2013;58(2):788-98.
40. Wardrop SL, Richardson DR. Interferon-gamma and lipopolysaccharide regulate the
expression of Nramp2 and increase the uptake of iron from low relative molecular mass
complexes by macrophages. Eur J Biochem. 2000;267(22):6586-93.
41. Xiao X, Yeoh BS, Vijay-Kumar M. Lipocalin 2: An Emerging Player in Iron Homeostasis
and Inflammation. Annu Rev Nutr. 2017;37:103-30.
42. Gkouvatsos K, Fillebeen C, Daba A, Wagner J, Sebastiani G, Pantopoulos K. Iron-
dependent regulation of hepcidin in Hjv-/- mice: evidence that hemojuvelin is dispensable
for sensing body iron levels. PLoS ONE. 2014;9(1):e85530.
43. Fillebeen C, Charlebois E, Wagner J, Katsarou A, Mui J, Vali H, et al. Transferrin receptor 1
controls systemic iron homeostasis by fine-tuning hepcidin expression to hepatocellular iron
load. Blood. 2019;133(4):344-55.
44. Esposito BP, Breuer W, Sirankapracha P, Pootrakul P, Hershko C, Cabantchik ZI. Labile
plasma iron in iron overload: redox activity and susceptibility to chelation. Blood.
2003;102(7):2670-7.

45. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. Methods. 2001;25(4):402-8.
46. Liang S, Bellato HM, Lorent J, Fernanda, Oertlin C, Vincent, et al. Polysome-profiling in
small tissue samples. Nucleic Acids Res. 2018;46(1):e3-e.
47. Panda A, Martindale J, Gorospe M. Polysome Fractionation to Analyze mRNA Distribution
Profiles. BIO-PROTOCOL. 2017;7(3).
48. Ross SL, Biswas K, Rottman J, Allen JR, Long J, Miranda LP, et al. Identification of
Antibody and Small Molecule Antagonists of Ferroportin-Hepcidin Interaction. Front
Pharmacol. 2017;8:838.
49. Maffettone C, Chen G, Drozdov I, Ouzounis C, Pantopoulos K. Tumorigenic properties of
iron regulatory protein 2 (IRP2) mediated by its specific 73-amino acids insert. PLoS ONE.
2010;5(4):e10163.
50. Michalke B, Willkommen D, Venkataramani V. Iron Redox Speciation Analysis Using
Capillary Electrophoresis Coupled to Inductively Coupled Plasma Mass Spectrometry (CE-
ICP-MS). Front Chem. 2019;7:136.
51. Michalke B, Willkommen D, Venkataramani V. Setup of Capillary Electrophoresis-
Inductively Coupled Plasma Mass Spectrometry (CE-ICP-MS) for Quantification of Iron
Redox Species (Fe(II), Fe(III)). J Vis Exp. 2020(159).

Figures legends

Fig. 1. Iron overload blunts hepcidin responsiveness to LPS-induced inflammation. Four-week-old male wt mice (n=12-14 per group) were placed on HID for five weeks. Conversely, age- and sex-matched isogenic Hjv-/- mice (n=12-14 per group) were placed on IDD for five weeks to prevent excessive iron overload. Controls from both genotypes were kept on SD. Half of the mice were injected with saline and the other half with 1 µg/g LPS; all animals were sacrificed 4 hours later. Sera were collected by cardiac puncture and analyzed for: (A) iron, (B) transferrin saturation, (C) NTBI, and (D) hepcidin. Livers were dissected and processed for LIC quantification by the ferrozine assay (E) and for histological detection of iron deposits by Perls' staining (magnification: 10x). Livers were also used for qPCR analysis of following mRNAs: (G) *Hamp*, (H) Id1, (I) Socs3, (J) *Fpn(+IRE)*, (K) *Dmt1*, (L) *Zip14*, (M) *Lcn2* and (N) *Tfrc*. The

dotted line in (A) and (B) indicates baseline serum iron and transferrin saturation, respectively,

of control wt mice on SD. Data in (A-E) are presented as the mean±SEM while in (G-N) are

presented as geometric mean±SD. Statistically significant differences ($p < 0.05$) compared to

values from saline- or LPS-treated wt control mice are indicated by a or b, respectively.

Fig. 2. Effects of LPS on hepatic and splenic ferroportin of iron-manipulated wt and Hjv-/-

mice. Livers and spleens from mice described in Fig. 1 were dissected and processed for

immunohistochemical and biochemical analysis of ferroportin. Immunohistochemical staining of

ferroportin in liver (A) and spleen (C) sections (magnification for liver is 20x and for spleen 5x).

Western blot for ferroportin and β-actin in liver (B) and spleen (D) extracts from four

representative mice in each condition. Blots were quantified by densitometry and ferroportin/β-

actin ratios are shown on the right. Densitometric data are presented as the mean±SEM.

Statistically significant differences ($p < 0.05$) compared to values from saline- or LPS-treated wt

control mice are indicated by a or b, respectively.

Fig. 3. Iron depletion of Hjv-/- mice improves the efficacy of synthetic hepcidin to promote

hypoferremia. Four-week-old wt male mice (n=12-14 per group) were placed on HID for five

weeks. Conversely, age- and sex-matched isogenic Hjv-/- mice (n=12-14 per group) were placed

on IDD for five weeks to prevent excessive iron overload. Controls from both genotypes were

kept on SD. Half of the mice were injected every 2 hours for a total of 4 injections with saline,

and the other half with 2.5 µg/g synthetic hepcidin. Sera were collected by cardiac puncture and

analyzed for: (A) iron, (B) transferrin saturation, and (C) NTBI. Livers and spleens were

dissected and processed for analysis of: (D) LIC and (E) SIC by the ferrozine assay. (F) qPCR

analysis of liver *Hamp* mRNA (F). (G) Western blot analysis of liver Tfrc, Tfr2, and β-actin; a

representative image (out of n=2 samples) is shown on the left; the blots were quantified by

densitometry and Tfrc/β-actin or Tfr2/β-actin ratios are shown on the right. The dotted line in

(A) and (B) indicates baseline serum iron and transferrin saturation, respectively, of control wt

mice on SD. Data in (A-E) are presented as the mean±SEM and in (F) as geometric mean±SD.

Statistically significant differences (p<0.05) compared to values from saline- or hepcidin-treated

wt control mice are indicated by a or b, respectively.

Fig. 4. Effects of synthetic hepcidin on hepatic and splenic ferroportin of iron-manipulated

wt and Hjv-/- mice. Livers and spleens from mice described in Fig. 3 were dissected and

processed for immunohistochemical and biochemical analysis of ferroportin.

Immunohistochemical staining of ferroportin in liver (A) and spleen (C) sections (magnification

for liver is 20x and for spleen 10x). Western blot for ferroportin and β-actin in liver (B) and

spleen (D) extracts from four representative mice in each condition. Blots were quantified by

densitometry and ferroportin/β-actin ratios are shown on the right. Densitometric data are

presented as the mean±SEM. Statistically significant differences (p<0.05) compared to values

from saline- or hepcidin-treated wt control mice are indicated by a or b, respectively.

Fig. 5. Dietary iron manipulations trigger IRP responses in the liver and spleen, as well as

in primary hepatocytes and non-parenchymal liver cells of wt and Hjv-/- mice. Whole liver

(A), whole spleen (B), isolated hepatocytes (C) or isolated non-parenchymal liver cells (D) from

the mice described in Fig. 3 were analyzed for IRE-binding activity by EMSA with a ^{32}P-labelled

IRE probe in the absence (top) or presence (bottom) of 2% mercaptoethanol (2-ME). Two or

three representative samples from each condition are shown. The positions of IRE/IRP1 and

IRE/IRP2 complexes are indicated by arrows. Shorter and longer exposures of the

autoradiograms are shown in the left and right panels, respectively. (E) Isolated hepatocytes and

isolated non-parenchymal liver cells were analyzed by Western blotting for expression of

ferroportin and β-actin. Blots were quantified by densitometry and ferroportin/β-actin ratios are

shown on the right. Densitometric data are presented as the mean±SEM.

Fig. 6. Iron regulation of Fpn(+IRE) mRNA translation in the mouse liver. Four-week-old

wt male mice (n=10-14 per group) were placed on HID for five weeks. Conversely, age- and sex-

matched isogenic Hjv-/- mice (n=10-14 per group) were placed on IDD for five weeks to prevent

excessive iron overload. Controls from both genotypes were kept on SD. At the endpoint, the

mice were sacrificed, and livers were used for polysome profile analysis and iron assays. (A)

Liver polysome profiles from n=3 mice in each experimental group. Top: Recording of

absorbance at 254 nm of representative samples. Fraction numbers and direction of the gradient

are indicated. Bottom left: Distribution of *Fpn(+IRE)*, *Fth1* and *Actb* mRNAs among light

monosomal and heavy polysomal fractions (separated by dashed line) was analyzed by qPCR.

Bottom right: Bar graph comparisons of pooled fractions. Numbers indicate the fold change

compared to wt mice on SD. (B and C) Analysis of total iron (B), and redox iron speciation (C)

in the liver by CE-ICP-MS. Data are presented as the mean±SEM. Statistical analysis in (A) was

performed by two-way ANOVA and in (B, C) by one-way ANOVA or unpaired t-test.

Statistically significant differences (p<0.05) compared to values from wt control mice on SD are

indicated by a.

Fig. 7. Elimination of ferroportin mRNA by prolonged LPS treatment potentiates hepcidin-

induced hypoferremia in mouse models of iron overload. Four-week-old wt male mice (n=10-

14 per group) were placed on HID for five weeks. Conversely, age- and sex-matched isogenic

Hjv-/- mice (n=10-14 per group) were placed on IDD for five weeks to prevent excessive iron

overload. Controls from both genotypes were kept on SD. (A and B) Half of the mice were

injected with saline and the other half with 1 µg/g LPS and sacrificed after 8 h. Livers were

dissected and processed for qPCR analysis of *Fpn(+IRE)* (A) and *Fpn(-IRE)* (B) mRNAs. (C-E)

All mice were injected with 1 µg/g LPS. Half of the animals were subsequently injected with

saline, and the other half with 2.5 µg/g synthetic hepcidin every two hours for a total of 4

injections. At the endpoint the mice were sacrificed. Sera were collected by cardiac puncture and

analyzed for: (C) iron, (D) transferrin saturation, and (E) NTBI. The dotted line in (C) and (D)

indicates baseline serum iron and transferrin saturation, respectively, of control wt mice on SD.

Data are presented as (A-B) geometric mean±SD or (C-E) mean±SEM. Statistically significant

differences (p<0.05) compared to values from saline-, LPS- or hepcidin-treated wt control mice

on SD are indicated by a, b or c, respectively.

Table S1. List of primers used for qPCR.

Gene	GenBank accession	Forward primer sequence	Reverse primer sequence
Hamp1	NM_032541.1	*AAGCAGGGCAGACATTGCGAT*	*CAGGATGTGGCTCTAGGCTATGT*
Fpn(+IRE)	NM_016917.2	*GGCATAAGGCTGTTGTGCTT*	*TCATGACACCAGGCGTTCTC*
Fpn(-IRE)	XM_006496137.4	*GCCGGTTGGAGTTTCAATGT*	*TCATGACACCAGGCGTTCTC*
Dmt1	NM_001146161.1	*CTTGGGATACTGACGGTGAC*	*GATTTGCAGTCTGGAGCAGT*
Zip14	NM_001135151.1	*TGGATAGTGAGGCTGCGTGG*	*ATGGTGAGGCCAAGGCTAAT*
Lcn2	NM_008491	*GGCCAGTTCACTCTGGGAAA*	*TGGCGAACTGGTTGTAGTCC*
Tfrc	NM_011638.4	*AGCCAGATCAGCATTCTCTAACT*	*GCCTTCATGTTATTGTCGGCAT*
Bmp6	NM_007556.2	*ACTCGGGATGGACTCCACGTCA*	*CACCATGAAGGGCTGCTTGTCG*
Id1	NM_010495.2	*GGTACTTGGTCTGTCGGAGC*	*GCAGGTCCCTGATGTAGTCG*
Socs3	NM_007707.3	*TGCGCCTCAAGACCTTCAG*	*GCTCCAGTAGAATCCGCTCTC*
Fth1	NM_010239.2	*AAGTGCGCCAGAACTACCAC*	*AGCCACATCATCTCGGTCAA*
Rpl19	NM_009078.2	*AGGCATATGGGCATAGGGAAGAG*	*TTGACCTTCAGGTACAGGCTGTG*
Actb	NM_007393.3	*GACGACATGGAGAAGATCTG*	*GTGAAGCTGTAGCCACGCTC*

Supplemental Figure legends

Fig. S1. Dietary iron loading does not disrupt inflammatory hepcidin induction in LPS-treated wild type mice but blunts hepcidin-mediated hypoferremia. Nine-week-old male mice (n=12-14 per group) were fed SD or HID for one day, one week, or five weeks prior to sacrifice. Half of the mice were injected intraperitoneally with saline and the other half with 1 µg/g LPS 4 hours before sacrifice. Sera were collected by cardiac puncture and analyzed for: (A) iron, (B) transferrin saturation, (C) ferritin, and (D) hepcidin. Livers were dissected and processed for biochemical analysis of: (E) liver iron content (LIC) by the ferrozine assay and (F) *Hamp* mRNA by qPCR. The dotted line in (A) and (B) indicates baseline serum iron and transferrin saturation, respectively, of control mice on SD. Data (A-E) are presented as the mean±SEM and in (F) as geometric mean±SD. Statistically significant differences (p<0.05) over time compared to values from saline- or LPS-treated control mice are indicated by a or b, respectively.

Fig. S2. Effects of dietary iron manipulations in splenic iron of wt and Hjv-/- mice. Spleen sections from mice described in Fig. 1 were analyzed histologically for iron deposits by Perls staining (magnification: 10x).

Fig. S3. Low magnification immunohistochemical images of ferroportin in liver sections of dietary iron-manipulated wild type and Hjv-/- mice following LPS treatment. Liver sections

775 from mice described in Fig. 1 were used for immunohistochemical analysis of ferroportin

776 (magnifications: 10x and 5x).

778 **Fig. S4. Low magnification immunohistochemical images of ferroportin in spleen sections**

779 **of dietary iron-manipulated wild type and Hjv-/- mice following LPS treatment.** Spleen

780 sections from mice described in Fig. 1 were used for immunohistochemical analysis of

781 ferroportin (magnification: 2x).

783 **Fig. S5. Perls staining for iron deposits in liver and spleen sections of dietary iron-**

784 **manipulated wild type and Hjv-/- mice following treatment with synthetic hepcidin.** Liver

785 and spleen sections from mice described in Fig. 3 were stained with Perls Prussian blue

786 (magnification: 10x).

788 **Fig. S6. Effects of LPS treatment on expression of mRNAs encoding iron transport proteins**

789 **and signaling endpoints in the liver of dietary iron-manipulated wild type and Hjv-/- mice.**

790 Livers from mice described in Fig. 3 were dissected and processed for qPCR analysis of mRNAs

791 encoding iron transport proteins and signaling endpoints. (A) *Fpn(+IRE)*, (B) *Socs3*, (C) *Id1*, (D)

792 *Bmp6*, (E) *Dmt1*, (F) *Zip14*, (G) *Lcn2* and (H) *Tfrc*. All data are presented as the geometric mean

793 ± SD. Statistically significant differences ($p<0.05$) compared to values from saline- or hepcidin-

794 treated control mice are indicated by a or b, respectively.

Fig. S7. Low magnification immunohistochemical images of ferroportin in liver sections of dietary iron-manipulated wild type and Hjv-/- mice following treatment with synthetic hepcidin. Liver sections from mice described in Fig. 3 were used for immunohistochemical analysis of ferroportin (magnifications: 10x and 5x).

Fig. S8. Low magnification immunohistochemical images of ferroportin in spleen sections of dietary iron-manipulated wild type and Hjv-/- mice following treatment with synthetic hepcidin. Spleen sections from mice described in Fig. 3 were used for immunohistochemical analysis of ferroportin (magnification: 2x).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. S1

Fig. S2

	wt		Hjv-/-	
	SD	HID	SD	IDD

+LPS (10x)

+LPS (5x)

Fig. S3

	wt		Hjv-/-	
	SD	HID	SD	IDD

+LPS

2x

Fig. S4

A   wt   Hjv-/-

Fig. S5

Fig. S6

Fig. S7

	wt		Hjv-/-	
	SD	HID	SD	IDD

+Hep

2x

Fig. S8