FINGER REPRESENTATIONS IN PRIMARY SOMATOSENSORY CORTEX ARE
MODULATED DURING VIBROTACTILE WORKING MEMORY

AUTHORS

Finn Rabe[1], Sanne Kikkert[1,2*], Nicole Wenderoth[1*]

AFFILIATIONS

[1]Neural Control of Movement Laboratory, Department of Health Sciences and Technology,
ETH Zürich, Zürich, Switzerland.

[2]Spinal Cord Injury Center, Balgrist University Hospital, University of Zürich, Zürich,
Switzerland.

* Both authors contributed equally

CORRESPONDING AUTHOR

rabef@ethz.ch & nicole.wenderoth@hest.ethz.ch

ABSTRACT

It is well-established that vibrotactile stimulations elicit Blood-oxygen-level-dependent
(BOLD) responses in somatotopically organized brain regions. Whether these somatotopic
maps are modulated by working memory (WM) is still unknown. In our WM experiment,
participants had to compare frequencies that were separated by a delay period. Vibrotactile
stimuli were sequentially applied to either their right index or little finger. Using functional
MRI, we investigated whether vibrotactile WM modulated neural activity in primary
somatosensory (S1), an area that is known to contain individual finger representations. Our
mass-univariate results revealed the well-described network of brain regions involved in WM.
Interestingly, our mass-univariate results did not demonstrate S1 to be part of this network.
However, when we parametrically modulated the time-binned regressors in our GLM we
found that the delay activity in S1 and secondary somatosensory cortex (S2) was reflected in
a U-shaped manner. Using multi-voxel pattern analysis (MVPA), an analysis technique that is

more sensitive to subtle activity differences, we found finger-specific patterns of activation in

the S1 hand area during the WM delay period. These results indicate that processes

underlying WM modulate finger-specific representations during our discrimination task.

1 INTRODUCTION

Topographic representations such as the somatotopic map in the somatosensory cortex
have been shown to be ubiquitous, to a certain extent, in the cerebral cortex of mammals.
They consist of orderly representations of receptor surfaces and are present in most of the
neocortex (Kaas, 1993, 1997; PENFIELD & BOLDREY, 1937; Silver & Kastner, 2009). These
somatotopic maps are incredibly specific to the point where the sensation of each finger can
be assigned to its own contralateral cortical region, so called finger representations (Besle et
al., 2013; Cunningham et al., 2013; Martuzzi et al., 2014; Sanchez Panchuelo et al., 2018;
Sanders et al., 2019).

Importantly, the selective activation of these finger representations is not exclusively
modulated by tactile stimulation or movement, but also through other mechanisms like (i)
through *attempted movements*, without producing overt motor output (i.e., without any tactile
or proprioceptive feedback) (Ariani et al., 2021; Guan et al., 2021; Kikkert et al., 2021) (ii)
observed touch (Kuehn et al., 2018), (iii) or directing *attention* to only one of multiple fingers
receiving tactile stimulation (Puckett et al., 2017).

Another mechanism that may modulate S1 activity is working memory (WM). In the
absence of sensory stimuli, decisions are made with information stored in WM (Christophel
et al., 2017). Support for S1 involvement in tactile WM stems from neurophysiological
research in non-human primates. Single-unit activity was recorded from the S1 hand area
while subjects had to match an object with a specific surface to previous presented surface
stimulus (Y. D. Zhou & Fuster, 1996; Y.-D. Zhou & Fuster, 2000). The authors observed cells
that were not only involved in perception of tactile stimuli, but also sustained their firing during
the WM delay period. This suggests that primary sensory cortices could serve as a memory
buffer for stimulus information (D'Esposito & Postle, 2015), an idea that has been
conceptualized in 'sensory recruitment' models of WM (Katus et al., 2015; Pasternak &
Greenlee, 2005). According to this model, WM is maintained in the same brain regions that
are also involved in sensory stimuli encoding. However, human neuroimaging studies on
vibrotactile WM do not suggest a selective modulation of activity in S1 during the delay period
of information storage (Christophel et al., 2017; Preuschhof et al., 2006). These studies
generally employed a mass-univariate approach, which is often not sensitive enough to reveal
(i) temporal changes in delay period activity and (ii) specific features of WM content.
Concerning the former, information retention in WM is not always reflected in constant delay
activity, especially when a temporal structure of the delay period can be anticipated (Rose et
al., 2016). Furthermore, it has been shown that persistent delay activity might not be crucial
to retain WM content (Stokes, 2015). One approach to more selectively identify brain regions

associated with inconsistent delay activity could be through using model predictors that are
graded according to the anticipated delay activity (Braver et al., 1997).

In addition, WM studies employing a mass-univariate approach might often not be
sensitive enough to patterns of WM content (Weaverdyck et al., 2020). Instead, multi-voxel
pattern analysis (MVPA) has the ability to detect stimulus information in spatially distributed
patterns of activation in a region of interest (ROI). In our case, somatotopic maps can be
investigated with high sensitivity through combining functional MRI (fMRI) and MVPA. Here,
above chance level classification accuracies indicate that brain activation patterns
representing single fingers in S1 are highly specific, in accordance with the idea of narrow
tuning curves in the S1 neuronal population (Detorakis & Rougier, 2014). Lower classification
accuracies, by contrast, indicate that S1 activity is less finger specific suggesting that tuning
curves are broader. Importantly, MVPA accuracies can reveal representational information
even if the average Blood-oxygen-level-dependent (BOLD) response does not surpass a
certain statistical threshold in univariate analyses.

Neuroimaging studies utilizing MVPA found that features like vibratory frequencies
were solely represented during the delay period in associative brain regions, i.e., posterior
parietal and frontal regions (Schmidt et al., 2017; Wu et al., 2018). Interestingly, features of
texture stimuli could be decoded from S1 delay period activity (Schmidt & Blankenburg,
2018). Similar to the aforementioned observations in non-human primates, decoded textures
were located in the contralateral S1 hand area, an area which has shown to possess fine-
grained finger representations (Besle et al., 2013; Cunningham et al., 2013; Martuzzi et al.,
2014; Sanchez Panchuelo et al., 2018; Sanders et al., 2019). This suggests that neural activity
in S1 during the delay period may not only reflect some stimulus feature, but also a resides
within somatotopically organized cortical areas.

Here we used fMRI combined with MVPA to explore whether finger representations
in contralateral S1 are selectively modulated through WM maintenance, i.e., in the absence
of any tactile input. Participants were asked to perform a vibrotactile frequency discrimination
task on the index or little finger while we collected fMRI data. We hypothesized that if WM
selectively activates S1, then we will observe finger selective activation patterns during the
WM delay period.

METHODS

PARTICIPANTS

Thirty young healthy volunteers (19 females; mean age= 24.48, SEM= 0.44) participated in
our study. Our sample size was comparable to those in previous reports on fMRI decoding
of WM content using discrimination tasks (Ester et al., 2009; Schmidt et al., 2017). All
participants were neurologically intact and reported to be right-handed. All of them gave
written informed consent and the study protocol was approved by the local ethics committee
(BASEC-Nr. 2018-01078). Three participants had to be excluded due to excessive head
motion based on our criterion (see 'Preprocessing of fMRI data' section for more detail).

EXPERIMENTAL PROCEDURE AND TASKS

TACTILE STIMULI

Vibrotactile stimuli (duration = 2s, sampling rate = 1kHz) were applied to the right index or
right little finger using a MR-compatible piezoelectric device (PTS-T1, Dancer Design, UK).
We selected these fingers as they have the largest inter-finger somatotopic distance (Besle
et al., 2013; Ejaz et al., 2015; Kolasinski et al., 2016; Sanders et al., 2019), allowing us to
robustly detect the modulation of somatotopic representations by the WM task. The one bin
piezoelectric wafers were mounted to the fingertips using custom 3D-printed retainers that
were fixed with a Velcro strap. Participants were asked to report any tingling sensation in
case the retainer was mounted too tightly. The stimulation consisted of mechanical sinusoids
that were transmitted from the testing computer to the piezoelectric device using a C Series
Voltage Output Module (National Instruments) and the in-house NI-DAQmx driver.

SENSORY DETECTION THRESHOLD ESTIMATION

To ensure similar task difficulty across runs of the main experiment (Harris et al., 2006), we
determined the sensory detection threshold (SDT) for both fingers prior to starting the main
experiment. SDT was defined as the stimulation intensity at which the participants detected
the stimulus 50% of the time. We stimulated each finger only once per trial at base frequency
(20 Hz) and participants were asked to press a button upon detection of a stimulation. To
reliably estimate SDT, we applied a conventional Bayesian-based Quest procedure
(QuestHandler in PsychoPy). After each detected or undetected stimulus the algorithm
searched for the most probable psychometric function via maximum likelihood estimation
over the course of 25 trials starting with a stimulation amplitude of 0.1 Volts (Watson & Pelli,
1983). The Weibull psychometric function was calculated using the following formula:
31
(1) $\Psi(x) = \delta\gamma + (1-\delta)[1-(1-\gamma)\exp(-10^{beta(x-T+\epsilon)})]$

where x is the stimulus intensity in Volts and T is the estimated sensory detection threshold.

This procedure was performed prior to the first run. If the percentage of correctly

discriminated memory trials in a run was below 60% or above 90%, then we redetermined

the SDT using a shortened version of the Quest procedure. In such a case we started the

Quest procedure with the previously determined stimulation intensity to reduce the number

of iterations (new iterations = 7 trials). This procedure was applied to keep task difficulty at

comparable levels throughout the experiment.

We analyzed changes in behavioral performance occurred across runs, potentially

due to perceptual learning, cooling of the fingertips, or fatiguing effects (**Fig. 2B**). A repeated

measures two-way ANOVA (2 fingers x 4 runs) did not show a significant difference in

behavioral performance across runs ($F(3, 2496) = 2.0$, $p = .11$, $\eta^2 < 0.01$), no significant

difference across fingers ($F(1, 2496) = .48$, $p = .49$, $\eta^2 < 0.01$), and no significant interaction

effect ($F(3, 2496) = .47$, $p = .47$, $\eta^2 < 0.01$).

MAIN EXPERIMENTAL TASK

The main experimental task was generated using PsychoPy (Peirce et al., 2019). The

experimental task consisted of memory and no memory trials. During a memory trial,

participants performed a two-alternative forced choice (2AFC) discrimination task. Two

vibrotactile stimuli were consecutively applied to the same finger (i.e., the index or the little

finger), separated by a jittered 6-8s delay. We targeted cutaneous mechanoreceptors that

respond to stimulations in the flutter range (Mountcastle et al., 1967). One of two stimuli

vibrated at 20 Hz (2s duration at SDT intensity) while the vibration frequency of the other

stimulus varied between 22, 24 or 26 Hz (same duration and intensity). Participants had to

indicate by means of a button press whether the first or the second stimulation was higher in

frequency (half of the participants) or whether the first or the second stimulation was lower in

frequency (the other half of the participants), following previously published procedures

(Pleger et al., 2006, 2008, 2009). Responses were recorded via index and middle finger button

presses of the other (left) hand using a MR-compatible fiber optic device. We randomized the

order how the response options (f1 and f2) appeared on the screen on a trial-by-trail basis to

prevent somatotopy-dependent anticipatory motor activity. After a 3s response period

participants received visual feedback (1s) indicating whether their response was correct

(highlighted by green color) or incorrect (red; see **Fig 1**). Participants were instructed to focus

their gaze on the fixation cross in the middle of the screen during the complete trial.

Vibrotactile stimuli trials targeted either the index or the little finger and which finger would

be stimulated per trial was counterbalanced across each run.

Fig 1. Vibrotactile frequency discrimination task. During memory trials (*top*) two vibrotactile stimuli

that differed in frequency were consecutively applied to the same finger (in this example the index

finger). Both stimulations were separated by a jittered delay period, during which participants had to

keep the first stimulus (f1) in memory in order to compare it to the subsequent one (f2). During the 3s

response period participants had to indicate by means of a left hand button press which of the two

stimulation frequencies (f1 and f2) was higher. The mapping between the discrimination response and

which button to press was randomized across trials. Subjects received feedback whether their

response was correct (*green*) or incorrect (*red*). The target finger (index or little finger) were intermixed

within a run and the inter trial interval (ITI) was jittered between 2-4s. During no memory trials (*bottom*),

vibrotactile stimulations and visual information remained the same. However, participants were

instructed not to focus on the stimulation and also not to compare the vibrotactile frequencies. They

simply had to press the button indicated by the arrow in the middle of the screen.

To disentangle WM processes from general responses to the stimuli, we also included

no memory trials. During a no memory trial participants received the exact same vibratory

stimulations as in during memory trials, but they were instructed not to focus on the stimuli

or on their vibration frequencies. During the response period subjects were informed by a

visual cue (pointing arrow) which button to press. To ensure participants did not switch

cognitive strategies, the indicated response was always contrary to the response that would

be expected when correctly discriminating both frequencies. Memory and no memory trials

conditions were separated in mini blocks of 4 trials. Participants were informed whether they

had to perform the memory or no memory task by means of a visual cue (1.5s) at the

beginning of each trial. Prior to the experiment, participants were familiarized with the memory and no memory tasks by completing 12 trials.

The order of stimulus sites (stimulated finger) was counterbalanced both within and across mini blocks. Stimulation frequencies were counterbalanced across the experiment. Each stimulus frequency was presented equally often in both memory and no memory condition. Jittered timings for Inter-stimulus-interval (ISI, 6-8s) and Inter-trial-interval (ITI, 2-4s) were randomly drawn from a uniform distribution. All participants completed 4 runs consisting of 48 trials each. Each run consisted of 6 memory and 6 no memory mini blocks in a counterbalanced order.

BEHAVIORAL ANALYSIS

We defined the discrimination accuracy per participant as the percentage of correctly discriminated trials separately for each condition. We expected that greater frequency differences would facilitate discrimination between both tactile vibrations while the stimulus site should have no effect. We therefore investigated whether behavioral performances differed across frequency differences and across fingers using a two-way repeated-measures ANOVA.

MRI DATA ACQUISITION

Functional as well as structural MRI images were acquired on a Philips Ingenia 3 Tesla MRI (Best, The Netherlands) using a 32-element head coil. fMRI data was collected using an echo-planar-imaging (EPI) sequence acquiring 36 transversal slices centred at the bicommissural line and with whole brain coverage, though excluding most of cerebellum (repetition time (TR): 2s, echo time (TE): 30ms, spatial resolution: $3mm^3$, FOV = 222 × $222mm^2$, 85° flip angle, slice orientation: transversal, SENSE factor (AP): 2, 472 functional volumes per run). Anatomical images were acquired during SDT estimation using a MPRAGE T1-weighted sequence (TR = 7.7ms, TE = 3.6ms, FOV = 240 × $240mm^2$, flip angle: 8°, resolution: $1mm^3$, number of slices: 160, slice thickness: 2.2mm, slice orientation: sagittal).

PREPROCESSING OF FMRI DATA

Conventional pre-processing steps for fMRI data were applied to each individual run in native three-dimensional space, using FSL's Expert Analysis Tool FEAT (v6.00; fsl.fmrib.ox.ac.uk/fsl/fslwiki). The following steps were included: Motion correction using MCFLIRT (Jenkinson, 2002), brain extraction using automated brain extraction tool BET

(Smith, 2002), high-pass filtering (100Hz), slice-time correction, and spatial smoothing using

a 3mm FWHM (full width at half maximum) Gaussian kernel using FEAT. Functional data was

aligned to structural images initially using FLIRT (Jenkinson & Smith, 2001), and optimised

using boundary based registration (Greve & Fischl, 2009). BOLD EPI data was assessed for

excessive motion using motion parameter estimates from MCFLIRT. If the functional data

from a participant showed greater than 1.5mm (half the voxel size) of absolute mean

displacement, this participant was excluded from all further analysis.

To reduce physiological noise artifacts, these CSF and white matter were used to

extract scan-wise time series which were then added to the model as nuisance regressors in

addition to the standard motion parameters.

Structural images were transformed to Montreal Neurological Institute (MNI) standard space

using nonlinear registration (FNIRT), and the resulting warp fields were applied to the

functional statistical images.

DEFINITION OF REGIONS OF INTEREST

We used each individual participant T1-weighted image to create a cortical surface

reconstruction by means of Freesurfer (Fischl et al., 1999). We identified regions of interest

(ROIs), specifically SI, anatomically for each subject based on the probabilistic Brodmann

area parcellation provided by Freesurfer (Fischl, 2012). More specifically, an S1hand ROI was

defined by combining Brodmann areas (BA) 1, 2, 3a, and 3b. We then converted this S1ROI

to volumetric space. Any holes were filled and non-zero voxels were mean dilated. Next, the

axial slices spanning 2cm medial/lateral to the hand area (Yousry et al., 1997) were identified

on the 2mm MNI standard brain (min-max MNI z-coordinates=40-62). This mask was non-

linearly transformed to each participant's native structural space. Finally, we used this mask

to restrict the S1ROI and extracted an S1hand area ROI. Similar ROI definition has been

previously used (Diedrichsen et al., 2013; Ejaz et al., 2015; Wiestler & Diedrichsen, 2013). The

S1hand area ROI was used to both extract time-binned estimates as well as to decode

information about the stimulus site during the delay period.

UNIVARIATE ANALYSIS

First-level parameter estimates were computed per run using a voxel-based general linear

model (GLM) based on the gamma hemodynamic response function. Time series statistical

analysis was carried out using FILM (FMRIB's Improved Linear Model) with local

autocorrelation correction.

To find neural correlates of WM we contrasted beta estimates from the delay period

during memory trials to those in no memory trials. We then used a fixed effects higher-level

analysis to average activity across runs for each individual participant. Finally, to make

inferences on the population level, we computed a mixed effects analysis (Flame 1). From

this we obtained statistical group maps (Z-statistic images) for each contrast of interest, e.g.

contrasting memory delay activity to no memory delay activity. Z-statistic images were

thresholded using clusters determined by $Z > 3.1$ and $p < .05$ family-wise-error-corrected

(FWE) cluster significance.

To further explore whether finger specific activity levels were maintained in a

somatotopic fashion, we first computed somatotopic ROIs by contrasting finger-specific

activity during the first stimulation. We did this by contrasting activity associated with right

index stimulations to right little finger stimulations, which elicited a finger specific map in the

lateral part of S1while the reverse contrast revealed more medially located activity (see

supplementary materials, Fig. 1). These S1activity maps were in line with previous findings

on finger somatotopy.

We then compared delay period activity levels between trials where either the index

or the little finger was stimulated within each finger ROI. Again, we computed a fixed-effects

analysis as mentioned before. We extracted the percent signal change (PSC) separately for

each participant using the Featquery tool in FSL.

Information retention in WM is not always reflected by constant delay activity,

especially when the duration of the delay period can be somewhat anticipated (Rose et al.,

2016). Indeed, persistent delay activity may not be crucial to retain WM content (Stokes,

2015). WM delay activity has been shown to decrease until shortly before memory retrieval

when the remembered stimulus information is reactivated as suggested by an increase of

neural oscillations in the theta band (Rose et al., 2016). We therefore hypothesized that the

BOLD activity level would vary in a U-shaped fashion over the delay period. To test this

hypothesis, we conducted a parametric modulation analysis. Our parametric modulation

regressor was modelled to predict activity in three consecutive time-bins of the delay period

delay period in a U-shaped manner (**Fig. 3A**). The length of each time-bin equalled one TR

(i.e., 2s). Since we jittered the delay period between 6 and 8 s, we only modelled the first

three time-bins of the delay period (2-6s). The remaining time of the delay was modelled as

a separate regressor of no interest. Z-statistic images were thresholded using clusters

determined by $Z > 3.1$ and a familywise error–corrected cluster significance threshold of $p <$

0.05 was applied to the suprathreshold clusters.

To further visualize the results of the parametric modulation, we extracted activity

estimates per time-bin. To do so, we modelled each time-bin of the delay period separately

in a voxel-based general linear model (GLM) based on the gamma hemodynamic response

function. The remaining time of the delay was modelled as a separate regressor of no interest.

We then extracted the z-scored estimates per time-bin within the previously defined S1hand

area ROI. All statistical maps were projected onto a cortical surface using Connectome's

Workbench (Marcus et al., 2011).

VARIANCE INFLATION FACTOR (VIF)

To test whether multicollinearity between the parameter estimates in our GLM was

reasonably low, we calculated the variance inflation factor (VIF). This represents how much

the variance of an individual regressor can be explained due to correlation to other regressors

in our model (Zuur et al., 2010). For each variable, VIF was computed by the following formula:

14
$$VIF = \frac{Var(E)}{Var(X)}$$

where *Var(E)* reflects the mean estimation variance of all regression weights (stimulation and

information storage regressors for each finger) while *Var(X)* reflects the mean estimation

variance in case all regressors would be estimated individually. A VIF of 1 indicates total

absence of collinearity between the regressor of interest and all other regressors in our GLM

while a large VIF signals a serious collinearity problem. There is no clear threshold for

acceptable multicollinearity. Previous literature however recommends that the VIF is ideally

smaller than 2.5 (Johnston et al., 2018). In our case, the VIF was on 1.45, averaging across

regressors reflecting stimulation one, delay and stimulation two. This indicates that we can

disentangle the stimulation and WM delay period activity with appropriate certainty. We

additionally computed the VIF for the regressors relating to the stimulation periods and the

time-binned delay periods (i.e. stimulation one, delay (0-2s), delay (2-4s), delay (4-6s) and

stimulation two). As expected, the resulting VIF was on average 11.38, signaling concerning

multicollinearity. Since the WM related activity within the time-bins is correlated, this result

was expected, but also means our findings of our time-binned analysis should be interpreted

with caution.

MULTIVARIATE PATTERN ANALYSIS (MVPA)

We used multivariate pattern analysis (MVPA) to decode which finger was stimulated based on activity during the delay period. This analysis was conducted for voxels within the S1 hand area mask that have been shown to possess fine-grained finger representations. First-level parameter estimates were computed for all events of each trial and each participant using a voxel-based general linear model (GLM) in SPM (v12) based on the gamma hemodynamic response function. This resulted in 192 beta estimates (48 trials x 4 runs) during the delay period per participant across both conditions. 96 beta estimates for each memory and no memory condition.

We trained a linear classifier (support vector machine, SVM) to predict which finger was stimulated in a specific trail based on the respective delay period activity using the nilearn toolbox (Abraham et al., 2014). We calculated classification accuracies using a leave-one-run-out cross-validation approach. The accuracies were averaged across folds, resulting in one accuracy per condition and per participant. To approximate the true chance level, we shuffled the condition labels using 1000 permutations (Ojala & Garriga, 2009), as implemented in the scikitlearn toolbox (Pedregosa, 2011). Then, we calculated permutation-based p-values based on the following formula:

$$\text{(2)} \quad (C + 1) / (n\_permutations + 1),$$

where C is the number of permutation scores that were higher than the true accuracy. To assess the statistical significance of each ROI's decoding accuracy, we used one sample t-tests against the approximated chance level (Stelzer et al., 2013). We further used a paired t-tests to compare classification accuracies between the WM and non-WM conditions.

Finally, to assess whether somatotopic representations of stimulus content determined participant's behavioral performance, we computed classification accuracies during both stimulation and the delay period for each run and the respective behavioral performance. Then we calculated a repeated-measures correlation coefficient (Bakdash & Marusich, 2017). For the purpose of exploration, we obtained coefficients for different time-points, during both stimulations as well as the delay period.

STATISTICAL DATA ANALYSIS

To detect outliers we implemented a state-of-the-art method using robustbase (Finger, 2010). $\mathbf{S_n}$ identifies an outlier (x_i) if the median distance of x_i from all other points, was greater than the outlier criterion ($\lambda=4$) times the median absolute distance of every point from every other point:

(1) $\frac{med_{j \neq i}|x_i - x_j|}{s_n} > \lambda \ where \ S_n = c_n \overset{med}{\underset{i=1:n}{}} \{\overset{med}{\underset{j \neq 1}{}}|x_i - x_j|\},$

where c_n is a bias correction factor for finite sample sizes (Rousseeuw & Croux, 1993). We

detected no outliers for behavioral data that had to be excluded from any further analysis.

Before conducting any ANOVA testing, we validated the assumptions for homogeneity of

variances and normal distributions using a Levene's and Shapiro-Wilk test. Effect sizes of

different variables were measured using eta squared. ANOVA analysis was done using the

bioinfo-kit and pingouin toolbox (Huang et al., 2014; Vallat, 2018).

When computing t-statistics, in order to correct for multiple comparisons we applied

a false discovery rate (FDR) method (Benjamini & Hochberg, 1995).

RESULTS

BEHAVIORAL PERFORMANCE WAS BETTER WITH GREATER FREQUENCY DIFFERENCES

A two-way ANOVA showed that behavioral performances differed significantly

between frequency differences ($F(2, 156) = 5.06$, $p < .001$, $\eta^2 = 0.06$); **Fig. 2A**), but not

between stimulated fingers ($F(1, 156) = .44$, $p = .51$, $\eta^2 < 0.01$). The frequency effect was not

significantly different between stimulated fingers (i.e., no interaction effect; $F(1, 156) =$

$.88$, $p = .42$, $\eta^2 = 0.01$). A post hoc test (Tuckey's HSD) on pairwise comparisons on frequency

differences pairs revealed that discrimination accuracy was significantly different between 2

and 6 Hz differences ($q = 4.48$, $p < .01$) and showed no significant difference for the rest of

the pairs (4 Hz vs 6 Hz: $q = 2.67$, $p = .15$ and 2 Hz vs 4 Hz $q = 1.81$, $p = .41$).

VIBROTACTILE WORKING MEMORY RECRUITS A DISTRIBUTED BRAIN NETWORK

We first determined brain areas that were more activate during the delay period in the
memory compared to the no memory condition. We found that WM processing involved a
distributed brain network: i.e., bilateral frontal lobe, bilateral (medial/inferior) frontal gyrus
(MFG, IFG), bilateral pre-motor cortex and supplementary motor area (PMC, SMA), bilateral
inferior parietal lobule (IPL), bilateral superior parietal lobule (SPL), bilateral supramarginal
gyrus (SMG), bilateral caudate, bilateral thalamus, bilateral nucleus accumbens, and bilateral
insula (**Fig. 3A** and **Table 1**). As in previous human fMRI studies, this univariate analysis did
not reveal any S1 activity.
Furthermore, we contrasted activity levels during vibrotactile stimulation between
fingers (i.e., index>little and little>index finger), and, as expected, observed separated finger
representations in contralateral S1(**Fig. 3B,** *top*). More specifically, there was stronger index
compared to little finger activity on the lateral part of S1, while there was stronger little
compared to index finger activity medial of S1. These S1 activity maps are in line with
previous findings on finger somatotopy which is highly specific within S1 (Besle et al., 2013;
Kikkert et al., 2021; Kolasinski et al., 2016; Martuzzi et al., 2014; Sanchez Panchuelo et al.,
2018; Sanders et al., 2019). Finally, we assessed whether finger-specific activity was
sustained within these finger-specific maps during the delay period. We did not find any
significant selective activity in these finger maps during the delay period (index > rest: $t = .6$,
$p = .52$, little finger > rest: $t = 1.5$, $p = .14$) (**Fig. 3B,** *bottom*).

Fig. 3. Univariate group results. A. We determined brain regions that were more activate during the

delay period of the memory compared to the no memory condition. A statistical map (Z > 3.1) was

obtained by contrasting delay period activity in memory trials to no memory trials. The map was

separately projected onto a cortical surface contralateral (*top*) and ipsilateral (*bottom*) to the stimulus

site. WM related activity resided in a network of brain regions (for more details, see **Table 1**). **B.** S1

areas activated during index (red) and little (blue) finger stimulation (*top*). Brain regions activated during

index and little finger stimulation Z-statistic images were thresholded using clusters determined by Z

> 3.1, p < .05 family-wise-error-corrected (FWE) cluster significance and were projected onto a cortical

surface. Finger maps were located in contralateral SI. We then extracted the percent signal change

(PSC) within these finger-specific S1 areas during the delay period by contrasting activity during each

finger stimulation to rest (bottom). The PSC was only extracted from the pre-defined finger maps in

left SI. Bar plots are centered at the mean and error bars reflect the standard deviation.

Anatomical region	Peak MNI Coordinates			Mean Fisher Z	U-shaped activity
	X	Y	Z		
Left Frontal Pole	-42	52	2	4.33	✓
Right Frontal Pole	40	46	32	3.86	✓
Left IFG	-52	16	2	5.14	✓
Right IFG	58	10	18	4.59	✓
Left MFG	-44	30	34	4.45	
Left PMC (+SMA)	-4	8	50	5.49	✓

Right PMC (+SMA)	8	20	46	5.78	✓
Left IPS	-42	-52	48	5.14	
Left IPL	-56	-20	28	4.78	✓
Right IPL	46	-48	52	4.86	✓
Left SMG	-42	-46	40	4.77	
Right SMG	51	-32	46	4.16	
Left SPL	-50	-48	58	4.64	
Left Insular cortex	-30	24	6	5.61	✓
Right Insular cortex	36	22	-2	5.41	✓
Left Nucleus Accumbens	-12	14	-6	5.27	
Right Nucleus Accumbens	10	12	-4	4.16	
Left Putamen	-18	6	-10	4.96	✓
Right Putamen	20	10	-8	4.34	✓
Left Caudate	-12	18	0	4.80	✓
Right Caudate	14	18	4	4.97	✓
Left Thalamus	-8	-24	-16	3.86	✓
Right Thalamus	2	-20	-12	3.91	✓

Table 1. Identified brain regions in which the local activity during the delay period reflected tactile WM processing. Sorted from anterior to posterior and from cortical to subcortical. A statistical map was obtained by contrasting BOLD responses in memory trials to no memory trials during the delay period. In a next step we parametrically modulated the delay activity, assuming a U-shaped activity during the delay period. Brain regions overlapping with the initial results are indicated by a check mark in the right column. All z-statistic images were thresholded using clusters determined by $Z > 3.1$ and $p < .05$ family-wise-error-corrected (FWE) cluster significance. Mean Fischer Z indicates peak z-values. Areas were labeled according to the Juelich Histological Atlas and Havard-Oxford (Sub-)cortcial Structural Atlas (Eickhoff et al., 2005). MFG= medial frontal gyrus IFG = inferior frontal gyrus, PMC = pre-motor cortex, SMA = supplementary motor area, IPL = inferior parietal lobule, SPL = superior parietal lobule, SMG = supramarginal gyrus.

RAMP-UP DELAY PERIOD ACTIVITY IN CONTRALATERAL S1

We then examined how brain activity temporally changed during the delay period. We parametrically modulated the delay period regressors by the hypothesized U-shaped activity changes and computed the associated statistical maps. A U-shaped activity modulation was found in a similar network of brain regions as in **Fig. 4A**. Interestingly, this modulation also elicited activity maps in S1 and secondary somatosensory cortex (SII). For visualization

purposes only, we extracted the activity at each time bin (i.e., from a separate GLM not involving a parametric modulation regressor) within the contralateral S1 hand area to demonstrate U-shaped activity across time bins (**Fig. 4B**).

Fig. 4. U-shaped parametric modulation of brain activity during the delay period. A Brain regions exhibiting u-shaped modulated delay activity patterns during the delay period (2-6s) (see insert at the top reflecting the parametric modulator entered into the GLM). The area highlighted in blue represents the S1 hand area. **B.** For visualization purposes only, each time point between 2 and 6s of the delay period (each time bin corresponding to one TR) was modeled as a separate regressor in an univariate GLM. Z scores of the beta estimates were extracted from the S1 hand area for each time bin of the delay period and averaged across participants. Blue shading reflects the 95% confidence interval.

FINGER-SPECIFIC S1 WM REPRESENTATIONS

We hypothesized that executing a WM task would modulate finger specific representations in S1. While univariate analysis may not allow us to find WM-related activity in S1 during the delay period, MVPA is more sensitive to detecting subtle differences in activity patterns (Weaverdyck et al., 2020), making it an interesting approach to further examine our hypotheses. We decoded the stimulated finger (e., index versus little finger) during stimulation 1(f1), during the delay period, and during stimulation 2 (f2) separately for memory and no memory trials (**Fig. 5**). We did this in contralateral S1, which has shown to possess fine-grained finger representations (Besle et al., 2013; Cunningham et al., 2013; Martuzzi et al., 2014; Sanchez Panchuelo et al., 2018; Sanders et al., 2019), and a control white matter ROI.

During the both stimulations (f1 and f2) of memory trials, classification accuracies in contralateral S1 hand area were significantly higher than chance level (f1: $t = 14.25$, $p < .001$,

f2: t = 13.79, p < .001) which was also the case for during no memory trials (f1: t = 11.81, p < .001, f2: t = 10.76, p < .001).

Classification accuracies differed significantly between timepoints (two-way ANOVA; F(2, 156) = 103.7, p < .001, η^2 = .57), and conditions (F(1, 156) = 12.14, p < .01, η^2 = .07). The time effect was not significantly different between conditions (i.e., no interaction; F(2, 156) = .11, p = .9, η^2 < 0.01). More specifically, during both stimulations (f1, f2) classification accuracies inferred from the S1 hand area were significantly higher during memory trials compared to no memory trials (f1: t = 4.28, p < .001; f2: t = 4.07, p < .001).

Classification accuracies derived from the control ROI (White matter mask) did not show any significant differences between timepoints (F(2, 156) = 1.84, p = .16, η^2 = .02) nor conditions (F(1, 156) = 2.11, p = p = .15, η^2 = .01), however exhibited an interaction effect (F(2, 156) = 4.8, p = p < .05, η^2 = .06).

During the WM delay period classification accuracy in the S1 hand area was significantly higher than chance level (t = 4.55, p < .001) which was not the case for the no memory delay period (t = 1.54, p = .14). Moreover, classification accuracies significantly varied between memory conditions in the S1 hand area (t = 2.91, p < .01). This was not the case in our control ROI where classification accuracies did not differ from chance level for both conditions (memory: t = .04, p = .97; no memory: t = -.36, p = .72) nor between the memory and no memory conditions (t = .02, p = .98). These results suggest that S1, which possess especially fine-grained finger somatotopy, stores information about the stimulus site during WM delay and in the absence of any tactile input.

Fig. 5. MVPA results on somatotopic modulations. We extracted activity patterns to predict what finger (index vs. little finger) information was kept in WM during different timepoints (f1, delay and f2) and conditions (memory vs. no memory). To approximate the true chance level we computed 1000

DISCUSSION

In the present study, we investigated S1 activity during a vibrotactile WM task through the lens of somatotopic representations. More specifically, we explored whether finger specific information in S1 is modulated by cognitive processes relevant for performing a WM vibrotactile discrimination task. Even though BOLD activity in S1 did not differ from rest, we could show more specific representational tuning when participants had to retain task-relevant information. Using MVPA we were able to decode for which finger information was kept in WM during the delay period, demonstrating that finger-specific activity patterns resided in the contralateral S1 hand area in the absence of any tactile stimulation. Importantly, finger-specific classification accuracies during the delay period were not merely driven by residue activity from the preceding tactile stimulation. Indeed, classification accuracies were not higher than chance in a no memory condition in which participants received identical stimulation as in the memory condition, but were not required to keep any tactile information in WM. Lower classification accuracies during no memory trials indicates that S1 activity was less finger specific suggesting that tuning curves were broader.

SOMATOTOPIC MODULATION IN S1 IS MAINTAINED DURING WM DELAY PERIOD

This modulation of finger representation in contralateral S1 appears to also occur during WM. Pasternak and Greenlee (2005) analyzed a wide range of non-human primate studies and found that sensory WM recruits a wide circuitry of brain areas including sensory brain regions, which is in line with our observations. Neurophysiological research in non-human primates have shown that S1 cells are not only involved in perception of tactile stimuli, but also sustain their firing during the WM delay period (Katus et al., 2015; Y. D. Zhou & Fuster, 1996; Y.-D. Zhou & Fuster, 2000). This suggests that sensory brain regions not only process sensory stimuli, but also retain and manipulate sensory information in the absence of any sensory input, also known as 'sensory recruitment' models. According to these models, vibrotactile

information is processed in S1, relayed to S2, which possesses a reciprocal connection with

S1 and disseminates information via insular projections to frontal and parietal regions (Wang

et al., 2013). This account is in line with our parametric mapping results. We found a whole

network to be active during WM retrieval including somatosensory and fronto-partial, insular

cortices as well as subcortical regions.

Furthermore, previous findings indicate that tactile WM operates within

somatotopically organized sensory brain regions (Harris et al., 2001; Katus et al., 2015;

Schmidt & Blankenburg, 2018). However, till date, it has remained unknow whether WM

modulates S1 in a somatotopic fashion. Our results indicate that finger-specific information

is retained during the delay period despite the absence of any tactile input, suggesting WM

content representation in contralateral S1. No such finger-specific information content was

observed during the delay period in the no memory control condition.

It is uncertain which specific cognitive mechanism might drive our result. Indeed, does

finger-specific representational tuning during the delay period reflect the maintenance of the

stimulus features in WM or is it driven by other cognitive processes? Attention is an integral

part of performing WM tasks (Cowan et al., 2013; Logie & Cowan, 2015; Schmidt &

Blankenburg, 2018, 2019). By directing attention, the brain controls which retained

information is in a state of availability and one important aspect of WM is to make information

temporally accessible to cognitive control instead of merely acting as an information buffer

system. With regard to representational tuning in S1, it has been shown that when a spot on

the body surface (e.g. fingertips) was being attended to, preparatory tuning of corresponding

area in S1 was observed (Nelson et al., 2004; Puckett et al., 2017; Roland, 1981; Sterr et al.,

2007). Similarly in our experiment, when participants were instructed to ignore vibrotactile

stimuli during no memory trials, no representational tuning was observed even though the

tactile stimuli were identical. Our findings further highlight that activity in S1 and S2

contralateral to the stimulated finger exhibits a U-shaped activity profile across the delay

period, which might provide a glimpse into the temporal modulation of a putative attentional

mechanism. Interestingly, S1 and S2 were not the only areas that exhibited a U-shaped

profile. BOLD responses in fronto-parietal and subcortical regions (i.e. caudate and thalamus)

also appeared to reflect this U-shaped activity profile. Task-relevant modulation could occur

through prefrontal cortex (PFC) by gating the flow of vibrotactile information to S1. Indeed,

the prefrontal-thalamic inhibitory system could play a key role in boosting relevant stimulus

information while reducing irrelevant ones (Staines et al., 2002). It is likely that the final

increase in delay activity might be driven by anticipation of increasing attentional demands,

although we jittered the length of the delay period to prevent for anticipatory activity (Rose et

al., 2016). Therefore, we cannot rule out that attentional mechanisms are driving our results. As mentioned before, WM and attention are thought to be closely intertwined (Cowan et al., 2013; Logie & Cowan, 2015) and fMRI studies might not be the appropriate approach to test these subtle differences on a synaptic level (Stokes, 2015).

DISENTANGELING WM AND PERCEPTION

Our results suggest that our S1 decoding results reflect representational tuning that did not merely arise from carry-over effects caused by stimulus encoding. Our memory delay period was relatively short (6-8s) compared to other fMRI studies investigating vibrotactile WM that often employ delays of 12s to avoid any carry-over effects (Schmidt & Blankenburg, 2018; Wu et al., 2018). However, it is unlikely that the WM related somatotopic activity patterns observed in S1 were merely a residue of stimulus evoked activity due to the sluggishness of the BOLD response. If we merely decoded a residue of the tactile stimulation per se, then we would expect similar decoding results for memory and no memory conditions during the delay period. Indeed, finger stimulations were identical for the memory and no memory conditions. The only difference between these conditions was the cognitive strategy participants employed. While decoding accuracy was higher than chance in the delay period for the memory condition, this was not the case for the no memory condition. Furthermore, we found a significant difference in decoding accuracies between conditions. Moreover, the low VIF of the regressors suggested that activity related to stimulus perception and WM storage could be disentangled by our model.

Together our results show that S1 plays a role in vibrotactile frequency discrimination. Recently, two transcranial magnetic stimulation (TMS) studies found that behavioral vibrotactile frequency discrimination performance was impaired if TMS pulses were applied to either contralateral S1 during the delay period (Harris et al., 2002). This hints at the idea that the involvement of S1 and modulation of its somatotopic maps might be crucial to discriminate between tactile stimuli perceived on different fingers.

CONCLUSION

Tactile decision making based on mentally represented information is an essential part of higher cognitive processes. In the absence of sensory stimuli, decisions are made with information stored in WM (Christophel et al., 2017). A more profound insight into how the brain stores information in WM is essential to understand human cognition. This includes not

only the involvement of sensory regions in WM but also how their topographic organization might be modulated for information storage. Together, our results show that vibrotactile WM (i) recruits a distributed network of brain regions, (ii) modulates delay activity in various brain regions and (iii) modulates somatotopic finger representations in contralateral S1. These insights aid our understanding on how cognitive processes involved in WM shape cortical sensory activation with potential implications for individuals with WM impairments.

ACHKNOLEDGMENTS

We thank Alain Plüss and Bianca Badii for help with data collection. This work was supported by the Swiss National Science Foundation (320030_175616) and by the National Research Foundation, Prime Minister's Office, Singapore under its Campus for Research Excellence and Technological Enterprise (CREATE) program (FHT). SK was supported by the ETH Zürich Postdoctoral Fellowship Program.

REFERENCES

Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., Gramfort, A., Thirion, B., & Varoquaux, G. (2014). Machine learning for neuroimaging with scikit-learn. *Frontiers in Neuroinformatics*, *8*. https://doi.org/10.3389/fninf.2014.00014

Ariani, P., Pruszynski, J. A., & Diedrichsen, J. (2021). Motor planning brings human primary somatosensory cortex into action-specific preparatory states. *BioRxiv*.

Bakdash, J. Z., & Marusich, L. R. (2017). Repeated Measures Correlation. *Frontiers in Psychology*, *8*. https://doi.org/10.3389/fpsyg.2017.00456

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1). https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Besle, J., Sánchez-Panchuelo, R.-M., Bowtell, R., Francis, S., & Schluppeck, D. (2013). Single-subject fMRI mapping at 7 T of the representation of fingertips in S1: a comparison of event-related and phase-encoding designs. *Journal of Neurophysiology*, *109*(9). https://doi.org/10.1152/jn.00499.2012

Braver, T. S., Cohen, J. D., Nystrom, L. E., Jonides, J., Smith, E. E., & Noll, D. C. (1997). A Parametric Study of Prefrontal Cortex Involvement in Human Working Memory. *NeuroImage*, *5*(1). https://doi.org/10.1006/nimg.1996.0247

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J.-D. (2017). The Distributed Nature of Working Memory. *Trends in Cognitive Sciences*, *21*(2). https://doi.org/10.1016/j.tics.2016.12.007

Cowan, N., Blume, C. L., & Saults, J. S. (2013). Attention to attributes and objects in working memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *39*(3). https://doi.org/10.1037/a0029687

Cunningham, D. A., Machado, A., Yue, G. H., Carey, J. R., & Plow, E. B. (2013). Functional somatotopy revealed across multiple cortical regions using a model of

complex motor task. *Brain Research*, *1531*.
https://doi.org/10.1016/j.brainres.2013.07.050

D'Esposito, M., & Postle, B. R. (2015). The Cognitive Neuroscience of Working
Memory. *Annual Review of Psychology*, *66*(1). https://doi.org/10.1146/annurev-
psych-010814-015031

Detorakis, G. Is., & Rougier, N. P. (2014). Structure of receptive fields in a
computational model of area 3b of primary sensory cortex. *Frontiers in
Computational Neuroscience*, *8*. https://doi.org/10.3389/fncom.2014.00076

Diedrichsen, J., Wiestler, T., & Ejaz, N. (2013). A multivariate method to determine the
dimensionality of neural representation from population activity. *NeuroImage*, *76*.
https://doi.org/10.1016/j.neuroimage.2013.02.062

Eickhoff, S. B., Stephan, K. E., Mohlberg, H., Grefkes, C., Fink, G. R., Amunts, K., &
Zilles, K. (2005). A new SPM toolbox for combining probabilistic
cytoarchitectonic maps and functional imaging data. *NeuroImage*, *25*(4).
https://doi.org/10.1016/j.neuroimage.2004.12.034

Ejaz, N., Hamada, M., & Diedrichsen, J. (2015). Hand use predicts the structure of
representations in sensorimotor cortex. *Nature Neuroscience*, *18*(7).
https://doi.org/10.1038/nn.4038

Ester, E. F., Serences, J. T., & Awh, E. (2009). Spatially Global Representations in
Human Primary Visual Cortex during Working Memory Maintenance. *Journal of
Neuroscience*, *29*(48). https://doi.org/10.1523/JNEUROSCI.4388-09.2009

Finger, R. (2010). Review of 'Robustbase' software for R. *Journal of Applied
Econometrics*, *25*(7). https://doi.org/10.1002/jae.1194

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2).
https://doi.org/10.1016/j.neuroimage.2012.01.021

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical Surface-Based Analysis.
NeuroImage, *9*(2). https://doi.org/10.1006/nimg.1998.0396

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using
boundary-based registration. *NeuroImage*, *48*(1), 63–72.
https://doi.org/10.1016/j.neuroimage.2009.06.060

Guan, C., Aflalo, T., Zhang, C. Y., Rosario, E. R., Pouratian, N., & Andersen, R. A.
(2021). Preserved motor representations after paralysis. *BioRxiv*.

Harris, J. A., Arabzadeh, E., Fairhall, A. L., Benito, C., & Diamond, M. E. (2006).
Factors Affecting Frequency Discrimination of Vibrotactile Stimuli: Implications
for Cortical Encoding. *PLoS ONE*, *1*(1).
https://doi.org/10.1371/journal.pone.0000100

Harris, J. A., Harris, I. M., & Diamond, M. E. (2001). The Topography of Tactile
Working Memory. *The Journal of Neuroscience*, *21*(20).
https://doi.org/10.1523/JNEUROSCI.21-20-08262.2001

Harris, J. A., Miniussi, C., Harris, I. M., & Diamond, M. E. (2002). Transient Storage of
a Tactile Memory Trace in Primary Somatosensory Cortex. *The Journal of
Neuroscience*, *22*(19). https://doi.org/10.1523/JNEUROSCI.22-19-08720.2002

Huang, Y., Wang, J. Y., Wei, X. M., & Hu, B. (2014). Bioinfo-Kit: A Sharing Software
Tool for Bioinformatics. *Applied Mechanics and Materials*, *472*.
https://doi.org/10.4028/www.scientific.net/AMM.472.466

Jenkinson, M. (2002). Improved Optimization for the Robust and Accurate Linear
Registration and Motion Correction of Brain Images. *NeuroImage*, *17*(2).
https://doi.org/10.1016/S1053-8119(02)91132-8

Jenkinson, M., & Smith, S. M. (2001). A global optimization method for robust affine
registration of brain images. *Medical Imaging Analysis*, *5*, 143–156.

Johnston, R., Jones, K., & Manley, D. (2018). Confounding and collinearity in regression analysis: a cautionary tale and an alternative procedure, illustrated by studies of British voting behaviour. *Quality & Quantity*, *52*(4). https://doi.org/10.1007/s11135-017-0584-6

Kaas, J. H. (1993). The functional organization of somatosensory cortex in primates. *Annals of Anatomy - Anatomischer Anzeiger*, *175*(6). https://doi.org/10.1016/S0940-9602(11)80212-8

Kaas, J. H. (1997). Topographic Maps are Fundamental to Sensory Processing. *Brain Research Bulletin*, *44*(2). https://doi.org/10.1016/S0361-9230(97)00094-4

Katus, T., Grubert, A., & Eimer, M. (2015). Electrophysiological Evidence for a Sensory Recruitment Model of Somatosensory Working Memory. *Cerebral Cortex*, *25*(12). https://doi.org/10.1093/cercor/bhu153

Kikkert, S., Pfyffer, D., Verling, M., Freund, P., & Wenderoth, N. (2021). Finger somatotopy is preserved after tetraplegia but deteriorates over time. *ELife*, *10*. https://doi.org/10.7554/eLife.67713

Kolasinski, J., Makin, T. R., Jbabdi, S., Clare, S., Stagg, C. J., & Johansen-Berg, H. (2016). Investigating the Stability of Fine-Grain Digit Somatotopy in Individual Human Participants. *Journal of Neuroscience*, *36*(4). https://doi.org/10.1523/JNEUROSCI.1742-15.2016

Kuehn, E., Haggard, P., Villringer, A., Pleger, B., & Sereno, M. I. (2018). Visually-Driven Maps in Area 3b. *The Journal of Neuroscience*, *38*(5). https://doi.org/10.1523/JNEUROSCI.0491-17.2017

Logie, R. H., & Cowan, N. (2015). Perspectives on working memory: introduction to the special issue. *Memory & Cognition*, *43*(3). https://doi.org/10.3758/s13421-015-0510-x

Marcus, D. S., Harwell, J., Olsen, T., Hodge, M., Glasser, M. F., Prior, F., Jenkinson, M., Laumann, T., Curtiss, S. W., & van Essen, D. C. (2011). Informatics and Data Mining Tools and Strategies for the Human Connectome Project. *Frontiers in Neuroinformatics*, *5*. https://doi.org/10.3389/fninf.2011.00004

Martuzzi, R., van der Zwaag, W., Farthouat, J., Gruetter, R., & Blanke, O. (2014). Human finger somatotopy in areas 3b, 1, and 2: A 7T fMRI study using a natural stimulus. *Human Brain Mapping*, *35*(1). https://doi.org/10.1002/hbm.22172

Mountcastle, V. B., Talbot, W. H., Darian-Smith, I., & Kornhuber, H. H. (1967). Neural Basis of the Sense of Flutter-Vibration. *Science*, *155*(3762). https://doi.org/10.1126/science.155.3762.597

Nelson, A. J., Staines, W. R., Graham, S. J., & McIlroy, W. E. (2004). Activation in SI and SII; the influence of vibrotactile amplitude during passive and task-relevant stimulation. *Cognitive Brain Research*, *19*(2). https://doi.org/10.1016/j.cogbrainres.2003.11.013

Ojala, M., & Garriga, G. C. (2009, December). Permutation Tests for Studying Classifier Performance. *2009 Ninth IEEE International Conference on Data Mining*. https://doi.org/10.1109/ICDM.2009.108

Pasternak, T., & Greenlee, M. W. (2005). Working memory in primate sensory systems. *Nature Reviews Neuroscience*, *6*(2). https://doi.org/10.1038/nrn1603

Pedregosa, F. and V. G. and G. A. and M. V. and T. B. and G. O. and B. M. and P. P. and W. R. and D. V. and V. J. and P. A. and C. D. and B. M. and P. M. and D. E. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830.

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made

easy. *Behavior Research Methods*, *51*(1). https://doi.org/10.3758/s13428-018-01193-y

PENFIELD, W., & BOLDREY, E. (1937). SOMATIC MOTOR AND SENSORY REPRESENTATION IN THE CEREBRAL CORTEX OF MAN AS STUDIED BY ELECTRICAL STIMULATION. *Brain*, *60*(4). https://doi.org/10.1093/brain/60.4.389

Preuschhof, C., Heekeren, H. R., Taskin, B., Schubert, T., & Villringer, A. (2006). Neural Correlates of Vibrotactile Working Memory in the Human Brain. *Journal of Neuroscience*, *26*(51). https://doi.org/10.1523/JNEUROSCI.2767-06.2006

Puckett, A. M., Bollmann, S., Barth, M., & Cunnington, R. (2017). Measuring the effects of attention to individual fingertips in somatosensory cortex using ultra-high field (7T) fMRI. *NeuroImage*, *161*. https://doi.org/10.1016/j.neuroimage.2017.08.014

Roland, P. E. (1981). Somatotopical tuning of postcentral gyrus during focal attention in man. A regional cerebral blood flow study. *Journal of Neurophysiology*, *46*(4). https://doi.org/10.1152/jn.1981.46.4.744

Rose, N. S., LaRocque, J. J., Riggall, A. C., Gosseries, O., Starrett, M. J., Meyering, E. E., & Postle, B. R. (2016). Reactivation of latent working memories with transcranial magnetic stimulation. *Science*, *354*(6316). https://doi.org/10.1126/science.aah7011

Rousseeuw, P. J., & Croux, C. (1993). Alternatives to the Median Absolute Deviation. *Journal of the American Statistical Association*, *88*(424). https://doi.org/10.1080/01621459.1993.10476408

Sanchez Panchuelo, R. M., Besle, J., Schluppeck, D., Humberstone, M., & Francis, S. (2018). Somatotopy in the Human Somatosensory System. *Frontiers in Human Neuroscience*, *12*. https://doi.org/10.3389/fnhum.2018.00235

Sanders, Z.-B., Wesselink, D. B., Dempsey-Jones, H., & Makin, T. R. (2019). Similar somatotopy for active and passive digit representation in primary somatosensory cortex. *BioRxiv*.

Schmidt, T. T., & Blankenburg, F. (2018). Brain regions that retain the spatial layout of tactile stimuli during working memory – A 'tactospatial sketchpad'? *NeuroImage*, *178*. https://doi.org/10.1016/j.neuroimage.2018.05.076

Schmidt, T. T., & Blankenburg, F. (2019). The Somatotopy of Mental Tactile Imagery. *Frontiers in Human Neuroscience*, *13*. https://doi.org/10.3389/fnhum.2019.00010

Schmidt, T. T., Wu, Y., & Blankenburg, F. (2017). Content-Specific Codes of Parametric Vibrotactile Working Memory in Humans. *The Journal of Neuroscience*, *37*(40). https://doi.org/10.1523/JNEUROSCI.1167-17.2017

Silver, M. A., & Kastner, S. (2009). Topographic maps in human frontal and parietal cortex. *Trends in Cognitive Sciences*, *13*(11). https://doi.org/10.1016/j.tics.2009.08.005

Smith, S. M. (2002). Fast robust automated brain extraction. *Human Brain Mapping*, *17*(3). https://doi.org/10.1002/hbm.10062

Staines, W. R., Graham, S. J., Black, S. E., & McIlroy, W. E. (2002). Task-Relevant Modulation of Contralateral and Ipsilateral Primary Somatosensory Cortex and the Role of a Prefrontal-Cortical Sensory Gating System. *NeuroImage*, *15*(1). https://doi.org/10.1006/nimg.2001.0953

Stelzer, J., Chen, Y., & Turner, R. (2013). Statistical inference and multiple testing correction in classification-based multi-voxel pattern analysis (MVPA): Random permutations and cluster size control. *NeuroImage*, *65*. https://doi.org/10.1016/j.neuroimage.2012.09.063

Sterr, A., Shen, S., Zaman, A., Roberts, N., & Szameitat, A. (2007). Activation of SI is modulated by attention: a random effects fMRI study using mechanical stimuli. *NeuroReport*, *18*(6). https://doi.org/10.1097/WNR.0b013e3280b07c34

Stokes, M. G. (2015). 'Activity-silent' working memory in prefrontal cortex: a dynamic coding framework. *Trends in Cognitive Sciences*, *19*(7). https://doi.org/10.1016/j.tics.2015.05.004

Vallat, R. (2018). Pingouin: statistics in Python. *Journal of Open Source Software*, *3*(31). https://doi.org/10.21105/joss.01026

Wang, L., Bodner, M., & Zhou, Y.-D. (2013). Distributed neural networks of tactile working memory. *Journal of Physiology-Paris*, *107*(6). https://doi.org/10.1016/j.jphysparis.2013.06.001

Watson, A. B., & Pelli, D. G. (1983). Quest: A Bayesian adaptive psychometric method. *Perception & Psychophysics*, *33*(2). https://doi.org/10.3758/BF03202828

Weaverdyck, M. E., Lieberman, M. D., & Parkinson, C. (2020). Tools of the Trade Multivoxel pattern analysis in fMRI: a practical introduction for social and affective neuroscientists. *Social Cognitive and Affective Neuroscience*, *15*(4). https://doi.org/10.1093/scan/nsaa057

Wiestler, T., & Diedrichsen, J. (2013). Skill learning strengthens cortical representations of motor sequences. *ELife*, *2*. https://doi.org/10.7554/eLife.00801

Wu, Y., Uluç, I., Schmidt, T. T., Tertel, K., Kirilina, E., & Blankenburg, F. (2018). Overlapping frontoparietal networks for tactile and visual parametric working memory representations. *NeuroImage*, *166*. https://doi.org/10.1016/j.neuroimage.2017.10.059

Yousry, T. A., Schmid, U. D., Alkadhi, H., Schmidt, D., Peraud, A., Buettner, A., & Winkler, P. (1997). Localization of the motor hand area to a knob on the precentral gyrus. A new landmark. *Brain*, *120*(1), 141–157. https://doi.org/10.1093/brain/120.1.141

Zhou, Y. D., & Fuster, J. M. (1996). Mnemonic neuronal activity in somatosensory cortex. *Proceedings of the National Academy of Sciences*, *93*(19). https://doi.org/10.1073/pnas.93.19.10533

Zhou, Y.-D., & Fuster, J. M. (2000). Visuo-tactile cross-modal associations in cortical somatosensory cells. *Proceedings of the National Academy of Sciences*, *97*(17). https://doi.org/10.1073/pnas.97.17.9777

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, *1*(1). https://doi.org/10.1111/j.2041-210X.2009.00001.x