1

Extracellular Vesicle (EV) Mitochondria and EV/HSP27 Mixtures

Protect the Ischemic Brain Endothelium

Kandarp M. Dave[1], Donna B. Stolz[2], Michael John Reynolds[3], Riyan Babidhan[1], Duncan X. Dobbins[1], Hannah Yankello[4], Rohan Reddy[1], Younsoo Bae[5], Sruti S. Shiva[3,6], Devika S Manickam[1]

[1]Graduate School of Pharmaceutical Sciences, Duquesne University, Pittsburgh, PA

[2]Center for Biologic Imaging, University of Pittsburgh Medical School, Pittsburgh, PA

[3]Pittsburgh Heart Lung Blood Vascular Institute, University of Pittsburgh Medical School, Pittsburgh, PA

[4]Departments of Chemical and Biomedical Engineering, Carnegie Mellon University, Pittsburgh, PA

[5]Department of Pharmaceutical Sciences, College of Pharmacy, The University of Kentucky, Lexington, KY

[6]Department of Pharmacology & Chemical Biology, University of Pittsburgh Medical School, Pittsburgh, PA

†**Corresponding author:**

Devika S Manickam, Ph.D.

600 Forbes Avenue, 453 Mellon Hall

Pittsburgh, PA 15282.

Email: soundaramanickd@duq.edu, Twitter: @manickam_lab

23   Phone: +1 (412) 396-4722, Fax +1 (412) 396-2501

Highlights

➢ Medium-to-large extracellular vesicles (m/lEVs), not small EVs contain mitochondria

➢ m/lEVs increased ATP and mitochondrial function in brain endothelial cells (BECs)

➢ m/lEV mitochondria colocalized with mitochondria in recipient BECs

➢ PEG-DET/EV/HSP27 complexes increase exogenous HSP27 protein loading

➢ EV mitochondria and exogenous HSP27 protect function and structure of ischemic BECs

Abstract

Ischemic stroke causes brain endothelial cell (BEC) death and damages tight junction integrity of the blood-brain barrier (BBB). We engineered endothelial cell-derived extracellular vesicles (EVs) for the delivery of exogenous heat shock protein 27 (HSP27) and harnessed the innate EV mitochondrial load as a one-two punch strategy to increase BEC survival (via mitochondrial delivery) and preserve their tight junction integrity (via HSP27 delivery). We demonstrated that the medium-to-large (m/lEV) but not small EVs (sEV) transferred their mitochondrial load that subsequently colocalized with the mitochondrial network of the recipient primary human BECs. This mitochondrial transfer increased the relative ATP levels and mitochondrial function in the recipient BECs. Compared to EV/HSP27 mixtures, PEG-DET/EV/HSP27 complexes increased HSP27 protein loading. (PEG-DET/HSP27)/EV ternary complexes and EV/HSP27 binary mixtures decreased the paracellular permeability of small and large molecular mass fluorescent tracers in oxygen glucose-deprived primary human BECs. This one-two punch approach to increase the metabolic function and structural integrity of BECs is a promising strategy for BBB protection and prevention of long-term neurological dysfunction post-ischemic stroke.

Graphical Abstract

Keywords

extracellular vesicles, mitochondria, heat shock protein, paracellular permeability, BBB protection, ischemic stroke

Abbreviations

BBB	Blood-brain barrier
BECs	Brain endothelial cells
Calcein AM	Calcein acetoxymethyl
ECAR	Extracellular acidification rate
EVs	Extracellular vesicles
EXOs	Exosomes
FT cycle	Freeze/Thaw cycle
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
HSP27	Heat shock protein 27
hCMEC/D3	human cerebral microvascular endothelial cell line
HBMEC	primary human brain microvascular endothelial cells
MVs	Microvesicles
m/lEV	medium-to-large EVs
MitoT-red-EV	Mitotracker deep red-labeled extracellular vesicles
OCR	Oxygen consumption rate
OGD	Oxygen-glucose deprivation
OGD/RP	Oxygen-glucose deprivation/reperfusion
PEI	Polyethylenimine
PEG-DET	poly (ethylene glycol)-*b*-poly (diethyl triamine)
ROS	Reactive oxygen species
sEV	small EVs
TRITC	Tetramethyl rhodamine iso-thiocyanate

1. Introduction

EVs are an emerging class of natural carriers for drug delivery due to their known roles in intercellular communication. They retain membrane signatures reminiscent of the donor cells from which they are derived, and therefore possess inherent homing capabilities to recipient cells of the same type (1, 2). They are also likely to be less immunogenic compared to synthetic nanoparticles. The smaller EVs (sEVs) range from 30-200 nm in particle diameter and their biogenesis involves the inward budding of endosomal membranes that transforms into multivesicular bodies followed by their fusion with the plasma membrane and sEV release into extracellular spaces (2). The biogenesis of the medium-to-larger EVs (m/lEVs) involves their outward budding from the cell's plasma membrane with particle diameters ranging from 100 – 1000 nm (2). The selective packaging of the functional mitochondria and mitochondrial proteins in the m/lEVs motivated us to harness the m/lEV mitochondrial load as a therapeutic modality. Mitochondria play a central role in cellular energy production and regulation of cell death including apoptosis and autophagy (3). Ischemia-induced mitochondrial dysfunction in the brain endothelial cells (BECs) lining the blood-brain barrier (BBB) initiates the generation of excessive reactive oxygen species, reduction in ATP levels, and consequently BEC death (3). Therefore, protection of mitochondrial function via exogenous mitochondria supplementation is a potent strategy to increase BEC survival post-ischemic stroke. Thus, we rationalize that mitochondria-containing m/lEVs derived from BECs can increase cellular bioenergetics and survival under hypoxic conditions.

Ischemic stroke-induced oxygen-glucose-deprivation (OGD) decreases ATP levels in the

BECs leading to the accumulation of cellular cations and excitatory neurotransmitters (4-6). The

cationic overload catalyzes enzymatic activities leading to generation of reactive oxygen species

(ROS), impairment of mitochondrial ROS defense mechanisms, and the subsequent

mitochondrial dysfunction. Therefore, restoring the mitochondrial function is a viable strategy to

rescue damage to the BECs. In addition, disruption of the BBB is a major hallmark of ischemic

stroke that is associated with altered expression of tight junctions, adherens junction proteins,

and BBB transporters (4, 5, 7). Early ischemia/reperfusion activates the polymerization of the

actin cytoskeleton in endothelial cells which disassemble and internalize the tight junction

proteins and consequently lead to the loss of barrier properties of the BECs lining the BBB (8,

9). Uncontrolled actin polymerization-induced breakdown of BBB leads to infiltration of

proinflammatory mediators, blood cells, circulatory immune cells, and toxins into the brain

parenchyma, and lead to the secondary injury cascade (9, 10). Hence, a combined strategy to

decrease BEC death and their paracellular permeability, ultimately leading to protection of the

BBB metabolic function and structural integrity is a potent approach for the treatment of

ischemia/reperfusion injury (stroke).

Preclinical studies have demonstrated that endothelial, but not neuronal overexpression of

heat shock protein 27 (HSP27), inhibited actin polymerization and elicited long-lasting

protection against stroke-induced BBB disruption and neurological deficits (9, 10). HSP27 binds

to actin monomer and inhibits tight junctional protein translocation in endothelial cells (9, 11-

14). Intravenous administration of cell-penetrating transduction domain (TAT)-HSP27 rapidly

enhanced HSP27 levels in brain microvessels, decreased infract volumes and attenuated

ischemia/reperfusion-induced BBB disruption (9). However, the systemic delivery of HSP27

using TAT peptide resulted in non-specific uptake-associated side effects in the brain and other

tissues (9). Therefore, the delivery of exogenous human recombinant HSP27 protein to BECs

using natural carriers like EVs can likely increase its specific uptake into BECs owing to

inherent membrane affinity of BEC-derived EVs to donor BECs (15, 16).

Surface engineering of BEC-derived EVs with exogenous HSP27 is a promising strategy that

can allow harnessing the inherent targeting capabilities of the EVs to the recipient BECs along

with the added benefits of their mitochondrial load. Here, we *hypothesize* that the engineering of

innate mitochondria-containing BEC-derived EVs with exogenous HSP27 protein is one-two

punch approach to increase BEC survival and protect its tight junction barrier via decreasing the

paracellular permeability post-ischemia. This approach will protect and strengthen the BBB that

in turn can ameliorate the long-term neurological damage and dysfunction. We tested the effects

of adding a cationic copolymer, poly (ethylene glycol)-*b*-poly (diethyltriamine) (PEG-DET) to

the EV/HSP27 mixtures to determine if the degree of HSP27 complexation can be further

improved. We have previously used PEG-DET polymer to form nanosized complexes with

superoxide dismutase protein and demonstrated a >50% reduction in brain infract volume in a

mouse model of acute ischemic stroke (17). PEG-DET is a cationic diblock copolymer known

for its safety and gene transfer efficacy in comparison to commercial transfection agents such as

lipofectamine, polyethyleneimine, and other cationic polymers (18-20).

In this study, we isolated sEVs and m/lEVs from hCMEC/D3: a human brain endothelial cell line using a sequential ultracentrifugation method (21) and characterized their particle diameter, zeta potential, and membrane integrity post-cold storage. While we collectively refer to both large (m/lEV) and small (sEV) vesicle fractions as EVs, we have studied the singular effects of both m/lEVs and sEVs. We showed the presence of mitochondrial components in m/lEVs using transmission electron microscopy and western blotting. We evaluated the effects of EV dose and incubation times on their uptake to the recipient BECs and demonstrated the colocalization of m/lEV-delivered mitochondria with the mitochondrial network of the recipient BECs. We studied the effects of EV exposure on the resulting relative ATP levels, mitochondrial respiration, and glycolytic capacity of the recipient BECs. The formation of EV/HSP27 mixtures and their physicochemical characteristics were studied using native polyacrylamide gel electrophoresis and dynamic light scattering. The effects of EV/HSP27 on the paracellular permeability of small and large molecule fluorescent tracers were evaluated under ischemic and ischemia/reperfusion conditions in primary human BECs.

2. Materials

Recombinant human HSP27 was purchased from Novus Biologicals (Centennial, CO). Cell Titer Glo 2.0 reagent (ATP assay) was procured from Promega (Madison, WI). Micro BCA and Pierce BCA protein assay kits were purchased from Thermo Scientific (Rockford, IL). PEG-DET polymer was synthesized by aminolysis of PEG-poly(β-benzyl L-aspartate) block copolymers with diethyltriamine as previously reported (18, 20, 22, 23). Bio-Safe Coomassie G-250 stain was purchased from Bio-Rad Laboratories Inc. (Hercules, CA). Collagen Type I was purchased from Corning (Discovery Labware Inc, Bedford, MA) and endothelial cell basal medium-2

(EBM-2) was procured from Lonza (Walkersville, MD). Hydrocortisone, human basic fibroblast

growth factor and ascorbic acid were purchased from Sigma-Aldrich (Saint Louis, MO).

Penicillin-Streptomycin solution and Chemically Defined Lipid Concentrate were procured from

Invitrogen (Carlsbad, CA). Polycarbonate centrifuge tubes were purchased from Beckman

Coulter, Inc. (Brea, CA). Electrophoresis sample buffer was purchased from Bio-Rad (Hercules,

CA). PET track-etched membrane Falcon Cell Culture inserts of 0.4 µm pore size were procured

from Corning (Discovery Labware Inc, Bedford, MA). TRITC 65-85 kD and 4.4 kD dextran was

procured from Sigma (St. Louis, MO). A low-volume disposable cuvette (Part no. ZEN0040,

Malvern) was used for particle size measurements. CellLight Mitochondria-GFP Backman 2.0

reagent, MitoTracker Green FM, and MitoTracker Deep Red FM, Dynabeads Protein G

(cat#10003D), and DynaMag-2 magnetic stand (cat#12321D) were procured from Invitrogen

(Eugene, OR). Mouse monoclonal antibodies against ATP5A, GAPDH, HSP27 were purchased

from Abcam. Mouse monoclonal antibody against CD9 was received from Life Technologies

corporation (Eugene, OR), whereas Alexa Fluor 790-conjugated donkey anti-mouse IgG was

received from Jackson ImmunoResearch Lab Inc (West Grove, PA).

2.1. Cell models

A human cerebral microvascular endothelial cell line (hCMEC/D3, catalog number.

102114.3C) was received from Cedarlane Laboratories (Burlington, Ontario) at passage number

(P) 25, and cells between P25 and P35 were used in all experiments (15, 21). hCMEC/D3 cells

were grown in tissue culture flasks, multiwell plates, or transwell inserts pre-coated using 0.15

mg/mL rat collagen I in a humidified 5% CO_2 incubator at $37 \pm 0.5°C$ (Isotemp, Thermo Fisher

Scientific). The cells were cultured in complete growth medium composed of endothelial cell

basal medium (EBM-2) supplemented with fetal bovine serum (5% FBS), penicillin (100

units/mL)-streptomycin (100 µg/mL) mixture, hydrocortisone (1.4 µM), ascorbic acid (5

µg/mL), Chemically Defined Lipid Concentrate (0.01%), 10 mM HEPES (pH 7.4), and bFGF (1

ng/mL). The complete growth medium was replenished every other day until the cells formed

confluent monolayers. Prior to passage, the cells were washed using 1x phosphate buffer saline

(PBS) and detached from the flask using 1x TrypLE Express Enzyme (gibco, Denmark). We

received primary human brain microvascular endothelial cells (HBMEC, catalog no. ACBRI

376) from Cell Systems (Manassas, VA) at P3, and cells below P11 were used in all

experiments. HBMECs maintained in Cell Systems Classic Culture Medium containing 1%

culture boost were cultured in tissue culture flasks, multiwell plates, or transwell inserts pre-

treated with attachment factors. The complete growth medium was replenished every day until

the cells formed confluent monolayers. For passage, HBMEC monolayers were washed with

Passage Reagent Group 1 (PRG 1, dPBS/EDTA solution), detached with PRG 2 (dPBS/trypsin-

EDTA solution) and PRG 3 (trypsin inhibitor solution).

2.2. Isolation of sEVs and m/lEVs from hCMEC/D3 cells

It should be noted that while we collectively refer to both large (m/lEVs) and small (sEVs)

vesicle fractions as EVs, we study both m/lEVs and sEVs. Wherever noted, a 1:1 w/w mixture of

sEVs and m/lEVs is collectively referred to as EVs. sEVs and m/lEVs were isolated from

conditioned medium supernatants of hCMEC/D3 cells using the differential ultracentrifugation

method (21). Briefly, tissue culture flasks with 175 cm^2 growth area (T175) containing confluent

hCMEC/D3 were washed with pre-warmed 1x PBS and incubated with serum-free medium for

48 h in a humidified 5% CO_2 incubator at 37 ± 0.5°C. Post-incubation, the EV-conditioned

medium was collected in polypropylene centrifuge tubes and centrifuged at 2000 ×g for 22 min

at 4°C to pellet down apoptotic bodies and cell debris using a Sorvall ST 8R centrifuge

(ThermoFisher Scientific, Osterode am Harz, Germany). The supernatant was transferred into

polycarbonate tubes and centrifuged at 20,000 ×g for 45 min at 4°C to pellet down m/lEVs using

an Optima XE-90 ultracentrifuge equipped with a 50.2 Ti rotor (Beckman Coulter, Indianapolis,

IN). Next, the supernatant was filtered through a 0.22 µm PES membrane syringe filter, and the

flow-through was centrifuged at 120,000 ×g for 70 min at 4°C to collect sEVs. Lastly, m/lEVs

and sEVs pellets were washed with 1x PBS and suspended in either 1x PBS for particle diameter

measurements and *in vitro* experiments or 10 mM HEPES buffer pH 7.4 for zeta potential

measurements. sEVs and m/lEVs samples were stored at -20°C until further use. The total

protein content in sEVs and m/lEVs was quantified using Pierce MicroBCA assay.

2.3. Dynamic light scattering (DLS)

The stability of naïve EVs under storage conditions was determined by their measuring

particle diameter and zeta potential using dynamic light scattering. EV samples at 0.5 mg

protein/mL in 1x PBS were frozen at -20°C for 24 h and thawed at room temperature for 30 min.

Post-thawing, the particle diameters, and dispersity indices were measured using Malvern

Zetasizer Pro (Worcestershire, UK). The freeze-thaw cycle was repeated three times and all

samples were run in triplicate. Average particle diameter, dispersity index, and zeta potential

values were reported as mean ± standard deviation.

2.4. Membrane integrity of EVs after isolation and upon storage conditions

The membrane integrity of sEV and m/lEV post-isolation and upon revival after their storage

conditions (described earlier in the DLS section) was determined using a calcein AM flow

cytometry assay. First, polystyrene sub-micron-sized beads ranging from 0.2 – 2 µm particle

diameters were used to calibrate the Attune NxT flow cytometer. The calibration beads, sEV,

and m/lEV samples were tested and events captured in forward scatter (FSC) and side scatter

(SSC) plots were analyzed using a small particle side scatter 488/10-nm filter (BL1) channel.

sEVs and m/lEVs were diluted to 20 µg protein/mL in PBS and incubated/stained with 10 µM

calcein AM for 20 min at room temperature in the dark. Unstained sEVs and m/lEVs were used

to gate the background signals, whereas samples treated with 2% v/v Triton X-100 followed by

staining with calcein AM were used as controls to determine if calcein-positive signals were

specifically associated with intact EVs. For each sample analysis, an aliquot of 100 µL was run

through Attune NxT Acoustic cytometer (Invitrogen, Singapore) and 50,000 events were

recorded during the run. The calcein-associated fluorescence intensity was detected at 488/10 nm

and percentage signal intensities were presented in histogram plots generated using Attune

software. Calcein AM-associated background signals were gated using the controls such as PBS

containing 10 µM calcein AM and PBS/2% Triton X-100/calcein AM mixture.

2.5. Transmission Electron Microscopy analysis of EVs

EVs were loaded on Formvar/carbon-coated grids, negatively stained with 1% uranyl

acetate and examined with a JEM-1400 Plus transmission electron microscope (JEOL, Peabody,

MA, USA) fitted with an AMT digital camera (Danvers, MA, USA) (24). Suspensions of EVs

were pelleted at 100,000xg in a Beckman airfuge for 20 min and the pellets were fixed in 2.5%

glutaraldehyde in PBS overnight. The supernatant was removed and the cell pellets were washed

3x in PBS and post-fixed in 1% OsO_4, 1% $K_3Fe(CN)_6$ for 1 hour. Following 3 additional PBS

washes, the pellet was dehydrated through a graded series of 30-100% ethanol. After several

changes of 100% resin over 24 hours, pellet was embedded in a final change of resin, cured at

37^oC overnight, followed by additional hardening at 65^oC for two more days. Ultrathin (70 nm)

sections were collected on 200 mesh copper grids, stained with 2% uranyl acetate in 50%

methanol for 10 minutes, followed by 1% lead citrate for 7 min. Sections were imaged using a

JEOL JEM 1400 Plus transmission electron microscope (Peabody, MA) at 80 kV fitted with a

side mount AMT 2k digital camera (Advanced Microscopy Techniques, Danvers, MA).

2.6. Uptake of Mitotracker-labeled EVs into HBMECs using flow cytometry

2.6.1. Isolation of mitochondria-labeled EVs

hCMEC/D3 cells (P32) were cultured in a T175 flask to confluency. The complete growth

medium was removed, cells were washed with 1x PBS, and cells were incubated with 100 nM

Mitotracker deep red (MitoT-red) diluted in a conditioned medium for 30 min in a humidified

incubator. Next, the medium was replaced with serum-free medium after washing the cells with

1x PBS, and cells were kept in a humidified incubator for 24 h. Post-incubation, the conditioned

medium was collected into centrifuge tubes. sEV (MitoT-red-sEV) and m/lEV (MitoT-red-

m/lEV) from Mitotracker Red-labeled cells were isolated from the conditioned medium using the

differential centrifugation method described in section 2.2. The EV protein content in MitoT-red-

sEV and MitoT-red-m/lEV were determined using MicroBCA assay and the samples were stored

at -80 °C until further use.

2.6.2. Quantification of MitoT-EV uptake into recipient hCMEC/D3 cells using flow cytometry

hCMEC/D3 cells were cultured in 48-well plates at 50,000 cells/well in complete growth medium. Unstained and untreated cells were used as a control, whereas cells stained with 100 nM of Mitotracker deep red (MitoT-red) for 45 min in complete growth medium were used as a positive control. The cells were treated with MitoT-red-sEV and MitoT-red-m/lEV at 30, 75, and 150 µg EV protein/well in complete growth medium for 72 h in a humidified incubator. The cells were also treated with unlabeled sEV and m/lEV at 150 µg EV protein/well in complete growth medium for 72 h. Post-treatment, the cells were washed with 1x PBS, dissociated using TrypLE Express, diluted with PBS, and collected into centrifuge tubes. For each sample, an aliquot of a 100 µL cell suspension was analyzed through Attune NxT Flow cytometer and 10,000 events were recorded in FSC vs. SSC plots. The Mitotracker deep red fluorescence intensity was detected at 670/14 nm and percentage signal intensities were presented in histogram plots generated using Attune software version 3.2.3. Mitotracker deep red background signals were gated using the controls including PBS and untreated cells.

2.7. Uptake of MitoT-EVs into primary human brain endothelial cells using fluorescence microscopy

2.7.1. Uptake of MitoT-EVs into primary HBMEC monolayers

MitoT-red sEV and MitoT-red-m/lEV were isolated from the conditioned medium of hCMEC/D3 cells as described in section 2.6.1. Primary HBMEC (P6) were cultured in 96-well plates at 16,500 cells/well in complete growth medium. Post-confluency, the cells were treated with MitoT-red-sEV and MitoT-red-m/lEV at 10, 25, and 50 µg EV protein/well in complete

growth medium for 24, 48, and 72 h in a humidified incubator. At each time-point, the cells were

observed under an Olympus IX 73 epifluorescent inverted microscope (Olympus, Pittsburgh,

PA) using the Cyanine-5 (Cy5, excitation 651 nm, and emission 670 nm) and bright-field

channels at 20x magnification. Images were acquired using CellSens Dimension software

(Olympus, USA).

2.7.2. Colocalization of MitoT-EVs with the mitochondrial network in the recipient

hCMEC/D3 and HBMEC monolayers

2.7.2.1. Mitotracker green staining of recipient cell mitochondria

HBMEC and hCMEC/D3 cells were seeded in a 96 well-plate at 16,500 cells/well and

incubated in a humidified incubator at 37°C. Post-confluency, the complete growth medium was

removed, cells were washed with 1x PBS, and cells were incubated with 100 µM of Mitotracker

green in a complete medium for 30 min. Post-treatment, the medium was replaced, washed with

PBS, and incubated with MitoT-red-sEV or MitoT-red-m/lEV at 50 µg EV protein/well in a

complete growth medium for 72 h in a humidified incubator. hCMEC/D3 cells stained with only

Mitotracker green were used as a control. Post-incubation, the medium was removed, cells were

washed and incubated with phenol-red free and serum-containing DMEM-high glucose medium.

The cells were then observed under an Olympus IX 73 epifluorescent inverted microscope

(Olympus, Pittsburgh, PA) using Cyanine-5 channel (Cy5, excitation 651 nm, and emission 670

nm) to detect MitoT-red-EV uptake and GFP channel to detect Mitotracker Green signals at 20x

magnification and images were acquired using CellSens Dimension software (Olympus, USA).

2.7.2.2. Staining of a mitochondrial matrix protein in the recipient cells using CellLight Mitochondria-GFP BackMam reagent

We used CellLight Mitochondria-GFP BacMam to label a structural mitochondrial matrix protein (25). HBMECs were seeded in a 96 well-plate at 16,500 cells/well and incubated in a humidified incubator at 37°C. Post-confluency, the complete growth medium was removed, cells were washed with 1x PBS, and cells were incubated with CellLight Mitochondria-GFP reagent (at a dilution of 2 µL/10,000 cells as recommended by the manufacturer) for 16 h. Post-transduction, the medium was removed and cells were washed with 1x PBS. Next, the cells were incubated with MitoT-red-sEV and MitoT-red-m/lEV at 50 µg EV protein/well in complete growth medium for 72 h in a humidified incubator. HBMECs transduced with only CellLight Mitochondria-GFP were used as a control. Post-incubation at 24 and 72 h, the medium was removed, cells were washed and incubated with phenol-red free and serum-containing DMEM-high glucose medium. The cells were then observed under Olympus IX 73 epifluorescent inverted microscope (Olympus, Pittsburgh, PA) using Cyanine-5 channel (Cy5, excitation 651 nm, and emission 670 nm) to detect MitoT-EV uptake and GFP channel for CellLight Mitochondria-GFP signals at 20x magnification and images were acquired using CellSens Dimension software (Olympus, USA).

2.8. Effects of naïve EVs on the relative ATP levels in primary HBMECs under normoxic and ischemic/oxygen-glucose deprivation (OGD) conditions

To simulate ischemic conditions *in vitro*, HBMECs were exposed to different OGD settings, and the resulting cell viability was evaluated using an ATP assay (21, 26, 27). HBMECs were cultured in 96-well plates at 16,500 cells/well. Confluent HBMECs were incubated with OGD

medium defined as follows: 120 mM NaCl, 5.4 mM KCl, 1.8 mM $CaCl_2$, 0.8 mM $MgCl_2$, 25

mM Tris-HCl, pH 7.4 for 0.5 to 24 h in either normoxic or hypoxic conditions (28). For

normoxic conditions, the culture plates were incubated in a humidified incubator with 95% air

and 5% carbon dioxide whereas for hypoxic conditions, the culture plates were incubated in an

OGD chamber (Billups Rothenberg, Del Mar, CA) pre-flushed with 5% carbon dioxide, 5%

hydrogen and 90% nitrogen at 37±0.5˚C. For normoxic conditions, the medium was removed

and the recipient HBMEC monolayers were incubated with hCMEC/D3-derived sEV, m/lEV,

and sEV+m/lEV at doses of 10, 25, and 50 µg EV protein/well in a humidified incubator for 24,

48, and 72 h. For hypoxic conditions, the medium was removed and the recipient HBMEC

monolayers were incubated with hCMEC/D3-derived sEV, m/lEV, and sEV+m/lEV at doses of

10, 25, and 50 µg EV protein/well in the OGD medium and a humidified incubator for 24 h.

Then, the treatment mixtures were replaced with pre-warmed complete growth medium, and an

equal quantity of Cell Titer Glo 2.0 reagent was added. The plates were incubated for 15 min at

RT on an orbital shaker in the dark and relative luminescence units were measured at 1 s

integration time.

2.9. Measurement of mitochondrial function using Seahorse Analysis

The oxidative phosphorylation and glycolytic functions of hCMEC/D3 cells treated with EVs

during normoxic conditions were evaluated using the Seahorse analysis by measuring oxygen

consumption rate (OCR) and Extracellular Acidification rate (ECAR) (15, 29). hCMEC/D3 cells

seeded at 20,000 cells/well were cultured in Seahorse XF96 plate for four days. The cells were

incubated with hCMEC/D3-derived sEV, m/lEV, and sEV+m/lEV for 24, 48, and 72h at 3.4 µg

EV protein/well equivalent to 30 µg EV protein/cm^2 in complete growth medium. Post-

incubation, the medium was replaced with pre-warmed DMEM and subjected to Seahorse

analysis (15). After measurement of baseline OCR, 2.5 µmol/L oligomycin A and 0.7 µmol/L

carbonyl cyanide-p-trifluoromethoxyphenyl-hydrazone were consecutively added to measure the

proton leak and maximal OCR, respectively (29). The total protein of the cells in each well was

measured using Pierce BCA assay.

2.10. Complexation of HSP27 with EVs, PEG-DET and EV-PEG-DET

PEG-DET/HSP27 complexes were prepared using a rapid mixing method. A PEG-DET

polymer solution prepared in 10 mM HEPES buffer, pH 7.4 was mixed with 2 µg HSP27 at

PEG-DET/HSP27 w/w ratio 0.05, 0.2, 1, 5, 10, and 20:1 for 30 s. The mixture was incubated at

room temperature (RT) for 30 min. For the preparation of sEV/HSP27 and m/lEV/HSP27

mixtures, hCMEC/D3-derived EVs were suspended in 1x PBS were incubated with 2 µg HSP27

at EV protein/HSP27 w/w ratios 5, 10, and 15:1 in centrifuge tubes. The mixture was vortexed

for 30 s and incubated at room temperature for 30 min. To prepare (PEG-DET/HSP27)/EV

complexes, PEG-DET/HSP27 complexes were prepared at 20:1 and 30:1 w/w ratios followed by

incubation with 10 µg EV protein for 30 min at RT. The different complexes were characterized

for electrophoretic mobility, particle diameter, and zeta potential.

2.11. Native Polyacrylamide Gel Electrophoresis (PAGE)

The complexation of HSP27 with PEG-DET, sEV, and m/lEV was confirmed using native

PAGE. A polyacrylamide gel consisting of 4% and 10% of acrylamide in the stacking and

resolving sections, respectively, was prepared using a gel casting assembly (Bio-Rad

Laboratories Inc., Hercules, CA) (21, 26). Native HSP27, PEG-DET/HSP27, and EV/HSP27

complexes containing 2 µg of HSP27 at the indicated w/w ratios were mixed with native sample

buffer and loaded into the gel lanes. Free PEG-DET polymer, naïve sEV, and m/lEV equivalent

to indicated w/w ratios were used as controls. In an independent experiment, PEG-DET/HSP27

at w/w 20:1 and 30:1 was complexed with sEV, m/lEV, and EVs (sEV: m/lEV 1:1) at PEG-DET

to EV protein 2:1 and 3:1 w/w ratios. The gel was run in 1x Tris-Glycine buffer, pH 8.3 at 100 V

for 2 h at 2-8 °C using PowerPac Basic setup (Bio-Rad Laboratories Inc., Hercules, CA). Post-

electrophoresis, the gel was washed with deionized water for 30 min and stained with 50 mL of

Biosafe Coomassie blue G-250 for 1 h on an orbital shaker at room temperature. The gel was

washed with deionized water for 30 min and scanned under Odyssey imager (LI-COR Inc.,

Lincoln, NE) at 800 nm channel and intensity setting 5. The band densities were quantified using

ImageStudio 5.2 software.

2.12. Dynamic light scattering analysis of HSP27 complexes/mixtures

The average particle diameters, dispersity indices, and zeta potentials of the HSP27

complexes were analyzed using a Malvern Zetasizer Pro (Worcestershire, UK). hCMEC/D3 cell

line-derived naïve EVs at 0.5 mg EV protein/mL were diluted in either 1x PBS for size and

dispersity index or 10 mM HEPES buffer pH 7.4 for zeta potential measurements. In addition,

particle sizes and dispersity indices of native HSP27 protein at 20 µg/mL, PEG-DET/HSP27

complexes at 10, 20, and 30:1 w/w ratios, sEV/HSP27 and m/lEV/HSP27 mixtures at 10:1 w/w

ratio were measured in a low-volume disposable cuvette. Free PEG-DET equivalent to 10, 20,

and 30:1 w/w ratios and naïve EVs equivalent to 10:1 w/w ratio were also analyzed. For the zeta

potential of the above complexes, a 50 µL sample of complexes was further diluted in zeta

cuvette containing 10 mM HEPES buffer at pH 7.4. The samples were run in triplicate. Data are

presented as average particle diameter±standard deviation. The reported data are representative

of 3 independent experiments.

2.13. Cytocompatibility of HSP27 complexes/mixtures with primary HBMEC and hCMEC/D3 monolayers

The cell viability of hCMEC/D3 and primary HBMEC monolayers treated with HSP27

complexes was measured using Cell Titer Glo (ATP) assay. hCMEC/D3 and HBMEC were

seeded at 16,500 cells/well in a 96 well-plate and cultured in a humidified incubator at 37±0.5˚C.

The growth medium was replaced with treatment mixtures containing either native HSP27,

sEV/HSP27 (10:1), m/lEV/HSP27 (10:1), sEV+m/lEV at 1:1/HSP27 (10:1), or PEG-

DET/HSP27 (20:1) at a dose of 2 µg HSP27 protein per well. Naïve sEV, m/lEV, equivalent

amounts of sEV+m/lEV, and free PEG-DET equivalent to 20:1 in complete medium were used

as controls. Polyethyleneimine (PEI) at 50 and 100 µg/mL in a complete growth medium was

used as a positive control. The complexes and controls were treated for 72 h in a humidified

incubator at 37±0.5˚C. Post-incubation, the ATP assay was performed using Cell Titer Glo 2.0

reagent as described earlier in section 2.8. The cell viability of HSP27 complexes treated

HBMECs was calculated using Equation 1.

$$Cell\ viability\ (\%) = \frac{Relative\ light\ units\ (RLU)\ of\ treated\ cells}{RLU\ of\ untreated\ cells} \times 100 \qquad \text{(Equation 1)}$$

2.14. Paracellular permeability of TRITC-labeled 4.4 and 65-85 kD Dextran in HBMEC monolayers pre-treated with EV/HSP27 mixtures, PEG-DET/HSP27 complexes and PEG-DET/EV/HSP27 ternary complexes.

2.14.1 Hypoxic conditions (OGD only)

Primary HBMEC (P4-P9) in a complete growth medium were seeded at 50,000 cells/cm^2 in a

24-well format cell culture insert (insert area: 0.33 cm^2) for four days to form a complete

endothelial monolayer. The medium was replaced every 48 h during this culturing period. The

abluminal wells were filled with complete growth medium throughout the culturing period. The

complete growth medium was replaced with 300 µL of growth medium containing 2 µg

HSP27/well (formulated as described in 2.10) for 72 h. Post-treatment, the complete growth

medium was replaced with 300 µL of OGD medium containing 1 µM TRITC-Dextran. The

OGD medium containing 1µM TRITC-Dextran alone was used as a control. The abluminal

chamber was filled with 0.5 mL of fresh complete growth medium. The untreated group was

incubated in a humidified incubator whereas OGD treatment groups were incubated in an OGD

chamber (as described earlier in section 2.8.). The concentration of TRITC-Dextran in the

abluminal medium was measured at 4, and 24 h post-treatment. A 50 µL aliquot was collected at

indicated time points from the abluminal side. An equal volume of fresh medium was replaced to

maintain the sink conditions. The concentration of TRITC-Dextran was measured using Synergy

HTX multimode plate reader at excitation 485/20 and emission 580/50 nm. The diffusion rate of

TRITC-Dextran at each time point was calculated by the amount of TRITC- Dextran per surface

area (0.33 cm^2) of the insert per time (30 min).

2.14.2. OGD/reperfusion

Post-24 h of OGD exposure, HBMECs in the culture inserts were washed with 1x PBS.

HBMECs were incubated with a complete growth medium containing 1 µM TRITC 65-85 or 4.4

kD Dextran in a 24-well format cell culture insert for 1h – 24h (reperfusion) in a humidified

incubator. The abluminal chamber was filled with 0.5 mL of fresh complete growth medium.

The concentration of TRITC-Dextran in the abluminal medium was measured at 1, 2, and 4, and

24 h during reperfusion. A 50 µL of samples from the abluminal side were collected at indicated

time points. An equal volume of fresh medium was replaced to maintain the sink conditions. The

concentration of TRITC-Dextran was measured using Synergy HTX multimode plate reader at

excitation 485/20 and emission 580/50 nm. The diffusion rate of TRITC-Dextran at each time

point was calculated as the amount of TRITC-Dextran present in the basolateral compartment

per unit surface area of the insert per unit time.

$Diffusion\ rate\ of\ TRITC - Dextran =$

$$\frac{Amount\ of\ TRITC - Dextran\ in\ basolateral\ compartment}{(Surface\ area\ of\ transwell\ insert)x(time\ at\ which\ basolateral\ sample\ was\ removed)}$$ (Equation 2)

2.15. Statistical analysis

Statistically significance among the mean of controls and treatment groups or within

treatment groups were analyzed using one-way analysis of variance (ANOVA) or two-way

ANOVA at 95% confidence intervals using GraphPad Prism 9 (GraphPad Software, LLC). The

notations for the different levels of significance are indicated as follows: $*p<0.05$, $**p<0.01$,

$***p<0.001$, $****p<0.0001$.

3. Results

3.1. EVs retained their physicochemical characteristics and membrane integrity upon

revival from storage conditions.

We collectively refer to sEVs and m/lEVs as EVs wherever applicable. We first studied the

effect of storage conditions on the retention of physicochemical characteristics of hCMEC/D3

BEC-derived sEVs and m/lEVs using dynamic light scattering (**Fig. 1a**). Freshly isolated sEVs

and m/lEVs showed particle diameters of about 122 and 185 nm with a dispersity index ~0.4

(**Fig. 1a**). We studied the morphology of naïve sEVs and m/lEVs using transmission electron

microscopy (TEM). Negative stain TEM analysis showed nearly spherical sEVs of about 50 nm

(**Fig. 1b**) whereas m/lEV were about 200 nm in size (**Fig. 1b**). In addition, there were no

significant differences in the particle diameters of sEVs between a freshly-isolated sample and

when measured after three freeze-thaw cycles (**Fig. S1a**). Similar to sEVs, m/lEVs retained their

particle diameters after three freeze-thaw cycles with a consistent dispersity index (**Fig. S1b,d**).

sEVs and m/lEVs showed an initial negative zeta potential of about -22 mV that ranged between

-15 to -30 mV after three consecutive freeze-thaw cycles (**Fig. 1a, S1e, f**).

Fig. 1. Physicochemical characteristics, membrane integrity, and mitochondrial contents in EVs. (**a**) Average particle diameters, and dispersity indices, and zeta potentials of hCMEC/D3 cell line-derived freshly isolated EVs were determined using dynamic light scattering on a Malvern Zetasizer Pro-Red. Samples were diluted to 0.1 mg protein/mL in 1x PBS for particle diameter and 10 mM HEPES buffer pH 7.4 for zeta potential measurements. Data are presented as mean±SD of n=3 measurements. (**b**) Negative stain TEM images of hCMEC/D3-derived sEV and m/lEV. (**c**) The histograms of calcein-positive events of freshly isolated intact sEVs and m/lEVs were detected using a small particle side scatter 488/10-nm filter in an Attune flow cytometer. Unstained EVs were used to gate the histograms for estimating percentage calcein-positive counts. Triton x-100 was used for EV lysis. (**d**) representative TEM images of sectioned sEVs and mitochondria containing m/lEVs. (**e**) Detection of mitochondrial proteins in sEVs and m/lEVs-derived from hCMEC/D3 cells using western blot analysis. Western blot showing mitochondria protein marker ATP5A, mitochondria marker TOMM20, exosome (sEVs) marker CD9, and cytosolic protein marker GAPDH expression. The total protein content in the cell lysates, sEV, and m/lEV was measured using a micro BCA assay. Fifty µg protein/sample was incubated with Laemmli sample buffer at 95°C for 5 min. The sample was run on sodium dodecyl sulfate polyacrylamide gel electrophoresis, proteins were transferred on nitrocellulose membrane, and non-specific proteins were blocked with blocking buffer. The blot was incubated with primary antibodies overnight at 4°C, and secondary antibodies at room temperature. The blots were imaged on the 800 nm channel using an Odyssey imager (LI-COR Inc. Lincoln, NE) at intensity setting 5 and processed using ImageStudio 5.2 software.

The membrane integrity of EVs upon revival from cold storage was determined using a previously reported calcein-based flow cytometry assay (21). Prior to analyzing the EVs, we first calibrated the flow cytometer using polystyrene beads of particle diameters ranging from 0.1 to 2 µm using a small particle side scatter filter (488/10 nm, BL1) (**Fig. S1g**). The relative position of EV clusters in the FSC/SSC overlay plots was proportional to the particle diameter since an increase in diameter showed a right-upward shift of the clusters. Notably, sEVs and m/lEVs clusters overlapped to a large extent in the area corresponding to 0.1-0.2 µm bead diameters suggesting that this protocol allowed us to detect EV-sized particles (**Fig. S1h**). The particle counts for PBS/calcein AM, unlabeled-sEVs and m/lEVs, PBS/Triton X-100/calcein AM mixture (sample processing controls) were acquired on SSC/BL1 density plots, and were used for gating calcein AM positive signals. About 90% of freshly-isolated sEVs and m/lEVs were

calcein-positive suggesting that EVs retained intact membranes after the ultracentrifugation and

resuspension processes (**Fig. 1c**). Importantly, >85-90% sEVs and m/lEVs maintained their

membrane integrity after three consecutive freeze-thaw cycles confirming the lack of significant

membrane damage during and upon revival from storage conditions (**Fig S1i,j**). In addition, EVs

lysed with Triton X-100 showed less than 10% calcein-positive particle counts demonstrating the

specificity of calcein signal intensities associated with the intact EVs.

We additionally performed an ATP assay to determine if EVs retained their functionality

upon revival from frozen storage conditions. HBMECs exposed to OGD conditions for 24 h

showed about a 60% reduction in relative ATP levels compared to normoxic HBMECs (**Fig. S2**).

Importantly, HBMECs treated with freshly isolated sEVs at 50 µg EV protein in OGD conditions

showed about a three-fold ($p<0.0001$) increase in relative ATP levels compared to untreated cells

(**Fig. S2**). OGD HBMECs treated with 50 µg sEV protein (sEVs were revived after three

subsequent freeze/thaw cycles) showed a similar increase in relative ATP levels compared to

untreated cells (**Fig. S2**). Importantly, there were no statistical ($p>0.05$) differences between

relative ATP levels of HBMECs treated with either freshly isolated sEVs or sEVs post-

freeze/thaw. Like sEV, OGD HBMECs treated with freshly isolated m/lEV showed about a four-

fold and statistically significant ($p<0.0001$) increase in relative ATP levels compared to

untreated cells (**Fig. S2**). Importantly, there was no statistical ($p>0.05$) difference between

relative ATP levels of HBMECs treated with either freshly isolated m/lEV or freeze/thaw-

subjected m/lEV. These observations suggest that EVs retain their functionality during frozen

storage conditions.

3.2. m/lEV contain mitochondria and transfer their mitochondrial load to recipient

primary HBMECs and hCMEC/D3 cells.

TEM images of cross-sectioned sEVs and m/lEVs are shown in **Fig. 1d.** The green arrow indicates m/lEV membranes, whereas the yellow arrow points to mitochondria in m/lEV. The high-speed centrifugation and extensive sample processing during the sample pelleting step prior to cutting ultrathin sections for TEM imaging likely resulted in the observed mitochondrial morphology. We noted mitochondria structures with a characteristic double membrane that were highly similar to published reports (30-33). It should also be noted that these previous reports have demonstrated functional activity of the m/lEV/extracellular mitochondria. TEM images of sectioned sEV showed vesicles with particle diameters ranging from 20-100 nm and heterogeneous shapes (**Fig. 1d**). The core of sEVs appeared less dense suggesting that sEVs may not contain organelles like mitochondria and may instead contain cytosolic components. Our data consistently showed that m/lEV- but not sEV sections contained mitochondria.

We performed western blot analysis of sEV and m/lEV lysates for determining the presence of mitochondrial proteins, including ATP5A (subunit of the mitochondrial ATP synthase) and TOMM20, an outer mitochondrial membrane protein. Our data showed that while both sEVs and m/lEVs contained ATP5A protein, TOMM20 was selectively present in m/lEVs. This suggests that while sEVs contain mitochondrial proteins, only m/lEVs contain mitochondria. The western blot aligned with the TEM data confirming the presence of mitochondria in m/lEVs. Collectively, our data demonstrate the presence of mitochondria in the m/lEVs but not sEVs. We showed presence of the characteristic exosomal marker protein CD9 tetraspanin in sEVs (**Fig.**

1e). GAPDH was present in both sEVs and m/lEVs indicating the presence of cytosolic proteins

in sEVs and m/lEVs (**Fig. 1e**).

Next, we wanted to determine if EVs can transfer their mitochondrial load into recipient

primary human brain microvascular endothelial cells (HBMECs) (**Fig 2a, b**). We isolated sEVs

and m/lEVs from hCMEC/D3 cells pre-stained with Mitotracker Red (MitoT-red). HBMECs

were treated with MitoT-red-sEV and MitoT-red-m/lEV at 30 to 150 µg EV protein/well for 72 h

and MitoT-red signals in the recipient HBMECs were measured using flow cytometry. The

intensity of MitoT-red-EVs in the recipient HBMECs (**Fig. 2a, b**) was analyzed using histogram

plots. Untreated HBMECs cells were used as control and were gated for data analysis (**Fig. 2a,**

b). Cells pre-stained with MitoT-red were used as a positive control and showed about 82%

MitoT signals suggesting the presence of polarized mitochondria (**Fig. 2a, b**). Cells treated with

unstained m/lEV did not show any MitoT-red signals in HBMECs suggesting the absence of

non-specific MitoT signals (**Fig. 2a**). Cells treated with MitoT-red-m/lEV at a low dose of 10 µg

dose showed about 32% MitoT-positive signals suggesting mitochondrial transfer into the

recipient HBMECs (**Fig. 2a, c**). The fraction of cells showing MitoT+ve signals increased from

61% at 75 µg to 68% at 150 µg m/lEV dose (**Fig. 2a, c**). sEVs at lower doses of 30 and 75 µg

protein per well showed <5% of mitochondrial transfer that increased to only about 13% at the

150 µg dose (**Fig. 2b, c**). Cells treated with EV (sEV: m/lEV 1:1) at 75 µg EV protein showed

about 54% MioT-red signal intensity suggesting that inclusion of m/lEVs in the EV mixture

treatment increased mitochondrial transfer to HBMECs (**Fig. 2b, c**). As expected, m/lEVs

showed a greater transfer of mitochondria into recipient HBMECs compared to sEVs consistent

with the presence of mitochondria in m/lEVs (**Fig. 1d, e**).

Fig. 2. Transfer of EV mitochondria into recipient HBMECs cells at varying doses. HBMECs cells were cultured in 48-well plates for 48 h in a humidified incubator. Cells were then incubated with the indicated amounts of MitoTracker red-labeled samples: (MitoT-red)-sEV, MitoT-red-EV (at a 1:1 sEV:m/lEV ratio, collectively referred to as EVs) and MitoT-red-m/lEV diluted in complete growth medium for 72 h. Post-incubation, the cells were washed, collected, and run through Attune NxT flow cytometer. The histograms of MitoT-red events of hCMEC/D3 cells treated at indicated doses of MitoT-red-m/lEVs (**a**) and sEV (**b**) were detected using a 674/10-nm side scatter filter in an Attune flow cytometer. Unstained EVs were used to gate the histograms for estimating percentage MitoT-red-positive counts. (**c**) Quantification of sEV and m/lEV-mediated mitochondria transfer in recipient HBMECs. Data represent mean±SD of n=3.

We also studied EV-meditated mitochondrial transfer into recipient hCMEC/D3 cells (**Fig. S3**). Consistent with our observations from the HBMECs, hCMEC/D3 cells treated with MitoT-red-m/lEV showed a dose-dependent increase in mitochondrial transfer from 48% at 30 µg to

68% at 150 µg dose (**Fig. S3a**). We noted a dose-dependent increase in the mitochondrial

transfer that increased from 13% at 30 µg sEV to about 33% at the 150 µg dose (**Fig. S3b**).

Interestingly, sEV-mediated a greater extent of mitochondrial transfer into hCMEC/D3 cells

compared to levels noted in primary HBMECs (33 vs. 13%) (**Fig. 2b and Fig. S3b**).

3.3. EVs transferred polarized mitochondria to recipient primary HBMECs.

We tested if the mitochondrial load in EVs can transfer into and subsequently colocalize with

the mitochondrial network of the recipient BECs. We first determined the uptake of EV-

associated mitochondria in recipient primary HBMECs and hCMEC/D3 cells. The recipient

HBMECs or hCMEC/D3 cells were treated with Mitotracker red-stained EVs (MitoT-red-EV)

for 72 h prior to observing under an epifluorescence microscope. No MitoT-red-associated

signals were observed in unstained/untreated primary HBMECs suggesting that only MitoT-red

specific signals were detected under the Cy5 channel (**Fig. 3a**). MitoT-red-sEV at 10 and 25 µg

doses did not show positive signals in HBMECs for 48 h, however, MitoT-red-sEV at 50 µg

doses showed faint intracellular Cy5 signals at 48 and 72 h suggesting low levels of uptake after

48 h (**Fig 3a**). In contrast, strong intracellular signals in HBMECs treated with MitoT-

sEV+m/lEV at the 25 µg (**Fig. 3a**) dose suggested that the inclusion of m/lEVs in the EV

mixture led to an efficient uptake of polarized mitochondria into HBMECs. Moreover, the

increase in sEV+m/lEV-mediated mitochondrial transfer was statistically significant (p<0.01) at

48 h compared to sEVs alone (**Fig. 3b**).

 EV-mediated mitochondrial transfer in recipient HBMECs

b. **EV-mediated mitochondrial transfer quantification**

Fig. 3. Transfer of EV mitochondrial load into the recipient HBMEC at varying doses and incubation times. (a) HBMECs were cultured in 96-well plates until 80% confluency in a humidified incubator. Cells were then incubated with the indicated amounts of MitoTracker red-labeled samples: (MitoT-red)-sEVs, MitoT-red-EV (at a 1:1 sEV: m/lEV ratio), MitoT-red-m/lEVs diluted in complete growth medium for 24, 48, and 72 h. Post-incubation, the cells were washed and incubated with phenol-red-free growth medium. Intracellular MitoT-red-sEV/sEV:m/lEV/m/lEV signals were observed under an Olympus IX 73 epifluorescent inverted microscope using Cy5 channel (purple puncta) at 20x magnification. Scale bar: 50 µm. **(b) EV-mediated mitochondrial transfer quantification**. HBMECs were treated with the indicated samples and doses for 24, 48, and 72 h. At each time point, from each control and treatment group, at least three images were acquired and the total sum of grayscale signal intensities in the Cy5 channel were estimated using Olympus CellSens software. The measured intensities were normalized with those of the untreated cells.

Interestingly, MitoT-red-m/lEV at a dose as low as 10 µg showed efficient uptake in HBMECs 72 h post-exposure. The levels of uptake increased significantly (p<0.0001) as the dose of MitoT-red-m/lEV increased from 10 to 50 µg at 48 h (**Fig. 3a,b**). The superior transfer of m/lEV-mediated mitochondria into HBMECs is likely due to a greater enrichment of functional mitochondria in the m/lEVs compared to sEVs. Therefore, we expected that m/lEVs may increase the cellular bioenergetics of the recipient BECs at lower doses compared to sEVs. We further confirmed the effect of dose and type of EV subtype (m/lEVs vs. sEVs) on the transfer of mitochondria into hCMEC/D3 cells (**Fig. S4a**). Similar to the observations noted in the primary HBMEC cultures at 72 h post-exposure, sEVs at 50 µg protein/well showed faint MitoT-red+ signals in the recipient hCMEC/D3 cells, whereas cells treated with m/lEVs showed a dose-dependent increase (p<0.01) in mitochondrial transfer compared to sEV-treated cells (**Fig. S4a,b**).

3.4. EV-transferred mitochondria colocalized with the mitochondrial network in the recipient BECs.

We wanted to determine if EV-mitochondria colocalized with the mitochondrial network of

the recipient BECs. We isolated MitoT-red-stained polarized mitochondria from the donor BECs

and the recipient BEC mitochondria were pre-stained using Mitotracker green (MitoT-green).

Post-treatment of the recipient BECs, the overlap of these fluorescent signals was observed under

an epifluorescent microscope. HBMECs and hCMEC/D3 cells pre-stained with Mitotracker

green were subsequently incubated with MitoT-red-sEVs and MitoT-red-m/lEVs at 10, 25, and

50 µg doses for 72 h. Cytosolic, diffuse MitoT-green signals were observed under the GFP

channel whereas punctate MitoT-red EV signals were captured under the Cy5 channel (purple

puncta). The pre-staining of HBMECs with Mitotracker green resulted in robust fluorescent

signals for 72 h and incubation with Mitrotracker red-stained EVs did not affect the Mitotracker

green signals.

a.

EV mitochondria colocalization with recipient HBMEC mitochondria

b. **Quantitative analysis of EV mitochondria colocalization with recipient HBMEC mitochondria**

Fig. 4. Colocalization of EV mitochondrial load with the recipient HBMEC mitochondria.
(a) HBMECs were cultured in 96-well plates until 80% confluency in a humidified incubator. HBMECs were stained with Mitotracker Green for 30 min. Post-staining, the cells were washed and treated with the indicated doses of MitoT-red-sEV, MitoT-red-EVs (at a 1:1 sEV: m/lEV ratio, collectively referred to as EVs), and MitoT-red-m/lEV for 72 h. Untreated cells and cells stained with MitoTracker Green only were used as controls. Post-incubation, the treatment mixture was replaced with phenol-red-free growth medium. The Mitotracker green staining in recipient HBMEC was acquired using the GFP channel, whereas the purple fluorescence associated with EV-transferred mitochondrial load was captured using Cy5 channel in an Olympus IX 73 epifluorescent inverted microscope. Colocalization of the mitochondrial signals was confirmed by the presence of yellow signals in the overlay images. Scale bar: 50 µm. (**b**) Pearson's correlation coefficient was obtained from the overlay images of Cy5 and GFP channels at constant signal intensities for both the channels using Cell Insight CX7 HCS microscope. Data are presented as mean±SD (n=3 images per treatment group). *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

The absence of GFP and Cy5 signals in the untreated cells suggested the absence of non-specific signals at the respective channel settings (**Fig. 4a**). The cells pre-stained with MitoT-green alone showed green cytosolic signals associated with the recipient mitochondria (**Fig. 4a**). MitoT-red-m/lEVs showed a greater intensity of Cy5 signals at all tested doses compared to MitoT-red-sEVs, once again demonstrating that m/lEV contain a greater mitochondrial load (**Fig. 4a**). The overlay images of recipient HBMEC mitochondria and EV-associated polarized mitochondria in the m/lEV-treated cells showed considerably higher colocalization compared to sEV-treated cells (**Fig. 4a**). The Pearson's correlation coefficient (PCC) of GFP and Cy5 channel intensities demonstrated that m/lEV exposure resulted in a statistically significant (p<0.0001), greater degree of mitochondria colocalization compared to sEVs (**Fig. 4b**). Similar to the primary HBMECs, hCMEC/D3 cells pre-stained with Mitotracker green and treated with MitoT-red-m/lEVs showed a greater degree of colocalization indicated by the overlap of MitoT-green and MitoT-red signals compared to sEV-treated cells (**Fig. S5**).

We further confirmed the colocalization of EV-mitochondria with mitochondria of the

recipient BECs using an orthogonal technique that allowed us to confirm that the EV-

mitochondria colocalized with the mitochondrial matrix in the recipient BECs. HBMECs were

first transduced with CellLight Mitochondria-GFP (CellLight-MitoGFP) before MitoT-red-EV

treatment. Despite the low absolute frequency of transduction (34), the cells showed strong GFP

fluorescence suggesting that CellLight-MitoGFP transduction effectively tagged the alpha

pyruvate mitochondrial matrix protein in the recipient HBMECs (**Fig. S6**). Similar to Figure 4,

MitoT-red-m/lEV at a dose of 50 µg showed efficient transfer of mitochondria (purple puncta)

into the recipient HBMECs. The EV MitoT-red puncta signals colocalized with recipient

mitochondria (CellLight-MitoGFP) at 72 h post-exposure. Notably, cells exposed to MitoT-red

m/lEVs showed stronger Cy5 signals compared to MitoT-red-sEV-treated cells confirming again

that m/lEVs contain more mitochondrial load compared to sEVs. The Cy5 signal intensity in

m/lEV-treated cells was considerably increased at 72 h post-exposure compared to the 24 h time

point suggesting that 72 h is an optimal exposure period for m/lEVs internalization into the

recipient HBMECs (**Fig. S6**). Importantly, the overlap of the structural protein-tagged recipient

BEC mitochondria with the MitoT-red-stained m/lEVs indicates that the m/lEV-delivered

mitochondria colocalized with the recipient cell mitochondria.

3.5. Naïve EVs increased HBMEC ATP levels under normoxic and hypoxic conditions.

The effect of BEC-derived EVs on the ATP levels in recipient primary HBMECs were first

evaluated under normoxic conditions using an ATP assay. Primary HBMECs were treated with

sEVs and m/lEVs at 10, 25, and 50 µg EV protein per well for 24, 48, and 72 h. **Fig 5a** shows

that the increase in HBMEC ATP levels upon sEV and m/lEV treatment for 24 h was not

statistically significant ($p > 0.05$). Importantly, HBMECs treated at a dose of 25 µg sEVs for 48 h

showed a significant ($p < 0.01$) increase in ATP levels compared to untreated cells. Interestingly,

cells treated with m/lEVs at 25 and 50 µg doses showed a dose-dependent and significant

($p < 0.0001$) increase in relative ATP levels compared to untreated cells. In addition, the m/lEV-

mediated increase in ATP levels was significantly ($p < 0.05$) higher compared to sEVs after 48 h

exposure (**Fig. 5b**). At 72 h post-exposure, m/lEVs at 25 and 50 µg doses showed a dose-

dependent and significant ($p < 0.01$) increase in relative ATP levels compared to untreated cells

under normoxic conditions (**Fig. 5c**). In contrast, sEVs did not show any significant increase in

relative HBMEC ATP levels compared to the control. In addition, m/lEV-treated HBMECs

showed significantly higher ATP levels compared to sEVs at 72 h at all tested doses. The data

suggested that the m/lEV-mediated significant increase in relative ATP levels at 48 h (**Fig. 7b**)

and 72 h (**Fig. 7c**) may likely be due to their innate mitochondrial load—including mitochondria,

mitochondrial DNA, and mitochondrial proteins.

Fig. 5. EV-mediated increase in HBMEC ATP levels and mitochondrial respiration during normoxic and hypoxic conditions. HBMEC cells were cultured in the 96-well plates until 80% confluency in a humidified incubator. (**a-c**) Confluent monolayers were treated with sEV, m/lEV, and EVs (sEV: m/lEV 1:1) at the indicated amounts for 24 (**a**), 48 (**b**), and 72h (**c**). Polyethyleneimine, PEI, at 50 µg/mL was used as a positive control for the ATP assay. Post-treatment, cells were incubated with a 1:1 mixture of fresh growth medium and Cell titer Glo reagent. The relative luminescence units (RLU) of the samples were measured using a SYNERGY HTX multimode plate reader at 1s integration time. Relative ATP levels were calculated by normalizing the RLU of treatment groups to the RLU of control, untreated cells. (**d**) Confluent HBMECs were treated with the indicated doses of sEV and m/lEV in OGD medium and cell viability was measured 24 h post-treatment while untreated cells were used as a control. Data represent mean ±SD (n=3). (**e-f**) *EV-mediated increase in recipient cell mitochondrial respiration and glycolysis capacity*: Cells were cultured in a Seahorse XF96 plate for 4 days at 20,000 cells/well. sEV, and m/lEV were diluted in complete growth medium at 3.4 µg EV protein/well and cells were incubated in a humidified incubator for 24, 48, and 72 h. Post-treatment at each time point, the medium was replaced with DMEM and maximum oxygen consumption rate (OCR) (**e**), and glycolytic capacity (**f**) by measuring extracellular acidification rate (ECAR) were analyzed using the Seahorse XFe96 analyzer. Data represents mean±SEM (n=3). * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

We then exposed primary HBMECs under oxygen-glucose deprived (OGD) conditions for

24 h to simulate ischemic conditions (**Fig. S7**). We studied the effects of naïve sEVs and m/lEVs

on the cell survival of OGD-exposed primary HBMECs. HBMECs in OGD medium were

incubated with sEVs and m/lEVs at 10, 25, and 50 µg EV protein/well for 24 h. The cell viability

of EV-treated HBMECs was compared with untreated HBMECs maintained in OGD medium

(**Fig. 5d**). sEV and m/lEV at all treated doses showed a significant (p<0.0001) increase in cell

viability compared to control, untreated HBMECs (**Fig. 5d**). In addition, an increase in sEV and

m/lEV dose from 10 to 25 µg EV protein/well showed a significant (p<0.0001) increase in ATP

levels. HBMECs treated with m/lEVs at 50 µg/well showed a maximum, ca. five-fold increase in

ATP levels compared to untreated cells. Importantly, m/lEVs showed a significantly (p<0.0001)

higher HBMEC cell survival rate compared to sEVs at the same dose (25 and 50 µg) suggesting

that m/lEVs outperformed sEVs in increasing HBMEC cellular energetics under ischemic

conditions. We also confirmed that the EV-mediated increase in endothelial cell viability under

OGD conditions in the recipient hCMEC/D3 cells (**Fig. S8**). Consistent with the primary

HBMECs, hCMEC/D3 cells treated with sEVs and m/lEVs at 10, 25, and 50 µg EV protein/well

showed about a three to four-fold increase in endothelial ATP levels compared to untreated cells.

Moreover, the EV-mediated increases in ischemic hCMEC/D3 cell survival were dose-

dependent. Lastly, m/lEV-treated ischemic hCMEC/D3 cells showed a greater increase in ATP

levels compared to sEV-treated cells (**Fig. S8**).

3.6. EVs increased the oxidative phosphorylation and glycolytic functions of recipient BECs

under normoxic conditions.

The mitochondrial function of hCMEC/D3 cells treated with EVs under normoxic conditions

was evaluated using Seahorse analysis by measuring their oxygen consumption rate (OCR).

hCMEC/D3 cells were treated with sEVs and m/lEVs at 3.4 µg protein/well in complete growth

medium for 24, 48, and 72 h. sEV or m/lEV-treated cells did not show changes in maximal OCR

compared to untreated cells at 24 and 48 h post-exposure (**Fig. 5e**). In contrast, BECs exposed to

sEVs and m/lEVs for 72 h showed a significant ($p<0.0001$) increase in maximum OCR

compared to control, untreated cells. Importantly, the m/lEV-mediated increase in OCR was

significantly ($p<0.05$) higher compared to sEV-treated cells suggesting that m/lEVs

outperformed sEVs in increasing the recipient BECs' mitochondrial function. The m/lEV-

mediated increase in mitochondrial function was consistent with the m/lEV-mediated increase in

cell viability (**Fig. 5c**), intracellular uptake of m/lEV mitochondria (**Fig. 2-3**), and the co-

localization of m/lEV-associated mitochondria with the recipient BEC's mitochondrial network

(**Fig. 4**).

We further evaluated the effects of EVs on non-mitochondrial energy generation pathways

such as glycolytic capacity in the recipient BECs. Extracellular acidification rate (ECAR) is a

key indicator of cellular glycolysis and can be determined in real-time by measuring free protons

in a Seahorse plate transient microchamber (35). ECAR (basal glycolysis rate and glycolytic

capacity) was measured in hCMEC/D3 cells treated with sEVs and m/lEVs (**Fig. 5f**). We did not

note any changes in the glycolytic capacity of hCMEC/D3 cells pre-treated with sEVs and

m/lEVs for 24 and 48 h compared to untreated cells. However, treatment with m/lEVs for 72 h

showed a significantly ($p<0.05$) greater glycolytic capacity compared to untreated cells and sEV-

treated cells (**Fig. 5f**). To summarize the first part of this study, our data demonstrated that (1)

m/lEVs but not sEVs contain mitochondria, (2) m/lEVs outperformed sEVs in transferring

mitochondrial components to recipient BECs and (3) m/lEVs resulted in a greater magnitude of

relative ATP levels and mitochondrial functions (OCR and ECAR) compared to sEV-treated

BECs.

The second goal of the present work was to evaluate the effects of an exogenous HSP27

protein formulated with EVs in an OGD-exposed BEC model of ischemia/reperfusion injury. We

formulated HSP27 protein with EVs and a synthetic cationic polymer, PEG-DET. We studied the

physicochemical characteristics of the formed mixtures/complexes and evaluated their effects on

the paracellular permeability of small and large molecular mass fluorescent tracers across

primary HBMECs exposed to OGD.

3.7. Exogenous HSP27 protein formed complexes with PEG-DET, EVs, and PEG-DET-

EV mixtures.

The complexation of HSP27 protein with PEG-DET and EVs was confirmed by studying the

electrophoretic mobility of HSP27 in a native polyacrylamide gel electrophoresis (PAGE) setup.

Native recombinant human HSP27 at the running buffer of pH 8.3 carries a net negative charge

(estimated charge: -4.2 mV) (36), and therefore, it migrated from the loading spot towards the

anode during electrophoresis (**Fig. 6a**). First, the complexation of PEG-DET with HSP27 was

studied by comparing the relative changes in HSP27 band densities at polymer: protein

weight/weight (w/w) ratios ranging from 0.05:1 to 20:1 (**Fig. 6a,b**). Compared to native HSP27

(100%, **Fig. 6b**), the relative band density of PEG-DET/HSP27 complexes at w/w 0.2:1 was

considerably reduced to about 43%. As the w/w ratios increased, there was a gradual and

significant decrease in HSP27 band densities (**Fig. 6a,b**). At PEG-DET/HSP27 w/w 10:1 and

20:1, the mean HSP27 band density decreased to nearly 25% suggesting that PEG-DET formed

electrostatic complexes with HSP27 at physiological pH. The free polymer did not show any

non-specific staining at w/w 1 and 20:1.

hCMEC/D3 BEC-derived sEVs and m/lEVs were mixed with HSP27 at EV protein/HSP27

protein w/w ratios of 10:1 and the electrophoretic mobility of HSP27 was studied using native

PAGE (**Fig. 6c-d**). Mixtures of HSP27 with EVs did not affect the migration of HSP27 at the

tested w/w ratios (**Fig. 6c,**). The signal intensity of free/native HSP27 was set as 100%, and the

relative band density of HSP27 in the different EV/HSP27 mixtures was compared with the band

density of native HSP27. Albeit the bands look similar, densitometry analysis showed that

sEV/HSP27 and m/lEV/HSP27 mixtures decreased HSP27 band densities to about 77.5±20%

and 84.1±2.4% compared to native HSP27 (100%), suggesting weak interactions between EVs

and HSP27 (**Fig. 6c-d, and Fig. S9**). We interpret this reduction in band intensity as a

consequence of HSP27 binding to EV membranes as opposed to intravesicular loading of

HSP27.

Fig. 6. Formation of EV- and PEG-DET/HSP27 complexes. (**a**) Native polyacrylamide gel electrophoresis (PAGE) for PEG-DET/HSP27 complexes. Native HSP27, PEG-DET/HSP27 at indicated weight ratios, and free PEG-DET polymers were mixed with 1x native sample buffer and loaded in an SDS-free 4-10% polyacrylamide gel at 1 μg HSP27 per lane. (**c**) Native PAGE

for hCMEC/D3-derived EV/HSP27 complexes: Native HSP27 and mixtures of sEV/HSP27 , and
m/lEV/HSP27 at 10:1 weight/weight (w/w) ratios were loaded in a SDS-free 4-10%
polyacrylamide gel at 1 µg HSP27 per lane. Free sEV and m/lEV equivalent to the amounts in
10:1 w/w complexes were used as controls. Native PAGE for (PEG-DET/HSP27)/EV ternary
mixtures. PEG-DET/HSP27 complexes were prepared at 20:1 and 30:1 w/w ratios followed by
incubation with 10 µg of EVs. The indicated samples were loaded in the gel at 1 µg HSP27/lane.
Each gel was run at 100 V for 2 h and stained using Biosafe Coomassie G250. The gel was then
scanned at 800 nm using an Odyssey imager at intensity setting 5. (**b,d**) Densitometry analysis
was performed by measuring band densities of HSP27 in the different experimental groups in
comparison to the band density of native HSP27 in the respective gel using Image Studio 5.0
software *p<0.05 (**e-h**) *Physicochemical characterization of HSP27 complexes with PEG-DET*
and EVs. Average particle diameters (**e**), and zeta potentials (**f**) of PEG-DET/HSP27 complexes
and EV/HSP27 mixtures at the indicated weight ratios. Average particle diameters (**g**), and zeta
potentials (**h**) of PEG-DET/EV and (PEG-DET/HSP27)/EV ternary mixtures at the indicated
weight ratios. The samples containing 1 µg HSP27 protein were diluted to 50 µL in 10 mM
HEPES buffer pH 7.4 for particle diameter measurements. The diluted samples were further
diluted to 800 µL in 10 mM HEPES buffer pH 7.4 for zeta potential measurements. Data
represent mean±SD (n=3). * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

We then formulated ternary mixtures of PEG-DET/HSP27 complexes with EVs, and the

resulting changes in HSP27 band intensity were studied using native PAGE followed by

densitometry analysis (**Fig. 6c**). sEVs, m/lEVs, and EV/HSP27 at w/w 10:1 showed about 20-

40% reduction in band density whereas PEG-DET/HSP27 at w/w 20:1 and 30:1 showed >90%

HSP27 complexation compared to native HSP27 (**Fig. 6d**). The HSP27 band density was

decreased by 40-50% when PEG-DET/HSP27 at w/w 20:1 were incubated with 10 µg of sEV,

m/lEV, and sEV+m/lEV 1:1 (EV). The % extent of reduction in HSP27 band density of (PEG-

DET/HSP27)/EV complexes ranged among the values noted in the case of PEG-DET/HSP27 and

EV/HSP27 mixtures suggesting the competitive binding of negatively-charged EVs and HSP27

with the positively charged PEG-DET. To summarize, HSP27 protein bound to EV membranes,

formed electrostatic complexes with PEG-DET, and formed ternary mixtures with PEG-DET and

EVs.

3.8. Physicochemical characterization of the formed HSP27 complexes/mixtures

Particle diameters and surface charge of the formed complexes/mixtures were measured using dynamic light scattering (**Fig. 6e-h**). The average diameter of native HSP27 protein was about 85 nm (**Fig. 6e**). The average zeta potential of native HSP27 (1 µg/mL in 10 mM HEPES buffer, pH 7.4) was about -9 mV suggesting that HSP27 exerts a net negative surface charge under physiological conditions (**Fig. 6f**). The average particle diameter of PEG-DET/HSP27 complexes at 10:1 w/w ratio was over 1000 nm with a broad dispersity index. As the w/w ratio increased from 20:1 to 30:1, the particle diameter significantly (p<0.05) decreased from about 359 nm to 205 nm (**Fig. 6e**). PEG-DET/HSP27 complexes showed a unimodal size distribution in the intensity plots (**Fig. S10a**). PEG-DET/HSP27 complexes at 10:1 w/w ratio shifted the zeta potential of the native HSP27 protein from -9 mV to +10 mV confirming the electrostatic complexation of PEG-DET with HSP27 (**Fig. 6f**). The zeta potentials, however, did not change for w/w ratios of 20:1 and 30:1 (**Fig. 6f**) suggesting that the excess polymer do not bind to HSP27 protein.

sEV and m/lEV showed diameters of about 136 and 225 nm, respectively (**Fig. 6e**). sEV showed a bimodal particle size distribution compared to m/lEVs (**Fig. S10c,e**). sEV/HSP27 or m/lEV/HSP27 mixtures at 10:1 w/w ratio did not change the particle diameter compared to naïve EVs (**Fig. 6e**). The zeta potential of m/lEV/HSP27 mixtures shifted towards near-neutral values compared to naïve m/lEVs and native HSP27 protein indicating weak interactions between m/lEVs and HSP27 (**Fig. 6f**).

Next, PEG-DET was mixed with sEVs and m/lEVs at PEG-DET/EV w/w ratios 2:1 and 3:1,

and the resulting changes in particle sizes and zeta potentials were compared with naïve sEVs

and m/lEVs (**Fig. 6g,h**). The complexation of PEG-DET to sEV showed a considerable increase

in particle size from 134 nm to 245 nm (**Fig. 6g**). The shift in mean zeta potential from -1.55 mV

to -0.18 mV suggested the electrostatic interactions of PEG-DET and sEV (**Fig. 6h**).

Furthermore, the z-average particle diameter of (PEG-DET/HSP27)/sEV significantly (p<0.01)

increased to about 400 nm with a near-neutral zeta potential suggesting the interactions of PEG-

DET/HSP27 and sEVs (**Fig. 6g,h**). A similar trend was observed for m/lEVs where the particle

diameter gradually and significantly (p<0.0001) increased from naïve m/lEVs (216 nm), PEG-

DET/m/lEV (431 nm), and (PEG-DET/HSP27)/m/lEV (991 nm) (**Fig. 6g**). A shift in zeta

potential was observed from -4 mV for naïve m/lEV to 1.29 mV for PEG-DET/m/lEV, and 0.75

mV for (PEG-DET/HSP27)/m/lEV confirming the electrostatic interactions of PEG-DET and

m/lEV (**Fig. 6h**). In addition, an increase in PEG-DET to EV w/w ratio from 2:1 to 3:1 showed a

reduction in z-average diameter (**Fig 6g**). A slight decrease in zeta potential at PEG-DET/EV at

w/w 3:1 suggested that increasing PEG-DET amounts may increase the extent of complexation

with EVs (**Fig. 6h**). The representative distribution plots of PEG-DET/EVs and (PEG-

DET/HSP27)/EVs were shown in **Fig. S11**.

To summarize, PEG-DET/HSP27 complexes showed a w/w ratio-dependent decrease in

particle diameter and an overall positive surface charge. EV/HSP27 mixtures showed

physicochemical characteristics more or less similar to naïve EVs. The observed changes in

particle diameter and zeta potential of (PEG-DET/HSP27)/EVs confirmed the complexation of

EVs with PEG-DET/HSP27 complexes.

3.9. PEG-DET/HSP27 complexes and EV/HSP27 mixtures were cytocompatible with primary HBMECs.

We performed an ATP assay to determine the cell viability of primary HBMECs treated with native HSP27, m/lEV, or sEV/HSP27 mixtures and PEG-DET/HSP27 complexes. The cell viability of treatment groups was calculated using *Equation 1*. The average viability of cells treated with native HSP27 was 108.7% and there were no significant ($p > 0.05$) differences between control, untreated cells, and native HSP27-treated groups suggesting that native HSP27 at 2 µg/well was well tolerated by HBMEC sunder normoxic conditions for 72 h (**Fig. S12**). The cell viability of HBMECs increased significantly ($p < 0.01$) when treated with sEV/HSP27 (124.7%), m/lEV/HSP27 (123.1%), and sEV+m/lEV/HSP27 (116.8%) mixtures at 10:1 w/w ratio compared to untreated cells suggesting that EV/HSP27 mixtures were well tolerated by HBMECs for 72 h (**Fig. S12**). The EV/HSP27 mixture-mediated increase in cell viability can be correlated with a significant increase in HBMEC ATP levels that was observed when cells were treated with naïve sEV (121%), m/lEV (126.4%), or sEV+m/lEV (120.1%) at amounts equivalent those present in the EV/HSP27 mixtures. HBMECs treated with PEG-DET/HSP27 complexes at w/w 20:1 showed an average cell viability of 100.4% suggesting that PEG-DET/HSP27 complexes were cytocompatible. Free PEG-DET polymer was also well tolerated by HBMECs for 72 h. Polyethyleneimine, a positive control, at 50 and 100 µg/mL concentrations showed a significant ($p < 0.0001$) reduction in HBMEC viability indicating that the assay was responsive to the toxicities.

3.10. HSP27 complexes attenuated the hypoxia-induced increase in tight junction permeability in primary HBMECs

3.10.1. Paracellular permeability of <u>4.4 kD TRITC-Dextran (a small molecule tracer)</u> in HBMEC culture inserts pre-treated with HSP27 complexes

We measured the paracellular permeability of 4.4 kD TRITC-Dextran to evaluate the effect of (PEG-DET/HSP27)/EV and EV/HSP27 pre-treatment on the diffusion of a small molecule-mimic across primary HBMECs during normoxic, OGD, and OGD/reperfusion conditions. HBMECs were treated at a dose of 2 µg/well HSP27 complexed with PEG-DET at 20:1 w/w ratio and sEV, m/lEV, EVs (sEV: m/lEV=1:1) at 10:1 w/w ratio. Native HSP27, free PEG-DET, and naïve sEV, m/lEV, and EVs were used as controls. The difference in the diffusion rate of 4.4 kD dextran between untreated and HSP27 mixtures-treated cells during normoxic conditions and pre-OGD phase was not statistically significant (p>0.05) with the exceptions of m/lEV/HSP27- and EV-treated groups (**Fig. S13a**). During the OGD phase, PEG-DET/HSP27, and (PEG-DET/HSP27)/EV-treated HBMECs showed a significant (p<0.05) reduction in the rates of 4.4 kD dextran diffusion compared to OGD, native HSP27, and free PEG-DET-treated HBMECs for 4 h (**Fig. 7a**). Besides, naïve sEV, m/lEV, and EV-treated HBMECs showed a significant (p<0.05) reduction in paracellular permeability for 4 h (**Fig. 7a**) of OGD exposure. (PEG-DET/HSP27)/EV, naïve sEV, m/lEV, and EV/HSP27 mixtures showed a consistent and significant (p<0.05) reduction in the diffusion rate of 4.4 kD dextran for 24 h of OGD exposure (**Fig. S13b**).

During the first hour of ischemia/reperfusion (OGD/reperfusion), HBMECs exposed to

native HSP27 and (PEG-DET/HSP27)/EV complexes showed a significant (p<0.05) reduction in

4.4 kD dextran diffusion rate compared to OGD and free PEG-DET polymer-treated cells (**Fig.**

7b). In addition, naïve sEV, EV, sEV/HSP27, m/lEV/HSP27, and EV/HSP27 pre-treated

HBMECs showed a highly significant (p<0.0001) decrease in 4.4 kD dextran diffusion rate

compared to control HBMECs (**Fig. 7b**). (PEG-DET/HSP27)/EV complexes, sEV/HSP27,

m/lEV/HSP27, and EV/HSP27 mixtures treatment resulted in a consistent and significant

reduction in the paracellular permeability of 4.4 kD dextran for 4 h of reperfusion compared to

the OGD control (**Fig. 7c,d**). In contrast, the dextran diffusion rate in other treatment groups at 2

and 4 h of OGD/reperfusion was non-significant compared to the untreated cells (**Fig. 7c,d**).

There was no significant difference in dextran diffusion rate between control and treated

OGD/reperfusion HBMECs at 24 h (**Fig. S13c**). The possible reason for the observed lack of

changes in the small molecule/4.4 kDa dextran diffusion rate during the reperfusion phase is

likely due to the reperfusion-mediated rapid recovery of ischemic HBMECs.

Paracellular permeability of TRITC 4.4 kD Dextran across primary HBMEC monolayers

Fig. 7. Paracellular permeability of TRITC 4.4 kD dextran under OGD, and OGD/reperfusion conditions in HBMEC transwell culture inserts pre-treated with HSP27 complexes. HBMECs were seeded in 24-well plates and maintained in a 37°C humidified incubator for a week. The complete growth medium was replaced with 300 µL of growth media containing indicated treatment groups for 72 h. Post-treatment, the treatment media was replaced with 300 µL of OGD medium containing 1 µM TRITC 4.4 kD dextran for 24 h. The abluminal chamber was filled with 0.5 mL of complete growth medium. Control, untreated cells were incubated in complete growth medium in a humidified incubator whereas OGD treatment groups were incubated in an OGD chamber. At 4 h post-OGD (**a**), a 500 µL volume was collected from the abluminal chamber and a fresh medium was added to the transwell inserts. Post-OGD treatment, HBMECs were washed with PBS and incubated with 300 µL of complete growth medium containing 1µM TRITC 4.4 kD dextran and incubated in a humidified incubator for 1-24h. At each time point, a 500 µL volume was collected from the abluminal chamber and fresh medium was added to the transwell inserts. The concentration of TRITC 4.4 kD Dextran was measured at 1 h (OGD/reperfusion 1h, **b**), 2 h (OGD/reperfusion 2h, **c**), and 4 h (OGD/reperfusion 4h, **d**) using a Synergy HTX multimode plate reader at 485/20 nm excitation

and 580/50 nm emission settings. The diffusion rate of TRITC 4.4 kD Dextran at each time point was calculated by normalizing the amount of TRITC Dextran per unit surface area of the insert per unit time. Data represent mean±SD (n=2). * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001, ns: non-significant.

3.10.2. Paracellular permeability of 65-85 kD TRITC-Dextran (a large molecule tracer) in HBMEC culture inserts pre-treated with HSP27 complexes

The effect of HSP27 complexed with PEG-DET and EVs on the paracellular permeability of 65-85kD TRITC-Dextran was evaluated as previously described for the small, 4.4 kDa tracer (section 3.10.1). Here, 65-85kD TRITC-Dextran was used a large molecular mass tracer simulating the diffusion of large molecules such as proteins through the damaged BBB (9). Prior to subjecting cells to OGD (pre-OGD) (**Fig. S13d**), cells treated with sEV/HSP27 mixtures showed a significant (p<0.05) reduction in TRITC-Dextran permeability compared to control, untreated cells. In addition, naïve sEV, m/lEV, and EVs showed a considerable decrease in 65-85 kD TRITC-Dextran permeability (**Fig. S13d**). The data suggested that naïve sEV and m/lEV and their HSP27 mixtures may protect the barrier properties of BEC tight junctions compared to untreated cells under normoxic conditions. Interestingly, cells treated with native HSP27, PEG-DET/HSP27, and (PEG-DET/HSP27)/EV complexes did not show any change in the TRITC-Dextran diffusion rate under normoxic conditions. The observed effects of decreased permeability was selective only for the EV-treated cells.

Fig. 8. Paracellular permeability of TRITC 65-85 kD Dextran under OGD, and OGD/reperfusion conditions in HBMEC transwell culture inserts pre-treated with HSP27 complexes. HBMECs were seeded in 24-well plates and maintained in a 37°C humidified incubator for a week. The complete growth medium was replaced with 300 µL of growth media containing indicated treatment groups for 72 h. Post-treatment, the treatment media was replaced with 300 µL of OGD medium containing 1 µM TRITC 65-85 kD dextran for 24 h. The abluminal chamber was filled with 0.5 mL of complete growth medium. Control, untreated cells were incubated in complete growth medium in a humidified incubator whereas OGD treatment groups were incubated in an OGD chamber. At 4 h post-OGD (**a**), a 500 µL volume was collected from the abluminal chamber and a fresh medium was added to the transwell inserts.

Post-OGD treatment, HBMECs were washed with PBS and incubated with 300 µL of complete growth medium containing 1µM TRITC 65-85 kD dextran and incubated in a humidified incubator for 1-24h. At each time point, a 500 µL volume was collected from the abluminal chamber and fresh medium was added to the transwell inserts. The concentration of TRITC 65-85 kD dextran was measured at 1 h (OGD/reperfusion 1h, **b**), 2 h (OGD/reperfusion 2h, **c**), and 4 h (OGD/reperfusion 4h, **d**) using a Synergy HTX multimode plate reader at 485/20 nm excitation and 580/50 nm emission settings. The diffusion rate of TRITC 65-85 kD dextran at each time point was calculated by normalizing the amount of TRITC dextran per unit surface area of the insert per unit time. Data represent mean±SD (n=2). * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$, ns: non-significant.

HBMECs pre-treated with PEG-DET/HSP27 and (PEG-DET/HSP27)/EV complexes showed a significant ($p<0.01$) reduction in the diffusion rate during 4 h OGD exposure, whereas native HSP27 and free PEG-DET did not affect the permeability of 65-85 kD TRITC-Dextran (**Fig. 8a**). sEV/HSP27 and m/lEV/HSP27 mixtures showed a significant ($p<0.0001$) reduction in 65-85 kD dextran diffusion rate compared to untreated OGD cells. Naïve sEV, m/lEV, and EVs also showed a statistically significant ($p<0.05$) reduction in dextran permeability under OGD conditions compared to untreated, OGD control (**Fig. 8a**). The data suggested that PEG-DET/HSP27 and (PEG-DET/HSP27)/EV complexes can limit the diffusion of large molecules post-ischemia. Importantly, naive sEV and m/lEV limited the dextran diffusion before and OGD conditions, whereas complexation of HSP27 with EVs showed a synergistic effect on decreasing dextran diffusion during ischemia. Notably, there was no difference in the 65-85 kD TRITC-Dextran diffusion rate between control and HBMECs treated with HSP27 complexes at 24 h of OGD exposure (**Fig. S13e**).

Post-OGD, the OGD medium was replaced with fresh complete growth medium to evaluate the effects of HSP27 on ischemia/reperfusion-mediated diffusion of 65kD TRITC-Dextran.

Exposure of cells to PEG-DET/HSP27 and (PEG-DET/HSP27)/EV complexes showed a

significant (p<0.05) reduction in dextran diffusion rate 1 h post-ischemia/reperfusion (**Fig. 8b**).

Importantly, sEV/HSP27, m/lEV/HSP27, and EV/HSP27 mixtures showed a significant

(p<0.0001) reduction in dextran diffusion rate compared to OGD control after 1 h

ischemia/reperfusion. Moreover, naïve sEVs and EVs also showed significant (p<0.05) reduction

in diffusion after reperfusion suggesting that naïve EVs increase BEC tight junction integrity

immediately after ischemia/reperfusion. In addition, their mixtures with HSP27 synergistically

reduce the large molecule infiltration across the BECs (**Fig. 8b**). Importantly, (PEG-

DET/HSP27)/EV complexes, m/lEV/HSP27 and EV/HSP27 mixtures showed a significant

(p<0.05) reduction in diffusion rate compared to OGD control 2 h post-OGD/reperfusion,

whereas changes in the diffusion rate of dextran were insignificant (p>0.05) in the other

treatment groups compared to OGD control (**Fig. 8c**). It can be inferred that in addition to the

naïve EV-mediated protection of tight junction integrity pre-OGD/normoxia, during OGD, and

OGD/reperfusion, their mixtures with HSP27 can protect BECs during OGD/reperfusion. There

were no further differences in dextran diffusion rate amongst the OGD control and treatment

groups during 4-24 h of ischemia/reperfusion (**Fig. 8d** and **S13f**).

1090

1091   **4. Discussion**

1092

The goal of this study was to study the effects of the innate EV mitochondrial load and to

determine if an exogenous protein, HSP27, can be formulated using EVs. This one-two punch

strategy allowed us to harness the innate EV mitochondrial load to increase cellular bioenergetics

and deliver an exogenous HSP27 protein to reduce paracellular permeability in ischemic BECs.

This novel approach to protect the ischemic BECs can potentially limit damage to the BBB, an

integral component of the neurovascular unit. Loss of mitochondrial function increases BBB

permeability leading to secondary injury that exacerbates post-stroke damage (37). Protection of

the BBB can limit early structural damage and prevent chronic neurological dysfunction post-

stroke (37-42).

1102

The present study investigated (1) the effects of the innate EV mitochondrial load on the

bioenergetics of recipient HBMECs under normoxic and ischemic conditions, and (2) the effects

of HSP27 formulated with EVs on the paracellular permeability of tracer molecules across

ischemic BEC monolayers. The results of our studies demonstrated that EV mitochondria,

specifically m/lEVs, transferred into and colocalized with the mitochondrial network of the

recipient BECs. As a result, EV-treated primary HBMECs demonstrated increased intracellular

ATP levels and mitochondrial respiration. Prophylactic treatment of EV/HSP27 mixtures and

(PEG-DET/HSP27)/EV complexes significantly reduced ischemia-induced paracellular

permeability of small and large tracer molecules across primary HBMEC monolayers.

1112

BECs form the foremost layer of the BBB and they contain about two to five-fold greater

mitochondrial content compared to peripheral endothelial cells (43). BEC-derived EVs contain

BBB receptors such as transferrin and insulin that enable EVs to cross the BBB for the treatment

of various neurovascular disorders (44-46). A greater mitochondrial load and their natural

affinity for BBB targeting motivated us to isolate sEVs and m/lEVs from the hCMEC/D3 BEC

cell line. We used a differential ultracentrifugation protocol, the most commonly used EV

isolation method (47), to isolate m/lEVs and sEVs from the EV-conditioned medium. Post-

ultracentrifugation and resuspension, sEVs and m/lEVs showed their characteristic particle

diameters (100-250 nm, **Fig. 1a,b**) that largely aligned with the previous reports (15, 21, 31, 48,

49).

1123

We have observed an apparent discrepancy between DLS and TEM sizing of sEVs and

m/lEVs. For sEV, TEM analysis showed a smaller particle size than DLS measurements,

whereas TEM sizing of m/lEVs was larger compared to DLS. It is important to note that sEVs

and m/lEVs are heterogenous EV subpopulations with polydispersity indices ranging from 0.3-

0.4 (**Fig. 1a**). Therefore, the observed particle size differences of EVs measured by different

techniques could be attributed to significant differences in the operating principles and sample

handling/processing conditions. DLS estimates particle hydrodynamic diameters in a solvent.

The hydrodynamic diameter includes the core of the particle and the surrounding liquid layer. On

the other hand, TEM measures the core size of an individual particle in a dried state under

vacuum. The environmental conditions may shrink the particles due to the osmotic diffusion of

solvent from the aqueous core to the surrounding space. Importantly, TEM represents the

morphology of only a small portion of the sample, which can further be affected by operator

bias.

1137

1138 Anionic phospholipid components such as phosphatidylinositol, phosphatidylserine, and

1139 glycosylated lipid derivatives exert a net negative zeta potential on the sEVs and m/lEVs

1140 membranes (**Fig. 1a**) (50, 51). The broad polydispersity indices demonstrate the natural

1141 heterogeneity of both EV sub-populations. Similar to cells and other biomolecules, it is

1142 important to preserve EV physicochemical characteristics and biological activities during storage

conditions that critically determine the scope of their therapeutic application. In our studies,

sEVs and m/lEVs isolated from conditioned medium retained their particle diameters, dispersity

indices and zeta potential after three consecutive freeze-thaw cycles (samples were frozen at -

20°C for 24 h and then were thawed at room temperature for 1 h, **Fig. S1a-f**). Jeyaram *et al.*

showed that EVs isolated from biofluids such as blood, milk, urine, and conditioned medium

preserved their physical and functional properties stored at -80°C compared to 4° and -20°C (52,

53). Moreover, freeze-thaw cycles at -20° and -80°C did not affect the stability of plasma

exosome miRNA (54).

Our TEM analysis demonstrated that m/lEVs, but not sEVs, contained mitochondria (**Fig.**
1d). Mitochondria-rich m/lEVs structures was consistent with published reports (30-33). Prior

studies have demonstrated that mitochondria or mitochondrial components such as mtDNA,

mitochondrial proteins were secreted into the extracellular milieu and transferred between cells

(31, 55-61). For instance, mesenchymal stem cell-derived-MVs transferred mitochondria into the

recipient macrophages leading to increased cellular bioenergetics (31). Guescini *et al.*

demonstrated that exosomes were released from glioblastoma and astrocytes transfer mtDNA

from glioblastoma to astrocytes (62).

We confirmed the presence of ATP5A protein (a subunit of mitochondrial adenosine

triphosphate synthase complex (15, 63)) in m/lEVs using western blotting (**Fig. 1e**) as we did in

our previous studies (15, 21). ATP5A plays important role in mitochondrial ATP production via

catalyzing the synthesis of ATP from ADP in the mitochondrial matrix during oxidative

phosphorylation (64). A considerably higher ATP5A band density in m/lEVs compared to sEVs

suggested that m/lEVs contain a greater mitochondrial load compared to sEVs. D'Acunzo *et al.*

demonstrated the presence of ATP5A in EVs isolated from mice brains using mass spectroscopy

based proteomic analysis (65). Sanson *et al.* also reported ATP5A expression as a mitochondrial

marker in stromal cell-derived EVs using western blot analysis (66). We performed western

blotting in sEVs and m/lEVs lysates to determine the presence of TOMM20, a 16 kDa outer

mitochondrial membrane protein, as an additional mitochondrial marker (**Fig. 1e**). TOMM20 is a

translocase of the outer membrane receptor that regulates the import of specific proteins from the

cytosol (67). The selective presence of TOMM20 in m/lEVs and cell lysates, but not in sEVs

lysates further confirmed that m/lEVs contains mitochondria. Silva *et al*. also demonstrated the

presence of TOMM20 in mesenchymal stromal cell-derived, mitochondria containing

extracellular vesicles (68). sEVs showed the presence of CD9 (**Fig. 1e**), a tetraspanin marker

associated with exosomal cargo selection, binding, and uptake of sEV by target cells (69).

Glyceraldehyde 3-phosphate dehydrogenase (37 kD) was used as an additional marker to

demonstrate the presence of cytosolic components in sEVs and m/lEVs (**Fig. 1e**). Collectively,

our TEM images of sectioned EVs and the presence of TOMM20 in m/lEVs indicated the

presence of mitochondria in m/lEVs, whereas sEVs contain mitochondrial proteins but not entire

mitochondria.

1183

We isolated sEVs and m/lEVs from the conditioned medium of Mitotracker deep red (MitoT-

red) pre-stained hCMEC/D3 cells. MitoT-red is a mitochondrion membrane potential-dependent

carbocyanine dye that selectively stains polarized mitochondria, and its fluorescence intensity is

reduced during mitochondrial depolarization (70, 71). We demonstrated that m/lEVs contain a

greater mitochondrial load compared to sEVs and are efficiently internalized into recipient BECs

(**Fig. 2-4 and Fig. S3,4-6**). m/lEV-mitochondria were transferred into the recipient BECs within

24 h of incubation. Increasing the m/lEV doses and incubation times significantly increased

uptake into the recipient BECs. Importantly, the m/lEV-mitochondria efficiently colocalized with

the mitochondria network in the recipient BECs. The colocalization of m/lEVs and recipient

BEC mitochondria was confirmed by the presence of overlapping signals of the EV-

mitochondria fluorescence signals with the recipient BEC mitochondria signals. We used two

orthogonal approaches to stain the mitochondrial network in the recipient BECs: Mitotracker

green and the CellLight mitochondria-GFP BacMam technique (**Fig. 4, and Fig. S6**). The

carbocyanine Mitotracker green dye stains the functional mitochondria whereas CellLight

Mitochondria-GFP BacMam comprising a fusion construct of α-pyruvate dehydrogenase and

emGFP packaged in the baculoviral vector stains a structural mitochondrial matrix protein (α-

pyruvate dehydrogenase) (34, 72). Thus, utilizing two orthogonal types of staining techniques,

we demonstrated an efficient colocalization of polarized EV mitochondria with the polarized

mitochondria in the recipient BECs (via Mito-T-green staining, **Fig. 4**) and colocalization of

functional, polarized EV mitochondria with the structurally intact mitochondria in the recipient

BECs (via CellLight Mitochondria-GFP staining, **Fig. S6**). m/lEVs and sEVs showed a dose-

dependent increase in colocalization at 72 h, specifically, the m/lEVs demonstrated a

significantly greater colocalization coefficient compared to sEVs (**Fig. 4b**). The selective

mitochondrial packaging into BEC-derived m/lEVs compared to sEVs was consistent with

published reports (31, 60, 62). Overall, our data demonstrated m/lEV-mediated transfer of

functional mitochondria and their colocalization with recipient BEC mitochondrial network.

1210

One of the main functions of mitochondria is to synthesize ATP from ADP during mitochondrial aerobic respiration, and therefore, we measured relative ATP levels in the recipient BECs treated with sEVs or m/lEVs using a Cell Glo ATP assay. Our results demonstrated that naïve sEVs and m/lEVs mediate a dose-dependent significant increase in the relative ATP levels at 48 h-post incubation (**Fig. 5a-c**). Importantly, m/lEVs outperformed sEVs in increasing recipient cell viability and the effects persisted for 72 h-post incubation (**Fig. 5c**). Islam *et al.* reported that mitochondria containing m/lEVs derived from bone marrow stromal cells increased ATP levels of alveolar epithelial cells (56). Guo *et al.* demonstrated that transfer of mitochondria isolated from donor bone marrow-derived mesenchymal cells (BMSC) into the recipient BMSCs increased cellular ATP production, proliferation, migration, and repaired bone defects *in vitro* and *in vivo* (73).

Numerous studies have demonstrated the ischemia-induced cerebral endothelial dysfunction/apoptosis and BBB breakdown (74-77). In our studies, BECs exposed to OGD exposure in a hypoxic chamber led to about 60% endothelial cell death at 24 h compared to untreated cells (**Fig 5d**). The observed data is consistent with the published reports (78, 79). Our results showed that primary HBMECs treated with naïve sEVs and m/lEVs resulted in a four to five-fold increase in endothelial ATP levels compared to control, untreated cells (**Fig. 5d**). Importantly, EV-mediated increases in ATP levels were dose-dependent and m/lEVs outperformed sEVs in rescuing the ATP levels and consequently the survival of ischemic HBMECs 24 h post-OGD (**Fig. 5d**).

We investigated the effects of m/lEV and sEV treatment on mitochondrial respiration and

glycolytic capacity function in the recipient BECs using a Seahorse setup. The state-of-art

Seahorse extracellular flux (XF) analyzer allows real-time analysis of extracellular acidification

rate (ECAR), an indicator of glycolysis, and the oxygen consumption rates (OCR), an indicator

of mitochondrial respiration in live intact cells (80, 81). We demonstrated that m/lEVs resulted in

increased OCR compared to sEVs suggesting that m/lEV mitochondria selectively increased the

recipient BEC mitochondrial function (**Fig. 5e**). Interestingly, we also noted that m/lEVs showed

a significantly ($p<0.05$) greater increase in recipient BEC glycolysis capacity compared to sEVs

(**Fig. 5f**). Phinney *et al.* demonstrated that MSC-derived EVs significantly increased maximum

OCR in the recipient macrophages compared to controls in MSC-macrophage cocultures (31).

Overall, through the use of these orthogonal tools (microscopy studies, ATP and Seahorse

assays), we have demonstrated that m/lEVs contain functional mitochondria compared to sEVs.

1245

We used EVs as a natural carrier and PEG-DET as a synthetic carrier to formulate HSP27.

The isoelectric point of human recombinant HSP27 is 5.89 (36), therefore, it exerts a net

negative charge at physiological pH 7.4 (**Fig. 6f**). The diethyltriamine side chain of PEG-DET

cationic diblock copolymer has two pKa values associated with its molecular conformation

(*gauche vs. anti*). The *gauche* conformation of DET exerts a pKa of 9.9 that induces the

formation of stable complexes with negatively charged polynucleotides at physiological pH 7.4

(17, 82). We confirmed the formation of PEG-DET/HSP27 complexes using native PAGE

followed by Coomassie staining (**Fig. 6a,b**) and dynamic light scattering (**Fig. 6e,f**). Despite the

negative surface charge of EVs and the native HSP27 protein at pH 7.4, EV/HSP27 mixtures

showed about 20% interactions in native PAGE analysis (**Fig. 6c,d**). Haney *et al.* also reported

that macrophage-derived exosomes incubated with bovine liver catalase protein did not affect the

particle diameter and dispersity indices of the EXO/catalase mixture (83). Notably, the zeta

potential of EV/HSP27 mixtures was significantly different compared to native HSP27. The

weaker interactions of HSP27 with EVs were particularly advantageous and avoided intrusive

modes of HSP27 loading (such as sonication, freeze/thaw cycles, saponin-mediated loading).

Such intrusive modes of loading may damage the EV membrane integrity and inversely impact

the functionality of innate EV cargo, specifically, their mitochondria. We further confirmed the

m/lEV and HSP27 interactions using an immunoprecipitation pull-down assay (**Fig. S14**). We

engineered ternary mixtures of EVs with PEG-DET/HSP27 complexes at different w/w ratios to

increase the HSP27 loading into hCMEC/D3-derived EVs. The positively-charged PEG-

DET/HSP27 complexes (+9 mV, **Fig. 6f**) formed an electrostatic complex with sEV and m/lEV

which was confirmed by an intermediate HSP7 band density between PEG-DET/HSP27

complexes and sEV/HSP27 mixtures (**Fig. 6c,d**) and increased resulting particle diameters (**Fig.**

6g). (PEG-DET/HSP27)/EV complexes showed a practically electroneutral zeta potential (**Fig.**

6h) that may allow longer systemic circulation *in vivo*. The inclusion of EVs in (PEG-

DET/HSP27)/EV complexes may facilitate interactions with the BBB and mediate endothelial

targeting (84) during *in vivo* delivery while the cationic PEG-DET can enhance the cellular

uptake and facilitate endosomal escape of HSP27 (17).

1274

We evaluated the prophylactic (PEG-DET/HSP27)/EVs and EV/HSP27 treatment-induced

protection of tight junction integrity by measuring the diffusion rate of hydrophilic tracers. We

used fluorescent tracers varying in molecular mass: we used 65-85 kDa TRITC-Dextran, a large

molecular weight tracer to simulate the infiltration of proteins and larger blood-borne molecules

and a 4.4 kDa TRITC-Dextran to simulate the diffusion of small molecules during

ischemia/reperfusion.

1281

We showed naïve EV, (PEG-DET/HSP27)/EV, and EV/HSP27-induced protection of tight

junction integrity against the paracellular flux of small molecules under OGD and

OGD/reperfusion conditions (**Fig. 7a-d**). (PEG-DET/HSP27)/EV complexes strengthened the

tight junctions to restrict 4.4 kD dextran entry up until 4 h of OGD exposure and as long as 4

hours of OGD/ reperfusion (**Fig. 7a-d**). Importantly, naïve EVs and their HSP27 mixtures

retarded small molecule permeability during OGD conditions and OGD/reperfusion. Notably, the

magnitude of EV/HSP27 complex-mediated protection is considerably greater than naïve EVs

demonstrating a synergistic effect of EVs and HSP27 in increasing BEC tight junction integrity

(**Fig. 7a-d**). In addition, HBMECs treated with (PEG-DET/HSP27)/EV complexes efficiently

decreased 65-85 kD TRITC-Dextran diffusion rate for 4 h of OGD exposure followed by

immediate two hours of reperfusion compared to untreated cells, native HSP27, and free PEG-

DET-treated groups (**Fig. 8a-d**). These results indicated that (PEG-DET/HSP27)/EV-mediated

efficient transfer of HSP27 into BECs restores the tight junction integrity and may protect the

BBB during ischemia/reperfusion injury. Importantly, naïve EVs and EV/HSP27 mixtures-

treated HBMECs showed efficient protection of tight junction integrity during OGD and the first

hour of ischemia/reperfusion (**Fig. 8b**).

1298

Interestingly, although ternary mixtures of (PEG-DET/HSP27)/EV showed prolonged

protection of the tight junction integrity compared to native HSP27 and PEG-DET/HSP27

complexes, the magnitude of protection is relatively lower than naïve EVs and EV/HSP27

mixtures. It is likely that the weak interactions between EV and HSP27 enable efficient EV and

HSP27 uptake and allow exerting the maximum therapeutic potential. In contrast, the stronger

electrostatic interactions in (PEG-DET/HSP27)/EV ternary complexes may impede the release of

HSP27 resulting in a slightly lower therapeutic effect compared to EV/HSP27 mixtures.

1306

5. Conclusion

This one-two punch approach using EVs increased the BEC mitochondrial function due

to the innate EV mitochondrial load, and EV/HSP27 protected tight junction integrity in

ischemic BECs. Naïve m/lEVs and sEVs increased ATP levels (albeit m/lEV showed a greater

magnitude of ATP increases), mitochondrial respiration, and glycolytic capacities in the recipient

BECs. (PEG-DET/HSP27)/EV and EV/HSP27 mixtures restored tight junction integrity in

primary human BECs by limiting the paracellular permeability of small and large molar mass

tracer molecules during ischemia/reperfusion injury. The outcome of the present study indicates

that this approach has potential to protect the BBB *in vivo* that in turn can ameliorate the long-

term neurological damage and dysfunction in rodent models of ischemic stroke.

1317

Acknowledgments

This work was supported via start-up funds for the Manickam laboratory from Duquesne

University (DU) and a 2021 Faculty Development Fund (Office of Research, DU) to the PI. We

would like to acknowledge the Neurodegenerative Undergraduate Research Experience (NURE)

R25NS100118 for funding DXD. The authors express their deep appreciation to Dr. Rehana

Leak (DU) for her brilliant insights and technical advice. The authors are thankful to Dr. Lauren

O'Donnell, and Mss. Manisha Chandwani and Yashika Kamte (DU) for flow cytometry support.

1325

References

1.   Murphy DE, de Jong OG, Brouwer M, Wood MJ, Lavieu G, Schiffelers RM, et al. Extracellular vesicle-based therapeutics: natural versus engineered targeting and trafficking. Experimental & Molecular Medicine. 2019;51(3):1-12.

2.   Doyle LM, Wang MZ. Overview of Extracellular Vesicles, Their Origin, Composition, Purpose, and Methods for Exosome Isolation and Analysis. Cells. 2019;8(7):727.

3.   He Z, Ning N, Zhou Q, Khoshnam SE, Farzaneh M. Mitochondria as a therapeutic target for ischemic stroke. Free Radical Biology and Medicine. 2020;146:45-58.

4.   Abdullahi W, Tripathi D, Ronaldson PT. Blood-brain barrier dysfunction in ischemic stroke: targeting tight junctions and transporters for vascular protection. Am J Physiol Cell Physiol. 2018;315(3):C343-C56.

5.   Bernardo-Castro S, Sousa JA, Brás A, Cecília C, Rodrigues B, Almendra L, et al. Pathophysiology of Blood–Brain Barrier Permeability Throughout the Different Stages of Ischemic Stroke and Its Implication on Hemorrhagic Transformation and Recovery. Frontiers in Neurology. 2020;11(1605).

6.   Jiang X, Andjelkovic AV, Zhu L, Yang T, Bennett MVL, Chen J, et al. Blood-brain barrier dysfunction and recovery after ischemic stroke. Prog Neurobiol. 2018;163-164:144-71.

7.   Nian K, Harding IC, Herman IM, Ebong EE. Blood-Brain Barrier Damage in Ischemic Stroke and Its Regulation by Endothelial Mechanotransduction. Front Physiol. 2020;11(1681).

8.   Stamatovic SM, Keep RF, Wang MM, Jankovic I, Andjelkovic AV. Caveolae-mediated internalization of occludin and claudin-5 during CCL2-induced tight junction remodeling in brain endothelial cells. J Biol Chem. 2009;284(28):19053-66.

9.   Shi Y, Jiang X, Zhang L, Pu H, Hu X, Zhang W, et al. Endothelium-targeted overexpression of heat shock protein 27 ameliorates blood-brain barrier disruption after ischemic brain injury. Proc Natl Acad Sci U S A. 2017;114(7):E1243-E52.

10.   Leak RK, Zhang L, Stetler RA, Weng Z, Li P, Atkins GB, et al. HSP27 protects the blood-brain barrier against ischemia-induced loss of integrity. CNS Neurol Disord Drug Targets. 2013;12(3):325-37.

11.   During RL, Gibson BG, Li W, Bishai EA, Sidhu GS, Landry J, et al. Anthrax lethal toxin paralyzes actin-based motility by blocking Hsp27 phosphorylation. EMBO J. 2007;26(9):2240-50.

12.   Rada CC, Mejia-Pena H, Grimsey NJ, Cordova IC, Olson J, Wozniak JM, et al. Heat shock protein 27 activity is linked to endothelial barrier recovery after proinflammatory GPCR-induced disruption. Sci Signal. 2021;14(698):eabc1044.

13.   Hirano S, Rees RS, Yancy SL, Welsh MJ, Remick DG, Yamada T, et al. Endothelial barrier dysfunction caused by LPS correlates with phosphorylation of HSP27 in vivo. Cell Biol Toxicol. 2004;20(1):1-14.

14.   Miron T, Vancompernolle K, Vandekerckhove J, Wilchek M, Geiger B. A 25-kD inhibitor of actin polymerization is a low molecular mass heat shock protein. J Cell Biol. 1991;114(2):255-61.

15.   D'Souza A, Burch A, Dave KM, Sreeram A, Reynolds MJ, Dobbins DX, et al. Microvesicles transfer mitochondria and increase mitochondrial function in brain endothelial cells. J Control Release. 2021;338:505-26.

16.   Banks WA, Sharma P, Bullock KM, Hansen KM, Ludwig N, Whiteside TL. Transport of
Extracellular Vesicles across the Blood-Brain Barrier: Brain Pharmacokinetics and Effects of
Inflammation. International journal of molecular sciences. 2020;21(12):4407.
17.   Jiang Y, Arounleut P, Rheiner S, Bae Y, Kabanov AV, Milligan C, et al. SOD1
nanozyme with reduced toxicity and MPS accumulation. J Control Release. 2016;231:38-49.
18.   Han M, Bae Y, Nishiyama N, Miyata K, Oba M, Kataoka K. Transfection study using
multicellular tumor spheroids for screening non-viral polymeric gene vectors with low
cytotoxicity and high transfection efficiencies. Journal of Controlled Release. 2007;121(1):38-
48.
19.   Xiong MP, Bae Y, Fukushima S, Forrest ML, Nishiyama N, Kataoka K, et al. pH-
Responsive Multi-PEGylated Dual Cationic Nanoparticles Enable Charge Modulations for Safe
Gene Delivery. ChemMedChem. 2007;2(9):1321-7.
20.   Lai TC, Bae Y, Yoshida T, Kataoka K, Kwon GS. pH-Sensitive Multi-PEGylated Block
Copolymer as a Bioresponsive pDNA Delivery Vector. Pharmaceutical Research.
2010;27(11):2260-73.
21.   Dave KM, Zhao W, Hoover C, D'Souza A, S Manickam D. Extracellular Vesicles
Derived from a Human Brain Endothelial Cell Line Increase Cellular ATP Levels. AAPS
PharmSciTech. 2021;22(1):18.
22.   Chen TH, Bae Y, Furgeson DY, Kwon GS. Biodegradable hybrid recombinant block
copolymers for non-viral gene transfection. Int J Pharm. 2012;427(1):105-12.
23.   Kanayama N, Fukushima S, Nishiyama N, Itaka K, Jang W-D, Miyata K, et al. A PEG-
Based Biocompatible Block Catiomer with High Buffering Capacity for the Construction of
Polyplex Micelles Showing Efficient Gene Transfer toward Primary Cells. ChemMedChem.
2006;1(4):439-44.
24.   Donker RB, Mouillet JF, Chu T, Hubel CA, Stolz DB, Morelli AE, et al. The expression
profile of C19MC microRNAs in primary human trophoblast cells and exosomes. Mol Hum
Reprod. 2012;18(8):417-24.
25.   Hough KP, Trevor JL, Strenkowski JG, Wang Y, Chacko BK, Tousif S, et al. Exosomal
transfer of mitochondria from airway myeloid-derived regulatory cells to T cells. Redox Biol.
2018;18(2213-2317):54-64.
26.   Dave KM, Ali L, Manickam DS. Characterization of the SIM-A9 cell line as a model of
activated microglia in the context of neuropathic pain. PLOS ONE. 2020;15(4):e0231597.
27.   Dave KM, Han L, Jackson MA, Kadlecik L, Duvall CL, D SM. DNA Polyplexes of a
Phosphorylcholine-Based Zwitterionic Polymer for Gene Delivery. Pharm Res. 2020;37(9):176.
28.   Dai X, Chen J, Xu F, Zhao J, Cai W, Sun Z, et al. TGFalpha preserves oligodendrocyte
lineage cells and improves white matter integrity after cerebral ischemia. J Cereb Blood Flow
Metab. 2020;40(3):639-55.
29.   Cardenes N, Corey C, Geary L, Jain S, Zharikov S, Barge S, et al. Platelet bioenergetic
screen in sickle cell patients reveals mitochondrial complex V inhibition, which contributes to
platelet activation. Blood. 2014;123(18):2864-72.
30.   Ikeda G, Santoso MR, Tada Y, Li AM, Vaskova E, Jung J-H, et al. Mitochondria-Rich
Extracellular Vesicles From Autologous Stem Cell–Derived Cardiomyocytes Restore Energetics
of Ischemic Myocardium. Journal of the American College of Cardiology. 2021;77(8):1073-88.
31.   Phinney DG, Di Giuseppe M, Njah J, Sala E, Shiva S, St Croix CM, et al. Mesenchymal
stem cells use extracellular vesicles to outsource mitophagy and shuttle microRNAs. Nature
Communications. 2015;6(1):8472.

32. Hayakawa K, Esposito E, Wang X, Terasaki Y, Liu Y, Xing C, et al. Transfer of mitochondria from astrocytes to neurons after stroke. Nature. 2016;535:551.

33. Hayakawa K, Chan SJ, Mandeville ET, Park JH, Bruzzese M, Montaner J, et al. Protective Effects of Endothelial Progenitor Cell-Derived Extracellular Mitochondria in Brain Endothelium. Stem Cells. 2018;36(9):1404-10.

34. CellLight™ Mitochondria-GFP, BacMam 2.0 (ThermoFisher website) [

35. https://www.agilent.com/en/products/cell-analysis/how-seahorse-xf-analyzers-work. [

36. http://protcalc.sourceforge.net/. [

37. Doll DN, Hu H, Sun J, Lewis SE, Simpkins JW, Ren X. Mitochondrial crisis in cerebrovascular endothelial cells opens the blood-brain barrier. Stroke. 2015;46(6):1681-9.

38. Shi Y, Jiang X, Zhang L, Pu H, Hu X, Zhang W, et al. Endothelium-targeted overexpression of heat shock protein 27 ameliorates blood–brain barrier disruption after ischemic brain injury. Proceedings of the National Academy of Sciences. 2017;114(7):E1243-E52.

39. Shi Y, Zhang L, Pu H, Mao L, Hu X, Jiang X, et al. Rapid endothelial cytoskeletal reorganization enables early blood–brain barrier disruption and long-term ischaemic reperfusion brain injury. Nature Communications. 2016;7:10523.

40. Sweeney MD, Zhao Z, Montagne A, Nelson AR, Zlokovic BV. Blood-Brain Barrier: From Physiology to Disease and Back. Physiological Reviews. 2019;99(1):21-78.

41. del Zoppo GJ. Inflammation and the neurovascular unit in the setting of focal cerebral ischemia. Neuroscience. 2009;158(3):972-82.

42. Knowland D, Arac A, Sekiguchi KJ, Hsu M, Lutz SE, Perrino J, et al. Stepwise recruitment of transcellular and paracellular pathways underlies blood-brain barrier breakdown in stroke. Neuron. 2014;82(3):603-17.

43. Oldendorf WH, Cornford ME, Brown WJ. The large apparent work capability of the blood-brain barrier: a study of the mitochondrial content of capillary endothelial cells in brain and other tissues of the rat. Ann Neurol. 1977;1(5):409-17.

44. Schiera G, Di Liegro CM, Di Liegro I. Extracellular Membrane Vesicles as Vehicles for Brain Cell-to-Cell Interactions in Physiological as well as Pathological Conditions. BioMed Research International. 2015;2015:152926.

45. Andras IE, Toborek M. Extracellular vesicles of the blood-brain barrier. Tissue Barriers. 2016;4(1):e1131804.

46. Yang T, Martin P, Fogarty B, Brown A, Schurman K, Phipps R, et al. Exosome delivered anticancer drugs across the blood-brain barrier for brain cancer therapy in Danio rerio. Pharm Res. 2015;32(6):2003-14.

47. Yang D, Zhang W, Zhang H, Zhang F, Chen L, Ma L, et al. Progress, opportunity, and perspective on exosome isolation - efforts for efficient exosome-based theranostics. Theranostics. 2020;10(8):3684-707.

48. Dozio V, Sanchez JC. Characterisation of extracellular vesicle-subsets derived from brain endothelial cells and analysis of their protein cargo modulation after TNF exposure. J Extracell Vesicles. 2017;6(1):1302705.

49. Kanada M, Bachmann MH, Hardy JW, Frimannson DO, Bronsart L, Wang A, et al. Differential fates of biomolecules delivered to target cells via extracellular vesicles. Proceedings of the National Academy of Sciences. 2015;112(12):E1433-E42.

50. Konoshenko MY, Lekchnov EA, Vlassov AV, Laktionov PP. Isolation of Extracellular Vesicles: General Methodologies and Latest Trends. Biomed Res Int. 2018;2018:8545347.

51.   El Andaloussi S, Mäger I, Breakefield XO, Wood MJA. Extracellular vesicles: biology and emerging therapeutic opportunities. Nature Reviews Drug Discovery. 2013;12(5):347-57.

52.   Jeyaram A, Jay SM. Preservation and Storage Stability of Extracellular Vesicles for Therapeutic Applications. AAPS J. 2017;20(1):1.

53.   Ge Q, Zhou Y, Lu J, Bai Y, Xie X, Lu Z. miRNA in Plasma Exosome is Stable under Different Storage Conditions. Molecules. 2014;19(2):1568-75.

54.   Witwer KW, Buzás EI, Bemis LT, Bora A, Lässer C, Lötvall J, et al. Standardization of sample collection, isolation and analysis methods in extracellular vesicle research. Journal of Extracellular Vesicles. 2013;2(1):20360.

55.   Spees JL, Olson SD, Whitney MJ, Prockop DJ. Mitochondrial transfer between cells can rescue aerobic respiration. Proceedings of the National Academy of Sciences. 2006;103(5):1283-8.

56.   Islam MN, Das SR, Emin MT, Wei M, Sun L, Westphalen K, et al. Mitochondrial transfer from bone-marrow–derived stromal cells to pulmonary alveoli protects against acute lung injury. Nature Medicine. 2012;18(5):759-65.

57.   Hayakawa K, Esposito E, Wang X, Terasaki Y, Liu Y, Xing C, et al. Transfer of mitochondria from astrocytes to neurons after stroke. Nature. 2016;535(7613):551-5.

58.   Sinha P, Islam MN, Bhattacharya S, Bhattacharya J. Intercellular mitochondrial transfer: bioenergetic crosstalk between cells. Current Opinion in Genetics & Development. 2016;38:97-101.

59.   Gao L, Zhang Z, Lu J, Pei G. Mitochondria Are Dynamically Transferring Between Human Neural Cells and Alexander Disease-Associated GFAP Mutations Impair the Astrocytic Transfer. Front Cell Neurosci. 2019;13(316):316.

60.   Puhm F, Afonyushkin T, Resch U, Obermayer G, Rohde M, Penz T, et al. Mitochondria Are a Subset of Extracellular Vesicles Released by Activated Monocytes and Induce Type I IFN and TNF Responses in Endothelial Cells. Circulation Research. 2019;125(1):43-52.

61.   van der Vlist M, Raoof R, Willemen H, Prado J, Versteeg S, Martin Gil C, et al. Macrophages transfer mitochondria to sensory neurons to resolve inflammatory pain. Neuron. 2022;110(4):613-26.e9.

62.   Guescini M, Genedani S, Stocchi V, Agnati LF. Astrocytes and Glioblastoma cells release exosomes carrying mtDNA. Journal of Neural Transmission. 2009;117(1):1.

63.   Goldberg J, Currais A, Prior M, Fischer W, Chiruta C, Ratliff E, et al. The mitochondrial ATP synthase is a shared drug target for aging and dementia. Aging Cell. 2018;17(2):e12715.

64.   Jonckheere AI, Smeitink JAM, Rodenburg RJT. Mitochondrial ATP synthase: architecture, function and pathology. Journal of Inherited Metabolic Disease. 2012;35(2):211-25.

65.   Pasquale D'Acunzo oP-G, Yohan Kim, Tal Hargash, Chelsea Miller, Melissa J. Alldred, Hediye Erdjument-Bromage, Sai C. Penikalapati, Monika Pawlik, Mitsuo Saito, Mariko Saito, Stephen D. Ginsberg, Thomas A. Neubert, Chris N. Goulbourne, Efrat Levy. Mitovesicles are a novel population of extracellular vesicles of mitochondrial origin altered in Down syndrome. Scientific Reports. 2021;7:eabe5085.

66.   Sansone P, Savini C, Kurelac I, Chang Q, Amato LB, Strillacci A, et al. Packaging and transfer of mitochondrial DNA via exosomes regulate escape from dormancy in hormonal therapy-resistant breast cancer. Proceedings of the National Academy of Sciences. 2017;114(43):E9066-E75.

67.   Yamamoto H, Itoh N, Kawano S, Yatsukawa Y-i, Momose T, Makio T, et al. Dual role of the receptor Tom20 in specificity and efficiency of protein import into mitochondria.

Proceedings of the National Academy of Sciences of the United States of America. 2011;108(1):91-6.

68.   Dutra Silva J, Su Y, Calfee CS, Delucchi KL, Weiss D, McAuley DF, et al. Mesenchymal stromal cell extracellular vesicles rescue mitochondrial dysfunction and improve barrier integrity in clinically relevant models of ARDS. Eur Respir J. 2021;58(1).

69.   Andreu Z, Yáñez-Mó M. Tetraspanins in extracellular vesicle formation and function. Front Immunol. 2014;5:442.

70.   Xiao B, Deng X, Zhou W, Tan EK. Flow Cytometry-Based Assessment of Mitophagy Using MitoTracker. Front Cell Neurosci. 2016;10(76):76.

71.   Chazotte B. Labeling mitochondria with MitoTracker dyes. Cold Spring Harb Protoc. 2011;2011(8):990-2.

72.   Hough KP, Trevor JL, Strenkowski JG, Wang Y, Chacko BK, Tousif S, et al. Exosomal transfer of mitochondria from airway myeloid-derived regulatory cells to T cells. Redox Biol. 2018;18:54-64.

73.   Guo Y, Chi X, Wang Y, Heng BC, Wei Y, Zhang X, et al. Mitochondria transfer enhances proliferation, migration, and osteogenic differentiation of bone marrow mesenchymal stem cell and promotes bone defect healing. Stem Cell Res Ther. 2020;11(1):245.

74.   Zhang J, Tan Z, Tran ND. Chemical hypoxia-ischemia induces apoptosis in cerebromicrovascular endothelial cells. Brain Res. 2000;877(2):134-40.

75.   Andjelkovic AV, Xiang J, Stamatovic SM, Hua Y, Xi G, Wang MM, et al. Endothelial Targets in Stroke. Arteriosclerosis, Thrombosis, and Vascular Biology. 2019;39(11):2240-7.

76.   Yang DI, Lin TN, Yin KJ, Hu CJ, Chen SD, Xu J, et al. Cerebral Endothelial Cell Reaction to Ischemic Insults. In: Lajtha A, Chan PH, editors. Handbook of Neurochemistry and Molecular Neurobiology: Acute Ischemic Injury and Repair in the Nervous System. Boston, MA: Springer US; 2007. p. 165-83.

77.   Keep R.F. AAV, Stamatovic S.M., Shakui P., Ennis S.R. Ischemia-induced endothelial cell dysfunction. Intracranial Pressure and Brain Monitoring XII. 2005;95.

78.   Park H-H, Han M-H, Choi H, Lee YJ, Kim JM, Cheong JH, et al. Mitochondria damaged by Oxygen Glucose Deprivation can be Restored through Activation of the PI3K/Akt Pathway and Inhibition of Calcium Influx by Amlodipine Camsylate. Scientific Reports. 2019;9(1):15717.

79.   Liao L-X, Zhao M-B, Dong X, Jiang Y, Zeng K-W, Tu P-F. TDB protects vascular endothelial cells against oxygen-glucose deprivation/reperfusion-induced injury by targeting miR-34a to increase Bcl-2 expression. Scientific Reports. 2016;6(1):37959.

80.   Smolina N, Bruton J, Kostareva A, Sejersen T. Assaying Mitochondrial Respiration as an Indicator of Cellular Metabolism and Fitness. Methods Mol Biol. 2017;1601:79-87.

81.   Rose S, Frye RE, Slattery J, Wynne R, Tippett M, Pavliv O, et al. Oxidative stress induces mitochondrial dysfunction in a subset of autism lymphoblastoid cell lines in a well-matched case control cohort. PLoS One. 2014;9(1):e85436.

82.   Miyata K, Oba M, Nakanishi M, Fukushima S, Yamasaki Y, Koyama H, et al. Polyplexes from Poly(aspartamide) Bearing 1,2-Diaminoethane Side Chains Induce pH-Selective, Endosomal Membrane Destabilization with Amplified Transfection and Negligible Cytotoxicity. Journal of the American Chemical Society. 2008;130(48):16287-94.

83.   Haney MJ, Klyachko NL, Zhao Y, Gupta R, Plotnikova EG, He Z, et al. Exosomes as drug delivery vehicles for Parkinson's disease therapy. J Control Release. 2015;207:18-30.

1551   84.   Yuan D, Zhao Y, Banks WA, Bullock KM, Haney M, Batrakova E, et al. Macrophage
1552   exosomes as natural nanocarriers for protein delivery to inflamed brain. Biomaterials.
1553   2017;142:1-12.

1554