1 Impact of Protein Identity on Tumor-Associated Antigen Uptake

2 into Infiltrating Immune Cells: A Comparison of Different

3 Fluorescent Proteins as Model Antigens

4 Short title: Tumor-Associated Antigen Uptake of Different

5 Fluorescent Proteins

6 Rulan Yi[1], Emily Chen[2], Edward W Roberts[3], Matthew F Krummel[2*], Nina K Serwas[2*]

7 [1]MCB Undergraduate Program, UC Berkeley, [2]Department of Pathology, UCSF, [3]Institute of Cancer Sciences University of Glasgow

8 *correspondence: matthew.krummel@ucsf.edu, nina.serwas@ucsf.edu

9 Abstract

Effective immune responses depend on efficient antigen uptake in the periphery,

transport of those antigens to, and presentation in draining lymph nodes (LNs). These processes

have been studied intensively using stable fluorescent proteins (FPs) as model antigens. To date,

ZsGreen is the only FP that can be tracked efficiently towards LNs, hence, it is difficult to

compare studies using alternated tracking proteins.

Here, we systematically compared six different FPs. We included ZsGreen, ZsYellow,

DsRed, AsRed, mCherry, and mRFP based on sequence homology and/or origin species, and

generated FP-expressing tumor cell lines. Stability of fluorescent signal was assessed *in vitro*

over time, across different pH environments, and *in vivo* through FP antigen uptake and transfer

to immune cells isolated from tumors and tumor-draining LNs.

ZsGreen could be detected in high percentages of all analyzed tumor-infiltrating immune

cells, with highest amounts in tumor-associated macrophages (TAMs) and type 2 conventional

dendritic cells (cDC2s). ZsYellow, AsRed, and DsRed followed a similar pattern, but

percentages of FP-containing immune cells in the tumor were lower than for ZsGreen.

Strikingly, mRFP and mCherry demonstrated a 'non-canonical' antigen uptake pattern where

percentages of FP-positive tumor-infiltrating immune cells were highest for cDC1s not TAMs

and cDC2s despite comparable stabilities and localization of all FPs. Analysis of antigen-

containing cells in the LN was hindered by intracellular degradation of FPs. Only ZsGreen could

be efficiently tracked to the LN, though some signal was measurable for ZsYellow and DsRed.

In summary, we find that detection of antigen uptake and distribution is subject to

variabilities related to fluorophore nature. Future experiments need to consider that these

processes might be impacted by protein expression, stability, or other unknown factors. Thus,

our data sheds light on potential under-appreciated mechanisms regulating antigen transfer and

highlights potential uses and necessary caveats to interpretation based on FP use.

Introduction

For a successful tumor immune response, the proper uptake and presentation of tumor-associated antigens (TAAs) by antigen presenting cells (APCs), such as dendritic cells (DCs), is essential. As such, antigens from peripheral tissues need to be taken up by phagocytic APCs which mature, upregulate costimulatory molecules, and present these antigens as processed peptides on MHC complexes. During this process, APCs traffic from the peripheral tissue where the tumor is located to the tumor-draining lymph node (tdLN). There, the APCs can interact with T and B cells to generate an effective long-term response (1,2). The interaction of the TAA-peptide:MHC complex and its accessory proteins with receptors on T and B cells determines downstream immune responses such as cell activation and differentiation. Exactly how TAAs are taken up and brought to the LN for presentation is not fully understood.

Previous studies showed that type 1 DCs (cDC1s) are essential for the transport of TAAs to the lymph node, and that this transport is dependent on C-C chemokine receptor 7 (Ccr7) (3,4). Upon entry into the LN, migratory DCs can transfer antigens to other lymph node resident DCs as membrane encapsulated vesicles through direct cell-cell contact (5). The choice of the receiving DCs may then dictate the downstream immune responses generated through subsequent T cell interaction causing activation, proliferation, memory induction, and/or effector responses in the T cell (5). Previous studies that analyzed the process of antigen uptake, presentation, and transfer have used fluorescent proteins (FPs) such as mCherry, green fluorescent protein (GFP), yellow fluorescent protein (YFP), tdTomato, and ZsGreen as model antigens (3–5). FPs used in these studies seemed to have different stabilities since only ZsGreen was substantially detected in resident dendritic cells. Importantly, other tumor antigens can be tracked in those resident DCs as well, confirming that ZsGreen presence was not an artifact, but

faithfully reported on transport (5). It is not clear why other FPs cannot be transported to the

same extend to the tdLNs and whether certain properties of FPs alter antigen uptake and

distribution or simply degrade more rapidly than others *en route* to the LN.

A wide range of FPs exists that are frequently used in imaging and tracking experiments.

These FPs have varying brightness and stabilities that may potentially impact the outcome of

antigen uptake, transfer, and presentation. Brightness is defined as extinction coefficient times

quantum yield at pH 7.4 (6) and can range from dim proteins, such as AsRed derived from

Anemonia sulcate, with a brightness index of 2.81 (7), to bright proteins with indices of 90

(tdTomato) (6). Stability is dependent on the specific environment in which the FP resides at the

time of analysis (6). In this study we tested antigen stability *in vitro* and *in vivo* for six different

FPs with a range of brightness indexes between 2.81 and 49. We tracked their stabilities during

uptake and intracellular storage in tumor-infiltrating immune cells and after transport to the

tdLNs. The combined results offer a comprehensive overview of the properties of the chosen

FPs with regards for their usefulness in assays tracking antigen uptake and transfer. This work

will facilitate the choice of FPs in studies that will use two or more fluorophores in parallel.

Material and Methods

Alignment

Amino acid sequences of all fluorescent proteins used were gathered via FPbase (7), then

aligned via Basic Local Alignment Search Tool (BLAST) (8). Percent similarity to ZsGreen and

DsRed were determined using the global align tool.

Mice

Mice were housed and bred under specific pathogen-free conditions at the University of

California, San Francisco Laboratory Animal Research Center and all experiments conformed to

ethical principles and guidelines approved by the UCSF Institutional Animal Care and Use

Committee. C57BL/6 mice were purchased from Jackson Laboratory or bred in-house. Male

mice between 8-12 weeks of age were used.

Cell lines

Vectors containing target fluorescent genes ZsGreen, ZsYellow, DsRed, AsRed,

mCherry, and mRFP (Takara Bio), were either transfected into B16-F10 cells using the calcium

chloride precipitation method (all FPs except ZsGreen), or retrovirally transduced into B16F10

cells (ZsGreen). Transformed cells were then harvested, washed, and repeatedly sorted through

fluorescent activated cell sorting (FACS) on a BD FACSAria^{TM} to select for stable fluorescent

positive populations. Sorted cells were then collected, frozen and/or propagated for use in future

experiments. Cells were cultured at 37°C with 5% CO_2 in DMEM (GIBCO) plus 10% heat-

inactivated FCS with penicillin, streptomycin, and L-glutamate on tissue-culture treated plastic

plates and split every other day.

Fluorescence stability assay

FP-expressing B16-F10 cells were harvested with trypsin, counted, and washed in ice

cold PBS. Then, cells were homogenized with mechanical force generated through repeated

passage of cells through a 22-gauge needle in hypotonic lysis buffer (20mM HEPES buffer,

2mM EGTA, 2mM MgCl2). Lysates were transferred to 96-well black microplates suitable for

fluorescent based assays (Invitrogen). Each well contained lysates of 1 Mio cells. pH was

adjusted with NaOH or HCl, controlled with a pH-meter before the assay, and controlled with

pH-indicator strips on the lysates after the experiment to ensure that the pH was stable overtime.

For each different fluorophore and pH condition, three technical replicates were performed.

Data was collected over varying time points using a microplate reader (SpectraMax) and

incubated at 37°C or 4C, respectively. Analysis of whole cells was done similarly in separate

plates. The experiment was done in biological triplicates.

Ectopic tumor injections

B16-F10 cells expressing the respective FP were grown to 80% confluency, then

harvested, washed, and counted. Cells were resuspended in a suitable amount of PBS and mixed

at a 1:1 (v:v) ratio with growth factor reduced Matrigel Matrix (BD Biosciences). A final

injection volume of 100ul containing 400k cells was injected subcutaneously on the flanks of

mice.

Harvest, collection of data, and analysis

Tumors, inguinal and axillary lymph nodes were harvested 2 weeks after injection. After

digestion with DNAse, Collagenase I and Collagenase IV, washing, and counting, 10 million

cells per FP were stained with fixable Live/Dead Zombie (BioLegend) and respective antibodies:

MHCII-BV421, F4/80-BV510, CD11b-BV605, CD11c-BV650, Ly6C-BV711, CD45R-BV785,

CD90.2-BV785, Ly6G-BV785, NK1.1-BV785, CD8alpha-PerCPCy5.5, CD103-APC, CD45-

AF700, CD24-PECy7 (Biolegend, eBioscience).

Flow cytometry was performed on a BD LSR Fortessa instrument at the ImmunoX Flow

Cytometry CoLab. Analysis of flow cytometry data was done using FlowJo (Treestar) software,

and graphs and figures were produced with GraphPad Prism software. All experiments were

done in triplicates.

Fluorescence microscopy

Cells from each respective fluorescent cell line were thawed and cultured. 50,000 cells

were seeded in a poly-L-lysine coated 8-well chambered coverslip and let rest overnight. For

live cell imaging, a Leica SP5 laser scanning confocal microscope with an incubation chamber

and a 20X oil objective was used to visualize adherent cells expressing respectively ZsGreen,

ZsYellow, AsRed, DsRed, mRFP, and mCherry. The experiment was done in triplicates.

Figures

Table 1. Fluorophore properties (7,8)

Fluorophore	Brightness	Excitation	Emission	% similarity to ZsGreen	% similarity to DsRed
AsRed	2.81	576 nm	592 nm	53.78%	61.54%
mRFP	12.5	584 nm	607 nm	56.65%	87.56%
ZsYellow	13	528 nm	539 nm	91.77%	58.87%
mCherry	15.84	587 nm	610 nm	53.53%	82.63%
ZsGreen	22.43	496 nm	506 nm	—	59.23%
DsRed	49.3	558 nm	583 nm	59.23%	—

Fig 1. Generation of cell lines with respective fluorescent genes for downstream analysis.

A) Lentiviral vectors expressing respective fluorescent genes ZsGreen, ZsYellow, mCherry,

mRFP, DsRed, and AsRed were transfected/transduced into B16-10 cells. **B)** Fluorescence-

activated cell sorting (FACS) was then used to select for FP positivity. ZsYellow and mCherry

cells display a bimodal pattern despite undergoing FACS multiple times with a selection

criterion for the top 5% of FP positive cells.

A) 37°C

B) Initial timepoint   137

Legend: Hypotonic lysis buffer | pH=7 | pH=6.5 | pH=6 | pH=5.5 | pH=5

Fig 2. *In vitro* stability assay shows comparable stabilities of selected fluorophores. Lysates

of 1 million respective FP-expressing B16-F10 cells were generated in hypotonic lysis buffer and

incubated at **A)** 37ºC in pH 7.4 or **B)** 37ºC in pH ranging from 5-7.4. Fluorescence intensity data

was collected over time at the indicated time points (**A**) or immediately following pH treatment

(**B**). Data was normalized to the maximum fluorescence intensity of each FP for every

independent experiment.

Fig 3. Growth and harvest of respective fluorescent cell lines A) Fluorescent cell lines were

grown, harvested, and injected subcutaneously on the flanks of B6 mice. After 14 days, tumors

and draining lymph nodes were harvested and processed for flow cytometry. **B)** Average sizes

of tumors (mm^3) at harvest are shown for all six respective fluorophores. Size was controlled to

ensure comparable *in vivo* growth of different FP-expressing B16-F10 cell lines.

Fig 4. TAA uptake by tumor-infiltrating immune cells identifies different patterns for different fluorophores. Accumulation of respective fluorescent proteins in immune cells, purified from the tumor microenvironment. A non-fluorescent control (B16-F10) was used to set the threshold for FP positivity. One representative example of three independent experiments is shown.

Fig 5. cDC2s and TAMs contain the most fluorescent TAAs per individual cell. Immune

cells were purified from the TME of B16 mice bearing B16 tumors with the indicated

fluorophors and analyzed for relative antigen uptake. Data was normalized to the maximum delta

MFI of each FP for every independent experiment.

Fig 6. Limited detection of other non-ZsGreen fluorescent proteins within immune cells of tumor draining lymph nodes. Tumor draining (inguinal and axillary) lymph nodes were harvested, processed, and analyzed through flow cytometry. Non-fluorescent control (B16-F10) was used to set the threshold for FP positivity. One representative example of three independent experiments is shown.

Results and Discussion

Characteristics of selected fluorescent protein

We compared five different fluorescent proteins (FPs) for antigen uptake and transfer

characteristics to ZsGreen. ZsGreen is to date the only fluorophore that can be efficiently

tracked from expressing tumors to lymph node resident dendritic cells. Our selection (Table 1)

was based on species origin (ZsYellow), brightness of the fluorophore (DsRed), sequence

homology (mCherry, mRFP) and availability of corresponding biochemical tools (AsRed,

mRFP, mCherry). ZsGreen, is derived from corals of the *Zoanthus sp*, and has been shown to

have high stability in various chemical environments (9). Its brightness index is 22.43 (7).

ZsYellow is a stable protein purified from the same species as ZsGreen, and is excited at a

wavelength around 528 nm with a brightness index of 13 (7,9). DsRed is derived from

Discosoma sp (10)*, excited by a wavelength of 558 nm, and has a brightness index of 49.3 (7).

Naturally occurring FPs such as DsRed have been codon-modified to achieve proteins with

optimized brightness and stability properties such as mRFP and mCherry, with brightness indices

of 12.5 and 15.84 respectively (7). In addition, these modified proteins exist as monomers in

contrast to the tetrameric nature of ZsGreen and DsRed (7). AsRed has the lowest brightness in

our collection (2.81) and is expressed in the coral species *Anemonia sulcate* (6,7,10).

FP-encoding vectors were transduced into B16-F10 cell lines. Transfected cells with the

highest signal for the respective FP were repeatedly sorted with a 5 %- cut-off using fluorescence

activated cell sorting (FACS) (Fig 1A and B). All generated cell lines reached high FP positivity

with a 2-3 log fluorescent intensity difference over un-transfected cells (Fig 1A and B). Despite

repeated and stringent selection for FP-positivity, both ZsYellow- and mCherry-transfected B16-

F10 cells kept a substantial amount of non-fluorescent cells in their cultures (Fig 1B). This might

be due to non-stable integration or epigenetic silencing of the transgene. In subsequent analyses

we considered this bimodal distribution as a potential factor for differences in antigen uptake or

distribution. Previous work has shown that differential subcellular localization has an effect on

antigen uptake and distribution into compartments of the immune system (11). All generated cell

lines had their respective transfected FPs evenly distributed in the cytoplasm (Fig S1).

Assessment of *in vitro* stability

The stability of fluorophores expressed in the generated cell lines was assessed in an *in*

vitro fluorescent stability assay. For this purpose, FP-expressing B16-F10 cell lines were lysed

in a hypotonic lysis buffer. Lysates were then kept at 37ºC to mimic the cellular environment.

No protease inhibitors were added. This allowed us to follow protein stability in the cytosolic

soup containing various proteases and other enzymes, through fluorescence intensity, over time.

As this assay investigates the stability of FPs, and not their individual brightness, we internally

normalized the brightness to timepoint 0 in neutral pH conditions. In general, ZsGreen

demonstrated the highest stability of all analyzed FPs, and showed only a minimal decrease in

fluorescent signal over the time frame of 6 hours (Fig 2A). This high stability of ZsGreen is in

concordance to previous published work (9). In contrast, all other FPs showed a severe

reduction of signal over the first two hours (Fig 2A). Acidification of the protein lysates showed

relative robustness of all FPs to pH alterations (Fig 2B). The highest fluorescent intensity of all

FPs could be detected at neutral pH conditions which is in line with previous published analyses

of pH stability for fluorescent proteins (12). Our data suggests that the impact of acidification in

certain intracellular compartments only plays a minimal role on FP fluorescence intensity and

stability. However, proteins might be degraded over time through intracellular proteases or other

enzymes, which was especially true for ZsYellow, AsRed, DsRed, mRFP, and mCherry.

ZsGreen markedly showed resistance towards degradation by proteins contained in the cellular

lysates which might explain its superior performance in tumor-associated antigen tracking

experiments. Differences between FPs other than ZsGreen in those assays seem to not be caused

by different fluorescence stabilities.

Tumor-associated antigen uptake varies for different FPs

To evaluate tumor-associated antigen (TAA) uptake of the six selected FPs, cells were

injected bilaterally into the flanks of B6 mice (Fig 3A). Tumors averaged at a size of 1300 mm^3

for all FP-containing cell lines two weeks after inocculation (Fig 3B). None of the generated FP-

expressing B16-F10 cell lines resulted in severe increase or reduction of tumor growth,

suggesting that the modification did not alter proliferation or fitness of the cells. For antigen

tracking analysis, only tumors of a size equal to or greater than 800 mm^3 were included.

We assessed distribution of FPs into different cellular populations of the tumor, namely

monocytes, neutrophils, tumor associated macrophages (TAMs), and DCs (Fig 4, Fig S2). Of

note, infiltrating immune cell distributions were similar for all injected tumors (Fig S3). Among

the selected FPs, percentages of FP-containing immune cells were highest in tumors expressing

ZsGreen. Interestingly, we identified two distinct antigen uptake pattern: The 'classical' pattern

showed highest percentages of FP-containing cells for cDC2s, TAM1s, and TAM2s. ZsGreen-

expressing tumor cells resulted in nearly 100% of these infiltrating immune cells to contain the

FP. ZsYellow-, DsRed- and AsRed-expressing tumor cells resulted in less FP-positive immune

cells, however, cDC2s, and TAMs were still the prime samplers of tumor associated FPs. For

this 'classical' uptake pattern, a smaller fraction of neutrophils, monocytes, and cDC1s displayed

FP signal than seen for cDC2s and TAMs. The 'non-canonical' antigen uptake pattern was

characterized by highest proportions of FP-positive cells for cDC1s and not cDC2s or TAMs.

Also for this pattern, the lowest amount of transferred antigen could be seen in neutrophils and

monocytes. Both, mRFP- and mCherry-expressing tumors resulted in such a distinct pattern of

FP uptake by immune cells (Fig 4). Despite comparable brightness, mCherry resulted in overall

higher percentages of FP-positive tumor-infiltrating immune cells than mRFP (Fig 4).

Remarkably, this was true although mCherry-expressing cells were a mixture of only two thirds

transformed and one third WT cells (Fig 1B).

Next, we analyzed the average amount of FPs within each of the individual populations.

We used normalized delta mean fluorescence intensity to visualize how much sampled FP each

cell population contained in comparison to other immune cells sampling from the same

fluorophore. Overall patterns of relative amount of uptake per cell (Fig 5) showed more

similarity than the percentages of FP- positive cells (Fig 4). For most of the tumors, cDC2s and

TAMs were the cells that contained most FP. The only exception from this was mRFP

containing neutrophils which on average contained more FP than TAM2s. Despite showing a

'non-canonical' uptake pattern, cDC1s infiltrating mRFP and mCherry-expressing tumors did

not contain an increased amount of antigen.

It remains unclear what causes the increased percentage of FP-positive tumor infiltrating

cDC1s in mRFP and mCherry-expressing tumors. Numbers of cDC1s are not different between

tumors expressing FPs irrespectively of whether they cause 'classical' or 'non-canonical' antigen

uptake patterns (Fig S3).

ZsGreen, ZsYellow, and DsRed can be tracked in the tdLN

All investigated FPs could be detected in tumor-infiltrating immune cells to a certain

extend when expressed in injected B16-F10 tumor cells. Next, we wanted to determine whether

selected fluorophores remained intact after trafficking of the containing immune cells to tumor-

draining lymph nodes (tdLN). We isolated these lymph nodes and analyzed migratory and

resident DCs as well as monocytes by flow cytometry. Migratory DCs and monocytes are

responsible for the transport of tumor-associated antigens to the tdLN (3), whereas resident DCs

can receive those antigens through synaptic transfer (5).

Analysis of lymph nodes harvested from mice injected with B16-ZsGreen tumor cells

revealed FP signal in both, migratory and resident DCs, as well as in monocytes. Almost 90 % of

monocytes, 10-20 % of cDC1s, and up to 50 % of cDC2s had detectable ZsGreen inside them

(Fig 6, Fig S4). Other FPs showed only a limited signal suggesting that proteins could have

already been degraded on their way to the draining lymph node. Percentages of ZsYellow and

DsRed positive immune cells in respective tdLNs where with up to 20 % highest in cDC2s and

monocytes, but severly lower than what has been seen for ZsGreen expressing tumors.

Fluorescent signal in cDC1s was barely detectable. No signal could be detected in tdLN from

mice injected with B16 tumors expressing AsRed, mRFP and mCherry. Interestingly, previous

reports showed that mCherry signal could be detected in tdLNs (3). This difference might be

due to different tumor models, different tumor sizes at harvest, and/or the mixture of wild type

and mCherry expressing cells in our cell line (Fig 1B).

In summary, ZsGreen was the only FP of the diverse groups of FPs tested that could be

tracked efficiently to the tdLN. ZsYellow and DsRed showed significantly less FP signal in

immune cells of the tdLN, but could still be considered as potential tools to analyze TAA

transfer between immune cells in tdLNs.

Conclusion

In summary, we tested six FPs: ZsGreen, ZsYellow, AsRed, DsRed, mRFP, and mCherry using an antigen uptake and transfer model. Despite comparable *in vitro* stabilities, those FPs showed vastly different loading patterns into tumor-infiltrating immune cells. ZsGreen is the only FP that can be efficiently tracked in lymph node resident DCs and remains the FP to be considered best for use in antigen uptake and transfer assays. We identified ZsYellow and DsRed as additional useful tools for these assays, however, tracking was not as efficient and consistent as for ZsGreen and might need optimization. Additionally, we identified two distinct patterns of FP-antigen loading into tumor-infiltrating immune cells. The 'classical' pattern showed preferential uptake of FPs by cDC2s and TAMs, whereas the 'non-canonical' pattern revealed highest antigen signal in tumor-infiltrating cDC1s. The underlying mechanisms for these distinct antigen uptake patterns remain elusive. Our data suggests that analyses of antigen uptake and transfer using different FPs should be carried out only after careful consideration of FP uptake patterns. Potential differences might arise as a result of the nature of the chosen FPs.

289 **Supplemental Figures**

290 **Fig S1. Similar subcellular localization of fluorescent proteins in FP-expressing B16-F10**

291 **cell lines.** Generated B16 melanoma cells with expression of fluorescent protein ZsGreen (**A**),

292 ZsYellow (**B**), AsRed, DsRed, mRFP, and mCherry (all **C**) respectively, overlayed with their

293 bright field images. Images of B16-F10 cell lines were taken to set background fluorescence.

A)

B)

Fig S2. Gating scheme for flow cytometry analysis of tumor and lymph node samples. A)

Cells collected from tumors were analyzed by flow cytometry using the shown gating methods.

Percentages of FP positive cells were analyzed in cDC1, cDC2, TAM1, TAM2, monocytes and

neutrophils. **B)** Lymph node cells were analyzed using these gating methods. The dump-

298   populations are B220, CD 90.2, NK 1.1 lineage marker negative cells. Percentage of FP positive

299   cells were analyzed in migratory cDC1, migratory cDC2, resident cDC1, resident cDC2 and

300   monocytes.

Fig S3. Tumor cell population distributions. All tumor cells collected were analyzed for their respective immune cell markers. The cell count for each type of immune cell is then plotted as percentages over the total number of live CD45+ cells.

304 **Fig S4. Lymph node cell population distributions.** All lymph node cells collected were

305 analyzed for their respective immune cell markers. The cell count for each type of immune cell is

306 then plotted as percentages over the total number of live CD45+ cells.

References

1.	Wang Y, Xiang Y, Xin VW, Wang XW, Peng XC, Liu XQ, et al. Dendritic cell biology and its role in tumor immunotherapy. J Hematol Oncol. 2020;13(1).

2.	Dersh D, Hollý J, Yewdell JW. A few good peptides: MHC class I-based cancer immunosurveillance and immunoevasion. Nat Rev Immunol [Internet]. 2021;21(2):116–28. Available from: http://dx.doi.org/10.1038/s41577-020-0390-6

3.	Roberts EW, Broz ML, Binnewies M, Headley MB, Nelson AE, Wolf DM, et al. Critical Role for CD103+/CD141+ Dendritic Cells Bearing CCR7 for Tumor Antigen Trafficking and Priming of T Cell Immunity in Melanoma. Cancer Cell [Internet]. 2016/07/19. 2016;30(2):324–36. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27424807

4.	Salmon H, Idoyaga J, Rahman A, Leboeuf M, Remark R, Jordan S, et al. Expansion and Activation of CD103+ Dendritic Cell Progenitors at the Tumor Site Enhances Tumor Responses to Therapeutic PD-L1 and BRAF Inhibition. Immunity. 2016;44(4):924–38.

5.	Ruhland MK, Roberts EW, Cai E, Mujal AM, Marchuk K, Beppler C, et al. Visualizing Synaptic Transfer of Tumor Antigens among Dendritic Cells. Cancer Cell [Internet]. 2020;37(6):786–99. Available from: https://doi.org/10.1016/j.ccell.2020.05.002

6.	Shaner NC, Steinbach PA, Tsien RY. A guide to choosing fluorescent proteins. Nat Methods. 2005;2(12):905–9.

7.	Lambert TJ. FPbase: a community-editable fluorescent protein database. Nat Methods. 2019;16(4):277–8.

8.	Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215(3):403–10.

9.	Matz M V., Fradkov AF, Labas YA, Savitsky AP, Zaraisky AG, Markelov ML, et al.

Fluorescent proteins from nonbioluminescent Anthozoa species. Nat Biotechnol.

1999;17(10):969–73.

10.   Shaner NC, Campbell RE, Steinbach PA, Giepmans BNG, Palmer AE, Tsien RY.

Improved monomeric red, orange and yellow fluorescent proteins derived from

Discosoma sp. red fluorescent protein. Nat Biotechnol. 2004;22(12):1567–72.

11.   Babała N, Bovens A, De Vries E, Iglesias-Guimarais V, Ahrends T, Krummel MF, et al.

Subcellular localization of antigen in keratinocytes dictates delivery of CD4 þ T-cell Help

for the CTL response upon therapeutic DNA vaccination into the Skin. Cancer Immunol

Res. 2018;6(7):835–47.

12.   Shinoda H, Shannon M, Nagai T. Fluorescent proteins for investigating biological events

in acidic environments. Int J Mol Sci. 2018;19(6).