Proteomics reveals unique identities of human TGF-β-induced and thymus-derived CD4+ regulatory T cells

Mark Mensink[1], Ellen Schrama[1], Eloy Cuadrado[2],

Maartje van den Biggelaar[3], Derk Amsen[2], Sander de Kivit[1,†,*] and Jannie Borst[1,†,*]

[1]Department of Immunology and Oncode Institute, Leiden University Medical Center, Leiden, The Netherlands

Department of Hematopoiesis[2] and Molecular and Cellular Homeostasis[3], Sanquin Research and Landsteiner

Laboratory, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

[†]These authors share senior authorship

* Correspondence: Dr Sander de Kivit (S.de_Kivit@lumc.nl) or Prof Jannie Borst (J.G.Borst@lumc.nl)

Keywords: conventional T cell, regulatory T cell, TGF-β, proteomics, signature

Abstract

The CD4$^+$ regulatory T (Treg) cell lineage, as defined by FOXP3 expression, comprises thymus-derived (t)Treg cells and peripherally induced (p)Treg cells. In human, naïve tTreg cells can be isolated from blood, while effector tTreg and pTreg cells cannot be purified reliably for lack of cell surface markers. As a model for Treg cells, studies often employ TGF-β-induced (i)Treg cells generated from CD4$^+$ conventional T (Tconv) cells *in vitro*. Here, we describe the relationship of iTreg cells to tTreg and Tconv cells, as optimally purified from human blood. Proteomic analysis revealed that each of these cell populations has a unique protein expression pattern before and after CD3/CD28-mediated activation. iTreg cells had limited overlap in protein expression with tTreg cells and exhibited markedly differential expression of proteins involved in signal transduction and metabolism. Whereas iTreg and Tconv cells responded strongly to CD3/CD28-mediated activation, tTreg cells showed a modest response, which reflects their adaptations in signal transduction pathways. As a benchmark, we used a previously defined proteomic signature that discerns *ex vivo* naïve and effector Treg cells from Tconv cells and includes conserved Treg cell properties. This Treg cell core signature was largely absent in iTreg cells, as exemplified by STAT4 expression that may contribute to pro-inflammatory functions. In addition, we used a proteomic signature that distinguishes *ex vivo* effector Treg cells from Tconv cells and naïve Treg cells. This effector Treg cell signature was partially present in iTreg cells, indicating that they do have protein expression features in common with *ex vivo* effector Treg cells. In conclusion, iTreg cells are distinct from tTreg cells and largely lack the Treg cell core proteomic signature, indicating that caution is warranted when using iTreg cells as a model for Treg cell populations found *in vivo*.

1   Introduction

Immune responses depend on a balance between the opposing activities of conventional (Tconv) and regulatory T (Treg) cells (1). While Tconv cells are key mediators of adaptive immunity, Treg cells are specialized in maintaining immune tolerance and controlling inflammation (1). Based on their origin, Treg cells can be classified into two main groups. Thymus-derived (t)Treg cells are primarily directed at self-antigens and stably express the Treg cell master transcription factor FOXP3, which is indispensable for their identity and suppressive function (2-6). In addition, they express the transcription factor Helios (IKZF2), which is consequently used as a marker of stable Treg cells (7, 8). Peripherally induced (p)Treg cells are converted from Tconv cells in response to foreign antigens (9) and reportedly have less stable FOXP3 expression (10-13), in particular recently after their development (14). Stable expression of FOXP3 is established epigenetically by demethylation of several regions within the *FOXP3* gene locus, including the so-called Treg-specific demethylated region (TSDR) in the CNS2 enhancer (5, 15).

Based on expression of CD45RA, human Treg cells from peripheral blood can be divided into naïve (CD45RA$^+$) and effector (CD45RA$^-$) Treg cells (16). Since conversion of Tconv cells into pTreg cells requires their activation, it stands to reason that the naïve Treg cell population is mainly composed of tTreg cells (17, 18). Such naïve Treg cells can differentiate into effector Treg cells in response to inflammatory signals (19). Therefore, the effector Treg cell population arguably consists of a mixture of pTreg cells and activated tTreg cells. Amongst human effector Treg cells, it is currently not possible to definitively discriminate tTreg and pTreg cells for a lack of cell surface markers (20).

Naïve Treg cells occur in low abundance in the blood (21, 22) and effector Treg cells from the blood are not amenable to expansion *in vitro* (16, 23). For these reasons, *in vitro* Treg cell studies often employ TGF-β-induced (i)Treg cells that are generated from Tconv cells (24-27). Such iTreg cells resemble Treg cells by their expression of FOXP3 and their suppressive capacity (24, 25). However, in terms of FOXP3 expression and suppressive function, iTreg cells are less stable than tTreg cells. For example, adoptively transferred iTreg cells in mice lost Foxp3 expression and failed

to prevent graft-versus-host disease, in contrast to tTreg cells (28). Similarly, others showed that iTreg

cells had reduced Foxp3 expression and suppressive capacity after removal of TGF-β *in vitro* and that

the majority of iTreg cells rapidly lost Foxp3 expression following adoptive transfer in wild-type mice

(29).

It is unclear whether iTreg cells represent an equivalent of *ex vivo* Treg cells. We recently

described that *ex vivo* naïve and effector Treg cells from human peripheral blood have a common

proteomic Treg cell signature, which sets them apart from other CD4[+] T cell subsets (22). This Treg

cell core signature reflects unique Treg cell properties regarding signal transduction, transcriptional

regulation and cell metabolism. In addition, we identified a proteomic signature that distinguishes *ex*

vivo effector Treg cells from both Tconv cells and naïve Treg cells. Using these proteomic signatures

as benchmarks, we compared in the current study the proteomes of iTreg, tTreg and Tconv cells to

determine to what extent *in vitro*-generated iTreg cells resemble *ex vivo* Treg cells.

2   Methods

2.1   Cell isolation and flow cytometric sorting

Human materials were obtained in compliance with the Declaration of Helsinki and the Dutch rules regarding the use of human materials from voluntary donors. Buffy coats from healthy anonymized male donors were obtained after their written informed consent, as approved by the internal ethical board of Sanquin, the Netherlands. Total CD4$^+$ T cells were collected from buffy coats by first performing a Ficoll-Paque Plus (GE Healthcare) density gradient centrifugation to isolate peripheral blood mononuclear cells (PBMC) and subsequently using CD4 magnetic MicroBeads (Miltenyi Biotec) according to the manufacturers' protocols. Alternatively, total CD4$^+$ T cells were directly isolated from buffy coats using the StraightFrom Buffy Coat CD4 MicroBead kit (Miltenyi Biotec) according to the manufacturer's instructions. To sort naïve Tconv and tTreg cells, CD4$^+$ T cells were stained with CD4-PE-Cy7, CD127-BV421 (BioLegend), CD25-PE (BD Biosciences), CD45RA-FITC (Immunotools) and GPA33-AF647 (30) monoclonal antibodies (mAbs) as indicated (**Supplementary Figure 1**). Cells were sorted on a MoFlo Astrios using Summit software version 6.2 (Beckman Coulter) or BD FACS Aria II using FACSDiva software version 8 (BD Biosciences). Either propidium iodide (Sigma) or near-IR dead cell stain kit (Invitrogen) was used as a live/dead marker.

2.1   T-cell expansion cultures and restimulation

Following cell sorting, human naïve Tconv and tTreg cells were cultured in 96-well round bottom plates (Greiner) at 1×10^4 cells per well using IMDM (Gibco, Life Technologies) supplemented with 8% FCS (Sigma), penicillin/streptomycin (Roche) and 300 IU/ml IL-2 (DuPont Medical), hereinafter referred to as T-cell medium, at 37°C and 5% CO_2. Cells were expanded using agonistic mAbs against CD3 (clone CLB-T3/4.E, IgE isotype, Sanquin, 0.1 µg/ml) and CD28 (clone CLB-CD28/1, Sanquin, 0.2 µg/ml) in solution. After four days, fresh T-cell medium was added. From day 7 to day 14, cells were cultured in 24-well plates (Greiner) at 5×10^5 cells per ml, while fresh T-cell medium was added on day 11. On day 14, cells were cultured in 6-well plates (Corning) at 1×10^6 cells per ml for four days

using fresh T-cell medium in the absence of agonistic mAbs. To generate iTreg cells, naïve Tconv

cells were cultured as described above in the presence of TGF-β (Peprotech, 10 ng/ml). Prior to

restimulation experiments, dead cells were removed by Ficoll-Paque Plus density gradient

centrifugation. T cells were restimulated as indicated using T-cell medium in the presence of agonistic

mAbs against CD3 (0.1 µg/ml) and CD28 (0.2 µg/ml), but without TGF-β.

2.2 Flow cytometry

For expression analysis of cell surface molecules, T cells were washed in PBS/1% FCS and stained

using the following mAbs in appropriate combinations: anti-OX40-PE-Cy7 (BioLegend), anti-GITR-

BV421 (BioLegend), anti-PD-1-APC-Cy7 (BioLegend), anti-CD39-BV510 (BioLegend), anti-ITGA4-

PE (CD49d) (BD Biosciences), anti-ICOS-PerCP-Cy5.5 (BioLegend), anti-CCR4-PE-Cy7

(BioLegend), anti-HLA-DR-APC-eFluor 780 (Invitrogen), anti-HLA-DR-BV605 (BD Biosciences) and

anti-FAS-BB700 (CD95) (BD Biosciences). Cell surface expression of CTLA-4 was measured by

adding anti-CTLA-4-PE-Dazzle594 mAb (BioLegend) in culture during restimulation for 24 h. Near-IR

dead cell stain kit (Invitrogen) was used as a live/dead marker. For intracellular staining, cells were

fixed and permeabilized using the FOXP3 transcription factor staining buffer set (Invitrogen) according

to the manufacturer's instructions. Next, cells were stained with combinations of anti-FOXP3-APC

(Invitrogen), anti-Helios-PE-Cy7, anti-CTLA-4-PE-Dazzle594 (BioLegend) and anti-NFATC2-AF488

(Cell Signaling Technology) mAbs. Flow cytometry was performed using a BD LSR Fortessa or BD

LSR II cell analyzer (BD Biosciences). Sample acquisition was performed using FACSDiva software

version 8 and data were analyzed using FlowJo software version 10.6.0.

2.3 Methylation assay of the TSDR

The methylation status of the TSDR within the *FOXP3* locus was determined in Tconv, tTreg and iTreg

cells as described previously (31). Only male donors were used for these analyses, given the X-

chromosomal location of the *FOXP3* locus. In brief, expanded T cells were resuspended in PBS for

proteinase K digestion, followed by bisulfite conversion of DNA using the EZ DNA Methylation-Direct

kit (Zymo Research) and methylation-specific quantitative PCR using iQTM SYBR Green Supermix

(Bio-Rad) (32). Methylation-specific primers were 5'-CGATAGGGTAGTTAGTTTTCGGAAC-3' and 5'-

CATTAACGTCATAACGACCGAA-3'. Demethylation-specific primers were 5'-TAGGGTAGT-

TAGTTTTTGGAATGA-3' and 5'-CCATTAACATCATAACAACCAAA-3'. Melt curve analysis was

performed on a LightCycler® 480-II (Roche). Methylation of the TSDR (%) was calculated using the

following formula: $100/(1+2^{Ct[CG]-Ct[TG]})$, where Ct[CG] is defined as Ct values obtained using

methylation-specific primers and Ct[TG] is defined as Ct values obtained using demethylation-specific

primers.

2.4   Suppression assay

Total human PBMCs were washed and resuspended in PBS, followed by incubation for 8 min using

5 μM CellTrace Violet dye (Invitrogen). Following fluorescent labeling, an equal volume of FCS was

added and cells were subsequently washed twice in IMDM supplemented with 8% FCS. CellTrace

Violet-labeled PBMCs were cocultured with expanded Tconv, tTreg or iTreg cells for 96 h at indicated

ratios, in the presence of anti-CD3 mAb (0.05 μg/ml). Proliferation of PBMCs was examined by tracing

CellTrace Violet dye dilution by flow cytometry on a BD LSR Fortessa or BD LSR II cell analyzer. Data

were analyzed using FlowJo software (version 10.6.0).

2.5   Proteomics

Proteomics data were generated as described before by Cuadrado et al. (22), including analysis of

expanded Tconv, tTreg and iTreg cells. In short, at least 1×10^6 expanded T cells were restimulated

for 24 h using T-cell medium in the presence of agonistic mAbs against CD3 (0.1 μg/ml) and CD28

(0.2 μg/ml) or not. Unstimulated and restimulated cells were washed, lysed and processed for mass

spectrometry. RAW mass spectrometry files were processed previously according to Cuadrado et al.

(22), here followed by the removal of potential contaminants and reverse hits using Perseus (version

1.6.12). Label-free quantification (LFQ) values were log_2-transformed and the biological replicates

were grouped based on cell type. Proteins with a minimum of 5 valid values in at least one cell type

were included for further analysis. For analysis of the effector Treg cell signature, a minimum of 4 valid

values in at least one cell type was used. Missing value imputation was performed by replacing

missing values by random numbers drawn from the lower part of the normal distribution (width = 0.3,

shift = 1.8). Next, proteomics data were analyzed by principal component analysis (PCA), ANOVA

and Student's *t*-test using Qlucore Omics Explorer (version 3.8). After z-score normalization, clusters

of proteins with similar expression patterns were identified using unsupervised hierarchical clustering

and visualized in heat maps. Protein clusters were subjected to overrepresentation analysis to assess

enrichment of Gene Ontology (GO) biological processes using the STRING app (version 1.6.0) in

Cytoscape (version 3.9.0). To refine the analysis of biological processes, a redundancy cutoff was

applied when significant GO terms contained large overlap based on enriched genes. The proteomic

response to restimulation was analyzed for each cell type using GO biological processes in gene set

enrichment analysis (GSEA) software (version 4.1.0). GSEA was performed using default parameters

and gene set permutations. To identify related groups of biological processes that were differentially

regulated following restimulation of T cells, the EnrichmentMap app (version 3.3.2) was used in

Cytoscape. The largest gene set clusters were displayed using manually assigned annotations that

describe the general biological processes represented by the clusters. To assess commonalities and

differences between the proteomic responses of the studied cell populations, the DiVenn tool (33) was

used.

2.6   Statistical analysis

Data were analyzed using GraphPad Prism (version 9.0.1), except for proteomics data. Statistical

analyses were performed as indicated in the figure legends. $\log_2$ transformation was performed when

data were not normally distributed. Data are presented as mean ± SEM and a two-sided $p < 0.05$ was

considered statistically significant.

3 Results

3.1 Validation of Tconv, tTreg and iTreg cell populations used in this study

To obtain the cell populations of interest, we first isolated CD4$^+$ T cells from human peripheral blood. These cells were flow cytometrically sorted based on a CD25^{lo}CD127^{hi}CD45RA$^+$GPA33^{int} phenotype to obtain naïve Tconv cells, or a CD25^{hi}CD127^{lo}CD45RA$^+$GPA33^{hi} phenotype to obtain naïve Treg cells, as described before (21, 31) (**Supplementary Figure 1**). After sorting, the cells were expanded using agonistic mAbs against CD3 and CD28 in the presence of IL-2. Cells derived from naïve Treg cells are hereinafter referred to as tTreg cells. To generate iTreg cells, TGF-β was added to Tconv cell cultures. After two weeks of expansion *in vitro*, cells were "rested" for four days prior to restimulation with anti-CD3/CD28 antibodies for 24 h and analysis (**Figure 1A**).

At this point, Tconv cells were FOXP3$^-$Helios$^-$ and largely retained this phenotype after CD3/CD28-mediated restimulation (**Figure 1B, C**). tTreg cells showed a uniform expression of FOXP3 and Helios (IKFZ2), both before and after restimulation, suggesting that the isolation and cell culture method generated pure tTreg cells as based on Helios expression (**Figure 1B, C**). The iTreg cells showed bimodal FOXP3 expression, with an upregulation of FOXP3 upon CD3/CD28-mediated restimulation, while Helios was minimally expressed, as described (7) (**Figure 1B, C**).

The TSDR of the *FOXP3* gene locus had the highly demethylated configuration in tTreg cells and was methylated in Tconv and iTreg cells, in agreement with published data (5, 34) (**Figure 1D**). Next, we functionally characterized the cell populations in a standard assay, testing their ability to suppress CD3-induced proliferation of CellTrace Violet-labeled Tconv cells from peripheral blood. In this assay, tTreg cells were suppressive and more so than iTreg cells, while Tconv cells were not, in agreement with earlier research (35) (**Figure 1E**). Together, these data validate the purification and cell culture protocols to generate tTreg, iTreg and Tconv cell populations for further characterization.

Figure 1

Figure 1. Characterization of Tconv, tTreg and iTreg cell populations. **(A)** Schematic overview of expansion protocols for Tconv, tTreg and iTreg cells. TGF-β was used to generate iTreg cells. Prior to analysis, expanded cells were restimulated for 24 h or not, as indicated in figure legends. **(B)** Flow cytometric analysis of FOXP3 and Helios (IKZF2) protein levels in Tconv, tTreg and iTreg cells at 24 h after restimulation with anti-CD3/CD28 mAbs or unstimulated control (representative of $n = 7$). **(C)** Quantification of the frequency of FOXP3+, FOXP3+Helios+ and FOXP3− cells in the resting Tconv, tTreg and iTreg cell populations after expansion ($n = 7$). **(D)** Analysis of the methylation status of the TSDR in the *FOXP3* locus in Tconv, tTreg and iTreg cells. One-way ANOVA with Tukey's *post hoc* test was used for statistical analysis ($n = 6-7$). **(E)** Assessment of the suppressive capacity of Tconv, tTreg and iTreg cells that were cocultured for four days with CellTrace Violet-labeled responder PBMCs in the presence of agonistic mAb to CD3. The percentage of dividing PBMCs is displayed ($n = 4$). One-way ANOVA with Bonferroni's *post hoc* test was used for statistical analysis ($n = 4$). **(C-E)** Data are presented as mean ± SEM. Sample size (n) represents cells from individual donors, analyzed in independent experiments. $*p < 0.05$, $**p < 0.01$, $****p < 0.0001$.

3.2 Expression of characteristic cell surface receptors on tTreg and iTreg cells

We next examined the expression of hallmark cell surface receptors that are associated with Treg

cells. Among these, CTLA-4 is a key mediator of suppression by tTreg cells (36) by reducing the

availability of CD80/86 for the costimulatory receptor CD28 on Tconv cells (37, 38). As CTLA-4

recycles between the cell surface and endosomes (39), we measured both total (cell surface plus

intracellular) CTLA-4 levels, as well as specifically the CTLA-4 levels on the cell surface only. Both

tTreg and iTreg cells showed higher total CTLA-4 expression than Tconv cells (**Figure 2A**,

Supplementary Figure 2). All three cell types upregulated total CTLA-4 expression upon CD3/CD28-

mediated restimulation, but levels were still higher in tTreg and iTreg cells than in Tconv cells. CTLA-

4 was only found on the cell surface of expanded cells after CD3/CD28-mediated restimulation, with

tTreg cells showing the highest levels (**Figure 2A**, **Supplementary Figure 2**). The costimulatory

receptors OX40 and GITR have received attention as Treg cell markers and mediators of Treg cell

function (40-42). These two receptors were expressed by all expanded cell types, but only following

CD3/CD28-mediated restimulation, with highest levels on tTreg cells (**Figure 2A**, **Supplementary**

Figure 2). Upon CD3/CD28-mediated restimulation, both iTreg and Tconv cells but not tTreg cells

upregulated PD-1, which is a co-inhibitory receptor for both Tconv and Treg cells (43) (**Figure 2A**,

Supplementary Figure 2). We also analyzed CD39 (ENTPD1), which facilitates extracellular ATP

hydrolysis that generates adenosine, a molecule with immunosuppressive effects on Tconv cells (44).

CD39 protein levels were highest on iTreg cells both before and after restimulation (**Figure 2A**,

Supplementary Figure 2). These data show that tTreg and iTreg cells exhibit different expression

patterns of these Treg cell-associated cell surface molecules (**Figure 2B**).

3.3 Proteomic analysis reveals that iTreg cells are distinct from tTreg and Tconv cells

We performed label-free quantitative proteomics to compare overall protein expression profiles in

resting Tconv, tTreg and iTreg cells in an unbiased manner. Based on the levels of the 4224 proteins

that were identified (**Supplementary Table 1**), PCA led to distinct grouping of expanded Tconv, tTreg

and iTreg cells (**Figure 3A**). PC1 explained most of the differences between the populations (32%). It

Figure 2

Figure 2. Expression of characteristic cell surface molecules associated with Treg cells. **(A)** Quantification of flow cytometric analysis of the protein expression of total CTLA-4 (cell surface plus intracellular) (n = 13), surface CTLA-4 (n = 8), surface OX40 (n = 8), surface GITR (n = 8), surface PD-1 (n = 8) and surface CD39 (ENTPD1) (n = 8) on indicated cell populations. Mean fluorescence intensity (MFI) is depicted on the y-axis. Two-way ANOVA with Tukey's *post hoc* test was used for statistical analysis. Data are presented as mean ± SEM. Sample size (n) represents cells from individual donors, analyzed in independent experiments. Asterisks without lines indicate comparisons between cells that were restimulated or not. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. **(B)** Table summarizing the data in **A**, indicating relatively low (–), intermediate (+/–) or high (+) expression levels.

revealed that iTreg and tTreg cells were distinct from one another and each more similar to Tconv

cells. In contrast, PC2 (16%) indicated commonalities between iTreg and tTreg cells that set them

apart from Tconv cells. Unsupervised hierarchical clustering, based on the 1290 proteins that are

differentially expressed between the three cell types (shown as a heat map of z-scores), also revealed

marked differences (**Figure 3B**, **Supplementary Table 2**). Each cell population exhibited a unique

pattern of clusters of proteins with relatively high or low expression. This unsupervised analysis

clustered iTreg and Tconv cells together in one branch of the dendrogram, whereas tTreg cells

clustered in a different branch, showing that iTreg cells are overall more related to Tconv cells than to

tTreg cells. In fact, only two small clusters of proteins were shared by iTreg and tTreg cells but not

Tconv cells: cluster 1 and cluster 6, which represented high or low expression of proteins compared

to Tconv cells, respectively. GO analysis indicated an overrepresentation of proteins associated with

protein folding and endoplasmic reticulum stress in these two clusters (**Figure 3C**, **Supplementary**

Figure 3). Notable were the shared high expression of Ikaros family member IKZF3 (Aiolos), a

cofactor for FOXP3-mediated transcriptional repression (45), and the shared low expression of YY1,

which inhibits FOXP3 expression and function (46).

Apart from this limited overlap in protein expression between iTreg and tTreg cells, a far greater

number of proteins (clusters 2–5 and 7–10) was differentially expressed between these cell types,

emphasizing their dissimilarity (**Figure 3B**). Proteins in clusters 2–5 were expressed at lower levels in

iTreg cells than in tTreg cells and were mainly associated with metabolic processes (**Figure 3D**). Part

of these clusters showed shared low protein expression by iTreg and Tconv cells (clusters 2–3),

whereas the other part showed low protein expression uniquely in iTreg cells (clusters 4–5)

(**Supplementary Figure 3**). Noteworthy was the lower expression of 31 proteins involved in oxidative

phosphorylation in iTreg cells compared to tTreg cells (**Figure 3E**). Of these, 23 proteins were located

in cluster 3, indicating that this lower expression was largely shared with Tconv cells. Clusters 7–10

consisted of proteins that were present at higher levels in iTreg cells than in tTreg cells and were

involved in signal transduction, cell division, vesicle-mediated transport and cell adhesion (**Figure 3F**).

These clusters included proteins related to lymphocyte activation, such as VAV1, ZAP70, STAT3 and

Figure 3

Figure 3. The iTreg, tTreg and Tconv cell proteomes are distinct. **(A)** PCA plot of the proteome of Tconv, tTreg and iTreg cells without restimulation (*n* = 3). **(B)** Heat map showing hierarchical clustering of the 1290 proteins that were differentially expressed between unstimulated Tconv, tTreg and iTreg cells (ANOVA, $p < 0.05$). Z-scores showing relative protein expression values are color-coded. Numbered boxes (1–10) indicate clusters of proteins. **(C)** STRING network of proteins that have shared high (cluster 1) or low (cluster 6) expression in iTreg and tTreg cells, relative to Tconv cells. Proteins with high or low expression are depicted as yellow or blue nodes, respectively. Protein associations according to STRING are connected with lines. **(D)** GO enrichment analysis of clusters 2–5 containing proteins with low expression in iTreg cells compared to tTreg cells, showing ten significantly enriched GO biological processes. **(E)** STRING network of proteins from clusters 2–5 that are involved in oxidative phosphorylation. Blue nodes indicate low expression in iTreg cells relative to tTreg cells. **(F)** GO enrichment analysis of clusters 7–10 containing proteins with high expression in iTreg cells compared to tTreg cells, showing ten significantly enriched GO biological processes. **(G)** STRING network of proteins from clusters 7–10 that are involved in lymphocyte activation. Yellow nodes indicate high expression in iTreg cells relative to tTreg cells.

STAT5B (**Figure 3G**). Part of these clusters indicated high protein expression shared by iTreg and

Tconv cells (clusters 7–8), while the other clusters indicated high protein expression uniquely in iTreg

cells (clusters 9–10) (**Supplementary Figure 3**). This analysis indicates that iTreg cells display a

protein expression profile related to that of Tconv cells that largely sets them apart from tTreg cells,

including metabolic features and potential responses to activation signals. However, iTreg cells do

share some features with tTreg cells, in particular regarding endoplasmic reticulum-associated

processes.

3.4   iTreg and tTreg cells respond differentially to CD3/CD28 stimulation

We also analyzed the proteome of Tconv, tTreg and iTreg cells following CD3/CD28-mediated

restimulation. Distinct clusters were identified in a PCA, showing that particularly Tconv and iTreg

cells altered their protein expression upon activation (**Figure 4A**). A much smaller response was

observed in tTreg cells, which clustered adjacent to each other regardless of stimulation. GSEA

emphasized the differential responses between the cell types (**Supplementary Figure 4A**), indicating

that iTreg cells, like Tconv cells and unlike tTreg cells, strongly responded, particularly by processes

associated with cell proliferation.

Linear regression analysis revealed that the response of Tconv and iTreg cells correlated more

than the response of tTreg and iTreg cells (**Supplementary Figure 4B**). Accordingly, a DiVenn

diagram revealed over 200 proteins that commonly altered their expression in both Tconv and iTreg

cells (**Figure 4B**). These proteins are primarily involved in cell activation, macromolecular biosynthesis

and responses to stress according to GO analysis (**Supplementary Figure 4C**). 46 proteins

commonly altered their expression in tTreg and iTreg cells, of which 17 also altered their expression

in Tconv cells (**Figure 4B**). These proteins are involved in a variety of cellular processes, including

lymphocyte activation (**Figure 4C**). For many proteins, however, levels were uniquely altered in either

Tconv, iTreg or tTreg cells. These proteins are involved in metabolism, RNA processing and

biogenesis (**Figure 4D**). Noteworthy, tTreg cells uniquely upregulated proteins involved in an

interferon-mediated defense response (**Figure 4E**). In conclusion, tTreg cells showed a relatively

Figure 4

Figure 4. Proteomic analysis of CD3/CD28-stimulated Tconv, tTreg and iTreg cells. **(A)** PCA plot of the proteome of Tconv, tTreg and iTreg cells that were restimulated or not ($n = 3$, $n = 2$ only for CD3/CD28-stimulated iTreg cells). **(B)** DiVenn diagram showing shared and unique proteins that were differentially expressed upon CD3/CD28 stimulation in Tconv, tTreg and iTreg cells (two-sided t-test, $p <$ 0.05). Upregliated proteins (yellow), downregulated proteins (blue) or proteins of which the expression changed in opposite directions (red) are shown. **(C)** STRING network of proteins that changed expression in both iTreg and tTreg cells upon restimulation. Proteins that also changed expression in Tconv cells are displayed as nodes with a grey border. Upregulated or downregulated proteins are depicted as yellow or blue nodes, respectively. Red nodes indicate proteins of which the expression changed in opposite directions, annotated with the cell type that showed upregulation. **(D)** GO enrichment analyses of unique differentially expressed proteins in restimulated Tconv, tTreg or iTreg cells, showing ten significantly enriched GO biological processes per cell type. **(E)** STRING analysis of proteins that uniquely changed expression in restimulated tTreg cells, showing a network of proteins involved in an interferon-mediated defense response.

modest response to CD3/CD28-mediated activation as compared to iTreg and Tconv cells.

Furthermore, all three cell types responded largely in a unique fashion, in particular regarding

processes related to macromolecular biosynthesis.

3.5 A core signature reflecting unique Treg cell properties is present in tTreg but not iTreg

cells

Recently, we established that *ex vivo* naïve and effector Treg cells from human peripheral blood have

a common proteomic signature that distinguishes them from CD4$^+$ Tconv cells, which we termed the

Treg cell core proteomic signature (22). This signature contains proteins with Treg cell-specific high

or low expression reflecting unique properties regarding signal transduction, transcriptional regulation,

iron storage, lysosomal processes and cell metabolism (**Figure 5A**, **B**). Here, we questioned whether

the Treg cell core signature is also present in iTreg cells. In our proteomics dataset, we detected 46

out of 51 proteins of the Treg cell core signature and evaluated their expression in expanded Tconv,

tTreg and iTreg cells. PCA indicated distinct clustering of these three cell types based on Treg cell

core signature protein levels, with iTreg cells clearly clustering away from tTreg cells (**Figure 5C**).

Hierarchical clustering based on this signature also separated the three cell types (**Figure 5D**). The

Treg cell core signature, with some exceptions, was preserved in tTreg cells following *in vitro*

expansion, as reported previously (22) (**Figure 5D**). tTreg and iTreg cells shared a number of the Treg

cell core signature proteins with characteristic high (cluster 1—including FOXP3, as expected) or low

(cluster 3) expression levels relative to Tconv cells. Moreover, iTreg cells, as opposed to *in vitro*-

expanded tTreg cells, expressed the core signature proteins MVP, HK1, ME2 and APOO at

characteristic levels relative to those in Tconv cells. Nevertheless, 26 out of 46 Treg cell core signature

proteins did not exhibit the characteristic expression pattern (compared to Tconv cells) in iTreg cells.

For instance, iTreg cells expressed high levels of the transcription factor STAT4, while low levels in

Treg cells are important to conserve Treg cell function (22). These proteomic data suggest that iTreg

cells share only a part of the unique characteristics of *ex vivo* Treg cells.

Figure 5

Figure 5. A Treg cell core signature is present in *in vitro*-expanded tTreg but not iTreg cells. **(A)** STRING network displaying 51 proteins of a previously defined Treg cell core proteomic signature (22). Upregulated and downregulated proteins are shown as yellow and blue nodes, respectively. **(B)** GO enrichment analysis of the Treg cell core signature proteins, showing ten significantly enriched GO biological processes. **(C)** PCA plot based on the expression of Treg cell core signature proteins in unstimulated Tconv, tTreg and iTreg cells (*n* = 3). **(D)** Heat map showing hierarchical clustering of 46 detected proteins of the Treg cell core signature in unstimulated Tconv, tTreg and iTreg cells. Z-scores depicting relative protein expression values are color-coded. Numbered boxes (1–5) indicate clusters of proteins. Protein names are colored according to high (yellow) or low (blue) relative expression in the Treg cell core signature. Proteins are marked (#) when the expression in the expanded and rested tTreg cells used here deviates from the described core signature (22).

3.6 iTreg cells do not acquire the Treg cell core signature upon CD3/CD28-mediated

activation

Since iTreg cells responded greatly to CD3/CD28-mediated restimulation in terms of altered protein

expression, we determined whether they obtained more of the Treg cell core signature under these

conditions. The core signature is present in both naïve and effector tTreg cells (22), indicating that

core signature proteins in tTreg cells do not respond majorly to cell activation. This was indeed

observed in a PCA, where levels of Treg cell core signature proteins did not alter in tTreg cells upon

CD3/CD28-mediated restimulation (**Figure 6A**). Interestingly, the levels of these proteins also did not

alter much in Tconv or iTreg cells upon restimulation (**Figure 6A**). Accordingly, hierarchical clustering

showed grouping of core protein expression levels based on cell type, with tTreg cells being most

distinct from Tconv and iTreg cells (**Figure 6B**). Part of the Treg cell core signature was shared by

iTreg and tTreg cells, including proteins with either high (cluster 2) or low (cluster 3) expression levels

compared to Tconv cells. Furthermore, core signature proteins HK1, APOO, ME2 and FKBP11 were

expressed at characteristic low levels in iTreg cells (cluster 1). However, the major part of the Treg

cell core signature proteins did not show characteristic expression levels in iTreg cells. Differential

expression of core signature proteins NFATC2, ITGA4 and ICOS between tTreg and iTreg cells was

confirmed by flow cytometry (**Figure 6C**, **Supplementary Figure 5**). Low expression of NFATC2 is a

critical feature of Treg cell function (22), whereas iTreg cells shared high expression with Tconv cells.

In conclusion, Treg cell core signature protein expression in iTreg cells was not more pronounced

after CD3/CD28-mediated activation.

3.7 iTreg cells acquire aspects of an effector Treg cell signature

Besides the Treg cell core signature, we previously observed a unique set of proteins that was

characteristically expressed at relatively high or low levels in *ex vivo* effector Treg cells, which sets

them apart from Tconv cells and naïve Treg cells (22). This was termed the effector Treg cell

signature, of which the highly expressed proteins are predominantly associated with DNA replication

(e.g. MCM proteins), apoptosis (e.g. FAS, CASP3), vesicular trafficking (e.g. STXBP2, NSF) and

Figure 6

Figure 6. Expression levels of Treg cell core signature proteins do not change in any cell type upon CD3/CD28-mediated restimulation. **(A)** PCA plot based on the expression of Treg cell core signature proteins in Tconv, tTreg and iTreg cells that were restimulated or not ($n = 3$, $n = 2$ only for CD3/CD28-stimulated iTreg cells). **(B)** Heat map showing hierarchical clustering of 46 detected proteins of the Treg cell core signature in Tconv, tTreg and iTreg cells that were CD3/CD28-stimulated or not. Z-scores depicting relative protein expression values are color-coded. Numbered boxes (1–5) indicate clusters of proteins. Protein names are colored according to high (yellow) or low (blue) relative expression in the Treg cell core signature. Marking (#) indicates the same as in Figure 5D. **(C)** Quantification of flow cytometric analysis of the protein expression of total NFATC2 ($n = 4$), surface ITGA4 (CD49d) ($n = 5$) and surface ICOS ($n = 5$) on indicated cell populations. MFI is depicted on the y-axis. Two-way ANOVA with Tukey's *post hoc* test was used for statistical analysis. Data are presented as mean ± SEM. Sample size (n) represents cells from individual donors, analyzed in independent experiments. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

cytoskeletal reorganization (e.g. TUBA1B, TUBB) (**Figure 7A**). Lowly expressed proteins of the

effector Treg cell signature primarily included those associated with signaling by the TCR/CD3

complex and CD28 (e.g. THEMIS, TRAT1, VAV1), phosphatidylinositol (e.g. INPP4A, INPP4B), NF-

κB (e.g. NFKB1, NFKB2) and JAK-STAT (e.g. STAT3, IL7R) (**Figure 7B**). Here, we examined the

effector Treg cell signature in unstimulated and restimulated Tconv, tTreg and iTreg cells and

questioned whether iTreg cells acquire effector Treg cell-like properties.

We detected all 41 signature proteins with high expression levels in effector Treg cells (effector

Treg^{hi}). Hierarchical clustering based on expression of these 41 signature proteins showed that iTreg

cells clustered away from Tconv and tTreg cells (**Figure 7C**). iTreg cells, as opposed to tTreg cells,

either before or after CD3/CD28-mediated activation, showed enrichment of the effector Treg^{hi}

signature with 21 out of 41 proteins exhibiting relatively high expression (clusters 1–2) (**Figure 7C**,

Supplementary Table 3). These proteins were primarily involved in DNA replication (e.g. MCM

proteins) and cytoskeletal processes such as cytokinesis (e.g. MYH9, CORO1C), indicating cell

proliferation. Additionally, the majority of 11 proteins in cluster 3 indicated relatively high expression

of effector Treg^{hi} signature proteins in both iTreg and tTreg cells but not Tconv cells. Proteins in this

cluster included CD39 (ENTPD1), a well-known effector molecule on Treg cells, which catalyzes a

critical step in the conversion of ATP into immunosuppressive adenosine (44). Flow cytometry verified

that CCR4 and HLA-DR had highest expression on iTreg cells, that CD39 was expressed on both

iTreg and tTreg cells (**Figure 2**), whereas FAS was highly expressed on tTreg cells, all in agreement

with the results from the proteome measurements (**Figure 7D**, **Supplementary Figure 6**). Thus,

altogether, 32 out of 41 Treg^{hi} signature proteins were also expressed at relatively high levels in iTreg

cells.

We detected 33 out of 39 signature proteins with characteristic low relative expression levels

in effector Treg cells (effector Treg^{lo}) (**Figure 7E**, **Supplementary Table 3**). Hierarchical clustering

based on expression of these proteins showed that iTreg cells clustered away from Tconv and tTreg

cells (**Figure 7E**). Among the 33 effector Treg^{lo} signature proteins, 16 exhibited relatively low

expression in iTreg cells (clusters 2–3). Proteins in cluster 2 were also expressed at low levels in tTreg

Figure 7

Figure 7. Aspects of an effector Treg cell signature are present in iTreg cells. **(A, B)** STRING networks showing 41 upregulated **(A)** or 39 downregulated proteins **(B)** of a previously defined effector Treg cell proteomic signature (22) as yellow and blue nodes, respectively. **(C)** Heat map showing hierarchical clustering of 41 detected proteins of the upregulated part of the effector Treg cell signature in Tconv, tTreg and iTreg cells that were restimulated or not ($n = 3$, $n = 2$ for CD3/CD28-stimulated iTreg cells). Z-scores showing relative protein expression values are color-coded. Numbered boxes (1–4) indicate clusters of proteins. CCR4, HLA-DRA and FAS are marked, as these molecules are also analyzed in **D**. **(D)** Quantification of flow cytometric analysis of the protein expression of surface CCR4 ($n = 4$), surface HLA-DR ($n = 4$) and surface FAS (CD95) ($n = 4$) on indicated cell populations. MFI is depicted on the y-axis. Two-way ANOVA with Tukey's *post hoc* test was used for statistical analysis. Data are presented as mean ± SEM. Sample size (n) represents cells from individual donors, analyzed in independent experiments. ***$p < 0.001$, ****$p < 0.0001$. **(E)** Heat map showing hierarchical clustering of 33 detected proteins of the downregulated part of the effector Treg cell signature in Tconv, tTreg and iTreg cells that were restimulated or not ($n = 3$, $n = 2$ for CD3/CD28-stimulated iTreg cells). Z-scores showing relative protein expression values are color-coded. Numbered boxes (1–3) indicate clusters of proteins.

cells and included INPP4A, THEMIS and members of the GIMAP family. Another group of proteins

was mainly expressed at low levels in iTreg cells, but not in tTreg cells (cluster 3). However, 17 out of

33 effector Treg[lo] signature proteins did not have the characteristic low expression in iTreg cells,

whereas they were lowly expressed in tTreg cells (cluster 1). These included proteins that play a role

in signaling downstream of immune receptors, such as NFKB1, NFKB2, CARD11, STAT3, VAV1,

STAM2 and INPP4B. These data show that the effector Treg cell signature is partially present in iTreg

cells, indicating that iTreg cells resemble *ex vivo* effector Treg cells to some extent.

4   Discussion

The Treg cell lineage can be subdivided into tTreg and pTreg cells based on their developmental

origin (15, 47). tTreg cells have escaped negative selection in the thymus and are mainly geared

towards self-antigen recognition (15). They comprise the majority of Treg cells *in vivo* and control

systemic and tissue-specific autoimmunity (48, 49). pTreg cells recognize foreign antigen and are

largely restricted to mucosal surfaces and the maternal-fetal interface (14, 50). Both cell types can be

distinguished in mice, where common and unique features in their phenotypes and transcriptome

profiles have been observed (11, 12). tTreg and pTreg cells complement each other in immune

homeostasis (51, 52), which may in part be due to their distinct TCR repertoires and differential

expression of certain transcriptional regulators (14, 53). In human, for lack of surface markers, effector

phenotype tTreg or pTreg cells cannot be discriminated, which complicates the interpretation of many

human Treg cell studies. Moreover, iTreg cells are often employed as an *in vitro* model for Treg cell

function, differentiation or metabolism, which increases the complexity of Treg cell literature.

Here, we questioned to what extent iTreg cells resemble tTreg cells in their protein expression,

focusing on conserved features of *ex vivo* Treg cells as previously defined (22). We performed a side-

by-side analysis of *in vitro*-generated iTreg cells with tTreg cells isolated to high purity from the blood

and also included Tconv cells as a reference. Proteomics gives more direct insight into the functional

cellular programs than transcriptomics, since protein and transcript levels often differ due to regulation

at protein and mRNA levels (22, 54). Our proteomic analysis pointed out that iTreg cells display a

protein expression profile that markedly sets them apart from tTreg cells.

Among the proteins with a relatively low expression in iTreg cells compared to tTreg cells in

the steady state, we found an enrichment of proteins involved in a variety of metabolic processes.

iTreg cells are frequently used to study Treg cell metabolism and reportedly utilize different metabolic

programs than other CD4[+] T cell subsets (26, 27, 55). Our findings now indicate dissimilarity between

iTreg and tTreg cells in terms of metabolism, suggesting that iTreg cells do not mirror (t)Treg cells

found *in vivo*. However, it remains to be resolved to what extent iTreg cells share metabolic features

with pTreg cells, which also derive from Tconv cells under the influence of TGF-β (14).

We also observed a substantial number of proteins with relatively high expression in iTreg cells

compared to tTreg cells in the steady state. Notable was the enrichment of signal transduction proteins

involved in lymphocyte activation. This different configuration of signaling molecules likely influences

the potential response of iTreg and tTreg cells to extracellular stimuli. Indeed, CD3/CD28-mediated

activation of these cells resulted in unique responses of different magnitudes. Whereas iTreg cells

strongly responded by differential expression of a large number of proteins, tTreg cells showed only

a modest response. Adaptations in signaling pathways downstream of immune receptors such as the

TCR and CD28 may underlie the attenuated signal transduction capacity in tTreg cells, which has

been shown for *ex vivo* Treg cells before (22). iTreg cells shared more features with Tconv cells than

with tTreg cells, which are associated with metabolism, responses to stress and lymphocyte activation.

Like iTreg cells, Tconv cells responded strongly to CD3/CD28-mediated activation, yet both cell types

responded largely in a unique manner.

The proteomic similarities between iTreg and tTreg cells were rather limited. GO analysis only

indicated an enrichment of proteins involved in endoplasmic reticulum-associated processes. We

confirmed that iTreg cells acquire Treg cell-like features such as increased expression of FOXP3 and

CTLA-4 (24), especially following CD3/CD28-mediated activation. FOXP3 expression in iTreg cells

was heterogeneous, as has been observed by others (26, 27, 55). We also observed shared high or

low expression of IKZF3 and YY1, respectively, which can interact with FOXP3 to regulate

transcription (45, 46). iTreg cells did not express Helios at similar high levels found in tTreg cells, but

minor upregulation was observed. It has been demonstrated in mice that, after recent egress from the

thymus, a small proportion of naïve CD4[+] Tconv cells has the capacity to develop into Helios[+] pTreg

cells (56). Nevertheless, Helios is preferentially expressed in tTreg cells and marks stable commitment

to the human Treg cell lineage (7, 8).

Stable expression of FOXP3 is established at the epigenetic level by demethylation of several

regions within the *FOXP3* gene locus. A critical region for sustained FOXP3 expression is the TSDR

in the CNS2 enhancer, which we confirmed to be highly demethylated in tTreg cells but not in iTreg

cells, as described (5, 15). However, a recent study comprehensively investigated mouse iTreg cells

at the epigenetic level after generation in the presence or absence of CD28 costimulation (57), as

strong CD28 signaling reportedly hampers iTreg cell development (58). iTreg cells deprived of CD28

costimulation showed increased DNA demethylation within Treg cell-specific genes, leading to more

stable Foxp3 expression and enhanced suppressive function, which even persisted after adoptive

transfer in mice (57).

We asked whether iTreg cells resemble Treg cells that are found *in vivo*. tTreg cells relocate

from lymphoid organs to peripheral tissues, where they are under the continuous influence of

environmental signals. They have a degree of plasticity and can adapt themselves to locally resident

effector T cells by responding to specific cytokines (49). This involves the gain of lineage-determining

transcription factors such as GATA-3 or T-bet in addition to FOXP3, which alters their functionality.

However, tTreg cells retain core features that safeguard their suppressive nature and prevent them

from exerting pro-inflammatory functions (22). To establish whether iTreg cells acquire such features,

we used proteomic signatures from human Treg cells that were analyzed directly from peripheral blood

(22).

The first benchmark we used was the Treg cell core signature, a conserved set of proteins

between naïve and effector Treg cells compared to other CD4[+] T cell subsets that emphasizes critical

Treg cell aspects (22). This signature strongly suggests differential wiring of signaling pathways, given

the constitutive differential expression of key signal transduction components between Treg and

Tconv cells. We observed that iTreg cells only share a limited part of these conserved characteristics,

while the signature was largely present in tTreg cells, regardless of whether the cells were resting or

had been restimulated via CD3/CD28. As iTreg cells do not possess the signal transduction profile of

the Treg cell core signature, this likely leads to differential wiring and a different response to stimuli.

The Treg cell core signature includes low expression of STAT4 and NFATC2, which are

mediators of inflammatory cytokine expression (59, 60). Low expression of these molecules may aid

Treg cells to protect their anti-inflammatory nature in a pro-inflammatory environment, while they can

still respond to pro-inflammatory cytokines through other pathways (22). It has been demonstrated

that deliberate elevation of STAT4 expression in tTreg cells increases their susceptibility to pro-

inflammatory cytokines, provoking effector cytokine production and loss of FOXP3 expression (22).

Both STAT4 and NFATC2 had relatively high expression in iTreg cells compared to tTreg cells,

suggesting that iTreg cells are more susceptible to respond to pro-inflammatory cues in a pro-

inflammatory manner, such as having the capacity to produce IFN-γ (61). In addition to the fragile

epigenetic stability of iTreg cells, our findings now suggest that the susceptibility of iTreg cells to

abandoning the Treg cell fate upon exposure to pro-inflammatory cytokines may also be caused by

their high expression of involved mediators, such as STAT4 and NFATC2.

The second benchmark we examined was an effector Treg cell signature, which sets effector

Treg cells apart from naïve Treg cells and other CD4$^+$ T cell subsets (22). Relative to protein

expression in tTreg and Tconv cells, iTreg cells acquired aspects of the effector Treg cell signature,

indicating that iTreg cells may resemble *ex vivo* effector Treg cells to some extent. The enrichment of

the effector Treg cell signature in iTreg cells may fit with the idea that the effector Treg cell population

contains both activated tTreg cells and pTreg cells (62, 63).

We propose that the core and effector Treg cell proteomic signatures (22) are useful reference

tools to define Treg cell populations. In conclusion, our global proteomic analysis showed that iTreg

and tTreg cells each share more protein expression with Tconv cells than with one another. Moreover,

part of the overall protein expression profile of iTreg cells is unique. Thus, despite the acquisition of

certain Treg cell-like features, the conversion of Tconv cells into iTreg cells generates cells that

become even more distinct from tTreg cells than Tconv cells are. Therefore, we propose that

experiments in which iTreg cells—cultured in commonly used conditions as in the present article—are

employed as a model for studying Treg cell differentiation, function or metabolism should be

interpreted with caution.

5   Data availability

Proteomics data on tTreg and Tconv cells have been deposited as outlined before (22). Proteomics data on iTreg cells will be deposited before publication of this manuscript.

6   Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7   Author contributions

MM, SdK and JB designed the study and wrote the manuscript, MM and SdK performed experiments and analyzed the data, ES provided technical assistance, EC provided data and insights, MvdB performed mass spectrometry measurements, DA provided conceptual advice in the study design and contributed to the manuscript.

8   Funding

This study was supported by Oncode, ZonMW-TOP grant 40-00812-98-13071, and grants ICI-00014 and ICI-00025 from the Institute for Chemical Immunology, funded by ZonMW Gravitation, and Strategic Funds from the LUMC.

9   Acknowledgements

We thank the staff members of the flow cytometry facility at the Netherlands Cancer Institute and the flow cytometry core facility at Leiden University Medical Center for their technical assistance.

10 References

1.	Sakaguchi S, Yamaguchi T, Nomura T, Ono M. Regulatory T cells and immune tolerance. *Cell* (2008) 133(5):775-87. Epub 2008/05/31. doi: 10.1016/j.cell.2008.05.009. PubMed PMID: 18510923.

2.	Jordan MS, Boesteanu A, Reed AJ, Petrone AL, Holenbeck AE, Lerman MA, et al. Thymic selection of CD4+CD25+ regulatory T cells induced by an agonist self-peptide. *Nat Immunol* (2001) 2(4):301-6. Epub 2001/03/29. doi: 10.1038/86302. PubMed PMID: 11276200.

3.	Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* (2003) 4(4):330-6. Epub 20030303. doi: 10.1038/ni904. PubMed PMID: 12612578.

4.	Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* (2003) 299(5609):1057-61. Epub 20030109. doi: 10.1126/science.1079490. PubMed PMID: 12522256.

5.	Floess S, Freyer J, Siewert C, Baron U, Olek S, Polansky J, et al. Epigenetic control of the foxp3 locus in regulatory T cells. *PLoS Biol* (2007) 5(2):e38. Epub 2007/02/15. doi: 10.1371/journal.pbio.0050038. PubMed PMID: 17298177; PubMed Central PMCID: PMCPMC1783672 authors S. Olek and U. Baron, who are a founder and an employee of the company Epiontis, respectively.

6.	Kieback E, Hilgenberg E, Stervbo U, Lampropoulou V, Shen P, Bunse M, et al. Thymus-Derived Regulatory T Cells Are Positively Selected on Natural Self-Antigen through Cognate Interactions of High Functional Avidity. *Immunity* (2016) 44(5):1114-26. Epub 2016/05/19. doi: 10.1016/j.immuni.2016.04.018. PubMed PMID: 27192577.

7.	Thornton AM, Korty PE, Tran DQ, Wohlfert EA, Murray PE, Belkaid Y, et al. Expression of Helios, an Ikaros transcription factor family member, differentiates thymic-derived from peripherally induced Foxp3+ T regulatory cells. *J Immunol* (2010) 184(7):3433-41. Epub 2010/02/26. doi: 10.4049/jimmunol.0904028. PubMed PMID: 20181882; PubMed Central PMCID: PMCPMC3725574.

8.	Lam AJ, Uday P, Gillies JK, Levings MK. Helios is a marker, not a driver, of human Treg stability. *Eur J Immunol* (2022) 52(1):75-84. Epub 20211013. doi: 10.1002/eji.202149318. PubMed PMID: 34561855.

9.	Josefowicz SZ, Lu LF, Rudensky AY. Regulatory T cells: mechanisms of differentiation and function. *Annu Rev Immunol* (2012) 30:531-64. Epub 2012/01/10. doi: 10.1146/annurev.immunol.25.022106.141623. PubMed PMID: 22224781; PubMed Central PMCID: PMCPMC6066374.

10.   Miyao T, Floess S, Setoguchi R, Luche H, Fehling HJ, Waldmann H, et al. Plasticity of Foxp3(+) T cells
reflects promiscuous Foxp3 expression in conventional T cells but not reprogramming of regulatory T cells.
Immunity (2012) 36(2):262-75. Epub 20120209. doi: 10.1016/j.immuni.2011.12.012. PubMed PMID: 22326580.

11.   Weiss JM, Bilate AM, Gobert M, Ding Y, Curotto de Lafaille MA, Parkhurst CN, et al. Neuropilin 1 is
expressed on thymus-derived natural regulatory T cells, but not mucosa-generated induced Foxp3+ T reg cells.
J Exp Med (2012) 209(10):1723-42, S1. Epub 20120910. doi: 10.1084/jem.20120914. PubMed PMID:
22966001; PubMed Central PMCID: PMCPMC3457733.

12.   Yadav M, Louvet C, Davini D, Gardner JM, Martinez-Llordella M, Bailey-Bucktrout S, et al. Neuropilin-1
distinguishes natural and inducible regulatory T cells among regulatory T cell subsets in vivo. *J Exp Med* (2012)
209(10):1713-22, S1-19. Epub 20120910. doi: 10.1084/jem.20120822. PubMed PMID: 22966003; PubMed
Central PMCID: PMCPMC3457729.

13.   Sawant DV, Vignali DA. Once a Treg, always a Treg? *Immunol Rev* (2014) 259(1):173-91. doi:
10.1111/imr.12173. PubMed PMID: 24712466; PubMed Central PMCID: PMCPMC4008876.

14.   Josefowicz SZ, Niec RE, Kim HY, Treuting P, Chinen T, Zheng Y, et al. Extrathymically generated
regulatory T cells control mucosal TH2 inflammation. *Nature* (2012) 482(7385):395-9. Epub 20120208. doi:
10.1038/nature10772. PubMed PMID: 22318520; PubMed Central PMCID: PMCPMC3485072.

15.   Ohkura N, Kitagawa Y, Sakaguchi S. Development and maintenance of regulatory T cells. *Immunity*
(2013) 38(3):414-23. doi: 10.1016/j.immuni.2013.03.002. PubMed PMID: 23521883.

16.   Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, et al. Functional delineation and differentiation
dynamics of human CD4+ T cells expressing the FoxP3 transcription factor. *Immunity* (2009) 30(6):899-911.
Epub 20090521. doi: 10.1016/j.immuni.2009.03.019. PubMed PMID: 19464196.

17.   Hoffmann P, Eder R, Boeld TJ, Doser K, Piseshka B, Andreesen R, et al. Only the CD45RA+
subpopulation of CD4+CD25high T cells gives rise to homogeneous regulatory T-cell lines upon in vitro
expansion. *Blood* (2006) 108(13):4260-7. Epub 20060817. doi: 10.1182/blood-2006-06-027409. PubMed PMID:
16917003.

18.   Canavan JB, Scotta C, Vossenkamper A, Goldberg R, Elder MJ, Shoval I, et al. Developing in vitro
expanded CD45RA+ regulatory T cells as an adoptive cell therapy for Crohn's disease. *Gut* (2016) 65(4):584-
94. Epub 20150224. doi: 10.1136/gutjnl-2014-306919. PubMed PMID: 25715355; PubMed Central PMCID:
PMCPMC4819603.

19.   Smigiel KS, Richards E, Srivastava S, Thomas KR, Dudda JC, Klonowski KD, et al. CCR7 provides
localized access to IL-2 and defines homeostatically distinct regulatory T cell subsets. *J Exp Med* (2014)
211(1):121-36. Epub 20131230. doi: 10.1084/jem.20131142. PubMed PMID: 24378538; PubMed Central
PMCID: PMCPMC3892972.

20.   Shevach EM, Thornton AM. tTregs, pTregs, and iTregs: similarities and differences. *Immunol Rev*
(2014) 259(1):88-102. Epub 2014/04/10. doi: 10.1111/imr.12160. PubMed PMID: 24712461; PubMed Central
PMCID: PMCPMC3982187.

21.   Opstelten R, de Kivit S, Slot MC, van den Biggelaar M, Iwaszkiewicz-Grzes D, Gliwinski M, et al. GPA33:
A Marker to Identify Stable Human Regulatory T Cells. *J Immunol* (2020) 204(12):3139-48. Epub 2020/05/06.
doi: 10.4049/jimmunol.1901250. PubMed PMID: 32366581.

22.   Cuadrado E, van den Biggelaar M, de Kivit S, Chen YY, Slot M, Doubal I, et al. Proteomic Analyses of
Human Regulatory T Cells Reveal Adaptations in Signaling Pathways that Protect Cellular Identity. *Immunity*
(2018) 48(5):1046-59 e6. Epub 2018/05/13. doi: 10.1016/j.immuni.2018.04.008. PubMed PMID: 29752063.

23.   Putnam AL, Brusko TM, Lee MR, Liu W, Szot GL, Ghosh T, et al. Expansion of human regulatory T-
cells from patients with type 1 diabetes. *Diabetes* (2009) 58(3):652-62. Epub 20081215. doi: 10.2337/db08-
1168. PubMed PMID: 19074986; PubMed Central PMCID: PMCPMC2646064.

24.   Chen W, Jin W, Hardegen N, Lei KJ, Li L, Marinos N, et al. Conversion of peripheral CD4+CD25- naive
T cells to CD4+CD25+ regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med* (2003)
198(12):1875-86. Epub 2003/12/17. doi: 10.1084/jem.20030152. PubMed PMID: 14676299; PubMed Central
PMCID: PMCPMC2194145.

25.   Fantini MC, Becker C, Monteleone G, Pallone F, Galle PR, Neurath MF. Cutting edge: TGF-beta induces
a regulatory phenotype in CD4+CD25- T cells through Foxp3 induction and down-regulation of Smad7. *J
Immunol* (2004) 172(9):5149-53. Epub 2004/04/22. doi: 10.4049/jimmunol.172.9.5149. PubMed PMID:
15100250.

26.   Michalek RD, Gerriets VA, Jacobs SR, Macintyre AN, MacIver NJ, Mason EF, et al. Cutting edge:
distinct glycolytic and lipid oxidative metabolic programs are essential for effector and regulatory CD4+ T cell
subsets. *J Immunol* (2011) 186(6):3299-303. Epub 2011/02/15. doi: 10.4049/jimmunol.1003613. PubMed PMID:
21317389; PubMed Central PMCID: PMCPMC3198034.

27.   Gerriets VA, Kishton RJ, Nichols AG, Macintyre AN, Inoue M, Ilkayeva O, et al. Metabolic programming
and PDHK1 control CD4+ T cell subsets and inflammation. *J Clin Invest* (2015) 125(1):194-207. Epub
2014/12/02. doi: 10.1172/JCI76012. PubMed PMID: 25437876; PubMed Central PMCID: PMCPMC4382238.

28.   Koenecke C, Czeloth N, Bubke A, Schmitz S, Kissenpfennig A, Malissen B, et al. Alloantigen-specific
de novo-induced Foxp3+ Treg revert in vivo and do not protect from experimental GVHD. *Eur J Immunol* (2009)
39(11):3091-6. Epub 2009/09/15. doi: 10.1002/eji.200939432. PubMed PMID: 19750478.

29.   Selvaraj RK, Geiger TL. A kinetic and dynamic analysis of Foxp3 induced in T cells by TGF-beta. *J
Immunol* (2007) 179(2):11 p following 1390. PubMed PMID: 17695668.

30.   Herbertson RA, Tebbutt NC, Lee FT, Gill S, Chappell B, Cavicchiolo T, et al. Targeted chemoradiation
in metastatic colorectal cancer: a phase I trial of 131I-huA33 with concurrent capecitabine. *J Nucl Med* (2014)
55(4):534-9. Epub 2014/02/22. doi: 10.2967/jnumed.113.132761. PubMed PMID: 24556590.

31.   de Kivit S, Mensink M, Hoekstra AT, Berlin I, Derks RJE, Both D, et al. Stable human regulatory T cells
switch to glycolysis following TNF receptor 2 costimulation. *Nat Metab* (2020) 2(10):1046-61. Epub 2020/09/23.
doi: 10.1038/s42255-020-00271-w. PubMed PMID: 32958937.

32.   Zhuo C, Li Z, Xu Y, Wang Y, Li Q, Peng J, et al. Higher FOXP3-TSDR demethylation rates in adjacent
normal tissues in patients with colon cancer were associated with worse survival. *Mol Cancer* (2014) 13:153.
Epub 2014/06/19. doi: 10.1186/1476-4598-13-153. PubMed PMID: 24938080; PubMed Central PMCID:
PMCPMC4074420.

33.   Sun L, Dong S, Ge Y, Fonseca JP, Robinson ZT, Mysore KS, et al. DiVenn: An Interactive and
Integrated Web-Based Visualization Tool for Comparing Gene Lists. *Front Genet* (2019) 10:421. Epub
2019/05/28. doi: 10.3389/fgene.2019.00421. PubMed PMID: 31130993; PubMed Central PMCID:
PMCPMC6509638.

34.   Baron U, Floess S, Wieczorek G, Baumann K, Grutzkau A, Dong J, et al. DNA demethylation in the
human FOXP3 locus discriminates regulatory T cells from activated FOXP3(+) conventional T cells. *Eur J
Immunol* (2007) 37(9):2378-89. doi: 10.1002/eji.200737594. PubMed PMID: 17694575.

35.   Tran DQ, Ramsey H, Shevach EM. Induction of FOXP3 expression in naive human CD4+FOXP3 T cells
by T-cell receptor stimulation is transforming growth factor-beta dependent but does not confer a regulatory
phenotype. *Blood* (2007) 110(8):2983-90. Epub 2007/07/24. doi: 10.1182/blood-2007-06-094656. PubMed
PMID: 17644734; PubMed Central PMCID: PMCPMC2018674.

36.   Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z, et al. CTLA-4 control over
Foxp3+ regulatory T cell function. *Science* (2008) 322(5899):271-5. doi: 10.1126/science.1160062. PubMed
PMID: 18845758.

37.   Qureshi OS, Zheng Y, Nakamura K, Attridge K, Manzotti C, Schmidt EM, et al. Trans-endocytosis of
CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science* (2011) 332(6029):600-3.
Epub 20110407. doi: 10.1126/science.1202947. PubMed PMID: 21474713; PubMed Central PMCID:
PMCPMC3198051.

38.   Walker LS, Sansom DM. Confusing signals: recent progress in CTLA-4 biology. *Trends Immunol* (2015)
36(2):63-70. Epub 2015/01/15. doi: 10.1016/j.it.2014.12.001. PubMed PMID: 25582039; PubMed Central
PMCID: PMCPMC4323153.

39.   Linsley PS, Bradshaw J, Greene J, Peach R, Bennett KL, Mittler RS. Intracellular trafficking of CTLA-4
and focal localization towards sites of TCR engagement. *Immunity* (1996) 4(6):535-43. Epub 1996/06/01. doi:
10.1016/s1074-7613(00)80480-x. PubMed PMID: 8673700.

40.   So T, Lee SW, Croft M. Immune regulation and control of regulatory T cells by OX40 and 4-1BB.
Cytokine Growth Factor Rev (2008) 19(3-4):253-62. Epub 20080527. doi: 10.1016/j.cytogfr.2008.04.003.
PubMed PMID: 18508403; PubMed Central PMCID: PMCPMC2486494.

41.   McHugh RS, Whitters MJ, Piccirillo CA, Young DA, Shevach EM, Collins M, et al. CD4(+)CD25(+)
immunoregulatory T cells: gene expression analysis reveals a functional role for the glucocorticoid-induced TNF
receptor. *Immunity* (2002) 16(2):311-23. doi: 10.1016/s1074-7613(02)00280-7. PubMed PMID: 11869690.

42.   Ndhlovu LC, Takeda I, Sugamura K, Ishii N. Expanding role of T-cell costimulators in regulatory T-cell
function: recent advances in accessory molecules expressed on both regulatory and nonregulatory T cells. *Crit
Rev Immunol* (2004) 24(4):251-66. doi: 10.1615/critrevimmunol.v24.i4.30. PubMed PMID: 15588225.

43.   Tan CL, Kuchroo JR, Sage PT, Liang D, Francisco LM, Buck J, et al. PD-1 restraint of regulatory T cell
suppressive activity is critical for immune tolerance. *J Exp Med* (2021) 218(1). doi: 10.1084/jem.20182232.
PubMed PMID: 33045061; PubMed Central PMCID: PMCPMC7543091.

44.   Borsellino G, Kleinewietfeld M, Di Mitri D, Sternjak A, Diamantini A, Giometto R, et al. Expression of
ectonucleotidase CD39 by Foxp3+ Treg cells: hydrolysis of extracellular ATP and immune suppression. *Blood*
(2007) 110(4):1225-32. Epub 2007/04/24. doi: 10.1182/blood-2006-12-064527. PubMed PMID: 17449799.

45.   Kwon HK, Chen HM, Mathis D, Benoist C. Different molecular complexes that mediate transcriptional
induction and repression by FoxP3. *Nat Immunol* (2017) 18(11):1238-48. Epub 20170911. doi: 10.1038/ni.3835.
PubMed PMID: 28892470; PubMed Central PMCID: PMCPMC5679728.

46.   Hwang SS, Jang SW, Kim MK, Kim LK, Kim BS, Kim HS, et al. YY1 inhibits differentiation and function
of regulatory T cells by blocking Foxp3 expression and activity. *Nat Commun* (2016) 7:10789. Epub 20160219.
doi: 10.1038/ncomms10789. PubMed PMID: 26892542; PubMed Central PMCID: PMCPMC4762897.

47.   Bilate AM, Lafaille JJ. Induced CD4+Foxp3+ regulatory T cells in immune tolerance. *Annu Rev Immunol*
(2012) 30:733-58. Epub 20120106. doi: 10.1146/annurev-immunol-020711-075043. PubMed PMID: 22224762.

48.   Burzyn D, Benoist C, Mathis D. Regulatory T cells in nonlymphoid tissues. *Nat Immunol* (2013)
14(10):1007-13. Epub 20130918. doi: 10.1038/ni.2683. PubMed PMID: 24048122; PubMed Central PMCID:
PMCPMC4708287.

49.   Liston A, Gray DH. Homeostatic control of regulatory T cell diversity. *Nat Rev Immunol* (2014) 14(3):154-
65. Epub 20140131. doi: 10.1038/nri3605. PubMed PMID: 24481337.

50.   Samstein RM, Josefowicz SZ, Arvey A, Treuting PM, Rudensky AY. Extrathymic generation of
regulatory T cells in placental mammals mitigates maternal-fetal conflict. *Cell* (2012) 150(1):29-38. doi:
10.1016/j.cell.2012.05.031. PubMed PMID: 22770213; PubMed Central PMCID: PMCPMC3422629.

51.   Haribhai D, Williams JB, Jia S, Nickerson D, Schmitt EG, Edwards B, et al. A requisite role for induced
regulatory T cells in tolerance based on expanding antigen receptor diversity. *Immunity* (2011) 35(1):109-22.
Epub 20110630. doi: 10.1016/j.immuni.2011.03.029. PubMed PMID: 21723159; PubMed Central PMCID:
PMCPMC3295638.

52.   Huang YJ, Haist V, Baumgartner W, Fohse L, Prinz I, Suerbaum S, et al. Induced and thymus-derived
Foxp3(+) regulatory T cells share a common niche. *Eur J Immunol* (2014) 44(2):460-8. Epub 20131030. doi:
10.1002/eji.201343463. PubMed PMID: 24170313.

53.   Ohnmacht C, Park JH, Cording S, Wing JB, Atarashi K, Obata Y, et al. The microbiota regulates type 2
immunity through RORgammat(+) T cells. *Science* (2015) 349(6251):989-93. Epub 20150709. doi:
10.1126/science.aac4263. PubMed PMID: 26160380.

54.   Vogel C, Marcotte EM. Insights into the regulation of protein abundance from proteomic and
transcriptomic analyses. *Nat Rev Genet* (2012) 13(4):227-32. Epub 20120313. doi: 10.1038/nrg3185. PubMed
PMID: 22411467; PubMed Central PMCID: PMCPMC3654667.

55.   Angelin A, Gil-de-Gomez L, Dahiya S, Jiao J, Guo L, Levine MH, et al. Foxp3 Reprograms T Cell
Metabolism to Function in Low-Glucose, High-Lactate Environments. *Cell Metab* (2017) 25(6):1282-93 e7. Epub
20170413. doi: 10.1016/j.cmet.2016.12.018. PubMed PMID: 28416194; PubMed Central PMCID:
PMCPMC5462872.

56.   Pratama A, Schnell A, Mathis D, Benoist C. Developmental and cellular age direct conversion of CD4+
T cells into RORgamma+ or Helios+ colon Treg cells. *J Exp Med* (2020) 217(1). doi: 10.1084/jem.20190428.
PubMed PMID: 31685531; PubMed Central PMCID: PMCPMC7037252.

57.   Mikami N, Kawakami R, Chen KY, Sugimoto A, Ohkura N, Sakaguchi S. Epigenetic conversion of
conventional T cells into regulatory T cells by CD28 signal deprivation. *Proc Natl Acad Sci U S A* (2020)
117(22):12258-68. Epub 20200515. doi: 10.1073/pnas.1922600117. PubMed PMID: 32414925; PubMed
Central PMCID: PMCPMC7275710.

58.   Semple K, Nguyen A, Yu Y, Wang H, Anasetti C, Yu XZ. Strong CD28 costimulation suppresses
induction of regulatory T cells from naive precursors through Lck signaling. *Blood* (2011) 117(11):3096-103.
Epub 20110118. doi: 10.1182/blood-2010-08-301275. PubMed PMID: 21245484; PubMed Central PMCID:
PMCPMC3062312.

59.   O'Shea JJ, Lahesmaa R, Vahedi G, Laurence A, Kanno Y. Genomic views of STAT function in CD4+ T
helper cell differentiation. *Nat Rev Immunol* (2011) 11(4):239-50. Epub 2011/03/26. doi: 10.1038/nri2958.
PubMed PMID: 21436836; PubMed Central PMCID: PMCPMC3070307.

60.   Macian F. NFAT proteins: key regulators of T-cell development and function. *Nat Rev Immunol* (2005)
5(6):472-84. Epub 2005/06/02. doi: 10.1038/nri1632. PubMed PMID: 15928679.

61.   Chen Q, Kim YC, Laurence A, Punkosdy GA, Shevach EM. IL-2 controls the stability of Foxp3
expression in TGF-beta-induced Foxp3+ T cells in vivo. *J Immunol* (2011) 186(11):6329-37. Epub 20110427.
doi: 10.4049/jimmunol.1100061. PubMed PMID: 21525380; PubMed Central PMCID: PMCPMC3098943.

62.   Vukmanovic-Stejic M, Zhang Y, Cook JE, Fletcher JM, McQuaid A, Masters JE, et al. Human CD4+
CD25hi Foxp3+ regulatory T cells are derived by rapid turnover of memory populations in vivo. *J Clin Invest*
(2006) 116(9):2423-33. doi: 10.1172/JCI28941. PubMed PMID: 16955142; PubMed Central PMCID:
PMCPMC1555646.

63.   Opstelten R, Amsen D. Separating the wheat from the chaff: Making sense of Treg heterogeneity for
better adoptive cellular therapy. *Immunol Lett* (2021) 239:96-112. Epub 20210304. doi:
10.1016/j.imlet.2021.03.002. PubMed PMID: 33676975.

Supplementary Figure 1

Supplementary Figure 1. Gating strategy to sort naïve Tconv and naïve tTreg cells from human peripheral blood by flow cytometry. Pre-enriched total CD4+ T cells from human peripheral blood were first separated into live CD4+CD25loCD127hi cells within gate 1 (G1) and live CD4+CD25hiCD127lo cells within gate 2 (G2). Subsequently, CD45RA+GPA33int naïve Tconv cells and CD45RA+GPA33hi naïve tTreg cells were selected within G1 and G2, respectively.

Supplementary Figure 2

Supplementary Figure 2. Representative flow cytometry histograms showing protein expression of total CTLA-4 (*n* = 13), surface CTLA-4 (*n* = 8), surface OX40 (*n* = 8), surface GITR (*n* = 8), surface PD-1 (*n* = 8) and surface CD39 (*n* = 8) on Tconv, tTreg and iTreg cells that were restimulated via CD3/CD28 or not.

Supplementary Figure 3

Shared high (cluster 1) or low (cluster 6) expression in iTreg and tTreg

Clusters 2–3: shared low expression in Tconv and iTreg

Clusters 4–5: unique low expression in iTreg

Clusters 7–8: shared high expression in Tconv and iTreg

Clusters 9–10: unique high expression in iTreg

Supplementary Figure 3. GO biological processes enrichment analyses of indicated clusters of proteins from Figure 3B that were differentially expressed between unstimulated Tconv, tTreg and iTreg cells ($p < 0.05$).

Supplementary Figure 4

Supplementary Figure 4. Analysis of the proteomic responses of tTreg, iTreg and Tconv cells to CD3/CD28-mediated restimulation. **(A)** GSEA enrichment maps of the proteomic responses of tTreg, iTreg or Tconv cells to CD3/CD28 restimulation, showing enriched GO biological processes in yellow (upregulated) and blue (downregulated). Gene sets are depicted as nodes, of which the size represents the number of genes in the gene set. Gene sets sharing a large number of genes are clustered and are connected by lines of which the thickness corresponds with the number of shared genes. Clusters of similar biological processes are annotated using general terms. **(B)** Linear regression analysis of the proteomic response of iTreg cells (x-axis) compared to the responses of either Tconv (red, y-axis) or tTreg (blue, y-axis) cells, revealing different regression lines ($p < 0.0001$). **(C)** GO biological processes enrichment analysis of shared changes in iTreg and Tconv cells, but not tTreg cells, upon CD3/CD28-mediated activation ($p < 0.05$).

Supplementary Figure 5

Supplementary Figure 5. Representative flow cytometry histograms showing protein expression of total NFATC2 ($n = 4$), surface ITGA4 (CD49d) ($n = 5$) and surface ICOS ($n = 5$) on Tconv, tTreg and iTreg cells that were restimulated via CD3/CD28 or not.

Supplementary Figure 6

 Unstimulated

 αCD3/CD28

Supplementary Figure 6. Representative flow cytometry histograms showing protein expression of surface CCR4 ($n = 4$), surface HLA-DR ($n = 4$) and surface FAS (CD95) ($n = 4$) on Tconv, tTreg and iTreg cells that were restimulated via CD3/CD28 or not.