Synthetic periphyton as a model system to understand species dynamics in complex microbial freshwater communities

Olga Lamprecht [a], Bettina Wagner [a], Nicolas Derlon [a] and Ahmed Tlili [a*]

[a] Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

* Corresponding author:

Dr. Ahmed Tlili

Department of Environmental Toxicology (Utox), Swiss Federal Institute of Aquatic Science and Technology (Eawag), Dübendorf, Switzerland.

Überlandstrasse 133

8600 Dübendorf

Switzerland

E-mail address: ahmed.tlili @eawag.ch

Phone: +41 58 765 53 30

Abstract Phototrophic biofilms, also known as periphyton, are microbial freshwater communities that drive crucial ecological processes in streams. Understanding species dynamics in such communities is challenging due to their complexity, yet fundamental to gain insights into the mechanisms of biofilm development and adaptation, such as under conditions of climate change and chemical pollution. To address this challenge, we rationally designed a workflow, based on knowledge about natural periphyton formation. This workflow allowed us to build a fully defined but diverse, stable and highly reproducible synthetic periphyton model system consisting of at least 22 phototrophic species. We demonstrate the ability to monitor microbial dynamics at the species level during periphyton development and how their abundances are impacted by stressors such as increased temperature and a herbicide, singly and in combination. Importantly, by knowing the exact composition of the periphyton we show that, besides specific sensitivities to stressors, species interactions are a major driver for community diversity.

Keywords

Freshwater biofilm, synthetic microbiology, microbial composition, amplicon sequencing, multiple stressors, temperature, pesticides.

Introduction

Periphyton are benthic freshwater communities of high diversity that are dominated by photoautotrophic microorganisms, such as diatoms, green algae and cyanobacteria [1-4]. Periphyton forms the basis of aquatic food-webs by primary production. it releases oxygen and influences the hydrodynamics of the water body [2, 5, 6]. Due to its ecological importance in stream functioning, periphyton is widely used as a biological community model to assess effects of biotic and abiotic factors, such as trophic interactions, man-made chemical pollution or eutrophication [7]. However, most of the data on periphyton are descriptive thus far, focusing on integrative endpoints that basically consider the community as a black box [8]. Such an integrative approach can be useful because it informs on the general community status. It does not, however, reveal mechanisms that govern the structural dynamics and functional properties of natural communities.

Gaining mechanistic understanding of the biological processes in periphyton, such as species interactions or responses to environmental factors, is challenging due to the complexity of the communities [9]. Every periphyton, and natural microbial community in general, is somewhat different owing to multiple and varying environmental factors. Hence, such microbial communities will, to a major part, remain undefined, despite the increasing power to annotate species by, for example, next generation sequencing [10-16]. These limitations have inspired the development of synthetic microbial ecology as a discipline [17-20]. It relies on the establishment of communities that are artificially created by co-culturing selected microbial genotypes while retaining key functionalities and properties of the more complex natural counterparts. These well-defined and relatively simplified consortia of interacting microorganisms can then be used as a model system for hypothesis testing in order to answer key ecological questions.

Synthetic microbial communities have been mainly constructed with bacterial biofilms [21, 22] and to a lesser extent with planktonic freshwater algae [23-25]. These communities were used to examine, for example, species interactions [26-28], biodiversity-productivity relationships [29-33] and effects of environmental stressors [34-41]. Despite the long and extensive history of research on periphyton and its ecological importance, attempts to design and construct a benthic synthetic community are virtually lacking. Here we describe the development of a workflow to establish a reproducible, complex and tenable synthetic periphyton, consisting of multiple freshwater phototrophic species. We applied a bottom-up approach [18] based on rational design and current knowledge on microbial dynamics during periphyton formation.

Relying on several criteria, we first selected twenty-six phototrophic microbial species and
organized them into functional groups based on microbial succession during periphyton
development. Then we defined the starting inoculum for each group, based on the growth rates
of each single species composing the groups. Finally, we experimentally determined the
colonization duration for each group. The established workflow of sequential addition of the
groups to colonize an artificial substrate, covered with an extracellular polymeric substance
(EPS)-producing bacteria, is summarized in Fig. 1. As a proof of principle, we used the
established periphyton to test for the single and combined effects of a herbicide and temperature
change on the community spatial structure, function and microbial composition.

Fig. 1: Schematic representation of the workflow to establish the synthetic periphyton. Twenty-six
phototrophic microbial species were selected based on their ability to grow under the same conditions
and their frequent detection in natural freshwater periphyton. The selected species were then assigned
to three groups (**I**). The cell density in the starting inoculum for each group was determined based on
the growth rates and reached maximum biomass of each single species (**II**). Groups were added
sequentially onto growth chambers covered with the EPS-producing bacteria *Sphingomonas elodea*.
Time points for the addition of the groups and colonization durations for each phototrophic group were
determined experimentally (**III**) (Illustration created with BioRender.com).

Results and Discussion

Species selection and grouping criteria

Following a literature search and screening of available databases that describe freshwater periphyton microbial composition, we selected a total of 34 phototrophic species, consisting of 18 diatoms (Bacillariophyta), 10 green algae (Chlorophyta) and 6 Cyanobacteria (Supplementary Table 1). The selection criteria included detection frequency in freshwater periphyton, cultivable species that are commercially available from algal banks, and a taxonomic composition that covers the different stages of periphyton development. Such composition implies various biological traits, such as nutrient requirements and sensitivity to stressors, and consequently different growth conditions. Therefore, we progressively adapted all the 34 phototrophic species for at least six months to grow in the same liquid medium (i.e., COMBO, Supplementary Table 2) as well as temperature and light conditions (see Methods for more details). Twenty-six out of the 34 photosynthetic species were successfully adapted and thus selected for the subsequent establishment of the synthetic periphyton (Table 1).

The initial cell density of single microbial species and their arrival order to a habitat can strongly influence community assembly and species succession dynamics, thus affecting functionality. Based on the current knowledge about periphyton development and microbial succession dynamics of phototrophs, we distributed the twenty-six selected species into three groups (A, B and C) (Table 1). Group A consists of *A. pyrenaicum*, a known pioneer diatom in periphyton formation, and *C. meneghiniana*, one of the most commonly found freshwater diatoms. These two species belong to the low profile guild (see also Supplementary Table 1), which comprises the most resistant species with regard to nutrient-poor conditions, and therefore dominate at low nutrient levels [42]. Group B consists of 9 high profile diatoms that contribute the most to the 3D structure development of the periphyton [42]. The high profile guild encompasses species of tall stature, filamentous, chain-forming, stalked diatoms and species with comparatively small cells that form long colonies and extend beyond the boundary layer next to the periphyton surface. This allows them to exploit resources that are unavailable to the low profile guild. The remaining 15 species, which compose group C, correspond to the green algae and cyanobacteria as well as the diatoms from the genera *Navicula*, *Nitzschia* and *Sellaphora* that belong to the motile guild. This guild includes relatively fast moving species, which allows them to physically avoid stress and to move towards resource-rich microhabitats within the benthic mat [43-45]. Green algae and cyanobacteria were added to the periphyton in the last group C, since they are most sensitive to nutrient depletion and light attenuation and might

produce unfavorable and harmful toxins to the other added species, respectively. An exception was made for *C. meneghiniana*, which was also included in Group C in addition to group A, as this species is not only low profile but also planktonic and can therefore colonize periphyton at any stage of development. This might explain its high abundance and frequent detection in natural periphyton [42].

Finally, we considered that the first stage of biofilm formation is the attachment of EPS-producing bacteria to the colonization substrate [46, 47]. With the development of genomics techniques, various bacterial strains belonging to the *Sphingomonas* family have been frequently identified as initiators of periphyton formation [48, 49]. We therefore used *Sphingomonas elodea* (ATCC 31461) for surface coating. *S. elodea* is a bacterium known to produce an exopolysaccharide gellan gum with unique colloidal and gelling properties [50]. This bacterium provided a solid and sticky attachment layer for the subsequently introduced phototrophic species, allowing them to bind the surface, multiply and presumably produce their own EPS.

Starting inoculum and colonization duration for each phototrophic group

As growth rates vary among microorganisms, one major challenge to obtain a complex community is to avoid that one or very few species dominate the community. In order to address this challenge, we defined the starting inoculum for each group, using suspension growth experiments that assumed that single species have similar growth in planktonic and benthic modes. Based on the growth rate and reached maximum biomass for each single species (details in Supplementary Tables 3 and 4), we added the least number of cells for species that grew fast and reached high biomasses, and the highest number of cells for species that grew slow and reached low biomasses (as shown in Fig. 1). The total starting cell concentration of all species within a group was 250'000 cells mL^{-1}, 999'000 cells mL^{-1} and 2'175'000 cells mL^{-1} for group A, B and C, respectively (see detailed cell concentrations for each species in Table 1).

Table 1. List of the 26 selected phototrophic species used to establish the periphyton.
Species were assembled into groups A, B and C, according to the order of addition to the community. Growth rate constant k and maximum reached biomass were derived from optical density measurements at 685 nm (Supplementary Table 3). Color code: green – fast growth and high biomass; yellow – fast growth and low/intermediate biomass; red – slow growth and low biomass. To ensure accurate grouping of species, growth rate constant k and maximum reached biomass were additionally derived from CASY cell counter for most of green algae and diatom species (see Supplementary Table 4). *Due to its cell morphology resulting in inaccurate measurements with the spectrophotometer, *N. vermicularis* was included in the intermediate (yellow) group based on the growth rate and maximum biomass measured by CASY cell counter (see Supplementary Table 4).

Group	Species	Phylum	Growth rate constant (k)	Maximum reached biomass	Starting inoculum (cells mL^{-1})
A	*Achnanthidium pyrenaicum*	Bacillariophyta	0.018	0.285	125 000
	Cyclotella meneghiniana	Bacillariophyta	0.023	0.194	125 000
B	*Gomphonema parvulum*	Bacillariophyta	0.011	0.231	75 000
	Gomphonema clavatum	Bacillariophyta	0.014	0.211	75 000
	Synedra sp.	Bacillariophyta	0.014	0.195	75 000
	Fragilaria crotonensis	Bacillariophyta	0.016	0.059	125 000
	Fragilaria capucina	Bacillariophyta	0.033	0.048	125 000
	Melosira sp.	Bacillariophyta	0.018	0.051	125 000
	Cymbella cistula	Bacillariophyta	0.018	0.034	133 000
	Tabellaria sp.	Bacillariophyta	0.007	0.142	133 000
	Ulnaria ulna	Bacillariophyta	0.01	0.081	133 000
C	*Scenedesmus acuminatus*	Chlorophyta	0.019	0.797	65 000
	Chlorella vulgaris	Chlorophyta	0.009	0.735	65 000
	Pediastrum boryanum	Chlorophyta	0.011	0.393	65 000
	Merismopedia glauca	Cyanobacteria	0.009	0.747	65 000
	Cyclotella meneghiniana	Bacillariophyta	0.018	0.285	125 000
	Nitzschia palea	Bacillariophyta	0.025	0.197	125 000
	Nitzschia vermicularis	Bacillariophyta	0.012	0.023*	125 000
	Craticula accomoda	Bacillariophyta	0.019	0.191	125 000
	Sellaphora minima	Bacillariophyta	0.025	0.198	125 000
	Scenedesmus vacuolatus	Chlorophyta	0.017	0.536	125 000
	Pediastrum duplex	Chlorophyta	0.021	0.660	125 000
	Chamaesiphon polonicus	Cyanobacteria	0.007	0.526	125 000
	Phormidium sp.	Cyanobacteria	0.022	0.295	125 000
	Pseudanabaena galeata	Cyanobacteria	0.020	0.520	190 000
	Botryococcus braunii	Chlorophyta	0.005	0.417	300 000
	Craticula cuspidata	Bacillariophyta	0.007	0.066	300 000

After determining the starting inoculum, we experimentally tested and set the optimal colonization duration for each of the formed groups. For this purpose, we monitored over time the increase of surface-attached cell number and photosynthetic efficiency as proxies for benthic biomass and community physiology, respectively. As well we measured surface colonization dynamics based on microscopy imaging. Adequate colonization time was especially important for group A in order to allow the low profile diatoms to colonize the EPS-covered surface but not to overgrowth the colonization substrate. Our results show that when grown separately, *C. meneghiniana* and *A. pyrenaicum* reached similar number of surface-attached cells (Fig. 2A) and the highest photosynthetic yield (Supplementary Fig. 1A) after 4 days of benthic growth. What is more, when both species were grown together at a one to one ratio for 4 days, the large majority of cells (approximately 6 to 7 times more cells than in the supernatant) were attached to the substrate surface (see Fig 2B and 2C). They also covered 9.6 ± 1.4% of the surface (Supplementary Fig. 2), leaving sufficient surface for colonization by the subsequently added groups B and C. Based on these results, we selected 4 days as the optimal colonization duration for group A.

To determine the colonization duration for groups B and C, we followed a slightly different strategy. Since group B species were added on top of group A after 4 days, we monitored over time the number of cells attached to the substrate and in the supernatant, as well as photosynthetic efficiency for the group AB together, thus accounting for potential species interaction effects. Our data show that 14 days after group B was added to group A, most of the cells were attached (Fig. 2B) and covered 47.4 ± 3.7 % of the surface (Supplementary Fig 3). Longer incubation, on the other hand, led to an increase of detached cells (Fig. 2B) and a decrease in the photosynthetic efficiency (Supplementary Fig 1B). Such results indicate that the periphyton formed by group AB has reached its maturity and that 14 days is the optimal colonization duration for group B, when added on top of group A. Subsequently, at 18 days (A 4 days + B 14 days) of colonization, we introduced group C. Twelve days after group C was added, most of the cells were attached to the colonization surface (Fig. 2C) with a coverage area of 91.8 ± 1.9 % (Supplementary Fig. 3) and relatively high photosynthetic efficiency (Supplementary Fig. 1C), indicating that we obtained a stable benthic community after a total colonization duration of 30 days. Such a duration is usually considered optimal for natural periphyton to reach its maturity when growing on artificial and natural substrates [4, 51] and has been applied in most of the field and microcosm studies focusing on periphyton (e.g., [52-56]).

Fig. 2: Increase of the benthic biomass during periphyton establishment.
The concentrations (means ± SD; n = 3) of attached and non-attached cells (supernatant) were quantified via optical density (OD) at 685 nm and cell count with a CASY cell counter throughout the periphyton development; separately for the two species of group A **(A)**, then for groups A and AB (upon addition of group B onto group A at 4 days of growth) **(B)** and for groups A, AB and ABC (upon addition of group C onto group AB at 18 days of growth) **(C)**. Red vertical arrows indicate selected time points when group B species were added on top of group A (t_1), group C on top of group AB (t_2) and periphyton was established (t_3). Cell concentrations in the supernatant for *C. meneghiniana* and *A. pyrenaicum* when grown separately **(A)** were below the quantification limit; hence no graphics for supernatant is shown.

3D structural changes during periphyton establishment

The spatial organization of periphyton is of importance as it provides microenvironments and niches for specific microorganisms as well as allows species that are located in close proximity to interact [57]. Here we used optical coherence tomography (OCT), which enables image acquisition at the meso-scale in the range of millimeters (mm) [58, 59] (Fig. 3A) and provides details of overall thickness, surface topology (relative roughness) and internal porosity changes during biofilm formation [60, 61]. At four days (t_1) of colonization by group A, the periphyton reached an average thickness of 101 ± 47 µm and a relative roughness of 0.4 (Fig. 3B). After species from group B were introduced and grown together with species from group A for another 14 days (t_2), the average periphyton thickness slightly decreased while the relative roughness and internal porosity increased. This suggests that high profile diatoms (group B) can modify the overall periphyton architecture from a rather homogeneous structure (Ra' < 0.5) formed by group A species towards a heterogeneous one with a higher surface roughness (Ra' > 0.5). Finally, following the addition of group C and after 30 days (t_3) of growth, the established periphyton was almost 200 µm thick and highly dense, as indicated by the sharp thickness increase and internal porosity decrease, respectively. A potential explanation of the increased structure density could be the presence of motile diatoms and small cyanobacteria in the last added group that fill the porous space between larger and filamentous cells within the periphyton matrix. Overall, these results provide additional detail on how physical structure parameters change during periphyton establishment and on the contribution of each phototrophic group to these changes.

Fig. 3. Changes of the periphyton physical structure during its establishment. Shown are representative OCT images obtained over 30 days of periphyton growth at time points t_1 (A 4 days), t_2 (AB 18 days) and t_3 (ABC 30 days) **(A)**. White pixel: Biomass accumulation, black pixel: absent/low biomass accumulation. Scale bar: 200 µm. Mean thickness, surface topology (relative roughness) and internal porosity (mean to max ratio of z, where z is the height of the biofilm) were determined from 20-30 images and 5 biological replicates per time point **(B)**. Error bars represent standard deviations. * $P<0.05$, ** $P<0.005$ *** $P<0.0001$, post hoc Tukey test.

Microbial diversity and composition during periphyton establishment

Analyses via confocal laser scanning microscopy (CLSM) of the intact periphyton at 30 days showed the presence of 12 to 15 different species that are co-existing in close proximity within the community (Supplementary Fig. 4). CLSM is a powerful technique to examine three-dimensional structure of the periphyton at the micrometer scale [62-64]. However, it does not allow precise identification of species with similar cellular morphologies (e.g., small and round or long filaments) or provide quantitative information on the relative abundance of each genotype within the community. To overcome these limitations, we performed high-throughput amplicon sequencing of 18S rRNA (green algae and diatom), *rbcL* (higher taxonomic resolution for diatoms) and 16S rRNA (cyanobacteria) genes (see Methods and Supporting Information for more details). For this we created our own reference database for the twenty-six single

species that were used to establish the synthetic periphyton with their respective amplicon gene sequences (Supplementary Table 5). We also validated the accuracy of the amplicon sequencing approach for single species identification and quantification by using a mock community (see Supplementary Discussion and Supplementary Table 6 for more details). Among the 26 species used to establish the periphyton, we were able to accurately identify 21 eukaryotic (diatoms and green algae) and 4 prokaryotic (cyanobacteria) species, with the only limitation to distinguish between *Pediastrum duplex* from *Pediastrum boryanum*.

Once the robustness of our approach was validated and single species were taxonomically assigned, we monitored community composition in the mature synthetic periphyton as well as changes over time during periphyton formation. In comparison to morphological analysis by CLSM, amplicon sequencing provided higher precision for single species identification. Indeed, characterization of the community composition revealed a diverse and complex mature periphyton composed of at least 22 different phototrophic species, corresponding to 13 diatoms, at least 5 green algae and 4 cyanobacteria, with various relative abundances (Fig. 4 A-C). For instance, *C. meneghiniana, A. pyrenaicum, S. acuminatus* and *S. vacuolatus* were the most abundant in the established periphyton at t_3, representing approximately 75% of the final community (Fig. 4A). Moreover, all four cyanobacteria species that were introduced into the community were also detected in the mature periphyton (Fig. 4C and Supplementary Fig. 6D). The highest abundances were measured for *C. polonicus* and *M. glauca*, and the lowest one for *Phormidium sp*. These findings are in line with previous studies that identified these phototrophic species in either natural *in-situ* periphyton or periphyton grown in microcosms [53, 65, 66], demonstrating the ecological relevance of our synthetic microbial community.

Our results also show that the development of periphyton follows a succession of changes that are mainly governed by species interactions. For instance while, as expected, only *C. meneghiniana* and *A. pyrenaicum* were detected at t_1 (Fig. 4A), we measured a significantly higher abundance of *C. meneghiniana* than *A. pyrenaicum* after 4 days (t_1) of benthic growth and also at 18 days (t_2) after addition of the nine high profile diatom species from group B (Supplementary Fig. 6A and 6E). The abundance of *C. meneghiniana*, on the other hand, decreased significantly at day 30 (t_3) in the mature periphyton (Fig. 4A and 4B, Supplementary Fig. 6A and 6E), suggesting that the presence and/or the activity of group C species may have suppressed its growth. Furthermore, *Tabellaria sp.* and *Melosira sp.* from group B, as well as *N. vermicularis* introduced with group C, were no longer detected in the mature periphyton, neither with 18S (Fig. 4A, Supplementary Fig. 6B and 6C, respectively) nor with *rbcL* (Fig.

4B, Supplementary Fig. 6F and 6G, respectively), probably as a result of species competition.

Therefore, from twenty-six sequentially introduced phototrophic species, three species were no

longer detected in the mature periphyton.

Fig. 4. Genetic composition of the synthetic periphyton at the species level. Community
composition profile and relative abundances (average; n = 9) were inferred from taxonomy-
based clustering at species level of assigned 18S rRNA **(A)** and *rbcL* **(B)** gene sequences and
at genus level of assigned 16S rRNA **(C)** at 4 (t_1), 18 (t_2) and 30 days (t_3) during periphyton
formation. NA: not assigned species.

Once the workflow to establish the synthetic periphyton was set, we tested its reproducibility.

For this purpose, we performed two independent experiments, with two months gap in between,

each time starting with planktonic cultures of the single species and following the established

workflow (summarized in Fig. 1). When comparing the experiments, our data on genetic

diversity shows the reproducibility of the obtained community (Supplementary Fig. 7). Despite

slight differences in relative abundances, community composition of the mature periphyton at

day 30 was well conserved. Most importantly, species succession during periphyton

establishment was also similar between both experiments and among biological replicates (n =

9). Taken together, these results demonstrate that by following a rational design and applying

existing knowledge on microbial dynamics during periphyton formation we were able to obtain

a reproducible and diverse synthetic community.

Case study: single and combined effects of the herbicide terbuthylazine and increased
temperature on periphyton structure and function

Understanding how the individual and combined effects of multiple environmental stressors
impact species interactions is a key challenge. In complex microbial communities, such as
periphyton, species have a large degree of freedom for interactions that further complicates the
evaluation of the outcomes. By using the established synthetic periphyton as a model system,
we aimed here to examine precisely how single species within the community and the
community as a whole respond during periphyton development to the single and combined
exposures to two major environmental stressors: the widely-used herbicide terbuthylazine at 0,
1, 10 and 100 nM and a change in temperature from 17°C to 20°C (see Methods for more details
on the experimental design).

Our results on community composition during periphyton establishment provide strong
evidence that increased temperature significantly impacted microbial diversity and altered
species succession in the community (see Supplementary Table 12 for statistical analysis).
When comparing between community composition profiles in the mature periphyton at 17°C
and 20°C at the phylum level, the average abundances of green algae (Chlorophyta) and
cyanobacteria decreased from 61% to 25% and from 65% to 49%, respectively, while the one
for diatoms (Bacillariophyta) increased from 37% to 74% (Supplementary Fig. 8A and 8B). A
detailed analyses at the single species level shows that the abundances of the green algae *C.*
vulgaris, *S. acuminatus* and *S. vacuolatus*, which dominated the community at 17°C, strongly
decreased with increased temperature (Fig. 5C and Supplementary Fig. 8C). Among
cyanobacteria, only the abundance of *C. polonicus* was negatively regulated by the elevated
temperature (Fig. 5D and Supplementary Fig. 8E). In sharp contrast, several diatoms, such as
C. meneghiniana, C. cistulla, Tabellaria sp. and *Melosira sp.*, were significantly favored by the
temperature increase (Fig. 5A and 5B and Supplementary Fig. 8C and 8D). Interestingly, *C.*
meneghiniana and *Melosira* sp. were not impacted by the temperature increase at t_2 (Fig. 5A
and 5B, respectively). On the other hand, at t_3 upon addition of group C species, their
abundances significantly decreased at 17 °C but increased at 20 °C (Fig. 5A and 5B,
Supplementary Fig. 8C and 8D, and Supplementary Table 12). Such results suggest that
temperature-sensitive species from the group C potentially balance the growth of *C.*
meneghiniana and *Melosira sp.* at 17 °C. However, when their own growth is reduced at 20 °C,
they can no longer compete, allowing *C. meneghiniana* and *Melosira sp.* both species to grow
within the community regardless of their specific tolerance to temperature.

Fig. 5. Abundance of single species during periphyton establishment. Shown is abundance
of group A (**A**), group B (**B**) and group C (**C**) eukaryotic species as well as group C prokaryotic
species (**D**) measured at t_1 (A 4 days), t_2 (AB 18 days) and t_3 (ABC 30 days) and grown under
different conditions. The conditions correspond to 4 levels of the herbicide terbuthylazine (0,
1, 10 and 100 nM) and 2 different temperatures (17 and 20 °C). Horizontal lines in the black
boxes correspond to the average values and the lower and higher limits of the standard deviation
(n = 3). Results of the pairwise comparison (post hoc Tukey test) among all treatments for each
species are shown in Supplementary Tables 13-15.

Exposure to increasing concentrations of terbuthylazine also led to changes in the microbial composition of the mature periphyton (Fig. 5 and Supplementary Fig. 8). The data on single species abundances shows that the large majority of green algae and some diatom species were negatively affected only by the highest terbuthylazine concentration (Fig. 5 and Supplementary Fig. 8). On the contrary, the relative abundance of the low profile diatom C. *meneghiniana* (Fig. 5A), green algae *Pediastrum sp.* (Fig. 5C) and the cyanobacterium *Phormidium sp.* (Fig. 5D) increased significantly (Supplementary Table 12) upon exposure to 100 nM terbuthylazine. This reflects their competitive advantage under unfavorable conditions for most of the other species within the community. Such findings are in line with those of previous studies that reported the presence of these species in natural periphyton that had been exposed to micropollutants such as herbicides [14, 53, 67, 68], indicating their ability to tolerate high levels of pollutants.

As conceptualized by the subsidy-stress concept, environmental factors, such as increased temperature, can be beneficial for microorganisms and consequently mitigate the negative effects of micropollutants [69, 70]. This is in line with our results on the relative abundances of single species which show significant beneficial interaction between temperature and terbuthylazine (Supplementary Table 12). Indeed, the negative effects of the highest concentration of terbuthylazine in the mature periphyton at 17°C were no longer observed at 20°C for several diatoms (e.g., A. *pyrenaicum* at t_1, *Synedra sp.* at t_3 and G. *parvulum* at t_3,) and green algae (e.g., S. *vacuolatus*, S. *acuminatus* and C. *vulagaris* at t_3) (Fig. 5). Interestingly, some of the high profile diatom species such as *Synedra sp.*, *Tabellaria sp.* and *C. cistula*, were not at all affected when exposed to terbuthylazine or temperature alone (Supplementary Table 12). Clearly, such results confirm that multiple interactions among species and temperature occur, circumstances that complicate the assessment of risks caused by chemicals in complex field situations. These can now be assessed and better understood under controlled conditions with our synthetic periphyton model.

One additional insight from our case study is that both increased temperature and exposure to the herbicide terbuthylazine affected negatively the 3D-structure and biomass but not the photosynthetic efficiency of the periphyton. Despite the observed growth during periphyton development in all treatments, the overall periphyton 3D-structure (Supplementary Fig. 9), thickness (Supplementary Fig. 10A) and concentration of attached cells to the colonization substrate (Supplementary Fig. 11A) were significantly lower at 20°C compared to 17°C. Such results can be explained by the sharp decrease in the relative abundances of most of the green

algae and diatom species composing group C upon increased temperature or exposure to terbuthylazine (Fig. 5, Supplementary Fig. 8). This conclusion is also supported by the fact that these negative effects were mainly observed after addition of group C species, confirming their important contribution to the benthic biomass and structural development of the periphyton. Despite the clear effects on benthic biomass and physical structure, but also microbial diversity and composition, exposure to terbuthylazine and increased temperature did not lead to notable effects at the functional level (Supplementary Fig. 12). Photosynthetic efficiency was similar in all treatments and time points, except for periphyton exposed to 100 nM terbuthylazine at 18 days (i.e., before addition of group C). Overall, these results from our synthetic periphyton confirm previous observations [37, 39, 41] about functional redundancy with regard to photosynthetic efficiency when less sensitive species of the same functional group compensate for the loss of other species in order to ensure important community functions [71].

Conclusion

In order to answer important ecological questions about how species interactions can translate into complex community-level processes and properties, we developed a workflow for a reproducible establishment of a synthetic freshwater periphyton. While the majority of previously established synthetic microbial communities consist of only two to four genotypes, our synthetic periphyton consists of at least 22 different microbial phototrophic species, growing and interacting in close proximity. Our results demonstrate that our protocol and established synthetic biofilm enables to overcome challenges encountered with complex natural periphyton while maintaining crucial ecological features, such as functional redundancy and varying species sensitivity to stressors. Species-level resolution within the community provided an accurate understanding of how species interactions, in addition to specific sensitivities to a given stressor, modulate the community properties at the functional and structural level. Although our focus was on phototrophs, future studies could also consider other groups of organisms, e.g. heterotrophic bacteria or (micro)grazers, in order to examine, for example, trophic networking and species interactions. Overall, the synthetic periphyton that we established offers great opportunities for theory testing through targeted manipulation and for exploring crucial ecological processes, such as diversity-function relationships or resistance and resilience to environmental stressors.

Methods

2.1.   Cultures of the phototrophic species

Non axenic planktonic cultures of the 34 selected single phototrophic species were obtained from (1) SAG Culture collection of algae (Goettingen university, Germany), (2) CCAC Culture collection (University of Cologne, Germany) and (3) TCC Thonon culture collection (INRAE, France) (Supplementary Table 1). Before starting the experiments, all species were adapted for six months to grow in the same freshwater medium COMBO (see Supplementary Table 2 for the detailed composition) [72] at 17°C and a light intensity of 40 to 60 µmol s^{-1} (Philips Master LED tube Value 1200 mm InstantFit 865, 16.5 W, MLT Moderne Licht-Technik AG, Wettingen, Switzerland) with a photoperiod of 16h light: 8h dark. Twenty-six out of the 34 species were successfully adapted and used for the establishment of the synthetic periphyton. All main planktonic cultures were kept in triplicate in sterile 300-mL Erlenmeyer flasks containing 150 mL of COMBO and placed on a shaker operated at 30 rpm. Every 3 weeks, 50 mL of these cultures were centrifuged at 2000 g, the supernatant removed and cells in the pellet resuspended in 1 mL fresh COMBO. 200 to 500 µl of this suspension were added to 75 mL fresh COMBO for growth measurements and 500 to 1000 µl were added to 150 mL of fresh COMBO and kept as backup cultures.

Depending on the species morphology, growth rates and maximum reached biomass of each single species were generated from optical density and/or cell count measurements of the planktonic cultures over time. The flasks were incubated in the same conditions as described above and 2 mL were taken every 48 h for the growth measurements. Optical density of the suspensions at 685 nm was measured with a spectrophotometer (UVIKON 930, Kontron Instruments, Basel, Switzerland). Cell concentration was determined by using an automated CASY cell counter Model TT (Roche Innovatis AG) with 60 µm capillary, 2.18 µm - 30 µm evaluation cursor and 1.5 – 29.83 µm normalization cursor to take into consideration the different cell sizes. The obtained data were fitted in a logistic model as function of time in order to derive the growth rate constant (*k*) and maximum reached biomass for each species (detailed results are provided in the Supplementary Tables 3 and 4).

2.2.   Experimental system and design for the establishment of the synthetic periphyton

The establishment of periphyton in our study was performed under static conditions in sterile 1-well Nunc^{TM} Lab-Tek^{TM} II Chamber Slides (74 x 25 x height mm, Thermo Scientific), which consists of a removable polystyrene media chamber attached to a standard glass slide. First, the bacteria *Sphingomonas elodea* were inoculated from the glycerol stock into a sterile Erlenmeyer flasks containing 100 mL ATCC rich broth medium (8.0 g Nutrient Broth (BD cat 234000) in

1 L deionized water). After 48 h of growth at 200 rpm and 30 °C, optical density of the bacterial

culture was measured at 600 nm. Chambers were then inoculated with 4 mL *Sphingomonas*

elodea in ATCC rich broth medium with an OD_{600} of 0.4 and incubated at room temperature

under static conditions for bacteria to colonize the surface and produce the EPS matrix. Three

days after inoculation, the medium was removed and chambers gently washed once with

COMBO before adding a mixture of the two group A species at a one to one ratio in 4 mL

COMBO. After 4 days of incubation, medium with non-attached cells was removed carefully

and 4 mL of COMBO containing a mixture of the nine species that compose group B were

introduced to the chambers. After another 14 days, the medium with non-attached cells was

again removed before addition of the 18 species of group C in 4 mL COMBO (see Table 1 and

Supplementary Table 2 for details). Finally, the chambers were incubated for another 12 days,

leading to a total duration of 30 days for periphyton establishment, starting from the

introduction of group A. In order to avoid potential effects of nutrient limitation, 2 mL of

COMBO were carefully replaced in the chambers every second day. All incubations were

performed in a temperature-regulated room at the same temperature and light conditions to

which all single species have been adapted to (i.e., 17°C and a light intensity of 40 to 60 µmol

s^{-1}).

2.3.   Periphyton characterization

2.3.1. Surface colonization area

Surface coverage of the different phototrophic species during periphyton establishment was

checked microscopically at low magnification on intact biofilms. Imaging was performed with

an inverted Leica microscope (Plan Neo Fluar Z 1 ×/0.25, FWD 56 mm objective) equipped

with a CCD camera (Axiocam 506 mono, Zeiss). Twenty-seven 9 × 3 tile acquisitions were

acquired, montaged and stitched together, covering large area of the chamber surface (10 ×

magnification, 1.2 µm pixel^{-1} resolution). A montage of images and analysis were performed

by using custom-written macro in ImageJ (version 1.53c). In brief, after thresholding with

"Huang dark" method, attached particles were analyzed and counted. An average value from

microscopy of three to five chambers was calculated per time point and condition.

2.3.2. Cell density and cell number

To quantify density or number of cells within the periphyton (attached to the surface) and in

suspension (supernatant) during periphyton establishment, supernatant was collected and the

attached cells scratched from the glass surface of the chambers in 4 mL fresh COMBO. Cell

suspensions of attached cells were diluted 1:20 and 1:400 for OD_{685} and CASY cell count

measurements, respectively. Three to five chambers (biological replicates) were retrieved per

time point and condition. A minimum of three measurements were performed per replicate.

2.3.3. Photosynthetic efficiency measurement

Photosynthetic efficiency was assessed by using an Imaging-PAM (pulse amplitude-

modulated) fluorimeter (Heinz Walz GmbH, Germany). Chlorophyll-a fluorescence from each

periphyton was measured at 665 nm after applying a single light saturation pulse to calculate

the effective quantum yield (ϕ') as:

488
$$\phi' = \frac{F'm - F'0}{F'm} \qquad Eq(3)$$

where F'_m is the maximum fluorescence after the saturation pulse and $F'0$ is the steady-state

fluorescence.

2.3.4. 3D-structure of periphyton

2.3.4.1. Optical Coherence Tomography (OCT)

Optical Coherence Tomography (model 930 nm Spectral Domain, Thorlabs GmbH, Dachau,

Germany) with a central light source wavelength of 930 nm was used to investigate the meso-

scale structure of periphyton. The use of long wavelength light allows to penetrate up to a depth

of 1.9 mm with axial and lateral resolutions of 4.4 µm and 15 µm, respectively. 20-30 scans

(XZ pane image) of 2x1 mm were acquired per replicate at t_1 (at day 4, just before addition of

group B species), t_2 (at day 18, just before addition of group C species) and t_3 (at day 30, the

end of periphyton establishment) during periphyton development. Three replicate samples per

timepoint (and condition) were examined. Image analysis software developed under Matlab®

(MathWorks, Natick, US) was used to analyze OCT image. Image analysis consisted of the

following steps: (1) surface-biofilm interface detection (grey-scale gradient analysis); (2) image

binarization (automatic threshold selection); (3) calculating physical properties of the biofilm.

Biofilm thickness was calculated based on the number of pixels found between the top edge of

the periphyton and the upper chamber surface of each OCT image, using a pixel scaling factor

to obtain the biofilm thickness in microns.

507
$$L_F = \frac{1}{N}\sum_{i=1}^{N} L_{F,i} \qquad Eq(1)$$

508   where $L_{F,i}$ (µm) is the biofilm thickness from a single A-scan (i.e., light reflected from each

509   optical interface) in the corresponding B-scan (i.e., cross-sectional reconstruction of a plane

510   from a series of A-scans across the structure) and N is the total number of A-scans.

511   The relative roughness coefficient (-) of the biofilm was calculated according to [73].

$$R_a^* = \frac{1}{N} \sum_{i=1}^{N} \frac{\left| L_{F,\ i} - \overline{L_F} \right|}{\overline{L_F}} \qquad Eq(2)$$

where *i* represents an A-scan and the overall number of A-scans. Biofilms with a smooth surface

and only a few variations from the mean biofilm thickness have low values close to 0. The

higher the roughness coefficient, the more variations are expected from the biofilm surface [74].

The roughness coefficient is based on the individual thickness measurement $L_{F,i}$, the sample

mean thickness L_F, and the number of thickness measurements N.

OCT images does provide some indication of how dense or loose a biofilm is structured, and

fraction of light/dark pixels or greyscale distribution can be quantified. Wagner *et al.* [75]

suggest zones of low signal intensity represent areas of absent biomass (i.e., voids, pores). The

distinction between solid materials and voids relies on the assumption that absence of OCT

signal equals voids. However, such an assumption and the resulting porosity quantification is

debatable.

2.3.4.2. Confocal laser scanning microscopy

At day 30, confocal laser scanning microscopy (CLSM) was used to visualize the periphyton

structure at the µm-resolution and to detect single species within a dense 3D-structure. For this

purpose, polystyrene media chambers of 1-well Nunc^{TM} Lab-Tek^{TM} II Chambers Slides were

removed with special adaptors and glass slides placed into a petri dish with 25 mL fresh

COMBO medium. Samples were examined with Leica SP5 confocal laser scanning microscopy

(Heidelberg GmbH, Mannheim, Germany). Algae and cyanobacteria were detected through the

autofluorescence of chlorophyll-a signal (excitation 630 nm, emission 650-720 nm). All

observations were performed using a 40×/0.75 NA objective (Leica, Plan-Apochromat®).

Periphytons were scanned at 2-µm intervals, starting at the glass surface, resulting in a series of

individual images at different depths (*z*-scan). CLSM data was analysed by Imaris 9.3.0. Three

replicate samples were examined per condition. For image analysis, three 3D-images per

replicate were recorded, producing nine 3D-images per periphyton sample.

2.3.5   Next generation sequencing for microbial community composition

2.3.5.1 DNA extraction, library preparation and sequencing

In order to compare the composition of prokaryotes (i.e. cyanobacteria) and eukaryotes (i.e. diatoms and green algae) in periphyton, total genomic DNA was extracted from an aliquot of 1 mL from each periphyton suspension at t_1 (4 days), t_2 (18 days) and t_3 (30 days) during the experiment. The samples were centrifuged at 14'000 g for 30 min at 4 °C and the pellets stored at -20 °C until their analyses. DNA extraction was performed by using the DNeasy Plant Mini Kit (Qiagen) following the manufacturer's instructions. Prior to DNA extraction, physical disruption of cells was performed using cooling BeadRuptor Elite (OMNI International, US) and bead tubes from the DNeasy PowerBiofilm Kit (0.5 mm glass, 0.1 mm glass and 2.8 mm ceramic beads). Total DNA was then quantified with a Qubit (1.0) fluorimeter following the recommended protocol for the dsDNA HS Assay (Life Technologies, Carlsbad, CA, USA).

Library construction consisted in a two-step PCR process (see Supplementary Methods for more details). The first PCR amplified the 16S rRNA from cyanobacteria, V4-V5 region of the 18S rRNA gene for eukaryotes and *rbcL* gene using two different primer sets with overhang adapters from [76], [77] and [78], respectively (Supplementary Table 8). The first PCR was performed in triplicate for each DNA sample, negative PCR controls, as well as positive PCR controls for 16S rRNA, 18S rRNA and *rbcL*, consisting of mock communities (Supplementary Tables 5, 9-11). The second PCR, consisting in a limited-cycle amplification, was carried out to add multiplexing indices and Illumina sequencing adapters. The libraries were then normalized and pooled to achieve a 3.2 nM concentration. Paired end (2×300 nt) sequencing was performed on an Illumina MiSeq (MiSeq Reagent kit v3, 300 cycles) following the manufacture's run protocols (Illumina, Inc.). The MiSeq Control Software Version 2.2, including MiSeq Reporter 2.2, was used for the primary analysis and the de-multiplexing of the raw reads. All raw sequences are available at the National Center for Biotechnology Information (NCBI) under the SRA accession ID PRJNA764864.

2.3.5.2 Sequencing data processing, amplicon sequence variants binning and taxonomic assignment

The reads were checked for quality and end-trimmed by using FastQC v0.11.2 [79] and seqtk (https://github.com/lh3/seqtk), respectively. For 16S rRNA and *rbcL*, the reads were merged using FLASH v1.2.11 (minimum and maximum overlap of 15 and 300 bp, respectively; maximum mismatch density of 0.25) [80] while only reads obtained with the forward primer were considered for 18S rRNA. The primers were trimmed by using cutadapt v1.12 (wildcards

allowed; full-length overlap; error rate 0.01) [81]. Quality filtering was performed with PRINSEQ-lite v0.20.4 (minimum quality mean 20; no ambiguous nucleotides; dust low-complexity filter with a threshold of 30) with a subsequent size and GC selection step (size selection range 330–600 bp; GC selection range 30–70%) [82]. The reads were processed with an Amplicon Sequencing Variants (ASV) analysis [83]. The sample reads were first denoised into ASVs with UNOISE3 in the USEARCH software v.11.0.667. The final predicted taxonomic ASV assignments were performed with Mega6 based on our own sequence database (see 2.3.5.3. and Supplementary Discussion) by applying a maximum likelihood phylogenetic tree using nearest-neighbor interchange (Supplementary Table 6). The total reads obtained at each step of bioinformatic filtration are reported in Supplementary Tables 9-11.

2.3.5.3. Generation of single species DNA barcodes database

In order to accurately assign species in a sample we created our own DNA barcode reference database containing all twenty-six single species that were used to establish the synthetic periphyton and their respective barcodes (Supplementary Table 6). For this purpose DNA from single species grown in planktonic mode was extracted as described in 2.3.5.1. The corresponding genes (18S rRNA, *rbcL*, 16S rRNA genes) were PCR amplified form the extracted DNA using primers and conditions listed in Supplementary Table 7. PCR purified fragments were sent for Sanger sequencing. Sequences were quality trimmed using SnapGene 5.3.2 and initially aligned using ClustalW as implemented in Mega 6. Sequences were deposited in GenBank under accession numbers OL304135-OL304156 (18S rRNA), OK669058-OK339073 (*rbcL*) and OK737797-OK737800 (16S rRNA).

2.4   *Case study: assessing the single and combined effects of the herbicide terbuthylazine and increased temperature on the synthetic periphyton structure and function*

Terbuthylazine is a widely used photosystem II inhibitor herbicide, with environmental concentrations that can vary from 2 to 600 ng/L (corresponding to 0.08 - 2.74 nM) in Swiss streams that are impacted by wastewater effluents or agriculture [52, 84] [85, 86]. In the context of global warming, increasing temperature is considered as one of the most critical threats to biodiversity preservation and ecosystem functioning. What is more, temperature can interact with micropollutants such as herbicides which complicates the prediction of potential outcomes. In this context and as a proof of principle, we experimentally tested the single and combined effects of these two stressors based on a full factorial replicated (n = 3) design. Synthetic periphyton communities were established, as described previously, in the presence of four levels of terbuthylazine (i.e., 0, 1, 10 and 100 nM) and under two temperatures (17°C and

20°C). During the 30 days of periphyton establishment, 2 mL COMBO, supplemented or not

with terbuthylazine, were exchanged every two days, to avoid nutrient limitation and maintain

a constant exposure to the herbicide. Periphyton was sampled at days 4, 18 and 30 to measure

benthic biomass, 3D-structure, photosynthetic yield and species composition, as described

above.

2.5　Data analysis

GraphPad Prism 9 was used for statistical tests and visualization of data. Significant differences

in phototrophic biomass, quantum yield, OCT and taxonomic abundance were assessed by one-

way ANOVA during the periphyton establishment and two-way ANOVA in the case study to

test for the interactions of the herbicide and temperature. When effects were significant, the

ANOVAs were followed by Tukey's post hoc tests. Normality and homogeneity of variance

were checked prior to the ANOVA analysis (Kolmogorov-Smirnov's and Levene's tests,

respectively). Data that were not normally distributed were log transformed. The significance

level for all tests was set at $\alpha = 0.05$.

Sequencing data analyses were performed with the R package Phyloseq version 1.32.0

(McMurdie and Holmes, 2013). After rarefaction, 16S, 18S rRNA and *rbcL* datasets were

composed of samples containing the same number of reads (17209, 26097 and 2652 reads,

respectively). Graphical representations were generated with the R package "ggplot2" or

GraphPad Prism9.

References

1.　Battin, T.J., et al., *The ecology and biogeochemistry of stream biofilms.* Nature Reviews
Microbiology, 2016. **14**(4): p. 251-263.
2.　Battin, T.J., et al., *Contributions of microbial biofilms to ecosystem processes in stream*
mesocosms. Nature, 2003. **426**(6965): p. 439-442.
3.　Vadeboncoeur, Y., et al., *Substratum as a driver of variation in periphyton chlorophyll and*
productivity in lakes. Journal of the North American Benthological Society, 2006. **25**(2): p.
379-392.
4.　Wetzel, R.G., *Periphyton of freshwater ecosystems : proceedings of the First International*
Workshop on Periphyton of Freshwater Ecosystems, held in Växjö, Sweden, 14-17 September
1982. Developments in hydrobiology. 1983, The Hague ; Boston

Hingham, MA: Junk ;

Distributors for the U.S. and Canada, Kluwer Boston. ix, 346 p.

5.   Haack, T.K. and G.A. Mcfeters, *Nutritional Relationships among Microorganisms in an*
639   *Epilithic Biofilm Community.* Microbial Ecology, 1982. **8**(2): p. 115-126.

640   6.   Romani, A.M., et al., *Biofilm structure and function and possible implications for riverine DOC*
641   *dynamics.* Microbial Ecology, 2004. **47**(4): p. 316-328.

642   7.   Montuelle, B., et al., *The periphyton as a multimetric bioindicator to assess the impact of land*
643   *use on river: an overview on the Ardières-Morcille experimental watershed (France).*
644   Hydribiologia, 2010. **657**: p. 123-147.

645   8.   Gubelit, Y.I. and H.-P. Grossart, *New Methods, New Concepts: What Can Be Applied to*
646   *Freshwater Periphyton?* Frontiers in Microbiology, 2020. **11**(1275).

647   9.   Gessner, M.O. and A. Tlili, *Fostering integration of freshwater ecology with ecotoxicology.*
648   Freshwater Biology, 2016. **61**: p. 1991-2001.

649   10.   Tardy, V., et al., *A pilot experiment to assess the efficiency of pharmaceutical plant*
650   *wastewater treatment and the decreasing effluent toxicity to periphytic biofilms.* Journal of
651   Hazardous Materials, 2021. **411**: p. 125121.

652   11.   Romero, F., et al., *Desiccation events change the microbial response to gradients of*
653   *wastewater effluent pollution.* Water Research, 2019. **151**: p. 371-380.

654   12.   Peng, Y., et al., *Characteristics of microbial community involved in early biofilms formation*
655   *under the influence of wastewater treatment plant effluent.* Journal of Environmental
656   Sciences, 2018. **66**: p. 113-124.

657   13.   Hladilek, M.D., et al., *Microbial community structure of a freshwater system receiving*
658   *wastewater effluent.* Environmental Monitoring and Assessment, 2016. **188**(11).

659   14.   Chonova, T., et al., *Benthic Diatom Communities in an Alpine River Impacted by Waste Water*
660   *Treatment Effluents as Revealed Using DNA Metabarcoding.* Front Microbiol, 2019. **10**: p.
661   653.

662   15.   Bricheux, G., et al., *Pyrosequencing assessment of prokaryotic and eukaryotic diversity in*
663   *biofilm communities from a F rench river.* MicrobiologyOpen, 2013. **2**(3): p. 402-414.

664   16.   Aubertheau, E., et al., *Impact of wastewater treatment plant discharge on the contamination*
665   *of river biofilms by pharmaceuticals and antibiotic resistance.* Science of The Total
666   Environment, 2017. **579**: p. 1387-1398.

667   17.   Dolinšek, J., F. Goldschmidt, and D.R. Johnson, *Synthetic microbial ecology and the dynamic*
668   *interplay between microbial genotypes.* FEMS Microbiology Reviews, 2016. **40**(6): p. 961-979.

669   18.   De Roy, K., et al., *Synthetic microbial ecosystems: an exciting tool to understand and apply*
670   *microbial communities.* Environ Microbiol, 2014. **16**(6): p. 1472-81.

671   19.   Ben Said, S. and D. Or, *Synthetic Microbial Ecology: Engineering Habitats for Modular*
672   *Consortia.* Frontiers in Microbiology, 2017. **8**(1125).

673   20.   Großkopf, T. and O.S. Soyer, *Synthetic microbial communities.* Current Opinion in
674   Microbiology, 2014. **18**: p. 72-77.

675   21.   Daims, H., S. Lucker, and M. Wagner, *daime, a novel image analysis program for microbial*
676   *ecology and biofilm research.* Environmental Microbiology, 2006. **8**(2): p. 200-213.

677   22.   Brenner, K. and F.H. Arnold, *Self-Organization, Layered Structure, and Aggregation Enhance*
678   *Persistence of a Synthetic Biofilm Consortium.* Plos One, 2011. **6**(2).

679   23.   Li, J.T., et al., *Cadmium pollution triggers a positive biodiversity-productivity relationship:*
680   *evidence from a laboratory microcosm experiment.* Journal of Applied Ecology, 2010. **47**(4): p.
681   890-898.

682   24.   Costello, D.M., et al., *Ranking stressor impacts on periphyton structure and function with*
683   *mesocosm experiments and environmental-change forecasts.* PLOS ONE, 2018. **13**(9): p.
684   e0204510.

685   25.   Godwin, C.M., et al., *Ecological Stoichiometry Meets Ecological Engineering: Using*
686   *Polycultures to Enhance the Multifunctionality of Algal Biocrude Systems.* Environmental
687   Science & Technology, 2017. **51**(19): p. 11450-11458.

26. Ueda, K., Saka, H., Ishikawa, Yoshiyuki, , et al., *Development of a membrane dialysis*
bioreactor and its application to a large-scale culture of a symbiotic bacterium,
Symbiobacterium thermophilum. Applied Microbiology and Biotechnology, 2002. **60**(3): p.
300-305.
27. Ohno, M., et al., *Establishing the Independent Culture of a Strictly Symbiotic*
BacteriumSymbiobacterium thermophilumfrom Its SupportingBacillusStrain. Bioscience,
Biotechnology, and Biochemistry, 1999. **63**(6): p. 1083-1090.
28. Dietz, D., W. Sabra, and A.-P. Zeng, *Co-cultivation of Lactobacillus zeae and Veillonella*
cricetifor the production of propionic acid. AMB Express, 2013. **3**(1): p. 29.
29. Godwin, C.M., et al., *Algal polycultures enhance coproduct recycling from hydrothermal*
liquefaction. Bioresource Technology, 2017. **224**: p. 630-638.
30. Zhang, H. and X. Wang, *Modular co-culture engineering, a new approach for metabolic*
engineering. Metabolic engineering, 2016. **37**: p. 114-121.
31. Burmolle, M., et al., *Enhanced biofilm formation and increased resistance to antimicro¬ bial*
agents and bacterial invasion are caused by synergistic interactions in multispeciesbiofilms.
Applied and Environmental Microbiology, 2006. **72**(6): p. 3916-3923.
32. Hays, S.G., et al., *Better together: engineering and application of microbial symbioses.*
Current Opinion in Biotechnology, 2015. **36**: p. 40-49.
33. Kim, J.-D. and C.-G. Lee, *Microbial degradation of polycyclic aromatic hydrocarbons in soil by*
bacterium-fungus co-cultures. Biotechnology and Bioprocess Engineering, 2007. **12**(4): p.
410-416.
34. Boyero, L., et al., *Biotic vs. Abiotic Control of Decomposition: A Comparison of the Effects of*
Simulated Extinctions and Changes in Temperature. PLoS ONE, 2014. **9**(1): p. e87426.
35. Holmstrup, M., et al., *Interactions between effects of environmental chemicals and natural*
stressors: A review. Science of The Total Environment, 2010. **408**(18): p. 3746-3762.
36. Laskowski, R., et al., *Interactions between toxic chemicals and natural environmental factors*
— A meta-analysis and case studies. Science of The Total Environment, 2010. **408**(18): p.
3763-3774.
37. Larras, F., et al., *The effect of temperature and a herbicide mixture on freshwater periphytic*
algae. Ecotoxicology and Environmental Safety, 2013. **98**: p. 162-170.
38. Morin, S., et al., *Changes in copper toxicity towards diatom communities with experimental*
warming. Journal of Hazardous Materials, 2017. **334**: p. 223-232.
39. Pesce, S., et al., *Experimental Warming Differentially Influences the Vulnerability of*
Phototrophic and Heterotrophic Periphytic Communities to Copper Toxicity. Frontiers in
Microbiology, 2018. **9**(1424).
40. Lambert, A.S., et al., *Temperature modulates phototrophic periphyton response to chronic*
copper exposure. Environmental Pollution, 2016. **208**: p. 821-829.
41. Chaumet, B., et al., *Light and temperature influence on diuron bioaccumulation and toxicity*
in biofilms. Ecotoxicology, 2020. **29**(2): p. 185-195.
42. Passy, S.I., *Diatom ecological guilds display distinct and predictable behavior along nutrient*
and disturbance gradients in running waters. Aquatic Botany, 2007. **86**: p. 171-178.
43. Johnson, R.E., N.C. Tuchman, and C.G. Peterson, *Changes in the Vertical Microdistribution of*
Diatoms within a Developing Periphyton Mat. Journal of the North American Benthological
Society, 1997. **16**(3): p. 503-519.
44. Fairchild, G.W., R.L. Lowe, and W.B. Richardson, *Algal Periphyton Growth on Nutrient-*
Diffusing Substrates: An in situ Bioassay. Ecology, 1985. **66**(2): p. 465-472.
45. Van der Grinten, E., et al., *Phosphate regime structures species composition in cultured*
phototrophic biofilms. Freshwater Biology, 2004. **49**(4): p. 369-381.
46. Irving, T.E., *Factors Influencing the Formation and Development of Microalgal Biofilms.* 2010.
47. Schnurr, P.J. and D.G. Allen, *Factors affecting algae biofilm growth and lipid production: A*
review. Renewable and Sustainable Energy Reviews, 2015. **52**: p. 418-429.

48. Di Gregorio, L., et al., *Unravelling the core microbiome of biofilms in cooling tower systems.*
Biofouling, 2017. **33**(10): p. 793-806.
49. Besemer, K., et al., *Unraveling assembly of stream biofilm communities.* ISME J, 2012. **6**(8): p.
1459-68.
50. Harding, N.E., Y.N. Patel, and R.J. Coleman, *Organization of genes required for gellan*
polysaccharide biosynthesis in Sphingomonas elodea ATCC 31461. Journal of Industrial
Microbiology and Biotechnology, 2004. **31**(2): p. 70-82.
51. Lower, R.L., et al., *An evaluation of periphyton community structure and function on tile and*
cobble substrata in experimental stream mesocosms. Hydrobiologia, 1996. **328**(2): p. 135-
146.
52. Tlili, A., et al., *Micropollutant-induced tolerance of in situ periphyton: Establishing causality in*
wastewater-impacted streams. Water Research, 2017. **111**: p. 185-194.
53. Carles, L., et al., *Impact of wastewater on the microbial diversity of periphyton and its*
tolerance to micropollutants in an engineered flow-through channel system. Water Res, 2021.
203: p. 117486.
54. Pesce, S., et al., *Towards simple tools to assess functional effects of contaminants on natural*
microbial and invertebrate sediment communities. Environmental Science and Pollution
Research, 2020. **27**(6): p. 6680-6689.
55. Muñoz, I., et al., *Effects of atrazine on periphyton under grazing pressure.* Aquatic Toxicology,
2001. **55**(3): p. 239-249.
56. Romero, F., et al., *Effects of multiple stressors on river biofilms depend on the time scale.*
Scientific Reports, 2019. **9**(1).
57. Tolker-Nielsen, T. and S. Molin, *Spatial Organization of Microbial Biofilm Communities.*
Microb Ecol, 2000. **40**(2): p. 75-84.
58. Wagner, M. and H. Horn, *Optical coherence tomography in biofilm research: A*
comprehensive review. Biotechnology and Bioengineering, 2017. **114**(7): p. 1386-1402.
59. Morgenroth, E. and K. Milferstedt, *Biofilm engineering: linking biofilm development at*
different length and time scales. Reviews in Environmental Science and Bio/Technology,
2009. **8**(3): p. 203-208.
60. Jafari, M., et al., *Biofilm compressibility in ultrafiltration: A relation between biofilm*
morphology, mechanics and hydraulic resistance. Water Research, 2019. **157**: p. 335-345.
61. Fortunato, L., et al., *In-situ assessment of biofilm formation in submerged membrane system*
using optical coherence tomography and computational fluid dynamics. Journal of
Membrane Science, 2017. **521**: p. 84-94.
62. Neu, T.R. and J.R. Lawrence, *Investigation of Microbial Biofilm Structure by Laser Scanning*
Microscopy. 2014, Springer International Publishing. p. 1-51.
63. Neu, T.R., et al., *Advanced imaging techniques for assessment of structure, composition and*
function in biofilm systems. FEMS Microbiol Ecol, 2010. **72**(1): p. 1-21.
64. Hartmann, R., et al., *Quantitative image analysis of microbial communities with BiofilmQ.*
Nature Microbiology, 2021. **6**(2): p. 151-156.
65. Morin, S., et al., *Recovery potential of periphytic communities in a river impacted by a*
vineyard watershed. Ecological Indicators, 2010. **10**(2): p. 419-426.
66. Konschak, M., et al., *Herbicide-Induced Shifts in the Periphyton Community Composition*
Indirectly Affect Feeding Activity and Physiology of the Gastropod Grazer Physella acuta.
Environmental Science & Technology, 2021.
67. Congestri, R., et al., *Diatoms (Bacillariophyta) in phototrophic biofilms colonising an Italian*
wastewater treatment plant. Diatom Research, 2005. **20**(2): p. 241-255.
68. Rimet, F. and A. Bouchez, *Use of diatom life-forms and ecological guilds to assess pesticide*
contamination in rivers: Lotic mesocosm approaches. Ecological Indicators, 2011. **11**(2): p.
489-499.

69. Wagenhoff, A., Townsend, Colin R., Phillips, N., Matthaei, Christoph, D., *Subsidy-stress and*
multiple-stressor effects along gradients of deposited fine sediment and dissolved nutrients in
a regional set of streams and rivers. Freshwater Biology, 2011. **56**(9): p. 1916-1936.
70. Aristi, I., et al., *Mixed effects of effluents from a wastewater treatment plant on river*
ecosystem metabolism: subsidy or stress? Freshwater Biology, 2015. **60**(7): p. 1398-1410.
71. Naeem, S., *Species Redundancy and Ecosystem Reliability.* Conservation Biology, 1998. **12**(1):
p. 39-45.
72. Kilham, S.S., et al., *COMBO: a defined freshwater culture medium for algae and zooplankton.*
Hydrobiologia, 1998. **377**(1): p. 147-159.
73. Murga, R., P.S. Stewart, and D. Daly, *Quantitative analysis of biofilm thickness variability.*
Biotechnology and Bioengineering, 1995. **45**(6): p. 503-510.
74. Blauert, F., H. Horn, and M. Wagner, *Time-resolved biofilm deformation measurements using*
optical coherence tomography. Biotechnology and Bioengineering, 2015. **112**(9): p. 1893-
1905.
75. Wagner, M., et al., *Investigation of the mesoscale structure and volumetric features of*
biofilms using optical coherence tomography. Biotechnology and Bioengineering, 2010.
107(5): p. 844-853.
76. Nübel, U., F. Garcia-Pichel, and G. Muyzer, *PCR primers to amplify 16S rRNA genes from*
cyanobacteria. Applied and Environmental Microbiology, 1997. **63**(8): p. 3327-3332.
77. Hugerth, L.W., et al., *Systematic Design of 18S rRNA Gene Primers for Determining Eukaryotic*
Diversity in Microbial Consortia. PLoS ONE, 2014. **9**(4): p. e95567.
78. Vasselon, V., et al., *Assessing ecological status with diatoms DNA metabarcoding: Scaling-up*
on a WFD monitoring network (Mayotte island, France). Ecological Indicators, 2017. **82**: p. 1-
12.
79. Andrews, S., *FastQC: a quality control tool for high throughput sequence data*. 2010.
80. Magoč, T. and S.L. Salzberg, *FLASH: fast length adjustment of short reads to improve genome*
assemblies. Bioinformatics, 2011. **27**(21): p. 2957-2963.
81. Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.*
EMBnet.journal, 2011. **17**(1): p. 10.
82. Schmieder, R. and R. Edwards, *Quality control and preprocessing of metagenomic datasets.*
Bioinformatics, 2011. **27**(6): p. 863-864.
83. Callahan, B.J., P.J. McMurdie, and S.P. Holmes, *Exact sequence variants should replace*
operational taxonomic units in marker-gene data analysis. The ISME Journal, 2017. **11**(12): p.
2639-2643.
84. Munz, N.A., et al., *Pesticides drive risk of micropollutants in wastewater-impacted streams*
during low flow conditions. Water Research, 2017. **110**: p. 366-377.
85. Wittmer, I., Junghaus, M., Singer, H., Stamm, C., *Mikroverunreinigungen*. 2014.
86. Moschet, C., et al., *Evaluation of in-situ calibration of Chemcatcher passive samplers for 322*
micropollutants in agricultural and urban affected rivers. Water Res, 2015. **71**: p. 306-17.

Acknowledgements

This work was supported by the project ToxAdapt funded by Eawag. We thank Alexa Agatiello
for the technical assistance in the DNA samples preparation for sequencing, as well as Dr. Louis
Carles and Prof. Kristin Schirmer for the fruitful discussions and their feedback on previous
drafts of the manuscript. We would also like to thank Silvia Kobel and Jean-Claude Walser
(Genetic Diversity Centre (GDC), ETH Zürich) for their support with community composition

analysis. Data on molecular diversity produced and analyzed in this paper were generated in
collaboration with the GDC, ETH Zürich.

Contributions

OL and AT conceived and designed the study. OL and BW performed the experiments. OL and
ND performed the quantitative OCT analysis. OL and AT analyzed the data and wrote the first
draft of the manuscript. All co-authors contributed to the data interpretation and subsequent
revisions of the manuscript, and they approved the final submitted version.