Overlapping Representation of Basic Tastes in the Human Gustatory Cortex

Du Zhang[a, #], Xiaoxiao Wang[a, #, *], Yanming Wang[a], Benedictor Alexander Nguchu[a], Zhoufang Jiang[a], Chenwei Shi[a], Bensheng Qiu[*]

a.Hefei National Lab for Physical Sciences at the Microscale and the Centers for Biomedical Engineering, University of Science and Technology of China, Hefei, China

*Corresponding author.

Bensheng Qiu, Hefei National Lab for Physical Sciences at the Microscale and the Centers for Biomedical Engineering, University of Science and Technology of China, Hefei, China. Email: bqiu@ustc.edu.cn

Xiaoxiao Wang, Hefei National Lab for Physical Sciences at the Microscale and the Centers for Biomedical Engineering, University of Science and Technology of China, Hefei, China. Email: wang@ustc.edu.cn

# Du Zhang, Xiaoxiao Wang contributed equally to this work

Abstract

The topological representation is a fundamental property of human primary sensory cortices. The human gustatory cortex (GC) responds to the five basic tastes: bitter, salty, sweet, umami, and sour. However, the topological representation of the human gustatory cortex remains controversial. Through functional magnetic resonance imaging(fMRI) measurements of human responses to the five basic tastes, the current study aimed to delineate the taste representations within the GC. During the scanning, the volunteers tasted solutions of the five basic tastes, then washed their mouths with the tasteless solution. The solutions were then sucked from the volunteers' mouths, eliminating the action of swallowing. The results showed that the bilateral mid-insula activated most during the taste task, and the active areas were mainly in the precentral and central insular sulcus. However, the regions responding to the five basic tastes are substantially overlapped, and the analysis of contrasts between each taste response and the averaged response to the remaining tastes does not report any significant results. Furthermore, in the gustatory insular cortex, the multivariate pattern analysis (MVPA) was unable to distinguish the activation patterns of the basic tastes, suggesting the possibility of weakly clustered distribution of the taste-preference neural activities in the human insular cortex. In conclusion, the presented results suggest overlapping representations of the basic tastes in the human gustatory insular cortex.

Keywords: Functional MRI, Taste, MVPA, insular cortex

1 Introduction

Much of the mammal primary sensory, such as somatosensory, visual and
auditory cortex, is organized into topological maps: nearby neurons have similar
sensory receptive fields (Romani, Williamson, & Kaufman, 1982; Sanchez-Panchuelo,
Francis, Bowtell, & Schluppeck, 2010; Wandell, Dumoulin, & Brewer, 2007). Taste
quality is generally acknowledged to contain five perceptual dimensions -- bitter, salty,
sour, sweet and umami (McBurney & Gent, 1979; Simon, de Araujo, Gutierrez, &
Nicolelis, 2006). For taste, the primary gustatory cortex was located in the middle
insula by previous works (Avery et al., 2020; Small, 2010; Veldhuizen et al., 2011;
Yeung, Goto, & Leung, 2018). However, the debate about its topography within the
gustatory cortex lasts for years. X. Chen, Gabitto, Peng, Ryba, and Zuker (2011)
observed highly separated clusters of neurons responding to taste qualities in the
mouse. In controversial, more recent researches observed overlapped clusters of
taste-responsive neurons in the mouse GC (K. Chen, Kogan, & Fontanini, 2021;
Fletcher, Ogg, Lu, Ogg, & Boughter, 2017). Most taste-responsive neurons within the
macaque insular cortex seemed to tune to multiple tastes, with no clear topographic
organization of taste sensitivity (Scott & Plata-Salaman, 1999). Many human
functional magnetic resonance imaging (fMRI) studies explored the question of a
gustotopic organization of the insular cortex. Some of them failed to find spatial
segregation of taste qualities (Dalenberg, Hoogeveen, Renken, Langers, & ter Horst,
2015; Schoenfeld et al., 2004). Some reported gustotopic organization but varied
across participants (Prinster et al., 2017). Over the past few years, the 7Tesla
high-field MRI offered the opportunity of investigating human brain activity in a
higher signal-to-noise ratio (SNR) and higher spatial resolution. The 7Tesla fMRI
studies also failed to find the brain regions that consistently preferred any individual
taste (Avery et al., 2020; Chikazoe, Lee, Kriegeskorte, & Anderson, 2019).
The results of these gustatory fMRI may be affected by some nuisance effects.
The head motion is one of the most significant confoundings and fMRI is extreme
sensitivity to head motion (Matthews & Jezzard, 2004). Swallowing would cause
inescapability task-related head motion and induce related somatosensory (S. L. Yang,
T. J. Ross, Y. Q. Zhang, E. A. Stein, & Y. H. Yang, 2005). The temperature of the
tastants could also contribute to the variability of blood oxygen level-dependent
(BOLD) response in the insular cortex (Guest et al., 2007). Thus, based on Canna et al.
(2019)'s low-cost taste delivery equipment, we built a gustometer equipment furtherly
including a pump for sucking liquid from the subject's mouth and water bathing to
maintain the tastants' temperature. It is also possible that the location of the gustotopic
map might vary across subjects, and it would mess up the results of group analysis. So
extensive fMRI measurements (180 trials per taste per subject) were conducted by our
self-made gustometer and novel taste task design, strictly controlling for the head
movement. Both univariate analyses and multivariate analysis were employed to
explore the taste coding in the human insular cortex.

2   Materials and Methods

2.1 Participants

Four subjects (2 female) aged between 21 and 26 (average: 24.5years) participated in the taste fMRI study. All participants were right-handed, and ethics approval for this study was granted by the University of Science and Technology of China. Participants were excluded from this experiment if they had any history of prenatal drug exposure, current usage of psychotropic medications, neurological illness, or any exclusion criteria for magnetic resonance imaging (MRI). The participants were asked to abstain from food for two hours before participating in experiments.

2.2 Taste solution

Five tastant solutions were administered to our participants during scanning, namely: bitter (0.5mM quinine hydrochloride), salty (180mM NaCl), sour (10mM citric acid), sweet (2mM sucralose), umami (100mM Monosodium Glutamate + 5mM Inosine monophosphate). The tasteless baseline solution was composed of 12.5mM KCL + 1.25mM NaHCO3. The choice of concentrations for each taste referred to previous neuroimaging studies of taste perception (Han, Mohebbi, Unrath, Hummel, & Hummel, 2018; Mascioli et al., 2015; Prinster et al., 2017). Every tastant solution was dissolved in the purified water. The tasteless solution was used as the control to remove the confounding effects of other feelings caused by introducing a fluid into the mouth.

2.3 Gustometer description

A self-made MRI-compatible system was designed to deliver tastants through the peristaltic pump. The uniformity of the temperature of tastant solutions was maintained using a water bath. A single-chip microcontroller STC12C5A60S2 was used to control the supply of the solutions to the subject. The solutions were flowing through the peristaltic pumps and delivered chronologically (Fig.1A). These peristaltic pumps are software-driven, capable of managing the flow rate, being set at 1.27ml/s in our case. Deliverance and removal of tastant solutions were achieved by a well-fabricated tooth socket (Fig.1B). The device was inserted correctly into the subject's mouth during experiment setup to ensure that participants need not swallow during taste fMRI. Visual cues for fMRI tasks were projected onto a screen located inside the scanner bore and viewed through a mirror system mounted on the head-coil. Tastant delivery and visual stimulus presentation were controlled and synchronized via a custom-built program developed in the Psychtoolbox-3 (Brainard, 1997; Kleiner, Brainard, & Pelli, 2007; Pelli, 1997) and Matlab (Mathworks, Natick, MA) environment.

Figure 1. Gustometer and taste fMRI design description. **A.** The taste solution delivery system. Six tastant solutions were delivered to the homemade tooth socket in the subjects' mouth by peristaltic pumps controlled by a single-chip microcontroller. The flow rate was controlled at 1.27 ml/s. Check valves ensure one-way liquid flow, and an independent peristaltic pump extracted the waste solution during the taste fMRI task. **B.** The homemade tooth socket. The six solution-inlet tubes were connected to the anterior inlet pipe and transferred the taste solution to the tip of the tongue. The outlet pipes were attached at the posterior side of the tooth socket and extracted the solution from the tongue base so that subjects were unnecessary to swallow the fluids. **C.** The diagram of the taste task in fMRI. A tasteless stimulus begins, and a queue of 30 trials whose average length was 16s with a taste stimulation for 2.6s and a tasteless inspiration for 11.4-15.4s. **D.** Image acquisition. An axial (left) and coronal (right) view of the image-acquisition areas of the brain. Actual scanning contained 30 slices.

2.4 Behavioral tests

Before the fMRI taste perception task, all subjects underwent behavioral tests to ensure that everyone could discriminate the tastes and correctly identify the solutions. The subjects rated each taste solution's perceived intensity and pleasantness using a visual analog scale from 0 for very weak to 10 for very intense, which was already used and validated in the previous study(Prinster et al., 2017). A paired t-test was used to analyze the behavioral data, with a significance threshold of $P<0.05$.

2.5 Taste task in fMRI

The taste task begins with a tasteless for 12s, then follows a queue of 30 trials with an average length of 16s with a taste stimulation for 2.6s and a tasteless inspiration for 11.4-15.4s (Fig. 1C). For maintaining alarm, the subjects were asked to press the response button through their right thumb once they felt the taste. The

143 sequence of tastes was pseudo-randomized, in which each taste was repeated six times,
144 and the tastes of neighboring trials were different during each scan. Each scan lasted
145 for 492 seconds (8min12sec). The scan was repeated six times per session in five
146 sessions, with one session per day. Thus, we obtained 180 trials of fMRI scan for each
147 of the five tastes of each participant.

148 2.6 FMRI data Acquisition

MRI data scanning was performed at the MRI center of the medical center of the
University of Science and Technology of China. MRI images were acquired with the
aid of a 3-T scanner (Discovery 750; GE Healthcare, Milwaukee, WI) and an
8-channel head coil. T1-weighted images were obtained to provide an anatomical
reference for the data analysis, with repetition time (TR) = 8.2ms, echo time (TE) =

3.2 ms, flip angle =12°, and voxel size =$1\times1\times1$mm^3. The human GC was found to be

in the insular cortex. To get a relatively high spatial resolution in the limited
acquisition time, the field of view (FOV) of the taste task fMRI covered partial brain,
i.e., the whole insular cortex and nearby brain areas(Fig. 1D), with 30 axial slices, TR

= 2000ms, echo time (TE) = 30ms, flip angle =90°, and voxel size =$2\times2\times2$mm^3. A

whole-brain fMRI scan with the same parameter (TR = 2000ms, TE = 30ms, flip

angle =90°, voxel size = $2\times2\times2$mm^3, NEX = 5) was also obtained as registration

reference for the taste task fMRI in the following pre-processing steps.

162 2.7 FMRI data analysis

2.7.1 Univariate analysis

Taste fMRI data analysis was performed using the AFNI software (Cox, 1996).
The first five volumes of each run were removed. Functional images were then slice
timing corrected and realigned within each session. Each TR's head movement
relative to the previous TR was calculated, and a threshold of outlier fraction of 0.2
was used for censoring in the following regression analysis. One run of fMRI data of
the Subject 3 was excluded from the data because of excessive head motion (the mean
outlier fraction of 28[th] to 39[th] TR was 0.424). Both the taste task fMRI volumes and
structural MRI were registered to the reference whole-brain fMRI volume. Spatial
smoothing with a 2mm full width at half maximum (FWHM) of Gaussian was used.

A General Linear Model was created to estimate the BOLD response to the
different conditions. The tasteless solution was the baseline solution whereby the
activeness of the brain region was directly determined by the statistical contrast, taste
(either bitter, salty, sour, sweet, or umami) versus tasteless. In this model, five
regressors of interest for taste (bitter, salty, sour, sweet and umami) trials were made
by the convolution of the trial events and the AFNI's SPMG1 canonical hemodynamic
response function. Movement regressors (3 translations, 3 rotations) were also added.

A high-pass filter was performed to remove slow-varying signal drifts less than 0.01Hz from the fMRI time series. Within-voxel auto-correlations of time-series data were controlled with AFNI's 3dREMLfit. Finally, using the ANTs program, the statistic maps were normalized to ICBM152 2009c space through the nonlinear warp from the registered structural data to the 1.0mm isotropic ICBM152 2009c template(Avants, Epstein, Grossman, & Gee, 2008).

At the subject level, to identify those voxels associated with five basic tastes across five sessions, AFNI's 3dMEMA was utilized with the mating t-statistics and regression coefficients as input. A one-tailed test was used to check for positive effects in five tastes responses. The statistic results were thresholded using a voxel-wise threshold of P<0.01 after the family-wise error (FWE) multiple comparisons, while the Subject3's threshold was set at P<0.02 due to his/her relatively weak taste reaction.

In addition, the statistical mean contrast:(0.2bitter + 0.2salty + 0.2sour + 0.2sweet + 0.2umami) was utilized to corroborate the tastes responsive areas of four subjects and the mean contrast results were FWE corrected thresholding at P<0.005. At the group level, taking within-subject variability into account because there are effect estimates from multiple sessions, regression coefficients for each session were entered into a mixed-effects ANOVA (3dLME in AFNI, z-stats reported) with the session as a random factor (Kaskan, Dean, Nicholas, Mitz, & Murray, 2019). First, we used the statistical contrast: (0.2bitter + 0.2salty + 0.2sour + 0.2sweet + 0.2umami) to identify taste-responsive areas, and the statistical map was corrected for multiple comparisons using the False Discovery Rate (FDR) algorithm (p<0.00001). Second, to identify the brain active areas for five basic tastes, we created group-level maps (bitter, salty, sour, sweet, umami), corrected for multiple comparisons using the False Discovery Rate correction (FDR) algorithm to acquire an overall alpha level of 0.01. We made five taste-responsive areas masks and added them up in a new probability template. Finally, we used the contrast between one taste minus the average responses of other tastes (e.g., bitter - 0.25 salty - 0.25 sour - 0.25 sweet -0.25 umami) to determine brain voxels showing specific taste preferences (3dLME in AFNI; z-stats reported).

To visualize the statistical maps of five taste responses in the insular cortex, freesurfer software (http://surfer.nmr.mgh.harvard.edu/) and Connectome Workbench (https://www.humanconnectome.org/software/connectome-workbench/) were used after multiple comparisons corrected in the volume space. The anatomical surface meshes of the MNI template were reconstructed using Freesurfer's recon-all program, and the five taste statistical map in MNI space were projected to the cortical surface using FreeSurfer's mri_vol2surf command.

2.7.2 Multivariate pattern analyses (MVPA)

MVPA was conducted to assess the difference between five basic taste representations of the GC by The Decoding Toolbox (Hebart, Gorgen, & Haynes, 2014). The classification algorithm used for this analysis was linear support-vector-machine (SVM). We modeled each run of taste fMRI tasks separately, leading to 20 beta-coefficient maps for each taste of every subject. Then we trained

and tested on subject-level regression coefficient maps using leave-one-out
cross-validation (LOOCV). The mid-insula regions-of-interest (ROIs) were identified
by mean contrast of five tastes (p<0.00001, FDR corrected for multiple comparisons),
as described in univariate analysis.

We also used searchlight (three-voxel radius, including 93 voxels) MVPA with
linear SVM kernel to analyze taste responses within the insular cortex. We employed
LOOCV to evaluate the activity patterns of different taste types in the searchlight
spheres. The output of this searchlight analysis was a map of average classification
accuracy minus chance (chanced level = 20%). Then we performed the AFNI program
3dttest++ to make group-level analyses to evaluate the classification results at the
group level and performed FDR corrected for multiple comparisons.

3　Results

3.1 Behavioral results

The range of the taste intensity (*mean±St.d.*) was 4.75±0.96 for bitter, 7.00±0.82
for salty, 4.50±1.00 for sour, 7.75±0.96 for sweet, 8.80±0.58 for umami, and
2.25±0.50 for neutral. As shown in Table 1, all tastes demonstrated significant
differences with the tasteless baseline solution in the intensity rating (P<0.05, paired
t-test). Participants enjoyed the sweet taste most (8.25±0.95), followed by umami
(5.50±1.91), neutral (5.25±0.50), and salty (4.75±0.96), with bitter (4.25±0.50) and
sour (4.50±1.29) being mostly disliked.

Table1.Rating results of the intensity and enjoyment of each taste.

Taste	Rating of the intensity		Rating of the enjoyment	
	Mean (St.d.)	vs. tasteless p-value	Mean (St.d.)	vs. tasteless p-value
bitter	4.75 (0.96)	0.030	4.25 (0.50)	0.092
salty	7.00 (0.82)	0.002	4.75 (0.96)	0.391
sour	4.50 (1.00)	0.037	4.50 (1.29)	0.391
sweet	7.75 (0.96)	0.002	8.25 (0.95)	0.005
umami	8.50 (0.58)	0.001	5.50 (1.91)	0.761
tasteless	2.25 (0.50)		5.25 (0.50)	

3.2 Brain activations by tastes

3.2.1 Gustatory activation areas with fMRI

We used the mean contrast of BOLD responses to five basic tastes for identifying taste-responsive areas. The gustatory activation areas in the group analysis are shown in Fig.2 and Table 2. The bilateral dorsal mid-insula activated most in the taste-responsive region of the brain. We also observed significant activation of the bilateral dorsal anterior insula, rolandic operculum, tongue somatosensory cortex, the right ventral thalamus, and left cerebellum of lobule VI, in agreement with previous human taste neuroimaging studies(Avery et al., 2020; Small, 2010; Yeung et al., 2018). The rolandic operculum has been identified as a cortical region integrating somatosensory and gustatory information from the tongue (Cerf-Ducastel, Ven de Moortele, MacLeod, Le Bihan, & Faurion, 2001; Mascioli et al., 2015). Our results were obtained using FDR corrected for multiple comparisons (P<0.00001).

Figure 2. The gustatory activation areas of the brain. The gustatory cortex exhibited more significant BOLD responses to five basic tastes relative to tasteless included bilateral dorsal mid-insula, dorsal anterior insula, ventral anterior insula, rolandic operculum, and right ventral thalamus (P<0.00001, FDR corrected).

Table 2.Gustatory activation areas responsive to taste

Regions	Voxels	Peak MNI coordinate(mm)			Z value	Volume/mm³
		X	Y	Z		
Right dorsal mid-insula	176	35.0	-9.0	13.0	15.42	1408
Left dorsal mid-insula	172	-35.0	-9.0	15.0	17.08	1376
Right rolandic operculum; Right somatosensory cortex	155	57.0	-17.0	21.0	16.08	1240
Left dorsal anterior insula	74	-29.0	21.0	11.0	13.21	592
Left cerebellum (Lobule VI)	54	-17.0	-69.0	-17.0	9.65	432
Right dorsal anterior insula	39	45.0	17.0	-3.0	11.86	312

Right ventral mid-insula	32	35.0	7.0	-17.0	13.49	256
Left somatosensory cortex	29	-65.0	-25.0	23.0	13.47	232
Left rolandic operculum	26	-47.0	-23.0	21.0	14.60	208
Right ventral thalamus	24	17.0	-11.0	9.0	7.84	192
Left ventral mid-insula	23	-39.0	3.0	-13.0	15.72	184

3.2.2 Five taste responses maps in the insular cortex

At the subject level, the five taste responses maps of four subjects in the insular
cortex were shown in Fig.3.In each subject, the five taste response maps showed an
important, roughly the same bilateral dorsal mid-insula location in the central insular
sulcus. Across four subjects, the five taste response maps demonstrated an
interindividual variability of response intensities. The mean statistic contrast map of
four subjects exhibited a significant activation of bilateral precentral and central
insular sulcus(Naidich et al., 2004; Richardson & Fridriksson, 2016; Uddin, Nomi,
Hebert-Seropian, Ghaziri, & Boucher, 2017).

Figure 3. Activation maps for four subjects' bitter, salty, sour, sweet, and umami
stimuli. **A.** The left hemisphere of five taste responses map in four subjects. **B.** The
right hemisphere of the five taste responses map in four subjects. L, left; R, right.

All tastes showed similar active areas at the group level, and three
taste-responding clusters were observed at each hemisphere. Two of the three clusters
were located in the dorsal mid-insula, with one in the mid of the dorsal mid-insula and
the other closed to the parietal operculum. Another cluster was located in the dorsal
left anterior insula (Fig. 4B).In addition, masks of taste-active voxels were generated
for each taste. The masks were highly overlapped within the central and precentral
insular sulcus (Fig. 4D), which meant the central sulcus was responsible for the
recognition of five tastes. In addition, the activations in the central insular sulcus
showed high symmetry and strong activation, suggesting taste is activated bilaterally.

Figure 4. Group analysis results. **A.** The left and right hemisphere insula is shown
with sulci labeled**. B**. The main three clusters are located in the dorsal middle insula,
ventral middle insula, and dorsal anterior insula in the left hemisphere. The main three
clusters are located in the parietal lobe-dorsal middle insula, mid-dorsal middle insula,
and dorsal anterior insula in the right hemisphere. **C.** taste preference maps of insular
cortex. Maps are thresholded at z = 1.96(P<0.05, cluster unclustered). Contrast
analyses of taste preference areas demonstrate no clear evidence of topographical

organization. **D**. The probability of overlapping of the taste active zone in the insula,
where the five tastes were reliably activated (P<0.01, FDR corrected).L, left; R, right.

3.2.3 Taste preference areas analysis

Contrasts between each taste and the average of the remaining tastes were
calculated (bitter-other, salty-other, sour-other, sweet-other, umami-other) to identify
areas within the insular cortex preferentially activated by either taste. Our voxel-wise
false discovery rate corrected (P<0.05) results did not provide evidence of any regions
of the insular cortex to be exhibiting a significant preference for either bitter, salty,
sour, sweet or umami (Fig. 4B; uncorrected z maps for display only).

3.2.4 MVPA results

We implemented multivariate pattern analysis using the 20 beta-coefficient maps
of each subject within the dorsal mid-insula and anterior insula identified by mean
taste contrast. The decoding results of every subject all failed to identify the five taste
activity patterns. We were unable to distinguish the taste-selective activity patterns
with an average classification accuracy of $20.18\pm5.70\%$ within the left dorsal
mid-insula and $22.24\pm4.97\%$ within the right dorsal mid-insula (Fig.5A). The
taste-selective activity patterns were also not distinguished with an average
classification accuracy of $21.33\pm2.98\%$ within the left dorsal anterior insula and
$20.48\pm3.56\%$ within the right dorsal anterior insula (Fig.5B). The MVPA
classification results near the random level revealed the variability of taste-responsive
patterns within different sessions.

Figure 5. Multivariate pattern analyses results of region-of-interest within the bilateral dorsal mid-insula and dorsal anterior insula. **A.**ROIs of the dorsal mid-insula and the dorsal anterior insula.**B.** The confusion matrix of the dorsal anterior insula.

C.The confusion matrix of the dorsal anterior insula.

Then we used the multivariate searchlight analysis to identify the regions which proved to have above-chance classification accuracy for discriminating between five tastes responses in the insular cortex. No voxel was found to survive after FDR

321 multiple comparisons (P<0.5). Our multivariate searchlight analysis exhibited little
322 discrimination of taste activation patterns in the insular cortex (Fig. 5; uncorrected
323 maps presented for display only).

324

325 **Figure 6.** Multivariate searchlight analysis results within the insular cortex. The
326 insular cortex shows little discrimination of taste activation patterns. The maps above
327 show uncorrected results at a low statistical threshold (P<0.05, cluster unclustered).L,
328 left; R, right.

4 Discussion

This study aimed to derive comprehensive gustatory topologic maps of the
human gustatory insular cortex. A self-developed taste delivery system was conducted
to deliver the five basic taste and tasteless solutions to the subjects' tongues,

controlling for the effect of swallowing and the tastant temperature. Four participants

were scanned over five sessions while performing the tasting task. Results show that
the tastes activate a wide range of brain areas within the dorsal mid-insula,
somatosensory cortex, dorsal anterior insula and ventral mid-insula. Both the
individual-level and group-level univariate analysis demonstrates that each basic taste
activates similar insular areas, which are mainly in the precentral and central insular
sulcus. A further MVPA was unable to distinguish the basic tastes either in the dorsal
anterior insula or the dorsal middle insula.

Another change of our experiment design compared to previous taste
experiments is the continuously flowing liquids within the subjects' mouth – taste (at
task state) or tasteless (at baseline state) liquids were continuously pumped into the
subjects' mouth. Thus, the somatosensory responses to the liquid flows were
minimized, and the bilateral gustatory insular cortex became the most activated area
in the present work. We used sucralose rather than sucrose for the sweet solutions.
Because sucrose solutions (for example, 0.6Mol/L sucrose) are so viscous that it is not
fully washed out during the washing stage of the experiment, leading to constant
sweetness feeling. Sucralose is about 600 times as sweetening compared to sucrose
and activates the human insular cortex in a similar pattern but lower intensity
compared to sucrose (Frank et al., 2008). The sucralose solutions (2mMol/L) used in
the present work were easily washed out, and eliminate the taste interference.

Human gustatory experience is a complex process referring to the perception of
taste quality, intensity, and pleasantness (Bender, Veldhuizen, Meltzer, Gitelman, &

Small, 2009; Small, 2010). Taste receptor cells (TRCs) clustered in taste buds on the
tongue are the first place where taste information processing begins. After taste
information discharges from the TRCs, it flows along the chorda tympani (the three
cranial nerves XII) and glossopharyngeal (cranial nerves IX), and lingual (cranial
nerves X) to the nucleus of the solitary tract (NST). In human beings, taste signal is
carried from the NST to the ventral posteromedial nucleus of the thalamus (VPMNT)
and then to the gustatory insular cortex (Flynn, 1999). The right ventral thalamus
cluster of the activation for five basic tastes further confirmed the experimental design
was reliable in locating the taste functional regions of the brain.

Our observation of taste responses in the bilateral dorsal anterior insula, dorsal
middle insula and parietal operculum is consistent with previous researches (Avery et
al., 2020; Veldhuizen et al., 2011; Yeung, Goto, & Leung, 2017; Yeung et al., 2018).
The dorsal mid-insula is thought to be the primary gustatory cortex (Bender et al.,
2009; Small, 2010). The human secondary somatosensory cortex is located at the
parietal operculum (Eickhoff, Schleicher, Zilles, & Amunts, 2006). The dorsal anterior
insula was likely to be an integration area of multiple sensory information and also
engaged in cognition and attention function (Kurth, Zilles, Fox, Laird, & Eickhoff,
2010). Our work furtherly shows that the taste response areas are within the central
and precentral insular sulcus of the human middle insula.

However, the gustatory insular cortex responding to each basic taste is
substantially overlapped. The taste preference areas analysis provides no evidence of
any regions of the insular cortex exhibiting a significant preference for either bitter,
salty, sour, sweet, or umami. The present work shows that the taste response areas are
overlapped at both the individual level and the group level. A further MVPA was also
unable to distinguish the basic tastes either in the dorsal anterior insula or the dorsal
middle insula. fMRI MVPA fails when neuronal responses are weakly
clustered(Dubois, de Berker, & Tsao, 2015). Thus, failure of the fMRI MVPA in the
present study suggests the possibility of weakly clustered distribution of the
taste-preference neurons in the human insular cortex. This is compatible with the most
recent mammal studies, which have observed taste-preference neurons spatially
distributing throughout the superficial layer of GC with no sign of spatial clustering or
topography (K. Chen et al., 2021; Fletcher et al., 2017). A previous MVPA work
distinguishes among the sweet, salty, and sour tastants above chance levels by the
signal from the middle insula(Avery et al., 2020). There are two noticeable differences
between Avery et al. (2020)'s and ours: First, the action of swallowing was avoided in
the present study, and so was the swallowing related neural activity (S. Yang, T. J.
Ross, Y. Zhang, E. A. Stein, & Y. Yang, 2005); Second, the duration of taste stimulus
is much shorter in our work (2.6 seconds *vs.*20 seconds), and it lessens the potential
effect of top-down affective responses or cognitive activities (Craig, 2002; Singer,
Critchley, & Preuschoff, 2009) in the insula.

Although taste quality information is not represented by spatially unique clusters
of neurons within the insular cortex, a microscale organization of taste selective
responses neurons in the human insular cortex is still possible. In addition, specific

temporal patterns of neuronal activity may represent both quality and valence. The temporal resolutions of fMRI may limit the study of the dynamic of taste perception, which could distinguish the taste quality information at the voxel level (Crouzet, Busch, & Ohla, 2015; Iannilli, Noennig, Hummel, & Schoenfeld, 2014). Imaging techniques with higher temporal and spatial resolution would be essential to explore the chemotopic organization of taste quality in the insular cortex.

5   Conclusions

The present study investigated the BOLD response of taste stimuli while controlling for effects induced by swallowing and tastant temperature. The results showed that the bilateral mid-insula activated most during the taste task, and the active areas were mainly in the precentral and central insular sulcus. But the areas responding to the five basic tastes were overlapped in the gustatory insular cortex. Furthermore, we did not identify any regions within the insula that exhibited a preference for a specific taste. Similarly, the multivariate pattern analysis results suggested it was difficult to catch the fixed pattern of five taste responses in the human insular cortex. In conclusion, our results support the idea that taste quality is not represented topographically in the human insula.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant nos. 81701665, 21876041) and the support of the Foundation Students' Innovation and Entrepreneurship Foundation of USTC (nos.CY2022X09).

Data and code availability statement

The data and code used in this paper are available via a request to the corresponding author.

References

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Med Image Anal, 12*(1), 26-41. doi:10.1016/j.media.2007.06.004

Avery, J. A., Liu, A. G., Ingeholm, J. E., Riddell, C. D., Gotts, S. J., & Martin, A. (2020). Taste Quality Representation in the Human Brain. *Journal of Neuroscience, 40*(5), 1042-1052. doi:10.1523/JNEUROSCI.1751-19.2019

Bender, G., Veldhuizen, M. G., Meltzer, J. A., Gitelman, D. R., & Small, D. M. (2009). Neural correlates of evaluative compared with passive tasting. *European Journal of Neuroscience, 30*(2), 327-338. doi:10.1111/j.1460-9568.2009.06819.x

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436. doi:10.1163/156856897x00357

Canna, A., Prinster, A., Fratello, M., Puglia, L., Magliulo, M., Cantone, E., . . . Esposito, F. (2019). A low-cost open-architecture taste delivery system for gustatory fMRI and BCI experiments. *J Neurosci Methods, 311*, 1-12. doi:10.1016/j.jneumeth.2018.10.003

Cerf-Ducastel, B., Ven de Moortele, P. F., MacLeod, P., Le Bihan, D., & Faurion, A. (2001). Interaction of gustatory and lingual somatosensory perceptions at the cortical level in the human: a functional magnetic resonance imaging study. *Chemical Senses, 26*(4), 371-383. doi:DOI 10.1093/chemse/26.4.371

Chen, K., Kogan, J. F., & Fontanini, A. (2021). Spatially Distributed Representation of Taste Quality in the Gustatory Insular Cortex of Behaving Mice. *Curr Biol, 31*(2), 247-256 e244. doi:10.1016/j.cub.2020.10.014

Chen, X., Gabitto, M., Peng, Y., Ryba, N. J., & Zuker, C. S. (2011). A gustotopic map of taste qualities in the mammalian brain. *Science, 333*(6047), 1262-1266. doi:10.1126/science.1204076

Chikazoe, J., Lee, D. H., Kriegeskorte, N., & Anderson, A. K. (2019). Distinct representations of basic taste qualities in human gustatory cortex. *Nature Communications, 10*. doi:ARTN 1048

10.1038/s41467-019-08857-z

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res, 29*(3), 162-173. doi:10.1006/cbmr.1996.0014

Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nat Rev Neurosci, 3*(8), 655-666. doi:10.1038/nrn894

Crouzet, S. M., Busch, N. A., & Ohla, K. (2015). Taste Quality Decoding Parallels Taste Sensations. *Current Biology, 25*(7), 890-896. doi:10.1016/j.cub.2015.01.057

Dalenberg, J. R., Hoogeveen, H. R., Renken, R. J., Langers, D. R., & ter Horst, G. J. (2015). Functional specialization of the male insula during taste perception. *Neuroimage, 119*, 210-220. doi:10.1016/j.neuroimage.2015.06.062

Dubois, J., de Berker, A. O., & Tsao, D. Y. (2015). Single-Unit Recordings in the Macaque Face Patch System Reveal Limitations of fMRI MVPA. *Journal of Neuroscience, 35*(6), 2791-2802. doi:10.1523/Jneurosci.4037-14.2015

Eickhoff, S. B., Schleicher, A., Zilles, K., & Amunts, K. (2006). The Human Parietal Operculum. I. Cytoarchitectonic Mapping of Subdivisions. *Cerebral Cortex, 16*(2), 254-267. doi:10.1093/cercor/bhi105

Fletcher, M. L., Ogg, M. C., Lu, L., Ogg, R. J., & Boughter, J. D., Jr. (2017). Overlapping Representation of Primary Tastes in a Defined Region of the Gustatory Cortex. *J Neurosci, 37*(32), 7595-7605. doi:10.1523/JNEUROSCI.0649-17.2017

Flynn, F. G. (1999). Anatomy of the insula functional and clinical correlates. *Aphasiology, 13*(1), 55-78. doi:10.1080/026870399402325

Frank, G. K., Oberndorfer, T. A., Simmons, A. N., Paulus, M. P., Fudge, J. L., Yang, T. T., & Kaye, W. H. (2008). Sucrose activates human taste pathways differently from artificial sweetener. *Neuroimage, 39*(4), 1559-1569. doi:10.1016/j.neuroimage.2007.10.061

Guest, S., Grabenhorst, F., Essick, G., Chen, Y., Young, M., McGlone, F., . . . Rolls, E. T. (2007). Human
cortical representation of oral temperature. *Physiol Behav, 92*(5), 975-984.
doi:10.1016/j.physbeh.2007.07.004

Han, P. F., Mohebbi, M., Unrath, M., Hummel, C., & Hummel, T. (2018). Different Neural Processing of
Umami and Salty Taste Determined by Umami Identification Ability Independent of Repeated
Umami Exposure. *Neuroscience, 383*, 74-83. doi:10.1016/j.neuroscience.2018.05.004

Hebart, M. N., Gorgen, K., & Haynes, J. D. (2014). The Decoding Toolbox (TDT): a versatile software
package for multivariate analyses of functional imaging data. *Front Neuroinform, 8*, 88.
doi:10.3389/fninf.2014.00088

Iannilli, E., Noennig, N., Hummel, T., & Schoenfeld, A. M. (2014). Spatio-temporal correlates of taste
processing in the human primary gustatory cortex. *Neuroscience, 273*, 92-99.
doi:10.1016/j.neuroscience.2014.05.017

Kaskan, P. M., Dean, A. M., Nicholas, M. A., Mitz, A. R., & Murray, E. A. (2019). Gustatory responses
in macaque monkeys revealed with fMRI: Comments on taste, taste preference, and internal
state. *Neuroimage, 184*, 932-942. doi:10.1016/j.neuroimage.2018.10.005

Kleiner, M., Brainard, D., & Pelli, D. (2007). What's new in Psychtoolbox-3?

Kurth, F., Zilles, K., Fox, P. T., Laird, A. R., & Eickhoff, S. B. (2010). A link between the systems:
functional differentiation and integration within the human insula revealed by meta-analysis.
Brain Struct Funct, 214(5-6), 519-534. doi:10.1007/s00429-010-0255-z

Mascioli, G., Berlucchi, G., Pierpaoli, C., Salvolini, U., Barbaresi, P., Fabri, M., & Polonara, G. (2015).
Functional MRI cortical activations from unilateral tactile-taste stimulations of the tongue.
Physiology & Behavior, 151, 221-229. doi:10.1016/j.physbeh.2015.07.031

Matthews, P. M., & Jezzard, P. (2004). Functional magnetic resonance imaging. *Journal of Neurology,
Neurosurgery & Psychiatry, 75*(1), 6-12.

McBurney, D. H., & Gent, J. F. (1979). On the nature of taste qualities. *Psychol Bull, 86*(1), 151-167.
Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/377351

Naidich, T. P., Kang, E., Fatterpekar, G. M., Delman, B. N., Gultekin, S. H., Wolfe, D., . . . Yousry, T. A.
(2004). The insula: Anatomic study and MR Imaging display at 1.5 T. *American Journal of
Neuroradiology, 25*(2), 222-232. Retrieved from <Go to ISI>://WOS:000189105600012

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7974606/pdf/0911.pdf

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers into
movies. *Spatial Vision, 10*(4), 437-442. doi:10.1163/156856897x00366

Prinster, A., Cantone, E., Verlezza, V., Magliulo, M., Sarnelli, G., Lengo, M., . . . Esposito, F. (2017).
Cortical representation of different taste modalities on the gustatory cortex: A pilot study. *Plos
One, 12*(12). doi:ARTN e0190164

10.1371/journal.pone.0190164

Richardson, J. D., & Fridriksson, J. (2016). The Insular Cortex. In *Neurobiology of Language* (pp.
115-127).

Romani, G. L., Williamson, S. J., & Kaufman, L. (1982). Tonotopic organization of the human auditory
cortex. *Science, 216*(4552), 1339-1340. doi:10.1126/science.7079770

Sanchez-Panchuelo, R. M., Francis, S., Bowtell, R., & Schluppeck, D. (2010). Mapping Human
Somatosensory Cortex in Individual Subjects With 7T Functional MRI. *Journal of*
Neurophysiology, 103(5), 2544-2556. doi:10.1152/jn.01017.2009

Schoenfeld, M. A., Neuer, G., Tempelmann, C., Schussler, K., Noesselt, T., Hopf, J. M., & Heinze, H. J.
(2004). Functional magnetic resonance tomography correlates of taste perception in the human
primary taste cortex. *Neuroscience, 127*(2), 347-353. doi:10.1016/j.neuroscience.2004.05.024

Scott, T. R., & Plata-Salaman, C. R. (1999). Taste in the monkey cortex. *Physiol Behav, 67*(4), 489-511.
doi:10.1016/s0031-9384(99)00115-8

Simon, S. A., de Araujo, I. E., Gutierrez, R., & Nicolelis, M. A. (2006). The neural mechanisms of
gustation: a distributed processing code. *Nat Rev Neurosci, 7*(11), 890-901.
doi:10.1038/nrn2006

Singer, T., Critchley, H. D., & Preuschoff, K. (2009). A common role of insula in feelings, empathy and
uncertainty. *Trends Cogn Sci, 13*(8), 334-340. doi:10.1016/j.tics.2009.05.001

Small, D. M. (2010). Taste representation in the human insula. *Brain Struct Funct, 214*(5-6), 551-561.
doi:10.1007/s00429-010-0266-9

Uddin, L. Q., Nomi, J. S., Hebert-Seropian, B., Ghaziri, J., & Boucher, O. (2017). Structure and Function
of the Human Insula. *Journal of Clinical Neurophysiology, 34*(4), 300-306.
doi:10.1097/Wnp.0000000000000377

Veldhuizen, M. G., Albrecht, J., Zelano, C., Boesveldt, S., Breslin, P., & Lundstrom, J. N. (2011).
Identification of Human Gustatory Cortex by Activation Likelihood Estimation. *Human Brain*
Mapping, 32(12), 2256-2266. doi:10.1002/hbm.21188

Wandell, B. A., Dumoulin, S. O., & Brewer, A. A. (2007). Visual field maps in human cortex. *Neuron,*
56(2), 366-383. doi:10.1016/j.neuron.2007.10.012

Yang, S., Ross, T. J., Zhang, Y., Stein, E. A., & Yang, Y. (2005). Head motion suppression using real-time
feedback of motion information and its effects on task performance in fMRI. *Neuroimage, 27*(1),
153-162. doi:10.1016/j.neuroimage.2005.02.050

Yang, S. L., Ross, T. J., Zhang, Y. Q., Stein, E. A., & Yang, Y. H. (2005). Head motion suppression using
real-time feedback of motion information and its effects on task performance in fMRI.
Neuroimage, 27(1), 153-162. doi:10.1016/j.neuroimage.2005.02.050

Yeung, A. W. K., Goto, T. K., & Leung, W. K. (2017). Basic taste processing recruits bilateral
anteroventral and middle dorsal insulae: An activation likelihood estimation meta-analysis of
fMRI studies. *Brain and Behavior, 7*(4), e00655. doi:10.1002/brb3.655

Yeung, A. W. K., Goto, T. K., & Leung, W. K. (2018). Affective value, intensity and quality of liquid
tastants/food discernment in the human brain: An activation likelihood estimation meta-analysis.
Neuroimage, 169, 189-199. doi:10.1016/j.neuroimage.2017.12.034

547