Antiviral activity of pacific oyster (*Crassostrea gigas*) hemolymph against a human coronavirus

Rebecca L. Pedler[a], James O. Harris[a], Peter G. Speck[a]#

[a]College of Science and Engineering, Flinders University, Bedford Park, South Australia, Australia.

Running Head: Antiviral activity of *C. gigas* against HCoV-229E

#Address correspondence to Peter G. Speck, peter.speck@flinders.edu.au

Mailing address:

College of Science and Engineering, Flinders University, Bedford Park,

GPO Box 2100

Adelaide 5001,

South Australia, Australia.

ABSTRACT

Coronaviruses can cause severe respiratory infections in humans. In this study we assessed

the antiviral activity of Pacific oyster (*Crassostrea gigas*) hemolymph against a human

coronavirus, HCoV-229E. An eight-fold reduction in infectivity of HCoV-229E on Huh-7

cells was observed in the presence of 10% *C. gigas* hemolymph. Antiviral activity of *C. gigas*

hemolymph positively correlated with its concentration and appears to be active during an

intracellular stage of HCoV-229E infection.

KEYWORDS: molluscan antivirals, HCoV-229E, marine invertebrates, *Crassostrea gigas*,

SARS-CoV-2

Human coronaviruses are enveloped, single stranded RNA viruses that are further classified

as alpha-coronaviruses (human coronavirus-229E (HCoV-229E) and HCoV-NL63) or beta-

coronaviruses (HCoV-OC43, HCoV-HKU1, middle eastern respiratory syndrome (MERS-

CoV) and severe acute respiratory syndrome (SARS-CoV and SARS-CoV-2) (1). SARS-

CoV-2 is a novel human coronavirus which emerged in December 2019 as the causative

agent of coronavirus disease 2019 (Covid-19) (2-4). Safe and effective antiviral treatments

for SARS-CoV-2 are yet to be identified, despite multiple drug repurposing attempts (5-7).

Marine molluscs represent an unexploited source of medicinal compounds (R.L. Pedler, and

P.G. Speck, submitted for publication) (8-11). Marine molluscs lack an adaptive immune

system and exclusively elicit innate immune responses (12-14), while living in an

environment containing virus particles in the order of $>10^7$ per ml (15, 16). This demonstrates

the success of their strategies to prevent viral infection, which includes production of potent

antiviral compounds (11, 13). *In vitro* inhibition of HSV-1 has been observed using extracts

from the common cockle (*Cerastoderma edule*), greenlip abalone (*Haliotis laevigata*) (17),

Japanese carpet shell (*Ruditapes philippinarum*), European flat oyster (*Ostrea edulis*),

common whelk (*Buccinum undatum*) (18), blacklip abalone (*Haliotis rubra*) (19, 20), veined

rapa whelk (*Rapanosa venosa*) (21) and Mediterranean mussel (*Mytilus galloprovincialis*)

(22). Extract from the flesh of the red abalone (*Haliotis rufescens*) protects mice against

poliovirus and influenza A (23, 24), while paolin II from the Eastern oyster (*Crassostrea*

virginica) inhibits poliovirus (25).

Hemolymph of the Pacific oyster (*Crassostrea gigas)* has *in vitro* antiviral activity against

HSV-1 and adenovirus respiratory strain 5 (AdV-5) (26-28). The major *C. gigas* hemolymph

protein, cavortin, exerts an antiviral effect against HSV-1 after entry into Vero cells (26).

Cavortin is a Mr 20,000 protein which acts as a metal chaperone (29). Intracellular zinc has

therapeutic potential for SARS-CoV-2 (30-32), and its efficacy is improved by coupling with

a metal chaperone (31, 33). *Crassostrea gigas* has a high zinc content (34) and given

cavortin's suggested role as a metal chaperone (29), it is possible that *C. gigas* cavortin has

potential antiviral activity against SARS-CoV-2 and may facilitate zinc transport into host

cells.

The discovery of antiviral agents for SARS-CoV-2 is challenged by the limited number of

laboratories with the appropriate biosafety containment level (35, 36). HCoV-229E can be

handled in lesser-rated laboratories making it more accessible for research on human

coronaviruses (37) and this virus could be used for initial screening for anti-coronavirus

activity. This study is the first, that we are aware of, which assesses antiviral activity of *C.*

gigas hemolymph against HCoV-229E.

Huh-7 cells, obtained from M. Beard (Adelaide University, South Australia), were grown in

Dulbecco's modified eagle medium (DMEM) (Gibco #11965118) supplemented with 10%

foetal bovine serum (FBS) (Gibco #10099141), according to standard methods (38). Twelve

C. gigas oysters, grown in Coffin Bay, South Australia, were purchased from local seafood

merchants. After opening, *C. gigas* hemolymph was extracted from the pericardial cavity

using a sterile syringe and 27g needle. Hemolymph was pooled, filter sterilised using a

0.2μm filter and stored at -20°C until required. Cytotoxicity of *C. gigas* hemolymph was

determined using a trypan blue exclusion assay (26). Huh-7 cells were seeded into a 24-well

plate with medium as above and 0%, 2%, 5%, 10% or 20% (v/v) *C. gigas* hemolymph. Cells

were incubated for two days at 37°C in 5% CO_2, before being stained *in situ* with 0.4%

trypan blue (Gibco #15250061) (26). The number of non-viable cells in three different fields

of view were then counted under an Olympus CK2 microscope at 40x magnification (39).

The mean number of non-viable cells was lowest for 0% (1.00 ± 0.00 cells) and 2% (1.00 ±

0.47 cells) (Fig. 1). Huh-7 cell death sharply increased as hemolymph concentration exceeded

10% (Fig. 1), therefore, 10% was considered an appropriate concentration for use in anti-

HCoV-229E assays. In Vero cells, *C. gigas* hemolymph can cause 10% cell death at a

concentration of 13% (v/v) (26) or 50% cell death at $750\mu g\ ml^{-1}$ (27). The acellular and

cellular fractions of *C. gigas* hemolymph can cause 50% cell death in Hep-2 cells at

concentrations of $0.32mg\ ml^{-1}$ and $0.19mg\ ml^{-1}$ respectively (28).

HCoV-229E was obtained from H. Whiley (Flinders Water Quality and Health Research

Consortium, South Australia). Virus titres were determined as 50% tissue culture infective

doses ($TCID_{50}$) (40). Huh-7 cells were seeded into 96-well plates and either 0 or 10% *C.*

gigas hemolymph. Three ten-fold dilutions, followed by eight two-fold dilutions were

prepared using HCoV-229E stock and DMEM and inoculated into 96-well plates. Cells were

incubated at 37°C in 5% CO_2 for five days before being fixed with 10% formaldehyde and

stained with 0.5% crystal violet (Thermo #S25275B). Wells illustrating cytopathic effect

were counted, allowing $TCID_{50}$ to be calculated using the Reed-Muench method (41). When

Huh-7 cells were assayed with 10% *C. gigas* hemolymph, an eight-fold reduction in the

HCoV-229E titre ($4.00 \times 10^5\ TCID_{50}\ ml^{-1}$) and an antiviral activity of 87.5% was observed

(Table 1). A dose-response curve was generated using 0%, 2%, 5%, 10% and 15% *C. gigas*

hemolymph which revealed that antiviral activity positively correlated with its concentration

(Table 1, Fig. 3). This is consistent with the dose-dependent antiviral activity of *C. gigas*

hemolymph protein, cavortin, against HSV-1 (26).

Time of addition assays were used to determine the stage of HCoV-229E infection targeted

by *C. gigas* hemolymph. In previous studies, the greatest antiviral protection of Vero cells

from HSV-1 infection was observed when *C. gigas* hemolymph was added 0-2 hours after

infection (26, 27), suggesting that the antiviral effect was most likely exerted after virus

attachment and entry. An intracellular mode of antiviral action has been observed for other

molluscan compounds, including lipophilic extract of *H. laevigata* (17) and myticin C

peptides from *M. galloprovincialis* (22). Here, Huh-7 cells were seeded into 24-well plates

with DMEM and 10% FBS. A series of dilutions (10^{-4}, 10^{-5}, 10^{-6} and 10^{-7}) were prepared

using HCoV-229E stock and DMEM and inoculated into plates. *C. gigas* hemolymph was

added either immediately or 60 minutes after HCoV-229E and a negative control plate was

also prepared. There was little difference in *C. gigas* hemolymph antiviral activity when it

was added to Huh-7 cells immediately (98.21%) or 60 minutes after HCoV-229E infection

(96.11%) (Table 2). This suggests that *C. gigas* hemolymph most likely acted during an

intracellular stage of HCoV-229E infection. Antiviral compounds which act during an

intracellular stage of HCoV-229E infection have been identified (42-44). The macrolide and

immunosuppressive compound, FK06 inhibits HCoV-229E replication in Huh-7 cells (44),

while the antimalarial drug chloroquine inhibits HCoV-229E replication in epithelial lung

cells (L132) by suppressing P38MAPK (43). Thapsigargin, from the Thapsia (*Thapsia*

garganica) plant acts during an intracellular stage of HCoV-229E infection either by

inhibiting replication or activating unknown antiviral effector systems in Huh-7 cells (42).

This study reveals that *C. gigas* hemolymph has *in vitro* antiviral activity against human

coronavirus HCoV-229E. This finding is relevant in the current pandemic and reinforces that

C. gigas hemolymph has broad-spectrum antiviral activity. Further research is required to

identify and characterise the antiviral compound(s) produced by *C. gigas.*

REFERENCES

1. Woo PCY, Huang Y, Lau SKP, Yuen K-Y. 2010. Coronavirus genomics and bioinformatics analysis. Viruses 2:1805-1820.
2. Wang C, Horby PW, Hayden FG, Gao GF. 2020. A novel coronavirus outbreak of global health concern. Lancet 395:470-473.
3. Ghanbari R, Teimoori A, Sadeghi A, Mohamadkhani A, Rezasoltani S, Asadi E, Jouyban A, Sumner SCJ. 2021. Existing antiviral options against SARS-CoV-2 replication in COVID-19 patients. Future Microbiol 15.
4. Triggle CR, Bansal D, Ding H, Islam MM, Farag EABA, Hadi HA, Sultan AA. 2021. A comprehensive review of viral characteristics, transmission, pathophysiology, immune response, and management of SARS-CoV-2 and COVID-19 as a basis for controlling the pandemic. Front Immunol 12:631139.
5. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, Shi Z, Hu Z, Zhong W, Xiao G. 2020. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell Res 30:269-271.
6. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, Hohmann E, Chu HY, Luetkemeyer A, Kline S, Lopez de Castilla D, Finberg RW, Dierberg K, Tapson V, Hsieh L, Patterson TF, Paredes R, Sweeney DA, Short WR, Touloumi G, Lye DC, Ohmagari N, Oh M-d, Ruiz-Palacios GM, Benfield T, Fätkenheuer G, Kortepeter MG, Atmar RL, Creech CB, Lundgren J, Babiker AG, Pett S, Neaton JD, Burgess TH, Bonnett T, Green M, Makowski M, Osinusi A, Nayak S, Lane HC. 2020. Remdesivir for the treatment of Covid-19 — final report. N Engl J Med 383:1813-1826.
7. Indari O, Jakhmola S, Manivannan E, Jha HC. 2021. An update on antiviral therapy against SARS-CoV-2: How far have we come? Front Pharmacol 12:632677.
8. Benkendorff K. 2010. Molluscan biological and chemical diversity: secondary metabolites and medicinal resources produced by marine molluscs. Biol Rev Camb Philos Soc 85:757-775.
9. Summer K, Browne J, Liu L, Benkendorff K. 2020. Molluscan compounds provide drug leads for the treatment and prevention of respiratory disease. Mar Drugs 18:570.
10. Riccio G, Ruocco N, Mutalipassi M, Costantini M, Zupo V, Coppola D, de Pascale D, Lauritano C. 2020. Ten-year research update review: Antiviral activities from marine organisms. Biomolecules 10:1-36.
11. Sagar S, Kaur M, Minneman KP. 2010. Antiviral lead compounds from marine sponges. Mar Drugs 8:2619-2638.
12. Agius JR, Corbeil S, Helbig KJ. 2020. Immune control of herpesvirus infection in molluscs. Pathogens (Basel) 9:1-11.
13. Dang VT, Benkendorff K, Green T, Speck P. 2015. Marine snails and slugs: A great place to look for antiviral drugs. J Virol 89:8114-8118.
14. Tiscar PG, Mosca F. 2004. Defense mechanisms in farmed marine molluscs. Vet Res Commun 28:57-62.
15. Bratbak G, Heldal M, BØrsheim KY, Bergh Ø. 1989. High abundance of viruses found in aquatic environments. Nature 340:467-468.
16. Suttle CA. 2007. Marine viruses - major players in the global ecosystem. Nat Rev Microbiol 5:801-812.
17. Dang VT, Benkendorff K, Speck P. 2011. *In vitro* antiviral activity against herpes simplex virus in the abalone *Haliotis laevigata*. J Gen Virol 92:627-637.

18. Defer D, Bourgougnon N, Fleury Y. 2009. Screening for antibacterial and antiviral
activities in three bivalve and two gastropod marine molluscs. Aquaculture 293:1-7.
19. Talaei Zanjani N, Miranda-Saksena M, Valtchev P, Diefenbach RJ, Hueston L,
Diefenbach E, Sairi F, Gomes VG, Cunningham AL, Dehghani F. 2016. Abalone
hemocyanin blocks the entry of herpes simplex virus 1 into cells: A potential new
antiviral strategy. Antimicrob Agents Chemother 60:1003-1012.
20. Dang VT, Speck P, Doroudi M, Smith B, Benkendorff K. 2011. Variation in the antiviral
and antibacterial activity of abalone *Haliotis laevigata*, *H. rubra* and their hybrid in
South Australia. Aquaculture 315:242-249.
21. Nesterova NV, Zagorodnya SD, Moshtanska V, Dolashka P, Baranova GV, Golovan AV,
Kurova AO. 2011. Antiviral activity of hemocyanin isolated from marine snail *Rapana*
venosa. Antiviral Res 90:A38-A38.
22. Novoa B, Romero A, Álvarez ÁL, Moreira R, Pereiro P, Costa MM, Dios S, Estepa A,
Parra F, Figueras A. 2016. Antiviral activity of myticin C peptide from mussel: An
ancient defense against herpesviruses. J Virol 90:7692-7702.
23. Li C, Precott B, Jahnes W. 1962. Antiviral activity of a fraction of abalone juice. Proc
Soc Exo Biol Med 109:534-538.
24. Li C. 1960. Antimicrobial effect of abalone juice. Proc Soc Exo Biol Med 103:522-524.
25. Prescott B, Li C, Caldes G, Martino E. Chemical studies of Paolin. II. An antiviral
substance from oysters. Proc Soc Exo Biol Med 123:460-464.
26. Green TJ, Robinson N, Chataway T, Benkendorff K, O'Connor W, Speck P. 2014.
Evidence that the major hemolymph protein of the Pacific oyster, *Crassostrea gigas*,
has antiviral activity against herpesviruses. Antiviral Res 110:168-174.
27. Olicard C, Renault T, Torhy C, Benmansour A, Bourgougnon N. 2005. Putative
antiviral activity in hemolymph from adult Pacific oysters, *Crassostrea gigas*. Antiviral
Res 66:147-152.
28. Carriel-Gomes MC, Kratz JM, Muller VDM, Barardi CRM, Simues CMO. 2006.
Evaluation of antiviral activity in hemolymph from oysters *Crassostrea rhizophorae*
and *Crassostrea gigas*. Aquat Living Resour 19:189-193.
29. Scotti PD, Dearing SC, Greenwood DR. 2007. Characterisation of cavortin, the major
haemolymph protein of the Pacific oyster (*Crassostrea gigas*). N Z J Mar Freshwater
Res 41:91-101.
30. Carlucci PM, Ahuja T, Petrilli C, Rajagopalan H, Jones S, Rahimian J. 2020. Zinc sulfate
in combination with a zinc ionophore may improve outcomes in hospitalized COVID-
19 patients. J Med Microbiol 69:1228-1234.
31. Brewer J, Gomez Marti JL, Brufsky A. 2021. Potential interventions for SARS-CoV-2
infections: Zinc showing promise. J Med Virol 93:1201-1203.
32. Asl SH, Nikfarjam S, Majidi Zolbanin N, Nassiri R, Jafari R. 2021.
Immunopharmacological perspective on zinc in SARS-CoV-2 infection. Int
Immunopharmacol 96:107630.
33. te Velthuis AJW, van den Worml SHE, Sims AC, Baric RS, Snijder EJ, van Hemert MJ.
2010. Zn2+ inhibits coronavirus and arterivirus RNA polymerase activity in vitro and
zinc ionophores block the replication of these viruses in cell culture. PLoS Pathogens
6:e1001176.
34. Kong N, Zhao Q, Liu C, Li J, Liu Z, Gao L, Wang L, Song L. 2020. The involvement of
zinc transporters in the zinc accumulation in the Pacific oyster *Crassostrea gigas*.
Gene 750:144759-144759.

35. Kaufer AM, Theis T, Lau KA, Gray JL, Rawlinson WD. 2020. Laboratory biosafety
measures involving SARS-CoV-2 and the classification as a Risk Group 3 biological
agent. Pathology 52:790-795.
36. Hall MD, Anderson JM, Anderson A, Baker D, Bradner J, Brimacombe KR, Campbell
EA, Corbett KS, Carter K, Cherry S, Chiang L, Cihlar T, de Wit E, Denison M, Disney M,
Fletcher CV, Ford-Scheimer SL, Götte M, Grossman AC, Hayden FG, Hazuda DJ,
Lanteri CA, Marston H, Mesecar AD, Moore S, Nwankwo JO, O'Rear J, Painter G,
Saikatendu KS, Schiffer CA, Sheahan TP, Shi P-Y, Smyth HD, Sofia MJ, Weetall M,
Weller SK, Whitley R, Fauci AS, Austin CP, Collins FS, Conley AJ, Davis MI. 2021.
Report of the national institutes of health SARS-CoV-2 antiviral therapeutics summit.
J Infect Dis 224:S1-S21.
37. Bracci N, Pan H-C, Lehman C, Kehn-Hall K, Lin S-C. 2020. Improved plaque assay for
human coronaviruses 229E and OC43. PeerJ 8:e10639-e10639.
38. Thermofisher. 2021. Gibco Cell Culture Basics.
https://www.thermofisher.com/au/en/home/references/gibco-cell-culture-
basics.html. Accessed October 24th.
39. Strober W. 2019. Trypan blue exclusion test of cell viability. Curr Protoc Immunol
111.
40. Smither SJ, Lear-Rooney C, Biggins J, Pettitt J, Lever MS, Olinger GG. 2013.
Comparison of the plaque assay and 50% tissue culture infectious dose assay as
methods for measuring filovirus infectivity. J Virol Methods 193:565-571.
41. Reed LJ, Muench H. 1938. A simple method of estimating fifty per cent endpoints.
Am J Epidemiol 27.
42. Shaban MS, Müller C, Mayr-Buro C, Weiser H, Meier-Soelch J, Albert BV, Weber A,
Linne U, Hain T, Babayev I, Karl N, Hofmann N, Becker S, Herold S, Schmitz ML,
Ziebuhr J, Kracht M. 2021. Multi-level inhibition of coronavirus replication by
chemical ER stress. Nature communications 12:5536-5536.
43. Kono M, Tatsumi K, Imai AM, Saito K, Kuriyama T, Shirasawa H. 2008. Inhibition of
human coronavirus 229E infection in human epithelial lung cells (L132) by
chloroquine: Involvement of p38 MAPK and ERK. Antiviral Res 77:150-152.
44. Carbajo-Lozoya J, Müller MA, Kallies S, Thiel V, Drosten C, von Brunn A. 2012.
Replication of human coronaviruses SARS-CoV, HCoV-NL63 and HCoV-229E is
inhibited by the drug FK506. Virus Res 165:112-117.
249
250

FIG. 1. Mean (± standard deviation) number of non-viable Huh-7 cells treated with varying

concentrations of Pacific oyster (*Crassostrea gigas*) hemolymph (0, 2, 5, 10, 20% v/v).

FIG. 2. non-viable (arrow) and viable (arrowhead) Huh-7 cells treated with 20% Pacific

oyster (*Crassostrea gigas*) hemolymph.

TABLE 1. Virus titre values ($TCID_{50}$ ml^{-1}) for human coronavirus 229E (HCoV-229E) in

Huh-7 cells treated with varying concentrations (0, 2, 5, 10, 15% v/v) of Pacific oyster

(*Crassostrea gigas*) hemolymph.

Extract/control	Virus titre ($TCID_{50}$ ml^{-1})	Antiviral activity: % Reduction in virus titre
Negative control (DMEM)	3.20×10^6	0.00
2% *C. gigas* hemolymph	8.00×10^5	75.00
5% *C. gigas* hemolymph	7.92×10^5	75.24
10% *C. gigas* hemolymph	4.00×10^5	87.50
15% *C. gigas* hemolymph	2.83×10^5	91.16

FIG. 3. Virus titre values ($Log(TCID_{50}\ ml^{-1})$) for human coronavirus 229E (HCoV-229E) in

Huh-7 cells treated with varying concentrations (0, 2, 5, 10, 15% v/v) of Pacific oyster

(*Crassostrea gigas*) hemolymph (HL).

TABLE 2. Virus titres ($TCID_{50}\ ml^{-1}$) and antiviral activity (% reduction in virus titre) for

human coronavirus 229E (HCoV-229E) in Huh-7 cells treated with either 0% (negative

control) or 10% Pacific oyster (*Crassostrea gigas*) hemolymph.

Extract/control	Time of extract addition	Virus titre ($TCID_{50}\ ml^{-1}$)	Antiviral activity: % reduction in virus titre
Negative control (DMEM)	-	2.79×10^8	0.00
10% *C. gigas* hemolymph	Immediately after infection	5.00×10^6	98.21
10% *C. gigas* hemolymph	60minutes after infection	1.08×10^7	96.11