Substitution load revisited:
a high proportion of deaths can be selective

Joseph Matheson[1,2], Moises Exposito-Alonso[3], Joanna Masel[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona 85721, USA

[2]Department of Ecology, Behavior, and Evolution, University of California San Diego, San Diego, CA, 92093, USA

[3]Departments of Plant Biology & Global Ecology, Carnegie Institution for Science, Stanford University, Stanford, California 94305, USA

Corresponding author: Joanna Masel, 1041 E Lowell St, Tucson AZ 85721, USA. masel@arizona.edu

Article Type: Letter

ABSTRACT

Haldane's Dilemma refers to the concern that the need for many "selective deaths" to complete a substitution (i.e. selective sweep) creates a speed limit to adaptation. However, discussion of this concern has been marked by confusion, especially with respect to the term "substitution load". Here we distinguish different historical lines of reasoning, and identify one, focused on finite reproductive excess and the proportion of deaths that are "selective" (i.e. causally contribute to adaptive allele frequency changes), that has not yet been fully addressed. We develop this into a more general theoretical model that can apply to populations with any life history, even those for which a generation or even an individual are not well defined. The actual speed of adaptive evolution is coupled to the proportion of deaths that are selective. The degree to which reproductive excess enables a high proportion of selective deaths depends on the details of when selection takes place relative to density regulation, and there is therefore no general expression for a speed limit. To make these concepts concrete, we estimate both reproductive excess, and the proportion of deaths that are selective, from a dataset measuring survival of 517 different genotypes of *Arabidopsis thaliana* grown in eight different environmental conditions. In this dataset, a much higher proportion of deaths contribute to adaptation, in all environmental conditions, than

the 10% cap that was anticipated as substantially restricting adaptation during historical discussions of

speed limits.

Keywords: cost of selection, adaptation rate, genetic load, fitness component, biological individual

LAY SUMMARY

The influential neutral theory of molecular evolution was predicated on theoretical arguments that

adaptation is subject to a speed limit. We resolve confusions regarding historical speed limit arguments,

which depend on differences in fitness, not variance (differences in fitness squared). We generalize the

underlying concepts of selective deaths and reproductive excess to populations with any life cycle, even

those for which an "individual" and hence generation and fitness, are poorly defined. We apply the

revised theory to Arabidopsis data, demonstrating the potential for future related experiments.

INTRODUCTION

During an adaptive sweep, new alleles must be substituted for old alleles across an entire population. This

means that individuals with the old alleles must leave no descendants that carry them, and individuals

with new alleles must produce enough offspring to replenish the population. These requirements limit the

number of sweeps that can happen within a given time. Haldane (1957) used this reasoning to propose a

rough estimate of the maximum speed of adaptation.

Haldane's arguments were the historical motivation for the development of Kimura's (Kimura 1968)

neutral theory. This speed limit became known as Haldane's dilemma (Van Valen 1963) because data on

rates of amino acid divergence between species seemed to exceed Haldane's speed limit. The

development of neutral theory resolved this apparent dilemma by suggesting that most amino acid

substitutions are neutral and do not count against the speed limit. However, the basis for this historical

argument is now on troubled ground, because recent literature argues that the fraction of amino acid

substitutions explained by adaptation can be high (Galtier 2016; Murga-Moreno et al. 2024; Sella et al.

2009; Uricchio, Petrov, and Enard 2019). On shorter timescales, recent experiments have shown rapid

seasonal adaptation in *Drosophila* (Bertram 2021; Bitter et al. 2024; Kelly 2022; Machado et al. 2021).

There are other possible resolutions — e.g. some estimates include substitutions of neutral alleles via

hitchhiking. Nevertheless, it is curious that the empirical collapse of historical arguments for neutral

theory has not yet led to a re-evaluation of related arguments by Haldane. Here we revise Haldane's

arguments for the modern era, finding them compatible with empirical evidence for abundant adaptation,

while still imposing upper limits that might matter in some contexts.

Haldane's logic has previously been challenged on multiple counts (Ewens 1970; Felsenstein 1971; Kern

and Hahn 2018; Maynard Smith 1968a). Here we first synthesize the historical literature, drawing out two

key quantities (see Results). First, a population has a "reproductive excess", meaning how many

individuals could be produced at a given life history stage, in excess of the minimum needed to maintain a

constant population size, if all individuals had the best genotype present (see Glossary for definitions of

the terms used here). The second key quantity is the proportion of deaths that are "selective deaths"

(including foregone fertility), meaning deaths that causally contribute to changes in allele frequencies.

The "cost of selection" is the number of selective deaths required for selection to effect the allele

frequency changes needed to achieve a given adaptation rate. Haldane (1957) assumed, based on the

limited evidence available at the time, that finite reproductive excess limited the budget for selective

deaths such that at most 10% of deaths were selective. This assumed 10% limit was not challenged by

subsequent work.

Haldane's calculation was criticized by Ewens (1970) for inappropriate reference to an optimal genotype

that is unlikely to exist. However, Nei (1971) and Felsenstein (1971) derived a near-identical speed limit

to Haldane's, without this flaw, using a model that more explicitly captures the finite nature of

reproductive excess.

Models of finite reproductive excess need at least two life history stages: adults and juveniles, where the

latter shows reproductive excess relative to the former. Selective deaths are then relative not to a single

adult population size N as in Haldane's (1957) model, but to the population size at the appropriate life

history stage (Kimura and Crow 1969). Once multiple life history stages are considered, assumptions

regarding when and how density regulation operates must be made explicit (Haldane 1956; Nicholson

1933). Standard relative fitness models have an implied infinite number of juveniles subject to perfect

density regulation in their transition to an adult population of size N (Bertram and Masel 2019).

An emphasis on life history transitions rather than on generations is a strength rather than a weakness of

the selective deaths view. One of the many flaws of the concept of "fitness" (Van Valen 1989) is the

difficulty of defining a "generation" for many species, especially colonial species for which an

"individual" is not well defined (Wilson and Barker 2021). Consider for example the budding yeast

Saccharomyces cerevisiae. Is each mitotic division a generation? Or each life cycle spanning from

meiosis to meiosis, with a variable number of mitoses in between? Or the span between outcrossing

events, with variable occurrences of selfing as well as mitoses in between? Or is a generation best defined

ecologically with respect to dispersal between resources that allow growth? Problems defining a

generation arise for a broad range of species (albeit not humans, nor many other animal species), but are

resolved when population dynamics are viewed as a series of life history transitions (Smith et al. 2024).

The "generation" that matters in this view is not one complete life cycle, but rather the "generation" of

reproductive excess, in contrast to other life history transitions that involve survival rather than

reproduction.

After synthesizing the literature, here we reformulate and generalize Nei's (1971) and Felsenstein's

(1971) ideas to selection on both fecundity and survival, to life cycles with selection at more than one

stage, and to life cycles with a variable number of stages. We clarify the concepts of reproductive excess

and selective deaths, and use our general theory to pose two empirically accessible questions. First, how

much reproductive excess does the best genotype have? Second, what fraction of deaths can be selective,

and how does this compare to the 10% limit assumed by Haldane? Posing questions in this form allows us

to make the first empirical estimates with which to ground Haldane's approach. We use data from

Exposito-Alonso et al. (2019), who counted or estimated every plant grown and seed produced of *A.*

thaliana cultivars from 517 different genotypes in one season, under 8 distinct environmental conditions.

These data are not representative of natural conditions, but they suffice to illustrate how such an analysis

can be done. Ours is the first direct application of selective death and reproductive excess arguments to

empirical data.

RESULTS

Synthetic historical review

Haldane's argument in his seminal 1957 paper contains two key parts, both novel at the time. In the first

part, he defined "selective deaths" as the subset of deaths $s(1-p)N$ that contribute to an increase in the

allele frequency p, assuming a haploid population where $1-s$ individuals without the beneficial allele

survive to reproductive maturity for every 1 individual with the beneficial allele. These deaths are

considered selective because the $(1-p)N$ individuals that lack a beneficial mutation expect s more

deaths than they would if they had the mutation, and those extra deaths are required for selection to have

its effects. Selective "deaths" can include absence of potential offspring, not just literal deaths, because

reduced fecundity is mathematically equivalent to increased mortality.

Haldane defined the "cost of selection" as the number of selective deaths occurring during a substitution

(i.e. a selective sweep from low allele frequency to fixation). He calculated this cost as the integral of

$s(1-p)N$ over the course of a sweep from allele frequency $p = p_0$ to close to 1 (Figure 1A). In a

haploid population of constant size N, one sweep requires $N \times D$ selective deaths, where $D =$

$-\ln(p_0)+O(s)$. For appropriately small s and p_0 (Haldane suggests $s < \frac{1}{3}$ and $p_0 = 10^{-4}$), the first term

dominates, making D nearly independent of s. For alternative assumptions about ploidy, dominance, and

degree of inbreeding, D is a different function of p_0, but s remains unimportant unless close to 1 (Haldane

1957). $D = 18.4$ for a representative case of $p_0 = 10^{-4}$ at a diploid autosomal locus with fitnesses 1, $1-$

s, and $1 - 2s$ for individuals homozygotic for the beneficial allele, heterozygotic, and homozygotic for

the wild-type allele, respectively. Haldane conservatively estimated that 20-30N selective deaths are

needed per sweep in natural populations.

In the second part of his argument, Haldane converts this estimate of selective deaths per sweep into an

upper bound on the number of sweeps per generation. Haldane considered independent sweeps at x loci,

such that the current allele frequency p_i at the i^{th} locus reduces population fitness by a factor of $1 - d_i =$

$1 - s_i(1 - p_i)$ relative to its post-sweep value. The fitness of the population is then lower than that of a

hypothetical population fixed for all segregating beneficial alleles by a factor of $\prod_{i=1}^{x}(1 - d_i) \approx$

$e^{-\sum_{i=1}^{x} d_i}$. Haldane observed that the mean individual therefore has $\sim e^{-\sum_{i=1}^{x} d_i}$ surviving descendants for

every 1 surviving descendant that an individual with all segregating beneficial alleles would have.

Assuming that this ideal individual exists, and that sites evolve independently, there are $\sum d_i N$ selective

deaths per generation. Comparing these to the $30N$ selective deaths needed per sweep, Haldane obtains a

minimum average spacing between fixation events of $n \geq \frac{30}{\sum d_i}$ generations.

Haldane argues that a maximum 'selection intensity' defines an upper bound on $\sum d_i$, and therefore on the

speed of adaptation. Haldane defines selection intensity $I = \ln\left(\frac{s_o}{S}\right)$, where s_o is the survival of the best

genotype in the population and S is the average survival of the population. Haldane's calculations proceed

with the implicit assumption that the best genotype in the population (referenced by s_o) will also be the

genotype with the beneficial version at all n segregating sites. Plugging in $n = \frac{30}{\sum d_i}$ yields a selection

intensity $I = \ln\left(\frac{s_o}{S}\right) = \frac{30}{n}$.

Finally, Haldane argued that 10% selection intensity was approximately the upper limit that could be

sustained long-term in most species, citing limited empirical knowledge of selection intensities available

at the time. From this estimated value, he derived a speed limit based on $I \leq 0.1 = \frac{30}{n}$, yielding $n \geq$

300 generations between sweeps. Subsequent discussion remained framed around this 10% figure.

However, if the optimal genotype is not present in the population, selection intensity will be lower than

Haldane calculated (Ewens 1970). An optimal genotype is likely present under stabilizing selection,

which Haldane's (1954, 1957) use of infant survival as a function of birth weight suggests he was

envisaging. However, his calculations entail directional selection.

Today, "selection intensity" has acquired a somewhat different definition within the Breeder's Equation

of quantitative genetics (Walsh and Lynch 2018). Haldane's (1957) selection intensity, defined with

respect to fitness (infant survival) rather than phenotype (birth weight), is better compared to load $L =$

$\frac{W_{opt} - \bar{W}}{W_{opt}}$, where $\bar{W}$ is population mean fitness relative to some optimal genotype's fitness W_{opt}.

Load relative to the best genotype present is now known as the 'lead', inspired by the degree to which the

fittest part of the population "leads" the bulk (Desai and Fisher 2007) (see Figure 1B and the Glossary).

We use the term 'lag load' (Maynard Smith 1976) for load relative to an optimal genotype $W_{opt} = 1$

possessing all currently segregating beneficial alleles (see Figure 1B and the Glossary).

The fact that there are so many amino acid substitutions, each requiring a sweep, was the original

evidence supporting neutral theory (Kimura 1968). Kimura and Ohta (1971) plugged in estimates of the

actual rate of substitution in mammalian lineages as n and rearranged Haldane's equation to get

$L = e^{-30/n}$, which produced what they considered to be an excessively large load. Their argument was

subtly different from Haldane's, arguing that a high load implies that typical individuals would need to

have a biologically implausible fraction of their offspring die (Kimura and Ohta 1971). It propagates

Haldane's (1957) problematic assumption that an individual with all beneficial variants is present in the

population (Ewens 1970).

Ewens (1970) adjusted Haldane's (1957) calculations to instead use the lead. Prior to recent travelling

wave models (Desai and Fisher 2007), approximations for the lead were derived from variance in fitness

(Ewens 1970; Kimura 1969). With many independent sweeps at once, variance in fitness (after

normalizing mean population fitness as 1) is approximately s/n, where s is the selection coefficient of an

adaptive allele and n is the number of generations between fixation events (Ewens 1970). The fittest

genotype likely to be present can be estimated using the statistics of extreme values, e.g. around 4.9

standard deviations above the mean for a population of size 10^6 (Ewens 1970). Using Haldane's 10% as

an upper bound on the lead instead of on the lag load yields $4.9\sqrt{s/n} = 0.1$. For $s = 0.01$, n is around 20,

much less than Haldane's estimate of 300, and n is lower still for lower s. In other words, Ewens (1970)

found that many simultaneous sweeps do not imply an implausibly large lead, and the corresponding

speed limit of $n \approx 20$ is not an obstruction with respect to observed rates of amino acid divergence.

Similar arguments have been applied to deleterious mutation load (Galeota-Sprung, Sniegowski, and

Ewens 2020).

Although Ewens' argument revolved around lead, which is a normalized difference between fitnesses, his

approach continued the traditional emphasis of evolutionary genetics on variance in fitness, which

describes the mean square of differences (Crow 1958; Ewens 2004; Fisher 1930). Modern traveling wave

theory instead derives the lead directly from s, N, and the beneficial mutation rate U, and obtains the

variance in fitness variance only downstream from that (Desai and Fisher 2007), rather than relying on

our ability to directly measure fitness and its variance as an input to the calculation of the lead.

Figure 1. Three different types of arguments have been used to argue for limits to the speed of adaptation.

A) The cost of selection for one sweep is the cumulative number of selective deaths that must occur to

complete that sweep (each independent sweep shown as a logistic curve). The cost of selection at one

timepoint is the sum of the instantaneous costs $s(1-p)N$ being incurred by each sweep at that timepoint,

illustrated as the slopes of the orange lines. The sample calculation shown in orange calculates the total

number of selective deaths for a set of allele frequencies arbitrarily chosen for simplicity of calculation (p

= 0.05, 0.1, 0.5, 0.8, 0.9, 0.95, 0.98, and 0.996). B) Load is a normalized difference between an optimal

fitness and mean population fitness: $L = \frac{W_{opt} - \bar{W}}{W_{opt}}$. Different loads can be defined by comparing to

different 'optimal' fitnesses. Lag load uses the difference between a hypothetical optimal genotype and

the population mean, while the lead uses the difference between the best currently living genotype and the

population mean. C) Finite reproductive excess imposes an upper limit on how many selective deaths per

generation are possible, which sets an upper bound to how fast substitutions can occur. Haldane (1957)

combined the cost of selection (A) with lag load (B) to produce his speed limit, while Ewens (1970)

corrected this calculation to use lead (B). In contrast, Nei (1971) and Felsenstein (1971) use the cost of

selection (A) and finite reproductive excess (C) to derive a speed limit.

Maynard Smith (1968b) made a quite different argument against speed limits, claiming that the reason

Haldane's dilemma is not a problem is pervasive synergistic epistasis. Synergistic epistasis increases

differences in fitness above those expected from differences in the numbers of beneficial mutations,

thereby making each selective death tend to count towards a larger number of sweeps at once. A

persistent source of confusion has been that Maynard Smith modeled epistasis using truncation selection,

where the least fit half of a population dies each generation, with the explicit assumption that the

remaining half always have enough reproductive capacity to reconstitute the population (Maynard Smith

1968b). The degree to which the relaxed limits on adaptation in Maynard Smith's model can be attributed

to epistasis vs. assuming infinite reproductive excess (see below) is thus unclear.

Although Ewens' lead-based approach negates the second part of Haldane's arguments that converted lag

load into a speed limit via selection intensity, this does not negate the first part of Haldane's argument,

namely that 20-30N selective deaths are needed for a sweep. Nei (1971) and Felsenstein (1971) made a

key advance building on just this first part of Haldane's argument. This advance has not received

substantial attention, perhaps partially due to confusion about terminology, and also because growing

acceptance of the neutral theory of molecular evolution had seemed to resolve the immediate issues posed

by speed limits.

The cost of selective deaths must be paid out of a finite budget of reproductive excess. The relative fitness

models that dominate population genetics (e.g. Wright-Fisher and Moran) implicitly assume inexhaustible

reproductive excess (Bertram and Masel 2019). This can be seen easily when simulating selection on

zygote viability using rejection sampling — when fitness is low, an absurd number of zygotes might be

generated and discarded prior to filling the N slots, potentially enabling the few survivors to include

exceptionally rare beneficial events such as double mutations. However, real populations have a finite

reproductive excess, e.g. human females do not easily give birth to more than 20 infants. This constrains

members of the next generation to options within a finite set of potential offspring. This concept has been

applied to lethal mutagenesis strategies for anti-viral drugs (Bull, Sanjuán, and Wilke 2007).

Nei (1971) and Felsenstein (1971) each modelled independently evolving sites in a haploid population

with adult population size N. Each adult has fecundity k, i.e. produces k offspring prior to juvenile deaths

(Figure 1C, first arrow). In their deterministic models, k is exact, but the theory readily generalizes to

interpreting k as an expectation. The raw reproductive excess is thus $(k-1)N$, with $k > 1$. Extrinsic

mortality that occurs prior to selective mortality, and hence at a fixed rate, can be treated as an effectively

lower value of k (Figure 1C, second arrow). Haldane's 10% selection intensity estimate is equivalent to

10% of deaths being selective, which corresponds to $k = 1.1$. Nei (1971) and Felsenstein (1971) retain

this 10% cap for illustrative purposes, where k includes the fact that extrinsic mortality substantially

reduces the fecundity available to be 'used' for selective deaths. Population size regulation after selective

mortality (Figure 1C, far right) will on average consume the remaining reproductive excess that was not

used by selection.

The population undergoes sweeps, all with the same initial frequency p_0 and selection coefficient s

applying to survival rather than fecundity. Each sweep follows the same trajectory with a mean delay of n

generations between sweeps (Figure 1A). Given independent sites, the cost of selection is summed across

loci at any given point in time (e.g. slopes of orange lines in Figure 1A); Haldane calculates the

expectation of this sum by integrating across time points for a single sweep. Comparing this cost to the

reproductive excess of the population produced the novel result that the minimum spacing n is

$-\ln{(p_0)}\big/\ln{(k)}$ (Felsenstein 1971; Nei 1971). For Haldane's estimates of $p_0 = 10^{-4}$ and $k = 1.1$, this

yields $n = 97$ generations between selective sweeps. This can be compared to Haldane's original spacing

of $-\ln{(p_0)}\big/\ln{(\frac{W_{max}}{\overline{W}})} = 92$, where Haldane's selection intensity $\ln{\left(\frac{W_{max}}{\overline{W}}\right)} = 0.1$ (by assumption) has

been replaced by $\ln(k) = \ln(1.1) \approx 0.095$.

Importantly, this speed limit calculation is not subject to the same criticisms as Haldane's original

argument. Where Haldane's calculation includes a comparison between the mean fitness of the population

and an optimal genotype which may or may not exist, Nei's (1971) and Felsenstein's (1971) approach

compares the available reproductive excess to the reproductive excess required to effect changes in allele

frequencies. Even if no individual exists who possesses the beneficial allele at every segregating site, each

sweep still requires a certain fraction of deaths to contribute to its selection. It is the finite nature of

reproductive excess that directly produces this limit on the rate of adaptation.

While an identical speed limit is derived by both papers, they interpret the speed limit differently. Nei

(1971) interprets k, and hence the speed limit, in the context of a strongly density-regulated population of

constant size. Under this interpretation, the rate at which mutations sweep, each affecting relative fitness,

is constrained, but this cannot threaten population persistence. Felsenstein (1971) considers the upper

bound on k applying at low population density and applies this to the context of ongoing environmental

change that degrades absolute fitness, requiring adaptive rescue at a rate constrained by a speed limit. A

population that lacks sufficient reproductive excess, even at low density, goes extinct.

Critical questions remain unanswered. Felsenstein (1971) considers a hypothetical population at the edge

of viability, while Nei (1971) considers a hypothetical population adapting at the maximum possible

speed — is it possible to relate their key theoretical quantities to fecundity and mortality data on real

populations? How do the speed limit and its consequences depend on the details of density regulation in

real populations, and the timing of selection relative to density regulation? Do populations with high

fertility have substantially more permissive speed limits, or is this balanced by greater loss of

reproductive excess to extrinsic mortality?

Theory

Life histories compatible with Nei's and Felsenstein's model

Life history goes through at least two transitions, which in Nei (1971) and Felsenstein (1971) are simple:

births and deaths, in that order. They assign all genotypes the same fertility k. They consider only the

limiting case in which all deaths are selective. Some violations of these obviously unrealistic assumptions

can be incorporated without changing the model.

Non-selective deaths can be incorporated as causing k to fall to $k_{effective}$, where Nei (1971) and

Felsenstein (1971) solve for $\text{Cost of selection} = k_{effective} - 1$. Density-independent non-selective

deaths, aligned with Felsenstein's (1971) interpretation, are well modelled as occurring prior to selection

on survival, reducing k in the current generation. Density-dependent non-selective deaths, aligned with

Nei's (1971) interpretation, are best incorporated into the framework after selection on survival. Here they

can balance out deaths that were "unused" by selection in the current generation, by reducing k in the

subsequent generation. Fig. 1C illustrates both kinds of non-selective deaths, proceeding first through

non-selective fecundity to produce raw reproductive excess, then the non-density-dependent component

of extrinsic mortality, then selective deaths, and finally density-dependent extrinsic mortality to cap the

population size at N adults.

Nei (1971) and Felsenstein (1971) emphasize literal deaths, but were presumably aware that differences

in fecundity also contribute selective 'deaths'. This is because mathematically, foregone fecundity is

equivalent to deaths that take place immediately after fecundity, and can be treated as:

$$\text{Selective 'deaths' during differential fecundity} = N_i(b_{best} - b_i)$$

where N_i is the number of reproductive mature adults, b_i is the fecundity of genotype i, and b_{best} is the

fecundity of the genotype with the highest fecundity in that environment.

Reproductive excess within a fixed life cycle

Next we generalize from just one life history transition experiencing selection, to multiple that occur in a

consistent order. We consider a life history transition j that starts with population size N_j and ends at

population size $N_{j+1} = k_j N_j$: We now define

$$\text{Reproductive excess after transition } j = k_j N_j - N_{min,j+1}$$

where $N_{min,j+1}$ is the minimum population size at the end of transition j that is required in order for the

population to achieve size N_j at the beginning of transition j in the next life history cycle. Note that $k_j >$

1 indicates fecundity, while $k_j \leq 1$ indicates survival.

For example, consider a microbial population in batch culture that alternates between exponential growth,

stationary phase, and dilution life history transitions. For stationary phase, k is not much less than 1 but

kN is relatively large because N is large. Many cells will be lost in the subsequent dilution (say 1:100),

but then exponential growth will typically yield N cells from any starting density, so long as seeding is

successful. $N_{min} > 100$ is thus required to recover N live cells at the beginning of the next stationary

phase. In contrast, for the dilution life history transition, $k \sim 0.01$ and N_{min} required in principle to regrow

the population in permissive media may not be much larger than 1.

Reproductive excess for transition j provides the maximum budget for how many selective deaths could

occur during transition j, without upsetting population dynamics by overwhelming the compensatory

capacity of density regulation during other transitions. The calculation of $N_{min,j+1}$ includes density

regulation at all life history transitions except j, while density regulation during transition j is held

constant in its representation by $k_j N_j$. Reproductive excesses specific to a particular life history transition

already take other transitions into account, and so reproductive excesses across different transitions do not

need to be further combined.

To produce selective deaths, $k_{i,j}$ must depend on genotype i. To capture density regulation, $k_{i,j}$ for at

least some values of j must depend on population size N_j. The values $k_{i,j}$ can also be functions of the

genotype frequencies and/or an absolute measure of time. Two life history transitions (survival and

fecundity) are the minimum, but each of these can be broken up into multiple transitions. For example,

survival ($k < 1$) can be broken into components representing survival at different ages, or into a selective

component depending only on genotype vs. a density-dependent extrinsic mortality component depending

only on N_j vs. an extrinsic mortality component occurring at a constant rate. The degree to which

selection and density regulation act on births vs. deaths has consequences, e.g. for rapid adaptation

(Draghi, McGlothlin, and Kindsvater 2024).

Reproductive excess is calculated for a focal population. The best choice of focal population depends on

the question that the model is intended to answer. For example, studying balancing selection calls for the

actual population, while studying evolutionary rescue (Bell 2017) in an asexual population calls for a

hypothetical population where each individual has the current best genotype present. These choices will

result in different values of k_j and $N_{min,j+1}$.

To calculate reproductive excess with respect to the actual population, we solve for $N_{min,j+1}$ in:

$$N_j = N_{min,j+1} \sum_i f_i \prod_{x \neq j} k_{i,x},$$

where f_i is the frequency of genotype i at the beginning of the transition. With respect to a population

where each individual has the best genotype, we instead solve for $N_{min,j+1}$ in:

$$N_j = N_{min,j+1} \prod_{x \neq j} k_{best,x}.$$

Speed limits with multiple life history stages

Under Nei's (1971) interpretation, a population has only one value of reproductive excess $RE =$

$(k - 1)N$, and one (adult) population size N, from which he derived a speed limit of

$-\ln{\left(\frac{RE}{N} + 1\right)}\Big/\ln{(p_0)}$ substitutions per generation. When selective deaths occur at only one life history

transition j, this expression applies with RE_j and N_j.

However, because each selective death reduces k for the transition at which it occurs, it reduces

reproductive excess at all other transitions, in a manner that depends on assumptions about when and how

density regulation occurs. E.g., the selective death of an adult tree creates a canopy opening into which

saplings may grow, in addition to reducing the number of seeds available to grow into it. Both these

density regulation effects dampen the degree to which selective deaths at one life history transition reduce

reproductive excess at other transitions. Reproductive excess calculations invoke N_{min} as a counterfactual

— how many losses at the focal life history transition could density regulation compensate for, holding all

else constant? There is no simple counterfactual for perturbing selective deaths at multiple transitions at

once, even when each invokes distinct, non-pleiotropic mutations. Life history trade-offs make matters

worse, because the same allele might cause selective deaths at multiple transitions, even when the allele

frequency does not change from one life cycle to the next. We thus reach the negative result that there is

no general formula for a speed limit when selection occurs during more than one life history transition —

it depends on the details of density regulation.

Reproductive excess beyond a fixed life cycle

Not all organisms proceed through the same sequence of life history transitions every cycle, e.g. budding

yeast experience a variable number of mitoses in between each meiosis, and a variable number of selfing

events between each outcrossing. In this case, we cannot take the product of an exact series of transitions.

Instead, we privilege the life history transition that produces the most severe bottleneck, assuming that the

population will spring back to vibrancy after. We define a minimum number of individuals N_{bot} who

need to make it through to the other side of the bottleneck, and define

Reproductive excess at transition $j = k_j N_j -$ min. needed to ensure N_{bot} after bottleneck

We now need to take the expectation over all possible series of life history transitions, and solve for

$N_{min,j+1}$ in

363
$$N_{bot} = \text{E}\left(N_{min,j+1} \overset{\substack{life\ history\ stages \\ between\ j\ and\ bot}}{\underset{x}{\prod}} k_{best,x} \right).$$

The precise value of N_{bot} will be informed by the ecology of the species in question. It may be small,

such as when just a modest number of new hosts, each colonized by just one infectious microorganism, is

sufficient to ensure the population's future. The appropriate value of N_{bot} is the smallest population size

that reliably escapes extinction.

Comparison to fitness

Haldane obtained selective deaths from $sN(1 - p)$, where s is the selection coefficient with respect to

per-generation relative fitness. Our k values are equivalent to absolute fitness components per life history

transition. Selective deaths can be derived from an underlying population dynamic model, without

requiring either generation or relative fitness to be defined first. E.g., in the simple case of an absolute

fitness version of the Wright-Fisher model, adults of genotype i produce an expected k_i offspring, then

die. Selective deaths per capita $k_{best} - k_{mean}$ is equal to the difference in absolute fitness. Total

population reproductive excess is $N k_{mean} - N_{min}$. By definition, $N = N_{min} * k_{best}$. Substituting $\frac{N}{k_{best}}$

for N_{min} and then dividing by N gives the per-capita reproductive excess $= k_{mean} - \frac{1}{k_{best}}$.

However, these simple relationships with fitness break down once selection occurs at multiple life history

stages. Per-generation fitness is the product of fitness components, such that when a genotype that

benefits fitness in one life history transition bears an antagonistically pleiotropic cost at another, costs and

benefits cancel out. In contrast, selective deaths never cancel out; indeed, incomplete density

compensation causes selective deaths to accrue across life history transitions. Similarly, there is no

cancelling out across generations, e.g. seasonally fluctuating selection must incur many selective deaths to

effect the large allele frequency fluctuations that have been observed around the long-term mean (Kelly

2022; Machado et al. 2021; Rudman et al. 2022).

Discrete time models such as Wright-Fisher typically emphasize the effective reproductive number rather

than the Malthusian parameter, i.e. equivalent to b/d rather than $b - d$ in a continuous time model with

births at rate b and deaths at rate d. Malthusian approaches are generally used to handle the complications

arising from multiple life history transitions (Charlesworth 1994). Our approach enables an effective

reproduction number framing in these more complex scenarios, while still avoiding the need to define one

"generation". This enables treatment of the finite nature of reproductive excess.

Comparison to travelling wave models

Desai and Fisher's (2007) Eq. 39 solves for the fitness difference or "lead" qs between the highest-fitness

"nose" with $r_{max} = 1 + qs$ and the average genotype $\bar{r} = 1$ in an asexual population, as a function of U,

N, and relative fitness difference per mutation s. The minimum reproductive excess of the best genotype

that is implicitly required to maintain constant population size is equal to the lead qs. This aligns closely

with the parameter $k - 1$ of Nei (1971) and Felsenstein (1971). The actual reproductive excess in Desai

and Fisher's (2007) model is infinite, as for all models that assume a constant population size and treat

only relative fitness.

We next calculate the proportion of deaths that are selective. The entire reproductive output of the best

genotype present, $1 + RE \geq 1 + qs$, represents non-selective deaths (or foregone fecundity) among its

offspring. Other genotypes all experience the same rate of non-selective deaths. The per-capita odds that

the next death hits the offspring of a specific average individual rather than a specific nose individual are

$1 : 1/(1 + qs)$. Consider a time interval in which an individual with the best genotype expects $1+RE$ non-

selective offspring deaths. An average parent expects $(1 + RE)(1 + qs)$ offspring deaths during this

time, 1+*RE* of which are non-selective. This makes the proportion of deaths that are selective equal to

$1 - 1/(1 + qs)$.

For a substantial range of parameters, especially with rapid adaptation with large *s, U,* and *N,* both the

minimum required reproductive excess, and the proportion of juvenile deaths that are selective, exceed

the value of 0.1 that was previously assumed for both (Figure 2). This application to the model of Desai

and Fisher (2007) helps clarify the relationship between these two properties. In this particular model,

with its explicit adults and implicit juveniles under selection, they are both functions of the lead qs, but

this need not be so simple when more complex life histories are considered. We note that the adaptation

speed has a one-to-one relationship with the proportion of deaths that are selective, whereas reproductive

excess imposes a speed limit without having a direct effect on speed.

Figure 2. As the adaptation rate goes up with increasing U, s, and N, following the multiple mutations

regime of Desai and Fisher (2007), so do the proportion of deaths selective and the minimum

reproductive excess required to sustain that rate of adaptation. Where not otherwise specified, $U = 0.001$,

$s = 0.01$, $N = 10^6$. Parameter ranges are truncated to avoid the regime $\frac{s}{U} < 3$, for which the assumptions

of the Desai and Fisher's (2007) model break down. The minimum required reproductive excess qs and

the corresponding proportion of selective deaths $1 - 1/(1 + qs)$ were calculated by numerically solving

Equation 39 for q in Desai and Fisher (2007). We note that Desai and Fisher (2007) use relative

Malthusian fitness $r = r' - \bar{r}$, rather than per-generation $w = W/\bar{W}$, but results converge for small s.

Experimental proof of concept

We measure reproductive excess and the proportion of deaths that are selective for the first time in an

empirical system using data from Exposito-Alonso et al. (2019) (Figure 3). Briefly (see Supplementary

Information for details), 517 homozygous lines of *Arabidopsis thaliana* were grown in $2 \times 2 \times 2$

environments of varying location, water availability, and adult density. For high density, thirty seeds of

the same genotype were planted per pot, and selective deaths are for the transition from seed to adult. For

low density, ~10 seeds were planted per pot, and only one seedling, chosen at random, was retained after

germination; selective deaths are for the transition from seedling to adult. Fitness differences in fecundity

were not significant (see Supplementary Results).

Figure 3. Worked example of reproductive excess for the life history transitions of *A. thaliana* in the experimental setup from Exposito-Alonso *et al* (2019). We found no evidence for between-genotype differences in fecundity and the experimental setup provides no information about seed dispersal, so we show no selective deaths during these transitions. Values of k are chosen for illustrative purposes.

The proportion of deaths that are selective exceeds 10% (Figure 4 x-axis, Supplementary Table 2). In Madrid with low water and high density, as many as 95% of deaths were selective.

Figure 4. <u>The proportion of deaths that are selective generally exceeds Haldane's 10% estimate (dotted vertical line), with ample reproductive excess (above zero, shown as dashed horizontal line) especially under high-water conditions.</u> Selective deaths shown on the x-axis apply either to seeds plus seedlings (A) or to seedlings alone (B). Reproductive excess is calculated (see Supplementary Information) for two life history transitions: seeds produced and seedlings surviving to adulthood. These are shown in red and blue, and correspond to values shown on the left and right y-axes, respectively. Note that reproductive excess of seedlings cannot exceed one per pot under low density conditions, and 30 per pot under high density conditions. Reproductive excesses are shown as a vertical range representing high and low values of seed survival during dispersal, a life history transition for which we lacked empirical data, and therefore considered a 10-fold plausible range. The proportions of deaths that are selective are adjusted for extreme value bias as shown in Supplementary Table 1. Values can be found in Supplementary Table 2.

A priori, we expect high water and low density to be more benign, which might increase reproductive

excess and/or reduce the proportion of deaths selective. While we cannot compare seedling deaths at low

density to seed plus seedling deaths at high density, these predictions are confirmed for high vs. low

water: strongly in the case of selective deaths and excess seeds, and weakly with respect to excess

seedlings (Figure 4, Supplementary Table 2). Estimated reproductive excess is fairly insensitive to the 10-

fold range we consider for the proportion of seeds that successfully disperse to suitable habitats (vertical

line length in Figure 4).

DISCUSSION

Population geneticists no longer exhibit obvious concern that Haldane's Dilemma poses a limitation to

the speed of adaptation, even when e.g. reporting high rates of adaptive sweeps (Galtier 2016; Murga-

Moreno et al. 2024; Sella et al. 2009; Uricchio et al. 2019). Here we began by reviewing the flaw in load-

based speed limit arguments; with appropriate comparisons to the best genotype currently present, rapid

adaptation is possible. However, there is a different and crucial type of limitation pointed out by Nei

(1971) and Felsenstein (1971), one that depends not on relative load but on the finite nature of

reproductive excess. Given an assumed 10% limit on the fraction of deaths that could be selective, and

allowing selection at just one life history transition, the rate of adaptation is significantly limited

(Felsenstein 1971; Nei 1971). We clarified and extended theoretical arguments regarding selective deaths

drawn from finite reproductive excess, to apply even to life histories for which a "generation" is poorly

defined, e.g. colonial organisms. Speed limits depend on when and how density regulation occurs, and

our proof-of-principle empirical analysis suggests that the fraction of selective deaths may not be limited

to 10%; these results might explain why adaptation is not significantly limited by reproductive excess.

Nei (1971) and Felsenstein (1971) solved for a hypothetical equilibrium where all 'effective' reproductive

excess was converted to selective deaths. They thus demonstrated that finite reproductive excess imposed

a speed limit, without referring to lag load. Our more complete treatment emphasizes that reproductive

excess varies by life history stage, and that the degree to which selective deaths during one life history

transition reduce reproductive excess at other transitions depends on how and when density regulation

occurs. While these theoretical results do not provide a general expression for a speed limit to adaptation,

they uphold the previous special case, which is compatible with some populations in some cases having

limits on their rate of adaptation.

We apply the concepts of reproductive excess and selective deaths to empirical data for the first time.

This proof-of-principle application was sufficient to suggest a novel possibility: Haldane's 10% limit to

selective deaths may be too low. The smallest proportion of selective deaths we observed across 8

environmental conditions was 8.5%, while in the more adverse environmental conditions within this

experiment, 95% of deaths were selective. Relaxing this auxiliary assumption about a critical parameter

value resolves Haldane's concerns. Future work under more natural conditions (e.g. with higher extrinsic

mortality) and in different species (e.g. less fecund) remains necessary to reach the conclusion that the

proportion of deaths that are selective is typically high. Our theoretical framework is flexible enough to

be customized for any species, using whichever life history transitions best describe that species' life

history. Uses might include understanding the limits to adaptation given high fecundity (Hutchings 2000).

We note that the method of Ewens (1970) might allow estimates of V_G for per-generation fitness,

combined with population size, to be repurposed to estimate the proportion of deaths selective.

Empirical demonstration of these concepts makes the concepts more concrete. One contribution of our

current work is simply to clarify the variety of lines of reasoning that produce limits on the rate of

adaptation. Our more specific theoretical and empirical analyses then develop a line of reasoning about

reproductive excess and selective deaths that has been neglected. The attention of 'creation science' to the

question of Haldane's dilemma (Remine 2005, 2006) highlights the importance of clearly resolving it.

Different papers use the term "substitutional load", which we avoid here, to mean very different things,

and the resulting confusion in terminology has obscured the consequences of formulating Haldane's

dilemma in different ways. 'Substitutional load' has been used to refer to what we here call the lag load

(Kimura and Ohta 1971; Maynard Smith 1976), the cost of selection (Kimura 1968), the number of

offspring that the most fit genotype must produce (Ewens 2004), the sum of lag load across all

generations involved in a substitution (Kimura 1960; Nei 1971), and even more broadly to refer to

variance-based rather than load-based arguments when made in the context of similar questions (Ewens

1970). The latter highlights standard modes of reasoning in genetics, especially quantitative genetics,

which follow Fisher (1930) to focus on variances — sums of differences squared — while selective

deaths and reproductive excess are, like load, both differences, with no square operation.

The optimal genotype re lag load is normally defined, as Haldane did, with respect to the best genotype

that can be constructed from segregating alleles. This is because a standard relative fitness model makes

environmental change implicit. When fitness is defined in absolute terms, the optimal genotype used to

define lag load can also include adaptive mutations that have not yet appeared in the population (Bertram,

Gomez, and Masel 2017; Osmond and Klausmeier 2017). While speed limits cannot threaten the

persistence of a species adapting in a constant environment, real species face rapidly changing

environments (biotic and abiotic) that can threaten population persistence. Even in a model of absolute

fitness, large lag load need not be a problem, although it must be stable rather than growing. I.e., for a

population to persist, adaptation must keep up with the speed of environmental change.

It is interesting that Ewens' (1970) key concept of relative load was later reinvented as the "lead", as part

of calculations that derived the actual speed of adaptation v (rather than limits to it) from the beneficial

mutation rate U, the population size N, and the per-mutation selection coefficient s (Desai and Fisher

2007). One reason this solution was not available to Haldane or Ewens was that population genetics had

not yet begun to treat origination processes (McCandlish and Stoltzfus 2014). Instead of treating a steady

input of beneficial new mutations, Haldane considered a scenario in which environmental change

activates beneficial variants within standing genetic variation. Indeed, a variant's initial frequency

$-\ln(p_0)$ is the primary factor in determining Haldane's maximum speed of adaptation. Some adaptation

comes not from activation of standing genetic variation, but from *de novo* mutations each appearing at

initial frequency $1/N$ or $1/2N$. A lead-based approach was used for *de novo* mutations to derive the rate

of beneficial sweeps in asexuals as $\frac{2s\ln[Ns]}{\ln^2[S/U]}$ for the simple case of constant *s* (Desai and Fisher 2007), for

parameter ranges in which the previously derived rate UNs does not apply. Here we relaxed the

assumption of infinite reproductive excess made by this relative fitness model, and calculated the minimal

reproductive excess required for the model to hold. We also reveal the model's implied fraction of deaths

(or foregone fecundity) that are selective. Both quantities are functions of the lead.

Yet another approach, starting with Kimura (1961), is to use the framework of information theory to set

bounds on the speed of adaptation. Natural selection increases the information stored in genomes (Adami

2012). Kimura calculates the amount of information acquired per sweep in terms of p_0 and then relates

this to the cost of selection using Haldane's equation that $D = -\ln(p_0)$ (Kimura 1961). More recent

approaches treat the bounds placed on the accumulation of information in much more detail, while

treating either the Malthusian parameter (McGee et al. 2022) or classic discrete time relative fitness

(Hledík, Barton, and Tkačik 2022). Both these approaches define an information "cost", but this is not

equal to our cost in terms of selective deaths.

Like Haldane (1957), Nei (1971), and Felsenstein (1971), we have neglected complexities from linkage

disequilibrium, which will typically make the conversion of selective deaths to adaptation less efficient.

Here we considered both the independent sites extreme, and the other extreme in the form of the asexual

model of Desai and Fisher (2007). Other limits to adaptation speed can be imposed by genetic map length

(Neher 2013; Weissman and Barton 2012; Weissman and Hallatschek 2014).

Excitingly, unlike most approaches in evolutionary genetics, the approach we describe does not require

the quantity 'fitness', which is deceptively difficult to define in a manner that can be generalized to all

circumstances (Ariew and Lewontin 2004; Bertram and Masel 2019; Doebeli, Ispolatov, and Simon 2017;

Smith et al. 2024; Van Valen 1989). Standard quantitative definitions of either relative or absolute

'fitness' require a clear definition of a 'generation' over which change is assessed, which in turn requires

a clear definition of an 'individual' whose life cycle a generation captures (Wilson and Barker 2021). Our

generalized selective deaths approach is derived from selection but does not require 'fitness' to be

defined. Rather, we measure selective deaths from pairwise differences in fecundity and survival between

each genotype and the best genotype present, during each life history transition. Reproductive excess

corresponding to that life history transition indicates how stringent selection can be, without triggering a

decline in population size. This approach applies at each life history transition and can therefore be

generalized to species with complex life histories, where it becomes difficult to define a 'generation' and

therefore fitness.

GLOSSARY

Absolute fitness of an adult: expected number of individual offspring surviving to adulthood (defined

here for hermaphrodite species — half this value, if reproducing sexually with males).

Absolute fitness of a juvenile: expected number of individual offspring (or half this value, if reproducing

sexually with males). Failure to survive to adulthood (reproductive maturity) implies zero offspring.

Adult: Reproductively mature individual. More than one adult life history stage may be defined.

Cost of selection: The number of selective deaths that must occur over time to accomplish defined

evolutionary change, e.g. to complete a single selective sweep.

Generation: A set of life history transitions that ends the first time it returns, with new individuals, to the

same life history stage (e.g. adult or juvenile) where it began.

Individual: An organism that meets a loosely defined set of criteria (Wilson and Barker 2021), including

a shared genome, and the degree of integration of parts. Whether e.g. a group of microbes is a closely

connected ecological community vs. an individual organism may be a matter of biological judgment.

Juvenile: Individuals that are not yet reproductively mature. More than one juvenile life history stage

may be defined, e.g. before vs. after dispersal.

Lag load: $\frac{W_{optimal}-\overline{W}}{W_{optimal}}$ where $W_{optimal}$ is a theoretical optimal genotype that might not be present in the

population and $\overline{W}$ is the average genotype present.

Lead: $\frac{W_{optimal}-\overline{W}}{\overline{W}}$ where $W_{optimal}$ is the best genotype present in the population and $\overline{W}$ is the average

genotype present.

Life history transition: Survival (i.e. persistence of an individual), reproduction (i.e. generation of new

individuals) and/or organismal growth from one life history stage to the next

Load: A difference in fitness between an actual genotype or population and a reference. See "lag load"

and "lead" as concrete examples.

Relative fitness: expected relative genetic contribution to the next generation.

Reproductive excess: The degree to which a hypothetical population of the optimal genotype would

conclude a life history transition with a larger population than the minimum required to complete a life

history cycle without the population shrinking in size.

Selective deaths: The subset of deaths (or foregone fertility) that contributes to selective changes in allele

frequency. This can be quantified as how many deaths each genotype experiences that would not have

been experienced if that genotype were replaced by the best genotype.

DATA AVAILABILITY

All analysis was performed in R, and our code is available on GitHub:

www.github.com/josephmatheson/selective_deaths.

ACKNOWLEDGEMENTS

This work was supported by the John Templeton Foundation [62028] and [62220]. We thank Taylor

Kessinger for sparking the last author's original interest in this topic many years ago by presenting

Haldane's 1957 paper in journal club, and Jason Bertram, Ryan Gutenkunst and Ulises Hernández for

helpful discussions. We thank Joe Felsenstein for his helpful review of a previous version of this

manuscript, which helped us understand the differences in interpretation between his paper and Masatoshi

Nei's, as well as Sally Otto and three other anonymous reviewers.

AUTHOR CONTRIBUTIONS

Initial conception of the paper by Joanna Masel and Moises Exposito-Alonso, with multiple rounds of

reconception by Joseph Matheson and Joanna Masel. Design and interpretation by Joseph Matheson and

Joanna Masel. Data and details of experimental methodology provided by Moises Exposito-Alonso. Data

analysis mostly by Joseph Matheson, with the MCMCglmm model analyzed by Moises Exposito-Alonso.

Manuscript first drafted by Joseph Matheson and substantively revised many times by all authors.

CONFLICT OF INTEREST

The authors declare no conflicting interests.

BIBLIOGRAPHY

Adami, Christoph. 2012. "The Use of Information Theory in Evolutionary Biology." *Annals of the New*
York Academy of Sciences 1256(1):49–65. doi: 10.1111/j.1749-6632.2011.06422.x.

Ariew, André, and R. C. Lewontin. 2004. "The Confusions of Fitness." *British Journal for the Philosophy*
of Science 55(2):347–63. doi: 10.1093/bjps/55.2.347.

Bell, Graham. 2017. "Evolutionary Rescue." *Annual Review of Ecology, Evolution, and Systematics*
48:605–27. doi: 10.1146/annurev-ecolsys-110316-023011.

Bertram, J., K. Gomez, and J. Masel. 2017. "Predicting Patterns of Long-Term Adaptation and Extinction
with Population Genetics." *Evolution*. doi: 10.1111/evo.13116.

Bertram, Jason. 2021. "Allele Frequency Divergence Reveals Ubiquitous Influence of Positive Selection
in Drosophila." *PLOS Genetics* 17(9):e1009833. doi: 10.1371/journal.pgen.1009833.

Bertram, Jason, and Joanna Masel. 2019. "Density-Dependent Selection and the Limits of Relative
Fitness." *Theoretical Population Biology* 129:81–92. doi: 10.1016/j.tpb.2018.11.006.

Bitter, M., S. Greenblum, S. Rajpurohit, A. Bergland, J. Hemker, N. Betancourt, S. Tilk, S. Berardi, H.
Oken, P. Schmidt, and D. Petrov. 2024. "Pervasive Fitness Trade-Offs Revealed by Rapid
Adaptation in Large Experimental Populations of Drosophila Melanogaster." *Biorxiv*. doi:
10.1101/2024.10.28.620721.

Bull, JJ, R. Sanjuán, and CO Wilke. 2007. "Theory of Lethal Mutagenesis for Viruses." *Journal of
Virology* 81(6):2930–39. doi: 10.1128/jvi.01624-06.

Charlesworth, Brian. 1994. *Evolution in Age-Structured Populations*. 2nd ed. Cambridge University
Press.

Crow, James F. 1958. "Some Possibilities for Measuring Selection Intensities in Man." *Human Biology*
30(1):1–13.

Desai, Michael M., and Daniel S. Fisher. 2007. "Beneficial Mutation-Selection Balance and the Effect of
Linkage on Positive Selection." *Genetics* 176(3):1759–98. doi: 10.1534/genetics.106.067678.

Doebeli, Michael, Yaroslav Ispolatov, and Burt Simon. 2017. "Towards a Mechanistic Foundation of
Evolutionary Theory." *ELife* 6:1–17. doi: 10.7554/elife.23804.

Draghi, Jeremy A., Joel W. McGlothlin, and Holly K. Kindsvater. 2024. "Demographic Feedbacks during
Evolutionary Rescue Can Slow or Speed Adaptive Evolution." *Proceedings of the Royal Society B:
Biological Sciences* 291(2016). doi: 10.1098/rspb.2023.1553.

Ewens, Warren. 1970. "Remarks on the Substitutional Load." *Theoretical Population Biology* 139:129–
39.

Ewens, Warren. 2004. *Mathematical Population Genetics*. 2nd ed. New York: Springer US.

Exposito-alonso, Moises, Oliver Bossdorf, Rasmus Nielsen, and Detlef Weigel. 2019. "Natural Selection
on the Arabidopsis Thaliana Genome in Present and Future Climates." *Nature* 573(7772):126–29.
doi: 10.1038/s41586-019-1520-9.

Felsenstein, Joseph. 1971. "On the Biological Significance of the Cost of Gene Substitution." *American
Naturalist* 105(941):1–11.

Fisher, Ronald. 1930. *The Genetical Theory of Natural Selection*. Oxford: Clarendon Press.

Galeota-Sprung, Benjamin, Paul Sniegowski, and Warren Ewens. 2020. "Mutational Load and the
Functional Fraction of the Human Genome." *Genome Biology and Evolution* 12(4):273–81. doi:
10.1093/gbe/evaa040.

Galtier, Nicolas. 2016. "Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis." *PLoS Genetics* 12(1):1–23. doi: 10.1371/journal.pgen.1005774.

Haldane, JBS. 1954. "The Measurement of Natural Selection." *Proceedings of the Ninth International Congress of Genetics, Supplement to Caryologia* 480–87.

Haldane, JBS. 1956. "The Relation between Density Regulation and Natural Selection." *Proceedings of the Royal Society of London. Series B-Biological Sciences* 145(920):306–8.

Haldane, JBS. 1957. "The Cost of Natural Selection." *Genetics* 55(3):511–24.

Hledík, Michal, Nick Barton, and Gašper Tkačik. 2022. "Accumulation and Maintenance of Information in Evolution." *Proceedings of the National Academy of Sciences of the United States of America* 119(36):1–10. doi: 10.1073/pnas.2123152119.

Hutchings, Jeffrey A. 2000. "Numerical Assessment in the Front Seat, Ecology and Evolution in the Back Seat:: Time to Change Drivers in Fisheries and Aquatic Sciences?" *Source: Marine Ecology Progress Series* 208:299–303. doi: 10.2307/24863827.

Kelly, John. 2022. "The Genomic Scale of Fluctuating Selection in a Natural Plant Population." *Evolution Letters* 506–21. doi: 10.1002/evl3.308.

Kern, Andrew D., and Matthew W. Hahn. 2018. "The Neutral Theory in Light of Natural Selection." *Molecular Biology and Evolution* 35(6):1366–71. doi: 10.1093/molbev/msy092.

Kimura, Motoo. 1960. "Optimum Mutation Rate and Degree of Dominance as Determined by the Principle of Minimum Genetic Load." *Genetics* 57(1):21–34.

Kimura, Motoo. 1961. "Natural Selection as the Process of Accumulating Genetic Information in Adaptive Evolution." *Genetical Research* 2(1):127–40. doi: 10.1017/S0016672300000616.

Kimura, Motoo. 1968. "Evolutionary Rate at the Molecular Level." *Nature* 217:624–26.

Kimura, Motoo. 1969. "The Number of Heterozygous Nucleotide Sites Maintained in a Finite Population Due to Steady Flux of Mutations." *Genetics* 61(4):893–903.

Kimura, Motoo, and James F. Crow. 1969. "Natural Selection and Gene Substitution." *Genetical Research* 13(2):127–41. doi: 10.1017/S0016672300002846.

Kimura, Motoo, and Tomoko Ohta. 1971. "On the Rate of Molecular Evolution." *Journal of Molecular Evolution* 1(1):1–17. doi: 10.1007/BF01659390.

Machado, Heather E., Alan O. Bergland, Ryan Taylor, Susanne Tilk, Emily Behrman, Kelly Dyer, Daniel K. Fabian, Thomas Flatt, Josefa González, Talia L. Karasov, Bernard Kim, Iryna Kozeretska, Brian P. Lazzaro, Thomas J. S. Merritt, John E. Pool, Katherine O'brien, Subhash Rajpurohit, Paula R. Roy, Stephen W. Schaeffer, Svitlana Serga, Paul Schmidt, and Dmitri A. Petrov. 2021. "Broad Geographic Sampling Reveals the Shared Basis and Environmental Correlates of Seasonal Adaptation in Drosophila." *ELife* 10. doi: 10.7554/eLife.67577.

Maynard Smith, J. 1968a. "'Haldane's Dilemma' and the Rate of Evolution." *Nature* 219(5159):1114–16. doi: 10.1038/2191114a0.

Maynard Smith, J. 1968b. "'Haldane's Dilemma' and the Rate of Evolution." *Nature* 219(5159):1114–16. doi: 10.1038/2191114a0.

Maynard Smith, J. 1976. "What Determines the Rate of Evolution?" *American Naturalist* 110(973):331–38.

McCandlish, David M., and Arlin Stoltzfus. 2014. "Modeling Evolution Using the Probability of Fixation: History and Implications." *Quarterly Review of Biology* 89(3):225–52. doi: 10.1086/677571.

McGee, Ryan Seamus, Olivia Kosterlitz, Artem Kaznatcheev, Benjamin Kerr, and Carl T. Bergstrom. 2022. "The Cost of Information Acquisition by Natural Selection." *Biorxiv* 2022.07.02. doi: 10.1101/2022.07.02.498577.

Murga-Moreno, Jesús, Sònia Casillas, Antonio Barbadilla, Lawrence Uricchio, and David Enard. 2024. "An Efficient and Robust ABC Approach to Infer the Rate and Strength of Adaptation." *G3: Genes, Genomes, Genetics* 14(4). doi: 10.1093/G3JOURNAL/JKAE031.

Neher, Richard A. 2013. "Genetic Draft, Selective Interference, and Population Genetics of Rapid Adaptation." *Annu. Rev. Ecol. Evol. Syst* 44:195–215. doi: 10.1146/annurev-ecolsys-110512-135920.

Nei, Masatoshi. 1971. "Fertility Excess Necessary for Gene Substitution in Regulated Populations." *Genetics* 68:169–84.

Nicholson, A. J. 1933. "Supplement: The Balance of Animal Populations." *Journal of Animal Ecology* 2(1):131–78.

Osmond, Matthew M., and Christopher A. Klausmeier. 2017. "An Evolutionary Tipping Point in a Changing Environment." 71(12):2930–41. doi: 10.2307/48575242.

Remine, Walter. 2006. "More Precise Calculations of the Cost of Substitution." *Creation Research Science Quarterly* 43(September).

Remine, Walter J. 2005. "Cost Theory and the Cost of Substitution — a Clarification." *Journal of Creation* 19(1):113–25.

Rudman, Seth M., Sharon I. Greenblum, Subhash Rajpurohit, Nicolas J. Betancourt, Jinjoo Hanna, Susanne Tilk, Tuya Yokoyama, Dmitri A. Petrov, and Paul Schmidt. 2022. "Direct Observation of Adapting Tracking on Ecological Time Scales in Drosophila." *Science* 375(6586). doi: 10.1126/science.abj7484.

Sella, Guy, Dmitri A. Petrov, Molly Przeworski, and Peter Andolfatto. 2009. "Pervasive Natural Selection in the Drosophila Genome?" *PLoS Genetics* 5(6):e1000495. doi: 10.1371/journal.pgen.1000495.

Smith, Daniel, Guilhem Doulcier, Pierrick Bourrat, Peter Takacs, and Joanna Masel. 2024. "Why There Are so Many Definitions of Fitness in Models." *Ecoevorxiv*. doi: 10.32942/X2V61T.

Uricchio, Lawrence H., Dmitri A. Petrov, and David Enard. 2019. "Exploiting Selection at Linked Sites to Infer the Rate and Strength of Adaptation." *Nature Ecology & Evolution* 3(6):977–84. doi: 10.1038/s41559-019-0890-6.

Van Valen, Leigh. 1963. "Haldane's Dilemma, Evolutionary Rates, and Heterosis." *American Naturalist* 97(894):185–90.

Van Valen, Leigh. 1989. "Three Paradigms of Evolution." *Evolutionary Theory* 9(2):1–17.

732   Weissman, Daniel B., and Nicholas H. Barton. 2012. "Limits to the Rate of Adaptive Substitution in
733   Sexual Populations." *PLoS Genetics* 8(6). doi: 10.1371/journal.pgen.1002740.

734   Weissman, Daniel B., and Oskar Hallatschek. 2014. "The Rate of Adaptation in Large Sexual Populations
735   with Linear Chromosomes." *Genetics* 196(4):1167–83. doi: 10.1534/genetics.113.160705.

736   Wilson, Robert A., and Matthew J. Barker. 2021. "Biological Individuals" edited by E. N. Zalta. *Stanford
737   Encyclopedia of Philosophy* 2021.