Substitution load revisited: reproductive excess, the proportion of deaths that are selective, and adaptation's speed limit

Joseph Matheson[1,2], Moises Exposito-Alonso[3], Joanna Masel[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona 85721, USA

[2]Department of Ecology, Behavior, and Evolution, University of California San Diego, San Diego, CA, 92093, USA

[3]Departments of Plant Biology & Global Ecology, Carnegie Institution for Science, Stanford University, Stanford, California 94305, USA

Corresponding author: Joanna Masel, 1041 E Lowell St, Tucson AZ 85721, USA. masel@arizona.edu

Article Type: Letter

ABSTRACT

Haldane's Dilemma refers to the concern that the need for many "selective deaths" to complete a substitution (i.e. selective sweep) creates a speed limit to adaptation. However, discussion of this concern has been marked by confusion over which features of adaptation produce speed limits, what those limits are, and the consequences of violating speed limits. The term "substitution load" has been particularly unhelpful in this regard. Here we distinguish different historical lines of reasoning that lead to speed limits. We identify one line of reasoning, focused on finite reproductive excess, that has not yet been fully addressed. We develop this into a more general theoretical model that can apply to populations with any life history, even those for which a generation or even an individual are not well defined. As proof of principle, we apply this model to a dataset measuring survival of 517 different genotypes of *Arabidopsis thaliana* grown in eight different environmental conditions. These data suggest highly permissive limits to the speed of adaptation in all environmental conditions. This is because a much higher proportion of deaths contribute to adaptation than were anticipated during historical discussions of speed limits.

Keywords: cost of selection, adaptation rate, genetic load, fitness component, biological individual

LAY SUMMARY

Neutral theory was predicated on theoretical arguments that adaptation is subject to a speed limit. We

resolve confusions regarding historical speed limit arguments, which depend on differences in fitness, not

variance (differences in fitness squared). We generalize the underlying concepts of selective deaths and

reproductive excess to populations with any life cycle, even those for which an "individual" and hence

generation and fitness, are poorly defined. We apply the revised theory to Arabidopsis data,

demonstrating the potential for future related experiments.

INTRODUCTION

During an adaptive sweep, new alleles need to be substituted for old alleles across an entire population.

This means that all individuals with the old alleles need to leave no descendants, and individuals with

new alleles must produce enough offspring to replenish the population. These requirements put a limit on

the speed at which sweeps can happen, which could be prohibitive if many sweeps need to occur quickly.

Haldane (1957) used this reasoning to propose a rough estimate of the maximum speed at which sweeps

could accumulate. This speed limit, later known as Haldane's dilemma (Van Valen 1963) motivated the

development of Kimura's (Kimura 1968) neutral theory. However, the underlying logic has been

challenged on multiple counts (Ewens 1970; Felsenstein 1971; Kern and Hahn 2018; Maynard Smith

1968). In these discussions, conceptually distinct approaches to quantifying the issue are often described

using identical terms, which apart from being confusing, leaves unresolved the critical question: what is

the upper limit on the speed of adaptation, and does it matter for natural populations?

Here we first synthesize the historical literature, drawing out several key quantities (see Results). First, a

population has a "reproductive excess", meaning how many individuals at a given life history stage are in

excess of the minimum number required to avoid a decline in population size over the next life cycle (see

Glossary for definitions of the terms used here). The second key quantity is the "selective deaths"

(including foregone fertility) required for selection to effect a change in allele frequency. These can be

quantified as the degree to which the mean individual survives and reproduces worse than the best

genotype present. The "cost of selection" is the number of selective deaths required to achieve a given

adaptation rate. Haldane (1957) assumed, without evidence, that at most 10% of deaths were selective, an

assumption repeated by all subsequent work.

Haldane (1957) confused the cost of selection, which is related to differences between the mean

individual and the best genotype present, with the lag load, which compares to an ideal genotype that is

unlikely to exist (Ewens 1970). However, Nei (1971) and Felsenstein (1971) derived a near-identical

speed limit to Haldane's, without this flaw, by examining the requirement for selective deaths within a

model in which reproductive excess is finite.

Explicit consideration of reproductive excess means abandoning relative fitness models, which consider

only one life history stage with adult population size N, with an implied infinite number of juveniles

(Bertram and Masel 2019). Models of finite reproductive excess need at least two life history stages:

adults and juveniles, where the latter shows reproductive excess relative to the former. Selective deaths

are then relative not to a single adult population size N as in Haldane's (1957) model, but to a

denominator describing the population size at the appropriate life history stage (Kimura and Crow 1969).

An emphasis on life history transitions rather than generations is a strength rather than a weakness of the

selective deaths view. One of the many flaws of the concept of "fitness" (Van Valen 1989) is the

difficulty of defining a "generation" for many species, especially colonial species for which an

"individual" is not well defined (Wilson and Barker 2021). Consider for example the budding yeast

Saccharomyces cerevisiae. Is each mitotic division a generation? Or each life cycle spanning from

meiosis to meiosis, with variable number of mitoses in between? Or the span between outcrossing events,

with variable occurrences of selfing as well as mitoses in between? Or is a generation best defined

ecologically with respect to dispersal between resources that allow growth? Problems defining a

generation arise for a broad range of species (albeit not humans, nor many other animal species), but are

resolved when population dynamics are viewed as a series of life history transitions. The "generation"

that matters in this view is not the concept of one complete life cycle, but rather the "generation" of

reproductive excess, in contrast to other life history transitions that involve survival rather than

reproduction.

After synthesizing the literature, here we reformulate and generalize Nei's (1971) and Felsenstein's

(1971) ideas to selection on both fecundity and survival, to life cycles with selection at more than one

stage, and to life cycles with a variable number of stages. We clarify the concepts of reproductive excess

and selective deaths, and use our general theory to pose two empirically accessible questions. First, how

much reproductive excess does a genotype or population produce, beyond what is needed to avoid

decline? Second, what fraction of deaths are selective (and how does this compare to the 10% guess made

by Haldane)? Posing questions in this form allows us to make the first empirical estimates with which to

ground Haldane's approach. We use data from Exposito-Alonso et al. (2019), who counted or estimated

every plant grown and seed produced of *A. thaliana* cultivars from 517 different genotypes in one season,

under 8 distinct environmental conditions. These data are not representative of natural conditions, but

they suffice to illustrate how such an analysis can be done. Ours is the first direct application of Haldane's

selective death arguments to empirical data, representing proof of principle.

METHODS

Environmental conditions

We re-analyze the data of Exposito-Alonso et al. (2019), who used a $2 \times 2 \times 2$ design for environmental

conditions, with the three treatments being climate, water availability, and adult density (see github for

raw data and analysis code). We treat these as eight separate populations. For climate, plants were grown

in outdoor field stations in either Tübingen, Germany (near the center of the species range of *A. thaliana*)

or in Madrid, Spain (at the southern edge of the range). Plants were all artificially watered. The high-

water treatment matched soil moisture levels near the station in Germany, and the low-water treatment

matched soil moisture levels near the station in Spain. To generate high adult density, thirty seeds of the

same genotype were planted per pot. For low density, several seeds (~10) were planted per pot, enough to

ensure that at least one seed would germinate, but few enough that the seeds were unlikely to inhibit each

other pre-germination. To avoid any competition between adult plants in the low density treatment, only

one seedling, chosen at random, was retained after germination, and the rest were plugged out and

discarded. We refer to each treatment with a three-letter abbreviation: M or T for Madrid or Tübingen, L

or H for low or high water, and I or P for a single individual plant or a population of thirty plants per pot.

For example, the treatment with thirty seeds per pot grown in Madrid with high water is abbreviated as

MHP.

Within each of the eight populations, seeds from 517 fully homozygous plant genotypes (taken from a

parental generation grown under controlled conditions to control for parental effects) were grown in pots

that included only plants of that genotype. The number of replicate pots per genotype per population was

occasionally as few as one due to experimental losses, but mostly ranged between five to seven replicates.

Our interest is in differences among genotypes, not among replicates. We therefore calculate key

quantities of interest for each genotype-environment combination by averaging across replicates.

Selective deaths

The experiment can be mapped reasonably easily onto theoretical treatments. In each environmental

treatment, a starting population of seeds grows into adult plants, experiencing both selective deaths and

non-selective deaths as they proceed from seeds to seedlings to adults. Plants which survive to become

adults then produce k seeds on average, some of which would normally constitute the next generation,

although the experiment concludes at the end of season. The experiment does not capture the life history

stage of seed dispersal to fertile ground, to complete the life cycle that began with seeds planted in a pot.

Juvenile deaths must be treated differently for the low- and high-density treatments. In the low-density

treatment, where exactly one seedling is retained after germination, we do not have access to data on

selective seed deaths, and so consider only seedling selective deaths. In the high-density treatment, any of

the thirty seeds that fail to survive to the end of the experiment are counted as deaths, whether due to seed

death before germination or to subsequent seedling death; our selective death calculations do not

differentiate between these two life history transitions. This means that across the two life history

transitions at which juvenile plants can die (as planted seeds before germination and as seedlings), only

one set of juvenile deaths is recorded in each density treatment, but they are not comparable. They are

combined seed and seedling deaths in the high-density case, and seedling deaths alone in the low-density

case. Histograms are shown in Supplementary Figures 1-2.

In each treatment, we score the observed juvenile death rate of the highest performing genotype as the

baseline extrinsic mortality for all genotypes (i.e. as non-selective deaths). Conceptually (ignoring a

correction for extreme value bias that is treated in the Supplement), for each life history transition in each

environmental condition we have:

$$\text{Selective deaths in the population } = \sum_i n_i(d_i - d_{best})$$

136   where n_i is the starting population of genotype i at that life history transition, d_i is the genotype's average

137   death rate during that life history transition, and d_{best} is the average death rate of the genotype with the

138   lowest death rate for that life history transition in that environment.

Test for genetic variance in fecundity

Selective "deaths" can also be defined for unrealized fecundity. We did not analyze this here, because of

lack of evidence for significant genetic differences in fecundity. In support of this, we performed an

ANOVA test on fecundity in each environmental condition. We only have information on fecundity as an

aggregate per replicate pot (rather than per individual plant in the high-density condition), so we compare

among-genotype variance to among-replicate variance. Note, however, that we remove replicates that had

no adults surviving to reproductive maturity, as well as genotypes with only a single replicate pot with

surviving adults (and therefore no way of estimating among-replicate variance). All surviving adults

produced at least some seeds. We Box-Cox transformed the data for each pot with surviving adults in

each environmental condition (see Supplementary Figure 3 for post-transform histograms) before

performing the ANOVA.

Proportion of juvenile deaths selective

Because *A. thaliana* is an annual plant, all juveniles will die by the end of the season, whether as selective

deaths during the experiment, non-selective deaths during the experiment, or non-selective deaths after

the end of the experiment. From this we obtain, for each environmental condition:

$$\text{Fraction of juvenile deaths selective} = \frac{Selective\ deaths\ of\ juveniles}{Starting\ population}$$

In the high-density populations, the starting population is 30 seeds per pot. In the low-density

populations, the starting population is 1 seedling per pot.

Pairwise genotype comparisons

For every possible pair of genotypes, we repeat the analysis above to estimate selective deaths and the

proportion of deaths which are selective, using the better genotype of the pair as the 'best' genotype in the

calculation of selective deaths. With only two genotypes, we do not adjust for extreme value bias. Using

whole-genome information, we calculated the total number of SNP differences between each pair

(Hamming distance, number of allele differences out of 1,353,386 biallelic SNPs) using PLINK v1.9.

RESULTS

Synthetic historical review

Haldane made two somewhat different arguments in his seminal 1957 paper, muddying the waters from

the beginning. In the first argument, he defined "selective deaths" as the subset of deaths $s(1-p)N$ that

contribute to a change in the allele frequency p, where s is the selection coefficient. This was on the basis

that the $(1 - p)N$ individuals that lack a beneficial mutation experience s more deaths than they would if

they had the mutation, and those extra deaths are required for selection to have its effects. Note that

reduced fecundity is mathematically equivalent to increased mortality, and selective "deaths" can thus

result from losing potential offspring, not just literal deaths.

Haldane defined the "cost of selection" as the number of selective deaths occurring during a substitution

(i.e. a selective sweep from low allele frequency to fixation). He calculated this cost as the integral of

$s(1 - p)N$ over the course of a sweep from allele frequency $p = p_0$ to close to 1 (Figure 1A). In a

haploid population of constant size N, one sweep requires $N \times D$ selective deaths, where $D =$

$-\ln(p_0)+O(s)$. For appropriately small s and p_0 (Haldane suggests $s < \frac{1}{3}$ and $p_0 = 10^{-4}$), the first term

dominates, making D nearly independent of the selection coefficient. For alternative assumptions about

ploidy, dominance, and degree of inbreeding, D is a different function of p_0, but s remains unimportant

unless close to 1 (Haldane 1957). In a representative case of $p_0 = 10^{-4}$ at a diploid autosomal locus with

no dominance, $D = 18.4$. Haldane conservatively estimated that 20-30N selective deaths are likely to be

typical for a sweep.

Haldane's second argument about adaptation rate limitations relied on load calculations. Load is a

reduction of a population's fitness relative to a reference optimal genotype (Figure 1B). Haldane

considered x loci independently undergoing sweeps, such that the current allele frequency at the i^{th} locus

reduces population fitness by a factor of $1 - d_i$ relative to its post-sweep value. The fitness of the

population is then lower than that of a hypothetical perfect population by a factor of $\prod_{i=1}^{x}(1 - d_i) \approx$

$e^{-\sum_{i=1}^{x} d_i}$. Haldane claimed that this load relative to an ideal genotype implies that the fraction of deaths

that are selective is $\sum d_i$. This was incorrect; the better reference point is the best genotype actually

present in the population. With 30N selective deaths required to complete each independent selective

sweep, and N deaths available per generation, Haldane obtained an average spacing between fixation

events $n \geq \frac{30}{\sum d_i}$ generations.

Haldane's fitness reduction relative to an ideal genotype was later named lag load (Maynard Smith 1976),

inspired by lagging adaptation to a changing environment, where new mutations are required to keep up

(Bertram, Gomez, and Masel 2017). (This is distinct from "evolutionary rescue" (Bell 2017) from

imminent population decline, without asking whether a similar disastrous scenario will promptly recur.)

Lag load can be defined even in a static environment, where innovative new adaptive alleles reveal the

possibility of an even better optimal genotype. The size of a lag load is not important *per se*; what matters

is that it is stable rather than growing. While speed limits do not threaten the persistence of a species

adapting in a constant environment, real species do face rapidly changing environments (biotic and

abiotic) that can threaten population persistence. I.e., for population persistence, the speed of adaptation

must keep up with the speed of environmental change.

Haldane argued *a priori* that species could probably only sustain about 10% selective deaths (which he

incorrectly equated with 10% lag load) for any serious length of time. From this assumption, he derived a

speed limit of around one sweep every 300 generations, later called "Haldane's dilemma" (Van Valen

1963). All subsequent authors have continued to assume a 10% figure.

The fact that there are so many amino acid substitutions, each requiring a sweep, was the original

evidence supporting neutral theory (Kimura 1968). Kimura and Ohta (1971) plugged in estimates of the

actual rate of substitution in mammalian lineages as n in Haldane's equation $L = e^{-30/n}$, which produced

what they considered to be an excessively large lag load. Although still a lag load argument, their

argument was subtly different from Haldane's, arguing that a high lag load implies that typical individuals

would need to have a biologically implausible fraction of their offspring die (Kimura and Ohta 1971).

Ewens (1970) pointed out that Haldane's and Kimura's load arguments improperly use the lag load

(comparison to an ideal genotype) to calculate selective deaths, where they should have mean fitness

relative to the most fit individual present. In a population with many sweeps occurring at once, the

likelihood that even a single individual has the ideal combination of alleles is vanishingly small (Figure

1B). More recent travelling wave theories have rediscovered the importance of this relative type of load,

and named it the "lead" (Desai and Fisher 2007).

Prior to modern travelling wave theory, approximations for the lead were derived from variance in fitness

(Ewens 1970; Kimura 1969). In the case of many independent sweeps at once, variance in fitness (after

normalizing mean population fitness as 1) is approximately s/n, where s is the selection coefficient of an

adaptive allele and n is the number of generations between fixation events (Ewens 1970). The fittest

genotype likely to be present can be estimated using the statistics of extreme values. E.g., for a population

of size 10^6, the most extreme fitness value likely to appear is around 4.9 standard deviations above the

mean (Ewens 1970). Using Haldane's 10% as an estimate of the lead (instead of the lag load) yields

$4.9\sqrt{s/n} = 0.1$. For $s = 0.01$, n is around 20, much less than Haldane's estimate of 300, and n is lower

still for lower s. In other words, Ewens (1970) found that many simultaneous sweeps do not imply an

implausibly large lead, and the corresponding speed limit of $n \approx 20$ is not an obstruction with respect to

observed rates of amino acid divergence. Similar arguments have been applied to deleterious mutation

load (Galeota-Sprung, Sniegowski, and Ewens 2020).

Although Ewens' argument revolved around lead, which is a difference between fitnesses, his approach

continued the traditional emphasis of evolutionary genetics on variance in fitness, which describes the

mean square of differences (Crow 1958; Ewens 2004; Fisher 1930). Modern traveling wave theory

instead derives the lead directly from s, N, and the beneficial mutation rate U, and obtains the variance in

fitness variance only downstream from that (Desai and Fisher 2007), rather than relying on our ability to

directly measure fitness and its variance as an input to the calculation.

Figure 1. Three different types of arguments have been used to argue for limits to the speed of adaptation. A) The cost of selection is the number of selective deaths that must occur over time to complete a single sweep (each sweep shown as a logistic curve). The cost of selection at one timepoint is the sum of the costs for each current sweep, illustrated as the slopes of the orange lines, each calculated as the subset of deaths $s(1 - p)N$ that contribute to a change in the allele frequency p. B) Load arguments calculate the reduction in mean fitness of a population from what it could be. C) Finite reproductive excess imposes an upper limit on how many selective deaths per generation a population can sustain, which sets an upper bound to how fast substitutions can occur.

Maynard Smith (1976) made a quite different argument against speed limits, claiming that the reason

Haldane's dilemma is not a problem is pervasive synergistic epistasis. Synergistic epistasis increases

differences in fitness above those expected from differences in the numbers of beneficial mutations,

thereby making each selective death more likely to count towards a larger number of sweeps at once. A

persistent source of confusion has been that in his model of truncation selection, Maynard Smith also

made the shift from Haldane's absolute fitness to a more standard population genetic relative fitness, and

hence from lag load to lead. The fact that Haldane's dilemma did not arise in Maynard Smith's model

might therefore be due to reasons put forth by Ewens, rather than due to epistasis.

Although Ewens' lead-based approach negates arguments that convert lag load into a speed limit (i.e.

Haldane's second argument), it doesn't address Haldane's first line of argument: the cost of natural

selection in terms of selective deaths. Confusion between these two disparate lines of argument was

exacerbated by the fact that different papers use the term "substitutional load", which we avoid here, to

mean very different things. 'Substitutional load' has been used to refer to what we here call the lag load

(Kimura and Ohta 1971), the cost of selection (Kimura 1968), the lead (Maynard Smith 1976), the

number of offspring that the most fit genotype must produce (Ewens 2004), the sum of lag load across all

generations involved in a substitution (Kimura 1960; Nei 1971), and even more broadly to refer to

variance rather than load-based arguments when made in the context of similar questions (Ewens 1970).

This confusion in terminology has obscured the consequences of formulating Haldane's dilemma in

different ways.

Nei (1971) and Felsenstein (1971) made a key advance that was perhaps not fully appreciated amidst the

confusion. The relative fitness models which dominate population genetics (e.g. Wright-Fisher and

Moran) implicitly assume inexhaustible reproductive excess (Bertram and Masel 2019). This can be seen

easily in simulations using a rejection sampling method – when fitness is low, an absurd number of

zygotes might be generated and discarded prior to filling the N slots. However, real populations have a

finite reproductive excess, e.g. human females do not easily have more than 20 infants. This constrains

members of the next generation to come from the options contained within that finite set of potential

offspring. This concept has been applied to lethal mutagenesis strategies for anti-viral drugs (Bull,

Sanjuán, and Wilke 2007).

Nei (1971) and Felsenstein (1971) each modelled independently evolving sites in a haploid population

with adult population size N. Each adult has fecundity k, i.e. produces k offspring prior to juvenile deaths

(Figure 1C, first arrow). In their deterministic models, k is exact, but the theory readily generalizes to

interpreting k as an expectation. The raw reproductive excess is thus $(k - 1)N$, with $k > 1$. Some

reproductive excess is lost to non-selective mortality, set not by an extrinsic rate, but derived from

population size regulation after selective mortality (Figure 1C, far right). Extrinsic mortality occurring

prior to selective mortality, and hence at a fixed rate, can be folded into a lower value of k. While the

number of selective deaths available for adaptation is then still denoted $(k - 1)N$, this no longer

represents raw reproductive excess (Figure 1C, second arrow). Haldane's estimate of $k = 1.1$ (resulting in

a maximum of 10% selective deaths), which Nei (1971) and Felsenstein (1971) retain, includes the fact

that extrinsic mortality substantially reduces the fecundity available to be 'used' for selective deaths.

The population then undergoes sweeps, all with the same initial frequency p_0 and selection coefficient s

applying to survival rather than fecundity. Each sweep follows the same trajectory with a mean delay of n

generations between sweeps (Figure 1A). Given independent sites, the cost of selection is summed across

loci at any given point in time (e.g. slopes of orange lines in Figure 1A); Haldane's integral is a method of

calculating the expectation of this sum. Comparing this cost to the reproductive excess of the population

produced the novel result that the minimum spacing n is $-\ln(p_0)/\ln(k)$ (Felsenstein 1971; Nei 1971).

For Haldane's estimates of $p_0 = 10^{-4}$ and $k = 1.1$, this yields $n = 97$ generations between selective

sweeps. This can be compared to Haldane's original spacing of $-\ln(p_0)/\ln\left(\frac{W_{max}}{\overline{W}}\right) = 92$ for a

denominator (somewhat oddly described by Haldane as a selection intensity) $= \ln\left(\frac{W_{max}}{\overline{W}}\right) = 0.1$.

This new limit based on the finite nature of reproductive excess is much slower than the speeds predicted

by lead-based arguments, but is similar in magnitude to Haldane's original result. Importantly, this speed

limit calculation is not subject to the same criticisms as Haldane's original argument. Where Haldane

compared the mean fitness of the population to the mean fitness of a hypothetical population, Nei's

(1971) and Felsenstein's (1971) approach compares the available reproductive excess to the reproductive

excess required to effect changes in allele frequencies. Even if no individual exists who possesses the

beneficial allele at every segregating site, each sweep still requires a certain fraction of deaths to

contribute to its selection. It is the finite nature of reproductive excess that directly produces this limit on

the rate of adaptation.

Felsenstein's (1971) and Nei's (1971) formulations of Haldane's dilemma define the amount of

reproductive excess that is available for selective deaths as $(k-1)N$ after controlling for extrinsic

mortality, where N is the population size prior to the generation of reproductive excess. But the value of k

is an effective value that can conceal much, the "-1" assumes that perfect density regulation demands no

excess individuals above N, and the N refers always to the same adult life history stage. This "effective"

reproductive excess parameter is better conceived of as a measure of the proportion of deaths that are

selective than as a true reproductive excess, and indeed they used Haldane's estimate for 10% deaths

being selective to set $k = 1.1$. Just because a species like *A. thaliana* has high fecundity (high raw

reproductive excess), this tells us nothing about the proportion of deaths that are selective.

Theory

Previous theoretical treatments by Nei (1971) and Felsenstein (1971) assume that all genotypes have the

same fecundity (k), i.e. that there is no selection on fecundity, only on the single life history transition

representing survival. They also assume that extrinsic mortality has a density-dependent component such

that the combination of selective and non-selective mortality is perfectly balanced with fecundity. These

are obviously not realistic assumptions. Next, we extend the theory in a variety of ways.

Selective deaths during generative life history transitions

Previous theoretical treatments by Nei (1971) and Felsenstein (1971) emphasize literal deaths. We can

also describe differences in fecundity as selective 'deaths'. This is because mathematically, foregone

fecundity is equivalent to deaths that take place immediately after fecundity, and can be treated as:

$$\text{Selective 'deaths' during differential fecundity} = N_i(b_{best} - b_i)$$

where N_i is the number of reproductive mature adults, b_i is the fecundity of genotype i, and b_{best} is the

fecundity of the genotype with the highest fecundity in that environment.

Reproductive excess within a fixed life cycle

Next we generalize from just one life history transition experiencing selection, to multiple that occur in a

consistent order. We consider a life history transition j that starts with population size N_j and ends at

population size $N_{j+1} = k_j N_j$. We now define

$$\text{Reproductive excess after transition } j = k_j N_j - N_{min,j+1}$$

where $N_{min,j+1}$ is the minimum population size at the end of transition j that is required in order for the

population to achieve size of N_j at the beginning of transition j in the next life history cycle. Note that

$k_j > 1$ indicates fecundity while $k_j \leq 1$ indicates survival.

To produce selective deaths, $k_{i,j}$ must depend on genotype i. To capture density regulation, $k_{i,j}$ for at

least some values of j must depend on population size N_j. The values $k_{i,j}$ can also be functions of the

genotype frequencies and/or an absolute measure of time. Two life history transitions (survival and

fecundity) is the minimum, but each of these can be broken up into multiple transitions. For example,

survival ($k < 1$) can be broken into components representing survival at different ages, or a selective

component depending only on genotype vs. a density-dependent extrinsic mortality component depending

only on N_j vs. an extrinsic mortality component occurring at a constant rate.

Reproductive excess can be calculated either with respect to the best genotype present (i.e. the one most

likely to become fixed), or with respect to the population mean, by using different values of k_j and

$N_{min,j+1}$. Reproductive excess with respect to the population mean is needed to avoid population decline

in the next generation, while reproductive excess with respect to the best genotype describes the ability to

avoid population decline that would continue even after the best genotype has swept to fixation. The best

choice depends on the particulars of the population in question. For example, studying balancing selection

calls for the population mean, while studying evolutionary rescue calls for the best genotype.

To calculate reproductive excess with respect to the population mean, we solve for $N_{min,j+1}$ in:

$$N_j = N_{min,j+1} \sum_i f_i \prod_{x \neq j} k_{i,x},$$

where f_i is the frequency of genotype i at the beginning of the transition. With respect to the best

genotype, we instead solve for $N_{min,j+1}$ in:

$$N_j = N_{min,j+1} \prod_{x \neq j} k_{best,x}.$$

Nei and Felsenstein as a special case

Nei (1971) and Felsenstein (1971) treated reproductive excess in the conceptually simple case of only two

alternating life history transitions: births and deaths. Deaths included *only* selective deaths, while the

"effective" fecundity transition was non-selective. They handled non-selective deaths by collapsing them

into the value of $k_{fecundity\_effective}$, either before selection on survival (in which case non-selective

deaths reduce $k_{fecundity\_effective}$ in the current generation), or after selection on survival (in such a way

as to exactly balance out any available deaths that were "unused" by selection in the current generation,

by reducing $k_{fecundity\_effective}$ in the subsequent generation). The product $k_{fecundity\_effective} \times$

$k_{selective\_mortality}$ was thus constrained to not exceed 1, via a fudge factor in the former. They solved for

equality to 1 in order to calculate the maximum amount of selective deaths. When this equality is

satisfied, $k_{fecundity\_effective}$ can be interpreted as the product of actual fecundity and non-selective

survival. Fig. 1C interprets this scheme in a temporal manner, proceeding first through non-selective

fecundity to produce raw reproductive excess, then the non-density-dependent component of extrinsic

mortality, then selective deaths, and finally density-dependent extrinsic mortality to cap the population

size at N adults. They score reproductive excess as subject to the first but not the second form of reduction

down to "effective" fecundity, minus the N individuals needed to replace the population.

Reproductive excess beyond a fixed life cycle

Not all organisms proceed through the exact same sequence of life history transitions every time, e.g. with

budding yeast experiencing a variable number of mitoses in between each meiosis, and a variable number

of selfing events between each outcrossing. In this case we cannot take the product of an exact series of

transitions. Instead, we privilege the life history transition that produces the most severe bottleneck,

assuming that the population will spring back to vibrancy after. We define a minimum number of

individuals N_{bot} who need to make it through to the other side of the bottleneck, and define

Reproductive excess at transition $j = k_j N_j - \text{min. needed to ensure } N_{bot} \text{ after bottleneck}$

We now need to take the expectation over all possible series of life history transitions, and solve for

$N_{min,j+1}$ in

379
$$N_{bot} = \text{E}\left(N_{min,j+1} \prod_{x}^{\substack{life\ histo\quad stages \\ between\ j\ and\ bot}} k_{best,x} \right).$$

380   The precise value of N_{bot} will be informed by the ecology of the species in question. It may be small,

381   such as when just a modest number of new hosts, each colonized by just one infectious microorganism, is

382   sufficient to ensure the population's future. The appropriate value of N_{bot} is the smallest population size

383   that reliably escapes extinction.

 Comparison to fitness

Values of k in our framework are equivalent to fitness components, with respect to absolute rather than

relative fitness. Haldane obtained selective deaths from $sN(1-p)$ over a time step of a complete

generation, where s is the selection coefficient with respect to relative fitness. We have shown how

selective deaths can be derived directly from the underlying population dynamic model, without requiring

either generation or relative fitness to be defined first.

Antagonistic pleiotropy is treated quite differently in a selective deaths framework than for fitness

components. Per-generation fitness is the product of fitness components, such that when a genotype that

benefits fitness in one life history transition bears an antagonistically pleiotropic cost at another, the costs

and benefits at least partially cancel out. In contrast, selective deaths accrue across life history transitions

– each selective death absorbs reproductive excess, and there is no reason for them to cancel out.

Similarly, there is no cancelling out across generations, e.g. seasonally fluctuating selection must incur

many selective deaths in order to effect the large allele frequency fluctuations that have been observed

around the long-term mean (Kelly 2022; Machado et al. 2021; Rudman et al. 2022). This high demand for

selective deaths also applies, given life history trade-offs, to unobserved effects that more quickly cancel

out between successive life history transitions.

In the simple case of just births at rate b and deaths at rate d, classic population genetic per-generation

fitness corresponds to the effective reproduction number b/d, while the Malthusian parameter gives an

alternative formulation of fitness as $b - d$. A Malthusian approach is generally preferred when dealing

with complications of age- or stage-structured populations. Our approach extends an effective

reproduction number framing to these more complex scenarios, while avoiding dependence on the

definition of one "generation". While the Malthusian approach is sufficient for many purposes, something

closer to an effective reproduction number approach is required to capture the finite nature of

reproductive excess and the corresponding limits to selective deaths and hence adaptation. We note that

often the question being asked is simply what will invade, in which case either approach can be used

(Lehmann et al. 2016; Metz, Geritz, and Nisbet 1992; Roff 2008).

Comparison to travelling wave models

It is instructive to calculate the required reproductive excess and the proportion of deaths selective in the

asexual relative fitness model treated by Desai and Fisher (2007), whose Eq. 39 solves for the lead qs as a

function of $s, U,$ and N. They define the lead qs in Malthusian fitness terms, but we approximate it here

in terms of per-generation fitness. With an approximately constant population size, the mean genotype has

absolute fitness ~ 1, and the best genotype present (the nose) has absolute fitness $\sim 1 + qs$. The lead qs

can thus be conceived of as the minimum reproductive excess, with respect to the optimal genotype, that

is required in order to avoid limits to adaptation. This aligns closely with the parameter $k - 1$ of Nei

(1971) and Felsenstein (1971). The actual reproductive excess in Desai and Fisher's (2007) model is

infinite, as for all models that assume a constant population size and treat only relative fitness.

We next calculate the proportion of deaths that are selective. The entire reproductive excess of the best

genotype present, $RE \geq 1 + qs$, represents non-selective deaths (or foregone fecundity) among its

offspring. Other genotypes all experience the same rate of non-selective deaths. The per-capita odds that

the next death hits a specific average individual rather than a specific nose individual are $1 : 1/(1 + qs)$.

An average parent therefore expects $RE(1 + qs)$ offspring deaths during the time in which it expects the

RE non-selective offspring deaths that represent a generation, making the proportion of deaths that are

selective equal to $1 - 1/(1 + qs)$.

For a substantial range of parameters, especially with rapid adaptation with large $s, U,$ and $N,$ both the

minimum required reproductive excess, and the proportion of juvenile deaths that are selective, exceed

the previously assumed value of 0.1 (Figure 2). This application to the model of Desai and Fisher (2007)

helps clarify the distinction between these two related properties, which were confounded into a single

value of k by Nei (1971) and Felsenstein (1971). In this particular model, with its explicit adults and

implicit juveniles under selection, they are both functions of the lead qs, but this need not continue to be

so simple when more complex life histories are considered.

Figure 2. As the adaptation rate goes up with increasing U, s, and N, following the multiple mutations

regime of Desai and Fisher (2007), so do the proportion of deaths selective and the minimum

reproductive excess required to sustain that rate of adaptation. Parameter ranges are truncated to avoid the

regime $\frac{s}{U} < 3$, for which the assumptions of the Desai and Fisher's (2007) model break down. The

minimum required reproductive excess qs and the corresponding proportion of selective deaths $1 -$

$1/(1 + qs)$ were calculated by numerically solving Equation 39 for q in Desai and Fisher (2007).

Application of theory to Arabidopsis *data*

The experiment analyzed here has a fixed life cycle of four life history transitions (adults producing

seeds, seeds successfully dispersing to suitable habitat, seeds surviving to be seedlings, seedlings

surviving to be adults). Matching this, we define reproductive excess (RE) four different ways, as

illustrated in Figure 3. Given the presence of many poorly adapted genotypes in the experiment, we

perform each calculation with respect to the best genotype (denoted by the prime symbol $'$), yielding:

448
$$\text{RE(seed survival)} = k'_{seed\_survival} N_{seeds} - \frac{N_{seeds}}{k'_{seedling\_survival} k'_{fecundity} k'_{dispersal}}$$

449
$$\text{RE(seedling survival)} = k'_{seedling\_survival} N_{seedlings} - \frac{N_{seedlings}}{k'_{fecundity} k'_{dispersal} k'_{seed\_survival}}$$

450
$$\text{RE(fecundity)} = k'_{fecundity} N_{adults} - \frac{N_{adults}}{k'_{dispersal} k'_{seed\_survival} k'_{seedling\_survival}}$$

451
$$\text{RE(dispersal)} = k'_{dispersal} N_{seeds\_produced} - \frac{N_{seeds\_produced}}{k'_{seed\_survival} k'_{seedling\_survival} k'_{fecundity}}$$

In the high density environmental conditions, we use a single survival transition to cover both seeds and

seedlings. Under low density conditions, where one seedling was chosen at random from the product of

10 planted seeds, we use $k_{seed\_survival} = 0.1$ for all genotypes. We treat all genotypes as having the same

$k_{fecundity}$ (due to lack of evidence for genetic variation – see Experimental Results below). Our

experiment provides no information about values of $k_{dispersal}$, so we consider a range from 0.01 to 0.1,

equal across genotypes. Because we lack data on the number of seeds after dispersal, we do not calculate

a reproductive excess for this transition. This means that we calculate only two reproductive excesses for

each environmental condition, one for survival and one for fecundity, although the calculations are

different between low- and high-density conditions. These reproductive excesses are given by:

$$\text{RE(fecundity, low density)} = k_{fecundity}N_{adults} - \frac{N_{adults}}{0.1k_{dispersal}k'_{seedling_{survival}}} \quad (1)$$

$$\text{RE(survival, low density)} = k'_{seedling\_survival}N_{seedlings} - \frac{N_{seedlings}}{0.1k_{dispersal}k_{fecundity}} \quad (2)$$

$$\text{RE(fecundity, high density)} = k_{fecundity}N_{adults} - \frac{N_{adults}}{k_{dispersal}k'_{survival}} \quad (3)$$

$$\text{RE(survival, high density)} = k'_{survival}N_{seeds} - \frac{N_{seeds}}{k_{dispersal}k_{fecundity}} \quad (4)$$

Figure 3. Worked example of reproductive excess for the life history transitions of *A. thaliana* in the

experimental setup from Exposito-Alonso *et al* (2019). We found no evidence for between-genotype

differences in fecundity and the experimental setup provides no information about seed dispersal, so we

show no selective deaths during these transitions. Specific values of k are chosen for illustrative purposes.

Experimental

Our empirical findings are restricted to selective deaths relating to juvenile survival. We did not analyze

selective 'deaths' attributable to differences in fecundity, because an ANOVA on fecundity within and

among genotypes showed no statistical support for any difference in mean fecundity among genotypes in

six of the eight experimental conditions. Even in the two conditions with statistical significance, among-

genotypic variance was three times smaller than among-replicate variance. A more sensitive

MCMCglmm model with Poisson errors and controlling for replicate found non-zero heritability in 6/8

environments, but still below 10%, which we consider low enough to neglect.

The proportion of *A. thaliana* deaths that are selective substantially exceeds Haldane's 10% estimate in

six out of eight experimental conditions, and is close to it in the other two (Figure 4 y-axis,

Supplementary Table 2). In Madrid with low water and high density, as many as 95% of deaths are

selective.

A priori, we expect high water and low density to be more benign, which might increase reproductive

excess and/or reduce the proportion of deaths selective. While we cannot compare seed deaths at low

density to seed plus seedling deaths at high density, these predictions are confirmed for high vs. low

water: strongly in the case of selective deaths and excess seeds, and weakly with respect to excess

seedlings (Figure 4, Supplementary Table 2). Estimated reproductive excess is fairly insensitive to the 10-

fold range we consider for the proportion of seeds that successfully disperse to suitable habitats (vertical

line length in Figure 4).

Figure 4. <u>The proportion of deaths that are selective generally exceeds Haldane's 10% estimate (dotted</u>

<u>vertical line), with ample reproductive excess (above zero, shown as dashed horizontal line) especially</u>

<u>under high-water conditions.</u> Selective deaths shown on the x-axis apply either to seeds plus seedlings (A)

or to seedlings alone (B). Reproductive excess in seeds produced vs. seedlings surviving is shown with

open vs. closed symbols, corresponding to values shown on the left and right y-axes, respectively.

Reproductive excess is with respect to what the genotype with the highest survival produced above what

would be required to replace the starting population of a pot. Note that reproductive excess of seedlings

cannot exceed one per pot under low density conditions, and 30 per pot under high density conditions.

Reproductive excess of seeds produced was calculated using equation 1 for low-density conditions and

equation 3 for high-density conditions. Reproductive excess of surviving seedlings was calculated using

equations 2 and 4. All reproductive excesses are shown as a vertical range, with the lower bound

calculated using $k_{dispersal} = 0.01$ and the upper bound calculated using $k_{dispersal} = 0.1$. The

proportions of deaths that are selective are adjusted for extreme value bias as shown in Supplementary

Table 1. Values can be found in Supplementary Table 2.

We expected *a priori* that harsher environmental conditions would have higher extrinsic mortality.

However, this wasn't the case. We can estimate extrinsic mortality as the death rate of the best genotype,

after correcting for extreme value bias (see columns 3 and 4 of Supplementary Table 1). Most

environmental conditions saw a highest-fitness genotype with perfect survival, or close to it. We saw the

most extrinsic mortality in the THP environmental condition, which is not one of the harsher conditions.

Nei (1971) and Felsenstein (1971) implicitly assume high extrinsic mortality via their choice of value for

reproductive excess; we explicitly account for extrinsic mortality during mortality transitions. Extrinsic

mortality might of course be much higher in natural conditions, lowering the proportion of deaths below

the high values observed here.

The artificially high genetic diversity in our experiment might inflate the proportion of deaths that are

selective. If this were the case, then we expect that competition between more similar genotypes should

lead to a smaller estimate for this proportion. We tested this prediction by repeating our analysis on every

pair of genotypes, as though they were the only two genotypes in the experiment, and looking for a

correlation between genetic distance and proportion of deaths selective. Note that we use genetic distance,

much of it presumably neutral, as a proxy for genetic differences related to adaptation. Although some

statistically significant correlations were observed in some environmental conditions, the direction of

correlation was evenly split between negative and positive (as seen in Table 1), and the highest R^2 value

observed in any environmental condition was $0.096^2 = 0.0092$, for seedling deaths in the MLI

environment, which we deem biologically insignificant. This is reassuring with respect to the artificially

high genetic diversity in our experiment.

Life history stage	Seedling deaths		Combined seed and seedling deaths	
Environmental condition	Spearman's rho	p-value	Spearman's rho	p-value
MLI	0.096	2.2e-16		
MHI	0.0004	0.88		
TLI	-0.0024	0.39		
THI	-0.0056	0.042		
MLP			0.055	2.2e-16
MHP			0.011	4.9e-5
TLP			-0.0081	0.0031
THP			-0.0085	0.002

Table 1: <u>Genetic distance between a pair of genotypes does not consistently correlate with proportion of</u>

<u>deaths selective</u>. Visualization of each of the eight relationships is available in Supplementary Figures 4-

5.

DISCUSSION

The prevailing consensus is that Haldane's Dilemma poses no real limitation to the speed of adaptation,

despite persistent confusion as to the reason. Here we began by clarifying the primary issue with load-

based arguments: correcting Haldane's comparison to the best theoretically possible genotype, to instead

compare to the best genotype actually present, enables far more rapid adaptation. However, the historical

consensus that adaptation is not significantly limited largely overlooks a different and crucial type of

limitation pointed out by Nei (1971) and Felsenstein (1971), one that depends not on relative load but on

the finite nature of reproductive excess. If we accepted Haldane's previously unchallenged 10% guess as

an estimate for the reproductive excess that is available for selective deaths to take place among, or as an

estimate of the proportion of deaths that are selective, then the rate of adaptation would be significantly

limited. We clarified and extended theoretical arguments regarding selective deaths drawn from finite

reproductive excess, along with a proof-of-concept application to an experimental dataset. Our extension

applies flexibly to different life histories, including those for which a "generation" is poorly defined, e.g.

colonial organisms.

Our experimental results suggest a possible resolution to Haldane's concern that the rate of adaptation

might be substantively limited. Our illustration of the model of Desai and Fisher (2007) confirms that the

finite nature of reproductive excess would indeed limit the speed of adaptation if only 10% of deaths were

available for selection. But the smallest proportion of selective deaths we observed across 8

environmental conditions was 8.5%, while in the most adverse environmental conditions, 95% of deaths

were selective. Relaxing this auxiliary assumption about a critical parameter value resolves Haldane's

concerns.

The data we use for our proof of concept measured selective deaths under artificial conditions. An

obvious concern with our setup is that with genotypes representing Europe-wide diversity of *A. thaliana*,

exaggerated differences between the best-adapted and worst-adapted genotypes would inflate estimates of

the proportion of deaths that were selective. These concerns are partially mitigated by our unexpected

finding that the genetic distance between genotypes is not consistently related to the proportion of deaths

which would be selective in a competition between genotypes. However, two similar genotypes in our

experiment represent more genetic distance than might be present within a typical natural population, and

even closely related genotypes might differ in important fitness-associated traits. Future work under more

natural conditions (e.g. with higher extrinsic mortality) and in different species (e.g. less fecund) remains

necessary to reach the conclusion that the proportion of deaths that are selective is typically high. Our

framework is flexible enough to be customized for any species, using whichever life history transitions

best describe that species' life history.

Empirical demonstration of the concepts, even with serious caveats about the generalizability of the

empirical example, makes the concepts more concrete. This is especially important because disparate

usage of the term 'substitutional load' in the literature, as well as the variety of underlying lines of

reasoning involved, has made this topic unnecessarily opaque. One aspect of our current work is simply to

clarify the variety of lines of reasoning that produce limits on the rate of adaptation. Our more specific

theoretical and empirical analyses then develop a line of reasoning about reproductive excess and

selective deaths that was not previously resolved. The attention of creation science to this matter (Remine

2005, 2006) highlights the importance of resolving it.

This approach, building on Haldane (1957), Nei (1971), and Felsenstein (1971), is not identical to

standard modes of reasoning in genetics. In particular, quantitative genetics approaches follow Fisher

(1930) to focus on variances — sums of differences squared — while selective deaths and reproductive

excess are, like load, both differences, with no square operation. We hesitate to call this aspect of our

approach 'novel', because it is clearly quite old, but it nevertheless comes across as novel with respect to

aspects of contemporary evolutionary genetics.

Interestingly, the concept of relative load was later reinvented as the "lead", as part of calculations that

derived the actual speed of adaptation v (rather than limits to it) from the beneficial mutation rate U, the

population size N, and the per-mutation selection coefficient s (Desai and Fisher 2007). One reason this

solution was not available to Haldane was that population genetics had not yet begun to treat origination

processes (McCandlish and Stoltzfus 2014). Instead of treating a steady input of beneficial new

mutations, Haldane considered a scenario in which environmental change activates beneficial variants

within standing genetic variation. Indeed, a variant's initial frequency $-\ln(p_0)$ is the primary factor in

determining the maximum speed of adaptation. Some adaptation comes not from activation of standing

genetic variation, but from *de novo* mutations each appearing at initial frequency $1/N$ or $1/2N$. A lead-

based approach was used for the latter to derive the rate of beneficial sweeps in asexuals as $\frac{2s \ln[Ns]}{\ln^2[s/U]}$ for

the simple case of constant s (Desai and Fisher 2007), for parameter ranges in which the previously

derived rate UNs does not apply. Here we relaxed the assumption of infinite reproductive excess made by

this relative fitness model, and calculated the minimal reproductive excess required for the model to hold.

We also reveal the model's implied fraction of deaths (or foregone fecundity) that are selective. Both

quantities are functions of the lead.

Another approach, starting with Kimura (1961), is to use the framework of information theory to set

bounds on the speed of adaptation. Natural selection increases the information stored in genomes (Adami

2012). Kimura calculates the amount of information acquired per sweep in terms of p_0 and then relates

this to the cost of selection using Haldane's equation that $D = -\ln(p_0)$ (Kimura 1961). More recent

approaches treat the bounds placed on the accumulation of information in much more detail, while

treating either the Malthusian parameter (McGee et al. 2022) or classic discrete time relative fitness

(Hledík, Barton, and Tkačik 2022). Both these approaches define an information "cost", but this is not

equal to our cost in terms of selective deaths.

The historical significance of Haldane's arguments about limitations to adaptation is that they were

convincingly used to support neutral theory. This was framed as a dilemma because data on the rates of

amino acid divergence among species seemed to exceed Haldane's speed limit. The development of

neutral theory resolved this apparent dilemma by suggesting that most amino acid substitutions are

neutral and do not count against the speed limit. However, the basis for this historical argument is now on

troubled ground, because recent literature argues that the fraction of substitutions explained by adaptation

can be high (Galtier 2016; Murga-Moreno et al. 2023; Sella et al. 2009; Uricchio, Petrov, and Enard

2019), and that on shorter timescales, as much as 37% of allele frequency change is attributable to

adaptation (Buffalo and Coop 2020). For example, recent experiments have shown rapid, pervasive

seasonal adaptation in *Drosophila* (Bertram 2021; Kelly 2022; Machado et al. 2021). There are other

possible resolutions — e.g. some estimates include substitutions of neutral alleles via hitchhiking.

Nevertheless, it is curious that the empirical collapse of historical arguments for neutral theory has not led

to a re-evaluation of related arguments by Haldane. Here we revise Haldane's arguments for the modern

era, finding that Haldane's revised arguments are compatible with empirical evidence for abundant

adaptation while still posing upper limits that might matter in some contexts.

We note, however, that Haldane (1957) used a one-locus model — linkage disequilibrium will typically

make the conversion of selective deaths to adaptation less efficient than implied by the assumption of

independence used both in his calculations, and in the subsequent calculations of Nei (1971) and

Felsenstein (1971). Here we also considered the other extreme, in the form of the asexual model of Desai

and Fisher (2007). The focus of the current work is to calculate the required reproductive excess and the

proportion of deaths that are selective; more work is required to quantify how this is converted into

adaptation in a broader range of models of linkage disequilibrium and epistasis, as well as life histories

and antagonistic pleiotropy.

Excitingly, unlike most approaches in evolutionary genetics, the approach we describe does not require

the quantity 'fitness', which is deceptively difficult to define in a manner that can be generalized to all

circumstances (Ariew and Lewontin 2004; Bertram and Masel 2019; Doebeli, Ispolatov, and Simon 2017;

Van Valen 1989). Standard quantitative definitions of either relative or absolute 'fitness' require a clear

definition of a 'generation' over which change is assessed, which in turn requires a clear definition of an

'individual' whose life cycle a generation captures (Wilson and Barker 2021). While "lineage fitness"

solves a number of problems (Akçay and Van Cleve 2016; Graves and Weinreich 2017; King and Masel

2007), it does so at the cost of defining the fittest genotype to be that which tends to eventually prevail,

sacrificing much of the quantitative benefit of 'fitness'. Our generalized selective deaths approach is

derived from selection, but does not require 'fitness' to be defined. Rather, we measure selective deaths

from pairwise differences in fecundity and survival between each genotype and the best genotype present,

during each life history transition. Reproductive excess corresponding to that life history transition

indicates how stringent selection can be, without triggering a decline in population size. This approach

applies at each life history transition and can therefore be generalized to species with complex life

histories, where it becomes difficult to define a 'generation' and therefore fitness.

GLOSSARY

Absolute fitness of an adult: expected number of individual offspring surviving to adulthood (defined

here for hermaphrodite species — half this value, if reproducing sexually with males).

Absolute fitness of a juvenile: expected number of individual offspring (or half this value, if reproducing

sexually with males). Failure to survive to adulthood (reproductive maturity) implies zero offspring.

Adult: Reproductively mature individual. More than one adult life history stage may be defined.

Cost of selection: The number of selective deaths that must occur over time to accomplish defined

evolutionary change, e.g. to complete a single selective sweep.

Generation: A set of life history transitions that ends the first time it returns, with new individuals, to the

same life history stage (e.g. adult or juvenile) where it began.

Individual: An organism that meets a loosely-defined set of criteria (Wilson and Barker 2021), including

a shared genome, and the degree of integration of parts. Whether e.g. a group of microbes is a closely

connected ecological community vs. an individual organism may be a matter of biological judgment.

Juvenile: Individuals that are not yet reproductively mature. More than one juvenile life history stage

may be defined, e.g. before vs. after dispersal.

Lag load: The difference in fitness between a theoretical best genotype that might not be present in the

population and the average genotype present.

Lead: The difference in fitness between the best genotype present in the population and the average

genotype present.

Life history transition: Survival (i.e. persistence of an individual), reproduction (i.e. generation of new

individuals) and/or organismal growth from one life history stage to the next

Load: A difference in fitness between an actual genotype or population and a reference. See "lag load"

and "lead" as concrete examples.

Relative fitness: expected relative genetic contribution to the next generation

Reproductive excess: The degree to which a hypothetical population concludes a life history transition

with a larger population than the minimum required to complete a life history cycle without the

population shrinking in size.

Selective deaths: The subset of deaths (or foregone fertility) that contributes to selective changes in allele

frequency. This can be quantified as how many deaths each genotype experiences that would not have

been experienced if that genotype were replaced by the best genotype.

DATA AVAILABILITY

All analysis was performed in R, and our code is available on GitHub:

www.github.com/josephmatheson/selective_deaths.

ACKNOWLEDGEMENTS

This work was supported by the John Templeton Foundation [62028] and [62220]. We thank Taylor

Kessinger for sparking the last author's original interest in this topic many years ago by presenting

Haldane's 1957 paper in journal club, and Jason Bertram and Ryan Gutenkunst for helpful discussions.

AUTHOR CONTRIBUTIONS

Conception of the paper by Joanna Masel and Moises Exposito-Alonso. Design and interpretation by

Joseph Matheson and Joanna Masel. Data and details of experimental methodology provided by Moises

Exposito-Alonso. Data analysis mostly by Joseph Matheson, with the MCMCglmm model analyzed by

Moises Exposito-Alonso. Manuscript drafted by Joseph Matheson and substantively revised by all

authors.

CONFLICT OF INTEREST

The authors declare no conflicting interests.

BIBLIOGRAPHY

Adami, Christoph. 2012. "The Use of Information Theory in Evolutionary Biology." *Annals of the New*
York Academy of Sciences 1256(1):49–65. doi: 10.1111/j.1749-6632.2011.06422.x.

Akçay, Erol, and Jeremy Van Cleve. 2016. "There Is No Fitness but Fitness, and the Lineage Is Its
Bearer." *Philosophical Transactions of the Royal Society B: Biological Sciences* 371(1687). doi:
10.1098/rstb.2015.0085.

Ariew, André, and R. C. Lewontin. 2004. "The Confusions of Fitness." *British Journal for the Philosophy*
of Science 55(2):347–63. doi: 10.1093/bjps/55.2.347.

Bell, Graham. 2017. "Evolutionary Rescue." *Annual Review of Ecology, Evolution, and Systematics*
48:605–27. doi: 10.1146/annurev-ecolsys-110316-023011.

Bertram, J., K. Gomez, and J. Masel. 2017. "Predicting Patterns of Long-Term Adaptation and Extinction
with Population Genetics." *Evolution*. doi: 10.1111/evo.13116.

Bertram, Jason. 2021. "Allele Frequency Divergence Reveals Ubiquitous Influence of Positive Selection
in Drosophila." *PLOS Genetics* 17(9):e1009833. doi: 10.1371/journal.pgen.1009833.

Bertram, Jason, and Joanna Masel. 2019. "Density-Dependent Selection and the Limits of Relative
Fitness." *Theoretical Population Biology* 129:81–92. doi: 10.1016/j.tpb.2018.11.006.

Buffalo, Vince, and Graham Coop. 2020. "Estimating the Genome-Wide Contribution of Selection to
Temporal Allele Frequency Change." *Proceedings of the National Academy of Sciences of the*
United States of America 117(34):20672–80. doi: 10.1073/pnas.1919039117.

Bull, JJ, R. Sanjuán, and CO Wilke. 2007. "Theory of Lethal Mutagenesis for Viruses." *Journal of*
Virology 81(6):2930–39. doi: 10.1128/jvi.01624-06.

Crow, James F. 1958. "Some Possibilities for Measuring Selection Intensities in Man." *Human Biology*
30(1):1–13.

Desai, Michael M., and Daniel S. Fisher. 2007. "Beneficial Mutation-Selection Balance and the Effect of
Linkage on Positive Selection." *Genetics* 176(3):1759–98. doi: 10.1534/genetics.106.067678.

Doebeli, Michael, Yaroslav Ispolatov, and Burt Simon. 2017. "Towards a Mechanistic Foundation of
Evolutionary Theory." *ELife* 6:1–17. doi: 10.7554/elife.23804.

Ewens, Warren. 1970. "Remarks on the Substitutional Load." *Theoretical Population Biology* 139:129–
39.

Ewens, Warren. 2004. *Mathematical Population Genetics*. 2nd ed. New York: Springer US.

Exposito-alonso, Moises, Oliver Bossdorf, Rasmus Nielsen, and Detlef Weigel. 2019. "Natural Selection
on the Arabidopsis Thaliana Genome in Present and Future Climates." *Nature* 573(7772):126–29.
doi: 10.1038/s41586-019-1520-9.

Felsenstein, Joseph. 1971. "On the Biological Significance of the Cost of Gene Substitution." *American
Naturalist* 105(941):1–11.

Fisher, Ronald. 1930. *The Genetical Theory of Natural Selection*. Oxford: Clarendon Press.

Galeota-Sprung, Benjamin, Paul Sniegowski, and Warren Ewens. 2020. "Mutational Load and the
Functional Fraction of the Human Genome." *Genome Biology and Evolution* 12(4):273–81. doi:
10.1093/gbe/evaa040.

Galtier, Nicolas. 2016. "Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis." *PLoS Genetics* 12(1):1–23. doi: 10.1371/journal.pgen.1005774.

Graves, Christopher J., and Daniel M. Weinreich. 2017. "Variability in Fitness Effects Can Preclude
Selection of the Fittest." *Annual Review of Ecology, Evolution, and Systematics* 48:399–417. doi:
https://doi.org/10.1146/annurev-ecolsys-110316-022722.

Haldane, JBS. 1957. "The Cost of Natural Selection." *Genetics* 55(3):511–24.

Hledík, Michal, Nick Barton, and Gašper Tkačik. 2022. "Accumulation and Maintenance of Information
in Evolution." *Proceedings of the National Academy of Sciences of the United States of America*
119(36):1–10. doi: 10.1073/pnas.2123152119.

Kelly, John. 2022. "The Genomic Scale of Fluctuating Selection in a Natural Plant Population." *Evolution
Letters* 506–21. doi: 10.1002/evl3.308.

Kern, Andrew D., and Matthew W. Hahn. 2018. "The Neutral Theory in Light of Natural Selection."
Molecular Biology and Evolution 35(6):1366–71. doi: 10.1093/molbev/msy092.

Kimura, Motoo. 1960. "Optimum Mutation Rate and Degree of Dominance as Determined by the
Principle of Minimum Genetic Load." *Genetics* 57(1):21–34.

Kimura, Motoo. 1961. "Natural Selection as the Process of Accumulating Genetic Information in
Adaptive Evolution." *Genetical Research* 2(1):127–40. doi: 10.1017/S0016672300000616.

Kimura, Motoo. 1968. "Evolutionary Rate at the Molecular Level." *Nature* 217:624–26.

Kimura, Motoo. 1969. "The Number of Heterozygous Nucleotide Sites Maintained in a Finite Population
Due to Steady Flux of Mutations." *Genetics* 61(4):893–903.

Kimura, Motoo, and James F. Crow. 1969. "Natural Selection and Gene Substitution." *Genetical
Research* 13(2):127–41. doi: 10.1017/S0016672300002846.

Kimura, Motoo, and Tomoko Ohta. 1971. "On the Rate of Molecular Evolution." *Journal of Molecular
Evolution* 1(1):1–17. doi: 10.1007/BF01659390.

King, Oliver D., and Joanna Masel. 2007. "The Evolution of Bet-Hedging Adaptations to Rare
Scenarios." *Theoretical Population Biology* 72(4):560–75. doi: 10.1016/j.tpb.2007.08.006.

Lehmann, Laurent, Charles Mullon, Erol Akçay, and Jeremy Van Cleve. 2016. "Invasion Fitness,
Inclusive Fitness, and Reproductive Numbers in Heterogeneous Populations." *Evolution;*
International Journal of Organic Evolution 70(8):1689–1702. doi: 10.1111/evo.12980.

Machado, Heather E., Alan O. Bergland, Ryan Taylor, Susanne Tilk, Emily Behrman, Kelly Dyer, Daniel
K. Fabian, Thomas Flatt, Josefa González, Talia L. Karasov, Bernard Kim, Iryna Kozeretska, Brian
P. Lazzaro, Thomas J. S. Merritt, John E. Pool, Katherine O'brien, Subhash Rajpurohit, Paula R.
Roy, Stephen W. Schaeffer, Svitlana Serga, Paul Schmidt, and Dmitri A. Petrov. 2021. "Broad
Geographic Sampling Reveals the Shared Basis and Environmental Correlates of Seasonal
Adaptation in Drosophila." *ELife* 10. doi: 10.7554/eLife.67577.

Maynard Smith, J. 1968. "'Haldane's Dilemma' and the Rate of Evolution." *Nature* 219(5159):1114–16.
doi: 10.1038/2191114a0.

Maynard Smith, J. 1976. "What Determines the Rate of Evolution?" *American Naturalist* 110(973):331–
38.

McCandlish, David M., and Arlin Stoltzfus. 2014. "Modeling Evolution Using the Probability of
Fixation: History and Implications." *Quarterly Review of Biology* 89(3):225–52. doi:
10.1086/677571.

McGee, Ryan Seamus, Olivia Kosterlitz, Artem Kaznatcheev, Benjamin Kerr, and Carl T. Bergstrom.
2022. "The Cost of Information Acquisition by Natural Selection." *Biorxiv* 2022.07.02. doi:
10.1101/2022.07.02.498577.

Metz, J. A. J., S. A. H. Geritz, and R. M. Nisbet. 1992. "How Should We Define 'Fitness' for General
Ecological Scenarios? The Simplest Textbook Model for Selection in Large Populations Relies N(t
Fl) = B(E(t)) N(t) (1) with B(E(t)) a Matrix Whose El-Ements Depend on E(t), Implying N(t) =
B(E(I-I)) B(E(t-2)). .." *Trends in Ecology & Evolution* 7(6):198–202.

Murga-Moreno, Jesús, Sònia Casillas, Antonio Barbadilla, Lawrence Uricchio, and David Enard. 2023.
"An Efficient and Robust ABC Approach to Infer the Rate and 1 Strength of Adaptation." *Biorxiv*.
doi: 10.1101/2023.08.29.555322.

Nei, Masatoshi. 1971. "Fertility Excess Necessary for Gene Substitution in Regulated Populations."
Genetics 68:169–84.

Remine, Walter. 2006. "More Precise Calculations of the Cost of Substitution." *Creation Research*
Science Quarterly 43(September).

Remine, Walter J. 2005. "Cost Theory and the Cost of Substitution — a Clarification." *Journal of*
Creation 19(1):113–25.

Roff, Derek A. 2008. "Defining Fitness in Evolutionary Models." *Journal of Genetics* 87(4):339–48. doi:
10.1007/s12041-008-0056-9.

Rudman, Seth M., Sharon I. Greenblum, Subhash Rajpurohit, Nicolas J. Betancourt, Jinjoo Hanna,
Susanne Tilk, Tuya Yokoyama, Dmitri A. Petrov, and Paul Schmidt. 2022. "Direct Observation of
Adapting Tracking on Ecological Time Scales in Drosophila." *Science* 375(6586). doi:
10.1126/science.abj7484.

Sella, Guy, Dmitri A. Petrov, Molly Przeworski, and Peter Andolfatto. 2009. "Pervasive Natural
Selection in the Drosophila Genome?" *PLoS Genetics* 5(6):e1000495. doi:
10.1371/journal.pgen.1000495.

Uricchio, Lawrence H., Dmitri A. Petrov, and David Enard. 2019. "Exploiting Selection at Linked Sites
to Infer the Rate and Strength of Adaptation." *Nature Ecology & Evolution* 3(6):977–84. doi:
10.1038/s41559-019-0890-6.

Van Valen, Leigh. 1963. "Haldane's Dilemma, Evolutionary Rates, and Heterosis." *American Naturalist*
97(894):185–90.

Van Valen, Leigh. 1989. "Three Paradigms of Evolution." *Evolutionary Theory* 9(2):1–17.

Wilson, Robert A., and Matthew J. Barker. 2021. "Biological Individuals" edited by E. N. Zalta. *Stanford
Encyclopedia of Philosophy* 2021.

SUPPLEMENTARY INFORMATION

Contents

Supplementary Methods

Correction for extreme value bias

The estimated best genotype is subject to extreme value bias, leading to overestimation of the number of selective deaths. I.e., the best genotype observed is likely not only to be a superior genotype, but also to have outperformed its own expected death rate by chance. The more uncertainty in estimated genotypic survival rates, relative to true genetic variance, the worse the extreme value bias problem. Here we attempt a rough estimate of the magnitude of extreme value bias using the observed noise among replicates and among genotypes. We then subtract a conservative estimate of bias from our best observed genotype, and redo our calculations of selective deaths.

We perform 10,000 simulations per environmental treatment. In each simulation, we assume that the observed 517 genotypic death rates are the 'true' values for each genotype (thus slightly overestimating genetic variance within the population). We then resample the number of surviving seeds 'observed' for each replicate of that genotype using a binomial distribution and calculate the resulting observed death rate.

In each simulated dataset, we took the best observed genotype and recorded the difference between its observed value and its actual genotypic value, then averaged these differences across the 10,000 simulations to obtain estimated bias. We then adjusted our estimate of the number of selective deaths to incorporate this estimated bias:

$$\text{Total adjusted selective deaths} = \sum_{i=1}^{517} n_i(d_i - (d_{best} + bias)).$$

Adjustments to selective deaths are shown in Supplementary Table 1. Across all environmental treatments, adjusting for extreme value bias leads to negligible change in the estimate of the proportion of deaths which are selective.

Supplementary Tables

Environmental condition	Among-genotype variance	Observed maximum survival rate	Estimated bias	Unadjusted proportion of deaths selective	Adjusted proportion of deaths selective
MLI	0.0618	1	0	0.664	0.664
MHI	0.0144	1	0	0.0847	0.0847
TLI	0.0337	1	0	0.564	0.564
THI	0.0139	1	0	0.109	0.109
MLP	0.0113	1	0	0.96	0.96
MHP	0.0433	1	0	0.564	0.564
TLP	0.023	0.947	0.00418	0.633	0.629
THP	0.0195	0.871	0.0195	0.391	0.372

Supplementary Table 1: Adjusting for extreme value bias has little effect on the estimated proportion of selective deaths.

In six out of eight environmental conditions, at least one genotype had all plants survive until the end of the experiment. This genotype then has an observed death rate of zero and an observed between-replicate variance of zero, and thus our approach will not find any bias. Determining bias in this case would require attempting to fit and then resample from a true distribution of genotypic values that includes death rates near but not equal to zero. We did not pursue this avenue, because resampling genotypes would add a new source of variance, and we need only correct for the extreme value bias pertaining to the genotypes actually studied. Given the small degree of bias observed in the two environmental conditions in which no genotype experienced perfect survival, we consider our approach sufficient to conclude that extreme value bias has little quantitative effect on our results.

	Proportion of seedling deaths selective	**Proportion of combined seed and seedling deaths selective**	**Excess seedlings per pot after seedling survival transition**	**Excess seedlings per pot after seed and seedling survival transition**	**Excess seeds produced per pot after fecundity transition**
MLI	0.664		0.742-0.974		967-1,269
MHI	0.0847		0.945-0.995		15,856-16,680
TLI	0.564		0.794-0.979		1,683-2,075
THI	0.109		0.908-0.991		8,848-9,651
MLP		0.960		20.51-29.05	257-364
MHP		0.564		28.16-29.82	20,029-21,207
TLP		0.633		13.71-22.97	833-1,396
THP		0.391		21.52-24.47	10,374-11,796

Supplementary Table 2. Proportion of deaths that are selective are above Haldane's 10% estimate for most environmental conditions. Reproductive excess after the survival and fecundity life history transitions are high. Note that reproductive excess per pot for low-density conditions cannot exceed one, and excess per pot for high-density conditions cannot exceed 30. For reproductive excess, lower bounds are calculated using $k_{dispersal} = 0.01$ and higher bounds are calculated using $k_{dispersal} = 0.1$. Reproductive excess values for fecundity are rounded to the nearest integer; approximate methods were used to estimate seed number.

Supplementary Figures

Supplementary Figure 1. Histograms of genotype mean seed production for every genotype with at least one surviving adult in each of the eight environmental conditions.

Supplementary Figure 2. Histograms of genotype mean seed survival for every genotype in each of the eight environmental conditions.

Supplementary Figure 3. Histograms of Box-Cox transformed fecundity values for every pot with surviving adults in each of the eight environmental conditions. Lambda values for the Box-Cox transformations are (A) -0.02, (B) 0.46, (C) 0.18, (D) -0.06, (E) 0.63, (F) 0.75, (G) -0.18, (H) -0.06

Supplementary Figure 4. Hamming distance does not substantially predict the proportion of deaths selective during the life history stage of combined seed and seedling survival at low density.

Supplementary Figure 5. Hamming distance does not substantially predict the proportion of deaths selective during the life history stage of seedling survival at high density.