The mitochondrial gene CMPK2 functions as a rheostat for macrophage homeostasis in

inflammation

Prabhakar Arumugam[1,2], Meghna Chauhan[1,2], Thejaswitha Rajeev[1], Rahul Chakraborty[1,2],

Deepthi Shankaran[1,2], Sivaprakash Ramalingam[1,2], Sheetal Gandotra[1,2], Vivek Rao[1,2]*

1-CSIR- Institute of genomics and Integrative Biology, Mathura Road, New Delhi-110025

India.

2- Academy of Scientific and Innovative Research (AcSIR), India

*Corresponding author: Tel: +91 11 29879229, E-mail: vivek.rao@igib.res.in

Abstract

In addition to their role in cellular energy production, mitochondria are increasingly recognized as

regulators of the innate immune response of phagocytes. Here, we demonstrate that altering

expression levels of the mitochondria associated enzyme, cytidine monophosphate kinase 2

(CMPK2), disrupts mitochondrial physiology and significantly de-regulates the resting immune

homeostasis of macrophages. Both CMPK2 silenced as well as constitutively over expressing

macrophage lines portray mitochondrial stress with marked depolarization of their membrane

potential, enhanced ROS and disturbed architecture culminating in the enhanced expression of

the pro-inflammatory genes- IL1β, TNFα and IL8. Interestingly, the long-term modulation of

CMPK2 expression resulted in an increased glycolytic flux of macrophages akin to the altered

physiological state of activated M1 macrophages. While infection induced inflammation for

restricting pathogens is regulated, our observation of a total dysregulation of basal inflammation

by bidirectional alteration of CMPK2 expression, only highlights a critical role of this gene in

mitochondria mediated control of inflammation.

Keywords: CMPK2, M1 macrophage, Immuno-metabolism, Mitochondria, Infection

Introduction

Intracellular microbial infections compel host macrophages to activate selective cellular pathways that are targeted towards the elimination of the pathogen. In response to bacterial infections like Mtb, in addition to the expression of pro-inflammatory response mediators that facilitate the activation of bactericidal pathways (M1 polarized), recent evidence has demonstrated the robust activation of type I IFN response in macrophages(Galvan-Pena and O'Neill, 2014; Kumar et al., 2019; Manzanillo et al., 2012; Patrick and Watson, 2021). While the importance of the individual genes of this response cascade (interferon stimulated genes- ISG) in restricting viral nucleic acids is well recognized, its contribution in controlling bacterial infections is still unclear (Alphonse et al., 2021; Boxx and Cheng, 2016; Snell et al., 2017).

One of the early and highly expressed ISG in macrophages following infection is the mitochondria associated kinase CMPK2. CMPK2 is actively induced in response to several viral infections like HEV(Zhang et al., 2014), HIV(El-Diwany et al., 2018), Dengue virus (DENV)(Lai et al., 2021b), Nipah virus(Hassan et al., 2019), SVCV(Liu et al., 2019) and in response to LPS and poly IC(Kim et al., 2021) in macrophages. A recent study has demonstrated the involvement of the fish homolog-3nCmpk2 in antibacterial response by controlling bacterial colonization and protecting the intestinal barrier (Feng et al., 2021). In addition to its role in protective responses, CMPK2, has recently been implicated in activating host cell inflammasomes (Zhong et al., 2018) by virtue of its role in mitochondrial DNA synthesis thereby contributing to liver tissue injury and fostering premature atherosclerosis in SLE(Lai et al., 2021a; Luo et al., 2021).

This inherent duality of the inflammatory response: 1) leading to infection control (host beneficial) and 2) extensive tissue damage on uncontrolled activation (detrimental) signifies the need to effectively regulate the onset and end of the inflammation. Several mechanisms have been characterized for the control of inflammation by host phagocytes (Cohen and Mosser, 2013; Lawrence and Natoli, 2011; Sugimoto et al., 2016; Yan and Horng, 2020). Studies in the last decade have implicated a determinant connection between mitochondria and the inflammation

balance of immune cells (Missiroli et al., 2020; Tiku et al., 2020; Wang et al., 2021). The enormous

metabolic flexibility offered by mitochondria resident processes forms the basis of cellular

plasticity facilitating fine-tuned responses to physiological conditions (Labbe et al., 2014; Picard

and McEwen, 2018). The importance of mitochondrial health in terms of physiology, network

architecture in immune cell function and response to infection is well recognized (Kikuchi et al.,

1988; Rambold and Pearce, 2018; Weinberg et al., 2015). Activated M1 polarized macrophages

demonstrate an important shift in the metabolic state relying completely on glycolysis for

increased and faster production of energy(Blagih and Jones, 2012; Kikuchi *et al.*, 1988;

Newsholme et al., 1986; Rodriguez-Prados et al., 2010; Viola et al., 2019). This transition is

associated with a distinctive change in mitochondrial structure and a diminished reliance on

mitochondrial oxidative phosphorylation and increased production of mitochondrial ROS(Kelly

and O'Neill, 2015; Labadarios et al., 1987; Mills et al., 2016; Van den Bossche et al., 2016).

In line with our previous study on the robust induction of type I IFN response in Mtb infected

macrophages (Shankaran et al., 2019), we show, that CMPK2, an ISG, is highly induced

immediately following infection and is rapidly regulated to baseline levels in human macrophages.

Altered expression of this gene is associated with a marked change in mitochondrial physiology

indicating acute organellar stress; resting macrophages with constitutive silencing or over-

expression of CMPK2 harbor significantly depolarized mitochondria with an atypical organellar

network and heightened elaboration of mitochondrial ROS. This is reflected in the heightened

levels of basal expression of inflammatory genes and a significant shift in macrophage

metabolism to reliance on glycolysis resembling the M1 macrophages in these cells. Our study

thus provides evidence for the importance of this gene in controlling inflammation in macrophages

and the importance of regulated CMPK2 expression in protective responses, thereby highlighting

an important link between mitochondrial metabolism and innate response kinetics of

macrophages.

Results

Macrophages respond to infection/ TLR4 stimulation by inducing CMPK2

Given the early induction of type I IFN response in Mtb infections (Boxx and Cheng, 2016; Kovarik et al., 2016; McNab et al., 2015), we sought to investigate the importance of this pathway in the response dynamics of macrophages. The recent developments on the importance of mitochondrial physiology in host cell infection responses directed our attention to the mitochondria associated ISG-*CMPK2* that was highly induced following Mtb infection of macrophages. Infected THP1 macrophages showed a steady increase in *CMPK2* expression from 15-fold by 6h and attaining peak levels of ~140 fold by 12h and thereafter maintained even at 24h of infection to greater than 100- fold (Fig. 1A). In line with previous reports (Zhong *et al.*, 2018), we also observed nearly 20- fold increase in *CMPK2* expression in response to LPS stimulation while the other TLR ligands failed to alter this gene (Fig. 1B). This was also evident as a steady increase in CMPK2 protein levels in THP1 macrophages even at 48h of LPS stimulation (Fig. 1C). The complete loss of LPS induced expression in the presence of a TLR4 signaling inhibitor-CLI095, confirmed the dependence of *CMPK2* expression on TLR4 signaling in macrophages (Fig. 1D). Further, macrophages responded to *Salmonella enterica subsp. enterica* serovar Typhimurium (STM) infection, by early induction of *CMPK2* expression with 15-16 -fold increase by 3h wherein escalated further to ~30 fold by 6h of infection (Fig. 1E) and this response was completely abolished with the addition of CLI095 confirming the TLR4 engagement dependent expression of this gene in macrophages (Fig. 1F). A decrease in expression by more than 50% was observed in the case of macrophages infected with Mtb and treated with CLI095 again supporting the importance of TLR4 signaling in *CMPK2* expression (Fig. S1).

Fig. 1

Fig. 1: CMPK2 expression is induced following TLR4 stimulation of infection.

[A-F] Analysis of CMPK2 expression in THP1 macrophages following: infection with Mtb at a MOI of 5 [A], stimulation with various TLR ligands- c: Untreated cells, L: LPS (10 ng/ml), P: Pam3CSK4 (20 ng/ml), Z: Zymosan (10 μg/ml), I: Poly IC (2 μg/ml) [B], LPS stimulation by immunoblotting with CMPK2-specific antibody [C], LPS stimulation with and without TLR4 inhibitor CLI095 [D], infection with STM [E], infection with STM in the presence of CLI095 [F]. Expression of *CMPK2* was quantitated by qPCR. Values are normalized with *GAPDH* and mean fold change compared to control cells $\pm$ SEM from N=2/ 3 independent experiments.

Modulation of CMPK2 activity affects the inflammation status of macrophages

To investigate the molecular function of CMPK2 in the macrophage response, we used *CMPK2* specific siRNAs (a-c) to specifically silence this gene in THP1 cells (KD) and achieved 60-75% decrease in gene expression in comparison to non-targeting siRNA (NT) (Fig. S2A). Only siRNA-c showed a significant decrease in protein levels in excess of 40% of the native CMPK2 expression and was used for further analysis (Fig. 2A). Given the critical role of proinflammatory cytokines in macrophage response to infection, we compared the levels of TNFα and IL1β in the NT and CMPK2 knockdown (KD) macrophages in response to Mtb infection. Mtb infection

induced *TNFα* and *IL1β* expression by 200-500- fold in KD macrophages while these levels were

10-20- fold lower in the case of NT macrophages (Fig. 2B). Interestingly, expression levels of

these genes were markedly elevated in the uninfected KD macrophages. This pattern was also

reflected by the enhanced expression levels of other immune effectors like *IL8*, *IP10*, *VEGF* and

decrease in *IL10* in naïve KD macrophages strongly alluding to an enhanced pro- inflammatory

status of these cells even in the resting state (Fig. 2C). This profile of increased *TNFα* and *IL1β*

gene expression was also observed in KD cells following LPS stimulation (Fig. S2B). This

enhanced expression also correlated well with the increase in secreted TNFα and IL1β in the

culture supernatant of LPS treated KD cells (Fig. 2D). Again, correlating with gene expression,

naïve KD cells also secreted increased amounts of both TNFα and IL1β in the supernatants as

compared to the non- detectable levels of the cytokine in naïve NT macrophages.

Given the inflammatory phenotype observed with decreased CMPK2 expression in THP1

macrophages, we attempted to reverse this by complementing a functional copy of the gene in

the silenced cells. We expressed the complete CMPK2 as a mCherry fusion protein in pCDNA3.1

and analyzed the basal expression levels of *IL1β* and *TNFα*. As expected, *CMPK2* expression

was ~30 folds higher than in the empty vector control macrophages (Fig. 2E). Contrary to our

expectations, this enhanced expression of the *CMPK2* could not reverse the alteration in basal

inflammation of these cells: nearly similar levels of *IL1β* and *TNFα* were observed in the

complemented cells as in the silenced cells (Fig. 2F). We hypothesized that similar to the situation

with CMPK2 silenced cells, an increase in expression of CMPK2 was also detrimental to the

inflammation homeostasis of macrophages. To test this, we analyzed the pro-inflammatory status

of wild type (Wt) THP1 cells stably expressing CMPK2-mCherry. Enhanced expression of *CMPK2*

transcript in excess of 100- fold basal levels was observed in the over expressing (OE) cells

compared to the vector control (VC) cells (Fig. S2C). The fusion protein of CMPK2 and mCherry

was also clearly detected in protein extracts of the OE strains by immunoblotting (Fig. S2D).

Again, this enhanced expression of CMPK2 resulted in heightened levels of *TNFα* (12-15- fold)

and *IL1β* (60-80- fold) in OE cells (Fig. 2G).

In an effort to understand the basis for the hyper-inflammatory state of KD cells, we compared

the immune signaling cascades of these cells with the NT cells in the basal state. Consistently,

we observed enhanced activation of ERK and NFκB with 3-fold higher levels of phospho-p42/44

(ERK) and p65 NFκB (Fig. 2H, S2E) without any change in phospho- JNK/SAPK and p38 MAP

kinase between the two cells (Fig. S2F). This pattern of enhanced ERK phosphorylation was also

observed in the case of OE cells in comparison to VC cells. The use of an ERK signaling specific

inhibitor U0126 not only reduced the levels of phospho-p42/44 in both the KD and OE cells

significantly (Fig. 2I), but also significantly reduced the elevated levels of both *IL1β* and *TNFα* in

both these cell types, further confirming the importance of ERK signaling in mediating the hyper

inflammatory phenotype of these macrophages (Fig. 2J, 2K).

Human mitochondrial CMPK2 has been shown to be a nucleotide kinase involved in mitochondrial

DNA synthesis (Fig. 2L). In an effort to probe the importance of the kinase domain, we substituted

the active site aspartate to alanine (D330A) as a kinase dead variant of CMPK2 (D330A) and

analyzed the functional consequence of this expression on cytokine gene expression. Despite the

strong and stable expression of D330A, comparable to the Wt, CMPK2 (OE), both at the transcript

(Fig. S2G) and protein (Fig. 2L) level, there was no evidence of increased basal inflammation.

Contrasting with the significantly elevated levels of *TNFα* and *IL1β* in the OE cells (~10 and ~80

fold, respectively over basal levels), THP1 cells transfected with D330A, harbored normal levels

of gene expression (Fig. 2M) emphasizing the importance of the kinase activity of CMPK2 in

regulating basal inflammation in macrophages.

Fig. 2

Fig. 2: CMPK2 regulates the basal inflammation of macrophages.

[A] Expression of CMPK2 in CMPK2 silenced macrophages by immunoblotting with specific antibodies. For immunoblotting, the expression of α-TUBULIN was used as control. One representative blot of 3 independent experiments is shown. The levels of CMPK2 expression are depicted as mean fold change $\pm$ SEM of N=3 experiments by densitometry. [B] Expression of the *IL1β* and *TNFα* transcripts following infection with Mtb for 6h was evaluated in NT or KD macrophages by qPCR. [C] Basal level expression of transcripts *CMPK2* and cytokine genes in KD macrophages was estimated by qPCR with specific primers. Values are normalized with *GAPDH* and mean fold change compared to control cells $\pm$ SEM from N=3 independent experiments. [D] Levels of TNFα and IL1β in the culture supernatants of THP1 stably

expressing non-targeting (NT) or siRNA specific to CMPK2 (KD) estimated by ELISA. Values are

represented as mean pg/ml of the cytokine in triplicate assays from N=3 experiments. [E] Expression of

CMPK2 in the CMPK2 silenced cells (KD) with stable expression of CMPK2 (KD-OE) or empty vector (KD-

VC). The levels in the corresponding NT and KD cells are also shown. Values are represented as mean $\pm$

SEM of the triplicate assays from N=3 experiments. [F] Expression of *TNFα* and *IL1β* in the CMPK2 silenced

cells with stable expression of CMPK2 (KD-OE) or empty vector (KD-VC). The levels in the corresponding

NT and KD cells are also shown. Values are represented as mean $\pm$ SEM of the triplicate assays from N=3

experiments. [G] Expression of *TNFα* and *IL1β* in THP1 cells after stable expression of CMPK2. Values

are represented as mean $\pm$ SEM of the triplicate assays from N=3 experiments. [H] Analysis of activation

of the ERK (p42/44) and NFκB (p65) signaling pathways in NT, KD, VC and OE macrophages by

immunoblotting with antibodies specific for the phosphorylated (active) and non- phosphorylated forms of

the proteins. Representative blot of one experiment out of 3 individual assays is shown. Expression of α-

TUBULIN was used as control. [I] ERK phosphorylation in macrophages with and without treatment with

ERK specific inhibitor U0126. Antibodies specific for the phosphorylated (active) and non-phosphorylated

forms of the proteins were used to probe cell extracts. Expression of α-TUBULIN was used as control. Blots

are representative of two independent experiments. [J-K] Expression of *IL1β* [J] and *TNFα* [K] in the 4 types

of macrophages after treatment with U0126 was analyzed by qPCR. Cells left untreated were used as

control. The relative gene expression folds in triplicate assay wells are represented with respect to *GAPDH*

as mean $\pm$ SEM for N=2. [L] The schematic of mutated catalytic site also depicted (D330A). Immunoblot

analysis of CMPK2-mCherry fusion protein using an antibody specific for mCherry protein in protein lysates

of VC, OE and D330A THP1 cells with mCherry specific antibody. GAPDH was used as control. [M]

Expression of *TNFα* and *IL1β* in THP1 cells after stable expression of vector alone (VC), full length CMPK2

(OE) and catalytic mutant of CMPK2 (D330A). The relative gene expression folds in triplicate assay wells

are represented with respect to *GAPDH* as mean $\pm$ SEM from N=4.

Increased inflammation of KD macrophages is associated with altered mitochondria and

augmented ROS

We presumed that identifying the precise localization of CMPK2 would reveal clues to its

molecular role in inflammation and mitochondrial membrane dynamics. We first validated the

mitochondrial localization of CMPK2 by immunoblotting subcellular fractions. With previous

reports indicative of the mitochondrial localization of CMPK2 (Xu et al., 2008) and our observation

of a majority of the protein in the mitochondrial fraction (Fig. 3A), we tested purified mitochondrial

fractions to pinpoint its location by standard biochemical and immunoblotting analyses. While

treatment with Proteinase K (removes the surface exposed proteins), completely removed the

outer membrane protein TOM20, nearly 30% of CMPK2 was present in the pellet fraction of intact

mitochondria, similar to that observed for the other mitochondrial membrane associated protein –

TIM50 (Fig. 3B) suggesting its association with the outer membrane of mitochondria. Most of the

matrix protein- SOD2 remained in the pellet and was lost only with complete solubilization with

Triton X-100. The peripheral association of CMPK2 with the mitochondrial membrane was also

revealed by a relatively higher amount of CMPK2 in the soluble fraction of mitochondria treated

with Na_2CO_3 in contrast with the integral membrane protein- TOM20 (Fig. 3C). We hypothesized

that two distinct possibilities could account for the enhanced basal levels of inflammation

observed in the CMPK2 expression altered cells (KD and OE) - 1) enhanced expression at all

times or 2) a temporal activation of the signaling with delayed decay kinetics. To test this, we

checked the expression of *IL1β* in the two cell lines during the monocyte to macrophage transition

upon PMA treatment. While the monocytes did not show any difference in *IL1β* expression,

activation with PMA induced distinct profiles of *IL1β* activation in the control (NT) or CMPK2

silenced (KD) cells. A rapid increase in *IL1β* within a day of activation (~700 fold) was followed by

a steady decline by day 3 of treatment further reaching basal levels by day 4 in the control cells

(Fig. 3D). CMPK2 silenced macrophages, however, displayed a protracted response kinetics with

a slower rise in expression levels ~1.5-2 folds lower than control cells at 24 h of activation with

PMA. Between 24h and 72h of treatment, the levels were 3-5 fold higher in KD with a delayed

decline of the levels to ~500 fold by day 4 and did not fall to basal levels even by day 5.

Macrophage inflammation is strongly associated with ROS activation. To check this, we compared

the levels of mitochondrial ROS in NT and KD macrophages by using a ROS specific stain-

MitoSOX Red. Mitotracker deep red was used as control for the levels of mitochondria in the two

cell types. While similar levels of mitotracker deep red staining hinted at unaltered mitochondrial

levels, the level of ROS (mitosox) was nearly 1.5-fold higher in KD cells as compared to control

NT macrophages (Fig. 3E). The importance of mito- ROS in enhanced inflammation in KD cells

was further evident when the cells were treated with a specific inhibitor- mitoquinone (MQ).

Addition of MQ early (immediately after PMA treatment) significantly reduced the expression of

IL1β in these cells (Fig. 3F); use of MQ at later stages of differentiation (48h and 72h after PMA

addition) did not alter the inflammation in the KD cells (Fig. S3A, B). Mitochondrial membrane

polarization is a critical component of organelle integrity and physiology with alterations in

membrane potential leading to ROS in macrophages (Suski et al., 2012; Vyssokikh et al., 2020).

To test if the ROS was associated with change in membrane potential, we compared the extent

of TMRE staining (a dye that shows a potential dependent differential partitioning across

mitochondrial membranes) in cells. As seen in Fig. 3G, the mitochondria of both KD and OE

macrophages exhibited significantly lower staining with TMRE, similar to the NT/VC cells treated

with the proton pump uncoupler- CCCP, suggestive of a loss of membrane integrity in the cells

even at the basal level.

Given the importance of mitochondrial fission-fusion dynamics as a stress mitigating factor we

tested if alteration in CMPK2 expression and the associated stress resulted in aberrant

mitochondrial replication dynamics. In line with the previous report (Youle and van der Bliek, 2012)

both KD and OE cells showed decreased levels of DNM1L (crucial for mitochondrial fission) in

comparison to NT and VC cells, respectively, (Fig. 3H), contrasting with an unaltered expression

of the mitochondrial fusion associated proteins (MFN1 and OPA1) in these cells (Fig. S3C, S3D).

This was further evident in the larger and more dense mitochondrial network of these organelles

in the KD and OE cells as compared to the small uniformly distributed mitochondria in the control

cells (Fig. 3I, 3J).

Fig. 3: Modulation of CMPK2 affects the mitochondrial physiology in macrophages.

[A] Expression of CMPK2 in subcellular fractions of THP1 macrophages by immunoblotting with specific

antibodies, T- total cell extract, C- cytoplasmic fraction and M- Mitochondria. Expression of cytosolic protein

α-TUBULIN and mitochondria resident protein VDAC are also represented. Blots are representative of three

independent experiments. [B] Mitochondrial fraction was subjected to Proteinase K treatment in the

presence or absence of Triton X-100 and given osmotic shock (OS) and analyzed by immunoblotting. Data

are representative of three independent experiments. [C] Mitochondrial fraction was incubated with

mitochondrial buffer with and without Na2CO3 and centrifuged at 13,000 rpm for 15 min. The pellet (P) and

supernatant (S) fractions were immunoblotted. Data are representative of three independent experiments.

[D] Kinetic profile of IL1β expression in NT and KD during differentiation of monocytes to macrophages.

Expression was checked at different time intervals after PMA treatment. Values are mean fold change in

expression with respect to GAPDH + SEM triplicate assays of N=2 experiments. [E] Analysis of

mitochondrial ROS in NT or KD macrophages. Cells were stained with MitoTracker Deep Red (as an

internal control) and ROS specific MitoSOX Red and specific populations were quantified by FACS. The

histogram plots of a representative experiment of N=4 is depicted. The extent of MitoSOX red mean

fluorescence intensity (MFI) + SEM is represented graphically in the inset. [F] Expression of IL1β in the

macrophages with the addition of a specific mitochondrial ROS inhibitor- MQ (1 μM) at day1 of PMA

treatment was analyzed by qPCR. Values are mean fold change in expression with respect to GAPDH +

SEM for triplicate assays of N=3 experiments. [G] Analysis of mitochondrial membrane potential in NT, KD,

VC and OE macrophages by TMRE staining. Fluorescence values normalized to protein in samples are

represented as mean fluorescent intensity + SEM for triplicate assays of N=3 experiments. [H] Expression

of DNM1L by immunoblotting with specific antibody was analyzed and is depicted along with the levels of

α-TUBULIN as a control. Relative intensity values are depicted as mean+ SEM of N=3. [I, J] Mitochondrial

architecture in control (NT) and KD [I] or VC and OE [J] macrophages were analyzed by confocal

microscopy. A representative image is depicted with the scale bar representing 10μm, the region shown in

higher magnification is depicted with the box.

Regulated levels of CMPK2 are critical for the normal metabolic activity of macrophages

In an attempt to understand the basis for this dysregulated inflammatory gene expression in both

the KD and OE macrophages, we performed global sequencing of transcripts and compared the

expression profiles with the control macrophages -NT and VC, respectively. Strikingly, again, on

comparison of the differentially expressed genes, inflammatory cytokine genes like *IL1β*, *IL8* and

TNFα were commonly upregulated in both the KD and OE macrophages (Fig. 4A). This

consistency in gene expression profiles was also evident with the significantly large numbers of

genes increased (515) or decreased (500) in macrophages with abnormal CMPK2 expression

(Fig. 4A inset). A significant deregulation of basal level inflammation was also evident in Gene

Set Enrichment Analysis (GSEA) with heightened inflammation response as a key gene family

common to the two macrophage lines (Fig. 4B). Interestingly, the hypoxia response was

highlighted as one of the highly upregulated gene family that was observed in both the KD and

OE macrophages. Most of the genes involved in the response were upregulated anywhere

between 2 and 16 folds in these cells (Fig. 4C). With hypoxia primarily regulated by HIF1α in cells,

we probed the levels of this protein in extracts of these macrophages. HIF1α levels were elevated

~2 folds in the OE cells as compared to the control VC cells, similarly higher level of HIF1α was

also observed in extracts of the KD cells (Fig. 4D).

Fig. 4

Fig. 4: Macrophages with dysregulated CMPK2 display increased hypoxia.

[A] Scatter plot of genes differentially expressed in KD or OE macrophages relative to the expression levels

in the control macrophages (NT or VC), respectively. Change in expression is depicted as $\log_2$ fold change

in expression. The number of genes up- and down- regulated in the macrophages are depicted as a Venn

diagram (inset). [B] GSEA hallmark pathway enrichment analysis of the commonly up and down regulated

genes in the KD and OE macrophages are represented as a bubble plot. X axis is the number of genes of

the pathway and size of the bubble depicts significance (-log P value). [C] Heat map of the hypoxia response

gene expression in the CMPK2 dysregulated macrophages KD and OE. The values represent $\log_2$ fold

change from the corresponding control cells. [D] Analysis of HIF1α in protein extracts of NT, KD, VC and

OE cells. Equal amounts of protein from the different extracts were probed with specific antibodies, a

representative blot of 3 independent experiments is depicted. The change in expression level of HIF1α with

respect to α-TUBULIN was calculated by densitometric analysis and is graphically represented.

Macrophages with dysregulated CMPK2 expression levels portray signatures of activated

M1 cells

The increased hypoxia, mitochondrial ROS and inflammation observed in the KD and OE cells

were reminiscent of M1 activated macrophages. To validate this, we analyzed the expression

profiles of previously identified signatures of M1 and M2 polarized macrophages in our gene sets

(Martinez et al., 2006). In agreement with our hypothesis, both the KD and OE macrophages

showed an increase in the expression of genes of M1 macrophages with a majority of the M2

specific signatures either unchanged or with reduced expression levels (Fig. 5A). Further

validation of a M1 bias was visible in the physiological state of the cells. Analysis of respiration

rates of cells revealed a distinct pattern with a considerable decrease in the oxygen consumption

rates of macrophages with altered expression of CMPK2. Both the KD and OE macrophages

displayed 2-2.5 folds lower oxygen consumption with delayed kinetics as opposed to the control

macrophages NT and VC (Fig. 5B) implicating a strong shift of macrophages with altered CMPK2

expression towards a pro-inflammatory phenotype and the important role for CMPK2 in

maintaining homeostasis of macrophages. Further, both these cell types harbored increased

expression of genes involved in glycolysis (Fig. S4A, 5C) that manifested as a significant increase

in the levels of glycolysis intermediates like fructose bis-phosphates, DHAP, culminating in

markedly high levels of lactate (Fig. 5C). Recent studies have demonstrated a strong link between

serine biosynthesis/ metabolism to the inflammatory profiles and mitochondrial physiology of cells

(Kurita et al., 2021; Wang et al., 2020; Yu et al., 2019). Strikingly, the serine- glycine biosynthetic

pathway was one of the dominant metabolic axes downregulated in these 2 cell types in

comparison with the control cells (Fig. S4B). The expression levels of the key enzymes involved

in this pathway was validated by qRTPCR. Consistently, in both the KD and OE cells, expression

levels of *PHGDH*, *PSAT1* and *PSMT1* were significantly lower in comparison to the control NT

and VC cells (Fig. 5C). This decrease in expression levels is reflected as ~2-fold decrease at the

metabolite level of serine in the KD and OE cells. Given an important role for PHGDH in controlling

inflammation under hypoxic conditions(Huang et al., 2021; Wilson et al., 2020) and our

observation of a significant decrease in expression levels combined with hypoxic conditions and

heightened pro-inflammatory gene expression in the KD and OE cells, we hypothesized an

important role for this axis in the observed phenotypes in this study. To investigate this, we

specifically inhibited PHGDH in THP1 Wt cells by the specific inhibitor- NCT-503. As a control,

cells were treated with an inactive inhibitor of PHGDH and evaluated the expression profiles of

IL1β and *TNFα* in these cells. Use of the inactive inhibitor did not alter the expression from the

untreated cells, in contrast to the significant increase in transcript levels by ~5 and 2.5 folds for

these genes with the inhibitor, substantiating our supposition (Fig. 5D).

Fig. 5

Fig. 5: Macrophages with dysregulated CMPK2 are metabolically similar to activated M1 macrophages with higher glycolytic flux.

[A] The expression patterns of genes specific to M1 and M2 activated macrophages in the CMPK2 dysregulated macrophages KD and OE are represented as heat maps. The values represent $\log_2$ fold change from the corresponding control cells. [B] Continual estimation of oxygen consumption in NT, KD and VC, OE macrophages until 3h with Oroboros oxygraph. The level of oxygen consumption was calculated and mean OCR $\pm$ SEM of N=3 experiments is shown. [C] Metabolite levels in the CMPK2

silenced (KD) and over-expression (OE) and their respective control macrophages were assayed from

cellular extracts by MS. The levels of individual metabolites in the KD and OE macrophages are represented

as relative abundance compared to control cells as mean $\pm$ SEM from 2/3 independent experiments (N=3).

Expression of genes involved in the glycolysis and serine to glycine biosynthetic pathway estimated by

qRT-PCR was also shown. [D] Expression of *IL1β* and *TNFα* in THP1 cells treated with the serine

biosynthesis inhibitor-NCT-503 and its inactive form-PHGDH-inactive. Expression values are mean fold

change in expression with respect to *GAPDH* $\pm$ SEM triplicate assays of N=2 experiments after 6h of

treatment.

CMPK2 plays a critical role in the macrophage ability to control infection

To understand the impact of altered CMPK2 expression on macrophage inflammatory properties,

we evaluated its ability to control the intracellular growth of Mtb. We observed a 2-3 fold decrease

in intracellular bacterial numbers in these macrophages as compared to NT late in infection (day

5), KD cells were able to control bacterial levels to input levels by this timepoint compared to

growth of about 5-6 folds in the NT macrophages (Fig. 6A). This pattern of increased bacterial

control was also evident in STM infected KD macrophages with 2-3 fold lower bacterial numbers

after 24 h of infection (Fig. 6B).

Fig. 6

398 **Fig. 6: CMPK2 regulates the bactericidal activity of macrophages**

399 [A] Growth kinetics of Mtb in NT and KD macrophages at different times post infection at a MOI of 5 for 6h.

400 Values are mean CFU $\pm$ SEM values in triplicate assays of N=4. [B] Growth kinetics of STM in NT and KD

401 macrophages at different times post infection at a MOI of 10 for 20 min. Values are mean CFU $\pm$ SEM

402 values in triplicate assays of N=4.

Discussion

Macrophages, the primary cells that sense and respond to any infectious agents or tissue damage, are critical components of vertebrate innate immune responses. Being endowed with the capacity to initiate and manifest strong infection control programs to a variety of pathogens, they often dictate both acute and chronic outcomes of infection in the host (Gong et al., 2020; Mosser et al., 2021). Recent evidences also hint at an important role for these cells in maintenance of tissue homeostasis during steady state and inflammation (Krenkel and Tacke, 2017; Mosser *et al.*, 2021; Oishi and Manabe, 2018; Watanabe et al., 2019). Numerous studies have identified functionally distinct macrophage sets with polarized pro-inflammatory (M1) or anti-inflammatory (M2) phenotypes with characteristic metabolic states (Galvan-Pena and O'Neill, 2014; Tan et al., 2015; Wang et al., 2018).

With a central role of mitochondria in cellular metabolism and physiology, it is logical to presume stringent control of organelle integrity in normal as well as in response to stress. Over the last few years, infection dependent modulation of mitochondrial dynamics and function have been demonstrated as key determinants of host cell response kinetics and immune signaling (Benmoussa et al., 2018; Mehta et al., 2017; Mills *et al.*, 2016; Ramond et al., 2019). A recent study has demonstrated a significant lowering of energy metabolism in macrophages on account of the strong type I IFN responses induced by Mtb infection (Olson et al., 2021). With the mitochondrial CMPK2, an ISG, one of dominantly expressed proteins in infected macrophages, we sought to investigate the importance of this gene in macrophage response dynamics in human macrophages.

We provide evidence for a critical role of the mitochondrial gene CMPK2 in regulating organelle integrity and macrophage physiology. Altered expression of this gene led to disruption of the normal mitochondrial physiology as evidenced by decreased membrane potential and increased ROS. This further translated into marked dysregulation of basal inflammation in the cells with enhanced activation of ERK signaling leading to a marked increase in production of pro-

inflammatory mediators like TNFα, IL8 and IL1β. In line with a recent study of the importance of

increased lactate production by M1 polarized macrophages in microbicidal capacity (Ó

Maoldomhnaigh et al., 2021), both the KD and OE cells also portrayed a significant shift in their

metabolic status towards rapid energy production via glycolysis and consequent accumulation of

lactate in the cells. Our observation of identical phenotypic characteristics of CMPK2 silenced as

well as overexpression cells (enhanced mitochondrial stress even at the basal state) hinted at the

possibility of a fundamental role of this gene in organelle function and integrity. It is also interesting

to assume an association of CMPK2 in a larger protein complex in cells, in which changes in

expression of one component would affect the overall stoichiometric composition of the complex

and its function as reported earlier in the case of SRPK1 in cancerous cells (Wang et al., 2014).

The association of CMPK2 with the mitochondrial membrane in proximity to the mitochondrial

ETC proteins coupled to our observed decrease in the levels of complex 1 in CMPK2 dysregulated

(KD and OE) macrophages (Fig. S5), provides initial evidence for this hypothesis. The observed

impairment in oxygen consumption in these macrophages and a concomitant increase of the

glycolytic metabolite- lactate further supports the involvement of CMPK2 in central metabolic

activities of macrophages.

We observed that the removal of MLS from CMPK2 destabilized the protein and resulted in

enhanced degradation by the cellular proteolytic machinery. This observation is in sync with

previous reports(Ravanelli et al., 2020) demonstrating that mitochondrial proteins without signals

for mitochondrial partitioning tend to form misfolded aggregates that are rapidly degraded by the

host proteasome (Wrobel et al., 2015). We also observed an absolute requirement of kinase

activity for CMPK2 function and the importance of the N- terminal domain in protein stability (data

not shown, part of another manuscript).

A previous study identified an important role for serine catabolism in the maintenance of cellular

redox under hypoxic conditions. Repression of serine catabolism by silencing SHMT2 in cancer

cells resulted in enhanced ROS under hypoxia (Ye et al., 2014). We also observed significant

downregulation of genes involved in serine biosynthesis and conversion to glycine in

macrophages with altered expression of CMPK2. In addition, we also see enhanced ROS and

hypoxia in these macrophages. Our observation of a significant decrease in SHMT1 expression

without any significant change in SHMT2 in the KD and OE macrophages highlights another

possible link between serine catabolism and cellular redox maintenance via the regulated

expression of CMPK2.

Serine metabolism has been shown to affect cellular physiology including inflammation in

macrophages. Studies have suggested an inflammation promoting role for serine biosynthesis in

macrophages with serine depletion in the culture media actively stunting the expression of pro-

inflammatory genes like TNF and IL1β (Chen et al., 2020; Rodriguez et al., 2019). While these

studies signify the importance of extracellular serine and its uptake by cells, intracellular serine

biosynthesis and its conversion to glycine by the serine glycine and one-carbon pathway

(SGOCP) genes in modulating macrophage phenotypes is yet not understood. A recent robust

study has demonstrated that Phgdh mediated serine synthesis is required for expression of anti-

inflammatory genes and inhibition of Phgdh resulted in enhanced expression of pro-inflammatory

genes (Wilson *et al.*, 2020).

To validate the PHGDH role in enhanced inflammatory gene expression, we inhibited PHGDH

and observed enhanced expression of pro-inflammatory genes in macrophages. This is in

concordance with a previous study by Wilson et al., wherein M2 (anti-inflammatory) macrophages

showed enhanced Phgdh and Psat1 expression (Wilson *et al.*, 2020). In fact, treatment with LPS

that polarizes the cells to an M1 phenotype reduced Phdgh expression in macrophages while IL4

enhanced the expression again emphasizing the association of serine biosynthesis with the anti-

inflammatory macrophages. Further inhibition of PHGDH resulted in heightened pro-inflammatory

response as observed in the KD and OE macrophages.

Given the importance of CMPK2 in maintaining mitochondrial architecture, function and in

enhancing overall inflammation, macrophage would actively induce this gene as a response to

intracellular bacterial infection. With bacterial contact immediately stimulating host cell surface

receptors, it is not surprising that CMPK2 would be induced in response to activation of more

central TLR4 signaling that primarily leads to activation of inflammatory programs in the infected

macrophages.   Moreover, our data of a long-term breakdown of inflammation control in

macrophages following prolonged disturbance of CMPK2 expression levels only highlights the

importance of rapid and robust regulation of this gene by cells for reverting to basal status. This

supports our hypothesis that CMPK2 is an important component of the cellular inflammation

rheostat controlling macrophage metabolic and transcriptional response to infections. A recent

study has defined the criticality of cellular pyrimidine levels in mitochondrial DNA mediated

regulated innate immunity (Sprenger et al., 2021). Given the putative role of CMPK2 in the

production of CTP/UTP, and the dysregulation of serine metabolism on CMPK2 imbalance,

another important biosynthetic node for nucleotide biosynthesis in the cell, one can envisage a

similar situation of an altered cellular *vis-a-vis* mitochondrial nucleotide levels. Comprehensive

analysis towards pinpointing the precise molecular mechanism of CMPK2 mediated regulation of

macrophage immuno- metabolism would help develop novel mechanisms of modulating innate

response control paving the way for future host cell directed therapeutics.

Acknowledgments: The authors thank CSIR (VR-BSC0123, MLP2012), and DBT (SG-

GAP0088) funding agency for supporting the study. The authors thank Mr. Manish Kumar and

CSIR- BSC0403 for the confocal microscopy facility, CSIR-STS0016 for BSL3 facility. The authors

wish to thank Dr. Soumya Sinha Roy, CSIR-IGIB for the plasmid pmtDsRed. The authors thank

PA's doctoral advisory committee members for useful discussions and comments on the project.

The mass spectrometry facilities are duly acknowledged. The student fellowships from CSIR: PA-

CSIR- SRF, RA, PA- CSIR-BSC0124, MC- CSIR-JRF, India. The authors thank Kanika Bisht and

Mahima Madan for proof reading of the manuscript.

Author Contributions: PA, SR, SG, and VR were involved in conceptualizing and design of the

work. Experiments were performed by PA, MC, TR and DS. PA and RC performed Mass

spectrometry, PA, SG, and VR wrote the manuscript.

Conflict of interest: The authors do not have any competing interests.

Materials and Methods

Reagents

The following chemicals were purchased from Sigma-Aldrich: U0126 (U120), RNAzol RT

(R4533), Zymosan (Z4250), Thermo Fisher Scientific: MitoSOX Red (M36008), MitoTracker Deep

Red (M22426) and Invivogen: LPS (tlrl-smlps), CLI095 (tlrl-cli095), poly I:C (tlrl-pic), Pam3CSK4

(tlrl-pms). Mitoquinol (89950), PHDGH-inactive (19717) and NCT-503 (19718) were purchased

from Cayman Chemical Company.

Cell culture

THP-1 cells were cultured in RPMI-1640 (Himedia laboratories, Mumbai, India) media

supplemented with 10% FBS and 1mM sodium pyruvate (Himedia laboratories, Mumbai, India).

HEK293T cells were cultured in DMEM supplemented with 10% FBS and 1mM sodium pyruvate.

For differentiation, THP-1 cells were treated with 100nM PMA for 24hr and rested for 48hr before

the next manipulation. Both the cell lines were confirmed for their identity by STR PCR analysis

and maintained mycoplasma free by routine monitoring. LPS treatment of macrophages was done

with 10 ng/ml LPS for 6h. TLR4 signaling was inhibited by treating with 10 µM of CLI095 before

infection or LPS treatment. For ERK inhibition, THP-1 macrophages were treated with 2 µM of

U0126 for 6h. To inhibit serine biosynthesis macrophages were treated with 10 µM NCT-503 for

6h and 10 µM PHGDH-inactive compound was used as negative control.

Generation of THP1 lines with CMPK2 silencing

To generate knockdown cells, HEK293T cells were co-transfected with CMPK2 siRNA plasmid

(Applied Biological Materials Inc, Richmond, Canada), packaging plasmid (pCMV-dR8.2) and

envelope plasmid (pCMV-VSV-G) using Lipofectamine-LTX reagent (Thermo Fisher Scientific

India Pvt. Ltd, Mumbai, India). After 6h of transfection, media was removed, and fresh media was

added to the cells. After 48h of transfection, the culture supernatant was collected, centrifuged at

500 rpm to remove cell debris. The virus particles in the supernatant were concentrated to 100µl

using Amicon® Ultra-15 Centrifugal Filters (Sigma-Aldrich Chemicals Private Limited, Bengaluru,

India) and used to infect THP1 cells. Cells were then selected with 0.6 µg/ml puromycin and GFP

positive clones were used for further studies.

Bacterial culturing and infection

Mycobacterium tuberculosis strain Erdman was grown in 7H9 Middlebrook media BD

Biosciences, USA) supplemented with Middlebrook ADC (BD, USA) at 37°C. *E. coli and*

Salmonella enterica serovar Typhimurium were grown in Luria-Bertani (LB) media (Himedia

Laboratories, Mumbai, India) at 37°C. Single cell suspension of mid-log phase bacteria was

adjusted to the required cell density and then used to infect the differentiated THP-1 macrophages

at multiplicity of infection (MOI) of 5. After 6h *p.i.* for Mtb, cells were washed with PBS to remove

extracellular bacteria, and at various times post infection, bacterial numbers were enumerated by

lysis of cells and plating for CFU in 7H10 agar plates. For *S. typhimurium/ E. coli*, post addition of

bacteria at a MOI of 10, the plates were centrifuged for 5 min and then left for 20 min at 37°C. The

media was removed and treated with 100µg/ml of gentamicin for 2h in RPMI at 37°C. The cells

were washed with PBS for 3 times, and infection was continued in media containing 12µg/ml of

gentamicin and at indicated time points cells were lysed with PBS-Triton-X-100 (1%) for CFU

plating on LB agar (Himedia laboratories, Mumbai, India) plates. The colonies were counted and

is represented as CFU/well.

Gene expression analysis by qPCR

Total RNA was isolated using RNAzol (Sigma-Aldrich, USA) method and the concentration was

quantified using Nanodrop 2000 UV-visible spectrophotometer. cDNA was prepared with 250-

1000 ng total RNA by a RT-PCR using Verso cDNA synthesis kit (Thermo Fisher Scientific, USA).

qPCR was performed on a Roche LC480 II system using DyNAmo Flash SYBR Green mix

(Thermo Fisher Scientific, USA). Primer sequences are given in Table 1.

Cloning of CMPK2 and the D330 variant.

The NFkB promoter from pNFkB-d2GFP was excised by digestion with *Ecl*136II and *Hind*III and

the mCherry fragment from pmCherry-N1was removed with *Hind*III and *Hpa*I and ligated to *Nru*I

and *Eco*RV digested pcDNA3.1(+) to get the plasmid pPRAM1. The NFkB promoter from

pPRAM1 was replaced with EF-1α from pTracer-EF/V5 His A by using *Mlu*I and *Eco*RI to get

pPRAM2. The CMPK2 fragments were amplified from THP1 cDNA using the primer sets PP1 and

PP2 and cloned into the pPRAM2 using *Kpn*I and *Bam*HI. The D330A catalytic site mutant was

prepared by site directed mutagenesis using the primers PP3 and PP4. The sequences were

verified by sequencing. The list of primers used in this study is given in table 1.

Western blotting and ELISA

For immunoblotting, cells were lysed in RIPA buffer, resolved by SDS-PAGE and transferred to

nitrocellulose membrane. After blocking with 5% BSA in Tris Buffered Saline Tween (TBST) for

1hr at room temperature, membranes were incubated with the appropriate dilution of the primary

antibody overnight, washed with PBS and probed with infrared dye conjugated secondary

antibody and developed in the LI-COR Odyssey platform. The following antibodies were

purchased from Cell Signaling Technology: Rabbit anti-α-tubulin (2125), Rabbit anti-phospho-

p42/44 (4377), Rabbit anti-p42/44 (4695), Rabbit anti-phospho-NFkB p65 (3033), Rabbit anti-

NFkB p65 (4764), Rabbit anti-phospho-JNK (4671), Rabbit anti-JNK (9258), Abcam: Rabbit anti-

mCherry (ab167453), Mouse anti-TOM20 (ab56783), Rabbit anti-TIM50 (ab109436), Mouse anti-

MFN1/2 (ab57602), Mouse anti-OPA1 (ab194830), Rabbit anti-DRP1 (ab140494), Rabbit anti-

HIF1α (ab51608) BD Bioscience: Rabbit anti-VDAC (D73D12), Mouse anti-SOD2 (611581) and

LI-COR Biotechnology: Goat anti-Rabbit IgG 800 (926-32211), Goat anti-Mouse IgG 680 (926-

68070). Rabbit anti-CMPK2 (HPA041430) was purchased from Sigma-Aldrich.

For ELISA, either cell supernatants or cell extracts were used for cytokine estimation with specific

kits- TNFα (cat no: 88-7346-77, eBioscience) and IL1β (cat no: 557953, BD Biosciences) ELISA

kits according to manufacturer's protocol.

Analysis of ROS by FACS

THP1 monocyte derived macrophages (MDM) were removed with PBS and 4mM EDTA, washed

with PBS, resuspended in RPMI media and used for staining with 5µM MitoSOX Red and 100nM

MitoTracker Deep Red at 37°C for 30 mins. Cells were then washed twice in HBSS and finally

resuspended in 500µl HBSS and analyzed in FACS ARIA II (BD Biosciences).

Metabolite measurement

Intracellular metabolites for MS-based targeted metabolomics were extracted using methanol-

water. Briefly, 1.2×10^6 cells from each condition were washed three times with ice-cold PBS, and

then quenched with methanol-water (4:1). The cell suspension was freeze thawed in liquid

nitrogen three times. The suspension was centrifuged at 15,000 g at 4°C for 10 min. Supernatant

was collected, vacuum dried and then reconstituted in 50 µl of 50% methanol. The reconstituted

mixture was centrifuged at 15,000 g for 10 min, and 5 µl was injected for LC–MS/MS analysis.

The data were acquired using a Sciex Exion LCTM analytical UHPLC system coupled with a triple

quadrupole hybrid ion trap mass spectrometer (QTrap 6500; Sciex) in negative ion mode.

Samples were loaded onto an Acquity UPLC BEH HILIC (1.7 µm, 2.1 × 100 mm) column, with a

flow rate of 0.3 ml/min. The mobile phases comprising of 10 mM ammonium acetate and 0.1%

formic acid (buffer A) and 95% acetonitrile with 5 mM ammonium acetate and 0.1% formic acid

(buffer B). The linear mobile phase was applied from 95% to 20% of buffer A. The gradient

program was used as follows: 95% buffer B for 1.5 min, 80%–50% buffer B in next 0.5 min,

followed by 50% buffer B for next 2 min, and then decreased to 20% buffer B in next 50 s, 20%

buffer B for next 2 min, and finally again 95% buffer B for next 2 min. After data acquisition, peaks

corresponding to each metabolite were extracted using Sciex Multiquant TM v.3.0 software and

the area was exported in an excel sheet. Normalization was performed using total area sum of

that particular run.

Oxygen Consumption

609   3-4 x 10^6 cells of THP1 MDMs in complete RPMI were added to the chambers of the Oxygraphy-

610   2k (O2k, Oroboros Instruments, Innsbruck, Austria). The oxygen concentration was measured

611   over time at 37⁰C under constant stirring. The oxygen consumption rate was calculated using XY

612   graph of oxygen consumption and time.

Transcriptome analysis

Total RNA was isolated using RNAzol (Sigma Aldrich, India) and transcript sequencing was done

commercially by Bencos research solutions private ltd., (Bengaluru, India). cDNA libraries were

generated using Truseq RNA Library Prep Kit (Illumina San Diego, USA) and sequenced in an

Illumina Novaseq 6000 platform. RNA-seq reads were aligned with hg38 genome using STAR

(Dobin et al., 2013). HTseq-count was used to count the transcripts (Anders et al., 2015). The

differential expression analysis across samples was analyzed using GSEA (Subramanian et al.,

2005).

Confocal Microscopy and Image analysis of stained cells

THP1 monocytes (stable transfectants of non-targeting or CMPK2 specific siRNA) were

transfected with plasmid DNA mtDsRed (kind gift by Dr. Soumya Sinha Roy, CSIR- IGIB) and

selected for stably expressing clones with 400µg/ml of G418. Cells were differentiated on

coverslips, fixed with 4% formaldehyde and imaged in a Leica 480 confocal microscopy. For live

imaging, monocytes overexpressing CMPK2 were plated at a cell density of 0.3 x 10^6 cells/ml in

the cell culture dish containing two chambers per dish (1 ml/ chamber). Mitotracker Green dye

was added to the media and the cells were kept in 37⁰C CO_2 incubator for 20mins. The cells were

imaged on a Leica SP8 confocal microscope.

Mitochondrial localization

Differential centrifugation was employed for mitochondria isolation from the cells (Frezza et al., 2007). Briefly, 2.5×10^6 THP1 MDMs were washed twice with PBS and resuspended in 3 ml ice-cold cell isolation buffer (IBc- containing 10mM Tris pH to 7.4, 1mM EGTA pH to 7.4 and 0.2M sucrose). The cells were homogenized using a glass Teflon pestle potter. The homogenate was then centrifuged at 600g for 10 min at 4ºC, the supernatant was collected and centrifuged at 7000g for 10 min at 4ºC. The pellet was then washed and resuspended in 200 µl IB_C and transferred to a 1.5ml Eppendorf tubes. The homogenate was again centrifuged at 7000g for 10 min at 4ºC. The supernatant was discarded and the pellet was resuspended to get the mitochondrial fraction.

For Proteinase K digestion, the mitochondria were resuspended in MS buffer (210mM mannitol, 70mM sucrose, 5mM Tris-HCL-pH 7.5, 1mM EDTA-pH 7.6)/ 20mMHEPES with or without 2 mg/ml Proteinase K (for 15 min) and/or 1% Triton-X-100. The reaction was terminated by the addition of 5 mM phenylmethylsulfonyl fluoride. The mixture was centrifuged at 15000g for 15 min and the pellet fraction was collected. For analysis of membrane proteins, the mitochondrial fraction was resuspended in MS buffer with and without 0.1M Na_2CO_3 and incubated in ice for 30 min. The pellet and supernatant fractions were collected at 15000g for 15 min. All fractions were analyzed by immunoblotting with respective antibodies.

Graphs and Statistical analysis

Statistical analyses were performed by using the two tailed Student's t-test. GraphPad Prism software was used for graphs and statistical analysis.

DATA availability statement: The RAW data of RNA-seq experiments of this study have been uploaded in NCBI-GEO with accession number -GSE199951.

658   **Table 1: List of primers used in this study**

Name	Primer sequence
CMPK2 F	CCAGGTTGTTGCCATCGAAG
CMPK2 R	CAAGAGGGTGGTGACTTTAAGAG
IL8 F	AGACAGCAGAGCACACAAGC
IL8 R	ATGGTTCCTTCCGGTGGT
GAPDH F	GAAGGTGAAGGTCGGAGTC
GAPDH R	GAAGATGGTGATGGGATTTC
IL1β F	CCTGTCCTGCGTGTTGAAAGA
IL1β R	GGGAACTGGGCAGACTCAAA
TNFα F	CCCCAGGGACCTCTCTCTAATC
TNFα R	GGTTTGCTACAACATGGGCTACA
IP10 F	TCCACGTGTTGAGATCATTGC
IP10 R	GGCCTTCGATTCTGGATTCAG
IL10 F	GCTGGAGGACTTTAAGGGTTACCT
IL10 R	CTTGATGTCTGGGTCTTGGTTCT
VEGFα F	CTGCTGTCTTGGGTGCATTG
VEGFα R	CCATGAACTTCACCACTTCG
PP1	ATGGTACCATGGCCTTCGCCCGCCGGCTC
PP2	CGTCTAGAGGATCCGCCGGTTCACTAAAACTATTCTGG
PP3	CAAATCTCCTGTGATTGTAGCCAGGTACTGGCACAGCAC
PP4	GTGCTGTGCCAGTACCTGGCTACAATCACAGGAGATTTG

660   **Supplementary Figures:**

Fig. S1

666   **Fig. S1:** Expression of *CMPK2* in THP1 macrophages upon Mtb infection in the presence or absence of

667   CLI095.

Fig. S2

Fig. S2: [A] Expression of *CMPK2* in THP1 macrophages stably expressing the non-targeting (NT) siRNA

or three different siRNAs (a-c) against CMPK2 by qPCR. The gene expression was normalized with *GAPDH*

and relative fold changes compared to NT are represented as mean ± SEM (N=2). [B] Expression of *TNFα*

and *IL1β* gene in NT and KD macrophages following LPS stimulation for 6h. The relative gene expression

folds in triplicate assay wells are represented with respect to *GAPDH* as mean ± SEM for N=2. [C]

Expression of *CMPK2* in OE or empty vector -VC cells was analyzed by qPCR. The relative gene

expression folds in triplicate assay wells are represented with respect to *GAPDH* as mean ± SEM for N=3.

[D] CMPK2 expression in VC and OE cells was analyzed by immunoblotting with tag specific (mCherry)

antibody. [E] Relative quantitation of p-ERK and p-NFκB in the NT, KD, VC, OE cells by densitometric

analysis of immunoreactivity is shown. Values are mean + SEM of triplicate (N=3) independent blots. [F]

Analysis of activation of the p38 MAPK and SAPK/JNK signaling pathways in NT and KD macrophages by

immunoblotting with antibodies specific for the phosphorylated (active) and non- phosphorylated forms of

the proteins. The relative intensities of the blots in the CMPK2 silenced cells w.r.t control (NT) cells are

represented as mean ± SEM of N=3. Expression of α-TUBULIN was used as control. [G] Expression of

CMPK2 in VC, OE and D330A cells was analyzed by qPCR and is depicted relative to GAPDH levels of

N=3 assays.

Fig. S3

Fig. S3: [A, B] Expression of *IL1β* in the macrophages with the addition of a specific mitochondrial ROS

inhibitor- MQ at day 2 [A] and day 3 [B] of PMA treatment intervals post activation was analyzed by qPCR.

Values are mean fold change in expression with respect to *GAPDH* $\pm$ SEM for triplicate assays of N=2/3

experiments. [C, D] Expression of proteins involved in mitochondrial fusion by immunoblotting with specific

antibodies. MFN1&2 and OPA1 were checked in NT, KD [C] or VC, OE [D] cells along with α-TUBULIN

levels as a control. Relative intensity values are depicted $\pm$ SEM of N=3.

Fig. S4

Fig. S4: [A] Expression of genes involved in glycolysis and TCA was represented as heatmap. The expression was extracted from the RNA sequencing data of CMPK2 dysregulated THP1 cells. The values are represented as $\log_2$ compared to respective control cells. [B] Panther gene family enrichment analysis of the commonly up and down regulated genes in the KD and OE macrophages is represented as a bubble plot. X axis is the number of genes of the pathway.

Fig. S5

Fig. S5: Analysis of expression of protein complexes of the mitochondrial respiratory chain in cell lysates of NT and KD macrophages by immunoblotting. α-TUBULIN (T) and VDAC1 (VD) were used as loading controls. The band intensities of the complex proteins were normalized to α-TUBULIN and the relative intensity of KD w.r.t. NT is represented as fold change $\pm$ SEM for N=2.

References

Alphonse, N., Dickenson, R.E., and Odendall, C. (2021). Interferons: Tug of War Between Bacteria and
Their Host. Front Cell Infect Microbiol *11*, 624094. 10.3389/fcimb.2021.624094.
Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics *31*, 166-169. 10.1093/bioinformatics/btu638

btu638 [pii].
Benmoussa, K., Garaude, J., and Acin-Perez, R. (2018). How Mitochondrial Metabolism Contributes to
Macrophage Phenotype and Functions. J Mol Biol *430*, 3906-3921. S0022-2836(18)30743-5 [pii]

10.1016/j.jmb.2018.07.003.
Blagih, J., and Jones, R.G. (2012). Polarizing macrophages through reprogramming of glucose metabolism.
Cell Metab *15*, 793-795. 10.1016/j.cmet.2012.05.008

S1550-4131(12)00201-X [pii].
Boxx, G.M., and Cheng, G. (2016). The Roles of Type I Interferon in Bacterial Infection. Cell Host Microbe
19, 760-769. 10.1016/j.chom.2016.05.016

S1931-3128(16)30212-8 [pii].
Chen, S., Xia, Y., He, F., Fu, J., Xin, Z., Deng, B., He, L., Zhou, X., and Ren, W. (2020). Serine Supports IL-
1beta Production in Macrophages Through mTOR Signaling. Front Immunol *11*, 1866.
10.3389/fimmu.2020.01866.
Cohen, H.B., and Mosser, D.M. (2013). Extrinsic and intrinsic control of macrophage inflammatory
responses. J Leukoc Biol *94*, 913-919. 10.1189/jlb.0413236

jlb.0413236 [pii].
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras,
T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.
10.1093/bioinformatics/bts635

bts635 [pii].
El-Diwany, R., Soliman, M., Sugawara, S., Breitwieser, F., Skaist, A., Coggiano, C., Sangal, N., Chattergoon,
M., Bailey, J.R., Siliciano, R.F., et al. (2018). CMPK2 and BCL-G are associated with type 1 interferon-
induced HIV restriction in humans. Sci Adv *4*, eaat0843. 10.1126/sciadv.aat0843

aat0843 [pii].
Feng, C., Tang, Y., Liu, X., and Zhou, Z. (2021). CMPK2 of triploid crucian carp is involved in immune defense
against bacterial infection. Dev Comp Immunol *116*, 103924. S0145-305X(20)30479-1 [pii]

10.1016/j.dci.2020.103924.
Frezza, C., Cipolat, S., and Scorrano, L. (2007). Organelle isolation: functional mitochondria from mouse
liver, muscle and cultured fibroblasts. Nat Protoc *2*, 287-295. nprot.2006.478 [pii]

10.1038/nprot.2006.478.
Galvan-Pena, S., and O'Neill, L.A. (2014). Metabolic reprograming in macrophage polarization. Front
Immunol *5*, 420. 10.3389/fimmu.2014.00420.
Gong, T., Liu, L., Jiang, W., and Zhou, R. (2020). DAMP-sensing receptors in sterile inflammation and
inflammatory diseases. Nat Rev Immunol *20*, 95-112. 10.1038/s41577-019-0215-7

10.1038/s41577-019-0215-7 [pii].

Hassan, M.Z., Ahmed, M.S., Khan, M.M., Uddin, M.A., Chowdhury, F., and Kamruzzaman, M. (2019).
Genomic profiling of Nipah virus using NGS driven RNA-Seq expression data. Bioinformation *15*, 853-862.
10.6026/97320630015853

97320630015853 [pii].
Huang, H., Liu, K., Ou, H., Qian, X., and Wan, J. (2021). Phgdh serves a protective role in Il1beta induced
chondrocyte inflammation and oxidativestress damage. Molecular medicine reports *23*.
10.3892/mmr.2021.12058.
Kelly, B., and O'Neill, L.A. (2015). Metabolic reprogramming in macrophages and dendritic cells in innate
immunity. Cell Res *25*, 771-784. 10.1038/cr.2015.68

cr201568 [pii].
Kikuchi, Y., Miyauchi, M., Iwano, I., Kita, T., Oomori, K., and Kizawa, I. (1988). Adjuvant effects of
prostaglandin D2 to cisplatin on human ovarian cancer cell growth in nude mice. Eur J Cancer Clin Oncol
24, 1829-1833. 10.1016/0277-5379(88)90093-4.
Kim, H., Subbannayya, Y., Humphries, F., Skejsol, A., Pinto, S.M., Giambelluca, M., Espevik, T., Fitzgerald,
K.A., and Kandasamy, R.K. (2021). UMP-CMP kinase 2 gene expression in macrophages is dependent on
the IRF3-IFNAR signaling axis. PLoS One *16*, e0258989. 10.1371/journal.pone.0258989

PONE-D-21-16514 [pii].
Kovarik, P., Castiglia, V., Ivin, M., and Ebner, F. (2016). Type I Interferons in Bacterial Infections: A Balancing
Act. Front Immunol *7*, 652. 10.3389/fimmu.2016.00652.
Krenkel, O., and Tacke, F. (2017). Liver macrophages in tissue homeostasis and disease. Nat Rev Immunol
17, 306-321. 10.1038/nri.2017.11

nri.2017.11 [pii].
Kumar, R., Singh, P., Kolloli, A., Shi, L., Bushkin, Y., Tyagi, S., and Subbian, S. (2019). Immunometabolism
of Phagocytes During Mycobacterium tuberculosis Infection. Front Mol Biosci *6*, 105.
10.3389/fmolb.2019.00105.
Kurita, K., Ohta, H., Shirakawa, I., Tanaka, M., Kitaura, Y., Iwasaki, Y., Matsuzaka, T., Shimano, H., Aoe, S.,
Arima, H., et al. (2021). Macrophages rely on extracellular serine to suppress aberrant cytokine
production. Scientific reports *11*, 11137. 10.1038/s41598-021-90086-w.
Labadarios, D., Cilliers, J., Visser, L., Van Stuijvenberg, M.E., Shephard, G.S., Wium, D., and Walker, R.
(1987). Vitamin A in acne vulgaris. Clin Exp Dermatol *12*, 432-436. 10.1111/j.1365-2230.1987.tb01942.x.
Labbe, K., Murley, A., and Nunnari, J. (2014). Determinants and functions of mitochondrial behavior. Annu
Rev Cell Dev Biol *30*, 357-391. 10.1146/annurev-cellbio-101011-155756.
Lai, J.H., Hung, L.F., Huang, C.Y., Wu, D.W., Wu, C.H., and Ho, L.J. (2021a). Mitochondrial protein CMPK2
regulates IFN alpha-enhanced foam cell formation, potentially contributing to premature atherosclerosis
in SLE. Arthritis Res Ther *23*, 120. 10.1186/s13075-021-02470-6

10.1186/s13075-021-02470-6 [pii].
Lai, J.H., Wu, D.W., Wu, C.H., Hung, L.F., Huang, C.Y., Ka, S.M., Chen, A., Chang, Z.F., and Ho, L.J. (2021b).
Mitochondrial CMPK2 mediates immunomodulatory and antiviral activities through IFN-dependent and
IFN-independent pathways. iScience *24*, 102498. 10.1016/j.isci.2021.102498

S2589-0042(21)00466-1 [pii].
Lawrence, T., and Natoli, G. (2011). Transcriptional regulation of macrophage polarization: enabling
diversity with identity. Nat Rev Immunol *11*, 750-761. 10.1038/nri3088

nri3088 [pii].

Liu, W., Chen, B., Chen, L., Yao, J., Liu, J., Kuang, M., Wang, F., Wang, Y., Elkady, G., Lu, Y., et al. (2019).
Identification of fish CMPK2 as an interferon stimulated gene against SVCV infection. Fish Shellfish
Immunol *92*, 125-132. S1050-4648(19)30588-1 [pii]

10.1016/j.fsi.2019.05.032.
Luo, Y., Zheng, D., Mou, T., Pu, J., Huang, Z., Chen, W., Zhang, Y., and Wu, Z. (2021). CMPK2 accelerates
liver ischemia/reperfusion injury via the NLRP3 signaling pathway. Exp Ther Med *22*, 1358.
10.3892/etm.2021.10793

ETM-0-0-10793 [pii].
Manzanillo, P.S., Shiloh, M.U., Portnoy, D.A., and Cox, J.S. (2012). Mycobacterium tuberculosis activates
the DNA-dependent cytosolic surveillance pathway within macrophages. Cell Host Microbe *11*, 469-480.
10.1016/j.chom.2012.03.007

S1931-3128(12)00125-4 [pii].
Martinez, F.O., Gordon, S., Locati, M., and Mantovani, A. (2006). Transcriptional profiling of the human
monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene
expression. J Immunol *177*, 7303-7311. 177/10/7303 [pii]

10.4049/jimmunol.177.10.7303.
McNab, F., Mayer-Barber, K., Sher, A., Wack, A., and O'Garra, A. (2015). Type I interferons in infectious
disease. Nat Rev Immunol *15*, 87-103. 10.1038/nri3787

nri3787 [pii].
Mehta, M.M., Weinberg, S.E., and Chandel, N.S. (2017). Mitochondrial control of immunity: beyond ATP.
Nat Rev Immunol *17*, 608-620. 10.1038/nri.2017.66

nri.2017.66 [pii].
Mills, E.L., Kelly, B., Logan, A., Costa, A.S.H., Varma, M., Bryant, C.E., Tourlomousis, P., Dabritz, J.H.M.,
Gottlieb, E., Latorre, I., et al. (2016). Succinate Dehydrogenase Supports Metabolic Repurposing of
Mitochondria to Drive Inflammatory Macrophages. Cell *167*, 457-470 e413. S0092-8674(16)31162-X [pii]

10.1016/j.cell.2016.08.064.
Missiroli, S., Genovese, I., Perrone, M., Vezzani, B., Vitto, V.A.M., and Giorgi, C. (2020). The Role of
Mitochondria in Inflammation: From Cancer to Neurodegenerative Disorders. J Clin Med *9*. E740 [pii]

10.3390/jcm9030740

jcm9030740 [pii].
Mosser, D.M., Hamidzadeh, K., and Goncalves, R. (2021). Macrophages and the maintenance of
homeostasis. Cell Mol Immunol *18*, 579-587. 10.1038/s41423-020-00541-3

10.1038/s41423-020-00541-3 [pii].
Newsholme, P., Curi, R., Gordon, S., and Newsholme, E.A. (1986). Metabolism of glucose, glutamine, long-
chain fatty acids and ketone bodies by murine macrophages. Biochem J *239*, 121-125. 10.1042/bj2390121.
Ó Maoldomhnaigh, C., Cox, D.J., Phelan, J.J., Mitermite, M., Murphy, D.M., Leisching, G., Thong, L.,
O'Leary, S.M., Gogan, K.M., and McQuaid, K. (2021). Lactate Alters Metabolism in Human Macrophages
and Improves Their Ability to Kill Mycobacterium tuberculosis. Frontiers in Immunology, 4104.
Oishi, Y., and Manabe, I. (2018). Macrophages in inflammation, repair and regeneration. Int Immunol *30*,
511-528. 10.1093/intimm/dxy054

5079207 [pii].

Olson, G.S., Murray, T.A., Jahn, A.N., Mai, D., Diercks, A.H., Gold, E.S., and Aderem, A. (2021). Type I
interferon decreases macrophage energy metabolism during mycobacterial infection. Cell Rep *35*,
109195. S2211-1247(21)00541-6 [pii]

10.1016/j.celrep.2021.109195.
Patrick, K.L., and Watson, R.O. (2021). Mitochondria: Powering the Innate Immune Response to
Mycobacterium tuberculosis Infection. Infect Immun *89*. e00687-20 [pii]

10.1128/IAI.00687-20

IAI.00687-20 [pii].
Picard, M., and McEwen, B.S. (2018). Psychological Stress and Mitochondria: A Conceptual Framework.
Psychosom Med *80*, 126-140. 10.1097/PSY.0000000000000544

00006842-201802000-00002 [pii].
Rambold, A.S., and Pearce, E.L. (2018). Mitochondrial Dynamics at the Interface of Immune Cell
Metabolism and Function. Trends Immunol *39*, 6-18. S1471-4906(17)30170-9 [pii]

10.1016/j.it.2017.08.006.
Ramond, E., Jamet, A., Coureuil, M., and Charbit, A. (2019). Pivotal Role of Mitochondria in Macrophage
Response to Bacterial Pathogens. Front Immunol *10*, 2461. 10.3389/fimmu.2019.02461.
Ravanelli, S., den Brave, F., and Hoppe, T. (2020). Mitochondrial Quality Control Governed by Ubiquitin.
Front Cell Dev Biol *8*, 270. 10.3389/fcell.2020.00270.
Rodriguez-Prados, J.C., Traves, P.G., Cuenca, J., Rico, D., Aragones, J., Martin-Sanz, P., Cascante, M., and
Bosca, L. (2010). Substrate fate in activated macrophages: a comparison between innate, classic, and
alternative activation. J Immunol *185*, 605-614. 10.4049/jimmunol.0901698

jimmunol.0901698 [pii].
Rodriguez, A.E., Ducker, G.S., Billingham, L.K., Martinez, C.A., Mainolfi, N., Suri, V., Friedman, A., Manfredi,
M.G., Weinberg, S.E., Rabinowitz, J.D., and Chandel, N.S. (2019). Serine Metabolism Supports Macrophage
IL-1beta Production. Cell Metab *29*, 1003-1011 e1004. S1550-4131(19)30014-2 [pii]

10.1016/j.cmet.2019.01.014.
Shankaran, D., Arumugam, P., Bothra, A., Gandotra, S., and Rao, V. (2019). Modern clinical
Mycobacterium tuberculosis strains leverage type I IFN pathway for a pro-inflammatory
response in the host. bioRxiv, 594655. 10.1101/594655.
Snell, L.M., McGaha, T.L., and Brooks, D.G. (2017). Type I Interferon in Chronic Virus Infection and Cancer.
Trends Immunol *38*, 542-557. S1471-4906(17)30093-5 [pii]

10.1016/j.it.2017.05.005.
Sprenger, H.G., MacVicar, T., Bahat, A., Fiedler, K.U., Hermans, S., Ehrentraut, D., Ried, K., Milenkovic, D.,
Bonekamp, N., Larsson, N.G., et al. (2021). Cellular pyrimidine imbalance triggers mitochondrial DNA-
dependent innate immunity. Nature metabolism *3*, 636-650. 10.1038/s42255-021-00385-9.
Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,
Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A
102, 15545-15550. 0506580102 [pii]

10.1073/pnas.0506580102.
Sugimoto, M.A., Sousa, L.P., Pinho, V., Perretti, M., and Teixeira, M.M. (2016). Resolution of Inflammation:
What Controls Its Onset? Front Immunol *7*, 160. 10.3389/fimmu.2016.00160.

Suski, J.M., Lebiedzinska, M., Bonora, M., Pinton, P., Duszynski, J., and Wieckowski, M.R. (2012). Relation
between mitochondrial membrane potential and ROS formation. Methods Mol Biol *810*, 183-205.
10.1007/978-1-61779-382-0_12.
Tan, Z., Xie, N., Cui, H., Moellering, D.R., Abraham, E., Thannickal, V.J., and Liu, G. (2015). Pyruvate
dehydrogenase kinase 1 participates in macrophage polarization via regulating glucose metabolism. J
Immunol *194*, 6082-6089. 10.4049/jimmunol.1402469

jimmunol.1402469 [pii].
Tiku, V., Tan, M.W., and Dikic, I. (2020). Mitochondrial Functions in Infection and Immunity. Trends Cell
Biol *30*, 263-275. S0962-8924(20)30018-0 [pii]

10.1016/j.tcb.2020.01.006.
Van den Bossche, J., Baardman, J., Otto, N.A., van der Velden, S., Neele, A.E., van den Berg, S.M., Luque-
Martin, R., Chen, H.J., Boshuizen, M.C., Ahmed, M., et al. (2016). Mitochondrial Dysfunction Prevents
Repolarization of Inflammatory Macrophages. Cell Rep *17*, 684-696. S2211-1247(16)31213-X [pii]

10.1016/j.celrep.2016.09.008.
Viola, A., Munari, F., Sanchez-Rodriguez, R., Scolaro, T., and Castegna, A. (2019). The Metabolic Signature
of Macrophage Responses. Front Immunol *10*, 1462. 10.3389/fimmu.2019.01462.
Vyssokikh, M.Y., Holtze, S., Averina, O.A., Lyamzaev, K.G., Panteleeva, A.A., Marey, M.V., Zinovkin, R.A.,
Severin, F.F., Skulachev, M.V., Fasel, N., et al. (2020). Mild depolarization of the inner mitochondrial
membrane is a crucial component of an anti-aging program. Proc Natl Acad Sci U S A *117*, 6491-6501.
10.1073/pnas.1916414117

1916414117 [pii].
Wang, F., Zhang, S., Vuckovic, I., Jeon, R., Lerman, A., Folmes, C.D., Dzeja, P.P., and Herrmann, J. (2018).
Glycolytic Stimulation Is Not a Requirement for M2 Macrophage Differentiation. Cell Metab *28*, 463-475
e464. S1550-4131(18)30513-8 [pii]

10.1016/j.cmet.2018.08.012.
Wang, F., Zhou, H., Deng, L., Wang, L., Chen, J., and Zhou, X. (2020). Serine Deficiency Exacerbates
Inflammation and Oxidative Stress via Microbiota-Gut-Brain Axis in D-Galactose-Induced Aging Mice.
Mediators of inflammation *2020*, 5821428. 10.1155/2020/5821428.
Wang, P., Zhou, Z., Hu, A., Ponte de Albuquerque, C., Zhou, Y., Hong, L., Sierecki, E., Ajiro, M., Kruhlak, M.,
Harris, C., et al. (2014). Both decreased and increased SRPK1 levels promote cancer by interfering with
PHLPP-mediated dephosphorylation of Akt. Mol Cell *54*, 378-391. 10.1016/j.molcel.2014.03.007

S1097-2765(14)00212-3 [pii].
Wang, Y., Li, N., Zhang, X., and Horng, T. (2021). Mitochondrial metabolism regulates macrophage biology.
J Biol Chem *297*, 100904. S0021-9258(21)00704-3 [pii]

10.1016/j.jbc.2021.100904.
Watanabe, S., Alexander, M., Misharin, A.V., and Budinger, G.R.S. (2019). The role of macrophages in the
resolution of inflammation. J Clin Invest *129*, 2619-2628. 10.1172/JCI124615

124615 [pii].
Weinberg, S.E., Sena, L.A., and Chandel, N.S. (2015). Mitochondria in the regulation of innate and adaptive
immunity. Immunity *42*, 406-417. 10.1016/j.immuni.2015.02.002

S1074-7613(15)00077-1 [pii].

Wilson, J.L., Nagele, T., Linke, M., Demel, F., Fritsch, S.D., Mayr, H.K., Cai, Z., Katholnig, K., Sun, X., Fragner,
L., et al. (2020). Inverse Data-Driven Modeling and Multiomics Analysis Reveals Phgdh as a Metabolic
Checkpoint of Macrophage Polarization and Proliferation. Cell reports *30*, 1542-1552 e1547.
10.1016/j.celrep.2020.01.011.
Wrobel, L., Topf, U., Bragoszewski, P., Wiese, S., Sztolsztener, M.E., Oeljeklaus, S., Varabyova, A., Lirski,
M., Chroscicki, P., Mroczek, S., et al. (2015). Mistargeted mitochondrial proteins activate a proteostatic
response in the cytosol. Nature *524*, 485-488. 10.1038/nature14951

nature14951 [pii].
Xu, Y., Johansson, M., and Karlsson, A. (2008). Human UMP-CMP kinase 2, a novel nucleoside
monophosphate kinase localized in mitochondria. J Biol Chem *283*, 1563-1571. S0021-9258(20)69650-8
[pii]

10.1074/jbc.M707997200.
Yan, J., and Horng, T. (2020). Lipid Metabolism in Regulation of Macrophage Functions. Trends Cell Biol
30, 979-989. S0962-8924(20)30175-6 [pii]

10.1016/j.tcb.2020.09.006.
Ye, J., Fan, J., Venneti, S., Wan, Y.W., Pawel, B.R., Zhang, J., Finley, L.W., Lu, C., Lindsten, T., Cross, J.R., et
al. (2014). Serine catabolism regulates mitochondrial redox control during hypoxia. Cancer Discov *4*, 1406-
1417. 10.1158/2159-8290.CD-14-0250

2159-8290.CD-14-0250 [pii].
Youle, R.J., and van der Bliek, A.M. (2012). Mitochondrial fission, fusion, and stress. Science *337*, 1062-
1065. 10.1126/science.1219855

337/6098/1062 [pii].
Yu, W., Wang, Z., Zhang, K., Chi, Z., Xu, T., Jiang, D., Chen, S., Li, W., Yang, X., Zhang, X., et al. (2019). One-
Carbon Metabolism Supports S-Adenosylmethionine and Histone Methylation to Drive Inflammatory
Macrophages. Molecular cell *75*, 1147-1160 e1145. 10.1016/j.molcel.2019.06.039.
Zhang, F., Qi, Y., Harrison, T.J., Luo, B., Zhou, Y., Li, X., Song, A., Huang, W., and Wang, Y. (2014). Hepatitis
E genotype 4 virus from feces of monkeys infected experimentally can be cultured in PLC/PRF/5 cells and
upregulate host interferon-inducible genes. J Med Virol *86*, 1736-1744. 10.1002/jmv.24014.
Zhong, Z., Liang, S., Sanchez-Lopez, E., He, F., Shalapour, S., Lin, X.J., Wong, J., Ding, S., Seki, E., Schnabl,
B., et al. (2018). New mitochondrial DNA synthesis enables NLRP3 inflammasome activation. Nature *560*,
198-203. 10.1038/s41586-018-0372-z

10.1038/s41586-018-0372-z [pii].