Muscle activation strategies of the vastus lateralis according to sex

Yuxiao Guo[1], Eleanor J. Jones[1], Thomas B. Inns[1], Isabel A.Ely[1], Daniel W. Stashuk[2], Daniel

J. Wilkinson[1], Kenneth Smith[1], Jessica Piasecki[3], Bethan E. Phillips[1], Philip J. Atherton[1],

Mathew Piasecki[1]

1. Centre of Metabolism, Ageing & Physiology (COMAP), MRC-Versus Arthritis Centre for

Musculoskeletal Ageing Research, National Institute for Health Research (NIHR)

Nottingham Biomedical Research Centre, University of Nottingham, Nottingham, UK

2. Department of Systems Design Engineering, University of Waterloo, Waterloo, ON,

Canada

3. Musculoskeletal Physiology Research Group, Sport, Health and Performance

Enhancement Research Centre, Nottingham Trent University, Nottingham, UK

Running title: Vastus lasteralis muscle activation in men and women

Corresponding author at

MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research and National Institute of

Health Research, NIHR Biomedical Research Centre, School of Medicine, Derby, DE22

3DT, UK.

E-mail address:

mathew.piasecki@nottingham.ac.uk (M. Piasecki)

Key points

- Increases in muscle force production are mediated by motor unit (MU) recruitment, and MU firing rate (FR).

- Women are underrepresented in studies of human neuromuscular research and markedly differ to men in a number of aspects of neuromuscular function, yet little is known of the recruitment strategies of each.

- Here we demonstrate men and women have similar vastus lateralis MU number estimates, yet women recruit smaller MUs with higher FR than men at normalised contraction levels. However, increases in force are achieved via similar trajectories of MU recruitment and MU FR in men and women.

- Although men and women exhibit divergent neuromuscular recruitment strategies to achieve normalised forces, increases in force are achived similarly and support the inclusion of mixed sex cohorts in studies of this nature.

Abstract

Aim: Despite men exhibiting greater muscle strength and fatigibility than women, it remains

unclear if there are sex-based differences in muscle recruitment strategies e.g. motor unit (MU)

recruitment and modulation of firing rate (FR) at normalised forces and during progressive

increases in force.

Methods: Twenty-nine healthy male and thirty-one healthy female participants (18-35 years)

were studied. Intramuscular electromyography was used to record individual motor unit

potentials (MUPs) and near fibre MUPs from the vastus lateralis (VL) during 10% and 25%

maximum isometric voluntary contractions (MVC), and spike-triggered averaging was used to

obtain motor unit number estimates (MUNE) of the VL. Multilevel mixed-effects linear

regression models were used to investigate the effects of sex at each contraction level.

Results: Men exhibited greater muscle strength ($p<0.001$) and size ($p<0.001$) than women,

with no difference in force steadiness at 10% or 25% MVC. Women had smaller MUs and

higher FR at 10% MVC (both $p<0.02$), similar to that at 25% MVC in MU size ($p=0.062$) and

FR ($p=0.031$). However, both sexes showed similar increases in MU size and FR when moving

from low- to mid-level contractions. There were no sex differences in any near fibre MUP

parameters or in MUNE.

Conclusion: In the vastus lateralis, women produce muscle force via different neuromuscular

recruitment strategies to men which is characterised by smaller MUs discharging at higher rates.

However, similar strategies are employed to increase force production from low to moderate

contractions. These findings of similar proportional increases between sexes support the use of

mixed sex cohorts in studies of this nature.

Keywords

Motor unit, neuromuscular junction, sex differences, vastus lateralis

Introduction

Skeletal muscle contraction is regulated by central and peripheral motor and sensory nerve function and excitation-contraction coupling of muscle fibres. The fundamental neuromuscular element regulating muscle contraction is the motor unit (MU), consisting of a motor neuron and the muscle fibres it innervates [1]. Increases in muscle force are largely mediated by two neuromuscular recruitment strategies, the recruitment of additional, progressively larger MUs, and an increase in MU firing rate (FR), referred to as MU recruitment and rate modulation, respectively [2]. Several studies have highlighted adaptive remodelling of MUs structure and function in response to exercise training, ageing and disease [3-6], which influences recruitment strategies, however the majority of data are only available in men.

Men generally possess greater muscle strength than women in upper and lower extremities, which is largely explained by greater muscle size [7]. Conversely, although task-specific, women are generally more resistant to neuromuscular fatigue when assessed at a normalised contraction level [8], which in the knee-extensors, is likely explained by differing fibre type ratios with a 7-23% greater proportion of type I fibres in vastus lateralis (VL) in women [9,10]. Sex differences of the hormonal milieu also influence neuromuscular function; testosterone and estrogen are the major sex hormones in males and females, respectively, and each exhibites a range of neuroprotective effects in motoneurons, such as dendritic maintenance and axonal sprouting [11]. Furthermore, hormonal metabolites are associated with the release of brain-derived neurotrophic factors (BDNF) [12], which are key mediators of synaptic plasiticity [13]. Acutely, differences in sex hormones partly explain the variability in fatigability in women across phases of the menstrual cycle [14]. Such differences in the hormonal milieu are difficult to experimentally control for and may explain why women are often underrepresented in studies of neuromuscular physiology [15].

Surface electromyography (EMG) has been commonly applied to study sex-based differences

of neuromuscular function and muscle recruitment strategies [16,17]. However, such approaches

are limited by the distance between activated MUs and recording electrodes [18], offering poor

MU yield in women, and in some cases, being influenced by adjacent muscles [19]. These

limitations can be overcome with the use of intramuscular EMG (iEMG), which also has the

added benefit of revealing further electrophysiological parameters relevant to MU size and

complexity [20]. Although we have previously reported the sex-based divergent trajectory of MU

FR from middle to older age in long-term trained master athletes [21], comparisons of normative

values in healthy young men and women at differing contraction levels are unknown. The aims

of the present study were to compare individual MU properties and muscle recruitment

strategies, as well as the MU number estimates (MUNEs) in the VL of healthy young men and

women. We hypothesised several parameters would differ at normalised contraction levels,

with no sex-based differences in recruitment strategies when moving from a low- to a mid-

level contraction.

Materials and Methods

Ethics approval

This research was approved by the University of Nottingham Faculty of Medicine and Health

Sciences Research Ethics Committee (C16122016, 160-0121, 186-1812, 103-1809, 302-1903)

and was conducted between 2019 and 2021 in accordance with the Declaration of Helsinki.

Participants

Twenty-nine healthy male and thirty-one healthy female participants, aged 18-35 years, were

recruited via advertisement posters in the local community. All the participants volunteered to

take part in the studies and provided written informed consent. Prior to enrolment, all

participants completed a comprehensive clinical examination and metabolic screening were

conducted at the School of Medicine, Royal Derby Hospital Centre. All articipants were

recreationally active. Participants with metabolic disease, lower limb musculoskeletal

abnormalities, acute cerebrovascular or cardiovascular disease, active malignancy,

uncontrolled hypertension, or those on medications that impact muscle protein metabolism or

modulate vascular tone were excluded.

Anthropometry

Body mass and height were measured using calibrated scales and a stadiometery, respectively

for the calculation of body mass index (BMI). Ultrasound was used to measure the cross-

sectional area of the VL using an ultrasound probe ((LA523 probe, B-mode, frequency range

26–32 Hz, and MyLabTM50 scanner, Esaote, Genoa, Italy) at the anatomical mid-point of the

muscle which was identified between the greater trochanter and the mid-point of the patella

with participants lying supine. Ultrasound images were acquired aligning the superior edge of

the probe following a middle-to-lateral direction position on the skin, beginning, and ending

the image capture at aponeurosis borders.   A water-based conductive gel was applied on the

surface of the ultrasound probe to enhance the fidelity of the image without causing excessive

contact pressure on the skin during the acquisition of the images. Images were subsequently

analysed using ImageJ software (National Institutes of Health, USA) to quantify CSA

measurements. The mean area of three images was taken as CSA. The CSA of eight female

participants was measured using magnetic resonance imaging (MRI) with a T1-weighted turbo

3D sequence on a 0.25-T G-Scan (Esaote, Genoa, Italy). Continuous transversal images with a

6-mm slice were acquired and analysed by using Osirix imaging software (Osirix medical

imaging, Osirix, Atlanta, GA, United States) through tracing around the VL following the

contour of the aponeurosis. VL CSA values are available for 23 men and 19 women.

Muscle strength and force steadiness

The maximum isometric voluntary contraction force (MVC) of the right knee extensor was assessed with the participants sitting in a custom-built chair with hips and knees flexed at ~90 degrees. The lower leg was securely attached to a force dynamometer with non-compliant straps (purpose-built calibrated strain gauge, RS125 Components Ltd, Corby, UK) slightly above the medial malleolus. A seat belt was fastened across the pelvis to avoid superfluous movement of the upper trunk during the test. To obtain the external knee joint moment arm, the distance from centre of the force strap to the lateral femoral condyle was measured. After a standardised warm-up of submaximal contractions, participants were instructed to perform a maximal isometric contraction with real-time visual feedback and verbal encouragement. This was repeated a further two times, with 60 second rest intervals between each, and the highest value was accepted as MVC. Peak torque during the selected MVC was also determined.

After determination of the MVC, participants were instructed to perform between four and six sustained isometric contractions at 10% and 25% MVC, respectively, each lasting 12-15 seconds with a target line displayed on the screen in front of the participants (with iEMG, below, Figure 1). Participants had 20-30 seconds rest between each contraction. Prior to the assessment, participants were allowed a single familiarisation practice at each contraction level. Force steadiness was quantified as the coefficient of variation of the force [CoV; (SD/mean) $\times$ 100]. The mean CoV at each contraction level was calculated from the middle two contractions.

Surface electromyography (sEMG)

An active recording sEMG electrode (disposable self- adhering Ag-AgCl electrodes; 95 mm^2, Ambu Neuroline, Baltorpbakken, Ballerup, Denmark) was placed over the motor point located around the mid-point of the VL, identified using a cathode probe (Medserve, Daventry, UK)

to apply percutaneous electrical stimulation at 400 V, pulse width of 50 µs and current of

around 8 mA (DS7A Digitimer, Welwyn Garden City, Hertfordshire, UK) with a self-adhesive

anode electrode (Dermatrode, Farmadomo, NL) placed over the right gluteus. A reference

electrode was placed over the patella tendon and a common ground electrode placed over the

patella. The common ground electrode served for both sEMG and iEMG measurements. sEMG

signals were sampled at 10kHz, and bandpass filtered between 5 and 5 kHz (1902 amplifier,

Cambridge Electronics Design Ltd., Cambridge, UK) and digitized with a CED Micro 1401

data acquisition unit (Cambridge Electronic Design) for offline analysis.

Compound muscle action potential (CMAP)

The CMAP of the VL was evoked by a manually triggered stimulator (model DS7A; Digitimer)

using percutaneous stimulation (Medserve, Daventry, UK) of the proximal femoral nerve

(approximately halfway between the anterior superior iliac spine and the pubic tubercle) with

a carbon-rubber anode electrode (Dermatrode self-adhering electrode, 5.08 cm in diameter;

Farmadomo Linde Homecare Benelux Bv, Leiden, The Netherlands) placed over the skin

overlying the gluteus muscle. The stimulator voltage was fixed at 400 V and the pulse width at

50 µS, with the current increased incrementally until the M-wave amplitude plateaued. At this

point, the current was increased again by ~30 mA to ensure supramaximal stimulation, ensuring

a sharp rise time of the negative peak of the m-wave.

Intramuscular electromyography (iEMG)

A 25-mm disposable concentric needle electrode (N53153; Teca, Hawthorne, New York, USA)

was inserted at the muscle belly of VL, adjacent to the recording surface electrode over the

motor point, to a depth of 1.5-2 cm depending on the muscle size. The iEMG shared the same

ground electrode as the sEMG, which was placed over the patella. iEMG signals were recorded

using Spike2 (Version 9.06), sampled at 50 kHz and bandpass filtered at 10 Hz to 10 kHz (1902

amplifier; Cambridge Electronic Design Ltd, Cambridge, UK) and stored for future off-line

analysis.

Prior to EMG and CMAP assessments, participants performed a series of voluntary, low-level

contractions once the needle was positioned to ensure adequate signal to noise ratio, thus

ensuring the recording needle electrode was close to depolarizing fibres. Each participant then

performed the sustained voluntary isometric contractions as detailed above (Figure 1). After a

10% and 25% MVC contraction, to avoid repeat sampling of the same MUs, the needle

electrode was repositioned by the combinations of twisting the bevel edge 180 degrees and

withdrawing by ~5 mm. This process was repeated until four to six contractions from spatially

distinct areas (from deep to superficial portions) were recorded [18]. Participants had ~30 seconds

rest between each contraction.

EMG analysis

Decomposition-based quantitative electromyography (DQEMG) software was used to detect

motor unit potentials (MUPs), extract motor unit potential trains (MUPTs) generated by

individual MUs from the iEMG signals and estimate, via ensemble averaging, their

corresponding surface MUPs (sMUPs) from the sEMG signals [22]. MUPTs that were composed

of MUPs from more than one MU or had fewer than 40 MUPs were excluded. The occurrence

times of individual MUPs within a MUPT were used to trigger and align sEMG signal epochs

for ensemble-averaging to produce an estimate of their corresponding sMUPs. The sMUPs

estimated from signals recorded during 25% MVC contractions were use to represent MU size.

All MUP and sMUP templates were visually inspected and their markers adjusted, where

required, to correspond to the onset, onset of negative phase (sMUP only), end, and positive

and negative peaks of the waveforms.

MUP amplitude was measured from the maximal positive and negative peaks and the MUP

area was taken as the total area within the MUP duration (onset to end) and is indicative of MU

size. The number of phases and turns are measures of MUP complexity and are classified as

the number of components above or below the baseline (phases) and a change in waveform

direction of at least 25 µV (turns), which indicates the level of temporal dispersion across

individual muscle fibre contributions to a single MUP. MU FR was assessed as the rate of MUP

occurrences within a MUPT, expressed as the number of occurrences per second (Hz).

A near fibre MUP (NFM) is defined as the acceleration of its corresponding MUP (Figure 1)

and calculated by applying a second-order, low-pass differentiator to the MUP which

effectively reduces the recording area of the needle electrode to within ~350 µm, thereby

ensuring only potentials from fibres closest to the needle electrode significantly contribute to

the NFM and reducing interference from distant active fibres of other MUs. NFM jiggle is a

measure of the shape variability of consecutive NFMs of an MUPT expressed as a percentage

of the total NFM area. NFM segment jitter is a measure of the temporal variability of individual

fibre contributions to the NFMs of a MUPT. It is calculated as a weighted average of the

absolute values of the temporal offsets between matched NFM segments of consecutive

isolated (i.e., not contaminated by the activity of other MUs) NFMs across an MUPT expressed

in microseconds. NFM dispersion is the time, in ms, between the first and last MU fibre

contributions [20].

Figure 1. Representative data from a male (A) and a female (B) participant. Top panels show knee extensor force
traces at 10% and 25% of MVC, and corresponding intramuscular electromyography (iEMG) raw data recorded
from the vastus lateralis. A representative MUP template and corresponding NFM shimmer plot isolated from
each contraction are shown below each iEMG signal. Vertical orange lines on MUPs and NFMs indicate the start
and end time of the NFM. Abbreviations: N, newtons; mV, millivolt; MVC, maximum voluntary contraction;
MUP; motor unit potential; Dur, duration; NF, near fibre; NFM, near fibre MUP; kV, kilovolt; μV, microvolt; ms,
millisecond.

Motor unit number estimates (MUNE)

The MUNE value was derived by dividing the negative peak area of the ensemble averaged

mean surface MUP (msMUP) from 25% MVC into the negative peak area of the CMAP [23]. A

msMUP is an ensemble average of the negative-peak onset aligned, sMUPs of the MUs

sampled from a muscle. The negative peak area of the msMUP was divided into the negative

peak area of the electrically evoked CMAP [24]. MUNE values are available for 15 men and 15

women.

Statistical analysis

All of the statistical analysis was performed using RStudio (Version 1.3.959 for macOS) [25].

Descriptive statistics of participant characteristics are presented as *mean ± standard deviation*

(SD). Student's unpaired t-test was used to compare physical parameters (age, BMI, MVC,

CSA, and force steadiness). As multiple MUs were recorded from each participant, *multi-level*

mixed-effect linear regression analysis was performed to investigate these MU parameters with

sex and contraction level as factors through the package *lme4* (Version 1.1.23) [26]. In the linear

mixed models, the first level was single motor unit; single motor units were clustered according

to each participant to form the second level, which was defined as the participant level. This

linear mixed-effect modelling framework is suitable for data of this nature as it: i) incorporates

the whole sample of extracted MUs not just the mean values obtained from each participant,

which preserves variability within and across participants simultaneously to the greatest extent;

ii) handles missing data better than an *analysis of variances (ANOVA)* framework as the

removal of a single missing observation has a much smaller effect in the mixed model; and iii)

provides coefficient estimates that indicate the magnitude and direction of the effects of interest

[27]. Interactions were first examined and where not present they were removed from the model,

sex and contraction level were explored individually. The results are displayed as coefficient

estimates, 95% confidence intervals, and *p*-values. Standardized estimates were calculated

through the package *effectsize* (Version 0.4.5) [28] for forest plotting. For data visualisation,

individual participant means and group means were shown in box-and-jitter plots. Statistical

significance was assumed when $p < 0.05$. Based on the models used, p values close to 0.05

were also addressed [29].

Results

The means and standard deviations for the participant's characteristics are shown in Table 1.

Significant differences between men and women were detected for weight, height, BMI, peak

torque and VL CSA (all *p*<0.05). There were no significant sex differences for age and force

steadiness (both *p*>0.05). Individual values are shown in Figure 2.

Table 1. Participant characteristics.

Measure	Men (n=29)	Women (n=31)	P value
Age (years)	23.7(5.0)	22.9(3.6)	0.490
Weight (kg)	81.2(11.6)	63.1(10.1)	**<0.001**
Height (cm)	180.3(7.5)	165.8(6.8)	**<0.001**
BMI (kg/m^2)	25.0(2.9)	23.0(3.2)	**0.012**
Peak Torque (Nm)	251.65(67.72)	158.95(46.69)	**<0.001**
VL CSA (cm^2)	28.35(6.43)	19.00(3.90)	**<0.001**
Force CoV-10%	4.51(1.21)	4.98(1.18)	0.139
Force CoV-25%	3.36(1.17)	3.24(0.74)	0.638

Data are reported as mean (standard deviation). Values in bold reflect statistically significant (*p*<0.05) results.
Abbreviations: BMI, body mass index; VL CSA, vastus lateralis cross-sectional area.

Figure 2. Box-and-jitter plots of the individual participant means, and group means (red dot) of **(A)** peak torque,
(B) vastus lateralis cross-sectional area, **(C)** motor unit number estimates (MUNE), and **(D)** force steadiness at
10% and 25% maximum voluntary contraction (MVC), in men (yellow) and women (blue). Abbreviations: CoV,
coefficient of variation.

Figure 3. Box-and-jitter plots of the individual participant means, and group means (red dot) of **(A)**motor unit
(MU) firing rate and **(B)** firing rate variability in men (yellow) and women (blue) at 10 and 25% maximum
voluntary contraction (MVC).

Figure 4. Box-and-jitter plots of the individual participant means, and group means (red dot) of motor unit potential (MUP) properties in men (yellow) and women (blue) at 10 and 25% maximum voluntary contraction (MVC).

Figure 5. Box-and-jitter plots of the individual participant means, and group means (red dot) of near fibre motor unit potential (NFM) properties in men (yellow) and women (blue) at 10 and 25% maximum voluntary contraction (MVC).

A total of 1645 MUs were analysed in men and 1207 in women. At 10% MVC, the mean number of MUs isolated per person was 26 in men and 17 in women; at 25% MVC, the mean number of MUs isolated per person was 31 in men and 22 in women. Individual mean values

for all functional, MU and NFM parameters are sown in Figures 2-5. There were no significant

interactions between sex and contraction level in any of the MU parameters. When interactions

were removed from the model, multilevel linear regression revealed women had greater MU

FR at both 10% (mean; M: 8.08 Hz; W: 8.79 Hz) and 25% (M: 8.62 Hz; W: 9.20 Hz) MVC

(both $p<0.05$, Table 2, Figure 3A). This was matched by a non-significant trend ($p<0.10$) for

greater MU FR vaiability in women at both contraction levels (Table 2, Figure 3B). MUP

duration was shorter at 10% (M: 8.37 ms; W: 6.61 ms) and 25% (M: 8.24 ms; W: 6.84 ms)

MVC in women when compared to men (both $p<0.01$, Table 2, Figure 4E). MUP area was

smaller in women at 10% (M: 741μV·ms; W: 531 μV ·ms) ($p=0.006$), with a non-significant

trend at 25% MVC (M: 1005 μV ·ms; W:775μV ·ms) ($p=0.062$). There were no significant

sex-based differences in any other MU characteristic (Table 2).

Table 2. Motor unit properties in different sexes

Parameter	Level	Beta	95%CI	P value
MU FR	10% MVC	0.73	0.14 to 1.32	**0.018**
	25% MVC	0.61	0.07 to 1.14	**0.031**
MU FR Variability	10% MVC	0.01	-0.001 to 0.02	0.080
	25% MVC	0.01	-0.001 to 0.03	0.066
MUP Area	10% MVC	-210.08	-352.90 to -67.27	**0.006**
	25% MVC	-215.41	-436.82 to 6.00	0.062
MUP Phases	10% MVC	-0.07	-0.46 to 0.31	0.707
	25% MVC	0.05	-0.38 to 0.48	0.823
MUP Amplitude	10% MVC	-62.83	-151.27 to 25.61	0.170
	25% MVC	-92.01	-206.31 to 22.29	0.120
MUP Turns	10% MVC	-0.19	-0.73 to 0.35	0.500
	25% MVC	-0.10	-0.60 to 0.40	0.691
MUP Duration	10% MVC	-1.85	-2.93 to -0.77	**0.001**
	25% MVC	-1.35	-2.33 to 0.37	**0.009**
NFM Jiggle	10% MVC	-0.06	-1.84 to 1.72	0.947
	25% MVC	0.09	-1.85 to 2.03	0.928
NFM Duration	10% MVC	-0.16	-0.60 to 0.29	0.486
	25% MVC	-0.02	-0.40 to 0.36	0.926
NFM Seg Jitter	10% MVC	-0.22	-4.29 to 3.84	0.915
	25% MVC	0.47	-3.42 to 4.36	0.814
NFM Dispersion	10% MVC	0.08	-0.22 to 0.38	0.622
	25% MVC	0.03	-0.25 to 0.32	0.825
NFM Area	10% MVC	0.001	-0.71 to 0.71	0.998

	25% MVC	-0.22	-0.97 to 0.54	0.575

Beta value and 95% confidence interval (CI) represents the model predicted change per unit from men to women, shown separately for 10 and 25% maximum voluntary contraction (MVC). All statistical analysis was based on multilevel mixed effect linear regression models with each subject as an independent cluster. The values in bold reflect statistically significant ($p<0.05$) results. Abbreviations: MU, motor unit; MUP, motor unit potential; FR, firing rate; NFM, near fibre motor unit potential; Seg, segment.

With increasing contraction level, both men and women exhibited higher MU FR and MU FR variability, as well as greater MUP amplitude, and larger MUP area (all $p<0.001$, Table 3, Figures 3 and 4). NFM segment jitter, NFM duration and NFM area were also larger with the higher contraction level, increasing to a similar extent in men and women (all $p<0.001$, Table 3, Figure 6). There were no interactions between sex and contraction level in any of the MU parameters, indicating the difference from 10 to 25% MVC did not differ between men and women (Figure 6).

Table 3. Motor unit properties at different contraction levels

Parameter	Sex	Beta	95%CI	P value
MU FR	Men	0.50	0.28 to 0.71	**<0.001**
	Women	0.43	0.18 to 0.69	**0.001**
MU FR Variability	Men	0.02	0.01 to 0.02	**<0.001**
	Women	0.02	0.01 to 0.02	**<0.001**
MUP Area	Men	273.62	215.81 to 331.43	**<0.001**
	Women	199.35	148.13 to 250.56	**<0.001**
MUP Phases	Men	-0.05	-0.14 to 0.04	0.302
	Women	0.10	-0.02 to 0.21	0.098
MUP Amplitude	Men	188.21	153.75 to 222.67	**<0.001**
	Women	142.40	106.27 to 178.53	**<0.001**
MUP Turns	Men	0.05	-0.11 to 0.20	0.565
	Women	0.09	-0.08 to 0.26	0.316
MUP Duration	Men	-0.16	-0.52 to 0.20	0.381
	Women	0.13	-0.16 to 0.42	0.371
NFM Jiggle	Men	3.08	2.38 to 3.79	**<0.001**
	Women	3.08	2.34 to 3.82	**<0.001**
NFM Duration	Men	-0.40	-0.54 to -0.26	**<0.001**
	Women	-0.34	-0.50 to -0.18	**<0.001**
NFM Seg Jitter	Men	0.77	-0.37 to 1.91	0.188
	Women	0.51	-0.85 to 1.87	0.463
NFM Dispersion	Men	-0.09	-0.23 to 0.05	0.228
	Women	-0.05	-0.28 to 0.18	0.655
NFM Area	Men	0.98	0.68 to 1.29	**<0.001**
	Women	0.72	0.39 to 1.05	**<0.001**

Beta value and 95% confidence interval (CI) represents the model predicted change per unit from 10 to 25% maximum voluntary contraction (MVC), shown separately for men and women. All statistical analysis was based

on multilevel mixed effect linear regression models with each subject as an independent cluster. The values in
bold reflect statistically significant ($p<0.05$) results. Abbreviations: MU, motor unit; MUP, motor unit potential;
FR, firing rate; NFM, near fibre motor unit potential; Seg, segment.

Figure 6. Forest plots of the standardised regression coefficient estimate for associations between motor unit
characteristics and contraction levels in men and women models. Beta value and 95% confidence intervals (CI)
represents the standardised model predicted change per unit from 10 to 25% maximum voluntary contraction
(MVC). All statistical analysis was based on multilevel mixed effect linear regression models with each subject
as an independent cluster, largely maintaining the motor unit variability within each subject. Standardised values
of each parameter make the comparisons executable between men and women. *** =$p<0.001$.

Discussion

This is the first study to compare muscle recruitment strategies and motor unit number

estimates of the VL using iEMG techniques in healthy young men and women. Despite men

having stronger and larger muscles, there were no differences in force steadiness at either lower

or moderate contraction levels between sexes. At each contraction level assessed, women

displayed smaller markers of MU size and greater MU FR, indicating differing recruitment

strategies to achieve a normalised force. When assessing the difference between contraction

levels, both men and women exhibited higher MU FR and greater MUP size, which differed to

a similar extent in both sexes, indicating a similar recruitment strategy to generate proportional

increases in force. In addition, there was no significant sex-based differences in motor unit

number estimates of the VL. These data reveal divergent neuromuscular recruitment strategies

between sexes to achieve a normlised force, which follow similar trajectories with increasing

force.

Consistent with previous studies, women exhibited 33% smaller muscle size (CSA of the VL),

which was reflected in a 31% lower strength [7,30,31]. The greater MU FR of women shown here

in VL is in an agreement with some but not all previously published data and again highlights

probable muscle specific confounders. For instance, women exhibited higher MU FR and MU

FR variability compared with men in elbow flexors, flexor digitorum indicis, biceps, knee

extensors and tibialis anterior [32-35]. However, others reported no sex difference in knee

extensors during 30% MVC [36] and significantly greater MU FR at 100% MVC in tibialis

anterior in men [37,38]. These differences in intrinsic motoneuron excitability may be explained

by sex-specific levels of persistent inward currents, which amplify and prolong synaptic input

to MUs and affected by the levels of central monoanmines [39], which are reported to be higher

in women [40]. As expected, in the current study MU FR increased with increasing force levels

to a similar extent in both men and women, accompanied by greater MU FR variability.

Recruitment threshold dominates at lower force levels, whereas MU FR is more significant at

higher forces [41,42], which may explain the increased variability with increased force. Although

differeng at each contraction level, the similar proportional increase in MU FR in men and

women indicates both sexes follow similar discharge pattern increases from low to mid-level

contractions.

Despite large differences in muscle strength, force steadiness - representing the ability to hold

a constant force, which is also influenced by MU FR and its variability [32,43-45], did not differ

between sexes at either contraction level. Differing from the current findings, Inglis [35] found

that women had a greater MU FR variability and greater fluctuation in steadiness than men

during dorsiflexion in tibiais anterior muscles, which may indicate a muscle specific sex

difference. In the current study, both men and women exhibited greater force steadiness at 25%

MVC compared 10% MVC, consistent with Inglis's finding that very high and low force

outputs have greater fluctuations compared to moderate force outputs [35,46].

The size of a MU can be estimated by the size of the MUP recorded using intramuscular

electrodes. As previously mentioned, men typically exhibit larger muscle size than women, [9,47],

with increases in force mediated by recruitment of additional larger MUs and increases in MU

FR. Here MUP area and duration were smaller in women which reflects smaller MU size.

When viewed alongside the greater MU FR in women, it suggests that at the normalised force

levels assessed here, women are more reliant on MU FR than on recruitment of larger MUs,

when compared to men. As expected, markers of MU size increased at larger contraction levels,

as larger MUs are recruited to produce larger forces. Again, the trajectory of each was similar

for men and women indicating MU recruitment strategies moving between these force levels

do not differ between sexes.

A NFM is derived from a MUP, such that is primarily composed of contributions from MU

fibres close to the intramuscular electrode [20]. Here there were no sex differences in any NFM

parameters at either contraction level. When comparing 10% MVC and 25% MVC contractions,

NFM area increased, while NFM duration decreased, albeit to a similar extent in both sexes.

These contraction induced alterations may be the result of the activation of larger MU fibres

with greater conduction velocity during higher level contractions.

Increases in NFM instability, as measured by NFM jiggle or NFM segment jitter can reflect

increases in neuromuscular junction (NMJ) transmission instability with age [21,48-50] and in

diabetic neuropathy [3]. In the current study NFM instability, as measured by NFM jiggle,

increased with contraction level for both sexes, and to similar extents. NFM jiggle is based on

variability in the amplitudes of NFM shapes, and although these amplitude changes are

normalized by the size of the NFM, it is possible that these increases with contraction level

may be due to the recruitment of larger MUs with more MU fibres contributing to larger NFMs

at 25% MVC. Combined with the lack of a sex difference in NFM segment jitter, it is clear

NMJ transmission instability in the VL is sensitive to contraction level and is similar in heathy

young men and women. However, there were no statistically significant contraction-based

differences in NFM segment jitter, which is based on variability in the occurrence times of

NFM segments and is not affected by NFM size, indicating it is less sensitive to the influences

of contraction level.

The mean values of MUNE in men (240 ± 66) and women (218 ± 68) reported here are similar

to those we have previously reported in male cohorts [5,50], and highlights the repeatability of

this method in this muscle group when applying identical experimental procedures. Although

the MUNE should be viewed as an index relative to the number of MUs within a muscle and

not a true anatomical count, the similar values reported here in men and women support

minimal sex-based differences. Combined with the small difference in MU size at 25% MVC,

which is lower than total muscle size difference, the current data support the notion that sex-

based differences in total muscle size are largely explained by greater individual fibre size in

men [47].

Although providing a high level of detail of MU structure and function via MUPs and NFMs

sampled in deep and superficial muscle regions, regardless of subcutaneous tissue amount,

iEMG is sensitive to contraction level and reliably identifying individual MU activity at high

levels in this muscle can be problematic. Therefore, data presented here were obtained during

low and mid-level contractions only. Secondly, we did not control for hormonal fluctuations

in women naturally occurring during the menstrual cycle, and although there is limited

evidence to confirm this, these fluctuations may influence several parameters included here.

This is a direct comparison of MU features in young men and women, and further

investigations concerning neural drive and influence of hormones on neural drive are still

required to further understand the sex-based differences in the motor nervous system.

In summary, when compared to men, women exhibited smaller VL MUs which exhibited

higher MU FR, when assessed at a single normalised contraction level. However, both men

and women showed similar increases in MU size and MU FR from a low- to a mid-level

contraction, indicating a similar neuromuscular recruitment strategies. These results suggest

that although sex-based neuromuscular differences are apparent at a single contraction level,

relative differences between levels are similar in this widely studied muscle group. These data

do not support the notion of excluding women from studies of this nature.

Acknowledgements
We thank all of the participants for their enthusiastic involvement in this study. We are grateful
to Mr Daniel McCormick for assistance with data collection.

Funding
This work was supported by the Medical Research Council [grant number MR/P021220/1] as
part of the MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research awarded to the
Universities of Nottingham and Birmingham, and was supported by the NIHR Nottingham
Biomedical Research Centre. The views expressed are those of the author(s) and not
necessarily those of the NHS, the NIHR or the Department of Health and Social Care. This
work was also supported by a Physiological Society Research Grant awarded to MP.

Author contributions

All authors contributed to the conception and design of the work. Y.G., E.J.J., T.B.I, I.A.E.,

J.P., and M.P. contributed to the acquisition and analysis of the data. YG analysed the data and

drafted the manuscript. B.E.P, P.J.A., D.J.W, K.S., D.W.S, and M.P. provided comments. All

authors have approved the final version of the submitted manuscript for publication and are

accountable for all aspects of the work. All persons designated as authors qualify for authorship,

and all those who qualify for authorship are listed.

Conflict of interest

The authors have no conflict of interest to declare.

Data availability statement

The datasets generated and analysed during the current study are available from the

corresponding author upon reasonable request.

Reference

1.   Heckman CJ, Enoka RM. Motor Unit. In: *Comprehensive Physiology.*2012:2629-2682.

2.   Enoka RM, Duchateau J. Rate Coding and the Control of Muscle Force. *Cold Spring*
Harb Perspect Med. 2017;7(10).

3.   Allen MD, Stashuk DW, Kimpinski K, Doherty TJ, Hourigan ML, Rice CL. Increased
neuromuscular transmission instability and motor unit remodelling with diabetic
neuropathy as assessed using novel near fibre motor unit potential parameters. *Clin*
Neurophysiol. 2015;126(4):794-802.

4.   Piasecki M, Ireland A, Jones DA, McPhee JS. Age-dependent motor unit remodelling in
human limb muscles. *Biogerontology.* 2016;17(3):485-496.

5.   Piasecki M, Ireland A, Piasecki J, et al. Long-Term Endurance and Power Training May
Facilitate Motor Unit Size Expansion to Compensate for Declining Motor Unit Numbers
in Older Age. *Frontiers in Physiology.* 2019;10(449).

6.   Del Vecchio A, Casolo A, Negro F, et al. The increase in muscle force after 4 weeks of
strength training is mediated by adaptations in motor unit recruitment and rate
coding. *The Journal of Physiology.* 2019;597(7):1873-1887.

7.   Hannah R, Minshull C, Buckthorpe MW, Folland JP. Explosive neuromuscular
performance of males versus females. *Experimental Physiology.* 2012;97(5):618-629.

8.   Hunter SK. Sex differences in human fatigability: mechanisms and insight to
physiological responses. *Acta Physiol (Oxf).* 2014;210(4):768-789.

9.   Haizlip KM, Harrison BC, Leinwand LA. Sex-based differences in skeletal muscle
kinetics and fiber-type composition. *Physiology (Bethesda).* 2015;30(1):30-39.

10.   Ansdell P, Thomas K, Hicks KM, Hunter SK, Howatson G, Goodall S. Physiological sex
differences affect the integrative response to exercise: acute and chronic implications.
Experimental Physiology. 2020;105(12):2007-2021.

11. Hyer MM, Phillips LL, Neigh GN. Sex Differences in Synaptic Plasticity: Hormones and
Beyond. *Front Mol Neurosci.* 2018;11:266.
12. Georgieva SDaK. Sex Hormones in Neurodegenerative Processes and Diseases. In:
Cellular and Molecular Mechanisms of the Effects of Sex Hormones on the Nervous
System. InTech; 2018.
13. Mantilla CB, Stowe JM, Sieck DC, et al. TrkB kinase activity maintains synaptic function
and structural integrity at adult neuromuscular junctions. *Journal of Applied*
Physiology. 2014;117(8):910-920.
14. Ansdell P, Brownstein CG, Škarabot J, et al. Menstrual cycle-associated modulations in
neuromuscular function and fatigability of the knee extensors in eumenorrheic
women. *Journal of Applied Physiology.* 2019;126(6):1701-1712.
15. Cowley ES, Olenick AA, McNulty KL, Ross EZ. "Invisible Sportswomen": The Sex Data
Gap in Sport and Exercise Science Research. *Women in Sport and Physical Activity*
Journal. 2021:1-6.
16. Clark BC, Collier SR, Manini TM, Ploutz-Snyder LL. Sex differences in muscle fatigability
and activation patterns of the human quadriceps femoris. *European Journal of Applied*
Physiology. 2005;94(1):196-206.
17. Bolgla L, Cook N, Hogarth K, Scott J, West C. Trunk and hip electromyographic activity
during single leg squat exercises do sex differences exist? *Int J Sports Phys Ther.*
2014;9(6):756-764.
18. Jones EJ, Piasecki J, Ireland A, et al. Lifelong exercise is associated with more
homogeneous motor unit potential features across deep and superficial areas of
vastus lateralis. *Geroscience.* 2021.
19. Christie A, Greig Inglis J, Kamen G, Gabriel DA. Relationships between surface EMG
variables and motor unit firing rates. *European Journal of Applied Physiology.*
2009;107(2):177-185.
20. Piasecki M, Garnés-Camarena O, Stashuk DW. Near-fiber electromyography. *Clinical*
Neurophysiology. 2021;132(5):1089-1104.
21. Piasecki J, Inns TB, Bass JJ, et al. Influence of sex on the age-related adaptations of
neuromuscular function and motor unit properties in elite masters athletes. *The*
Journal of Physiology. 2021;599(1):193-205.
22. Stashuk DW. Decomposition and quantitative analysis of clinical electromyographic
signals. *Med Eng Phys.* 1999;21(6-7):389-404.
23. Brown WF, Strong MJ, Snow R. Methods for estimating numbers of motor units in
biceps-brachialis muscles and losses of motor units with aging. *Muscle Nerve.*
1988;11(5):423-432.
24. Piasecki M, Ireland A, Piasecki J, Stashuk DW, McPhee JS, Jones DA. The reliability of
methods to estimate the number and size of human motor units and their use with
large limb muscles. *European Journal of Applied Physiology.* 2018;118(4):767-775.
25. *RStudio: Integrated Development for R* [computer program]. RStudio, PBC, Boston,
MA2020.
26. Bates D MM, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software. 2015;67(1):1-48.
27. Brown VA. An Introduction to Linear Mixed-Effects Modeling in R. *Advances in*
Methods and Practices in Psychological Science. 2021;4(1):2515245920960351.
28. Ben-Shachar MS LD, Makowski D. effectsize: Estimation of Effect Size Indices and
Standardized Parameters. *Journal of Open Source Software.* 2020;5(56):2815.

29. Greenland S, Senn SJ, Rothman KJ, et al. Statistical tests, P values, confidence intervals,
and power: a guide to misinterpretations. *Eur J Epidemiol.* 2016;31(4):337-350.

30. Miller AEJ, MacDougall JD, Tarnopolsky MA, Sale DG. Gender differences in strength
and muscle fiber characteristics. *European Journal of Applied Physiology and*
Occupational Physiology. 1993;66(3):254-262.

31. Jeon Y, Choi J, Kim HJ, Lee H, Lim JY, Choi SJ. Sex- and fiber-type-related contractile
properties in human single muscle fiber. *J Exerc Rehabil.* 2019;15(4):537-545.

32. Taylor AM, Christou EA, Enoka RM. Multiple Features of Motor-Unit Activity Influence
Force Fluctuations During Isometric Contractions. *Journal of Neurophysiology.*
2003;90(2):1350-1361.

33. Harwood B, Cornett KMD, Edwards DL, Brown RE, Jakobi JM. The effect of tendon
vibration on motor unit activity, intermuscular coherence and force steadiness in the
elbow flexors of males and females. *Acta Physiologica.* 2014;211(4):597-608.

34. Peng Y-L, Tenan MS, Griffin L. Hip position and sex differences in motor unit firing
patterns of the vastus medialis and vastus medialis oblique in healthy individuals.
Journal of Applied Physiology. 2018;124(6):1438-1446.

35. Inglis JG, Gabriel DA. Sex differences in the modulation of the motor unit discharge
rate leads to reduced force steadiness. *Appl Physiol Nutr Metab.* 2021;46(9):1065-
1072.

36. Tenan MS, Peng Y-L, Hackney AC, Griffin L. Menstrual Cycle Mediates Vastus Medialis
and Vastus Medialis Oblique Muscle Activity. *Medicine & Science in Sports & Exercise.*
2013;45(11).

37. Christie A, Kamen G. Short-term training adaptations in maximal motor unit firing
rates and afterhyperpolarization duration. *Muscle Nerve.* 2010;41(5):651-660.

38. Inglis JG, Gabriel D. Sex Differences in Motor Unit Discharge Rates at Maximal and
Submaximal Levels of Force Output. *Applied Physiology, Nutrition, and Metabolism.*
2020;45.

39. Heckman CJ, Johnson M, Mottram C, Schuster J. Persistent inward currents in spinal
motoneurons and their influence on human motoneuron firing patterns.
Neuroscientist. 2008;14(3):264-275.

40. Nishizawa S, Benkelfat C, Young SN, et al. Differences between males and females in
rates of serotonin synthesis in human brain. *Proceedings of the National Academy of*
Sciences. 1997;94(10):5308.

41. De Luca CJ, LeFever RS, McCue MP, Xenakis AP. Control scheme governing
concurrently active human motor units during voluntary contractions. *J Physiol.*
1982;329:129-142.

42. Contessa P, De Luca CJ. Neural control of muscle force: indications from a simulation
model. *J Neurophysiol.* 2013;109(6):1548-1570.

43. Yao W, Fuglevand RJ, Enoka RM. Motor-Unit Synchronization Increases EMG
Amplitude and Decreases Force Steadiness of Simulated Contractions. *Journal of*
Neurophysiology. 2000;83(1):441-452.

44. Enoka RM, Christou EA, Hunter SK, et al. Mechanisms that contribute to differences in
motor performance between young and old adults. *Journal of Electromyography and*
Kinesiology. 2003;13(1):1-12.

45. Farina D, Negro F, Dideriksen JL. The effective neural drive to muscles is the common
synaptic input to motor neurons. *The Journal of Physiology.* 2014;592(16):3427-3441.

46. Yoon T, Vanden Noven ML, Nielson KA, Hunter SK. Brain areas associated with force
steadiness and intensity during isometric ankle dorsiflexion in men and women.
Experimental Brain Research. 2014;232(10):3133-3145.
47. Staron RS, Hagerman FC, Hikida RS, et al. Fiber type composition of the vastus lateralis
muscle of young men and women. *J Histochem Cytochem.* 2000;48(5):623-629.
48. Hourigan ML, McKinnon NB, Johnson M, Rice CL, Stashuk DW, Doherty TJ. Increased
motor unit potential shape variability across consecutive motor unit discharges in the
tibialis anterior and vastus medialis muscles of healthy older subjects. *Clin
Neurophysiol.* 2015;126(12):2381-2389.
49. Piasecki M, Ireland A, Coulson J, et al. Motor unit number estimates and
neuromuscular transmission in the tibialis anterior of master athletes: evidence that
athletic older people are not spared from age-related motor unit remodeling.
Physiological Reports. 2016;4(19):e12987.
50. Piasecki M, Ireland A, Stashuk D, Hamilton-Wright A, Jones DA, McPhee JS. Age-related
neuromuscular changes affecting human vastus lateralis. *J Physiol.*
2016;594(16):4525-4536.