Increased RNA Polymerase Activity and Pausing at CTCF binding sites on the Epstein Barr Virus Genome During Reactivation from Latency

Laura E. M. Dunn[1,2], Fang Lu[3], Paul M. Lieberman[3], Joel D. Baines[1,2*]

[1] Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, New York, USA

[2] Department of Pathobiology, School of Veterinary Medicine, Louisiana State University, Baton Rouge, Louisiana, USA

[3] The Wistar Institute, Philadelphia, Pennsylvania, USA

*Corresponding author

Email: jdb11@cornell.edu

Abstract

The ability of Epstein-Barr Virus (EBV) to switch between latent and lytic infection is key to its long-term persistence, yet the molecular mechanisms behind this switch remain unclear. To investigate transcriptional events during the latent to lytic switch we utilized Precision nuclear Run On followed by deep Sequencing (PRO-Seq) to map cellular RNA polymerase (Pol) activity to single-nucleotide resolution on the host and EBV genome in latently infected Mutu-I cells at 1, 4 and 12 h post-reactivation. During latency, Pol activity was primarily limited to the EBNA1 transcript initiating at the Qp promoter, the EBER and RPMS1/BART regions and the BHLF1 transcript. Unexpectedly at early time-points post-reactivation, the EBV transcripts with the largest increase in Pol activity were LMP-2A, EBER-1 and RPMS1. Closer analysis of the PRO-Seq data at these regions revealed a distinct pattern of high Pol activity with bidirectional transcription and strong peaks indicative of Pol pausing. Alignment to ChIP-Seq data revealed a strong correlation with CTCF binding sites on the EBV genome. In addition, alignment to ATAC-Seq data indicated that many of these transcription regulatory regions were sites of accessible chromatin. Similar features were observed in Akata cells activated from latency with anti-IgG. Overall, these data suggest that during reactivation, EBV recruits RNA polymerase to CTCF binding sites where it transcribes short distances and pauses. These activities likely help open chromatin on the viral genome to initiate productive replication.

Author summary

The ability of EBV to switch between latent and lytic infection is key to its long-term persistence in memory B-cells and its ability to persist in proliferating cells is strongly linked to oncogenesis. During latency, most viral genes are epigenetically silenced,

and the virus must overcome this repression to reactivate lytic replication.

Reactivation occurs once the immediate early (IE) EBV lytic genes are expressed.

However, the molecular mechanisms behind the switch from the latent transcriptional

program to begin transcription of the IE genes remains unknown. In this study, we

were able to precisely map RNA polymerase positioning and activity during latency

and reactivation. Unexpectedly, RNA polymerase activity was not enriched at the IE

genes during early reactivation but accumulated at distinct regions, characteristic of

enhancers, on the EBV genome previously shown to be associated with CTCF and

open chromatin. We speculate that Pol accumulation at these sites helps maintain

open chromatin locally, eventually leading to changes in the quaternary structure of

the EBV genome to promote lytic replication.

Introduction

Epstein-Barr virus (EBV) is a gammaherpesvirus that establishes a persistent

infection in 95% of adults worldwide. EBV maintains this persistence through a

biphasic lifecycle in which it switches between latent and lytic phases. The virus

establishes latency in memory B-cells where it exists as a circular episome and the

majority of genes are epigenetically silenced via repressive chromatin and DNA

methylation (1).

There are four distinct latency types (0-III), which are classified based on the

expression of the EBNA proteins, LMP proteins and non-coding RNAs (2). EBNA

transcript expression is regulated by promoter selection from the Qp, Cp and Wp

latency promoters (2). This promoter selection is mediated by the chromatin

architecture of the EBV episome, which is dependent on host CCCTC-binding factor

(CTCF) (3). CTCF is a multifunctional DNA binding protein with known roles in

transcriptional activation/repression, chromatin loop and boundary formation and

promoter-enhancer blocking activity (4). The EBV genome contains multiple CTCF

binding sites (5) and a number of these sites have been linked to the maintenance of

the latent chromatin state (6-8).

Reactivation is known to be induced by stimulation of the B cell receptor (BCR) and

various stress-related signals (9). These events lead to activation of the viral Zp and

Rp promoters to drive transcription of the immediate early (IE) EBV genes, BZLF1

and BRLF1, which in turn induce early lytic gene expression (10). A number of host-

cell factors including BLIMP1 (11), PARP1 (8) and MYC/MAX (12) have been shown

to be involved in regulating the induction of IE expression. However, the exact

molecular mechanisms as to how EBV overcomes the repressive latent state to

promote transcription of the IE genes remains unknown.

EBV relies on cellular RNA polymerase II (Pol II) for transcription, although it also

uses RNA Polymerase III (Pol III) to transcribe Epstein-Barr virus-encoded small

RNA 1 and 2 (EBER1, 2) (13). Pol II contains an extra C-terminal domain (CTD) and

the phosphorylation pattern of the CTD is a key factor in the regulation of

transcription initiation, elongation and termination (14). EBV is known to modulate

Pol II transcription elongation by promoting serine 2 CTD phosphorylation to drive B-

cell immortalisation. The virus achieves this by inducing Pol II pausing at the Cp

promoter via association with the negative elongation factor (NELF), which facilitates

recruitment of the positive transcription elongation factor (pTEFb). pTEFb maintains

CTD serine 2 phosphorylation, leading to strong expression of EBV immortalisation

genes (15).

Pausing of Pol II at promoter regions, known as promoter proximal pausing (PPP), is

a mechanism known to be used by herpesviruses to regulate transcription. The

alphaherpesvirus, Herpes simplex virus-1 (HSV-1) utilizes PPP to regulate temporal

gene expression during lytic replication (16). Most notably, Kaposi's sarcoma

associated herpesvirus (KSHV), another gammaherpesvirus, exploits PPP to

regulate the latent-lytic switch. During latency, NELF is recruited to pause Pol II on

lytic KSHV gene promoters allowing for prompt expression upon stimulation of

reactivation (17).

The aim of this study was to investigate changes in Pol II activity and positioning

during the transition from a highly restricted latent state to the transcriptionally active

lytic state. We hypothesized that Pol II remains paused on the promoters of the IE

genes during latency and that upon reactivation, Pol is rapidly released into

elongation. To address this hypothesis we used Precision nuclear Run On followed

by deep Sequencing (PRO-Seq) (18) to map cellular Pol activity to single-nucleotide

resolution on the EBV genome in both latently infected Mutu-I cells and Akata cells

and at 1, 4 and 12 h post-reactivation.

Using PRO-Seq, we found no evidence of PPP on IE genes during EBV latency or

reactivation. Instead, Pol activity and pausing upon reactivation was found to be

enriched at distinct regions that were often not associated with any known viral

transcript. These regions correlated with CTCF binding sites on the EBV genome,

and the pattern of Pol activity suggested that these CTCF binding sites induced Pol

pausing. Alignment to ATAC-Seq data indicated that this stalling of Pol was linked to

sites of open chromatin. Overall, this study suggests that one of the first steps during

EBV reactivation from latency involves increased bidirectional Pol activity and

pausing with an associated opening of chromatin near CTCF binding sites on the

viral genome.

Results

Distribution of active polymerase on the EBV episome in latent Mutu-I cells.

In order to assess reactivation, the extent of RNA Pol activity during latency needed

to be first established. Nuclei were harvested from the EBV positive Burkitt's

lymphoma cell-line, Mutu-I (type 1 latency), and PRO-Seq was performed as

previously described (16, 19). Drosophila nuclei were spiked into the run-on reaction

and corresponding sequencing reads were used to normalize each sample for

sequencing depth. EBV sequences were aligned to the EBV Mutu genome build (20)

lacking all but one W repeat to allow appropriate read assignment.

To get an overview of Pol activity on the entire EBV episome during latency, the data

was first visualised using the Integrative Genomics Viewer (IGV) (21) (Fig. 1A).To

highlight peak regions the IGV windowing function was to set to maximum. Overall,

there was a relatively low level of RNA Pol activity across the EBV episome, with the

most substantial peak visible at the Pol III-transcribed EBERs and the BHLF1

transcript. To identify the transcripts with the most active Pol, SeqMonk software

(22) was used to align and quantify PRO-Seq reads to the viral genome. Most

transcripts aligned with between 10 - 1000 normalised reads (Fig. 1B), indicating a

low background level of Pol activity on the viral genome during latency.

The only transcripts with over 1000 reads were BHLF1, RPMS1, both EBERs and

multiple EBNA transcripts. The EBERs had the highest level of Pol activity with a

mean of 29,061 normalised reads for EBER-1 (Log10 4.46) and 24,062 for EBER-2

(Log10 4.38); this level was over 4-fold higher than any other EBV transcript. As the

EBNA genes have overlapping mRNAs we were unable to assign the exact EBNA

transcript to reads from this region. However, we were able to quantify reads from

the Cp, Qp and Wp latency promoters (Fig. 1C). This analysis revealed that only the

Qp promoter, which is known to drive EBNA1 expression, had substantial Pol activity

during latency. In summary, PRO-Seq analyses indicated that the highest level of

Pol activity in latent Mutu-I cells was on the EBERs and the EBNA transcripts (driven

from the Qp promoter). This result is highly consistent with type 1 latency and thus

confirmed the validity of this technique for studying EBV transcription.

Distribution and quantification of active polymerase on the EBV episome

during reactivation in Mutu-I cells.

We next investigated the transcriptional profile of EBV reactivation using PRO-Seq.

Lytic replication in latently infected Mutu-I cells was initiated by treatment with

NaB/TPA. Nuclei were harvested at 1, 4 and 12 h post-reactivation, followed by

PRO-Seq analysis. Representative whole genome IGV views for all time points are

shown in Fig. 2A. To highlight peak regions the IGV windowing function was to set to

maximum. Using IGV visualization of the data at the whole genome level, there were

only small differences visible overall. However, at 1h post-reactivation, noticeable

peaks emerged at LMP-2A, RPMS1-OriLyt and antisense to the EBERs and these

remained at 4 and 12h post reactivation.

To determine whether the changes in Pol activity were significant, the reads for each

EBV transcript in latent and reactivated samples were quantified using DeSeq2 Fold

change. At 1h post-reactivation, the strongest and most significant increase in Pol

activity mapped to LMP-2A, while EBER-1 also had a significant increase (Fig. 2B).

Interestingly, most EBV genes had a reduction in Pol activity during this first hour of

reactivation. At 4h post-reactivation, Pol activity on LMP-2A remained significantly

up-regulated and activity also increased on other LMP genes, LMP-1 and LMP-2B

(Fig. 2C). Activity on the RPMS1 transcript also increased significantly at this time

point. By 12h post-reactivation, Pol activity was increased on most genes but still

remained highest on LMP-2A and RPMS1 (Fig. 2D). The transcriptional activators

BZLF1 and BRLF1 are also highlighted in Fig. 2D but significant increases of Pol

activity on these genes was only observed starting at 12h post-reactivation.

180 PRO-Seq identifies distinct patterns of Pol activity on the EBV episome during

181 reactivation.

Because the regions with the strongest increases in Pol activity during reactivation

(LMP-2A, RPMS1, EBER) were unexpected, we examined these regions in more

detail using IGV. This revealed that the increase in Pol activity on LMP-2A at 1 h

post-reactivation was not at the promoter region, but instead was primarily limited to

the first exon, located approximately 80 bp downstream of the promoter (Fig. 3A). At

4 and 12 h post-reactivation another peak appeared 150 bp downstream of the initial

peak but on the opposite strand, and in a region that does not map to any known

transcript. An additional broad Pol peak appeared on the LMP-2A intronic region at 4

and 12 h post-reactivation, approximately 500 bp downstream of the exon 1 peak.

EBER-1 was the only other transcript to have a significant upregulation of Pol activity

at 1h post-reactivation. Closer analysis of this region revealed that most of the Pol

activity increase was upstream of EBER-1 on both strands, in an area of no known

transcripts (Fig. 3B). The RPMS1 region also had an interesting pattern of Pol

activity, with Pol activity appearing during reactivation on both strands of the

genome, and mapping to introns and regions with no known transcripts (Fig. 3C).

Overall, PRO-Seq analysis has indicated that upregulation of Pol activity during EBV

reactivation occurs not at gene promoters (as might be expected if promoter

proximal pausing was a prominent feature), but more commonly it occurs

bidirectionally at intronic or intergenic regions.

dREG detection of transcription regulatory elements from PRO-Seq

corresponds with CTCF binding sites on the EBV episome.

The regions with the highest levels of Pol activity during reactivation were

unexpected and so we sought to analyze the data in an alternative way. We chose to

use the software package dREG (Detection of regulatory elements using GRO-seq

and other run-on and sequencing assays), which is trained to detect transcription

regulatory elements including transcription initiation regions (TIRs) (23), and applied

this to our PRO-Seq data. Novel TIRs discovered by this package have been shown

to be associated with transcription factor binding sites. We therefore utilised the EBV

portal database (5) to align EBV transcription factor binding ChIP-Seq data to the

EBV dREG peaks generated from our PRO-Seq data (Fig. 4A). There were certain

regions on the EBV episome where multiple transcription factor binding sites aligned

to the significant dREG pol peaks (highlighted in green, false discovery rate (FDR) $\leq$

0.05). For example, CTCF, c-FOS, c-JUN, MAX and RAD21 all had peaks aligning to

the EBER dREG peak. CTCF, Sp1, EGR1 and GCN5 also had peaks mapping to

dREG peaks at both origins of lytic replication (OriLyt). However, the most striking

observation was that of the 8 significant dREG peaks (highlighted in green), a CTCF

peak also aligned. Notably, the regions found to be of relevance from DeSeq/PRO-

Seq analysis, LMP-2A, EBER and RPMS1, all had aligned dREG/CTCF peaks. With

the exception of the Cp peak, all of the significant dREG peaks were also present

during latency despite low PRO-Seq reads at most of these regions.

To assess if this was specific to Mutu-I cells and the NaB/TPA method of reactivation

we repeated the experiment using another EBV positive Burkitt's lymphoma cell-line,

Akata, reactivated with anti-IgG. Reactivation was followed by extraction of nuclei at

1, 4 and 12 hours, and PRO-Seq/dREG analysis was performed as above. It should

be noted that EBV copy number varies between the cell types, resulting in the Akata

PRO-Seq data having lower read counts (PRO-Seq IGV browser images and

volcano plots in Fig. S1). dREG analysis improved the ability to compare peak

regions between the cell types. This data showed an association of 7 CTCF peaks

with significant dREG pol peaks during latency and reactivation (Fig. 4B) and 6 of

these were in common with Mutu-I cells. These were located at; EBERs, OriLyt (left),

Qp promoter, RPMS1, OriLyt (right) and LMP-2A (exon 1). All significant dREG peak

locations are listed in Table S1. A noticeable difference from Mutu-I was that two

significant dREG peaks emerged during reactivation in the middle of the genome

near BZLF1 and BKRF3 (overlapped with a CTCF site). In addition, no peaks were

apparent close to the Cp and Wp promoters in Akata, as seen in Mutu-I. In

conclusion, dREG analysis has detected a number of significant transcriptional

regulatory elements on the EBV genome during latency and reactivation. Though

there was a slight variation between the total and location of peaks between Mutu-I

and Akata cells, there was a consistent association of dREG peaks with CTCF

binding sites in both cell types.

Alignment of CTCF ChIP-Seq and PRO-Seq/dREG indicates pausing of Pol at

CTCF sites on EBV genome.

It was noticeable that most dREG/CTCF peak alignments in Fig. 4 were slightly

shifted from each other. To further investigate this observation, we used IGV to look

at regions of interest at higher resolution using the Mutu-I data. Peaks of Pol

accumulation were visible on both strands of the genome next to or over the CTCF

peak. Peaks in PRO-Seq are associated with pausing of the polymerase, which

leads to an abundance in transcripts adjacent to the pause site (18). This is shown in

detail at the LMP-2A region in Fig. 5A, with clear Pol peaks emerged on either side

of the CTCF peak during reactivation. A similar pattern was also seen at the RPMS1

CTCF peak, with strong bidirectional Pol activity accumulation around this CTCF

peak during reactivation (Fig. 5B). Further examples of bidirectional Pol activity

accumulation adjacent to CTCF sites during reactivation are shown for the EBER

region (Fig. 5C), at the Qp promoter (Fig. 5D) and the Cp promoter (Fig. 5E). The

shifted dREG/CTCF peak pattern was also apparent in Akata cells during

reactivation, with these sites also showing strong Pol activity during reactivation (Fig.

S2 note: due to the lower EBV genome copy number in Akata, the PRO-Seq signal

is weaker and often not visible until 12 h post-reactivation). Overall, these data

suggest that Pol initiates, accumulates, and pauses bidirectionally at regions

adjacent to CTCF binding sites during EBV reactivation.

Pol accumulates at sites of accessible chromatin on the EBV episome during

early reactivation.

The extension of Pol activity near CTCF binding sites at only 1 h post-reactivation

suggests a localized increase in accessible chromatin on the EBV episome. To

assess this, we used data from a previous Assay for Transposase-accessible

Chromatin Sequencing (ATAC-Seq) experiment (GEO: GSE172476) in 2 sub-

populations of latent Mutu-I cells, MUN14 and M14, and aligned this to the Mutu-I

dREG and CTCF ChIP-Seq data in IGV (Fig. 6). Five ATAC-Seq peaks aligned

strongly with CTCF peaks and with dREG Pol peaks at regions already identified to

be of importance, specifically; EBER, Qp, RPMS1, LMP-2A and both OriLyts. This

alignment indicates that Pol activity localizes in specific regions of open chromatin

during reactivation.

CTCF binding alterations on the EBV episome during reactivation

Finally, we investigated what happens to CTCF binding on the EBV episome during

early reactivation. ChIP was performed on latent (untreated) and 4h reactivated

(NaB/TPA treated) Mutu-I cells and also latent (untreated) and 4h reactivated (anti-

IgG treated) Akata cells using antibodies against CTCF, Pol II and RAD21. qPCR

was used to assess binding at multiple CTCF sites on both the EBV and human

genome. The Pol II ChIP confirmed an increase in Pol II binding on the EBV genome

during reactivation in both Mutu-I and Akata cells, though the reactivation treatment

in both cell types also led to an increase in Pol II at sites on the human genome (Fig.

7A, 7B). At 4h post-reactivation in Mutu-I, there was a decrease in CTCF binding at

all 3 sites examined on the EBV genome (LMP-CTCF, Cp-CTCF, Qp-CTCF) (Fig.

7C). In contrast, at 4h post-reactivation in Akata cells, CTCF binding was increased

at the LMP-CTCF and Qp-CTCF binding sites and was unchanged at the Cp-CTCF

site (Fig. 7D). As CTCF can function alongside the cohesin complex, binding of the

cohesin component RAD21 was also assessed. RAD21 binding at the LMP-CTCF

locus has already been shown (Fig. 4A) and the ChIP results revealed an increase in

RAD21 binding at this site in Mutu-I cells at 4h post-reactivation (Fig. 7E). RAD21

binding at this site was unchanged at 4h post-reactivation in Akata cells (Fig. 7D).

Overall, this ChIP study indicates that CTCF binding on the EBV episome is altered

during reactivation and could be linked to the cohesin complex. However, the

changes are dependent on the cell-line/method of reactivation and the subsequent

kinetics of reactivation.

Discussion

In this study we used PRO-Seq to investigate how Pol activity changes on the EBV

episome during the switch from the highly restricted latent transcriptional state to

productive lytic replication. Despite the well documented activity of BZLF1 as the

viral lytic transcriptional activator (24), the molecular mechanisms that initiate

transcription of BZLF1 remain unknown. Here we describe important insights as to

how Pol activity responds to reactivation signals to aid the transcriptional switch.

We initially speculated that EBV regulates the latent to lytic switch at the elongation

step of transcription via release from PPP of Pol II. However, the PRO-Seq data from

EBV latency (Fig. 1A) showed no evidence of PPP on any EBV genes normally

expressed during productive infection. Instead, the data documented a highly

restricted transcriptional state during latency. Pol activity (above background) was

limited to the EBERs, EBNA1 (from Qp promoter) and RPMS1 transcripts (Fig. 1B

and 1C), consistent with transcription of these genes in type I latency (25).

At 1 h post-reactivation, peaks of Pol activity did emerge, although none were

located at gene promoters. Quantification of the PRO-Seq reads using Deseq2 fold

change analysis (26) indicated that the region in Mutu-I cells with the largest

increase in Pol activity was the first exon of the latent gene, LMP-2A. In Akata cells,

activity at LMP-2A was also strongly increased (Akata volcano plot data shown in

Fig. S1) during the first hour of reactivation but due to low EBV read counts in these

cells, it was difficult to visualize the precise region of activity increase. EBER-1 and

the RPMS1 transcript also had strong increases in Pol activity during early

reactivation in Mutu-I cells, and similar to LMP-2A Pol activity at these sites was

bidirectional. Bidirectional transcriptional initiation has been shown to be associated

with active enhancers on the human genome (27). This feature and recognition by

the dREG tool (23) suggest that the increased Pol activity occurs at previously

unrecognized transcriptional enhancer regions. The ATAC-Seq data (Fig. 6) also

supports the enhancer hypothesis as accessible chromatin is linked to *cis*-regulatory

elements, predominantly enhancers and promoters (28).

EBV latency is known to be regulated by enhancer-promoter interactions. Multiple

EBV-encoded transcription factors (EBNA-LP, EBNA2, EBNA3C) are associated

with *B-cell super-* enhancer regions important for EBV-induced lymphoproliferation

(29). On the EBV genome, *oriP* acts as an enhancer for efficient transcription from

the Cp and Wp promoter (30) and *oriLyt* contains an enhancer region required for

viral DNA replication (31) that mediates late gene transcription (32). Recently, it has

been shown that MYC binds to the *oriLyt* enhancer during latency and MYC

depletion leads to reactivation (12). Our data supports the conclusion that enhancer

elements at EBV *oriLyt* regions have important regulatory functions in EBV

reactivation.

A curious finding was that the IE transcriptional activators, BZLF1 and BRLF1, did

not have any significant increase in Pol activity until 12h post-reactivation in Mutu-I

cells. This is despite RT-qPCR validation of lytic cycle induction confirming

expression of BZLF1 (Fig S3). This could be explained by the enhancer hypothesis

as it is possible the PRO-Seq peaks are due to transcription at these potential

enhancer regions, leading to the generation of enhancer RNAs (eRNAs) (33). As

PRO-Seq measures Pol activity and not stable transcripts, eRNAs, should be

detected readily by PRO-Seq. IE gene transcription could then be stimulated through

long-range interactions between these enhancer regions and the IE gene promoters

(34). Notably, CTCF has been linked to involvement in the activity of enhancer-

promoter function (35, 36).

We discovered a strong correlation between Pol activity and CTCF binding sites on

the EBV genome. CTCF is known to be important for maintenance of the chromatin

state during EBV latency (6, 37) and interestingly, mutation of only the LMP-CTCF

locus disrupted the epigenetic state of the latent viral episome (7). We also found

that CTCF binding changed at 3 EBV sites upon stimulation of reactivation in Mutu-I

(Fig. 7C). This contrasts with a previous study that indicated CTCF binding is

unchanged during EBV reactivation (8). A possible explanation for this difference is

that we looked at an earlier time point, 4h, compared to 24h in the above study.

Indeed, ChIP-qPCR at 24h post-reactivation in Mutu-I cells showed a recovery in

CTCF binding at both the LMP and Cp CTCF loci (Fig. S4). In addition, the cell/virus

background and method of reactivation leads to variation in the kinetics of

reactivation (38). This could therefore also explain differences between the two

studies and the differences we found here between Mutu-I and Akata cells. Our

PRO-Seq data indicate that Mutu-I and Akata cells have different reactivation

kinetics as Pol activity on IE genes and other lytic genes increased earlier in Akata

cells (Fig. S1).

CTCF is involved in maintaining the distinct chromatin loops found in different EBV

latency stages (3) and so may also be involved in reactivation by alteration of

chromatin conformation. It has already been shown that EBV reactivation involves

the recruitment of cellular chromatin remodeling enzymes, mediated by BZLF1 (39).

The authors of this study noted an association between BZLF1, CTCF and open-

chromatin (ATAC-Seq), supporting a link between CTCF and chromatin modification

during EBV reactivation. The effects we observe on CTCF-cohesin DNA binding may

reflect changes in EBV chromatin conformation that are likely to contribute to the

regulation of EBV reactivation.

The pattern of alignment between CTCF and PRO-Seq peaks was indicative of Pol

pausing adjacent to, and at the CTCF sites on the EBV genome. It has recently been

shown that CTCF binding sites lead to Pol II stalling on the human genome but that

stalling was independent of CTCF binding (40). Therefore, the association of Pol and

CTCF binding sites on the EBV genome during reactivation may not be due to CTCF

binding itself. Our ChIP-Seq data did suggest a role for CTCF at certain binding

sites, but more studies are needed to confirm whether CTCF loss is functionally

important for reactivation. Paused Pol is associated with maintaining DNA

accessibility (41) and stalling of Pol has previously been implicated in the prevention

of nucleosome assembly on latent EBV promoters (15). This previous work adds

support to the hypothesis that Pol accumulation at CTCF binding sites is involved in

altering accessibility of the EBV genome during reactivation.

In summary, the PRO-Seq data presented here show that Pol activity during early

EBV reactivation occurs at distinct sites on the EBV episome that show similar

characteristics to enhancer regions. A strong correlation was identified between

CTCF binding sites and Pol activity, and it appeared that these CTCF sites lead to

Pol stalling. Our current hypothesis is that Pol activity is initiated at specific enhancer

regions during reactivation due to the accessibility of the genome at these sites. Pol

accumulates and stalls to aid maintenance of an open chromatin conformation,

allowing for increased promoter accessibility eventually leading to productive

replication.

Materials and methods

Cells

EBV positive Burkitt's lymphoma cell line Mutu I and Akata were maintained in RPMI

1640 containing 12% FBS and antibiotics (penicillin and streptomycin). Mutu I was

obtained from Dr. Jeffrey Sample, Penn State University Hershey Medical School,

PA. Akata was obtained from Dr. Elena Mattia, University of Rome SAPIENZA, Italy.

Drosophila melanogaster S2 (ATCC) cells were grown in Schneider's medium

(Lonza) containing 10% foetal bovine serum (FBS) and maintained at 23°C

EBV reactivation

418 For PRO-seq, Mutu I cells were treated with NaB (1 mM) and TPA (20 ng/ml) for 0,

419 1, 4, or 12 hrs, and Akata cells were treated with 10 μg/ml of goat anti-human IgG

420 (Sigma I1886) for 0, 1, 4, or 12 hrs. For ChIP assays, Mutu I cells were untreated or

421 treated with NaB (1 mM) and TPA (20 ng/ml) for 4 or 24 hrs, and Akata cells were

422 untreated or treated with 10 μg/ml of goat anti-human IgG for 4 hrs.

Nuclei isolation

Nuclei were isolated following method previously described (42). In brief, cells were

washed 2x with ice-cold PBS and then incubated for 10 min on ice with swelling

buffer (10mM Tris-HCl [pH 7.5], 10% glycerol, 3mM $CaCl_2$, 2mM, $MgCl2_2$, 0.5mM

DTT, protease inhibitors [Roche] and 4 U/ml RNase inhibitors [RNaseOUT,

ThermoFisher]). Cells were scraped from plate, pelleted via centrifugation at 600 x g

for 10 min (4°C), resuspended in lysis buffer (swelling buffer + 0.5% Igepal) and then

incubated on ice for 20 min to release nuclei. Nuclei were pelleted by centrifugation

at 1500 x g for 5 min (4°C), washed 2x with lysis buffer with a final wash in storage

buffer (50mM Tris-HCl [pH 8.0], 25% glycerol, 5mM MgAcetate, 0.1mM EDTA, 5mM

DTT). Nuclei were resuspended in storage buffer, flash frozen in LN_2 and stored at -

80°C.

Precision nuclear run-on assay

The nuclear run-on assays were performed as described in (18, 42, 43). Frozen

nuclei were thawed on ice and S2 nuclei spiked-in to Mutu-I/Akata cells at a ratio of

1:1000. An equal volume of run-on buffer (10mM Tris-HCl [pH 8.0], 5mM $MgCl_2$,

1mM DTT, 300 mM KCl, 20μM biotin-11-ATP, -UTP, -GTP, -CTP, 0.4 U/ml RNase

inhibitor [RNaseOUT ThermoFisher] and 1% Sarkosyl) was added to thawed nuclei.

Run-on was performed under constant shaking for 3 min on a vortex shaker at 37°C.

Reaction was ended by the addition of TRIzol LS (ThermoFisher) and vortexed to

homogenise.

Library preparation

RNA was extracted from run-on nuclei via TRIZOL extraction. RNA underwent base

hydrolysis by the addition of 0.2N NaOH and incubation on ice for 20 min.

Unincorporated nucleotides were removed using a P-30 column (Bio-Rad).

Biotinylated RNA was purified using streptavidin M280 Dynabeads (ThermoFisher)

with a series of washes. Bead-bound RNA was washed 2x in ice-cold high salt wash

(50mM Tris-HCl [pH 7.4], 2M NaCl, 0.5% Triton X-100, 0.4 U/ml RNase inhibitor

[RNaseOUT ThermoFisher]), 2x in ice-cold medium salt wash (10mM Tris-HCl [pH

7.4], 300mM NaCl, 0.1% Triton X-100, 0.4 U/ml RNase inhibitor [RNaseOUT

ThermoFisher]) and once in ice-cold low salt wash (5mM Tris-HCl [pH 7.4], 0.1%

Triton X-100, 0.4 U/ml RNase inhibitor [RNaseOUT ThermoFisher]). RNA was eluted

from beads with two TRIzol extractions.

The 3'-RNA adapter with a 5' phosphate (5'-Phos) and 3' inverted dT (InvdT) (5'-

Phos-GAUCGUCGGACUGUAGAACUCUGAAC-3'-InvdT) was ligated to the 3' end

of the RNA using T4 RNA ligase I (NEB). RNA was purified as above by binding to

streptavidin beads, washing and TRIzol extracted followed by removal of the 5' cap

with 10 U of 5'-pyrophosphohydrolase (RppH) (NEB) and 5' end repair with T4 PNK

(NEB). The 5'-RNA adapter (5'-CCUUGGCACCCGAGAAUUCCA-3') was ligated to

the 5' end of the RNA using T4 RNA ligase I (NEB). RNA was purified as above by

binding to streptavidin beads, washing and TRIzol extracted.

RNA was reverse transcribed with SuperScript III reverse transcriptase

(ThermoFisher) using the RNA PCR primer 1 (5'-

AATGATACGGCGACCACCGAGATCTACACGTTCAGAGTTCTACAGTCCGA-3').

The cDNA was PCR amplified with Phusion high-fidelity DNA Polymerase (NEB)

using barcoded Illumina PCR index primers. Libraries were purified on an 8%

polyacrylamide-TBE gel and sequenced on an Illumina NextSeq 500 (GeneLab at

Louisiana State University School of Veterinary Medicine).

PRO-Seq analysis

FastQ files were processed using the PRO-Seq pipeline developed by the Danko lab

(Cornell) https://github.com/Danko-Lab/utils/tree/master/proseq. First, adapters were

trimmed and reads aligned to a concatenated genome file containing hg38, dm3,

EBV (Mutu-I or Akata (20)) and rRNA. Note: the EBV genomes were edited to

remove all but one W repeat region to aid read alignment. SeqMonk software (22)

was used to align the output .bam files (containing the full-length read) to the

drosophila genome for normalisation and to the hg38 and EBV genomes for read

quantification. The IGV genome browser (21) was used for visualisation of the output

.bw files (containing only the 3' final read position). Fold change analysis was

performed using the R package DESeq2 (26).

dREG analysis

The PRO-Seq output .bw files were merged for each replicate using the

mergeBigWigs script from the Danko PRO-Seq pipeline https://github.com/Danko-

Lab/utils/tree/master/proseq. The merged .bw files were then imported into dREG

web-based server, freely available at https://django.dreg.scigap.org/ and dREG

analysis performed (23).

ChIP-Seq and ATAC-Seq data

WIG files of ChIP-Seq data from the EBV genome were downloaded from the EBV

portal (5), freely available at https://ebv.wistar.upenn.edu/. ATAC-Seq data is

available from the NCBI GEO database under accession GSE172476.

Chromatin immunoprecipitation (ChIP)-qPCR Assays.

ChIP-qPCR assays were performed as described previously (44). Quantification of

precipitated DNA was determined using real-time PCR and the delta Ct method for

relative quantitation (ABI 7900HT Fast Real-Time PCR System). Rabbit IgG (Cell

Signaling, 2729S), rabbit anti-CTCF (EMD Millipore, 07-729) and rabbit anti-Rad21

(Abcam, ab992) were used in ChIP assays. Primers for ChIP assays are listed in

Supplement Table S2.

Data availability

Raw data and processed BigWig files have been submitted to the GEO database

and will be made public upon publication. Peer reviewers can request a secure token

to view the data.

Acknowledgements

Portions of this research were conducted with the SuperMic supercomputer

maintained at Louisiana State University (http://www.hpc.lsu.edu), and we thank Dr

Le Yan for his help in maintaining the PRO-Seq pipeline. We thank Thaya Stoufflet

and Dr. Vladimir Chouljenko at the LSU SVM GeneLab for NextSeq 500 sequencing.

We also thank Dr. Claire Birkenheuer for her assistance with PRO-Seq technique

and interesting discussions.

This work was supported by public health service grants R01 DE017336 and R01

CA093606 to PM and R01 AI141968 and R21 AI 148926 to JDB from the National

Institutes of Health.

References

1. Hammerschmidt W. 2015. The Epigenetic Life Cycle of Epstein-Barr Virus.
Curr Top Microbiol Immunol 390:103-17.
2. Minarovits J. 2006. Epigenotypes of Latent Herpesvirus Genomes, p 61-80. *In*
Doerfler W, Böhm P (ed), DNA Methylation: Development, Genetic Disease
and Cancer doi:10.1007/3-540-31181-5_5. Springer Berlin Heidelberg, Berlin,
Heidelberg.
3. Tempera I, Klichinsky M, Lieberman PM. 2011. EBV latency types adopt
alternative chromatin conformations. PLoS pathogens 7:e1002180-e1002180.
4. Phillips JE, Corces VG. 2009. CTCF: master weaver of the genome. Cell
137:1194-1211.
5. Arvey A, Tempera I, Tsai K, Chen H-S, Tikhmyanova N, Klichinsky M, Leslie
C, Lieberman Paul M. 2012. An Atlas of the Epstein-Barr Virus Transcriptome
and Epigenome Reveals Host-Virus Regulatory Interactions. Cell Host &
Microbe 12:233-245.
6. Tempera I, Wiedmer A, Dheekollu J, Lieberman PM. 2010. CTCF prevents
the epigenetic drift of EBV latency promoter Qp. PLoS Pathog 6:e1001048.

7.   Chen H-S, Martin KA, Lu F, Lupey LN, Mueller JM, Lieberman PM, Tempera
544   I. 2014. Epigenetic deregulation of the LMP1/LMP2 locus of Epstein-Barr virus
545   by mutation of a single CTCF-cohesin binding site. Journal of virology
546   88:1703-1713.
547   8.   Lupey-Green LN, Moquin SA, Martin KA, McDevitt SM, Hulse M, Caruso LB,
548   Pomerantz RT, Miranda JL, Tempera I. 2017. PARP1 restricts Epstein Barr
549   Virus lytic reactivation by binding the BZLF1 promoter. Virology 507:220-230.
550   9.   McKenzie J, El-Guindy A. 2015. Epstein-Barr Virus Lytic Cycle Reactivation, p
551   237-261. *In* Münz C (ed), Epstein Barr Virus Volume 2: One Herpes Virus:
552   Many Diseases doi:10.1007/978-3-319-22834-1_8. Springer International
553   Publishing, Cham.
554   10.   Kenney SC, Mertz JE. 2014. Regulation of the latent-lytic switch in Epstein-
555   Barr virus. Seminars in cancer biology 26:60-68.
556   11.   Reusch JA, Nawandar DM, Wright KL, Kenney SC, Mertz JE. 2015. Cellular
557   differentiation regulator BLIMP1 induces Epstein-Barr virus lytic reactivation in
558   epithelial and B cells by activating transcription from both the R and Z
559   promoters. Journal of virology 89:1731-1743.
560   12.   Guo R, Jiang C, Zhang Y, Govande A, Trudeau SJ, Chen F, Fry CJ, Puri R,
561   Wolinsky E, Schineller M, Frost TC, Gebre M, Zhao B, Giulino-Roth L,
562   Doench JG, Teng M, Gewurz BE. 2020. MYC Controls the Epstein-Barr Virus
563   Lytic Switch. Molecular Cell 78:653-669.e8.
564   13.   Arrand JR, Rymo L. 1982. Characterization of the major Epstein-Barr virus-
565   specific RNA in Burkitt lymphoma-derived cells. J Virol 41:376-89.
566   14.   Buratowski S. 2009. Progression through the RNA Polymerase II CTD Cycle.
567   Molecular Cell 36:541-546.
568   15.   Palermo RD, Webb HM, West MJ. 2011. RNA polymerase II stalling promotes
569   nucleosome occlusion and pTEFb recruitment to drive immortalization by
570   Epstein-Barr virus. PLoS pathogens 7:e1002334-e1002334.
571   16.   Birkenheuer CH, Baines JD. 2020. RNA Polymerase II Promoter-Proximal
572   Pausing and Release to Elongation Are Key Steps Regulating Herpes
573   Simplex Virus 1 Transcription. Journal of virology 94:e02035-19.
574   17.   Toth Z, Brulois KF, Wong L-Y, Lee H-R, Chung B, Jung JU. 2012. Negative
575   elongation factor-mediated suppression of RNA polymerase II elongation of
576   Kaposi's sarcoma-associated herpesvirus lytic gene expression. Journal of
577   virology 86:9696-9707.
578   18.   Kwak H, Fuda NJ, Core LJ, Lis JT. 2013. Precise maps of RNA polymerase
579   reveal how promoters direct initiation and pausing. Science 339:950-3.
580   19.   Birkenheuer CH, Danko CG, Baines JD, Sandri-Goldin RM. 2018. Herpes
581   Simplex Virus 1 Dramatically Alters Loading and Positioning of RNA
582   Polymerase II on Host Genes Early in Infection. Journal of Virology
583   92:e02184-17.
584   20.   Lin Z, Wang X, Strong MJ, Concha M, Baddoo M, Xu G, Baribault C, Fewell
585   C, Hulme W, Hedges D, Taylor CM, Flemington EK. 2013. Whole-genome
586   sequencing of the Akata and Mutu Epstein-Barr virus strains. J Virol 87:1172-
587   82.
588   21.   Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics
589   Viewer (IGV): high-performance genomics data visualization and exploration.
590   Brief Bioinform 14:178-92.
591   22.   Institute B. SeqMonk.
592   http://www.bioinformatics.babraham.ac.uk/projects/seqmonk/. Accessed

593   23.   Wang Z, Chu T, Choate LA, Danko CG. 2019. Identification of regulatory
594   elements from nascent transcription using dREG. Genome Res 29:293-303.
595   24.   Murata T. 2014. Regulation of Epstein–Barr virus reactivation from latency.
596   Microbiology and Immunology 58:307-317.
597   25.   Price AM, Luftig MA. 2015. To Be or Not IIb: A Multi-Step Process for Epstein-
598   Barr Virus Latency Establishment and Consequences for B Cell
599   Tumorigenesis. PLOS Pathogens 11:e1004656.
600   26.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
601   dispersion for RNA-seq data with DESeq2. Genome Biology 15:550.
602   27.   Andersson R, Gebhard C, Miguel-Escalada I, Hoof I, Bornholdt J, Boyd M,
603   Chen Y, Zhao X, Schmidl C, Suzuki T, Ntini E, Arner E, Valen E, Li K,
604   Schwarzfischer L, Glatz D, Raithel J, Lilje B, Rapin N, Bagger FO, Jørgensen
605   M, Andersen PR, Bertin N, Rackham O, Burroughs AM, Baillie JK, Ishizu Y,
606   Shimizu Y, Furuhata E, Maeda S, Negishi Y, Mungall CJ, Meehan TF,
607   Lassmann T, Itoh M, Kawaji H, Kondo N, Kawai J, Lennartsson A, Daub CO,
608   Heutink P, Hume DA, Jensen TH, Suzuki H, Hayashizaki Y, Müller F, Forrest
609   ARR, Carninci P, Rehli M, Sandelin A, et al. 2014. An atlas of active
610   enhancers across human cell types and tissues. Nature 507:455-461.
611   28.   Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E,
612   Sheffield NC, Stergachis AB, Wang H, Vernot B, Garg K, John S, Sandstrom
613   R, Bates D, Boatman L, Canfield TK, Diegel M, Dunn D, Ebersol AK, Frum T,
614   Giste E, Johnson AK, Johnson EM, Kutyavin T, Lajoie B, Lee B-K, Lee K,
615   London D, Lotakis D, Neph S, Neri F, Nguyen ED, Qu H, Reynolds AP, Roach
616   V, Safi A, Sanchez ME, Sanyal A, Shafer A, Simon JM, Song L, Vong S,
617   Weaver M, Yan Y, Zhang Z, Zhang Z, Lenhard B, Tewari M, Dorschner MO,
618   Hansen RS, et al. 2012. The accessible chromatin landscape of the human
619   genome. Nature 489:75-82.
620   29.   Zhou H, Schmidt SCS, Jiang S, Willox B, Bernhardt K, Liang J, Johannsen
621   EC, Kharchenko P, Gewurz BE, Kieff E, Zhao B. 2015. Epstein-Barr virus
622   oncoprotein super-enhancers control B cell growth. Cell host & microbe
623   17:205-216.
624   30.   Puglielli MT, Woisetschlaeger M, Speck SH. 1996. oriP is essential for EBNA
625   gene promoter activity in Epstein-Barr virus-immortalized lymphoblastoid cell
626   lines. J Virol 70:5758-68.
627   31.   Hammerschmidt W, Sugden B. 1988. Identification and characterization of
628   oriLyt, a lytic origin of DNA replication of Epstein-Barr virus. Cell 55:427-33.
629   32.   Djavadian R, Chiu Y-F, Johannsen E. 2016. An Epstein-Barr Virus-Encoded
630   Protein Complex Requires an Origin of Lytic Replication In Cis to Mediate
631   Late Gene Transcription. PLoS pathogens 12:e1005718-e1005718.
632   33.   Lam MTY, Li W, Rosenfeld MG, Glass CK. 2014. Enhancer RNAs and
633   regulated transcriptional programs. Trends in Biochemical Sciences 39:170-
634   182.
635   34.   Higgs DR. 2020. Enhancer–promoter interactions and transcription. Nature
636   Genetics 52:470-471.
637   35.   Kubo N, Ishii H, Xiong X, Bianco S, Meitinger F, Hu R, Hocker JD, Conte M,
638   Gorkin D, Yu M, Li B, Dixon JR, Hu M, Nicodemi M, Zhao H, Ren B. 2021.
639   Promoter-proximal CTCF binding promotes distal enhancer-dependent gene
640   activation. Nat Struct Mol Biol 28:152-161.
641   36.   Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang
642   Y, Lu Y, Wu Y, Jia Z, Li W, Zhang MQ, Ren B, Krainer AR, Maniatis T, Wu Q.

2015. CRISPR Inversion of CTCF Sites Alters Genome Topology and
Enhancer/Promoter Function. Cell 162:900-10.
37. Lupey-Green LN, Caruso LB, Madzo J, Martin KA, Tan Y, Hulse M, Tempera
I, Jung JU. 2018. PARP1 Stabilizes CTCF Binding and Chromatin Structure
To Maintain Epstein-Barr Virus Latency Type. Journal of Virology 92:e00755-
18.
38. Countryman J, Gradoville L, Bhaduri-McIntosh S, Ye J, Heston L, Himmelfarb
S, Shedd D, Miller G. 2009. Stimulus duration and response time
independently influence the kinetics of lytic cycle reactivation of Epstein-Barr
virus. Journal of virology 83:10694-10709.
39. Schaeffner M, Mrozek-Gorska P, Buschle A, Woellmer A, Tagawa T,
Cernilogar FM, Schotta G, Krietenstein N, Lieleg C, Korber P,
Hammerschmidt W. 2019. BZLF1 interacts with chromatin remodelers
promoting escape from latent infections with EBV. Life Science Alliance
2:e201800108.
40. Luan J, Xiang G, Gómez-García PA, Tome JM, Zhang Z, Vermunt MW,
Zhang H, Huang A, Keller CA, Giardine BM, Zhang Y, Lan Y, Lis JT,
Lakadamyali M, Hardison RC, Blobel GA. 2021. Distinct properties and
functions of CTCF revealed by a rapidly inducible degron system. Cell reports
34:108783-108783.
41. Jonkers I, Kwak H, Lis JT. 2014. Genome-wide dynamics of Pol II elongation
and its interplay with promoter proximal pausing, chromatin, and exons. Elife
3:e02407.
42. Birkenheuer CH, Danko CG, Baines JD. 2018. Herpes Simplex Virus 1
Dramatically Alters Loading and Positioning of RNA Polymerase II on Host
Genes Early in Infection. Journal of virology 92:e02184-17.
43. Mahat DB, Kwak H, Booth GT, Jonkers IH, Danko CG, Patel RK, Waters CT,
Munson K, Core LJ, Lis JT. 2016. Base-pair-resolution genome-wide mapping
of active RNA polymerases using precision nuclear run-on (PRO-seq). Nat
Protoc 11:1455-76.
44. Chau Charles M, Lieberman Paul M. 2004. Dynamic Chromatin Boundaries
Delineate a Latency Control Region of Epstein-Barr Virus. Journal of Virology
78:12308-12319.
676

Figure Legends

Figure 1: Distribution of active polymerase on the EBV episome in latent Mutu-

I cells. (A) Genome browser view of the distribution of PRO-Seq reads (normalised

to *Drosophila melanogaster* spike-in) along the EBV episome in latent Mutu-I cells.

Representative sample of 2-3 biological replicates. The W repeat region was deleted

to include only 1 repeat for the sequencing alignment. The IGV window was set to

maximum for each track. (B) Violin plot of the Log10 transformed mean normalised

read count of each EBV transcript during latency. The transcripts with a mean of

>1000 normalised reads are labelled. (C) Normalised sequencing read count of each

EBV latency promoter during latency in Mutu-I cells. Data represents mean of 3

biological replicates, error bars indicate SEMs.

Figure 2: Distribution and quantification of active polymerase on the EBV

episome during reactivation in Mutu-I cells. (A) Genome browser view of the

distribution of PRO-Seq reads (normalised to *Drosophila melanogaster* spike-in)

along the EBV episome in latent Mutu-I cells and at 1, 4 and 12h post-reactivation

with NaB/TPA. Representative sample of 2-3 biological replicates. The W repeat

region was deleted to include only 1 repeat for sequencing alignment purposes. The

IGV window was set to maximum for each track. Volcano plots showing the DeSeq2

calculated fold changes of PRO-Seq reads for each EBV transcript relative to latency

at (B) 1h post-reactivation, (C) 4h post-reactivation and (D) 12h post-reactivation.

Figure 3: PRO-Seq identifies distinct patterns of Pol activity on the EBV

episome during reactivation. High resolution IGV view of PRO-Seq tracks from

EBV genome regions with the most significant increase in Pol activity during early

reactivation in Mutu-I cells. Cells were treated with NaB/TPA to induce reactivation.

(A) LMP-2A, exon 1 region. (B) EBER region. (C) RPMS1 region. Representative

sample of 2-3 biological replicates. Sequencing reads were normalised to *Drosophila*

melanogaster spike-in. PRO-Seq peaks of interest are highlighted in purple. Solid

red indicates coding regions, black lines indicate non-coding regions of transcripts.

Transcription start sites and promoter regions are in green.

Figure 4: dREG detection of transcription regulatory elements from PRO-Seq

corresponds with CTCF binding sites on the EBV episome. (A) IGV view of the

PRO-Seq results from latent and reactivated (NaB/TPA treated) Mutu-I cells after

analysis with the dREG software package to identify transcription initiation regions

(TIRs), aligned to transcription factor ChIP-Seq on the latent EBV episome. (B) IGV

view of PRO-Seq results from latent and reactivated (anti-IgG treated) Akata cells

after analysis with dREG aligned to CTCF ChIP-Seq on the EBV episome.

Significant dREG peaks are highlighted in green (false discovery rate (FDR) ≤0.05).

ChIP-Seq data from (5).

Figure 5: Alignment of CTCF ChIP-Seq peaks, dREG and PRO-Seq indicates

pausing of Pol at CTCF sites on EBV genome. IGV view of aligned dREG, PRO-

Seq and CTCF ChIP-Seq peaks from latent and reactivated EBV in Mutu-I cells

(TPA/NaB). (A) LMP-2A, exon 1 region. (B) RPMS1 region. (C) EBER region. (D) Qp

promoter. (E) Cp promoter. Representative sample of 2-3 biological replicates.

Significant dREG peaks are highlighted in green (false discovery rate (FDR) ≤0.05).

Figure 6: Pol accumulates at sites of accessible chromatin on the EBV

episome during early reactivation. IGV view of the PRO-Seq results from latent

and reactivated (NaB/TPA treated) Mutu-I cells after analysis with the dREG

software package to identify transcription initiation regions (TIRs). Aligned in

comparison to CTCF ChIP-Seq and ATAC-Seq from 2 sub populations of latent

Mutu-I cells on the EBV episome. Significant dREG peaks are highlighted in green

(false discovery rate (FDR) ≤0.05).

Figure 7: CTCF binding alterations on the EBV episome during reactivation.

ChIP-qPCR at CTCF binding sites on the EBV and human genomes in latent and at

4h post-reactivation Mutu-I cells (NaB/TPA treated) (A, C and E) and latent 4h post-

reactivation Akata cells (anti-IgG treated) (B, D and F). Using antibodies against (A,

B) Pol II, (B, C) CTCF and (E, F) RAD21.

Supporting Information

Figure S1: Distribution and quantification of active polymerase on the EBV

episome during reactivation in Akata cells. (A) Genome browser view of the

distribution of PRO-Seq reads (normalised to *Drosophila melanogaster* spike-in)

along the EBV episome in latent Akata cells and at 1, 4 and 12h post-reactivation

with anti-IgG. Representative sample of 3 biological replicates. The W repeat region

was deleted to include only 1 repeat for sequencing alignment purposes. The IGV

window was set to maximum for each track. Volcano plots showing the DeSeq2

calculated fold changes of PRO-Seq reads for each EBV transcript relative to latency

at (B) 1h post-reactivation, (C) 4h post-reactivation and (D) 12h post-reactivation.

Figure S2: Alignment of CTCF ChIP-Seq peaks, dREG and PRO-Seq from Akata

cells indicates pausing of Pol at CTCF sites on EBV genome. IGV view of

aligned dREG, PRO-Seq and CTCF ChIP-Seq peaks from latent and reactivated

EBV in Akata cells (Anti-IgG treated). (A) LMP-2A, exon 1 region. (B) RPMS1 region.

(C) EBER region. (D) Qp promoter. (E) BKRF region. Representative sample of 3

biological replicates. Significant peaks are highlighted in green (false discovery rate

(FDR) ≤0.05).

Figure S3: RT-qPCR validation of EBV reactivation. Relative expression of EBV

transcripts in untreated and reactivated cells. (A) Mutu-I cells untreated and 24 h

post-treatment with NaB/TPA. (B) Akata cells untreated and at 4 and 24 h post-

treatment with anti-IgG. Zta: BZLF1, RTA: BRLF1, EA-D: early antigen D.

Figure S4: CTCF binding alterations on the EBV episome during reactivation.

ChIP-qPCR at CTCF binding sites on the EBV and human genomes in latent and at

4 and 24 h post-reactivation Mutu-I cells (NaB/TPA treated) using antibody against

CTCF.

Table S1: Location of dREG peaks on the NC_007605.1 EBV genome in Mutu-I

and Akata cells

Table S2: Primers used for ChiP-qPCR