1 **Novel Feline Papillomavirus Isolate P20 Assembled from Metagenomic Data**
2 **Isolated from Human Skin of a House Cat Owner**
3

4 Ema H. Graham [1], Michael S. Adamowicz [2], Peter C Angeletti [3,4], Jennifer L. Clarke [5,6],
5 Samodha C. Fernando [7], Joshua R. Herr [4,8,9,#]
6
[1] PhD Program in Complex Biosystems, University of Nebraska, 68583, Lincoln, NE,
USA
[2] College of Agricultural Sciences & Natural Resources, University of Nebraska, Lincoln,
NE 68583, USA
[3] Nebraska Center for Virology, University of Nebraska, Lincoln, NE 68583
[4] School of Biological Sciences, University of Nebraska, Lincoln, NE 68588
[5] Department of Statistics, University of Nebraska, Lincoln, NE, 68588, USA
[6] Food Science and Technology Department, University of Nebraska, Lincoln, NE
68588 USA
[7] Department of Animal Science, University of Nebraska, Lincoln, NE, 68583, USA
[8] Department of Plant Pathology, University of Nebraska, Lincoln, NE 68503, USA
[9] Center for Plant Science Innovation, University of Nebraska, Lincoln, NE 68503, USA
19
[#] Corresponding Author
Joshua R. Herr, PhD;
422 Plant Sciences Hall
University of Nebraska
Lincoln NE 68583
Email: jherr@unl.edu
Phone: +1 (402) 472-1352

27
28 **RUNNING TITLE**: Feline papillomavirus from human skin metagenome
29
30 **KEYWORDS**
31 Virome, Metagenome, Human Skin, Papilloma Virus, Feline Papillomaviridae
32
33

ABSTRACT

A novel feline papillomavirus isolate was assembled from metagenomic sequencing

data collected from the human skin of a house cat owner. This circular papillomavirus

isolate P20 is 8069 bp in length and displays genome organization typical of feline

papillomaviruses, but only exhibits approximately 75% synteny to other feline

papillomaviruses.

ANNOUNCEMENT

Papillomaviruses are most well-known for infecting humans, but they also broadly infect

mammals, reptiles, and birds (1). Classification of papillomaviruses has largely focused

on the sequence divergence of the Major Capsid Protein L1, which is conserved across

the papilloma viruses (1). The broad phylogenetic relationships of mammalian papilloma

viruses suggest that their evolution acutely mirrors their host relationships (2). In recent

years, numerous studies have reported on the discovery and diversity of

papillomaviruses associated with domestic and wild cats (3,4,5). Currently, seven types

of feline papillomaviruses are recognized that may or may not produce both skin and

oral lesions (5,6). Multiple studies in the last year have tied feline papilloma viruses to

skin and oral squamous cell carcinomas in house cats (7,8,9,10).

In a previous study addressing the diversity and stability of viruses on human skin, we

sampled three locations – left hand, right hand, and scalp – over a longitudinal 6-month

period (11). Viral enriched metagenomic DNA was extracted from skin swabs and

sequenced on an Illumina Hi-Seq sequencer as previously described (11). In that study,

a questionnaire was given to participants to collect pertinent information about subject

lifestyle and environmental variables, such that we could understand how the human

skin virome is shaped and maintained. Questions on the survey included the frequency

and locations of travel, skin care habits, and recent contact with animals (i.e. types of

pets). When evaluating viral diversity of the virome samples it was noted that one study

participant had a metagenomic virome profile that consistently showed synteny to a

feline papilloma virus. That individual had also identified in their questionnaire as being

an owner of a domesticated house cat.

To further investigate this observation, a dataset containing the viral metagenomic

assembly of compiled data from that specific human individual was generated. This

included 15 different metagenomic samples from across three body locations and five

sampling times. In our previous study, a combination of laboratory techniques and

bioinformatic tools to isolate the viral component of human skin was employed (11).

Here, utilizing the virome sample reads from that study, sequencing reads were first

mapped to Papillomaviridae reference database material from NCBI using BBMap

v38.94 (12) with flags and parameters described for high precision mapping as

suggested in the BBMap Guide. Mapped reads were the further processed using

KHMER v2.0.0 (13) to remove singleton reads and high copy number reads using k-mer

profiles. All filtered papilloma virus reads were then assembled using MEGAHIT v1.2.8

(14) and the subsequent assembly quality was assessed using QUAST v5.0.2 (15). To

double check the assembly, the putative viral contigs – designated as contigs containing

annotated papilloma virus genes – were identified using the tool CheckV v0.7.0 (16).

Contigs were further classified using both nucleotide-based classification tools, such as

Kraken2 v2.0.8-beta (17), Demovir (https://github.com/feargalr/Demovir), Blastn (with a

>10% query coverage cut-off) (18), and using a classification tool based on protein

coding sequences, Kaiju v1.7 (19). Based on these assembly and contig taxonomic

identification methods, a feline papilloma virus genome assembly was identified which

showed 74% percent similarity to feline papilloma virus type 2. Genes of this single virus

were annotated with PROKKA v1.14.5 (20) using the embedded viral annotation

database and Cenote-Taker 2 v. 2.1 (21).

We identified a complete, circular feline papilloma virus that was 8069 bp in length

(Figure 1) which we identified as isolate P20. Through annotation using PROKKA

v1.14.5, six open reading frames which included the E6 Protein, E7 protein, E1 protein,

E2 protein, Late Protein L2, and Major capsid protein L1 were predicted (Figure 2).

These six open reading frames are shared among papillomaviruses. Genome maps

(Figure 1, Figure 2) were visualized using the SnapGene software (GSL Biotech LLC,

San Diego, CA; https://snapgene.com). Using a BLASTn search in NCBI GenBank, we

found that the papilloma virus showing the greatest synteny to our genome assembly

was isolated from the skin of a domestic Maine Coon house cat in 2007 (GenBank

accession number NC038520). Our isolate showed 70.3% amino acid synteny for all

coding regions when compared to that accession.

To determine the phylogenetic placement of isolate P20 identified in this study, we

downloaded all papilloma viruses identified as having a feline host on NCBI (along with

two outgroups from closely related hosts – domestic dog and Weddell Seal) and

annotated open reading frames using PROKKA v1.14.5 and aligned the concatenated

amino acid sequences using MUSCLE v3.8.1551 (22) for the six genes (E6 Protein, E7

protein, E1 protein, E2 protein, Late Protein L2, and Major capsid protein L1) shared

among all the feline papilloma available on NCBI. After removal of uninformative amino

acid regions using GBLOCKS v0.91b (23), a phylogenetic tree (Figure 3) of the isolates

was generated using IQTREE v1.6.12 (24) under the LG+F+I+G4 substitution model

optimally determined by MODEL-FINDER v1.5.4 (25). This phylogenetic tree (Figure 3),

visualized with FIGTREE v1.4.4 (https://github.com/rambaut/figtree), placed our isolate,

in close proximity to other feline papilloma type 2 accessions.

DATA AVAILABILITY

All raw sequencing data has been deposited in the NCBI Short Read Archive (SRA)

under the accession code PRJNA754140. The complete feline papillomavirus isolate

P20 genome has been submitted to NCBI Genbank with the accession number of

OL310516. All metadata, sequences, annotation files, and scripts used here are publicly

available and archived at:

https://github.com/HerrLab/Graham_2021_feline_papilloma_P20.

FUNDING SOURCES

This work was supported by the Department of Justice (grant numbers 2017-IJ-CX-

0025, 2019-75-CX-0075, and 2019-R2-CX-0048). This funding agency had no role in

study design, data collection and interpretation, or the decision to submit the work for

publication.

DECLARATION OF COMPETING INTERESTS

None to declare.

ACKNOWLEDGMENTS

This work was completed using the Holland Computing Center of the University of

Nebraska, which receives support from the Nebraska Research Initiative.

We thank all the anonymous participants for their contribution to this study. We also

thank members of the Fernando Lab who assisted with advice on laboratory methods.

REFERENCES

1. Rector, A. and Van Ranst, M., 2013. Animal papillomaviruses. *Virology*, *445*(1-2), pp.213-223.

2. Rector, A., Lemey, P., Tachezy, R., Mostmans, S., Ghim, S.J., Van Doorslaer, K., Roelke, M., Bush, M., Montali, R.J., Joslin, J. and Burk, R.D., 2007. Ancient papillomavirus-host co-speciation in Felidae. *Genome biology*, *8*(4), pp.1-12.

3. Sundberg, J.P., Van Ranst, M., Montali, R., Homer, B.L., Miller, W.H., Rowland, P.H., Scott, D.W., England, J.J., Dunstan, R.W., Mikaelian, I. and Jenson, A.B., 2000. Feline papillomas and papillomaviruses. *Veterinary Pathology*, *37*(1), pp.1-10.

4. Munday, J.S., 2014. Papillomaviruses in felids. *The Veterinary Journal*, *199*(3), pp.340-347.

5. Munday, J.S. and Thomson, N.A., 2021. Papillomaviruses in Domestic Cats. *Viruses*, *13*(8), p.1664.

6. Kaynarcalidan, O. and Oğuzoğlu, T.Ç., 2021. The oncogenic pathways of papillomaviruses. *Veterinary and Comparative Oncology*, *19*(1), pp.7-16.

7. Altamura, G., Cardeti, G., Cersini, A., Eleni, C., Cocumelli, C., Bartolomé del Pino, L.E., Razzuoli, E., Martano, M., Maiolino, P. and Borzacchiello, G., 2020. Detection of *Felis catus* papillomavirus type-2 DNA and viral gene expression suggest active infection in feline oral squamous cell carcinoma. *Veterinary and comparative oncology*, *18*(4), pp.494-501.

8. Carrai M, Van Brussel K, Shi M, Li C-X, Chang W-S, Munday JS, Voss K, McLuckie A, Taylor D, Laws A, Holmes EC, Barrs VR, Beatty JA. 2020. Identification of a Novel Papillomavirus Associated with Squamous Cell Carcinoma in a Domestic Cat. Viruses 12(1):124. https://doi.org/10.3390/v12010124

9. Chu, S., Wylie, T.N., Wylie, K.M., Johnson, G.C., Skidmore, Z.L., Fleer, M., Griffith, O.L. and Bryan, J.N., 2020. A virome sequencing approach to feline oral squamous cell carcinoma to evaluate viral causative factors. *Veterinary microbiology*, *240*, p.108491.

10. Ito, S., Chambers, J.K., Sumi, A., Yamashita-Kawanishi, N., Omachi, T., Haga, T., Nakayama, H. and Uchida, K., 2021. Involvement of *Felis catus* papillomavirus type 2 in the tumorigenesis of feline Merkel cell carcinoma. *Veterinary pathology*, p.03009858211045440.

11. Graham, E.H., Fernando, S., Clarke, J., Herr, J.R. and Adamowicz, M., 2021. The Application of The Skin Virome for Human Identification. *bioRxiv*. doi: https://doi.org/10.1101/2021.09.10.459834

12. Bushnell, B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. https://sourceforge.net/projects/bbmap/

13. Crusoe M.R., Alameldin H.F., Awad S., Boucher E., Caldwell A., Cartwright R., Charbonneau A., Constantinides B., Edvenson G., Fay S., Fenton J., Fenzl T., Fish J., Garcia-Gutierrez L., Garland P., Gluck J., González I., Guermond S., Guo J., Gupta A., Herr J.R., Howe A., Hyer A., Härpfer A., Irber L., Kidd R., Lin D., Lippi J., Mansour T., McA'Nulty P., McDonald E., Mizzi J., Murray K.D., Nahum J.R., Nanlohy K., Nederbragt A.J., Ortiz-Zuazaga H., Ory J., Pell J., Pepe-Ranney C., Russ Z.N., Schwarz E., Scott C., Seaman J., Sievert S., Simpson J., Skennerton C.T., Spencer J., Srinivasan R., Standage D., Stapleton J.A., Steinman S.R., Stein J., Taylor B., Trimble W., Wiencko H.L., Wright M., Wyss B., Zhang Q., zyme e., Brown C.T., 2015. The khmer software package: enabling efficient nucleotide sequence analysis. F1000Res. 2015 Sep 25; 4:900. doi: 10.12688/f1000research.6924.1.

14. Li, D., Liu, C.M., Luo, R., Sadakane, K. and Lam, T.W., 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, *31*(10), pp.1674-1676.

15. Gurevich, A., Saveliev, V., Vyahhi, N. and Tesler, G., 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics 29*(8), pp.1072-1075.

16. Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S. and Kyrpides, N.C., 2021. CheckV assesses the quality and completeness of metagenome-assembled viral genomes. *Nature biotechnology*, *39*(5), pp.578-585.

17. Wood, D.E. and Salzberg, S.L., 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. *Genome biology*, *15*(3), pp.1-12.

18. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. and Madden, T.L., 2009. BLAST+: architecture and applications. *BMC bioinformatics*, *10*(1), pp.1-9.

19. Menzel, P., Ng, K.L., Krogh, A., 2016. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nature communications*, *7*(1), pp.1-9.

20. Seemann T. 2014. *Prokka: rapid prokaryotic genome annotation*. Bioinformatics 2014 Jul 15;30(14):2068-9.

21. Tisza M.J., Belford A.K., Domínguez-Huerta G., Bolduc B., Buck C.B., 2021. Cenote-Taker 2 democratizes virus discovery and sequence annotation, Virus

Evolution, Volume 7, Issue 1, January 2021, veaa100,
https://doi.org/10.1093/ve/veaa100

22. Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research*, *32*(5), pp.1792-1797.

23. Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Molecular Biology and Evolution 17, 540-552.

24. Minh B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear R. (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.*, 37:1530-1534. https://doi.org/10.1093/molbev/msaa015

25. Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nat. Methods*, 14:587-589. https://doi.org/10.1038/nmeth.4285

248 **Figure 1**. Circular genome map, which is 8069 bp in length, of Feline papillomavirus
249 type 2 like virus. From outside to center: open reading frame annotations of E6 Protein,
250 E7 protein, E1 protein, E2 protein, Late Protein L2, and Major capsid protein L1 (inside
251 purple arrows)
252

Figure 2. Linear genome map, 8069 bp in length, of Feline papillomavirus type 2 like
virus. Top portion of the figure shows %GC content for regions of the circular genome.
Higher GC content are shown in warmer colors (red) and lower GC content are shown
in cooler colors (blue). Below %GC content is the location of the six gene regions in the
viral genome consisting of open reading frame annotations of E6 Protein, E7 protein, E1
protein, E2 protein, Late Protein L2, and Major capsid protein L1 (inside purple arrows)
266

Feline papillomavirus type 2 virus isolate P20
8069 bp

267
268
269

270 **Figure 3.** Phylogenetic tree of the placement of the feline type 2 papilloma virus
271 identified in this study along with nine other feline papilloma viruses. The tree was
272 rooted with Canine papilloma virus type 2 and *Leptonychotes weddelii* (Wedell Seal)
273 papilloma virus type 5.
274
275

Feline papillomavirus type 2 virus isolate P20

8069 bp

LC612600 *Felis* papillomavirus 19191gA

NC038520 Feline papillomavirus type 2 isolate Maine-Coon-2007

Feline papillomavirus type 2 isolate P20

KY853656 *Felis catus* papillomavirus type 5

LC333412 *Felis catus* papillomavirus 4 isolate 13136

LC333413 *Felis catus* papillomavirus 4 isolate 141110

MZ357115 *Felis catus* papillomavirus isolate Co064PV01-10

KF147892 *Felis catus* papillomavirus 4 isolate MY-3

MN857145 *Felis catus* papillomavirus type 6

JX972168 *Felis catus* papillomavirus type 3

KY825188 *Felis catus* papillomavirus type 3 isolate Missouri

OUTGROUP AY722648 Canine papillomavirus type 2

OUTGROUP MG571094 *Leptonychotes weddellii* papillomavirus type 5 isolate 12975