1

5   **High-Speed Imaging Reveals the Bimodal Nature of Dense Core**

6   **Vesicle Exocytosis and Jet Flow through the Fusion Pore**

8   Pengcheng Zhang[1], David Rumschitzki[2] and Robert H. Edwards[1],*

[1] Departments of Neurology and Physiology

School of Medicine

University of California, San Francisco

San Francisco, CA 94158

U.S.A.

[2] Department of Chemical Engineering

City College of New York

New York, NY 10031

U.S.A.

* Correspondence: robert.edwards@ucsf.edu

ABSTRACT

During exocytosis, the fusion of secretory vesicle with plasma membrane forms a pore that

regulates release of neurotransmitter and peptide. Osmotic forces contribute to exocytosis but

release through the pore is thought to occur by diffusion. Heterogeneity of fusion pore behavior

has also suggested stochastic variation in a common exocytic mechanism, implying a lack of

biological control. Imaging at millisecond resolution to observe the first events in exocytosis, we

find that fusion pore duration is bimodal rather than stochastic. Loss of calcium sensor

synaptotagmin 7 increases the proportion of slow events without changing the intrinsic properties

of either class, indicating the potential for independent regulation. In addition, dual imaging shows

a delay in the entry of external dye relative to release that indicates discharge at high velocity

rather than strictly by diffusion.

KEYWORDS

1 INTRODUCTION

Regulated exocytosis provides a mechanism to control the timing of signals important for

development, physiology and behavior. The latency from stimulus to membrane fusion, or

synchrony, makes an important contribution to the time course of signaling. In neurons, synaptic

vesicle exocytosis within milliseconds of stimulation confers the synchronous release of

neurotransmitter essential for brain function (Kaeser and Regehr, 2014). In endocrine cells,

release from dense core vesicles (DCVs) exhibits a biphasic time course with initial, rapid release

followed by slow, sustained secretion (Parsons et al., 1995; Voets et al., 1999). The biphasic

release of insulin from pancreatic β-cells plays a critical role in glucose homeostasis, and loss of

the first phase is considered a hallmark of type 2 diabetes (Rorsman and Renstrom, 2003). The

initial rapid phase of release involves a readily releasable pool of vesicles at the plasma

membrane, whereas the slower phase requires activation of additional vesicles through docking

or priming. The variation in synchrony within individual cells is thus thought to reflect vesicles at

different steps along a common pathway to exocytosis.

After membrane fusion, release from individual vesicles also varies in time course. On

fusion, a channel known as the fusion pore connects the vesicle lumen to the extracellular space

(Breckenridge and Almers, 1987). The properties and behavior of the fusion pore affect the time

course of release in multiple ways. Pore size influences the rate of release: narrow pores slow

secretion and wide pores accelerate (Bao et al., 2018; Chow et al., 1992; Shin et al., 2018). The

fusion pore can also act as a molecular sieve that differentially affects the release of cargo based

on their size or charge (Fulop et al., 2005; Perrais et al., 2004). In addition, the fusion pore can

dilate, leading to the discharge of all contents and vesicle collapse into the plasma membrane.

Alternatively, the fusion pore can reseal, resulting in a form of exocytosis known as 'kiss-and-run'

that preserves the vesicle for additional rounds of release (Alabi and Tsien, 2013), thus prolonging

the response. The fusion pore influences release from DCVs in many cell types including neurons

as well as endocrine cells (Breckenridge and Almers, 1987; Logan et al., 2017; Xia et al., 2009).

The fusion pore also exhibits remarkable heterogeneity within individual cells. Since the analysis

of synchrony has suggested a single pathway to exocytosis, the heterogeneity in pore behavior

has been attributed to stochastic variation, which implies a lack of biological control.

Previous work has characterized the diversity of fusion pore behavior by classifying

exocytic events based on specific characteristics. For example, DCV exocytosis has been imaged

using fluorescent reporters, and the time course of fluorescence classified based on the shape of

the curve (Chiang et al., 2014; Logan et al., 2017). Similarly, the amperometric measurement of

catecholamine release shows spikes indicating rapid pore expansion and 'feet' suggestive of

narrow pores, and these are analyzed separately (Zhou et al., 1996). The properties of each

class are thus quantified using different metrics, making it difficult to determine whether the

classes differ in mechanism or reflect only stochastic variation.

In addition, the available methods reveal only specific features of the exocytic process.

Amperometry has high temporal resolution but lacks spatial information and does not monitor

DCV behavior before fusion. In addition, diffusion makes the measurement of event kinetics

sensitive to the distance between release site and electrode (Albillos et al., 1997). In contrast,

imaging by total internal reflection fluorescence (TIRF) microscopy provides high spatial

resolution and can monitor DCV behavior before exocytosis. However, the weak emission of

conventional fluorophores requires long exposure times, limiting temporal resolution. The inability

to monitor simultaneously multiple features of an individual exocytic event produces a fragmentary

description, potentially failing to capture a relationship between features that has important

implications for mechanism.

We now describe a method for imaging that detects secretory vesicles before fusion and

monitors release at high temporal resolution, providing a holistic picture of exocytosis that enables

us to characterize the diversity of fusion pore behavior. We identify two distinct modes of fusion

pore behavior, showing that stochasticity alone cannot account for the diversity.

RESULTS

Imaging Exocytic Events at Millisecond Resolution

Fluorescent false neurotransmitters (FFNs) are fluorescent catecholamine analogues

recognized as substrates by the vesicular monoamine transporter (VMAT) and hence

concentrated into neurosecretory vesicles (Gubernator et al., 2009; Hu et al., 2013). The pH-

insensitive FFN206 (Figure 1—figure supplement 1A,B), which is not quenched by the acidic

lumenal environment (Figure 1—figure supplement 1C,D), enables us to detect vesicles before

fusion. When loaded into primary adrenal chromaffin cells, FFN206 accumulates in chromaffin

granules, presumably due to the catalytic activity of VMAT, conferring a high signal-to-noise ratio

that enables imaging at millisecond resolution. Using TIRF microscopy, we observe a punctate

distribution of FFN-loaded vesicles illuminated by the evanescent field (Figure 1A), presumably

vesicles docked at the plasma membrane. Stimulation produces a large number of transient

events that we imaged at an acquisition rate of 800 Hz (Figure 1B–D, Figure1—figure supplement

2, and Figure 1—video 1). For each event, we observe an increase in fluorescence suggestive of

dye approaching the TIRF plane at the coverslip surface due to exocytosis. Consistent with an

exocytic mechanism, all FFN events require stimulation (Figure 1—video 1) and external Ca^{++}

(Figure 2A). Cleavage of v-SNAREs by tetanus toxin (McMahon et al., 1993; Schiavo et al., 1992)

abolishes the FFN response (Figure 2B), further supporting an exocytic origin for all events.

The majority of FFN events involve a characteristic, brief spike in fluorescence (Figure 1B,

Figure 1—figure supplement 2A, and Figure 1—video 2) followed by dispersion (Figure 2H),

consistent with release and subsequent diffusion of a small molecule. These fast events

presumably reflect release through a large fusion pore. Almost all events result in reduction of

baseline fluorescence (Figure 1B,C, and Figure 1—figure supplement 2A,B), which we interpret

as the release of all FFN from vesicles docked at the plasma membrane. However, a small fraction

(1–10%) of exocytic events do not involve a change in baseline fluorescence (Figure 1D, Figure

1—figure supplement 2C, and Figure 1—video 3), suggesting a vesicle that arrives from the cell

interior (new arrival). To exclude the possibility of exocytosis occurring outside the TIRF plane,

we counted only events that begin as a point source. Using the fluorescence intensity of cell-

impermeant Alexa dye in the evanescent field as a proxy for the tightness of cell attachment

(Figure 1—figure supplement 3A–C), we also found no difference in attachment tightness at the

sites of new arrivals compared to docked events (Figure 1—figure supplement 3D), confirming

that both event types occurred within the evanescent field. It remains possible that invagination

of the plasma membrane enables fusion outside the TIRF plane, and this might be expected to

reduce the magnitude of fluorescence increase. However, the peak amplitude of fast events

showed no difference whether a docked vesicle was detected or not (Figure 1—figure supplement

3E). Despite the elaborate choreography of proteins required for neurotransmitter release (Jahn

and Fasshauer, 2012), tethering, docking, priming and fusion can apparently occur within 1–2 ms

of vesicle arrival at the plasma membrane.

A substantial fraction of events persists from hundreds of milliseconds to seconds,

exhibiting highly variable fluorescence traces without a recognizable spike (Figure 1C, Figure 1—

figure supplement 2B, and Figure 1—video 4). These slow events presumably reflect the efflux of

FFN through a narrow fusion pore that limits diffusion of even a small molecule (Shin et al., 2018).

Amperometric recording of catecholamine release has identified 'stand-alone feet' that reflect the

slow efflux of catecholamine through a narrow fusion pore (van Kempen et al., 2011; Zhou et al.,

1996).   However, most slow FFN events last longer (up to 10-fold) than stand-alone feet,

indicating the extreme stability of the narrow fusion pore. Monitoring chromaffin granules before

exocytosis, we find that slow events derive exclusively from docked vesicles. In contrast, FFN

release by new arrivals is invariably fast. The slow mode of release thus requires a stably docked

intermediate, indicating that the state of docking influences subsequent behavior of the fusion

pore.

Dual Imaging of FFN and Peptide Release

The pH-insensitivity of FFN206 raises the possibility that slow FFN events may arise from vesicle movement within the TIRF plane rather than exocytosis. To monitor opening of the fusion pore by an independent method, we labeled chromaffin granules by expressing brain-derived neurotrophic factor (BDNF) fused to the ecliptic pHluorin, a particularly pH-sensitive form of GFP (Miesenböck et al., 1998). BDNF-pHluorin sorts to DCVs where it is quenched by the low lumenal pH. Opening of the fusion pore exposes lumenal pHluorin to the neutral external pH, increasing its fluorescence. FFN and pHluorin are excited at different wavelengths, enabling us to image both fluorophores at the same events (Figure 2C–G and Figure 1—video 5). The weakly fluorescent BDNF-pHluorin required longer exposure times, reducing the rate of image acquisition and resulting in the loss or truncation of FFN fluorescence increase associated with fast events (Figure 2D,F,G). By dual imaging, we find that the vast majority of FFN events are accompanied by exocytosis of BDNF-pHluorin (147/166, Figure 2C). BDNF-pHluorin unquenching invariably follows release of the much smaller FFN (Figure 2D–G), due at least in part to the effects of lumenal buffering (Holz, 1978). A small number of FFN events are not associated with BDNF-pHluorin unquenching (19/166, Figure 2C), presumably reflecting FFN loaded into vesicles made before exogenous expression of BDNF-pHluorin. A fraction of BDNF-pHluorin events are not associated with FFN release (66/213, Figure 2C), indicating that FFN206 does not label all chromaffin granules.

Fast and slow events are accompanied by BDNF-pHluorin exocytosis, confirming they are both exocytic, consistent with their dependence on stimulation, external Ca^{++}, and v-SNAREs. Corelease of BDNF-pHluorin also shows that FFNs release predominantly if not exclusively from peptidergic DCVs. Thus, DCVs are responsible for the full range of FFN event kinetics, and a class of vesicle other than DCVs does not contribute to the observed variation.

The Fusion Pore Can Limit Entry of External Dye

Since lumenal buffering delays the unquenching of pHluorin reporters (Holz, 1978), the precise moment when the fusion pore opens in slow FFN events remains unclear. We therefore monitored the entry of external Alexa Fluor 488, a small, cell-impermeant fluorophore, as an alternative reporter for fusion pore opening (Chiang et al., 2014; Zhao et al., 2016). Chromaffin cells were loaded with FFN206 and bathed in solution containing Alexa Fluor 488 to monitor both Alexa dye entry and FFN release at the same events. The vast majority of FFN events are accompanied by Alexa dye entry (144/179, Figure 3A–G), consistent with exocytosis. Events identified by Alexa dye entry alone, without FFN release, may represent endocytosis (6/150, Figure 3G), whereas FFN-only events (35/179, Figure 3G) may represent rapid vesicle collapse that prevents dye entry (Chiang et al., 2014). All fast FFN events coincide with simultaneous Alexa dye entry (Figure 3A), as expected. For slow events detectable with both reporters, however, we were surprised to find a difference in timing between the onset of FFN fluorescence increase and Alexa dye entry (Figure 3B–F). In some cases, the FFN event and Alexa entry occur over the same extended period (Figure 3B). More often, dye entry lags the onset of FFN signal increase (Figures 3C,D) and in some cases occurs only toward the end of FFN discharge (Figure 3E,F).

Figure 3H shows the temporal relationship between the onset of Alexa dye entry and peak FFN fluorescence. Although Alexa dye entry often follows the peak of FFN fluorescence, it can precede the peak by over 500 ms. The fusion pore can thus open during the gradual increase of FFN fluorescence, well before the peak. Since slow FFN events are exocytic, the decrease of FFN fluorescence after the peak reflects release, requiring that the fusion pore open at or before the peak of FFN fluorescence. The delay in Alexa dye entry relative to peak FFN fluorescence thus suggests that cargo efflux can prevent external dye entry.

If cargo release involved only diffusion, external dye would enter exocytic DCVs instantaneously upon full membrane fusion. On the other hand, if the velocity of efflux exceeds the rate of external dye diffusion, dye cannot enter. External dye entry should only be possible

when the rate of efflux drops below the rate of dye diffusion, as cargo release progresses. In this

case, the delay in entry should be prolonged for a dye that diffuses more slowly. To test this

possibility, we examined the kinetics of entry for CF488A, a cell-impermeant dye with spectral

properties similar to Alexa fluor 488 but larger (MW ~914 Da compared to 643 Da for Alexa).

CF488A is thus expected to diffuse more slowly and experience a longer delay in dye entry.

Indeed, compared to Alexa dye entry events, a much larger fraction of CF488A dye entry events

start after peak FFN fluorescence. The extent to which dye entry events can precede peak FFN

fluorescence is also significantly reduced for CF488A events (Figure 3H). These observations are

consistent with a delay in external dye entry due to the rapid efflux of lumenal cargo through a

narrow pore.

What accounts for the gradual increase of FFN fluorescence in slow events? Since the

strength of the evanescent field increases exponentially with approach to the coverslip, the

increase in fluorescence could reflect either vesicle movement or FFN release. In either of these

cases, the fluorescence should not increase with excitation by a field of uniform strength. To test

this idea, we imaged FFN events at the lateral edge of the cell by confocal microscopy. Using

Alexa dye entry, we identified 25 events from nine cells as exocytic. Twenty of these events

showed slow FFN release, and all of these exhibited an increase in fluorescence similar to that

observed by TIRF microscopy (Figure 1—figure supplement 2D). The increase in fluorescence

for slow events thus cannot be explained by FFN movement, either within vesicles or by release.

Vesicle swelling provides an alternative explanation: Since FFN206 self-quenches at high

concentrations (Figure 1—figure supplement 1E), an increase in DCV volume would lower FFN

concentration, relieve self-quenching and thereby increase fluorescence. Indeed, both mast cell

and chromaffin granules swell upon exocytosis (Breckenridge and Almers, 1987; Terakawa et al.,

1991). Taken together, these observations suggest that the fusion pore opens at the onset of

fluorescence increase for slow as well as fast events.

Quantitative Analysis of FFN Event Kinetics Reveals Distinct Modes of Fusion

High-speed imaging with FFN206 captures events with diverse fluorescence profiles. Some exhibit a characteristic spike whereas the rest vary widely, suggesting distinct mechanisms. To distinguish between this possibility and the prevailing assumption of a single, common mechanism with stochastic variation, we characterized the kinetics of all FFN events by their full width at half-maximum (FWHM), a simple metric broadly applicable to any curve with a peak that increases monotonically before and decreases monotonically after the maximum. However, variation in event profile required different methods to calculate FWHM. Events with a spike were fit by non-linear least-squares regression, whereas those without spikes were fit using free-knot B-splines (Dung and Tjahjowidodo, 2017). FWHM was then computed from the fitted curves (Figure 4A,B). To avoid classifying individual events into discrete categories, which prevents comparison across categories, we modeled the FWHM of individual events as a continuous random variable (Figure 4C). With no *a priori* classification, maximum likelihood estimation shows that the observed data are best explained by a mixture of two lognormal distributions (Figure 4D,E). The analysis of event half-life yields similar results (Figure 4—figure supplement 1). This is inconsistent with stochastic variation as the sole source of fusion pore heterogeneity and strongly suggests two distinct mechanisms for exocytosis.

Previous work has shown that membrane tension is a primary determinant of fusion pore behavior (Amatore et al., 2000; Bretou et al., 2014). We also observed that the enzyme used to digest extracellular matrix (ECM) of the adrenal gland when isolating chromaffin cells and the substrate used for cell attachment can have profound effects on exocytosis specific to the site of attachment. Extensive removal of ECM with papain or the exclusive use of polylysine for substrate attachment can both result in tight adherence that eliminates FFN events within the TIRF footprint (data not shown). To determine whether local differences in substrate attachment account for bimodal fusion pore behavior, we again used the intensity of Alexa dye fluorescence in the TIRF field as a proxy for tightness of attachment. We found that FFN release kinetics do not correlate

1   with the local tightness of attachment at the sites of fusion (correlation coefficient −0.075, Figure

2   1—figure supplement 3E).   Thus, the tightness of substrate attachment does not account for

3   bimodal behavior of the chromaffin granule fusion pore.

5   **Differential Regulation of Fast and Slow Events by Synaptotagmin 7**

6   A number of proteins have been shown to influence the rate of fusion pore dilation

(Anantharam et al., 2011; Graham et al., 2002; Doreian et al., 2008; Berberian et al., 2009; Han

et al., 2004; Logan et al., 2017). These include the synaptotagmins, Ca^{++} sensors for regulated

exocytosis (Sudhof, 2013). Synaptotagmin (syt) 1 and 7 are the major Ca^{++} sensors in chromaffin

cells (Schonn et al., 2008), and loss of either isoform slows dilation of the chromaffin granule

fusion pore, promoting kiss-and-run exocytosis (Das et al., 2020; Lynch et al., 2008; Segovia et

al., 2010; Wang et al., 2001; Zhang et al., 2010). Syt 7 alone is responsible for ~50% of all

regulated exocytosis in chromaffin cells (Schonn et al., 2008), and accumulating evidence has

suggested that it localizes to a distinct subset of chromaffin granules, conferring slow pore dilation

(Bendahmane et al., 2018; Bendahmane et al., 2020; Jaiswal et al., 2004; Rao et al., 2014; Zhang

et al., 2011).

To determine whether the two classes of fusion pore behavior depend on different Ca^{++}

sensors, we compared FFN release kinetics in chromaffin cells from wild-type and syt 7 knockout

(KO) mice. The cumulative frequency distribution of event FWHM differs significantly between

wild-type and syt 7 KO cells ($p < 10^{-5}$ by Kolmogorov-Smirnov) with an overall slowing of FFN

release in the KO (Figure 5A), consistent with previous reports. Using maximum likelihood

estimation, we then examined the two kinetic components of fusion pore behavior. Surprisingly,

wild-type and KO show no significant difference in the kinetics of the corresponding components

(Figure 5B). However, the proportion of each component differs significantly between wild-type

and KO (Figure 5C). Wild-type cells exhibit ~70% fast events and ~30% slow, consistent with

earlier observations (Figure 4E). In contrast, fast and slow events each account for ~50% of the

events in syt 7 KO cells, and the results do not change with analysis using half-life (Figure 5—

figure supplement 1). We conclude that syt 7 alters the proportion of fast and slow events without

changing their intrinsic properties. Thus, syt 7 plays a regulatory rather than direct role in fusion

pore behavior. The two classes of fusion pore behavior respond differently to syt 7, supporting

the involvement of distinct mechanisms and indicating the potential for independent regulation.

DISCUSSION

The use of FFNs has enabled us to image DCV exocytosis at millisecond resolution. Catalytic accumulation of FFN within chromaffin granules increases the fluorescent signal, reducing the required exposure time. The small size of FFNs also makes their release kinetics a direct measure of fusion pore opening, unlike pHluorin unquenching which is delayed by dissolution of the dense core and lumenal buffering (Holz, 1978). In addition, the pH insensitivity of FFN206 enables detection of vesicle docking before fusion, in contrast to amperometry, which monitors release, and pHluorin imaging, in which the fluorophore is quenched before exocytosis. Taking advantage of these properties, we observe a range of FFN events from fast to slow, and all of these depend on stimulation, external Ca^{++} and v-SNARE. Moreover, all event types associate with the unquenching of BDNF-pHluorin and external dye entry, consistent with an exocytic mechanism.

The majority of FFN events derive from docked vesicles. However, a small number of vesicles appear to fuse immediately upon arrival at the plasma membrane, and this does not reflect fusion outside the evanescent field. Fusion within 1–2 ms of vesicle arrival indicates that tethering, docking, priming and fusion can occur almost instantaneously. Previous work has shown that synaptic vesicles can fuse ~60 ms after docking (Chen et al., 2013) but FFN imaging now further constrains the time needed even by DCVs, which have been thought to require more time for fusion (Steyer et al., 1997). We also observe very few vesicles arriving at the plasma membrane that do not undergo fusion, consistent with a large population of stably docked granules.

Although all the observed events are exocytic, the time of pore opening in slow events remains unclear because lumenal buffering delays BDNF-pHluorin unquenching and cargo efflux delays external dye entry. How does cargo efflux delay dye entry? Previous work has shown that osmotic forces drive catecholamine efflux (Troyer and Wightman, 2002). The mechanism remains unclear but presumably involves an increase in intralumenal pressure produced by an osmotic

gradient across the vesicle membrane (Holz, 1986; Sudhof, 1982). The increased pressure would

result in high-velocity outflow through a narrow fusion pore, and we hypothesize that this velocity

exceeds the rate of inward Alexa dye diffusion, which may also be impeded by the ECM (see

Supplementary Text): the method we use to prepare mouse chromaffin cells primarily digests

collagen, leaving other ECM components largely intact. As efflux slows due to release of soluble

cargo and dissipation of the osmotic gradient that drives water into the DCV, the velocity falls

below the rate of external dye diffusion, thus enabling dye entry. Consistent with this, the entry

of CF488A, a larger dye with slower diffusion, shows a longer delay. Previously, a delay in external

dye entry was interpreted as evidence for a stable, hemifused state preceding full fusion (Chiang

et al., 2014; Zhao et al., 2016). Our results suggest that this may instead reflect slow cargo efflux

through a narrow pore.

When does the fusion pore open? If chromaffin granules swell upon exocytosis and

membrane fusion precedes swelling, as reported for mast cell as well as chromaffin granules

(Breckenridge and Almers, 1987; Terakawa et al., 1991), then swelling would account for the

increase in FFN fluorescence observed by confocal as well as TIRF microscopy and hence

independent of the evanescent field. In this case, membrane fusion occurs no later than the onset

of FFN fluorescence increase, well before Alexa dye entry.

The ability to monitor the duration of all exocytic events makes the full distribution of fusion

pore behavior amenable to quantitative methods. Previous work assigning exocytic events to

different classes based on the presence or absence of certain features (Chiang et al., 2014; Logan

et al., 2017; van Kempen et al., 2011) made it difficult to determine the relationship between

classes and assess the mechanistic basis for differences between them. In this work, we use a

simple kinetic parameter applicable to all events, FWHM. By treating event FWHM as a

continuous random variable, the analysis accounts for the inherent stochasticity of biological

systems while avoiding the *a priori* classification of individual events. The analysis of FWHM

distribution reveals two lognormal components, indicating two distinct modes of release that are

inconsistent with stochastic variation in a common pathway.

What accounts for the difference between the two classes of release event? A possibility

is that one of them does not involve a chromaffin granule. However, exocytosis of a peptide

reporter accompanies both event classes, indicating that both involve DCVs. The local tightness

of attachment also appears to have no role in the variability of event kinetics. Loss of syt 7 reduces

the average rate of release, in agreement with some previous studies (Li et al., 2009; Segovia et

al., 2010; Tsuboi and Fukuda, 2007; Zhang et al., 2010) but not all (Bendahmane et al., 2018;

Bendahmane et al., 2020; Rao et al., 2014; Zhang et al., 2010; Zhang et al., 2011). This has

generally been interpreted as evidence of a direct role for syt 7 in pore dilation as well as

membrane fusion (Figure 5D). However, we find that loss of syt 7 slows overall release by

increasing the proportion of DCVs that undergo exocytosis with small pores, suggesting a role in

the regulation of exocytosis rather than direct participation in the fusion pore. Differences in fusion

pore behavior may reflect the exocytosis of different DCVs. Recent work has indeed suggested

the segregation of syt 1 and 7 to chromaffin granules of different size (Rao et al., 2014; Zhang et

al., 2011). Syt 7 may thus play a role upstream of exocytosis, in DCV biogenesis or targeting to

the release site, producing vesicles that undergo fast release (Figure 5E, green arrow).

Alternatively, syt 7 may act during exocytosis to promote specifically the fusion of DCVs destined

for fast release, a role consistent with its known function as a calcium sensor for exocytosis

(Figure 5E, red arrow). It will now be important to identify the factors that confer the two modes of

release and to understand their differential regulation by synaptotagmin and other proteins. The

analysis of biological variation presented here thus provides a new perspective on the function of

syt 7 and should also serve to guide future work on the molecular mechanisms that underlie the

bimodal nature of regulated DCV exocytosis.

Slow events originate only from docked vesicles whereas new arrivals release with

invariably fast kinetics. Stable docking must therefore confer the potential for a long-lived small

fusion pore. Docked vesicles release with both fast and slow kinetics, suggesting different states

of docking or priming. We do not know whether the divergent behavior reflects different vesicles,

fusion sites or cytosolic machinery.

The demonstration of two distinct exocytic mechanisms suggests a biological role for the

diversity of DCV fusion pore behavior which has important implications for the release of peptides,

monoamines and other neuromodulators. A global change due to regulation of a common exocytic

pathway would in principle affect all release events, making it difficult to regulate signaling

independently over different time scales. In contrast, two modes of release enable differential

control of temporally distinct signals. The fast events associated with rapid pore dilation release

more cargo acutely. A narrow pore slows transmitter efflux and also enables the trapping of

residual peptide cargo for subsequent release. Bimodal fusion pore kinetics thus confer the

potential for independent regulation of fast versus slow release, adding a new dimension to the

temporal control of signaling. The defect in fusion pore dilation that underlies impaired insulin

release in type 2 diabetes (Collins et al., 2016) indicates the potential for subversion of this

mechanism in disease.

ACKNOWLEDGMENTS

This work was supported by grants from the NIH (R01 NS109295) and the Michael J. Fox

Foundation to R.H.E. We thank Todd Logan for developing the use of FFNs in chromaffin cells

and members of the Edwards lab for helpful discussion and suggestions. We thank Wade Regehr

for sending the synaptotagmin 7 knockout mice. We thank DeLaine Larson, Kari Herrington and

SoYeon Kim at the UCSF Center for Advanced Light Microscopy for technical assistance and

advice. The algorithm for Dual-View calibration was generously provided by Dr. Damien Jullié.

AUTHOR CONTRIBUTIONS

Conceptualization: R.H.E.; Methodology: P.Z., R.H.E.; Software: P.Z.; Validation: P.Z.; Formal

Analysis: P.Z., D.R.; Investigation: P.Z.; Resources: R.H.E.; Data Curation: P.Z.; Writing –

Original Draft: R.H.E.; Writing – Review & Editing: P.Z., D.R. and R.H.E.; Visualization: P.Z. and

R.H.E.; Supervision: R.H.E.; Funding Acquisition: R.H.E.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Alabi, A.A., and Tsien, R.W. (2013). Perspectives on kiss-and-run: role in exocytosis, endocytosis, and neurotransmission. Annu Rev Physiol *75*, 393–422.

Albillos, A., Dernick, G., Horstmann, H., Almers, W., Alvarez de Toledo, G., and Lindau, M. (1997). The exocytotic event in chromaffin cells revealed by patch amperometry. Nature *389*, 509–512.

Amatore, C., Bouret, Y., Travis, E.R., and Wightman, R.M. (2000). Adrenaline release by chromaffin cells: constrained swelling of the vesicle matrix leads to full fusion at the ENS. Angew Chem Int Ed Engl *39*, 1952–1955.

Bao, H., Das, D., Courtney, N.A., Jiang, Y., Briguglio, J.S., Lou, X., Roston, D., Cui, Q., Chanda, B., and Chapman, E.R. (2018). Dynamics and number of trans-SNARE complexes determine nascent fusion pore properties. Nature *554*, 260–263.

Bendahmane, M., Bohannon, K.P., Bradberry, M.M., Rao, T.C., Schmidtke, M.W., Abbineni, P.S., Chon, N.L., Tran, S., Lin, H., Chapman, E.R.*, et al.* (2018). The synaptotagmin C2B domain calcium-binding loops modulate the rate of fusion pore expansion. Mol Biol Cell *29*, 834–845.

Bendahmane, M., Morales, A., Kreutzberger, A.J.B., Schenk, N.A., Mohan, R., Bakshi, S., Philippe, J.M., Zhang, S., Kiessling, V., Tamm, L.K.*, et al.* (2020). Synaptotagmin-7 enhances calcium-sensing of chromaffin cell granules and slows discharge of granule cargos. J Neurochem.

Breckenridge, L.J., and Almers, W. (1987). Final steps in exocytosis observed in a cell with giant secretory granules. Proc Natl Acad Sci USA *84*, 1945–1949.

Bretou, M., Jouannot, O., Fanget, I., Pierobon, P., Larochette, N., Gestraud, P., Guillon, M., Emiliani, V., Gasman, S., Desnos, C.*, et al.* (2014). Cdc42 controls the dilation of the exocytotic fusion pore by regulating membrane tension. Mol Biol Cell *25*, 3195–3209.

Chen, M., Van Hook, M.J., Zenisek, D., and Thoreson, W.B. (2013). Properties of ribbon and non-ribbon release from rod photoreceptors revealed by visualizing individual synaptic vesicles. J Neurosci *33*, 2071–2086.

Chiang, H.C., Shin, W., Zhao, W.D., Hamid, E., Sheng, J., Baydyuk, M., Wen, P.J., Jin, A., Momboisse, F., and Wu, L.G. (2014). Post-fusion structural changes and their roles in exocytosis and endocytosis of dense-core vesicles. Nat Commun *5*, 3356.

Chow, R.H., von Rüden, L., and Neher, E. (1992). Delay in vesicle fusion revealed by
electrochemical monitoring of single secretory events in adrenal chromaffin cells. Nature *356*, 60–
63.

Collins, S.C., Do, H.W., Hastoy, B., Hugill, A., Adam, J., Chibalina, M.V., Galvanovskis, J.,
Godazgar, M., Lee, S., Goldsworthy, M*., et al.* (2016). Increased Expression of the Diabetes Gene
SOX4 Reduces Insulin Secretion by Impaired Fusion Pore Expansion. Diabetes *65*, 1952–1961.

Das, D., Bao, H., Courtney, K.C., Wu, L., and Chapman, E.R. (2020). Resolving kinetic
intermediates during the regulated assembly and disassembly of fusion pores. Nat Commun *11*,
231.

Dung, V.T., and Tjahjowidodo, T. (2017). A direct method to solve optimal knots of B-spline
curves: An application for non-uniform B-spline curves fitting. PLoS One *12*, e0173857.

Fulop, T., Radabaugh, S., and Smith, C. (2005). Activity-dependent differential transmitter release
in mouse adrenal chromaffin cells. J Neurosci *25*, 7324–7332.

Gubernator, N.G., Zhang, H., Staal, R.G., Mosharov, E.V., Pereira, D.B., Yue, M., Balsanek, V.,
Vadola, P.A., Mukherjee, B., Edwards, R.H*., et al.* (2009). Fluorescent false neurotransmitters
visualize dopamine release from individual presynaptic terminals. Science *324*, 1441–1444.

Holz, R.W. (1978). Evidence that catecholamine transport into chromaffin vesicles is coupled to
vesicle membrane potential. Proc Natl Acad Sci USA *75*, 5190–5194.

Holz, R.W. (1986). The role of osmotic forces in exocytosis from adrenal chromaffin cells. Annu
Rev Physiol *48*, 175–189.

Hu, G., Henke, A., Karpowicz, R.J., Jr., Sonders, M.S., Farrimond, F., Edwards, R., Sulzer, D.,
and Sames, D. (2013). New fluorescent substrate enables quantitative and high-throughput
examination of vesicular monoamine transporter 2 (VMAT2). ACS Chem Biol *8*, 1947–1954.

Jahn, R., and Fasshauer, D. (2012). Molecular machines governing exocytosis of synaptic
vesicles. Nature *490,* 201–207.

Jaiswal, J.K., Chakrabarti, S., Andrews, N.W., and Simon, S.M. (2004). Synaptotagmin VII
restricts fusion pore expansion during lysosomal exocytosis. PLoS Biol *2*, e233.

Kaeser, P.S., and Regehr, W.G. (2014). Molecular mechanisms for synchronous, asynchronous,
and spontaneous neurotransmitter release. Ann Rev Physiol *76*, 333–363.

Li, J., Xiao, Y., Zhou, W., Wu, Z., Zhang, R., and Xu, T. (2009). Silence of Synaptotagmin VII inhibits release of dense core vesicles in PC12 cells. Sci China C Life Sci *52*, 1156–1163.

Logan, T., Bendor, J., Toupin, C., Thorn, K., and Edwards, R.H. (2017). alpha-Synuclein promotes dilation of the exocytotic fusion pore. Nat Neurosci *20*, 681–689.

Lynch, K.L., Gerona, R.R., Kielar, D.M., Martens, S., McMahon, H.T., and Martin, T.F. (2008). Synaptotagmin-1 utilizes membrane bending and SNARE binding to drive fusion pore expansion. Mol Biol Cell *19*, 5093–5103.

McMahon, H.T., Ushkaryov, Y.A., Edelmann, L., Link, E., Binz, T., Niemann, H., Jahn, R., and Sudhof, T.C. (1993). Cellubrevin is a ubiquitous tetanus-toxin substrate homologous to a putative synaptic vesicle fusion protein. Nature *364*, 346–349.

Miesenböck, G., De Angelis, D.A., and Rothman, J.E. (1998). Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. Nature *394*, 192–195.

Nenadic, Z., and Burdick, J.W. (2005). Spike detection using the continuous wavelet transform. IEEE Trans Biomed Eng *52*, 74–87.

Newville, M., Stensitzki, T., Allen, D.B., Rawlik, M., Ingargiola, A., and Nelson, A. (2016). LMFIT: non-linear least-square minimization and curve-fitting for Python. Astrophysics Source Code Library.

Parsons, T.D., Coorssen, J.R., Horstmann, H., and Almers, W. (1995). Docked granules, the exocytic burst, and the need for ATP hydrolysis in endocrine cells. Neuron *15*, 1085–1096.

Perrais, D., Kleppe, I.C., Taraska, J.W., and Almers, W. (2004). Recapture after exocytosis causes differential retention of protein in granules of bovine chromaffin cells. J Physiol *560*, 413–428.

Rao, T.C., Passmore, D.R., Peleman, A.R., Das, M., Chapman, E.R., and Anantharam, A. (2014). Distinct fusion properties of synaptotagmin-1 and synaptotagmin-7 bearing dense core granules. Mol Biol Cell *25*, 2416–2427.

Rorsman, P., and Renstrom, E. (2003). Insulin granule dynamics in pancreatic beta cells. Diabetologia *46*, 1029–1045.

Schiavo, G., Benfenati, F., Poulain, B., Rossetto, O., Polverino de Laureto, P., DasGupta, B.R., and Montecucco, C. (1992). Tetanus and botulinum-B neurotoxins block neurotransmitter release by proteolytic cleavage of synaptobrevin. Nature *359*, 832–835.

Schonn, J.S., Maximov, A., Lao, Y., Sudhof, T.C., and Sorensen, J.B. (2008). Synaptotagmin-1 and -7 are functionally overlapping Ca2+ sensors for exocytosis in adrenal chromaffin cells. Proc Natl Acad Sci USA *105*, 3998–4003.

Segovia, M., Ales, E., Montes, M.A., Bonifas, I., Jemal, I., Lindau, M., Maximov, A., Sudhof, T.C., and Alvarez de Toledo, G. (2010). Push-and-pull regulation of the fusion pore by synaptotagmin-7. Proc Natl Acad Sci USA *107*, 19032–19037.

Shin, W., Ge, L., Arpino, G., Villarreal, S.A., Hamid, E., Liu, H., Zhao, W.D., Wen, P.J., Chiang, H.C., and Wu, L.G. (2018). Visualization of membrane pore in live cells Reveals a dynamic-pore theory governing fusion and endocytosis. Cell *173*, 934–945.e12.

Steyer, J.A., Horstmann, H., and Almers, W. (1997). Transport, docking and exocytosis of single secretory granules in live chromaffin cells. Nature *388*, 474–478.

Sudhof, T.C. (1982). Core structure, internal osmotic pressure and irreversible structural changes of chromaffin granules during osmometer behaviour. Biochim Biophys Acta *684*, 27–39.

Sudhof, T.C. (2013). Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. Neuron *80*, 675–690.

Taylor, M.J., Perrais, D., and Merrifield, C.J. (2011). A high precision survey of the molecular dynamics of mammalian clathrin-mediated endocytosis. PLoS Biol *9*, e1000604.

Terakawa, S., Fan, J.H., Kumakura, K., and Ohara-Imaizumi, M. (1991). Quantitative analysis of exocytosis directly visualized in living chromaffin cells. Neurosci Lett *123*, 82–86.

Troyer, K.P., and Wightman, R.M. (2002). Temporal separation of vesicle release from vesicle fusion during exocytosis. J Biol Chem *277*, 29101–29107.

Tsuboi, T., and Fukuda, M. (2007). Synaptotagmin VII modulates the kinetics of dense-core vesicle exocytosis in PC12 cells. Genes Cells *12*, 511–519.

van Kempen, G.T., vanderLeest, H.T., van den Berg, R.J., Eilers, P., and Westerink, R.H. (2011). Three distinct modes of exocytosis revealed by amperometry in neuroendocrine cells. Biophys J *100*, 968–977.

Voets, T., Neher, E., and Moser, T. (1999). Mechanisms underlying phasic and sustained secretion in chromaffin cells from mouse adrenal slices. Neuron *23*, 607–615.

Wang, C.T., Grishanin, R., Earles, C.A., Chang, P.Y., Martin, T.F., Chapman, E.R., and Jackson,

M.B. (2001). Synaptotagmin modulation of fusion pore kinetics in regulated exocytosis of dense-

core vesicles. Science *294*, 1111–1115.

Xia, X., Lessmann, V., and Martin, T.F. (2009). Imaging of evoked dense-core-vesicle exocytosis

in hippocampal neurons reveals long latencies and kiss-and-run fusion events. J Cell Sci *122*,

75–82.

Zhang, Z., Hui, E., Chapman, E.R., and Jackson, M.B. (2010). Regulation of exocytosis and fusion

pores by synaptotagmin-effector interactions. Mol Biol Cell *21*, 2821–2831.

Zhang, Z., Wu, Y., Wang, Z., Dunning, F.M., Rehfuss, J., Ramanan, D., Chapman, E.R., and

Jackson, M.B. (2011). Release mode of large and small dense-core vesicles specified by different

synaptotagmin isoforms in PC12 cells. Mol Biol Cell *22*, 2324–2336.

Zhao, W.D., Hamid, E., Shin, W., Wen, P.J., Krystofiak, E.S., Villarreal, S.A., Chiang, H.C.,

Kachar, B., and Wu, L.G. (2016). Hemi-fused structure mediates and controls fusion and fission

in live cells. Nature *534*, 548–552.

Zhou, Z., Misler, S., and Chow, R.H. (1996). Rapid fluctuations in transmitter release from single

vesicles in bovine adrenal chromaffin cells. Biophys J *70*, 1543–1552.

1 **FIGURES**

3 **Figure 1.** High frequency imaging reveals distinct classes of exocytic event. (**A**) Micrographs of

4 an isolated primary mouse adrenal chromaffin cell, illuminated by bright field (left) or total internal

5 reflection, revealing FFN puncta (right). Scale bar: 5 µm. (**B–D**) Fluorescence time traces showing

fast (**B**) and slow (**C**) release of FFNs with reduced post-event baseline (indicating docked

vesicles), as well as rapid release with unchanged baseline (new arrivals) (**D**). Insets in each

panel show select frames of the corresponding event, marked by the time relative to initial FFN

signal increase (time 0).

Figure 1—figure supplement 1. FFN properties.

Figure 1—figure supplement 2. Sample fluorescence time traces of FFN release events.

Figure 1—figure supplement 3. Tightness of cell attachment does not affect FFN release

kinetics.

Figure 1—video 1. High-speed imaging of FFN release from an isolated primary mouse adrenal

chromaffin cell in culture.

Figure 1—video 2. Fast FFN release from a docked vesicle.

Figure 1—video 3. Fast FFN release from a new arrival.

Figure 1—video 4. Slow FFN release from a docked vesicle.

Figure 1—video 5. Dual imaging of FFN release and BDNF-pHluorin exocytosis.

2   **Figure 2.** FFN release is exocytic in nature. (**A**) Removal of Ca^{++} from the external solution

3   abolishes FFN release from chromaffin cells. (**B**) Lentivral transduction of tetanus toxin light

4   chain (TeTX-LC) abolishes FFN release. Error bars indicate standard deviation. (**C**) Quantitative

Venn diagram shows the vast majority of FFN release events coincide with unquenching of

BDNF-pHluorin. (**D–G**) Fluorescence time traces show that unquenching of BDNF-pHluorin

(green) follows FFN release (blue) for both fast (**D** and **F**) and slow events (**E**) with reduced

post-event baseline, as well as for release with an unchanged post-event baseline (**G**). The

prolonged exposure time required for dual imaging of BDNF-pHluorin and FFN limited capture

of the FFN event to a single (**D** and **G**) or no (**F**) frame. (**H**) Kymographs showing the spread of

FFN following rapid release.

2 **Figure 3.** FFN and alexa fluor 488 dual imaging. (**A–F**) Fluorescence time traces showing Alexa

3 (green) dye entry occurs simultaneously with a fast FFN event (blue) (**A**) but during (**B**) or after a

4 slow FFN event (**C–F**). (**G**) Quantitative Venn diagram showing that the vast majority of FFN

1   events are accompanied by Alexa dye entry. (**H**) Onset of external dye entry relative to peak FFN

2   fluorescence. Negative delay means external dye entry occurred before FFN reached peak

3   fluorescence.

2   **Figure 4.** Quantitative analysis of FFN events. (**A**) Events with a spike were fit by least-squares

3   regression. (**B**) Events without spikes were fit using free-knot B-splines. Event kinetics are

4   characterized using the full width at half-maximum (FWHM), defined as the time difference

5   between the half-rise and half-decay points (green crosses). (**C**) The observed cumulative

6   frequency of FWHM and the cumulative probability function determined by fitting the data using

1   maximum likelihood estimation (MLE). Inset: P-P plot showing the goodness of fit. (**D**) The

2   probability density functions of the two lognormal distributions used to fit the data in **C**. (**E**) Relative

3   contribution of the two components shown in **D**. Error bars: 95% confidence interval.

4   **Figure 4—figure supplement 1.** Characterizing FFN release kinetics using full width at half

5   maximum.

2 **Figure 5.** Synaptotagmin 7 knockout increases the proportion of slow events without affecting

3 their kinetics. (**A**) Cumulative frequencies and MLE best fit of event full width half maximum

4 (FWHM) in wild-type (left) and syt 7 knockout (right) cells. Insets show the respective P-P plots.

5 (**B**) Loss of syt 7 does not alter the properties of individual kinetic components. Error bars

6 represent the 2.5% to 97.5% quantile region of each distribution, calculated using best-fit

parameters. (**C**) Loss of syt 7 significantly reduces the relative contribution of the fast component

to the total. Error bars indicate 95% confidence interval. (**D**) Traditional view that syt 7 mediates

both membrane fusion and pore expansion. (**E**) The results suggest two distinct pathways to

exocytosis, leading to fast (solid black arrows) or slow release (dashed arrows). Syt 7 may either

act to promote the targeting of vesicles for fast release (green arrow), or to increase specifically

the fusion probability of vesicles forming large pores (red arrow).

Figure 5—figure supplement 1. The effects of synaptotagmin 7 knockout measured using full

width at half maximum.

METHODS

Primary chromaffin cell culture. Primary mouse adrenal chromaffin cells were isolated as

previously described (Logan et al., 2017), with slight modification. Briefly, adrenal glands were

harvested from animals 4–6 weeks old and placed in ice cold Ca^{++}-, Mg^{++}-free (CMF) Hank's

balanced salt solution (HBSS, UCSF cell culture facility [CCF] media production) supplemented

with 10 U ml^{-1} penicillin G and 10 μg ml^{-1} streptomycin (pen-strep, UCSF CCF media production).

Adrenal medullae were isolated and digested in CMF-HBSS containing collagenase type I (2.9

mg ml^{-1}, Worthington), bovine serum albumin (BSA) (3 mg ml^{-1}, Sigma), DNase I (0.15 mg ml^{-1},

Sigma), penicillin G (10 U ml^{-1}), and streptomycin (10 μg ml^{-1}) at 37° C for 40 min with shaking

at 800 rpm in a thermomixer (Eppendorf). Digested medullae were resuspended and the

enzymatic reaction quenched by addition of 10 volumes cold CMF-HBSS. Isolated chromaffin

cells were collected by centrifugation at 300 g for 10 min at 4° C and resuspended in pre-warmed

DMEM / Ham's F-12 1:1 mix (UCSF CCF media production) supplemented with 10% FBS

(HyClone), penicillin G (4 U ml^{-1}), and streptomycin (4 μg ml^{-1}). Cell suspensions were plated

drop-wise onto chambered coverglass (μ-Slide, Ibidi) coated with 100 μg ml^{-1} collagen (type I,

Corning), 50 μg ml^{-1} fibronectin (Innovative Research), and 1 μg ml^{-1} BSA, and incubated at 37°

C / 5% CO_2 for 45 min before addition of fresh culture media or lentiviral supernatant for overnight

incubation. Medium was replaced the next morning.

Lentiviral production. Low passage Lenti-X 293T cells (Clontech) were seeded onto 6-well

plates coated with 0.1 mg ml^{-1} poly-L-lysine and transfected using FuGENE HD transfection

reagent (Promega) with a mixture of third-generation lentiviral backbone vector pJHUMCS that

encodes BDNF-pHluorin, as well as accessory plasmids pVSV-G, pREV, and pRRE. Cells were

switched into chromaffin cell culture medium the next day. The culture medium was collected 48

h after transfection and cell debris sedimented by centrifugation at 1,000 g for 10 min at RT. The

resulting lentiviral supernatant was either used immediately or stored at −80° C.

Fluorescent false neurotransmitter (FFN) loading. Chromaffin cells were loaded with FFNs

immediately before imaging. The culture medium was replaced with DMEM high glucose (UCSF

CCF media production) containing 10 µM FFN (Abcam) filtered through a 0.2 µ filter (FFN loading

buffer). Cells were incubated in FFN loading buffer at 37° C / 5% CO_2 for 1 h before switching into

modified Tyrode's solution (in mM, 140 NaCl, 10 HEPES-NaOH, pH 7.4, 10 glucose, 4.5 KCl, 5

$CaCl_2$, 1 $MgCl_2$) for imaging. Exocytosis was evoked by adding an equal volume of stimulation

buffer (in mM, 54.5 NaCl, 10 HEPES-NaOH, pH 7.4, 10 glucose, 90 KCl, 5 $CaCl_2$, 1 $MgCl_2$). For

experiments using Alexa dyes, 60 µM Alexa Fluor 488 carboxylic acid, tris(triethylammonium) salt

(Thermo Fisher) was included in modified Tyrode's solution and stimulation buffer.

Microscopy. Chromaffin cells were imaged at room temperature using an inverted microscope

(Ti-E, Nikon). TIRF microscopy was done using a MLC400 monolithic laser combiner (Agilent

Technologies), ZT405/488/561/640rpc dichroic filter (Chroma), and Plan Apo Lambda 100✕ 1.45

N.A. oil objective (Nikon). Rapid FFN imaging was done using ET455/50m emission filter

(Chroma) and Orca Flash 4.0 C11440-22C sCMOS camera (Hamamatsu). Dual imaging of FFN

and BDNF-pHluorin was done using ZET488/561/635m emission filter (Chroma) and iXon DU-

897 EMCCD camera (ANDOR). Dual imaging of FFN with Alexa Fluor 488 (ThermoFisher) or

CF488A (Biotium) was done using Dual-View filter cube (Optical Insights) mounted with

T510lpxrxt beam splitter (Chroma), ET460/36m and ET545/40m emission filters (Chroma), and

ORCA Flash 4.0 C11440-22C sCMOS camera (Hamamatsu). Confocal imaging of FFN and Alexa

Fluor 488 was done using an integrated laser engine (ANDOR), Borealis CSU-W1 spinning disk

(ANDOR), ZT405/488/561/640rpcv2 dichroic filter (Chroma), Plan Apo VC 100✕ 1.4 N.A. oil

objective (Nikon), ZET405/488/561/635 emission filter (Chroma), and iXon DU-888 EMCCD

camera (ANDOR). Cells were imaged in modified Tyrode's solution and exocytosis was stimulated

with 45 mM K$^+$ by adding an equal volume of stimulation buffer (in mM, 54.5 NaCl, 10 HEPES-

NaOH, pH 7.4, 10 glucose, 90 KCl, 5 CaCl$_2$, 1 MgCl$_2$).

Dual-View calibration. Images acquired using the Dual-View (Optical Insights) beam-splitter

were calibrated to minimize distortion between the two channels. The optical configurations of the

beam-splitter were mechanically calibrated using TetraSpeck microspheres (0.1 µm,

ThermoFisher) that fluoresce in both channels. To correct for the remaining small shifts, images

were aligned using a modified version of a previously described algorithm (Taylor et al., 2011).

Briefly, before each imaging session, a calibration image of TetraSpeck microspheres was

captured. At least eight beads that appear in both channels were manually selected and the

coordinates of their respective centers refined by 2-D Gaussian fit. A third-degree polynomial

spatial transform was then used to align the corresponding centers of each bead in the two

channels and the coefficients were calculated by interpolation. The calculated coefficients were

then used for a third-degree polynomial spatial transform of the Alexa channel for the images

collected in the subsequent imaging session.

Data analysis. Individual exocytotic events were identified manually in ImageJ by placing 5×5-

pixel regions of interest (ROIs) over the center of events. Average intensity profiles were extracted

using the Time Series Analyzer plugin (V3) and further processed using Python. Peak detection

was done using continuous wavelet transform (Nenadic and Burdick, 2005) and verified by

manual inspection. Pre- and post-event baselines were estimated by analyzing consecutive 20-

frame sliding windows for statistical equivalency, and refined by manual inspection. Events for

which both pre- and post-event baselines were identified underwent further processing. The null

hypothesis that the fluorescence of the pre- and post-event baselines are equivalent was tested

using the two-tailed t-test, treating each imaging frame as an independent observation. If the null

hypothesis was rejected, the vesicle was categorized as docked, otherwise as a new arrival. If a

peak was detected, the trace was fitted to six models (linear / exponential rise and linear / exponential / two-phase exponential decay) by non-linear least-squares minimization using the lmfit package (Newville et al., 2016), and the best-fit model chosen using the Akaike information criterion corrected for limited sample size (AICc). If no peak was detected, the trace was fitted using free-knot non-uniform B-splines, using a direct method to optimize knot placement (Dung and Tjahjowidodo, 2017). Goodness of fit was evaluated by testing the normality of the residuals and verified by manual inspection. For some events that were not well fitted by either approach, the trace was smoothed by taking the 11-frame walking average and the smoothed curve was interpolated using a cubic spline. Interpolation results were also verified by manual inspection. From the processed curves, the peak time point as well as the time points at which the trace reaches the mid-point of fluorescence increase (half-rise) and the mid-point of fluorescence decay (half-decay) were determined. The half-life is defined as the time from peak to half-decay, whereas the full-width at half-maximum (FWHM) was defined as the time from half-rise to half-decay. The cumulative frequency distribution of the half-life or FWHM was then obtained and fitted to a log-normal or two-component log-normal mixture distribution by maximum likelihood estimation. The best-fit model was selected by performing a likelihood-ratio test. The confidence intervals of the best-fit parameters were determined by finding the 95% likelihood ratio confidence interval. The mean and 95% confidence interval of each kinetic component were computed from the best-fit parameters.

Evaluating cell attachment. The TIRF footprint of individual chromaffin cells, visualized by Alexa Fluor 488 fluorescence, was subtracted from the image of an empty field of view. Relative attachment was then normalized to the maximum pixel value of the resulting image, with 0.0 representing the loosest attachment (beyond the evanescent field) and 1.0 representing the tightest attachment (directly against the cover glass).

1   **SUPPLEMENTAL MATERIAL**

2   **SUPPLEMENTAL FIGURES**

4   **Figure 1—figure supplement 1. FFN properties.**

5   (**A–D**) pH sensitivity of FFN206 and FFN202. *In vitro* (**A** and **B**), FFN fluorescence was measured

6   at 10 μM in pH 5.5 (dashed lines) and 7.4 (solid lines). Low pH does not affect FFN206

fluorescence (**A**), whereas FFN202 is quenched at pH 5.5 (**B**). In cells (**C** and **D**), FFN-loaded vesicles were unquenched by addition of NH_4Cl. Cells loaded with FFN206 show much less unquenching (**C**) than cells loaded with FFN202 (**D**). (**E** and **F**) FFN206 exhibits self-quenching at higher concentrations. Measured *in vitro* at pH 7.4, FFN206 peak emission at 465 nm does not increase linearly with concentration (**E**). The emission spectrum does not change with concentration (**F**). (**G**) TIRF micrographs of an isolated primary chromaffin cell loaded with FFN206. Distinct FFN puncta are visible before addition of NH_4Cl (left). Dissipation of the pH gradient with NH_4Cl (50 mM) results in dispersion of the punctate fluorescence and an increase in overall fluorescence (right) presumably due to FFN206 leakage from vesicles and less self-quenching.

Figure 1—figure supplement 2. Sample fluorescence time traces of FFN release events.

(**A–C**) FFN release events imaged by TIRF microscopy. Release events are broadly divided into

three categories: rapid release with reduced post-event baseline (**A**), slow release with reduced

post-event baseline (**B**), and rapid release with unchanged post-event baseline (**C**). Note the

variation in event duration and profile among slow events. (**D**) Slow FFN release and Alexa dye

entry imaged by confocal microscopy. Cells were loaded with FFN and bathed in imaging buffer

containing cell-impermeant Alexa fluor 488. FFN release and Alexa dye entry were monitored by

dual imaging using confocal microscopy. The confocal plane was set above the glass coverslip

to image exocytic events that do not release toward the objective. FFN signal increases slowly

before declining, similar to slow release events imaged by TIRF microscopy (**B**).

Figure 1—figure supplement 3. Tightness of cell attachment does not affect FFN release

kinetics.

(**A** and **B**) TIRF micrographs of an isolated primary mouse adrenal chromaffin cell overlaid with

ROIs marking sites of FFN release, showing FFN (**A**) and Alexa Fluor 488 (**B**). The image in **A**

shows the average of 10 frames, which is how events were identified (see Methods). Sites of FFN

release with unchanged post-event baseline are indicated in red. The variation in background

Alexa 488 fluorescence occurs over a larger spatial dimension likely reflecting unevenness of the

glass coverslip. The brighter area to the upper right was observed in the same position relative

to all cells and thus reflects refraction of the incident light. Scale bar: 2 µm. (**C**) Cumulative

frequency distribution of relative attachment tightness at sites of FFN release (292 events from

21 cells). A relative tightness of 1.0 represents the tightest attachment (to the cover glass)

whereas a value of 0.0 represents the loosest attachment (beyond the evanescent field). (**D**)

Comparison of relative attachment tightness for sites of new arrivals vs. docked vesicles. p =

0.7953 by two-tailed Mann-Whitney test. Error bars: standard deviation. (**E**) Comparison of peak

amplitude for new arrivals vs. docked vesicles. p = 0.0286 by two-tailed Mann-Whitney test. Error

bars: standard deviation. (**F**) Scatter plot showing the relationship between peak amplitude and

FFN release kinetics. (**G**) Scatter plot showing the relation between relative attachment tightness

and FFN release kinetics. Covariance analysis yields a correlation coefficient of −0.008.

2   **Figure 4—figure supplement 1. Characterizing FFN release kinetics using half-life.**

3   (**A** and **B**) The same FFN release data from Figure 3 was alternatively characterized using the

4   half-life, defined as the time difference between peak and half-decay (green crosses). (**C**) The

5   observed cumulative frequency of half-lives for 257 events from 12 cells in an experimental group,

6   and the cumulative probability function determined by fitting the data using maximum likelihood

1   estimation (MLE). Inset: P-P plot depicting the goodness of fit. (**D**) The probability density

2   functions of the two lognormal distributions used to fit the data in (**C**). (**E**) Relative contribution of

3   the two components shown in (D). Error bars, 95 % confidence interval.

Figure 5—figure supplement 1. The effects of synaptotagmin 7 knockout measured using

half-life.

(**A**) Cumulative frequencies and MLE best fit of event half-lives in wild-type (left, 303 events

from 4 cells) and syt 7 knockout (right, 157 events from 10 cells). Insets show the respective P-

P plots depicting the goodness of fit. (**B**) Loss of syt 7 does not alter the properties of individual

kinetic components. Error bars represent the 2.5 % to 97.5 % quantile region of each

distribution, calculated using best-fit parameters. (**C**) Loss of syt 7 significantly reduces the

relative contribution of the fast component to the total. Error bars indicate 95 % confidence

interval.

Figure 1—video 1. High-speed imaging of FFN release from an isolated primary mouse

adrenal chromaffin cell in culture.

After loading with FFN (10 µM) for one hour, the cell was imaged at rest for ~5 s before stimulation

with 45 mM K$^+$. Imaging was performed at 800 Hz.

Figure 1—video 2. Fast FFN release from a docked vesicle.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as

described for Video S1.

Figure 1—video 3. Fast FFN release from a new arrival.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as

described for Video S1.

Figure 1—video 4. Slow FFN release from a docked vesicle.

An example event is marked by the arrow. The cell was loaded, stimulated and imaged as

described for Video S1.

Figure 1—video 5. Dual imaging of FFN release and BDNF-pHluorin exocytosis.

The cell was loaded as described for Video S1 and imaged at rest for ~10 s before stimulation

with 45 mM K$^+$. The FFN channel is shown on the left and the BDNF-pHluorin channel on the

right. The arrowhead marks a fast FFN release event in both channels. An arrow similarly marks

a slow FFN event. (The slow event will be better visualized by increasing the frame rate.)