Maize AFP1 confers antifungal activity by inhibiting chitin deacetylases from a broad

range of fungi

Lay-Sun Ma[1]*, Wei-Lun Tsai[1], Raviraj M. Kalunke[1,2], Meng-Yun Xu[1], Yu-Han Lin[1],

Florensia Ariani Damei[1], and Hui-Chun Lee[1]

[1]Institute of Plant and Microbial Biology, Academia Sinica, Taipei 11529, Taiwan.

[2]Present address: Donald Danforth Plant Science Center, 975 N Warson Rd, Olivette, St

Louis, MO 63132, USA

Correspondence:

Dr. Lay-Sun Ma

Institute of Plant and Microbial Biology, Academia Sinica

128 Sec. 2, Academia Rd, Nankang, Taipei

11529 Taiwan.

Email: laysunma@gate.sinica.edu.tw

Tel: (+886) 02-2787-1145

Fax: (+886) 02-2782-7954

Abstract

Adapted plant pathogenic fungi deacetylate chitin to chitosan to avoid host perception and

disarm the chitin-triggered plant immunity. Whether plants have evolved factors to counteract

this fungal evasion mechanism in the plant-pathogen interface remains obscure. Here, we

decipher the underlying mechanism of maize cysteine-rich receptor-like secreted proteins

(CRRSPs)- AFP1, which exhibits mannose-binding dependent antifungal activity. AFP1

initials the action by binding to specific sites on the surface of yeast-like cells, filaments, and

germinated spores of the biotrophic fungi *Ustilago maydis*. This could result in fungal cell

growth and cell budding inhibition, delaying spore germination and subsequently reducing

fungal viability in a mannose-binding dependence manner. The antifungal activity of AFP1 is

conferred by its interaction with the PMT-dependent mannosylated chitin deacetylases (CDAs)

and interfering with the conversion of chitin. Our finding that AFP1 targets CDAs from

pathogenic fungi and nonpathogenic budding yeast suggests a potential application of the

CRRSP in combating fungal diseases and reducing threats posed by the fungal kingdom.

Key words: *Ustilago maydis*, chitin deacetylases, CRRSP, DUF26 domain,

mannosyltransferase PMT4, antifungal activity

Introduction

Plants safeguard the apoplast environment by deploying cell-surface localized pattern recognition receptors (PRRs) to sense potential danger signals via recognition of pathogen-associated molecular patterns (PAMP) or plant-derived damaged-associated molecular patterns (DAMP)[1,2]. This alerts the plant immune system and leads to the delivery of a mix of plant defense molecules to the apoplast to ward off pathogenic intruders[3]. Adapted pathogens are able to breach this defense barrier by shielding their cell surfaces, sequestering PAMPs, and modifying cell-surface glycans to avoid recognition and escape from the host attack at the initial steps of pathogenesis[4]. In return, plants evolve new weapons to counteract the fungal camouflage strategies.

The biotrophic fungus *Ustilago maydis* causes smut disease in maize by inducing large tumors in which the fungal hyphae proliferate and form spores[5]. The complex interplay between *U. maydis* and maize at the interface is mainly governed by an arsenal of fungal effector proteins induced in consecutive waves after the fungus contacts and colonizes maize[6]. To date, a few functionally characterized effectors have been reported to act in the apoplasts to suppress plant immunity and to protect hyphae from the attack of plant defense molecules. *U. maydis* Fly1 prevents the release of chitin by destabilizing plant chitinases[7], Rsp3 shields fungal hyphae to block the antifungal activity of maize AFP1 proteins[8], and Pit2 and Pep1 inhibit the activities of plant cysteine protease and peroxidases respectively[9,10]. *U. maydis* also possesses six active chitin deacetylases to convert chitin to chitosan to sustain fungal cell viability and evade recognition by chitin receptors and promote virulence[11].

Plant cysteine-rich receptor-like kinase (CRK) is one type of the transmembrane PRR proteins featured with domains of unknown function 26 (DUF26; stress-antifungal domain PF01657) which harbors a conserved cysteine motif ($C-X_8-C-X_2-C$) in their ectodomain[12,13]. CRKs have been implicated in plant defense, oxidative/salt stress and salicylic acid responses[14-

[17]. DUF26 domain has also been found in some plasmodesmata-localized proteins (PDLPs)

and cysteine-rich receptor-like secreted proteins (CRRSPs)[12,13]. PDLPs have a configuration

similar to CRKs but lack a kinase domain. They regulate plasmodesmata permeability and

immunity and modulate callose deposition in plasmodesmata[17,18]. CRRSPs harbor one or two

copies of the DUF26 domain and are secreted to the apoplast upon pathogen infection[8,19]. Two

of the maize CRRSPs (AFP1 and AFP2), display mannose-binding-dependent antifungal

activity and are significantly upregulated upon the perception of *U. maydis*[8]. AFP1/2-silenced

plants are more susceptible to the infection by *U. maydis* and *Colletotrichum graminicola*[8],

suggesting that maize AFP1/2 are involved in plant defense against a broad range of fungal

pathogens. So far, the antifungal activities of CRRSPs have been demonstrated for maize

AFP1[8] and *Ginkgo biloba* Gnk2[20] but not for cotton CRR1 which was shown to stabilize plant

chitinases [21]. Despite the importance of the DUF26-containing proteins in regulating biotic and

abiotic stress responses, their biological function in plant immunity is not clear, and the

antifungal mechanism of the CRRSP has not yet been elucidated.

In this work, we study maize AFP1 to provide mechanistic insights into the mode of action

of plant CRRSP. The provided evidence supports that maize AFP1 proteins could inhibit fungal

growth by interfering with the activity of chitin deacetylases which are essential for fungal

development and virulence. Interfering with the alteration of a fungal cell wall important

component highlights the potential use of CRRSPs as antifungal agents to act against a broad-

range of pathogenic and non-pathogenic fungi.

Results

AFP1 localizes on the growing tips and bud necks of yeast-like cells in *Ustilago maydis*

AFP1 displays a mannose-binding dependent antifungal activity[8]. However, the underlying

molecular mechanism has not yet been deciphered. To investigate the initial action mode of

AFP1, we examined the binding of AFP1 to *U. maydis* cells using immunolocalization. SG200

budding cells incubated with purified wild-type AFP1-His and mannose-binding-deficient

AFP1*-His proteins (contains S34A, R115A, E126A, N144A, Q227A, and E238A mutations

in the two mannose-binding sites)[8] were subjected to immunostaining using the anti-His

antibody and AF488-conjugated secondary antibody without any permeabilization and fixation

(Fig. 1A). The fluorescence of AFP1-His was detected throughout the cell body and showed

some accumulation in speckles on the cell periphery. Predominant fluorescence accumulation

was seen at cell division zones and in the growing tips of emerging buds (Fig.1A, central panel

and enlarged panel on the right). Besides that, AFP1 fluorescence was also detected at growing

tips and on the surface of SG200 filaments induced by hydroxy-fatty acids on a hydrophobic

surface without any permeabilization (Fig. S1A). AFP1*-His fluorescence could neither be

detected on yeast-like cells nor on filaments of SG200 (Fig. 1A and Fig. S1A), suggesting that

the AFP1 binding is dependent on its mannose-binding ability.

***U. maydis* CDAs display similar localization patterns as the maize AFP1 proteins**

The localization pattern of AFP1 in *U. maydis* cells is similar to that of *U. maydis* chitin

synthases (CHS) and chitinases (CTS), and is also correlated with the sites in the cell wall

where chitin and chitosan are accessible to antibodies in yeast-like cells[11,22-24]. This prompted

us to examine if AFP1 target the fungal CHS or CTS proteins. To test this idea, we examined

the interaction of AFP1 with either CHS or CTS using a yeast-two hybrid (Y2H) assay.

However, we could not detect the interactions between AFP1 and CHS or CTS in the Y2H

analysis (Fig. S2).

The conversion of chitin to chitosan is important for fungal development and is one of

the fungal evading strategies to avoid recognition by host plants. Chitin deacetylases might co-

localize with CHS to access to nascent chitin substrates. We posit that AFP1 might target chitin

deacetylases (CDAs) to block the deacetylation of chitin. The *U. maydis* genome contains seven *cda* genes that encode six C-terminal GPI (glycosylphosphatidylinositol) anchored CDAs and one secreted CDA protein (Cda4). While *cda6* is a pseudogene and not expressed[11], all active *cda* genes were differentially expressed in the axenic culture of SG200 cells (Fig. S3). Since the deletion of all *cda* genes is lethal to *U. maydis*, we initially examined the AFP1-His localization in a *cda* sextuple deletion mutant (Δcda2-7) that lacks six out of the seven *cda* genes[11]. Surprisingly, instead of detecting a weak AFP1 accumulation, we observed a more intense AFP1-His fluorescence in the Δcda2-7 mutant comparing to the parental SG200 cells (Fig. 1A). Since a significant increase in the expression of the *cda1* gene was detected in the deletion mutants Δcda2-6 and Δcda2-5 (Yanina S. Rizzi, personal communication), the *cda1* gene expression might also be upregulated in Δcda2-7 to compensate for the loss of other *cda* genes. This prompts us to speculate that the remaining Cda1 could be a potential binding target of AFP1.

Since the localization of CDA proteins in *U. maydis* has not been reported, we next explored the immunolocalization of the two highly expressed Cda1 and Cda2 proteins in *U. maydis* cells. We generated an N-terminal HA-tagged Cda1 protein (HACda1). However, we failed to immunolocalize HACda1 proteins on the cell surface without permeabilization even though the protein was detected in cell extracts (Fig. S4), suggesting that the N-terminus of Cda1 protein might undergo processing before it becomes membrane-anchored. An HA-tag was then inserted at the amino acid 76 of Cda1 (HA$_{76}$Cda1), and overexpressed in the *cda* quintuple mutant (Δcda1,3-6)[11] and *cda1* deletion mutant (Δcda1) to avoid the interference of endogenous Cda1 to compete with HA$_{76}$Cda1 for the localization. In both backgrounds, HA$_{76}$Cda1 fluorescence appeared in speckles at the growing tips and around the cell periphery of emerging cells but rarely accumulated in the cell periphery of mother cells (Fig. 1B and Fig. S1B). Occasionally, we also visualized the HA$_{76}$Cda1 speckles on the cell division zones (Fig.

1B; Enlarged panels). In filaments of Δcda(1,3-6)-HA$_{76}$Cda1 strain, the HA$_{76}$Cda1

fluorescence was also detected on the surface and growing tips of filamentous cells (Fig. S1B),

which was similar to the localization of AFP1-His on filaments of SG200 and Δcda(1,3-6)-

HA$_{76}$Cda1 (Fig. S1A and S1C). In comparison to the HA$_{76}$Cda1 localization pattern,

HA$_{118}$Cda2 speckles mainly appeared in the cell periphery of the *cda2* deletion mutant that

overexpressed HA$_{118}$Cda2 proteins (Δcda2-HA$_{118}$Cda2) (Figure 1B). Additionally, few

HA$_{118}$Cda2 speckles appeared at the cell tips and cell division zones (Fig. 1B; Enlarged panels).

Further localization analysis of the secreted Cda4 proteins expressed in Δcda(1,3-6)-Cda4HA

strain also revealed that the Cda4HA proteins exclusively localized to the cell periphery (Fig.

1B). These results show that the CDAs display similar localization patterns as the AFP1 in

yeast-like cells and filaments of *U. maydis* cells.

Maize AFP1 colocalizes and interacts with *U. maydis* Cda1

We next examined the possible interaction between AFP1 and Cda1 by analyzing their co-

localization. Δcda(1,3-6)-HA$_{76}$Cda1 cells were subject to immunostaining before being

incubated with the fluorescent probe DyLight550-labeled AFP1-His proteins (AFP1-His[550]).

We found that the fluorescence of HA$_{76}$-Cda1 and AFP1-His[550] overlapped at bud necks and in

the growing tips of the cells (Fig. 2A). To examine the interaction between AFP1 and Cda1 in

Δcda (1,3-6)-HA$_{76}$Cda1 cells at suboptical resolutions, we measured acceptor photobleaching

FRET (Fluorescence resonance energy transfer) efficiency[25-27] by bleaching the acceptor -

AFP1-His[550] to eliminate or reduce energy transfer from the donor HA$_{76}$Cda1, and thereby

yielding an increase in the donor fluorescence (Fig. 2B). The FRET efficiency of HA$_{76}$Cda1

after bleaching AFP1-His[550] increased significantly with an average of 23%, compared to 4%

in the negative control (WGA-AF488, wheat germ agglutinin conjugated to Alexa-Fluor 488

that recognizes chitin). This result suggests that Cda1 could be an AFP1-interacting target in

the *U. maydis*.

AFP1 targets more than one *U. maydis* chitin deacetylases and reduces CDA activity

The interaction of AFP1 with Cda1 prompted us to investigate whether AFP1 also directly

interacts with other *U. maydis* CDA proteins. Using yeast-two hybrid (Y2H) assay, we found

that AFP1 interacted with Cda1, 3, 5, and 7 but not with Cda2 and Cda4 even though the

proteins expressed at similar levels (Fig. 3A and Fig. S5A). When adding a competitive

inhibitor 3-AT (3-amino-1, 2, 4-triazole) to increase the stringency of the selection, Cda1 and

Cda7 were the only two targets left in the interaction and AFP1 displayed a stronger interaction

with the Cda1 (Fig. S5B). Furthermore, the CDA domain of Cda1 was sufficient to mediate the

interaction with AFP1 in the Y2H (Fig. S5C).

194   The AFP1-CDA interaction was further validated by immobilizing secreted C-terminal

Strep-tagged Cda1, 2, 3, 5, and 7 deleting respective GPI-anchor proteins (CDA-Strep) on

Strep-Tactin agarose beads and followed by incubation with purified AFP1 proteins.

Unexpectedly, the wild-type AFP1-His proteins were pulled down by all CDA-Strep proteins

but not by an unrelated protein control- 1204-Strep (UMAG01204, an *U. maydis* signal-peptide

containing protein) and the buffer control (Fig. 3B). Notably, the Cda1-AFP1 interaction also

required the mannose-binding ability of AFP1 as the Cda1-Strep could not pull down AFP1*-

His proteins (Fig. 3C). To analyze if AFP1 also interacts with the secreted Cda4 proteins

lacking GPI anchor, secreted Cda4HA (C-terminal HA-tagged Cda4) or Cmu1HA (*U. maydis*

secreted chorismate mutase)[28] from culture supernatants of strains Δcda(1,3-6)-Cda4HA or

SG200-Cmu1HA was collected and incubated with the immobilized AFP1-His proteins on Ni-

NTA agarose beads. AFP1-His proteins could pull down Cda4HA but not Cmu1HA proteins,

and non-specific binding of Cda4HA proteins to the beads was also not detected (Fig. 3D).

These results indicate that AFP1 could interact with all six *U. maydis* CDA proteins. Since the

CDA proteins share a conserved NodB homolog domain[11,29], AFP1 likely recognizes the

conserved structural fold of CDA proteins.

We next studied the impact of AFP1-CDA interaction on the chitin deacetylase activity

by measuring the release of acetate. The Cda1 or Cda2-Strep proteins were immobilized on

beads and incubated with either AFP1 proteins or buffer before adding the substrate GlcNAc5

(A5). The released acetate was significantly reduced when the Cda1-Strep and Cda2-Strep

proteins were incubated with AFP1-His, but not when incubated with AFP1*-His or buffer

control (Fig. 3E). Altogether, the results suggest that AFP1 binds to the CDA proteins and acts

to reduce the deacetylase activity of CDA in a mannose-binding-dependent manner.

Cda1 and Cda2 but not Cda4 are the substrates of PMT4

Considering AFP1-binding is the mannose-binding dependent, we reasoned that CDAs might

be mannosylated. We thus analyzed the AFP1-binding in a mannosyltransferase deletion

mutant of *U. maydis*, *pmt4*, which is defective in early infection-related development[30]. In the

Δpmt4 strain, the prominent AFP1 fluorescence on the bud necks and scars almost diminished,

whereas the intensity of speckles on the cell periphery became more evident (Fig. 4A and 4B).

These results indicate that the CDAs localized at the cell division zone and the growing tips

require the mannosylation activity of PMT4 and imply that the cell division localized Cda1 and

Cda2 but not the cell-periphery localized Cda4 could be the targets of PMT4.

We overexpressed $HA_{76}Cda1$, $HA_{118}Cda2$, and Cda4HA in SG200 and

the *pmt4* deletion mutant strain and examined the migration patterns of the proteins in both

backgrounds (Fig. 4C). Cda1 and Cda2 proteins displayed a smear and faster migration pattern

when expressed in the *pmt4* mutant compared to the wild-type SG200 background (Fig. 4C).

In contrast, the migration patterns of the cell-periphery localized Cda4HA in SG200 and

the *pmt4* mutant background were similar. The result illustrates that Cda1 and Cda2 but not

Cda4 are the substrates of PMT4. Further immunolocalization analysis of HA$_{76}$Cda1 and

HA$_{118}$Cda2 revealed no difference in the localization patterns of the CDA proteins expressed

in wild-type SG200 and the *pmt4* mutant (Fig. S6). Therefore, the deficiency in the PMT4-

dependent mannosylation does not alter the localization of Cda1 and Cda2 but reduces the

AFP1-binding. The incomplete abrogate AFP1-binding to cells of the *pmt* mutant suggests that

either AFP1 binds to the additional CDA targets that are not mannosylated by PMT4, e.g., Cda4.

Or, the decrease in the Cda1/Cda2 mannosylation may significantly reduce but not abolish the

interaction of Cda1/2 and AFP1. The result links the mannose-binding of AFP1 to

mannosylated CDAs.

AFP1 binds and blocks spore germination by inhibiting cell budding and growth of fungi

Since spores are the likely propagules in which the infections occur in the field and the *cda*

genes in *U. maydis* are upregulated and expressed during spore germination[11], we hypothesized

that AFP1 can bind *U. maydis* spores and inhibit its germination. To investigate this, we

examined the AFP1 localization on germinated spores of *U. maydis*. Asynchronous germinated

spores on agar were harvested and incubated with the AFP1 proteins before immunostaining.

While background noises were found in cell debris and mucilage but not in promycelium of

the AFP1*-His-treated spores (Fig. 5A), the AFP1-His fluorescence was seen at the bulged

regions of spores which presumably are the sites of promycelium emergence (Fig. 5B) and the

base of promycelium where it had protruded from spores (Fig. 5C-D). In addition, the

fluorescence also appeared at the tips of promycelium (Fig. 5E-F) and the regions between two

basidial cells (Fig. 5G-I). The results illustrate that AFP1 binds to germinating spores at distinct

morphological stages during germination, as early as the promycelium first protrusion.

Owing to asynchronous spore germination and low germination rate in *U. maydis*[31-34],

quantitative inhibitory assays of AFP1 on spore germination is challenging. To dissect the

inhibitory activity of AFP1 on spore germination, we performed time-lapse live-cell imaging

to monitor the germination process of individual spores in the presence or absence of

DyLight550 labeled AFP1^{550}-His proteins (Fig. 5J). In the absence of AFP1 proteins, most of

the spores remained dormant and only a few promycelium emerged from spores, elongated,

produced basidiospores, and then fell off. The germination process was slow down 5 hours

later and was at a halt in 7 hours after nutrients depleted. The emergence and elongation of

promycelium were detected between 20 minutes and 60 minutes, depending on the germination

stages of spores (Fig. 5J and Fig. S7). Compared to germinated spores with no AFP1^{550} added,

AFP1^{550}-bound promycelium that showed stronger fluorescence signal was either completely

stopped elongating (Fig. 5J; Upper panel) or the promycelium abandoned the AFP1^{550}-bound

basidial cells and produced a new basidial cell from another side to generate basidiospores and

escaped the inhibitory effect of AFP1. However, this escaping process could delay germination

by one to two hours (Fig. 5J; Lower panel). The results indicate that AFP1 binding could delay

and block spore germination.

The inhibitory effect of AFP1 in yeast-like cells was also observed (Fig. 5K). Budding

and growth of yeast-like cells were inhibited by the AFP1-His treatment but not by the AFP1*-

His treatment where cells continued growing and budding. Therefore, this result reinforces the

antifungal activity of AFP1 towards *U. maydis* yeast-like cells.

277   AFP1 targets pathogenic and non-pathogenic fungi by acting on conserved CDAs

Having shown that AFP1 acts on the *U. maydis* CDAs to inhibit spore germination and cell

growth, we next asked whether AFP1 also exerts inhibitory effects on other pathogenic and

nonpathogenic fungi. We localized AFP1 proteins on budding yeast *Saccharomyces*

cerevisiae (Sc) and the maize pathogen *Colletotrichum graminicola* (CgM2), which increases

virulence in AFP1-silenced plants[8]. We visualized AFP1 fluorescence on the conidial cell poles

of CgM2, and the bud sites and the cell division zones of budding yeast, whereas AFP1*-His

fluorescence was not detected in these regions (Fig. 6A). While yeast Sc*cda1 and cda2* genes

are expressed and required for spore wall assembly[35,36], six out of the seven *cda* genes were

expressed in the CgM2 conidial cells (Fig. S3B). In the Y2H assay, AFP1 interacted with two

of the most highly expressed CgM2 CDAs (GLRG7915 and GLRG386) and with a yeast

ScCda2 (Fig. 6B and 6C). The binding of the AFP1 to budding yeast cells resulted in reduced

cell viability, as evidenced by a significant decrease in the yeast cell numbers when treated with

the AFP1-His proteins. The same reduction in cell viability was not observed in the AFP1*-

His-treated yeast cells and mock controls (Fig. 6D). This data is consistent with our previous

finding that the AFP1 exhibits antifungal activity against the *U. maydis*[8]. These finding reveals

that AFP1 has the ability to inhibit a wide range of fungi by acting on the conserved fungal

CDA proteins.

Discussion

In this study, we demonstrate that AFP1 acts on fungal chitin deacetylases (CDAs) to interfere

with the conversion of chitin to chitosan. Given the importance and multifaceted roles of CDAs

in fungal cell growth across developmental stages and virulence[11,37-40], AFP1's action on the

CDAs could be an effective strategy to stop fungal colonization by inhibiting fungal cell growth

and spore germination, activating chitin-triggered plant immunity, and ultimately blocking the

fungal invasion (Fig. 7). To our knowledge, this is the first report revealing the molecular

mechanism of plant CRRSP proteins in plant-pathogen interactions. The maize CRRSP

proteins AFP1 displays inhibitory activities against a wide range of fungi, and counteracts the

strategy used by *U. maydis* to avoid recognition.

Mannose-binding defective mutant (AFP1*) proteins diminish the binding ability

supports that mannose-binding is the prerequisite for the localization of AFP1 on the fungal

cell surface. Together with the subsequent finding of a remarkable reduction in AFP1-binding

to the bud tips/necks in the Δpmt4 cells where the mannosylated Cda1 and Cda2 localize

provide a direct link between mannose-binding property of AFP1 and mannosylated CDA

proteins. We envisage that AFP1 could bind to the cell periphery-localized mannosylated CDAs,

e.g., Cda4 and Cda2 and GPI-anchored mannosylated CDAs. But, how AFP1 proteins bypass

the multilayers of cell walls to reach the GPI-anchored CDAs on the plasma membrane. We

anticipated that the CDA-located sites, i.e. cell division, cell poles and hyphal tips are actively

growing regions not well covered by cell-wall components, which might expose GPI-anchored

CDA proteins and facilitate the access of AFP1 to the targets.

PMT forms a heterodimer or homodimer in *Saccharomyces cerevisiae*[41]. We consider

that three PMTs in *Ustilago maydis* likely form a dimer complex to mannosylated their CDA

substrates. The smear migration pattern of Cda1/Cda2 in the *pmt4* mutant background could

indicate the mannosylation of the proteins by other PMTs (Fig. 4C). PMT1 and PMT2 could

also participate in the mannosylation of Cda1, Cda2, and Cda4. At present, we do not know in

addition to Cda1 and Cda2 whether other CDAs could also occupy the bud tips/necks and are

mannosylated by PMT4. We speculate that Cda7 could be another potential substrate of PMT4

since PMT4 is essential for pathogenesis, and the single deletion of *cda7* reduced fungal

virulence [11]. Although the Cda1/2 localization does not alter in the *pmt4* mutant, we could not

conclude that the mannosylation of CDA plays no role in the protein localization. A partial

mannosylation in Cda1/2 reduces the interaction with AFP1 but may be sufficient to mediate

the protein localization. Due to the essential role of PMT2 in fungal viability[30], we could not

delete all *pmt* genes to study the functionality of mannosylation in CDA localization and

understand the PMT-CDA specificity. However, we provide the evidence to support that the

mannose-binding of AFP1 and the PMT-dependent mannosylated CDAs are coupled (Fig. 3C

and Fig. 4).

In view of the AFP1 interaction with all CDAs in *Ustilago maydis* and other fungal

CDAs but not chitinases (CTS) and chitin synthases (CHS) that are putative mannosylated

proteins (as predicted by NetOGlyc Server, unpublished analysis), the mannose-binding

property of AFP1 appears not to play a role in target selection. The CDA domain of Cda1 is

sufficient in mediating the AFP1-Cda1 interaction suggesting that AFP1 likely recognizes the

conserved distorted (β/α)8-barrel structure of NodB homologous domain-containing active

catalytic sites of CDAs shared by the Carbohydrate Esterase family 4 (CE4) members.

According to the phylogenetic analysis of CDAs in the CE4 family, bacterial CDAs and CODs

(chitin oligosaccharide deacetylases) are clustered into a different clade and placed at a more

related distance to the fungal CDAs[42]. Whether the AFP1 inhibitory could extend to bacterial

species will be worth investigating. Based on the observation that AFP1 exhibited the strongest

interaction with the Cda1 in the Y2H assay, we reason that AFP1 might have different affinities

for each CDA target. The various degrees of mannosylation on CDAs might determine the

AFP1-CDA interaction affinity. Alternatively, a specific mannosylation pattern by a different

combination of PMT dimers may contribute to the interaction affinity.

Although the interaction of CDA proteins with any fungal membrane-associated

proteins has not been reported elsewhere, the localization patterns of CHS[23,43] and CDAs are

similar in *U. maydis* yeast-like cells. It is tempting to hypothesize that CDAs interact with

chitin synthases (CHS) to access de novo nascent chitin substrates and effectively convert them.

If this hypothesis holds true, AFP1 targeting on the CDAs might also have an additional

negative impact on the interaction between CDAs and CHS, therefore blocking the CDA access

to chitin. In the future, it would be interesting to investigate the biological roles of

mannosylation on the biophysical properties of CDAs in terms of protein distribution, enzyme

activity, and interaction with fungal cell wall proteins.

Our maize AFP1 study provides molecular insights into the understanding of the

antifungal mechanisms of plant CRRSP proteins. The finding that AFP1 targets conserved

fungal CDAs and displays antagonistic effects against several fungi tested makes its high

potential to be developed into an antifungal agent in controlling fungal diseases to reduce

threats posed by the fungal kingdom.

Methods

Strain construction and growth conditions. The haploid solopathogenic *Ustilago maydis*

SG200 strain was used as a reference strain in this study[44,45]. Strains used and generated in this

study are listed in Supplementary Table 1. Plasmid construction using either Gibson Assembly

or standard cloning methods as described in Supplementary Table 2. Primers used in each

generated plasmid are listed in Supplementary Table 3. A PCR-based approach was used to

generate mutants[46]. For gene integration into the *ip* locus or *mig2-6* locus, plasmids containing

a carboxin resistant *ip* allele (*ip*R)[47] or a *mig2-6* allele were linearized with the restriction

enzymes SspI or AgeI or EcoN1 and subsequently inserted via homologous recombination.

Transformation of *U. maydis* and genomic DNA isolation were performed as described.

Positive *U. maydis* transformants were verified by Southern blot analysis.

U. maydis strains were grown at 28 °C in a liquid YEPSL medium (0.4% yeast extract,

0.4% peptone, and 2% sucrose) and on potato dextrose agar (PDA; 2.4% potato dextrose broth

and 2% agar). Yeast cells were grown on YPD agar (BD Difco, USA) at 28 °C and *C.*

graminicola CgM2 was grown on oatmeal agar (BD Difco, USA) with continuous exposure to

daylight at room temperature.

Gene expression analysis. SG200 cell pellet was collected after cell grown in liquid YESPL

medium reached an OD_{600} of 0.6. *C. graminicola* (CgM2) conidial cells were harvested from

oatmeal agar plates. The cell pellet was washed with sterilized water and frozen in liquid

nitrogen. Total RNA extracted using TRIzol (Invitrogen) was subject to DNase-treatment

(Promega; Cat#M6101) according to the manufacturer's recommendation. The cDNA

preparation and quantitated RT-PCR analysis were performed as described previously [8]. The

expression of *U. maydis ppi* (peptidyl-prolyl isomerase) was used for normalization. To analyze

CgM2 *cda* gene expression using semiquantitative RT-PCR, the *actin* gene expression was used

for normalization. PCR amplification was programmed as follows: 95°C for 25 sec, 60°C for

25 sec, and 72°C for 45 sec. After 33 cycles for *cda* amplification and 28 cycles for *actin*, PCR

reactions were further incubated at 72°C for 3 min and then chilled at 10°C. Gel images were

acquired using the Molecular Imager Gel Doc XR+ system (Bio-RAD) and densitometric

analysis was performed using ImageJ[48].

Immunolocalization of AFP1 and CDA. The purification of AFP1 proteins from *Nicotiana*

benthamiana was performed as described [8]. To visualize the localization of AFP1-His on yeast-

like cells, a final volume of 200 µl of 1x PBS buffer (pH 7.2) containing 4 µg AFP1-His

proteins and 1 OD of cells (pre-washed with MES buffer) was incubated for 4 hours at 28 °C.

To visualize AFP1-His localization on germinated spores of FB1xFB2, spores were collected

from infected maize leaves and allowed to germinate on PD agar containing 150 µg/ml

ampicillin and 35 µg/ml chloramphenicol at 28°C as described[34]. Germinated spores on PD

agar were removed, washed, and incubated with 20 µg AFP1-His proteins in a final volume of

200 µl for 4 hours at 28 °C. The cells/spores were washed with PBS buffer before being

subjected to the immunostaining using the primary anti-His antibody and Alexa Fluor 488

(AF488)-conjugated secondary antibody as described[8].

To visualize the localization of HA-tagged CDA proteins, yeast-like cells were pre-washed

before being subjected to immunostaining using primary anti-HA antibody and Alexa Fluor

488 (AF488)-conjugated secondary antibody[8] in 10mM MES buffer (pH5.5). To visualize the

protein localization on filaments of *U. maydis*, cells were induced using hydroxyl fatty acid on

parafilm as described[8] and followed by either an initiated incubation with AFP1-His proteins

as described above or directly subjected to immunostaining. AF488 fluorescence was excited

at 488 nm and subsequently detected at 500-540 nm using Axio Observer fluorescence

microscope equipped with Axiocam 702 Monochrome camera (ZEISS, Germany). Images

were processed using ZEN 3.2 imaging software (ZEISS).

Co-localization and Acceptor photobleaching. The fluorescence-labeled AFP1-His proteins

were prepared according to the manufacturer's instruction of the DyLight 550 Antibody

Labeling kit (Thermo Fisher; Cat#84530). One OD_{600} of cells washed with MES buffer was

subjected to immunostaining using the primary anti-HA antibody and AF488-conjugated

secondary antibody or wheat germ agglutinin conjugated antibody (WGA-A488; Molecular

Probes, Karlsruhe, Germany) to localize CDA proteins or chitin respectively. The

immunostained cells or WGA-488 stained cells were washed with MES buffer, followed by

adding 4 µg of DyLight 550 labeled AFP1-His proteins (AFP1^{550}) to a final volume of 200 µl,

and then continued the incubation at room temperature for 4 hours. The AFP1^{550}-treated cells

were washed and visualized using epi-fluorescence microscopy.

Acceptor photobleaching measurements were performed using a confocal microscope

(ZEISS LSM880). The donor (AF488) channel was excited at 488 nm and detected through a

500-540 nm emission filter. The acceptor (DyLight 550) channel was excited at 561 nm and

detected through a 580-630 nm emission filter. The images in the donor and acceptor channels

were acquired before and following acceptor fluorescent bleaching. Regions of interest (ROI)

that overlap with the donor-acceptor fluorescence were selected for photo-destruction by

applying 100% laser power at a wavelength of 561 nm for 120 iterations. Evaluation of images

was analyzed using the ZEN 3.0 software. The FRET efficiency was calculated according E_{FRET}

$= (I_{Donor(post)} - I_{Donor(pre)})/I_{Donor(post)}$ where $I_{Donor(pre)}$ and $I_{Donor(post)}$ are the donor fluorescence

intensities prior to and following photo-destruction of the acceptor.

Live cell time lapse imaging. Spores germinated on PD agar were harvested, incubated with

0.1 µg/µl AFP1-His[550] proteins in 1x PBS buffer for 4 hours, and mixed with agar to a final

concentration of 2% which containing 2% PD medium before being spotted on a chambered

coverslip (Ibidi, Catalog #: 80827). Image series were acquired using Olympus DeltaVision

Core microscope system with a 63x water immersion objective. Frame time was set at 10 min

and a total of 42 frames were acquired. For monitoring AFP1 inhibitory on yeast-like cells,

SG200 cells (OD_{600} = 0.002) were incubated with AFP1-His proteins for 4 hours at room

temperature before being spotted on a chambered coverslip to acquire images. Frame time was

set at 5 min and a total of 85 frames was acquired in a 7-hour duration.

CDA activity assay. Cells expressing secreted CDA-Strep proteins were cultured in YEPSL

liquid medium until OD_{600} reached 0.6. A total 75 ml culture supernatant was concentrated and

exchanged with buffer W using 10 kD cutoff centrifugal filters (Sartorius, Germany). A 50 µl

of Strep-Tactin®XT agarose beads were added to the concentrated sample in a final volume of

1 ml containing 1x protease inhibitor cocktail and incubated overnight at 4 °C. The beads were

washed with 50 mM TEA (Triethanolamine) buffer (pH 8) and aliquoted evenly into three tubes

before 10 µg of AFP1-His, AFP1*-His, or TEA buffer was added to a final volume of 350 µl.

The incubation was carried out at room temperature for 4 hours and followed by adding

GlcNAc$_5$ (A5; Megazyme Cat# O-CHI5) to a final concentration of 0.625 mg/ml and continued

the incubation at 37 °C for 16 hours. The reaction mixture was centrifuged. The release of

acetate in the supernatant was measured by following the manufacturer's instruction of the

Acetic Acid Test Kit (R Biopharm Inc. Cat#10148261035).

Pull down assay. Cells expressing secreted Strep-tag Cda proteins were grown to OD_{600} of

0.6 at 28 °C. The 50 ml culture supernatant containing Strep-tagged CDA proteins was

concentrated to 1 ml and exchanged to buffer W (100 mM Tris/HCl, 150 mM NaCl, 1 mM

EDTA, pH 8) using 10 kD cutoff centrifugal filters (Sartorius, Germany) before being

incubated with 25 µl of Strep-Tactin®XT agarose beads (IBA Lifesciences, Germany)

overnight at 4 °C in presence of protease inhibitor cocktail (Roche, Switzerland). The beads

were washed with buffer W, and incubated with 0.1 µg AFP1-His proteins in a final volume of

500 µl binding buffer (100 mM Tris/HCl, 0.3 M NaCl, 1 mM EDTA, 0.05% NP-40, 1x protease

inhibitor cocktail, pH 8) at 4 °C for 2 hours, and then washed with binding buffer. The bound

proteins were removed by boiling in sample buffer and subjected to immunoblotting analysis

using mouse monoclonal anti-6xHis (Yao-Hong Biotech. Inc., TW) and Strep-tagII (IBA

Lifesciences, Germany) primary antibodies and the goat anti-mouse IgG-HRP-conjugated

secondary antibody (Thermo Scientific, USA). For the interaction of Cda4-AFP1, 0.2 µg of

AFP1-His proteins was incubated with 20 µl Ni-NTA agarose beads (Qiagen, Germany) in

buffer (25 mM Tris-Cl, 0.15 M NaCl, 10 mM Imidazole, pH 7.5) for 1 hour at 4 °C. The beads

were washed with the same buffer, incubated with the concentrated culture supernatant

containing Cda4HA in buffer (25 mM Tris-Cl, 0.3 M NaCl, 0.1% NP-40, 10 mM Imidazole,

pH 7.5) at 4 °C for 2 hours. The beads were washed and boiled in a sample buffer to remove

the bound proteins for the immunoblotting analysis using mouse monoclonal anti-HA and anti-

6xHis antibodies (Yao-Hong Biotech. Inc., TW).

Yeast two-hybrid assay. Yeast two-hybrid assays were performed as described[49]. Yeast

(AH109) transformants containing the desired plasmids were screened on a selective dropout

(SD) medium lacking tryptophan (W) and leucine (L) (Clontech). Protein interactions were

483 assessed on SD selection medium lacking LW, histidine (H), and Adenine (A) or lacking LWH

484 and containing 3-amino-1, 2, 4-triazole (3-AT) after three to five days incubation at 28 °C. To

485 detect the expression of proteins in yeast transformants, one OD_{600} of yeast cells was lysed in

486 a buffer containing 1% β-mercaptoethanol and 0.25 M NaOH. The supernatant fraction was

487 TCA-precipitated and then centrifuged. The protein pellet was dissolved in HU-buffer (100

488 mM Tris-Cl pH 6.8, 8 M urea, 5% SDS, 0.1 mM EDTA, and 0.1% bromophenol blue) and

489 analyzed by immunoblotting.

491 **Yeast growth inhibition assay**. For the yeast growth inhibition assay, a 100 µl final volume

492 of a reaction containing yeast cells BY4742 (OD_{600} =0.001) and 20 µg AFP1 proteins were

493 incubated at 28°C for 4 hours in 10mM MES buffer (pH5.5). The cells were serial-diluted,

494 spread on YPD plates (3 plates per each dilution), and incubated at 28°C for two days until

495 colonies appeared. Colony numbers were then manually counted.

497 **Author contributions**

498 LSM conceived and designed the study and wrote the manuscript. WLT performed

499 immunostaining and FRET analysis. WLT and RMK did live-cell imaging and time-lapse

500 microscopy. RMK performed the initial AFP1 binding analyses. MYX did protein purification.

501 MYX, WLT, and YHL generated constructs and CDA overexpression strains. MYX, WLT,

502 YHL, FAD, and HCL performed Y2H assays. YHL analyzed *cda* gene expression. WLT and

503 LSM did pull-down assays. WLT and FAD analyzed CDA activity inhibition assay and yeast

504 growth-inhibition assay. All authors discussed the results and commented on the manuscript.

506 **Acknowledgements**

507 We are indebted to Regine Kahmann for supply of the *cda* deletion mutants and other strains

(Import permit# 109-F-500 and 107-F-002) and critically reading the manuscript. We are

grateful to Erh-Min Lai, Chih-Hang Wu, and Ooi-Kock Teh for suggestions and comments on

the manuscript. We thank Prof. Dr. Bruno Moerschbacher and Mounashree J. Urs for providing

technical advice on CDA activity analysis. We acknowledge Ji-Ying Huang in the Plant Cell

Biology Core Laboratory for assisting in live-cell imaging and FRET microscopy and the staff

in the DNA Sequencing Core Laboratory at the Institute of Plant and Microbial Biology for

technical assistance. We thank Chibbhi K. Bhaskar and Cuong V. Hoang for their comments

on an early version of the manuscript. Our work is supported by funds from the Ministry of

Science and Technology, Taiwan (MOST# 107-2311-B-001 -044 -MY3).

Conflict of interest

519   All authors declare no conflict of interests regarding the publication of this article.

References

522   1   Jones, J. D. & Dangl, J. L. The plant immune system. *Nature* **444**, 323-329,
523   doi:10.1038/nature05286 (2006).

524   2   Monaghan, J. & Zipfel, C. Plant pattern recognition receptor complexes at the plasma
525   membrane. *Curr Opin Plant Biol* **15**, 349-357, doi:10.1016/j.pbi.2012.05.006 (2012).

526   3   Doehlemann, G. & Hemetsberger, C. Apoplastic immunity and its suppression by
527   filamentous plant pathogens. *New Phytol* **198**, 1001-1016, doi:10.1111/nph.12277
528   (2013).

529   4   Buscaill, P. & van der Hoorn, R. A. L. Defeated by the nines: nine extracellular
530   strategies to avoid MAMP recognition in plants. *Plant Cell*,
531   doi:10.1093/plcell/koab109 (2021).

532   5   Lanver, D. *et al.* Ustilago maydis effectors and their impact on virulence. *Nat Rev*
533   *Microbiol* **15**, 409-421, doi:10.1038/nrmicro.2017.33 (2017).

534   6   Lanver, D. *et al.* The Biotrophic Development of Ustilago maydis Studied by RNA-
535   Seq Analysis. *Plant Cell* **30**, 300-323, doi:10.1105/tpc.17.00764 (2018).

7   Okmen, B. *et al.* Dual function of a secreted fungalysin metalloprotease in Ustilago

maydis. *New Phytol* **220**, 249-261, doi:10.1111/nph.15265 (2018).

8   Ma, L. S. *et al.* The Ustilago maydis repetitive effector Rsp3 blocks the antifungal

activity of mannose-binding maize proteins. *Nat Commun* **9**, 1711,

doi:10.1038/s41467-018-04149-0 (2018).

9   Mueller, A. N., Ziemann, S., Treitschke, S., Assmann, D. & Doehlemann, G.

Compatibility in the Ustilago maydis-maize interaction requires inhibition of host

cysteine proteases by the fungal effector Pit2. *PLoS Pathog* **9**, e1003177,

doi:10.1371/journal.ppat.1003177 (2013).

10   Hemetsberger, C., Herrberger, C., Zechmann, B., Hillmer, M. & Doehlemann, G. The

Ustilago maydis effector Pep1 suppresses plant immunity by inhibition of host

peroxidase activity. *PLoS Pathog* **8**, e1002684, doi:10.1371/journal.ppat.1002684

(2012).

11   Rizzi, Y. S. *et al.* Chitosan and Chitin Deacetylase Activity Are Necessary for

Development and Virulence of Ustilago maydis. *mBio* **12**, doi:10.1128/mBio.03419-

20 (2021).

12   Chen, Z. A superfamily of proteins with novel cysteine-rich repeats. *Plant Physiol*

126, 473-476 (2001).

13   Vaattovaara, A. *et al.* Mechanistic insights into the evolution of DUF26-containing

proteins in land plants. *Commun Biol* **2**, 56, doi:10.1038/s42003-019-0306-9 (2019).

14   Czernic, P. *et al.* Characterization of an Arabidopsis thaliana receptor-like protein

kinase gene activated by oxidative stress and pathogen attack. *Plant J* **18**, 321-327

(1999).

15   Yadeta, K. A. *et al.* A Cysteine-Rich Protein Kinase Associates with a Membrane

Immune Complex and the Cysteine Residues Are Required for Cell Death. *Plant*

Physiol **173**, 771-787, doi:10.1104/pp.16.01404 (2017).

16   Ohtake, Y., Takahashi, T. & Komeda, Y. Salicylic acid induces the expression of a

number of receptor-like kinase genes in Arabidopsis thaliana. *Plant Cell Physiol* **41**,

1038-1044 (2000).

17   Hunter, K. *et al.* CRK2 Enhances Salt Tolerance by Regulating Callose Deposition in

Connection with PLDalpha1. *Plant Physiol* **180**, 2004-2021, doi:10.1104/pp.19.00560

(2019).

18   Lee, J. Y. *et al.* A plasmodesmata-localized protein mediates crosstalk between cell-to-

cell communication and innate immunity in Arabidopsis. *Plant Cell* **23**, 3353-3373,

doi:10.1105/tpc.111.087742 (2011).

19   Kim, S. G. *et al.* In-depth insight into in vivo apoplastic secretome of rice-

572   Magnaporthe oryzae interaction. *J Proteomics* **78**, 58-71,

573   doi:10.1016/j.jprot.2012.10.029 (2013).

574   20   Miyakawa, T. *et al.* A secreted protein with plant-specific cysteine-rich motif

575   functions as a mannose-binding lectin that exhibits antifungal activity. *Plant Physiol*

576   **166**, 766-778, doi:10.1104/pp.114.242636 (2014).

577   21   Han, L. B. *et al.* The Cotton Apoplastic Protein CRR1 Stabilizes Chitinase 28 to

578   Facilitate Defense against the Fungal Pathogen Verticillium dahliae. *Plant Cell* **31**,

579   520-536, doi:10.1105/tpc.18.00390 (2019).

580   22   Treitschke, S., Doehlemann, G., Schuster, M. & Steinberg, G. The myosin motor

581   domain of fungal chitin synthase V is dispensable for vesicle motility but required for

582   virulence of the maize pathogen Ustilago maydis. *Plant Cell* **22**, 2476-2494,

583   doi:10.1105/tpc.110.075028 (2010).

584   23   Weber, I., Assmann, D., Thines, E. & Steinberg, G. Polar localizing class V myosin

585   chitin synthases are essential during early plant infection in the plant pathogenic

586   fungus Ustilago maydis. *Plant Cell* **18**, 225-242, doi:10.1105/tpc.105.037341 (2006).

587   24   Langner, T. *et al.* Chitinases Are Essential for Cell Separation in Ustilago maydis.

588   *Eukaryot Cell* **14**, 846-857, doi:10.1128/EC.00022-15 (2015).

589   25   Kenworthy, A. K. Imaging protein-protein interactions using fluorescence resonance

590   energy transfer microscopy. *Methods* **24**, 289-296, doi:10.1006/meth.2001.1189

591   (2001).

592   26   Kenworthy, A. K., Petranova, N. & Edidin, M. High-resolution FRET microscopy of

593   cholera toxin B-subunit and GPI-anchored proteins in cell plasma membranes. *Mol*

594   *Biol Cell* **11**, 1645-1655, doi:10.1091/mbc.11.5.1645 (2000).

595   27   Karpova, T. S. *et al.* Fluorescence resonance energy transfer from cyan to yellow

596   fluorescent protein detected by acceptor photobleaching using confocal microscopy

597   and a single laser. *J Microsc* **209**, 56-70, doi:10.1046/j.1365-2818.2003.01100.x

598   (2003).

599   28   Djamei, A. *et al.* Metabolic priming by a secreted fungal effector. *Nature* **478**, 395-

600   398, doi:10.1038/nature10454 (2011).

601   29   Zhao, Y., Park, R. D. & Muzzarelli, R. A. Chitin deacetylases: properties and

602   applications. *Mar Drugs* **8**, 24-46, doi:10.3390/md8010024 (2010).

603   30   Fernandez-Alvarez, A., Elias-Villalobos, A. & Ibeas, J. I. The O-mannosyltransferase

604   PMT4 is essential for normal appressorium formation and penetration in Ustilago

maydis. *Plant Cell* **21**, 3397-3412, doi:10.1105/tpc.109.065839 (2009).

31 Banuett, F. & Herskowitz, I. Discrete developmental stages during teliospore
formation in the corn smut fungus, Ustilago maydis. *Development* **122**, 2965-2976
(1996).

32 Sacadura, N. T. & Saville, B. J. Gene expression and EST analyses of Ustilago maydis
germinating teliospores. *Fungal Genet Biol* **40**, 47-64, doi:10.1016/s1087-
1845(03)00078-1 (2003).

33 Ostrowski, L. A., Seto, A. M. & Saville, B. Investigating Teliospore Germination
Using Microrespiration Analysis and Microdissection. *J Vis Exp*, doi:10.3791/57628
(2018).

34 Fukada, F., Rossel, N., Munch, K., Glatter, T. & Kahmann, R. A small Ustilago
maydis effector acts as a novel adhesin for hyphal aggregation in plant tumors. *New*
Phytol **231**, 416-431, doi:10.1111/nph.17389 (2021).

35 Christodoulidou, A., Briza, P., Ellinger, A. & Bouriotis, V. Yeast ascospore wall
assembly requires two chitin deacetylase isozymes. *FEBS Lett* **460**, 275-279,
doi:10.1016/s0014-5793(99)01334-4 (1999).

36 Christodoulidou, A., Bouriotis, V. & Thireos, G. Two sporulation-specific chitin
deacetylase-encoding genes are required for the ascospore wall rigidity of
Saccharomyces cerevisiae. *J Biol Chem* **271**, 31420-31425,
doi:10.1074/jbc.271.49.31420 (1996).

37 Upadhya, R. *et al.* Cryptococcus neoformans Cda1 and Its Chitin Deacetylase Activity
Are Required for Fungal Pathogenesis. *mBio* **9**, doi:10.1128/mBio.02087-18 (2018).

38 Gao, F. *et al.* Deacetylation of chitin oligomers increases virulence in soil-borne
fungal pathogens. *Nat Plants* **5**, 1167-1176, doi:10.1038/s41477-019-0527-4 (2019).

39 Geoghegan, I. A. & Gurr, S. J. Chitosan Mediates Germling Adhesion in Magnaporthe
oryzae and Is Required for Surface Sensing and Germling Morphogenesis. *PLoS*
Pathog **12**, e1005703, doi:10.1371/journal.ppat.1005703 (2016).

40 Baker, L. G., Specht, C. A., Donlin, M. J. & Lodge, J. K. Chitosan, the deacetylated
form of chitin, is necessary for cell wall integrity in Cryptococcus neoformans.
Eukaryot Cell **6**, 855-867, doi:10.1128/EC.00399-06 (2007).

41 Girrbach, V. & Strahl, S. Members of the evolutionarily conserved PMT family of
protein O-mannosyltransferases form distinct protein complexes among themselves. *J*
Biol Chem **278**, 12554-12562, doi:10.1074/jbc.M212582200 (2003).

42 Grifoll-Romero, L., Pascual, S., Aragunde, H., Biarnes, X. & Planas, A. Chitin

639   Deacetylases: Structures, Specificities, and Biotech Applications. *Polymers (Basel)*
640   **10**, doi:10.3390/polym10040352 (2018).

641 43 Weber, I., Gruber, C. & Steinberg, G. A class-V myosin required for mating, hyphal
642   growth, and pathogenicity in the dimorphic plant pathogen Ustilago maydis. *Plant*
643   *Cell* **15**, 2826-2842, doi:10.1105/tpc.016246 (2003).

644 44 Muller, P., Aichinger, C., Feldbrugge, M. & Kahmann, R. The MAP kinase kpp2
645   regulates mating and pathogenic development in Ustilago maydis. *Mol Microbiol* **34**,
646   1007-1017, doi:10.1046/j.1365-2958.1999.01661.x (1999).

647 45 Kamper, J. *et al.* Insights from the genome of the biotrophic fungal plant pathogen
648   Ustilago maydis. *Nature* **444**, 97-101, doi:10.1038/nature05248 (2006).

649 46 Kamper, J. A PCR-based system for highly efficient generation of gene replacement
650   mutants in Ustilago maydis. *Mol Genet Genomics* **271**, 103-110, doi:10.1007/s00438-
651   003-0962-8 (2004).

652 47 Broomfield, P. L. & Hargreaves, J. A. A single amino-acid change in the iron-sulphur
653   protein subunit of succinate dehydrogenase confers resistance to carboxin in Ustilago
654   maydis. *Curr Genet* **22**, 117-121, doi:10.1007/BF00351470 (1992).

655 48 Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
656   image analysis. *Nat Methods* **9**, 671-675, doi:10.1038/nmeth.2089 (2012).

657 49 Tanaka, S. *et al.* A secreted Ustilago maydis effector promotes virulence by targeting
658   anthocyanin biosynthesis in maize. *Elife* **3**, e01355, doi:10.7554/eLife.01355 (2014).

Fig. 1. AFP1 and *U. maydis* CDAs display similar localization patterns. (A) Yeast-like cells of indicated strains were incubated with either AFP1-His or AFP1*-His (defects in mannose-binding) proteins and followed by immunostaining with an anti-His antibody and an AF488-conjugated secondary antibody. Bars, 20 μm. (B) Cells expressing HA-tagged CDA proteins under constitutive promoter *otef* in indicated deletion strains were subjected to immunostaining using anti-HA antibody and an AF488-conjugated secondary antibody to localize CDA proteins. White arrows indicate cells enlarged in right panels. Bars, 20 μm.

Fig. 2. The co-localization of Cda1 and AFP1. Cells of Δcda(1,3-6)-HA$_{76}$cda1 strain were immunostained to localize HA$_{76}$cda1 (A, B) or stained with WGA-AF488 to locate chitin (B) before incubation with DyLight 550 labeled AFP1-His proteins (AFP1^{550}) and visualized the fluorescence by confocal microscopy. (B) FRET efficiency was calculated using the indicated equation. Number of spots from three biological replicates used in the analysis as indicated above column. Values represent mean ± standard deviation. Representative images of cells showing fluorescence of AFP1^{550} and HA$_{76}$Cda1/WGA-AF488 were taken and measured before and after the photobleaching of acceptor AFP1^{550} (middle panel). Cell images with overlapped fluorescence of HA$_{76}$Cda1 or WGA-AF488 and AFP1^{550} and the signal intensity in overlapping regions are shown (right panel).

Fig. 3. AFP1 acts on CDA proteins to interfere chitin deacetylase activity. (A) Yeast two-hybrid assay to detect interaction of AFP1 and CDA proteins. Yeast cells with indicated plasmids were grown on SD/-Leu/-Trp (LW) and SD/-Leu/-Trp/-His/-Ade (LWHA) plates for 3 days. Self-interaction of chorismate mutase (Cmu1) was served as the positive control and AD-AFP1/BD as negative control. Similar results were observed in at least two independent experiments. (B-C) In vitro pull down assay of CDA-Strep and AFP1 proteins. Indicated C-terminal Strep-tagged CDA or 1204 (*U. maydis* secreted proteins; unrelated protein control) proteins were immobilized on Strep-Tactin agarose beads and served as baits to pull down AFP1-His or AFP1*-His proteins. (D) AFP1-His proteins were immobilized on Ni-NTA agarose beads and incubated with C-terminal HA-tagged Cda4 and Cmu1 (negative control). (E) Enzyme activity of Cda1 and Cda2 in presence of AFP1-His and AFP1*-His. Cda1-Strep and Cda2-Strep proteins immobilized on Strep-Tactin agarose beads were treated with or without AFP1 proteins before incubated with GlcNAc$_5$ (A5) substrates. Data given represent the mean ± SD values of the results from three independent experiments. Immunoblots showing similar amount of proteins in each reaction and could be detected after incubation (lower panel). Asterisks indicate significant differences of CDA activity in respective reaction compared with buffer control (-) determined by a two-tailed Student's *t*-test. *P* values are shown.

Fig. 4. Pmt4 mannosylates Cda1 and Cda2. (A) Yeast-like cells of indicated strains were incubated with AFP1-His and followed by immunostaining with an anti-His antibody and an AF488-conjugated secondary antibody. Bars, 20 µm. Arrows indicate the bud neck/cell pole localization of AFP1. (B) Total number of cells (n) from three biological replicates in experiment a as indicated at above each column was analyzed. The percentage of cells with AFP1 binding at bud necks/cell poles was calculated. Values represent mean ± sd of three biological replicates. Arrows indicate AFP1 fluorescence on cell pole /bud neck of cells of Δpmt4 strain. (C) PMT4 mannosylates Cda1 and Cda2. $HA_{76}Cda1$ and $HA_{118}Cda2$ were overexpressed in either SG200 or Δpmt4 strains. The total protein extracts collected from the cell pellet of indicated strain were treated with (+) or without (-) deglycosylation enzyme mix (NEB Cat# P6039) according to the manufacturer's protocol, and the glycosylation pattern of CDA proteins were analyzed by immunoblotting using anti-HA antibody. (D) Immunolocalization of $HA_{76}Cda$ and $HA_{118}Cda2$ overexpressed in indicated strains. Bars, 20 µm.

K

Fig. 5. AFP1 inhibits spore germination, cell budding and the growth of budding cells. *U. maydis* FB1x FB2 germinated spores and SG200 yeast-like cells were treated with AFP1-His and AFP1*-His proteins in agar or liquid medium containing 2% nutrients. Bars,10 µm. (A-I) AFP1-treated germinated spores were immunostained with an anti-His antibody and an AF488-conjugated secondary antibody to visualize AFP1 fluorescence. j, Images of germinated spores with or without AFP1^{550} added were acquired at indicated time points. Yellow arrows in two Z-stack images indicate AFP1^{550}-His binding sites on the germinated spore. (K) Images of AFP1-treated SG200 yeast-like cells were acquired at indicated time points. White arrows track the growth of cells over time in medium with AFP1-His or AFP1*-His proteins added.

Fig. 6. AFP1 acts on CDA of pathogenic and non-pathogenic fungi to affect cell survival. (A) *C. graminicola* and *S. cerevisiae* cells were treated with AFP1-His and AFP1*-His proteins, and followed by immunostaining. Bars:10 μm. (B-C) Yeast two-hybrid assays to detect interaction of AFP1 and CDA proteins of *C. graminicola* (B) or *S. cerevisiae* (C). Yeast transformants containing indicated plasmids were grown on SD/-Leu/-Trp (LW) for growth control and on a SD/-Leu/-Trp/-His/-Ade (LWHA) or LWH plates containing 10 mM of 3AT (3-amino-1,2,4-triazole) to assess protein interaction. Self-interaction of chorismate mutase (Cmu1) was served as positive control, and AD-AFP1/BD and AD/BD-CDA were used as negative controls. Similar results were observed in two independent experiments. Sc: *S. cerevisiae.* (D) *S. cerevisiae* cells incubated with AFP1-His, AFP1*-His and mock (buffer) for 4 hours, serial diluted, and plated on YPAD agars. Colony forming units (CFUs) of yeast cells plating on agar plates were quantified. Values represent mean ± sd of three biological replicates. ***$p < 0.001$ indicates significant differences compared to the CFUs in mock determined by a two-tailed Student's *t*-test.

Spores germinate and produce basidial cells

CDA

AFP1

C. graminicola

AFP1 targets fungal CDAs which are either secreted or GPI-anchored proteins to interfere conversion of chitin

Spore germination and cell growth and budding are blocked by AFP1

Fig.7. Antifungal action mode of AFP1. Upon pathogen invasion, plants deliver AFP1 proteins to the apoplast to stop the fungal colonization. It interferes with conserved fungal CDA proteins by blocking the conversion of chitin to chitosan, which is necessary for fungal cell development and virulence[11]. By acting on fungal CDAs, AFP1 could prevent spore germination, stop cell budding, and inhibit fungal growth, ultimately leading to fungal cell death and stopping the fungal invasion.

Supplementary Information

Maize CRRSP acts against a broad range of fungi by inhibiting chitin deacetylases

Lay-Sun Ma[1]*, Wei-Lun Tsai[1], Raviraj M. Kalunke[1,2], Meng-Yun Xu[1], Yu-Han Lin[1], Florensia Ariani Damei[1], and Hui-Chun Lee[1]

[1]Institute of Plant and Microbial Biology, Academia Sinica, Taipei 11529, Taiwan.

[2]Present address: Donald Danforth Plant Science Center, 975 N Warson Rd, Olivette, St Louis, MO 63132, USA

Fig. S1. Immunostaining of AFP1 and Cda1 in *U. maydis* cells. (A) Filaments of SG200 induced by hydroxyl fatty acids on hydrophobic surface were treated with AFP1-His and AFP1*-His proteins, and followed by immunostaining with an anti-His antibody and an AF488-conjugated secondary antibody. Bars, 20 μm. (B) Yeast-like cells and filaments of Δcda(1,3-6):HA_{76}Cda1 strain which constitutively expressed HA_{76}Cda1 proteins under promoter *otef* in Δcda(1,3-6) deletion mutant were subject to immunostaining using an anti-HA antibody and an AF488-conjugated secondary antibody. (C) Filaments of Δcda(1,3-6):HA_{76}Cda1 strain treated with AFP1-His proteins before subjected to immunostaining using an anti-His antibody and an AF488-conjugated secondary antibody. Bars, 20 μm. Arrow heads indicate AFP1-His or HA_{76}Cda1 fluorescence on hyphal tips.

A

Fig. S2. Yeast-two hybrid analysis of AFP1- CHS (chitin synthases) and AFP1-CTS (chitinase) interactions. Yeast two-hybrid assay to detect interaction of AFP1 and CHS/CTS proteins. Yeast cells containing the indicated plasmids were grown on SD/-Leu/-Trp (LW) and SD/-Leu/-Trp/-His/-Ade (LWHA) plates for 2-3 days. Self-interaction of chorismate mutase (Cmu1) was served as the positive control and AD-AFP1/BD and BD-AFP1/AD and BD/CHS as negative controls. Similar results were observed in at least two independent experiments.

A

B

Fig. S3. *cda* gene expression of *U. maydis and C. graminicola*. (A) Total RNA was extracted from SG200 cells grown in YEPSL liquid medium and subjected to quantitative RT-PCR. Expression levels of *U. maydis cda* gene were normalized relative to the constitutively expressed peptidyl-prolyl isomerase (ppi). Three biological replicates were analyzed. Values represent mean ± sd. Right panel showed the schematic drawings of CDA proteins with indicated signal peptides (grey boxes) and predicted GPI anchors (black boxes). a. a., amino acids. (B) Total RNA was extracted from *C. graminicola* conidial cells grown on agar plates and subjected to semi-quantitative RT-PCR. Values were obtained by quantified intensities of PCR bands in the right panel using ImageJ software, and were normalized relative to the constitutively expressed *actin* gene. Data represent mean ± sd of the three biological replicates shown in right panel.

Fig. S4. Immunoblot analysis of HA-tagged CDA protein expression. HA-tagged CDA proteins were expressed under constitutive promoter *otef* in the indicated deletion mutants. The indicated strains grown in YEPSL liquid medium to OD_{600} of 0.6-0.7 were harvested and adjusted to OD_{600} of 20. Proteins from cell pellets were prepared and subjected to immunoblot analysis using anti-HA or anti tubulin antibodies as indicated.

Fig. S5. Yeast-two hybrid analysis of AFP1-CDA interaction. (A) Immunoblot analysis of CDA and AFP1 protein expression in yeast transformants. Yeast cells containing indicated plasmids were grown in YPD liquid medium until OD_{600} of 0.7-0.8. One OD of cell pellet was collected, lysed, and TCA precipitated and analyzed by immunoblots using indicated antibodies. Two independent clones were selected for analysis. (B, C) Y2H assay to detect interaction of AFP1 with (B) CDA proteins and (C) CDA domain of Cda1 protein. Yeast cells with indicated plasmids were grown on SD-LW and SD-LWHA or SD-LWH containing 1mM of 3AT plates for 3 days. Self-interaction of chorismate mutase (Cmu1) was served as the positive control. Similar results were observed in at least two independent experiments.

Fig. S6. Immunolocalization of HA$_{76}$Cda and HA$_{118}$Cda2. Cells expressing HA-tagged CDA proteins under constitutive promoter *otef* in indicated strains were subjected to immunostaining using anti-HA antibody and an AF488-conjugated secondary antibody to localize CDA proteins. Bars, 20 μm. (Right) Wild-type SG200 cells subjected to immunostaining using anti-HA antibody and AF488-conjugated secondary antibody was served as negative control.

Fig. S7. Spore germination in absence of AFP1 proteins. *U. maydis* FB1x FB2 germinated spores in agar containing 2% PD medium. Bars,10 µm. Images of germinated spores without AFP1 proteins added were acquired at indicated time points. White arrows track emergence of promycelium over incubation time.

Supplementary Table 1: Strains used in this study

Strain	Genotype	Resistance[*]	References
SG200	*a1 mfa2 bW2 bE1, ble;*	P	1
SG200Δpmt4	*a1 mfa2 bW2 bE1, ble; umag05433(pmt4)::cbx*	P, C	2
SG200Δcda2,3,4,5,6[em],Δ7	*a1 mfa2 bW2 bE1, ble; cda2Δ121 - 134; cda3Δ21 - 28; cda4Δ146 - 155; cda5 Δ101 -113; cda6 Δ74 - 83; umag02381(cda7)::hyg*	P, HY	3
SG200Δcda1,3,4,5,6 [em]	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83*	P	3
SG200_P$_{otef}$-cmu1HA	*a1 mfa2 bW2 bE1, ble; ip^r[P$_{otef}$-cmu1-HA]ip^s*	P, C	4
SG200Δcda1	*a1 mfa2 bW2 bE1, ble; umag00638(cda1)::hyg*	P, HY	This work
SG200Δcda2	*a1 mfa2 bW2 bE1, ble; umag01143(cda2)::hyg*	P, HY	This work
SG200Δcda1_ P$_{otef}$-HA$_{76}$cda1	*a1 mfa2 bW2 bE1, ble; cda1::hyg; ip^R[P$_{otef}$-HA$_{76}$cda1]ip^S*	P, HY, C	This work
SG200Δcda2_ P$_{otef}$-HA$_{118}$cda2	*a1 mfa2 bW2 bE1, ble; cda2::hyg; ip^R[P$_{otef}$-HA$_{118}$cda2]ip^S*	P, HY, C	This work
SG200Δcda1,3,4,5,6[em] P$_{otef}$-HAcda1	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{otef}$-HAcda1]ip^S;*	P, C	This work
SG200Δcda1,3,4,5,6 [em] P$_{otef}$-HA$_{76}$cda1	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{otef}$-HA$_{76}$cda1]ip^S;*	P, C	This work
SG200Δcda1,3,4,5,6[em] P$_{act}$-cda1-Strep(ΔGPI)	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{act}$-cda1-Strep(ΔGPI)]ip^S;*	P, C	This work
SG200Δcda 2,3,4,5,6 [em],Δ7 P$_{act}$-cda2-Strep(ΔGPI)	*a1 mfa2 bW2 bE1, ble; cda2Δ121-134; cda3Δ21-28; cda4Δ146-155; cda5 Δ101-113; cda6 Δ74-83; cda7::hyg; ip^R[P$_{act}$-cda2-Strep(ΔGPI)]ip^S;*	P, HY, C	This work
SG200Δcda1,3,4,5,6[em] P$_{act}$-cda3-Strep(ΔGPI)	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{act}$-cda3-Strep(ΔGPI)]ip^S;*	P, C	This work
SG20Δcda1,3,4,5,6[em] P$_{otef}$-cda5-Strep(ΔGPI)	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{otef}$-cda5-Strep(ΔGPI)]ip^S;*	P, C	This work
SG200Δcda2,3,4,5,6[em],Δ7 P$_{otef}$-cda7-Strep(ΔGPI)	*a1 mfa2 bW2 bE1, ble; cda2Δ121-134; cda3Δ21-28; cda4Δ146-155; cda5 Δ101-113; cda6 Δ74-83;cda7:: hyg; ip^R[P$_{otef}$-cda7-Strep(ΔGPI)]ip^S;*	P, HY, C	This work
SG200Δcda1,3,4,5,6[em] P$_{cda4}$-cda4HA	*a1 mfa2 bW2 bE1, ble; cda1Δ121-127; cda3Δ21-28; cda4Δ146-155; cda5Δ101-113; cda6Δ74-83; ip^R[P$_{cda4}$-cda4HA]ip^S;*	P, C	This work
SG200Δpmt4 P$_{otef}$-HA$_{76}$cda1(mig2-6 locus)	*a1 mfa2 bW2 bE1, ble; um05433::cbx; [P$_{otef}$-HA$_{76}$cda1]mig2-6;*	P, C, G	This work
SG200Δpmt4 P$_{otef}$-HA$_{118}$cda2	*a1 mfa2 bW2 bE1, ble; um05433::cbx; [P$_{otef}$-HA$_{118}$cda2]mig2-6;*	P, C, G	This work
SG200Δpmt4 P$_{otef}$-cda4HA	*a1 mfa2 bW2 bE1, ble; um05433::cbx; [P$_{otef}$-cda4HA]mig2-6;*	P, C, G	This work
SG200 P$_{otef}$-HA$_{76}$cda1	*a1 mfa2 bW2 bE1, ble; ip^R[P$_{otef}$- HA$_{76}$cda1] ip^S;*	P, C	This work
SG200 P$_{otef}$-HA$_{118}$cda2	*a1 mfa2 bW2 bE1, ble; ip^R[P$_{otef}$- HA$_{118}$cda2] ip^S;*	P, C	This work
SG200 P$_{otef}$-cda4HA	*a1 mfa2 bW2 bE1, ble; ip^R[P$_{otef}$- cda4HA] ip^S;*	P, C	This work

* Phleomycin (P), Hygromycin (HY), Carboxin (C), Geneticin (G)

Supplementary Table 2: Plasmids used in this study

Plasmids	Description	Reference
p123	Plasmid containing the *gfp* gene controlled by constitutive promoter *otef, nos* terminator, the *U. maydis* carboxin resistant *ip* allele (ip^R), and ampicillin resistance gene. This plasmid served as backbone to insert gene of interest ectopically into the *U. maydis ip* locus.	5
pHwtFRT	Plasmid containing the hygromycin resistance cassette (Hyg^R).	6
pJET-cda1KO	The left and right borders of cda1 gene were amplified from SG200 gDNA using the primer pairs #608/609 and #610/611. The hygromycin resistance cassette was obtained from SfiI digestion of pHwtFRT. The three DNA fragments were merged with the EcoRV digested pJET plasmid via Gibson assembly.	This study
pJET-cda2KO	The left and right borders of cda2 gene were amplified from SG200 gDNA using the primer pairs #612/613 and #614/615. The hygromycin resistance cassette was obtained from SfiI digestion of pHwtFRT. The three DNA fragments were merged with the EcoRV digested pJET plasmid via Gibson assembly.	This study
P$_{act}$-mcherry	p123-derived plasmid containing the *mcherry* gene under the control of the *actin* promoter (UMAG_11232).	7
P$_{otef}$-vp1HA	p123-derived plasmid used in constitutively expressing secreted Vp1HA in *U. maydis*	8
P$_{otef}$-CAP-strep	A PCR fragment amplified using the primer pairs #102/65 from SG200 gDNA was digested with XmaI/XbaI and ligated into with the XmaI/XbaI digested P*act*-cda2strep (ΔGPI).	This study
P$_{act}$-cda1-strep (ΔGPI)	Two PCR fragments amplified using the primer pairs #409/410 from SG200 gDNA and #76/411 from p123 plasmid, were merged with the NcoI/EcoRV digested P*act*-mcherry via Gibson assembly.	This study
P$_{act}$-cda2-strep (ΔGPI)	A XmaI/XbaI digested PCR fragment containing cda2 gene amplified using the primer pairs #453/454 from SG200 gDNA and a XbaI/EcoRV digested fragment containing strepII from P*act*-cda1strep (ΔGPI) were ligated with XmaI/EcoRV digested P*act*-mcherry vector.	This study
P$_{act}$-cda3-strep (ΔGPI)	A XmaI/XbaI digested PCR fragment amplified from SG200 gDNA with the primer pairs #455/456 was ligated with the XmaI/XbaI digested P*act*-cda2strep (ΔGPI) plasmid.	This study
P$_{otef}$-cda5-strep (ΔGPI)	A XmaI/XbaI digested PCR fragment amplified from SG200 gDNA using primer pair #436/473 and a XbaI/EcoRV digested fragment from P*act*-cda2strep (ΔGPI) were ligated with the XmaI/EcoRV digested p123 plasmid.	This study
P$_{otef}$-cda7-strep (ΔGPI)	A XmaI/XbaI digested PCR fragment amplified from SG200 gDNA using primer pair #438/457 was ligated into the XmaI/XbaI digested P*otef*-cda5strep (ΔGPI) plasmid.	This study
P$_{cda4}$-cda4HA	A KpnI/XbaI digested PCR fragment containing the native promoter and ORF of *cda4* gene amplified from SG200 gDNA using primer pair #1/2 was ligated into the KpnI/XbaI digested P*otef*-vp1HA backbone.	This study
P$_{otef}$-HAcda1	Two PCR fragments amplified using the primer pairs #358/370 and #368/369 from SG200 gDNA, were merged with the BamHI/NotI digested p123 vector via Gibson assembly.	This study
P$_{otef}$-HA$_{76}$cda1	Two fragments of cda1 amplified using the primer pairs #358/391 and #368/369 from SG200 gDNA, were merged with BamHI/NotI digested p123 vector via Gibson assembly.	This study
P$_{otef}$-HA$_{118}$cda2	Two fragments of cda2 amplified using the primer pairs #453/5 and #3/4 from SG200 gDNA were digested with XmaI/NgoMIV and NgoMIV/NotI respectively and ligated into the XmaI/NotI digested p123 vector.	This study
Yeast two-hybrid constructs		
pGADT7	Yeast expression vector that is designed to express a protein of interest fused to a GAL4 activation domain	Clontech
pGBKT7	Yeast expression vector that is designed to express a protein of interest fused to a GAL4 DNA binding domain	Clontech
pAD-cmu1/pBD_cmu1	A NdeI/EcoRI digested PCR fragment amplified from SG200 gDNA using primer pair #354/355 was ligated into the NdeI/EcoRI digested pGADT7 and pGBKT7 vectors.	This study
pAD-AFP1/BD-AFP1	A NdeI/BamHI digested-PCR fragment amplified from maize cDNA using primer pair #288/289 was ligated into the NdeI/BamHI digested pGADT7 and pGBKT7 vectors.	This study
pBD-cda1	A NdeI/EcoRI digested-PCR fragment amplified from SG200 gDNA using primer pair #352/356 was ligated into the NdeI/EcoRI digested pGBKT7 vector.	This study
pBD-cda2	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #374/375 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study
pBD-cda3	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #376/377 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study

pBD-cda4	A NdeI/EcoRI digested-PCR fragment amplified from SG200 gDNA using primer pair #378/379 was ligated into the NdeI/EcoRI digested pGBKT7 vector.	This study
pBD-cda5	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #380/381 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study
pBD-cda7	A NdeI/EcoRI digested-PCR fragment amplified from SG200 gDNA using primer pair #401/402 was ligated into the NdeI/EcoRI digested pGBKT7 vector.	This study
pBD-cts1	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #290/291 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study
pBD-cts2	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #292/293 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study
pBD-cts3	A NdeI/BamHI digested-PCR fragment amplified from SG200 gDNA using primer pair #294/295 was ligated into the NdeI/BamHI digested pGBKT7 vector.	This study
pBD-cts4	A SfiI/XmaI digested-PCR fragment amplified from SG200 gDNA using primer pair #296/297 was ligated into the SfiI/XmaI digested pGBKT7 vector.	This study
pBD-Sccda1	A NcoI/NotI digested-PCR fragment amplified from AH109 cDNA using primer pair #474/475 was ligated into the NcoI/NotI digested pGBKT7 vector.	This study
pBD-Sccda2	A NcoI/NotI digested-PCR fragment amplified from AH109 cDNA using primer pair #476/477 was ligated into the NcoI/NotI digested pGBKT7 vector.	This study
pBD-Cg386	A NdeI/EcoRI digested-PCR fragment amplified from *C. graminicola* CgM2 cDNA using primer pair #488/489 was ligated into the NdeI/EcoRI digested pGBKT7 vector.	This study
pBD-Cg7915	Two PCR fragments amplified from *C. graminicola* CgM2 cDNA using the primer pairs #490/559 and #560/491 were combined and performed the overlap extension PCR. The overlap extension PCR fragment was digested with NdeI and EcoRI, and ligated into the NdeI/EcoRI digested pGBKT7 vector.	This study
pAD-chs5	A SfiI/BamHI digested-PCR fragment amplified from SG200 using primer pair #323/324 was ligated into the SfiI/BamHI digested pGADT7 vector.	This study
pAD-chs6	A SfiI/XmaI digested-PCR fragment amplified from SG200 using primer pair #325/326 was ligated into the SfiI/XmaI digested pGADT7 vector.	This study
pAD-chs7	A NdeI/SacI digested-PCR fragment amplified from SG200 using primer pair #327/328 was ligated into the NdeI/SacI digested pGADT7 vector.	This study

Supplementary Table 3: Oligonucleotides used in this study

Name	Sequence (5' to 3')
Oligonucleotides for plasmid construction	
#1	CTCGGTACCATCGCGGTTGTCGTCGTACT
#2	GGTCTAGAGTTAGCCGCTCCAGGAGC
#3	CCTGCCGGCTACCCCTACGACGTGCCCGACTATGCCACCGACATGACCAACGTCTG
#4	TAAGCGGCCGCTCAGAGCAAGAGAGCAAAAGCG
#5	TAGCCGGCAGGGTAGACGGTG
#65	ATATCTAGAGTTGGGCGAGACGTTCTC
#76	CCTTTGAGTGAGCTGATACC
#102	CCCCCGGGATGAAAGTCACATCTGTGATCG
#288	CTGCATATGGCGGACTCCATCGGCAGCTAC
#289	GACGGATCCTTAAGGACGCACGACGATCTTG
#290	GCTCATATGTTTGGACGTCTTAAGCACAGG
#291	GATGGATCCTACTTGAGGCCGTTCTTG
#292	GCTCATATGGTGCCTCACGAGCAGAGC
#293	GATGGATCCTAGCTCAATCCGGCAGCGTC
#294	GCTCATATGGCGCTGAACAATGATGGATC
#295	GTGGAATTCCTAGGAAGAGATAGCACCTG
#296	TATGGCCATGGAGGCCCTTTGGCCTCACCCAACCAC
#297	CCACCCGGGCTAATGCATGTTGCACATGCC
#323	TATGGCCATGGAGGCCAATCCTTTCGAATCTCTTCC
#324	GATGGATCCTCAGTCAAAGCTTGGAGGAG
#325	TATGGCCATGGAGGCCTCGACCAAAGACGCCCACG
#326	TTACCCGGGTCAGGCTTGTGCGCCACCGGAC
#327	GCTCATATGCCCGCAGTTGAGCGCAAC
#328	CTCGAGCTCAACTGAATCGATCATGATAG
#352	GCTCATATGGGCGACTTCACCGTCAAGATCC
#354	GCTCATATGGCTGTATCTGGCAAGTCG
#355	GTGGAATTCTTAGGTGCACTTGTTGGCGTGG
#356	GTGGAATTCTCAAAGCAGAGTTGCCAC
#358	CACAGACAACATCATCCACGGGATCCATGCTGCGTTTGCTACTTC
#368	TACCCATACGACGTACC
#369	CGATCTGCAGCCGGGCGGCCGCTCAAAGCAGAGTTGCCAC
#370	TCTGGTACGTCGTATGGGTAAGCAAGAGCGCTGCCAGCTA
#374	GCTCATATGGCTGGACGCCACGAGCGAGGCCTTGT
#375	GATGGATCCTCAGAGCAAGAGAGCAAAAGCG
#376	GCTCATATGAACATCGGCCGTGGCGTAC
#377	GATGGATCCTCAGGCGAGCATGGCGATGG
#378	GCTCATATGTCCCCCACTACAATGAGCAC
#379	GTGGAATTCTCAGTTAGCCGCTCCAGGAG
#380	GCTCATATGAGTCCCGACTTTGAGAAGCG
#381	GATGGATCCTTAGACGAGAAGACCGAAGAAG
#391	AGCGTAATCTGGTACGTCGTATGGGTAGCTGCCGCCTTGATTGAG
#401	GCTCATATGCATGGTGGCGACCTTCTC
#402	TGGAATTCTTAGATAAAGACCATCATAGCACCG
#409	AAAACCTAGTTCACAGTCATCCCATGCTGCGTTTGCTACTTC
#410	TTTTTCAAACTGCGGATGTGACCATCTAGAGCTGGTGGCGATGCCCTGTTTG
#411	TGGTCACATCCGCAGTTTGAAAAATAGAGCGGCCGCCCGGCTGCA
#436	TTTTTCAAACTGCGGATGTGACCATCTAGAGGTGCCAGCACCCCCAGTC
#438	TTTTTCAAACTGCGGATGTGACCATCTAGAGCTGCCGAACGTCCTGTAG
#453	CATCCCGGGATGCGTCTCTCCGTCTCCG
#454	ACCATCTAGAACTCGACTGACTCTTG
#455	CATCCCGGGATGAAGCTTTCCTCGACAGC
#456	CCATCTAGACTTATCTGCGGACTG
#457	CATCCCGGGATGAAGTCTACCACGGTATTCTCG
#473	CATCCCGGGATGGTCAAAATCACTTTTGC
#474	TGGCCATGGAATCAAATGGGAGTACCGCATTGATGGG
#475	TATGCGGCCGCCTAGTCGTAGCGTTCGATG
#476	TGGCCATGGCCGAAGCTAATAGGGAAGATTTAAAG
#477	GCCGCGGCCGCTTAGGACAAGAATTCTTTTATG
#488	GCTCATATGGGCCCGGTCACCCGCCGTCAAAG
#489	GGGAATTCTTAGGACTTGTACCAGTTCTC
#490	CGCCATATGACTCCTTTCCGCCGCCAAAC
#491	GCCGAATTCTCAGGCACAGGTACCGAAAAG

#559	CGGTGTTTTGGTGAACATCGTGGGCCAGAGCGATG
#560	TCTGGCCCACGATGTTCACCAAAACACCGCCGAGT
#608	GCTCGAGTTTTTCAGCAAGATAAATATTGCACGAGTATATTTTCGAGAGG
#609	AGAATAGGAACTTCTGGCCATCTAGGCCGTCGGAAAAAAAAGCGATCTCTC
#610	AGTATAGGAACTTCTGGCCTGAGTGGCCACTTAGCATTGCCAACTGCATTC
#611	AGGAGATCTTCTAGAAAGATAATATTGCAGATGTTGATTCTTTGAGG
#612	GCTCGAGTTTTTCAGCAAGATAATATTCTTGGTTTGCCTCAGATGTTC
#613	AGAATAGGAACTTCTGGCCATCTAGGCCGATATGAAAAGTCGATGAGGG
#614	AGTATAGGAACTTCTGGCCTGAGTGGCCGCTTCACCATACGGCTCAGC
#615	AGGAGATCTTCTAGAAAGATAATATTGCAGCGCAACTCGCACCATAC
Oligonucleotides for RT-PCR	
ppi-F	ACATCGTCAAGGCTATCG
ppi-R	AAAGAACACCGGACTTGG
cda1F (UMAG_00638)	CGAGTACAAGTGCTACTACC
cda1R	AGGCCGAAGGTAGCAGGAAC
cda2F (UMAG_01143)	ACCAGAACGCGTGGTTCATC
cda2R	CGACAGCCAAAGCGCCAAAG
cda3F (UMAG_11922)	CCACGGGTACCAACACCAAC
cda3R	ATGGCCGAGAGCAAGACGAC
cda4F (UMAG_01788)	CAACCACGAGACCGTCCAAG
cda4R	CTGCAGGTCCAGGTGGAATC
cda5F (UMAG_02019)	TTCCCGCCGCTGCTCAAATG
cda5R	TTGGTGGAAGTGAGCTGATG
cda6F (UMAG_05792)	TGGCTCTGGGAACCGTCAAG
cda6R	TACCCATCACGGTGGCAACAG
cda7F (UMAG_02381)	ACTTGTGCGTTTGACGACTC
cda7R	ACACGGTCCAGCCAATCAAG
GLRG_07915_F	TCTGGCCCACGATGTTCAC
GLRG_07915_R	TAGGAGTAGCGGTGCTAGTG
GLRG_11148_F	CAATTGCTGCTCTCACAGGTC
GLRG_11148_R	TAGAAGCCGAGCCGGATGTAG
GLRG_11238_F	TGCTGTTCCAGTGCGGGATGGTG
GLRG_11238_R	AGCCTTGGCATTGTACGCCTTG
GLRG_11241_F	TCGTCAAACGAGCCGAGTG
GLRG_11241_R	GGTGGTAATTATTGAGCCATAAG
GLRG_09099_F	CGAGCAGTACTTGAGGAAC
GLRG_09099_R	GTATGGGAAGTGGGATCTC
GLRG_00386_F	CGTCACCGACCTCACTAAC
GLRG_00386_R	ATCTTCTGCACGAGGGAGATG
GLRG_03854_F	GGCTTCGTACACTGACGACTTG
GLRG_03854_R	GCCGAGGATGTTGCGAATAGCC
GLRG_02208_F	TTGCTGCTCCTCGTACGGATG
GLRG_02208_R	CCGGCAACAGCAGTCGACAAACAG
Actin F (GLRG_05255)	GTGGCAGCACTCTGTACAAG
Actin R (GLRG_05255)	GCCGTATTCCTCGTACTCAG

References

Kamper, J. *et al.* Insights from the genome of the biotrophic fungal plant pathogen Ustilago maydis. *Nature* **444**, 97-101, doi:10.1038/nature05248 (2006).

Fernandez-Alvarez, A., Elias-Villalobos, A. & Ibeas, J. I. The O-mannosyltransferase PMT4 is essential for normal appressorium formation and penetration in Ustilago maydis. *Plant Cell* **21**, 3397-3412, doi:10.1105/tpc.109.065839 (2009).

Rizzi, Y. S. *et al.* Chitosan and Chitin Deacetylase Activity Are Necessary for Development and Virulence of Ustilago maydis. *mBio* **12**, doi:10.1128/mBio.03419-20 (2021).

Djamei, A. *et al.* Metabolic priming by a secreted fungal effector. *Nature* **478**, 395-398, doi:10.1038/nature10454 (2011).

Aichinger, C. *et al.* Identification of plant-regulated genes in Ustilago maydis by enhancer-trapping mutagenesis. *Mol Genet Genomics* **270**, 303-314, doi:10.1007/s00438-003-0926-z (2003).

Khrunyk, Y., Munch, K., Schipper, K., Lupas, A. N. & Kahmann, R. The use of FLP-mediated recombination for the functional analysis of an effector gene family in the biotrophic smut fungus Ustilago maydis. *New Phytol* **187**, 957-968, doi:10.1111/j.1469-8137.2010.03413.x (2010).

Lanver, D. *et al.* The Biotrophic Development of Ustilago maydis Studied by RNA-Seq Analysis. *Plant Cell* **30**, 300-323, doi:10.1105/tpc.17.00764 (2018).

Cuong V. Hoang, C. K. B., Lay-Sun Ma. A Novel Core Effector Vp1 Promotes Fungal Colonization and Virulence of *Ustilago maydis*. *J. of Fungi* **7**, doi:10.3390/jof7080589 (2021).