1 ***Pseudomonas zeiradicis* from corn and *Pseudomonas soyae* from soybean, two**
2 **new *Pseudomonas* species and endophytes from agricultural crops**
3

4 Jacqueline Lemaire[1], Sarah Seaton[1], Patrik Inderbitzin[1], Martha E. Trujillo[2]
5

6 [1]Indigo Ag, Inc., 500 Rutherford Avenue, Boston, MA 02129, United States
7 [2]Departamento de Microbiología y Genética, Campus Miguel de Unamuno,
8 University of Salamanca, Salamanca, Spain

9
10

ABSTRACT

Two novel *Pseudomonas* species associated with healthy plants and other habitats
are described from the United States. They are *Pseudomonas zeiradicis* strain PI116 from
corn in Missouri, compost from Massachusetts, urban soil from Iowa and water of Lake
Erie; and *Pseudomonas soyae* strain JL117 from soybean in Indiana and Wisconsin, and
soil in Wyoming. No plant pathogenic strains are known for any of the novel species based
on genome comparisons to assemblies in GenBank.

Keywords: novel species, taxonomy, bacteria, prokaryotes, endophytes, agriculture
20
21

INTRODUCTION

Pseudomonas is a large and diverse genus in the *Gammaproteobacteria* comprising
more than 180 named species (Hesse et al. 2018). *Pseudomonas* species are rod-shaped,
Gram-type negative chemoheterotrophs, which are motile by means of polar flagella, and
can utilize an array of small organic molecules as sources of carbon and energy. This
nutritional diversity is reflected in the high abundance of *Pseudomonas* species in nature,
with as many as 10^6 fluorescent pseudomonads residing in a single gram of soil
(Tarnawski et al. 2003; Vančura 1980). *Pseudomonas* species are strictly aerobic, but some
can utilize NO_3 as an electron acceptor in place of O_2. In addition, nitrate can be used as a
nitrogen source for all known species (Stanier et al. 1966).
Pseudomonads are ubiquitous in the rhizosphere, often living in a commensal relationship
with plants and utilizing plant-exuded nutrients. *Pseudomonas* species also impact plant
health by suppressing phytopathogens (Couillerot et al. 2009; Gerbore et al. 2014),

enhancing local access to nutrients, and inducing systemic resistance in the plant host

(Bakker et al. 2007; De Vleesschauwer et al. 2008). A subset of rhizosphere

pseudomonads are further adapted to life as endophytes, residing within cells, the

intercellular spaces or the vascular system of host plants (Mitter et al. 2013). We describe

two such endophytic strains, representing two novel species of the genus *Pseudomonas*.

Pseudomonas zeiradicis sp. nov. strain PI116, which was isolated from healthy, field-

grown corn in Missouri, United States, and *Pseudomonas soyae* sp. nov. strain JL117 from

healthy, field-grown soybean in Indiana, United States. We provide phenotypic and

phylogenomic details describing these two species, and report geographic distribution and

additional substrates from conspecific, cultured strains by comparison to whole genomes at

GenBank.

METHODS

Isolation. Strain PI116 was isolated from healthy field-grown *Zea mays* roots in

Missouri, United States, and strain JL117 from seedlings of *Glycine max* in Indiana,

United States. Plant tissue was washed with a mild detergent to remove particulates,

surface-sterilized with bleach (1% v/v sodium hypochlorite) and ethanol (70% v/v), and

homogenized. Serial dilutions of tissue homogenate were plated on a panel of media types

for endophyte cultivation. All strains were streaked to purity and stored in glycerol (20%

v/v) at -80°C until subjected to further testing.

Motility. Strains were tested for flagellar-dependent swimming and swarming

motility on R2A plates solidified with 0.3% and 0.6% agar, respectively. Three

independent colonies were inoculated onto R2A broth and grown for 36 hr at 24°C. Broth

cultures were normalized to an OD600 of 0.1, and 1.5 µl of culture was spotted directly

onto the surface of the motility agar. The diameter of colony expansion was measured for

5 days.

Carbon source utilization. Substrate utilization was assessed using Biolog GenIII

Microplates (Catalogue No. 1030) (Biolog Inc., Hayward, CA). Each bacterium was

inoculated in duplicate plates using Protocol A described by the manufacturer, with the

exception that plates were incubated at 30°C for 36 hr. Respiration leading to reduction of

the tetrazolium indicator was measured by absorbance at 590 nm.

Biochemical analyses. Catalase activity was evaluated by immediate effervescence

after the application of 3% (v/v) hydrogen peroxide solution via the tube method, a

positive reaction was indicated by the production of bubbles. *Staphylococcus aureus*

NCIMB 12702 and *Streptococcus pyogenes* ATCC 19615 were used as positive and

negative controls, respectively. Oxidase activity was evaluated via the oxidation of Kovács

oxidase reagent, 1% (w/v) tetra-methyl-p-phenylenediamine dihydrochloride in water, via

the filter-paper spot method. A positive reaction was indicated when the microorganism's

color changed to dark purple. *Pseudomonas aeruginosa* NCIMB 12469 and *Escherichia*

coli ATCC 25922 were used as positive and negative controls, respectively.

Phylogenetic and genomic analyses. DNA was extracted from pure cultures using

the Omega Mag-Bind Universal Pathogen Kit according to manufacturer's protocol with a

final elution volume of 60µl (Omega Biotek Inc., Norcross, GA). DNA samples were

quantified using Qubit fluorometer (ThermoFisher Scientific, Waltham, MA) and

normalized to 100 ng. DNA was prepped using Nextera DNA Flex Library Prep kit

according to manufacturer's instructions (Illumina Inc., San Diego, CA). DNA libraries

were quantified via qPCR using KAPA Library Quantification kit (Roche Sequencing and

Life Science, Wilmington, MA) and combined in equimolar concentrations into one 24-

sample pool. Libraries were sequenced on a MiSeq using pair-end reads (2x200bp). Reads

were trimmed of adapters and low-quality bases using Cutadapt (version 1.9.1) and

assembled into contigs using MEGAHIT (version 1.1.2) (Li et al. 2015). Reads were

mapped to contigs using Bowtie2 (version 2.3.4) (Langmead and Salzberg 2012), and

contigs were assembled into scaffolds using BESST (2.2.8) (Sahlin et al. 2014).

16S rRNA gene sequences were extracted from genome assemblies using barrnap

(Seemann 2019) and submitted to GenBank (OL306320, OL306321).

Phylogenomic analyses were performed using GToTree (Lee 2019) with default

settings. Taxon sampling for each species is described in the respective phylogenetic tree

figure legend.

Average nucleotide identity analyses were performed using the pyani ANIm

algorithm (Richter and Rosselló-Móra 2009) implemented in the MUMmer package

(Kurtz et al. 2004) retrieved from https://github.com/widdowquinn/pyani.

Geographic distribution and host range of novel species were inferred by ANI to

assemblies from unidentified strains from GenBank (Ciufo et al. 2018) and the Indigo

internal collection. An ANI threshold of ≥95% indicated conspecificity (Chun et al. 2018;

Richter and Rosselló-Móra 2009). Distribution maps were generated in R version 4.0.4 (R

Core Team 2021) with tigris 1.5 (Walker 2021), sf 1.0-2 (Pebesma 2018), ggplot2 3.3.2

(Wickham 2016), cowplot 1.1.1 (Wilke 2020) and other packages.

RESULTS

Genomic analyses

Pseudomonas zeiradicis sp. nov. strain PI116

Strain PI116 shared 99.9% 16S rRNA gene sequence identity with *Pseudomonas*

mohnii CCUG 53115[T] and less with the remaining *Pseudomonas* species. A phylogenomic

tree using GToTree (Lee 2019) confirmed the affiliation of strain PI116 with the genus

Pseudomonas. PI116 was most closely related to *P. umsongensis* DSM 16611[T] with 100%

bootstrap support (Figure 1). The top average nucleotide identity (ANI) value of PI116

was 90.8% with *P. umsongensis*. This value was well below the threshold for species

demarcation (Richter and Rosselló-Móra 2009; Chun et al. 2018) providing further

genomic support that strain PI116 represents a new genomic species of *Pseudomonas*.

Pseudomonas soyae sp. nov. strain JL117

Strain JL117 shared 99.5% 16S rRNA gene sequence identity with *Pseudomonas*

reinekei CCUG 53116[T] and less with the remaining *Pseudomonas* species. A

phylogenomic tree using GToTree (Lee 2019) confirmed the affiliation of strain JL117

with the genus *Pseudomonas*. JL117 was most closely related to *P. neuropathica* P155[T]

with 97% bootstrap support (Figure 1). The top average nucleotide identity (ANI) value of

JL117 was 93.2% with *P. crudilactis* UCMA 17988[T]. This value was below the threshold

for species demarcation (Richter and Rosselló-Móra 2009; Chun et al. 2018) providing

further genomic support that strain JL117 represents a new genomic species of

Pseudomonas.

Geographic distribution and host range

Geographic distribution and host range of the novel species was inferred by

comparison to congeneric genome assemblies of unidentified strains from GenBank. Hits

to *Pseudomonas zeiradicis* sp. nov. strain PI116 included assembly GCF_016756945.1

from water of the western basin of Lake Erie (ANI: 99.2%; query coverage: 91.1%),

assembly GCF_001006135.1 from urban soil in Iowa (ANI: 99.2%; query coverage:

91.3%), and assembly GCF_001976065.1 from compost in Massachusetts (ANI: 99.1%;

query coverage: 92.6%). Hits to *Pseudomonas soyae* sp. nov. strain JL117 included

assembly GCF_002003425.1 from the rhizosphere of *Glycine max* in Wisconsin (ANI:

96.2%; query coverage: 92.4%) and assembly GCF_003732405.1 from soil in Wyoming

(ANI: 96.1%; query coverage: 93.3%). Known geographic distributions of the novel

species from cultures is illustrated in Figure 2 and Figure 3, and substrates are compiled in

Table 1.

Morphology, physiology and biochemical characteristics

Pseudomonas zeiradicis sp. nov. strain PI116 and *Pseudomonas soyae* sp. nov.

strain JL117

Strains PI116 and JL117 stained Gram-negative and had a rod shape (Figure 4,

Figure 5). Growth was observed after 2 days on Tryptic soy (TSA) and R2A agars, with

abundant biomass obtained after 5 days, at 30°C. No growth was observed at 37°C on any

of the media tested. On TSA, colonies had a beige-yellow color, changing to opaque-white

on R2A agar. Colonies were circular with regular edges, and smooth with a mucoid

texture. An oxidative metabolism was recorded for both strains with growth at pH 5 and 6.

In addition, the strains grew in 1 and 4% NaCl, but not at 8%. Catalase and oxidase were

produced by the two bacteria, but urease activity was only observed for strain PI116.

Physiological tests using Biolog GenIII Microplates showed that the new bacteria use a

variety of organic compounds as carbon sources. These results are found in Table 1 and

the species description.

Strain PI116 showed resistance to lincomycin, troleandomycin, rifamycin RV,

vancomycin, nalidixic acid and aztreonam, and susceptibility to minocycline according to

Biolog GenIII microplate assays. The strain could grow in the presence of guanidine HCl,

niaproof 4, but not in lithium chloride or potassium tellurite.

Production of urease and the utilization of D-galacturonic acid, D-glucuronic acid,

glucuronamide and α-keto-butyric acid (Table 1) can be used to differentiate strain PI116

from its closest phylogenetic neighbor, *P. umsongensis* Ps 3-10^T.

Similar to strain PI116, JL117 also had the same antibiotic resistance profile except

that in addition to minocycline, it was susceptible to aztreonam. Growth of strain JL117

was inhibited by lithium chloride but not by potassium tellurite guanidine HCl or niaproof

4.

Strain JL117 can be differentiated from its closest phylogenetic neighbors *P.*

neurophatica P155^T and *P. crudilactis* UCMA 17988^T by a combination of several

physiological characteristics such as growth at 37°C, pH 5 and 6, and urease production.

Several carbon source substrates that include trehalose, sucrose, glycerol, L-histidine and

formic acid can also help to differentiate between strain JL117 and these species (Table 1).

DESCRIPTION OF *PSEUDOMONAS ZEIRADICIS* SP. NOV. STRAIN

PI116.

Pseudomonas zeiradicis sp. nov. ze.i.ra'di.cis. L. fem. N. *zea* spelt, also the genus name of corn, *Zea mays;* L. fem. N. *radix* root; N.L. gen. n. *zeiradicis* of corn root

Motile, short rod-shaped cells (1-2 μm long x 0.2-0.4 μm wide). Gram-stain negative. Colonies are opaque white to beige yellow, round, and mucoid. Aerobic and mesophilic. Catalase, oxidase, and urease positive. Growth is abundant at 30°C after 5 days on TSA and R2A agars, but not at 37°C. Growth is also observed at pH 5-7 and up to 4% NaCl. The following substrates are used as carbon sources: α-D-glucose, D-mannose, D-fructose, D-galactose, sodium lactate (1%), fusidic acid, DL-serine, D-mannitol, D-arabitol, glycerol, L-alanine, L-arginine, L-glutamic acid, L-pyroglutamic acid, D-glucuronic acid, D-gluconic acid, glucuronamide, mucic acid, quinic acid, D-saccharic acid, methyl pyruvate, L-lactic acid, citric acid, α-keto-glutaric acid, DL-malic acid, bromo-succinic acid, Tween 40, γ-amino-butyric acid, β-hydroxy-D,L-butyric acid, propionic acid, acetic acid and formic acid. Does not use the following sugars as carbon sources: dextrin, D-maltose, D-trehalose, D-cellobiose, gentibiose, sucrose, D-turanose, stachyose, D-raffinose, α-D-lactose, D-melibiose, β-methyl-D-glucoside. D-salicin, N-acetyl-D-glucosamine, N-acetyl-β-D- mannosamine, N-acetyl-D-galactosamine, N-acetyl neuraminic acid, 3-methyl-glucose, DL-fucose, L-rhamnose and inosine. Does not hydrolyze gelatin or pectin.

The type strain, PI116^T, was isolated from the roots of healthy field-grown *Zea mays* in Missouri, United States.

DESCRIPTION OF *PSEUDOMONAS SOYAE* SP. NOV. STRAIN JL117.

Pseudomonas soyae sp. Nov., so'yae. N.L. gen. N. *soyae*, of soybean, referring to the isolation source of the strain.

Motile, short rod-shaped cells (1-2 μm long x 0.2-0.4 μm wide). Colonies are opaque white to beige yellow, round, and mucoid. Gram-stain negative. Growth is

abundant at 30°C after 5 days on TSA and R2A agars, but not at 37°C. Aerobic and

mesophilic. Catalase and oxidase positive; urease negative. Grows at pH 5-7 and in the

presence of 4% NaCl but not at 8%. The following substrates are used as carbon sources:

D-trehalose, sucrose, N-acetyl-D-glucosamine, α-D-glucose, D-mannose, D-fructose, D-

galactose, sodium lactate (1%), fusidic acid, DL-serine, D-mannitol, D-arabitol, glycerol,

L-alanine, L-arginine, L-glutamic acid, L-pyroglutamic acid, D-gluconic acid,

glucuronamide, mucic acid, quinic acid, D-saccharic acid, methyl pyruvate, L-lactic acid,

citric acid, α-keto-glutaric acid, L-malic acid, Tween 40, γ-amino-butyric acid, β-hydroxy-

D,L-butyric acid, propionic acid, and acetic acid. Does not use the following sugars as

carbon sources: dextrin, D-maltose, D-cellobiose, gentibiose, D-turanose, stachyose, D-

raffinose, α-D-lactose, D-melibiose, β-methyl-D-glucoside. D-salicin, N-acetyl-D-

glucosamine, N-acetyl-β-D- mannosamine, N-acetyl-D-galactosamine, N-acetyl

neuraminic acid, 3-methyl-glucose, DL-fucose, and L-rhamnose. Does not hydrolyze

gelatin or pectin.

The type strain, JL117^T, was isolated from *Glycine max* in Indiana, United States.

ACKNOWLEDGEMENTS

We would like to thank Professor Aharon Oren, The Hebrew University of

Jerusalem, for checking Latin species names.

LITERATURE CITED

Bakker, P. A., Pieterse, C. M., and Van Loon, L. 2007. Induced systemic resistance by
fluorescent *Pseudomonas* spp. Phytopathology. 97:239–243.
Chun, J., Oren, A., Ventosa, A., Christensen, H., Arahal, D. R., da Costa, M. S., et al.
2018. Proposed minimal standards for the use of genome data for the taxonomy of
prokaryotes. Int. J. Syst. Evol. Microbiol. 68:461–466.
Ciufo, S., Kannan, S., Sharma, S., Badretdin, A., Clark, K., Turner, S., et al. 2018. Using
average nucleotide identity to improve taxonomic assignments in prokaryotic
genomes at the NCBI. Int. J. Syst. Evol. Microbiol. 68:2386–2392.
Couillerot, O., Prigent-Combaret, C., Caballero-Mellado, J., and Moënne-Loccoz, Y. 2009.
Pseudomonas fluorescens and closely-related fluorescent pseudomonads as
biocontrol agents of soil-borne phytopathogens. Lett. Appl. Microbiol. 48:505–
512.
De Vleesschauwer, D., Djavaheri, M., Bakker, P. A., and Höfte, M. 2008. *Pseudomonas
fluorescens* WCS374r-induced systemic resistance in rice against *Magnaporthe*

oryzae is based on pseudobactin-mediated priming for a salicylic acid-repressible
multifaceted defense response. Plant Physiol. 148:1996–2012.
Duman, M., Mulet, M., Altun, S., Saticioglu, I. B., Gomila, M., Lalucat, J., et al. 2021.
Pseudomonas piscium sp. nov., *Pseudomonas pisciculturae* sp. nov.,
Pseudomonas mucoides sp. nov. and *Pseudomonas neuropathica* sp. nov. isolated
from rainbow trout. Int. J. Syst. Evol. Microbiol. 71:004714.
Gerbore, J., Benhamou, N., Vallance, J., Le Floch, G., Grizard, D., Regnault-Roger, C., et
al. 2014. Biological control of plant pathogens: advantages and limitations seen
through the case study of *Pythium oligandrum*. Environ. Sci. Pollut. Res.
21:4847–4860.
Hesse, C., Schulz, F., Bull, C. T., Shaffer, B. T., Yan, Q., Shapiro, N., et al. 2018.
Genome-based evolutionary history of Pseudomonas spp. Environ. Microbiol.
Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., et al.
2004. Versatile and open software for comparing large genomes. Genome Biol.
5:R12.
Kwon, S. W., Kim, J. S., Park, I. C., Yoon, S. H., Park, D. H., Lim, C. K., et al. 2003.
Pseudomonas koreensis sp. nov., *Pseudomonas umsongensis* sp. nov. and
Pseudomonas jinjuensis sp. nov., novel species from farm soils in Korea. Int. J.
Syst. Evol. Microbiol. 53:21–27.
Langmead, B., and Salzberg, S. L. 2012. Fast gapped-read alignment with bowtie 2. Nat
Methods. 9 Available at: https://doi.org/10.1038/nmeth.1923.
Lee, M. D. 2019. Applications and considerations of GToTree: a user-friendly workflow
for phylogenomics. Evol. Bioinforma. 15:1176934319862245.
Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. 2015. MEGAHIT: an ultra-fast
single-node solution for large and complex metagenomics assembly via succinct
de Bruijn graph. Bioinformatics. 31:1674–1676.
Mitter, B., Brader, G., Afzal, M., Compant, S., Naveed, M., Trognitz, F., et al. 2013.
Advances in elucidating beneficial interactions between plants, soil, and bacteria.
Adv. Agron. 121:381–445.
Pebesma, E. 2018. Simple features for R: standardized support for spatial vector data. R J.
10:439–446.
R Core Team. 2021. *R: A language and environment for statistical computing*. R
Foundation for Statistical Computing, Vienna, Austria. Available at:
https://www.r-project.org/.
Richter, M., and Rosselló-Móra, R. 2009. Shifting the genomic gold standard for the
prokaryotic species definition. Proc. Natl. Acad. Sci. 106:19126–19131.
Sahlin, K., Vezzi, F., Nystedt, B., Lundeberg, J., and Arvestad, L. 2014. BESST-efficient
scaffolding of large fragmented assemblies. BMC Bioinformatics. 15:281.
Schlusselhuber, M., Girard, L., Cousin, F. J., Lood, C., De Mot, R., Goux, D., et al. 2021.
Pseudomonas crudilactis sp. nov., isolated from raw milk in France. Antonie Van
Leeuwenhoek. 114:719–730.
Seemann, T. 2019. *barrnap 0.9: rapid ribosomal RNA prediction*. Available at:
https://github.com/tseemann/barrnap.
Stanier, R. Y., Palleroni, N. J., and Doudoroff, M. 1966. The aerobic pseudomonads a
taxonomic study. Microbiology. 43:159–271.
Tarnawski, S., Hamelin, J., Locatelli, L., Aragno, M., and Fromin, N. 2003. Examination
of Gould's modified S1 (mS1) selective medium and Angle's non-selective
medium for describing the diversity of *Pseudomonas* spp. in soil and root
environments. FEMS Microbiol. Ecol. 45:97–104.

Vančura, V. 1980. Fluorescent pseudomonads in the rhizosphere of plants and their
relation to root exudates. Folia Microbiol. (Praha). 25:168–173.
Walker, K. 2021. *tigris: Load Census TIGER/Line Shapefiles*. Available at:
https://CRAN.R-project.org/package=tigris.
Wickham, H. 2016. *ggplot2: elegant graphics for data analysis*. New York: Springer
Verlag. Available at: https://ggplot2.tidyverse.org.
Wilke, C. O. 2020. *cowplot: streamlined plot theme and plot annotations for "ggplot2."*
Available at: https://CRAN.R-project.org/package=cowplot.

Figure 1. Phylogenomic tree of *Pseudomonas zeiradicis* sp. nov. strain PI116 and

Pseudomonas soyae sp. nov. strain JL117 and relatives inferred by GToTree (Lee 2019).

The species with the highest ANI to the new *Pseudomonas* species were included in the

tree. Alignment consists of 38,231 amino acid positions from 160 – 170 genes depending

on taxon. Species described in this study are in bold. Strain numbers and GenBank

accession numbers follow species names, T stands for 'type'. Support values above 50%

are given by the branches. *Pseudomonas zeiradicis* sp. nov. strain PI116 is most closely

related to *P. umsongensis* and *Pseudomonas soyae* sp. nov. strain JL117 to *P.*

neuropathica, with 100% and 97% support, respectively. Branch lengths are proportional

to the changes along the branches, a scale bar is provided at the bottom.

Figure 2. Geographic distribution of *Pseudomonas zeiradicis* sp. nov. strain PI116

based on culturing. Dark blue indicates state of origin for type strain, light blue are states

of origin for additional conspecific strains based on GenBank genome assemblies. The

strain isolated from Lake Erie water by the University of Toledo was assigned to Ohio. See

text for details.

Figure 3. Geographic distribution of *Pseudomonas soyae* sp. nov. strain JL117 based on culturing. Dark blue indicates state of origin for type strain, light blue indicates states of origin for additional conspecific strains based on GenBank genome assemblies.

Figure 4. Morphology of *Pseudomonas zeiradicis* sp. nov. strain PI116 depicted

following Gram stain using bright field microscopy. Bar = 10 µm.

Figure 5. Morphology of *Pseudomonas soyae* sp. nov. strain JL117 depicted

following Gram stain using bright field microscopy. Bar = 10 µm.

TABLES

Table 1. Substrates of novel species based on culturing.

Species	Substrate[1]
Pseudomonas zeiradicis sp. nov. strain PI116	Compost, lake water, urban soil, *Zea mays*
Pseudomonas soyae sp. nov. strain JL117	*Glycine max*, soil

339 [1] No disease symptoms reported for strains isolated from plants.

341 **Table 2.** Physiological characteristics of newly described *Pseudomonas zeiradicis* sp.

342 nov. strain PI116 and *P. soyae* sp. nov. strain JL117 and related *Pseudomonas* type strains.

	P. zeiradicis PI116[T]	*P. soyae* JL117[T]	*P. crudilactis* UCMA 17988[T]	*P. neuropathica* P155[T]	*P. umsongensis* Ps 3-10[T]	*P. reinekei* MT-1[T]
General Characteristics:						
Colony Morphology	Beige-yellow, circular, regular edges, mucoid	Beige-yellow, circular, regular edges, mucoid	Beige, circular, convex, regular edges, mucoid	Opaque, beige, circular, regular edges, mucoid	White-yellow, circular	White-yellow, circular, mucoid
Growth at 37°C	-	-	-	+	-	+
Growth at pH 5	+	+	-	-	nr	
Growth at pH 6	+	+	+	-	nr	
Enzymatic Activity:						
Oxidase	+	+	+	+	+	+
Catalase	+	+	+	nr	+	nr
Urease	+	-	+	nr	-	-
Utilization of:						
D-Maltose	-	-	+	-	-	-
D-Trehalose	-	+	-	+	-	-
Sucrose	-	+	-	-	-	-
N-Acetyl-D-Glucosamine	-	+	+	+	-	-
α-D-Glucose	+	+	+	+	+	+
Inosine	-	+	+	+	-	-
Glycerol	+	+	+	-	+	+
L-Arginine	+	+	+	+	nr	nr
L-Histidine	+	+	-	+	+	-
D-Galacturonic Acid	-	-	+	-	+	+
D-Glucuronic Acid	+	-	-	w	-	-
Glucuronamide	+	+	+	+	-	-
Formic Acid	+	+	+	-	+	+
Propionic Acid	+	+	-	+	+	w
Acetic Acid	+	+	-	+	+	+
α-Keto-Glutaric Acid	+	+	-	+	+	+
α-Keto-Butyric Acid	-	-	-	-	+	+

344
345 +, Positive; w, Weakly positive; -, Negative; nr, Not reported.

346
347 Data from this study and Cámara et al. (2021), Duman et al. (2021), Kwon et al. (2003)
348 and Schlusselhuber et al. (2021).