Title: A flexible method for estimating tip diversification rates across a range of speciation and
extinction scenarios

Authors: Thais Vasconcelos[1], Brian C. O'Meara[2], Jeremy M. Beaulieu[1]

[1] Department of Biological Sciences, University of Arkansas, Fayetteville, AR 72701, USA.

[2] Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN
37996, USA.

Abstract

Estimates of diversification rates at the tips of a phylogeny provide a flexible approach for
correlation analyses with multiple traits and to map diversification rates in space, while also
avoiding the uncertainty of deep time rate reconstructions. Available methods for tip rate
estimation make different assumptions, and thus their accuracy usually depends on
characteristics of the underlying model generating the tree. Here we introduce MiSSE, a trait-
free, state-dependent speciation and extinction approach that can be used to estimate varying
speciation, extinction, net-diversification, turnover rates, and extinction fraction at the tips of the
tree. We compare the accuracy of tip rates inferred by MiSSE against similar methods and
demonstrate that, due to certain characteristics of the model, the error is generally low across a
broad range of speciation and extinction scenarios. MiSSE can be used alongside regular
phylogenetic comparative methods in trait related diversification hypotheses, and we also
describe a simple correction to avoid pseudoreplication from sister tips in analyses of
independent contrasts. Finally, we demonstrate the capabilities of MiSSE, with a renewed focus
on classic comparative methods, to examine correlation between plant height and turnover rates
in eucalypts, a species-rich lineage of flowering plants.

Key words: diversification, *Eucalyptus*, hidden-states, plant height, turnover, state-dependent
speciation and extinction.

INTRODUCTION

Molecular phylogenies, when scaled in relation to time, are powerful sources of data to
understand the diversification dynamics of organisms and have become crucial in multiple areas
of ecology and evolution (Wiens & Donoghue, 2004). As the trees themselves become bigger,
more robust, and increasingly comprehensive (e.g. Beaulieu & O'Meara, 2018; Smith & Brown,
2018), the statistical tools used to infer macroevolutionary patterns from them also become

more biologically realistic (e.g. Beaulieu & O'Meara, 2016). Modelling the dynamics of lineage
origination and extinction through time have allowed us to understand, for example, how shifts
in pollination and dispersal strategies are connected to changes in diversification rates of
angiosperms (e.g. Lagomarsino et al., 2016; Vasconcelos et al., 2019; Reginato et al., 2020) or
the role of environmental instability in the diversification of several vertebrate groups (e.g.
Harvey et al., 2020; Morales-Barbero et al., 2021).

Recently, however, a renewed wave of criticisms regarding these methods calls into
question whether diversification rates from time-calibrated trees of extant-only organisms should
even be estimated at all (Louca & Pennell, 2020). While it is true that phylogenies are often
used to address problems beyond their capabilities (Losos, 2011; Cooper et al., 2016; Uyeda et
al., 2018), there is still a considerable amount of information that extant-only phylogenies can
provide about the diversification process (Helmstetter et al., 2021; Morlon et al., 2022). For
instance, O'Meara and Beaulieu (2021) demonstrated that state-dependent speciation and
extinction (SSE) models can identify different likelihoods for the generating parameters of trees
with different topologies but identical lineage through time plots. However, they also discuss
how the uncertainty around parameter estimates increases as one moves from tip to root in the
tree, due to the decreasing amount of information in ancestral rate reconstructions (O'Meara
and Beaulieu, 2021).

A solution, then, could be to continue to model diversification dynamics, but focus only
on rates estimated near the present or at the tips of the tree, rather than deep in time. Tip rate
estimates, also referred to as species-specific diversification rates (Maliet et al., 2019), have
additional advantages in their flexibility when testing for correlations between multistate discrete
and continuous traits (Harvey & Rabosky, 2018; Title & Rabosky, 2019) and for providing a
straightforward way to map diversification rates in space (e.g. Sun et al., 2020; Suissa et al.,
2021). Several methods for estimating tip diversification rates have been proposed in the last
decade (e.g. Jetz et al., 2012; Rabosky 2014; Maliet et al., 2019) and they tend to provide
different levels of accuracy for the estimates depending on the underlying model generating the
tree (Title & Rabosky, 2019). There are also multiple views on how to use tip rate estimates in
trait-based correlation analyses, depending on how the tip rates themselves were estimated
(Freckleton et al., 2008; Rabosky and Huang, 2016; Maliet et al., 2019). Of course, the
underlying processes generating empirical phylogenies are unknown, and therefore developing
a flexible method that estimates accurate tip rates under a broad range of speciation and
extinction scenarios is desirable.

Here we formally describe MiSSE ("Missing State Speciation and Extinction"), an

extension of the SSE framework that provides accurate estimates of various metrics of

diversification rates at the tips of a tree under various speciation and extinction scenarios. We

also show how MiSSE estimates can be used alongside regular phylogenetic comparative

methods after a simple correction for pseudoreplication of tip rates, contributing to its flexibility in

tip correlation analyses. We compare the accuracy of MiSSE against other popular approaches

for estimating tip diversification rates, and further demonstrate its capabilities with an empirical

example that examines the correlation between turnover rates and plant height in eucalypts

(Eucalypteae, Myrtaceae), a diverse lineage of flowering plants. Finally, we argue why focusing

on tip estimates can be advantageous in testing complex hypotheses of diversification and

discuss some caveats of this approach and possible ways forward for modeling diversification.

MATERIALS AND METHODS

The MiSSE model

State-dependent speciation and extinction models expand the birth-death process to

account for speciation (λ), extinction (μ), and trait evolution (q, the transition rates between

character states), estimating parameters that maximize the likelihood of observing both the

character states at the tips of the tree and the tree itself (Maddison et al., 2007). SSE models

were initially developed to overcome three main perceived shortcomings in the field: (1) the

need for greater flexibility in tests of key-innovation hypotheses, which, at the time, typically

involved sister-clade comparisons that could only measure differences in net-diversification and

assumed constant rates within the clades under comparison (Barraclough et al., 1998); (2) the

need to incorporate differences in speciation and/or extinction rates associated with a particular

character state in analyses of trait and geographical range evolution (Goldberg et al., 2010;

Goldberg & Igic, 2012; Ree & Sanmartin, 2018); and (3) the potentially confounding effects of

different transition and diversification rates when looking at diversification or character evolution,

respectively (Maddison, 2006).

All discrete SSE models exist within the following generalized ordinary differential

equations (corresponding to eq. 1a-b in FitzJohn, 2012):

$$\frac{dE_i}{dt} = \mu_i - \left(\lambda_i + \mu_i + \sum_{i \neq j} q_{ij}\right) E_i(t) + \lambda_i E_i(t)^2 + \sum_{j \neq i} q_{ij} E_j(t)$$

96 (1a)

$$\frac{dD_i}{dt} = -(\lambda_i + \mu_i + \sum_{i \neq j} q_{ij}) D_i(t) + 2 \lambda_i D_i(t) E_i(t) + \sum_{j \neq i} q_{ij} D_j(t)$$

97   (1b)

The probability $E_i(t)$ is the probability that a lineage starting at time t in state i leaves no

descendants at the present day ($t = 0$), and $D_i(t)$ is the probability of a lineage in state i at time t

before the present ($t > 0$) evolved the exact branching structure as observed.

With these ordinary differential equations, any number of states can be included in an

SSE model. For example, in character-based models, such as BiSSE ("Binary-State Speciation

and Extinction"; Maddison et al., 2007), i and j represent two observed states of a focal

character (e.g. observed states 0 and 1). A potential issue in this case is that when comparing a

simple model where there is no variation in rates among states against a model of trait-

dependent diversification there is almost always strong support for a trait-dependent

diversification process in empirical settings (Maddison & Fitzjohn, 2014; Rabosky & Goldberg,

2015). The HiSSE model ("Hidden-State Speciation and Extinction"; Beaulieu & O'Meara, 2016)

partially corrects this issue by harnessing the properties of hidden Markov models to allow rate

heterogeneity to depend not only on the focal trait, but also to be correlated with other factors

that were not explicitly scored as character observations at the tips (see also Caetano et al.,

2018; Nakov et al., 2019; Boyko & Beaulieu, 2021, 2022). Thus, with HiSSE, i and j represent

the different observed and hidden states combinations specified in the model (e.g. observed

states 0 and 1, hidden states *A* and *B*).

Although hidden state SSE models made hypothesis testing in diversification studies

more realistic, existing SSE models are still generally used to understand the correlation of a

particular observed trait or geographical range distribution and the diversification dynamics of a

group. However, in reality, it is not a single factor but, rather, a combination of circumstances

that are responsible for the heterogeneous diversification rates among clades in a phylogenetic

tree (Donoghue & Sanderson, 2015; Nürk et al., 2020). To understand heterogeneous

processes that arise from the effect of multiple traits on the dynamics of speciation, extinction,

and trait evolution is one of the main utilities of hidden Markov models in phylogenetic

comparative methods (Caetano et al., 2018).

It is natural, then, to completely drop the observed trait from the analysis and focus only

on the impact of the "unobserved" traits, or the hidden states, in the diversification dynamics of a

clade. This is what our MiSSE model, an extension of the HiSSE framework, is intended to do

(see also Barido-Sottani et al. 2020 for a similar SSE extension in a Bayesian framework).

MiSSE is a direct extension of HiSSE with the main difference between the two models being
that with HiSSE, we have an observed character with states 0 or 1, as in BiSSE (Fig.1a), *and*
hidden states *A* and *B* (Fig. 1b). Essentially, MiSSE operates in the same way, but simply
ignores the observed states altogether and performs the calculations of the hidden rate classes
directly (Fig.1c).

135
Figure 1: Diagrammatic representations of state-dependent speciation and extinction models
that: (a) use only observed trait data in parameter estimation (e.g. BiSSE), (b) use both
observed trait data and hidden-states in parameter estimation (e.g. HiSSE, where observed and
hidden are coded as numbers and letters and are combined to yield a four-state system); and
(c) use only hidden states in parameter estimation (e.g. MiSSE).

The assignment of particular hidden state to each tip is based on probabilities of the
data, which, in the case of MiSSE, is just the structure of the tree. This process is like the
maximum likelihood calculations in other hidden state methods. These calculations are

described thoroughly in Caetano et al. (2018), so here we provide only a brief overview. The
first step is to search for parameter values that maximize the probability of observing the tree at
the root by marginalizing over all possible assignment of rate classes at internal nodes and
along branches. This search consists of a bounded stochastic simulated annealing algorithm,
run for 5000 iterations, followed by a bounded subplex routine that runs until the maximum
likelihood is found. Alternating between a stochastic optimization routine, followed by a "greedy"
hill climbing routine, helps ease MiSSE away from finding local optima.

Suppose that, in an iteration, rate class A has $\lambda = 0.1$ and $\mu = 0.05$, and rate class B has
$\lambda = 0.2$ and $\mu = 0.1$. The node subtending the branch subtending a tip could have started in A
giving rise to an A or B, or it could have started in B giving rise to an A or a B. Each of these
scenarios has a probability associated with them. The sister edge has the same set of scenarios
and its own set of probabilities. These probabilities are combined at the nodes (which includes
the speciation rate to account for the speciation event at a node) and carried down the tree in
the same way as in other SSE models. Once at the root, marginal probabilities for whether the
root is in A or B based on a set of rates can be calculated, as well as the overall log-likelihood
that these rates produced the observed tree.

At this point, all other branches, nodes, and tips are assumed to be in all possible
states. The second step, then, is to take the MLE of the rates and determine which of the states
at a given node and given tip is more likely than any of the other possible states. For that,
MiSSE simply chooses a node (or tip), fixes it to be in rate class A, traverse the tree, and
calculates the overall likelihood. Then it does the same for that same node, but this time it fixes
it to be in rate class B. At the end, MiSSE calculates the marginal probabilities by dividing the
probability that a node was in each state divided by the sum of the probabilities across all
states. In that way, every tip and node have a probability of being in both states A and B, but
often one state will be more probable than the other. For example, in a clade where speciation
occurred more rapidly, the higher λ associated with B will be a better fit to the shape of that part
of the tree, and so the marginal probabilities will reflect higher support for B. In another clade,
where speciation is slower, lower λ will be the better fit, so the marginal probabilities will reflect
the higher support for hidden state A in that part of the tree. And in other clades they may be
uncertain, given frequent transitions in and out each hidden state.

Note that A and B are arbitrary labels that can and will shift positions at the tips in
different runs. What really matters is the parameter combination that underlies each label in
each run, which should not change their MLE between runs. Note too that MiSSE is still a model
of tree and character states like other SSE models, and so the topology matters to assign tips to

the correct (hidden) states. Because it uses data from the topology as well as branch lengths,
MiSSE likelihood can distinguish between trees that have the same lineage through time plot,
but different topologies (O'Meara & Beaulieu, 2021). Also note that, contrary to previous SSE
models, transition rates (q) among rate classes are always set equal. They are informed in the
output, but they have no direct interpretable biological meaning like the transition rates in
BiSSE, which represent the frequency of state changes among observed character states, or in
GeoSSE, which can be interpreted as the frequency of dispersal out of a biogeographic area.
Our current implementation will continue to treat these rates as fixed until it is clearer whether
transition rates are practically feasible in MiSSE. An implementation where transition rates are
allowed to vary would be ideal given that shifts between rate classes are likely to occur at
different speeds throughout the evolution of a group. Differential diversification rates have a
major effect on the tree (since this affects exponential growth of clades) while differential
transition rates are likely (but not guaranteed) to have a less substantial effect, so for now we
use the data only to estimate differences in the former.

MiSSE is available within the R package *hisse* (Beaulieu & O'Meara, 2016). Some
details of MiSSE's implementation differ from other SSE models also implemented in *hisse*.
These differences are summarized at SM1 (Supplementary Material) and readers are also
encouraged to follow the example code available as SM2 when using MiSSE for their empirical
analyses.

**Comparing the accuracy of tip rate metrics across different speciation and extinction
scenarios**

We compare the accuracy of tip diversification rates estimated by popular methods using
a set of simulated trees extracted from Title and Rabosky (2019) and Maliet et al. (2019). Title
and Rabosky (2019) used 5200 trees compiled from previous literature and simulated under
eight different models as a benchmark set to test the accuracy of different tip rate metrics
across a range of speciation and extinction scenarios. For speed, we selected a random sample
of 10% of their original set of simulations, i.e. 521 trees, for our analyses. We then excluded
trees simulated under the "multi-regime, constant-rate birth–death" of Meyer and Wiens (2018)
because those were represented by only two trees in our set. From Maliet et al. (2019), we
extracted their 40 trees simulated under the ClaDS2 scenario for the comparisons, the same
model available in the recent data-augmentation implementation of the ClaDS model (Maliet
and Morlon 2022). We then excluded five trees where the median height of branch lengths was
above a threshold of one trillion units of time because they could be out of the numerical limits

for some methods (e.g. BAMM's MCMC chains struggled to reach convergence in those cases).
Finally, seven additional trees were excluded due to modelling limitations in one of the methods
(see details in SM3; Supplementary Material). The final dataset comprises 547 trees of different
sizes, though most of them are < 500 tips (Table 1).
Six different tip-rate metrics were compared: DR statistics (Redding & Mooers, 2006),
BAMM (Rabosky, 2014), node density (ND; Freckleton et al., 2008), the inverse of terminal
branch length (TB; Steel & Mooers, 2010), ClaDS (Maliet et al. 2019; Maliet and Morlon 2022),
and our MiSSE model. For MiSSE, we summarized tip rates in the following ways: (1) tip-rates
estimated from the model with overall lowest AICc (i.e. the "best" model; $MiSSE_{best}$); and (2) tip-
rates estimated by averaging all models according to their Akaike weight ($MiSSE_{average}$) (see
SM3, Supplementary Material, for details on the settings for other methods).
223
224 **Table 1:** Simulated trees used in comparisons between tip rate metrics.

Simulation model	Number of trees	Tree-size	Source
Speciation rate evolves via diffusion process	120	25 - 662	Rabosky (2010); Beaulieu and O'Meara (2015); Rabosky (2016); Title and Rabosky (2019)
Single- and multi-regime, constant-rate birth–death	18	9 - 3458	Mitchell et al. (2019)
Single-regime, constant-rate birth–death, lambda uniform	100	100	Title and Rabosky (2019)
Single-regime, constant-rate birth–death	10	100	Mitchell and Rabosky (2017)
Single- and multi-regime, constant-rate birth–death	10	100	Moore et al. (2016)
Single-regime, constant-rate birth–death, net diversification uniform	100	100	Title and Rabosky (2019)
Pure birth root regime, 1–4 discrete shifts to diversity-dependent regimes	117	54 - 794	Rabosky (2014); Mitchell and Rabosky (2017)
Single- and multi-regime,	40	10 - 3157	Rabosky et al. (2017)

constant-rate birth–death			
Rate changes at every speciation event, constant extinction-fraction	32	100-105	Maliet et al. (2019)

We followed the same statistical metric as Title and Rabosky (2019), namely, we

compared mean absolute error, given by the formula $\sum_{i=1}^{N_i} |\text{rate}_i - \text{rate}_{TRUE_i}| / N$, and the RMSE

is given by $\sqrt{\sum_{i=1}^{N_i} (\text{rate}_i - \text{rate}_{TRUE_i})^2 / N}$, where *i* is a single tip and N is the total number of tips

in all trees of a given type of simulation. In all cases, the lower the error values the more

accurate is the metric. We assessed the accuracy of five parameters: speciation (λ), extinction

(μ), and the orthogonal transformations of these, namely, net-diversification (r = λ - μ), turnover

(τ = λ + μ) and extinction fraction (ε = μ / λ). Note that TB, ND and DR are non-parametric (i.e.

not based on an underlying model) and only estimate λ. We then used the function

posthoc.kruskal.conover.test from the R package PMCMR (Pohlert, 2014) to perform a pairwise

test for multiple comparisons of mean rank sums and calculate whether errors are significantly

different between metrics, assuming a significance value of p < 0.05.

Empirical example: plant height and turnover in Eucalypts

We demonstrate the capabilities of MiSSE estimates for analyses of tip correlation with

an empirical example. Body size is considered an important trait in studies of animal evolution

(Cooper & Purvis, 2010) and it has been interpreted as a potential correlate of diversification

rates in vertebrates (Cope's rule; FitzJohn, 2010). A similar argument can be made for plants

(Boucher et al., 2017). Through a complex link between rates of molecular evolution, fecundity,

and population size, life span and generation time are expected to be negatively correlated with

the number of speciation and extinction events on a per time basis (Stebbins, 1974; Petit &

Hampe, 2006); i.e. slower turnover rates. Although these correlations are expected in theory,

surprisingly few studies have compared them in practice using model-based approaches (e.g.

Boucher et al., 2020). The MiSSE framework allows us to easily test this correlation by using

regular comparative phylogenetic methods and tip-rates as a response variable.

We examined turnover rates in relation to plant height, a proxy for lifespan (Westoby,

1998), in eucalypts, one of the most distinctive components of the Australian landscape (Wilson

et al., 2005). Eucalypts are members of the tribe Eucalypteae in the flowering plant family

Myrtaceae, a group that includes some of the tallest trees in all angiosperms (e.g. *Eucalyptus*

regnans can reach up to 120 m height) (Wilson et al., 2005; Nevill et al., 2010). Eucalypts are

also unusual for being considered a relatively speciose (c. 800 species) lineage of trees. It is

thought that the tree habit tends to decrease speciation rates, which is offered as reason that

clades comprised of predominantly tree growth forms are frequently found to be less diverse

than their herbaceous or shrubby relatives (Petit & Hampe, 2006). However, many eucalypts

species are also large shrubs to treelets between 1.5 and 5m height (EUCLID, 2015), which led

to the hypothesis that the radiation is driven primarily by the smaller representatives of the clade

(Petit & Hampe, 2006). Here we test this idea by contrasting tip turnover rates with plant height

in this diverse clade of angiosperms.

We used the "ML1" time calibrated phylogenetic tree from Thornhill et al. (2019), which

covers 716 out of the c. 800 species of eucalypts. We then collected data on plant height for

673 species available at EUCLID Eucalypts of Australia Edition 4 (2015) (EUCLID, 2015). All

measurements represent plant maximum height and are given in meters. Again, we use the

default implementation of MiSSE, that is, stop.deltaAICc=10 and chunk.size=10, to estimate tip

turnover rates in Eucalypteae, pruning redundant models with the function

PruneRedundantModels. Next, we estimated tip turnover rates averaging all models according

to their AICc weights with the function GetModelAveRates.

To test correlations between plant height and tip turnover rates, we used the function

TipCorrelation, which performs a regression-through-the-origin of positivized phylogenetic

independent contrasts (PICs, Felsenstein, 1985) between tip rates and a continuous trait. This

function also gives the option of pruning out PICs from two tips that are sister to each other and

share the same branch length to the direct ancestral node (i.e. "cherries"), before regressions.

We reasoned that because these sister tips theoretically inherit the exact same rate class

probabilities in MiSSE they may: (1) present identical tip rates, forcing the slope of the

regression to be close to 0 since their PICs will be 0; and therefore (2) constitute

pseudoreplicates in the analyses. Note that, in this approach, we prune PICs and not individual

sister tips. Pruning tips might not be adequate because it would affect all other PIC calculations

in the tree; it also can make the results sensitive to which tip is pruned and gives less

information for computing contrasts deeper in the tree. All functions mentioned above are

available in the R package *hisse* (Beaulieu and O'Meara, 2016)*.* Rates and maximum plant

height were log scaled before analyses so that they conformed with Brownian motion evolution

(Felsenstein, 1985; Garland et al., 1992).

RESULTS

Simulation studies

Comparisons between tip rate metrics using simulated data show that, in general, MiSSE estimates accurate tip rates across a range of speciation and extinction scenarios (Fig. 2). We present the results for mean absolute error, though RMSE results are practically the same (SM4; Supplementary Material). Our results show that TB is significantly less accurate than other methods in all simulation scenarios, and, in eight out of the nine simulation scenarios, there was no significant difference in error measurements between ND and DR. In all simulated scenarios, results show that the model-based approaches MiSSE, BAMM and ClaDS tended to be significantly ($p < 0.01$) more accurate than the non-parametric methods TB, ND, and DR for speciation rates, the only parameter estimated by the latter three metrics (see SM5; Supplementary Material for all pairwise comparisons).

Accuracy among the three model-based approaches for the five diversification parameters estimated at the tips varies depending on the model used to generate the simulated tree. Significant differences in accuracy are observed for extinction fraction in the "speciation rate evolves via diffusion process" scenario (Fig. 2a), where ClaDS is significantly more accurate ($p = 0.04$) than MiSSE$_{average}$ and BAMM, but not than MiSSE$_{best}$ ($p = 0.32$). Significant differences in accuracy between the estimates of the MiSSE$_{best}$ model and the AICc-weighted average of the MiSSE$_{average}$ model were not observed in any other simulated scenario; therefore, the two results are henceforward treated as a single MiSSE category. In both the "single-regime, constant-rate birth–death, lambda uniform" (Fig. 2c) and the "single-regime, constant-rate birth–death, net diversification uniform" (Fig. 2f) scenarios, ClaDS estimates for speciation at the tips are significantly less accurate ($p < 0.01$) than the other model-based approaches. ClaDS estimates also perform significantly poorly than MiSSE ($p < 0.01$ and $p = 0.049$ when compared to MiSSE$_{best}$) for estimates of net-diversification and turnover rates in those scenarios. BAMM and ClaDS are significantly more accurate than MiSSE for speciation in the "Pure birth root regime, 1–4 discrete shifts to diversity-dependent regimes" scenario ($p < 0.01$; Fig. 2g). In the same scenario, ClaDS also tends to be more accurate than other methods for extinction rates ($p < 0.01$) and BAMM for net-diversification ($p < 0.01$), but BAMM is also significantly less accurate for turnover rates ($p < 0.01$) than the other methods in this scenario. ClaDS is significantly more accurate than BAMM for estimates of extinction fraction in trees simulated under the model "rate changes at every speciation event, constant extinction-fraction" ($p < 0.01$; Fig. 2i), but does not significantly differ from MiSSE for any parameter in that scenario.

Figure 2: Boxplots showing the distribution of mean absolute error between true and estimated
tip rates using five previously described metrics and our MiSSE model. Each data point
corresponds to the mean absolute error between all tips in one tree. λ = speciation rate, μ =

extinction rate,   = turnover rate, and r = net-diversification rate,   = extinction fraction. Letters

between brackets highlight significantly different error measurements.

Tip-correlation between plant height and turnover

Our empirical example using tip rate correlations with MiSSE show that, despite the idea

that smaller plants have driven the radiation in eucalypts, maximum plant height appears

uncorrelated with turnover across the tips of the tree ($R^2 < 0.001$, Fig. 3c). The general lack of

correlation is mainly because: (1) higher turnover rates are not always restricted to clades of

small shrubs and treelets, as it can be seeing by comparing the distribution of tip-rates and plant

height (Fig. 3a,b); and (2) plant height (Fig. 3b) appears to be a much more labile trait than

turnover rates, regardless of the methods used to infer tip-rates (Fig. 3a,b). We note that

pruning PICs from sister tips from the tree minimally changes the slope of the regression line in

this example and it does not change our main conclusion that turnover and plant height are

likely uncorrelated in eucalypts, given that contrasts of maximum plant height explain less than

0.1% of the contrasts of turnover. In any event, the inclusion of PICs from sister tips can force

the slope of the PIC regression towards 0 in other tip rate comparisons and therefore they

should be pruned before analyses.

Figure 3: Comparison between maximum plant height and tip turnover rates estimated by

MiSSE in Eucalypteae. (a) MiSSE turnover rates at the tips; (b) Maximum plant height in

meters; warmer colors indicate higher values in both cases. (c) Regression through the origin

between PICs of MiSSE turnover rates and plant height, with PICs from sister tips included and

pruned. Solid line: smoothed line using the loess method; dashed line: regression line.

DISCUSSION

MiSSE as a flexible model-based approach to estimate tip diversification rates

Here, we describe our "trait-free" MiSSE framework and show how it can be used to estimate accurate diversification rates at the tips of the tree. Our results show that the difference in accuracy in parameters estimated by averaging all MiSSE models or using only the best MiSSE model seem to be minimal and may depend on the tree (Fig. 2). However, as discussed in previous publications (Beaulieu & O'Meara, 2016; Caetano et al., 2018), the user is strongly encouraged to focus mainly on parameter estimation rather than which model fits "best" when using MiSSE, since the former is more informative about the biology of a clade. We also note that, although extinction fraction and turnover are sometimes synonymous in the literature (see SM6, Supplementary Material, for a short terminology survey), we use turnover rate as an explicit measure of events per units of time, and a parameter that is explicitly orthogonal to extinction fraction. We emphasize this difference because many diversification hypotheses may be better described by turnover rates rather than net-diversification or extinction fraction (e.g. Vrba, 1993; Vasconcelos et al., 2022; our empirical example). Also, given that extinction and speciation should tend to correlate through time (Marshall, 2017), net-diversification should tend to approach 0. In those scenarios, and as long as extinction is different from 0, turnover can naturally become a better metric for understanding diversification dynamics nuances than its isolated components. We note, however, that when extinction rate is equal to 0, turnover rate collapses into speciation rate (i.e. there is no turnover). We also note that our concept of turnover is different from the one used in the island biogeography literature (e.g. Simberloff 1974).

Key differences between MiSSE and analogous methods are highlighted by our analyses of simulated data. Although estimates from the three model-based approaches, MiSSE, BAMM and ClaDS, tend to provide generally accurate estimates of diversification parameters at the tips, the three methods make very different assumptions. BAMM is based on inferences of discrete rate shifts, so that all tips after a shift inherit a similar rate (Rabosky et al. 2014). ClaDS, on the other hand, assumes that changes in diversification dynamics occur at every speciation event, so that the estimates at the tips are allowed to be highly heterogeneous in relation to one another (Maliet et al. 2019). MiSSE may also have some shift-like properties, as in BAMM, in areas where the probabilities of being in different rate classes change abruptly. However, because at every moment in the tree, including at the tips, there is a combined probability of being in all possible rate classes, the evolution of diversification rates becomes

smoother, allowing it to pick up some rate heterogeneity at the tips, as in ClaDS. Closely related
tips do have similar rates because it is likely that they will be in the same rate class, but they do
not inherit the absolute same rate.

A visual inspection of individual results (Fig.4; all plots in SM7, Supplementary Material)
show that differences in accuracy observed in our comparisons probably result from these
operational differences between model-based approach. For instance, because BAMM works
with discrete shifts, this method tends to be more accurate when the true model assumes that
few large discrete shifts of diversification rates occurred in the tree. In those scenarios, models
like MiSSE and ClaDS may perform comparatively worse because they may infer rate
heterogeneity where there is none (Fig. 4d). On the other hand, BAMM can underestimate rate
heterogeneity when the true model produces highly heterogeneous tip rates in the tree. In those
cases, even if the mean error is low, this method may fail to capture rate variation in small
clades with particularly high or particularly low rates (Fig.4a,c). Similarly, ClaDS may
overestimate variation in scenarios where rate heterogeneity is nonexistent, resulting in higher
error (Fig,4b). Because MiSSE works as an intermediate between BAMM and ClaDS, we argue
that it can become a model-based approach that is more flexible in identifying areas of rate
heterogeneity or homogeneity at the tips of the tree. Importantly, even in scenarios where
MiSSE is less accurate, it tends to capture areas with faster and lower rates in the tree correctly.
In "diversity dependent" scenarios, for example, MiSSE may be less accurate because it can
overestimate speciation rates of clades with higher speciation rates, since it does not explicitly
allow speciation rates to slow down along a given terminal branch (as in BAMM). However, it
tends to be similar to other methods in identifying where variation in rates occurs (Fig. 4d).

Figure 4.: A comparison between rates estimated using different model-based approaches (colored dots, warmer colors indicate higher values) and true rates (gray dots) at the tips of the tree in four different simulated scenarios: (a) "rate changes at every speciation event"; (b) "speciation rate evolves via diffusion process"; (c) "single-regime, constant-rate birth–death, lambda uniform"; (d) "pure birth root regime, 1–4 discrete shifts to diversity-dependent regimes".

Plant height and turnover rates are uncorrelated in eucalypts

Our empirical analysis demonstrated that the correlation between plant height and turnover rates is weak within Eucalypteae. The use of plant height as a proxy for longevity is common in plants, but it may be that plant height is not the best proxy for longevity in eucalypts.

Tall eucalypt trees are often native to productive environments with fertile soils (Pryor, 1976;
Thornhill et al., 2019) and it may be that they present fast life cycles for their size, as observed
in other parts of the world where soils are also fertile (Russo et al., 2008). In fact, many eucalypt
trees are known to be fast-growing plants (e.g. Barnard et al., 2003; Almeida et al., 2004) and
fast growth is linked to higher mortality through the growth-survival trade-off (Reich, 2014),
which could potentially lead to higher turnover rates even in lineages of tall trees (Baker et al.,
2014). If that is the case, perhaps plant height is not the trait that best captures the variation that
we sought to explore in our hypothesis. Other proxies, such as leaf mass area and seed size
(Westoby, 1998; Wright et al., 2004) may be a better proxy of longevity in these scenarios and
should also be tested.

Alternatively, if plant height is, in fact, a good proxy for longevity in eucalypts, it may be
that longevity is generally uncorrelated with turnover in the clade. A similar trend of uncorrelated
body size and speciation rates has been observed in other groups (FitzJohn, 2010; Boucher et
al., 2020), but it is difficult to ascertain if the lack of correlation is a particularity of the groups
that have been analyzed so far or a general rule. Given the heterogeneity of evolution,
identifying what is a rule and what is an exception in macroevolution requires analyses of
multiple natural replicates. Similar tests in several clades would be ideal to rule out this
possibility. A different approach would be to use QuASSE (FitzJohn, 2010) to construct a model
of turnover and height; the disadvantage of this is the need to model the relationship and the
(current) impossibility of including additional observed or unobserved factors.

One benefit of MiSSE (and similar methods) is that, even though our hypothesis was not
supported, we still have new information about the tips: turnover rate (and the other
diversification parameters if one wishes). In the same way that looking at the correlation of two
traits may generate new hypotheses to test (Boyko and Beaulieu, 2021), having estimates of
turnover rate at tips as in Figure 3 may also lead to new ideas about the factors leading to
higher turnover rates in some eucalypts.

Best practices in tip rate estimation

MiSSE is intended to be used to estimate diversification rates at the tips of the tree. Tip
rates are, technically, an emergent property of how diversification dynamics are modeled along
the branches of the tree (Freckleton et al., 2008; Title & Rabosky, 2019). However, rate
estimates are not unique per tip species, so species should not be analyzed in isolation. In other
words, a tip with extinction-fraction of 0.9, for instance, does not necessarily indicate that that tip
is about to go extinct, but rather that a lot of closely related species are relatively short-lived in

comparison to the rest of the tree. Tip rates can then be seen as a "snapshot" of the
diversification potential of the species in a clade. In the MiSSE framework, this diversification
potential may be interpreted as the combined effect of the many (hidden) states evolving in the
tree, which are represented by the different rate classes.

From a mathematical standpoint, there is an advantage in using tip rates because it is at
the tips where the certainty around the probability of being in a particular rate class is the
highest (O'Meara & Beaulieu, 2021). The certainty around these probabilities decreases as one
moves towards the root of the tree, and so the ability to point out changes in rate classes also
decreases. Of course, the model *does* use information from the whole phylogenetic tree in the
calculations of tip-rates, but as one moves towards the present in the tree, the more information
is available to reconstruct those rates and thus we are more likely to be certain about the
probabilities of being in different rate categories (see O'Meara & Beaulieu, 2021).

The intuitive question that follows is how far back in the past one can go to interpret how
the observable diversification potential at the tips came to be and how far into the future can we
use it to make predictions about how clades will diversify. The answer is, again, not
straightforward and it may depend on the size and age of the phylogenetic tree under analysis.
Tip rates will not be able to tell us about mass extinction events deep in the past as these are
arguably only measurable from the fossil record (Barnosky et al., 2011), but they may represent
a good reflection of diversification dynamics above the species level in times slices close to the
present. We therefore recommend great caution with literal interpretations of rate classes deep
in time. That is why although MiSSE can "paint" the averaged rate reconstruction along the
branches of the tree (i.e. with the function plot.misse.states in the R package *hisse*), inferences
of rates through time should not be interpreted literally. The "painting" should be used just to
visualize the rates inferred with MiSSE and should not be used to support narratives of past
diversification dynamics for a group. Interpreting past rates is risky given limited information.
Similarly, modern drivers of diversification dynamics (climate change, invasive species, habitat
destruction, and more), which affect especially extinction rates at the present, will not be
reflected in the rate estimates returned by MiSSE or similar methods.

Another issue is the non-independence of the tip rates. The identical rates for sister tips
have been discussed above, but even non-sister tips may have somewhat correlated rates. It is
important to realize that though a phylogenetic model was used in the estimation of tip rates,
they themselves are not corrected for phylogeny (which is a reason we used independent
contrasts above to compare turnover and plant height). There is also limited information: a
resolved tree of 100 taxa has 198 edges. Though MiSSE, BAMM, ClaDS, and other methods

give 100 tip estimates, there is not enough information to take each one as known with great
certainty, nor to draw a strong conclusion based on a rate shared by a few taxa.

Caveats and persistent issues in modeling diversification dynamics

Tip-diversification rates are advantageous for their flexibility in trait related diversification
analyses and for avoiding the uncertainty of rate reconstructions in deep time. However, there
are several caveats that users should consider when using these methods, including MiSSE: (1)
Clade-specific sampling fraction was found to lead to an incorrect likelihood behavior in *any*
diversification method despite its appeal (Beaulieu, 2020). Contrary to other similar methods,
MiSSE deliberately does not include an implementation for clade-specific sampling fraction, so
all sampling fractions are global and only one sampling fraction is given for the whole tree.
Imputation using stochastic polytomy resolvers can work as an alternative solution when dealing
with incomplete phylogenies (e.g. Chang et al. 2020; but see Rabosky 2015) (2) How species
are defined is the most relevant when looking at tip rates. For diversification rate models, the
data come from the distribution of branching events across the phylogeny. Since most of these
events are nearer the present, lumping and splitting taxonomic entities at the species level will
have a greater impact on tip-rate estimation. We suspect that the methods discussed herein are
particularly sensitive to taxonomic subjectivity. (3) There may be still issues related to
ascertainment bias and underestimation of extinction rates (Beaulieu & O'Meara, 2018). In that
sense, the larger and broader taxonomic sample one uses to test diversification hypotheses, the
more biologically realistic extinction rates will tend to be. Ideally, one would be able to correct
the extinction estimates in smaller trees, but that is still not possible with the current
implementation of MiSSE nor in any other model we are aware of. So, even though extinction
rates are estimated accurately relative to the simulated data, they may still be biologically
unrealistic in small and young trees. (4) Finally, given that these models do not account for
mass-extinction events, it is unclear what impact such events will have on tip-rate estimation in
any approach currently available.

CONCLUSIONS

Time-calibrated phylogenetic trees have been important sources of data to understand
fundamental aspects of the evolution of organisms. MiSSE provides a novel tool to explore this,
and its flexibility in estimating accurate tip rates across a broad range of speciation and
extinction scenarios makes MiSSE a powerful alternative for diversification analyses with correct
likelihoods. There remain many caveats and cautions about its use, but it is an additional tool to

understand diversification processes, including a focus on parameter of perhaps great biological relevance such as turnover rate. We suggest that tip-rates estimated by MiSSE will be useful to several questions that were previously addressed by other SSE models. Its versatility is appealing to explore integrative questions linking traits, geographical range distribution, or both in time slices close to the present.

Author contributions

T.V., B.C.O. and J.M.B. designed the study. J.M.B. led software development, T.V. and B.C.O. led the simulation study and T.V. led the empirical example. T.V. wrote the first draft and all authors contributed to the final writing of the manuscript.

Acknowledgements

The authors thank members of the Beaulieu lab for discussion. The authors are also grateful to D. Rabosky and two anonymous reviewers for comments that greatly improved an earlier version of this manuscript. This work was funded by the National Science Foundation (grants DEB–1916558 and DEB–1916539). The authors have no conflict of interest to declare.

Data Accessibility Statement: All code and data required to reproduce the analyses in this manuscript can be found at https://github.com/bomeara/missecomparison and in the Dryad folder linked to this publication: [ ]

References

Almeida, A. C., Landsberg, J. J., & Sands, P. J. (2004). Parameterisation of 3-PG model for fast-growing *Eucalyptus grandis* plantations. *Forest Ecology and Management*, *193*(1-2), 179–195. https://doi.org/10.1016/j.foreco.2004.01.029

Baker, T. R., Pennington, R. T., Magallon, S., Gloor, E., Laurance, W. F., Alexiades, M., et al. (2014). Fast demographic traits promote high diversification rates of Amazonian trees. *Ecology letters*, *17*(5), 527–536. https://doi.org/10.1111/ele.12252

Barido-Sottani, J., Vaughan, T. G., & Stadler, T. (2020). A multitype birth–death model for Bayesian inference of lineage-specific birth and death rates. *Systematic Biology*, 69(5), 973-986.

Barnard, H. R., & Ryan, M. G. (2003). A test of the hydraulic limitation hypothesis in

fast-growing *Eucalyptus saligna*. *Plant, Cell & Environment*, *26*(8), 1235–1245.

https://doi.org/10.1046/j.1365-3040.2003.01046.x

Beaulieu, J. M., & O'Meara, B. C. (2015). Extinction can be estimated from moderately sized

molecular phylogenies. *Evolution*, *69*(4), 1036–1043. https://doi.org/10.1111/evo.12614

Beaulieu, J. M., & O'Meara, B. C. (2016). Detecting hidden diversification shifts in models of

trait-dependent speciation and extinction. *Systematic biology*, *65*(4), 583–601.

https://doi.org/10.1093/sysbio/syw022

Beaulieu, J. M., & O'Meara, B. C. (2018). Can we build it? Yes we can, but should we use it?

Assessing the quality and value of a very large phylogeny of campanulid angiosperms.

American Journal of Botany, 105(3), 417–432. https://doi.org/10.1002/ajb2.1020

Beaulieu, J. M. (2020) The Problem with Clade-specific Sampling Fractions. Available at:

https://rdrr.io/cran/hisse/f/inst/doc/Clade-specific-sampling.pdf.

Barraclough, T. G., Vogler, A. P., & Harvey, P. H. (1998). Revealing the factors that promote

speciation. *Philosophical Transactions of the Royal Society of London. Series B: Biological*

Sciences, *353*(1366), 241–249. https://doi.org/10.1098/rstb.1998.0206

Boucher, F. C., Verboom, G. A., Musker, S., & Ellis, A. G. (2017). Plant size: a key determinant

of diversification? *New Phytologist*, *216*(1), 24–31. https://doi.org/10.1111/nph.14697

Boucher, F. C., Quatela, A. S., Ellis, A. G., & Verboom, G. A. (2020). Diversification rate vs.

diversification density: Decoupled consequences of plant height for diversification of Alooideae

in time and space. *PloS one*, *15*(5), e0233597. https://doi.org/10.1371/journal.pone.0233597

Boyko, J. D., & Beaulieu, J. M. (2021). Generalized hidden Markov models for phylogenetic

comparative datasets. *Methods in Ecology and Evolution*, *12*(3), 468–478.

https://doi.org/10.1111/2041-210X.13534

Boyko, J., & Beaulieu, J. (2022). A potential solution to the unresolved challenge of false

correlation between discrete characters. EcoEvoRxiv https://doi.org/10.32942/osf.io/e2kj8

Burnham, K.P., & Anderson, D.R. (2002). Model selection and multimodel inference: a practical

information-theoretic approach. New York: Springer.

Chang, J., Rabosky, D. L., & Alfaro, M. E. (2020). Estimating diversification rates on

incompletely sampled phylogenies: theoretical concerns and practical solutions. *Systematic*

Biology, 69(3), 602-611. https://doi.org/10.1093/sysbio/syz081

Caetano, D. S., O'Meara, B. C., & Beaulieu, J. M. (2018). Hidden state models improve

state-dependent diversification approaches, including biogeographical

models. *Evolution*, *72*(11), 2308–2324. https://doi.org/10.1111/evo.13602

Cooper, N., & Purvis, A. (2010). Body size evolution in mammals: complexity in tempo and

mode. *The American Naturalist*, *175*(6), 727–738. https://doi.org/10.1086/652466

Cooper, N., Thomas, G. H., & FitzJohn, R. G. (2016). Shedding light on the 'dark side'of

phylogenetic comparative methods. *Methods in Ecology and Evolution*, *7*(6), 693–699.

https://doi.org/10.1111/2041-210X.12533

Donoghue, M. J., & Sanderson, M. J. (2015). Confluence, synnovation, and depauperons in

plant diversification. *New Phytologist*, *207*(2), 260–274. https://doi.org/10.1111/nph.13367

EUCLID Eucalypts of Australia Edition 4 (2015). Internet based, hosted by the Identic Pty Ltd,

Brisbane at: https://apps.lucidcentral.org/euclid

FitzJohn, R. G. (2010). Quantitative traits and diversification. *Systematic biology*, *59*(6), 619–

633. https://doi.org/10.1093/sysbio/syq053

FitzJohn, R. G. (2012). Diversitree: comparative phylogenetic analyses of diversification in

R. *Methods in Ecology and Evolution*, *3*(6), 1084–1092. https://doi.org/10.1111/j.2041-

210X.2012.00234.x

Felsenstein, J. (1985). Phylogenies and the comparative method. *The American*

Naturalist, *125*(1), 1–15. https://doi.org/10.1086/284325

Freckleton, R. P., Phillimore, A. B., & Pagel, M. (2008). Relating traits to diversification: a simple

test. *The American Naturalist*, *172*(1), 102–115. https://doi.org/10.1086/588076

Goldberg, E. E., Kohn, J. R., Lande, R., Robertson, K. A., Smith, S. A., & Igić, B. (2010).

Species selection maintains self-incompatibility. *Science*, *330*(6003), 493–495.

https://doi.org/10.1126/science.1194513

Goldberg, E. E., & Igić, B. (2012). Tempo and mode in plant breeding system

evolution. *Evolution: International Journal of Organic Evolution*, *66*(12), 3701–3709.

https://doi.org/10.1111/j.1558-5646.2012.01730.x

Harvey, M. G., & Rabosky, D. L. (2018). Continuous traits and speciation rates: Alternatives to
state-dependent diversification models. *Methods in Ecology and Evolution*, *9*(4), 984–993.
https://doi.org/10.1111/2041-210X.12949

Harvey, M. G., Bravo, G. A., Claramunt, S., Cuervo, A. M., Derryberry, G. E., Battilana, J., ... &
Derryberry, E. P. (2020). The evolution of a tropical biodiversity hotspot. *Science*, 370(6522),
1343–1348. https://doi.org/10.1126/science.aaz6970

Helmstetter, A. J., Glemin, S., Käfer, J., Zenil-Ferguson, R., Sauquet, H., de Boer, H., Dagallier,
L. M. J., Mazet, N., Reboud, E.L., Couvreur, T. L. P., Condamine, F. L. (2022). Pulled
diversification rates, lineages-through-time plots and modern macroevolutionary modelling.
Systematic Biology. https://doi.org/10.1093/sysbio/syab083

Lagomarsino, L. P., Condamine, F. L., Antonelli, A., Mulch, A., & Davis, C. C. (2016). The
abiotic and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae). *New
Phytologist*, *210*(4), 1430–1442. https://doi.org/10.1111/nph.13920

Losos, J. B. (2011). Seeing the Forest for the Trees: The Limitations of Phylogenies in
Comparative Biology: (American Society of Naturalists Address). *The American
Naturalist*, *177*(6), 709–727. https://doi.org/10.1086/660020

Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of
diversification histories. *Nature*, *580*(7804), 502–505. https://doi.org/10.1038/s41586-020-2176-
1

Maddison, W. P. (2006). Confounding asymmetries in evolutionary diversification and character
change. *Evolution*, 60(8), 1743–1746. https://doi.org/10.1111/j.0014-3820.2006.tb00517.x

Maddison, W. P., Midford, P. E., & Otto, S. P. (2007). Estimating a binary character's effect on
speciation and extinction. *Systematic biology*, *56*(5), 701–710.
https://doi.org/10.1080/10635150701607033

Maddison, W. P., & FitzJohn, R. G. (2015). The unsolved challenge to phylogenetic correlation
tests for categorical characters. *Systematic biology*, *64*(1), 127–136.
https://doi.org/10.1093/sysbio/syu070

Maliet, O., Hartig, F., & Morlon, H. (2019). A model with many small shifts for estimating
species-specific diversification rates. *Nature ecology & evolution*, 3(7), 1086–1092.
https://doi.org/10.1038/s41559-019-0908-0

Maliet, O., & Morlon, H. (2022). Fast and accurate estimation of species-specific diversification

rates using data augmentation. *Systematic Biology*, 71(2), 353-366.

https://doi.org/10.1093/sysbio/syab055

Marshall, C. R. (2017). Five palaeobiological laws needed to understand the evolution of the

living biota. *Nature Ecology & Evolution*, *1*(6), 1–6. https://doi.org/10.1038/s41559-017-0165

Meyer, A. L., & Wiens, J. J. (2018). Estimating diversification rates for higher taxa: BAMM can

give problematic estimates of rates and rate shifts. *Evolution*, *72*(1), 39–53.

https://doi.org/10.1111/evo.13378

Mitchell, J. S., & Rabosky, D. L. (2017). Bayesian model selection with BAMM: effects of the

model prior on the inferred number of diversification shifts. *Methods in Ecology and*

Evolution, *8*(1), 37–46. https://doi.org/10.1111/2041-210X.12626

Mitchell, J. S., Etienne, R. S., & Rabosky, D. L. (2019). Inferring diversification rate variation

from phylogenies with fossils. Systematic Biology, 68(1), 1–18.

https://doi.org/10.1093/sysbio/syy035

Moore, B. R., Höhna, S., May, M. R., Rannala, B., & Huelsenbeck, J. P. (2016). Critically

evaluating the theory and performance of Bayesian analysis of macroevolutionary

mixtures. *Proceedings of the National Academy of Sciences*, *113*(34), 9569–9574.

https://doi.org/10.1073/pnas.1518659113

Morales-Barbero, J., Gouveia, S. F., & Martinez, P. A. (2021). Historical climatic instability

predicts the inverse latitudinal pattern in speciation rate of modern mammalian biota. *Journal of*

Evolutionary Biology, *34*(2), 339–351. https://doi.org/10.1111/jeb.13737

Morlon, H., Robin, S., & Hartig, F. (2022). Studying speciation and extinction dynamics from

phylogenies: addressing identifiability issues. *Trends in Ecology & Evolution*.

https://doi.org/10.1016/j.tree.2022.02.004

Nakov, T., Beaulieu, J. M., & Alverson, A. J. (2019). Diatoms diversify and turn over faster in

freshwater than marine environments. *Evolution*, *73*(12), 2497–2511.

https://doi.org/10.1111/evo.13832

Nevill, P. G., Bossinger, G., & Ades, P. K. (2010). Phylogeography of the world's tallest

angiosperm, *Eucalyptus regnans*: evidence for multiple isolated Quaternary refugia. *Journal of*

Biogeography, *37*(1), 179–192. https://doi.org/10.1111/j.1365-2699.2009.02193.x

Nürk, N. M., Linder, H. P., Onstein, R. E., Larcombe, M. J., Hughes, C. E., Piñeiro Fernández,

et al. (2020). Diversification in evolutionary arenas—Assessment and synthesis. *Ecology and*

Evolution, *10*(12), 6163–6182. https://doi.org/10.1002/ece3.6313

O'Meara, B. C., & Beaulieu, J. M. (2021). Potential survival of some, but not all, diversification

methods. Available at https://ecoevorxiv.org/w5nvd/.

Petit, R. J., & Hampe, A. (2006). Some evolutionary consequences of being a tree. *Annual*

Review of Ecology, Evolution, and Systematics, *37*, 187–214.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110215

Pohlert, T. (2014). The Pairwise Multiple Comparison of Mean Ranks Package (PMCMR). R

package, available at: https://CRAN.R-project.org/package=PMCMR.

Pryor, L. D. (1976). *The biology of Eucalypts.* Edward Arnold, London, UK.

Rabosky, D. L. (2010). Extinction rates should not be estimated from molecular

phylogenies. *Evolution*, *64*(6), 1816–1824. https://doi.org/10.1111/j.1558-5646.2009.00926.x

Rabosky, D. L. (2016). Challenges in the estimation of extinction from molecular phylogenies: a

response to Beaulieu and O'Meara. *Evolution*, *70*(1), 218–228.

https://doi.org/10.1111/evo.12820

Rabosky, D. L., Grundler, M., Anderson, C., Title, P., Shi, J. J., Brown, J. W., Huang, H., &

Larson, J. G. (2014). BAMM tools: an R package for the analysis of evolutionary dynamics on

phylogenetic trees. *Methods in Ecology and Evolution*, *5*(7), 701–707.

https://doi.org/10.1111/2041-210X.12199

Rabosky, D. L. (2014). Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PloS one*, *9*(2), e89543.

https://doi.org/10.1371/journal.pone.0089543

Rabosky, D. L. (2015). No substitute for real data: a cautionary note on the use of phylogenies

from birth–death polytomy resolvers for downstream comparative analyses. *Evolution*, 69(12),

3207-3216. https://doi.org/10.1111/evo.12817

Rabosky, D. L., & Goldberg, E. E. (2015). Model inadequacy and mistaken inferences of trait-

dependent speciation. *Systematic biology*, *64*(2), 340–355.

https://doi.org/10.1093/sysbio/syu131

Rabosky, D. L., Mitchell, J. S., & Chang, J. (2017). Is BAMM flawed? Theoretical and practical

concerns in the analysis of multi-rate diversification models. *Systematic biology*, *66*(4), 477–

498. https://doi.org/10.1093/sysbio/syx037

Redding, D. W., & Mooers, A. Ø. (2006). Incorporating evolutionary measures into conservation

prioritization. *Conservation Biology*, *20*(6), 1670–1678. https://doi.org/10.1111/j.1523-

1739.2006.00555.x

Ree, R. H., & Sanmartín, I. (2018). Conceptual and statistical problems with the DEC+ J model

of founder‐event speciation and its comparison with DEC via model selection. *Journal of*

Biogeography, *45*(4), 741–749. https://doi.org/10.1111/jbi.13173

Reich, P. B. (2014). The world‐wide 'fast–slow'plant economics spectrum: a traits manifesto.

Journal of Ecology, 102(2), 275–301. https://doi.org/10.1111/1365-2745.1221

Reginato, M., Vasconcelos, T. N., Kriebel, R., & Simoes, A. O. (2020). Is dispersal mode a

driver of diversification and geographical distribution in the tropical plant family

Melastomataceae? *Molecular phylogenetics and evolution*, *148*, 106815.

https://doi.org/10.1016/j.ympev.2020.106815

Russo, S. E., Brown, P., Tan, S., & Davies, S. J. (2008). Interspecific demographic trade‐offs

and soil‐related habitat associations of tree species along resource gradients. Journal of

Ecology, 96(1), 192–203. https://doi.org/10.1111/j.1365-2745.2007.01330.x

Simberloff, D. S. (1974). Equilibrium theory of island biogeography and ecology. *Annual review*

of Ecology and Systematics, 5(1), 161–182.

https://doi.org/10.1146/annurev.es.05.110174.001113

Smith, S. A., & Brown, J. W. (2018). Constructing a broadly inclusive seed plant

phylogeny. *American journal of botany*, *105*(3), 302–314. https://doi.org/10.1002/ajb2.1019

Stebbins, G. L. 1974. Flowering plants: evolution above the species level. Harvard University

Press, Cambridge, MA.

Steel, M., & Mooers, A. (2010). The expected length of pendant and interior edges of a Yule

tree. *Applied Mathematics Letters*, *23*(11), 1315–1319.

https://doi.org/10.1016/j.aml.2010.06.021

Suissa, J. S., Sundue, M. A., & Testo, W. L. (2021). Mountains, climate and niche heterogeneity

explain global patterns of fern diversity. *Journal of Biogeography*, 48(6), 1296-1308.

https://doi.org/10.1111/jbi.14076

Sun, M., Folk, R. A., Gitzendanner, M. A., Soltis, P. S., Chen, Z., Soltis, D. E., & Guralnick, R. P. (2020). Recent accelerated diversification in rosids occurred outside the tropics. *Nature Communications*, 11(1), 1-12. https://doi.org/10.1038/s41467-020-17116-5

Title, P. O., & Rabosky, D. L. (2019). Tip rates, phylogenies and diversification: what are we estimating, and how good are the estimates? *Methods in Ecology and Evolution*, *10*(6), 821–834. https://doi.org/10.1111/2041-210X.13153

Thornhill, A. H., Crisp, M. D., Külheim, C., Lam, K. E., Nelson, L. A., Yeates, D. K., & Miller, J. T. (2019). A dated molecular perspective of eucalypt taxonomy, evolution and diversification. *Australian Systematic Botany*, *32*(1), 29–48. https://doi.org/10.1071/SB18015

Uyeda, J. C., Zenil-Ferguson, R., & Pennell, M. W. (2018). Rethinking phylogenetic comparative methods. *Systematic Biology*, *67*(6), 1091–1109. https://doi.org/10.1093/sysbio/syy031

Vasconcelos, T. N., Chartier, M., Prenner, G., Martins, A. C., Schönenberger, J., Wingler, A., & Lucas, E. (2019). Floral uniformity through evolutionary time in a species-rich tree lineage. *New Phytologist*, *221*(3), 1597–1608. https://doi.org/10.1111/nph.15453

Vasconcelos, T., O'Meara, B. C., & Beaulieu, J. M. (2022). Retiring "cradles" and "museums" of biodiversity. *The American Naturalist*, 199(2), 194-205. https://doi.org/10.1086/717412

Vrba, E. S. (1993). Turnover-pulses, the Red Queen, and related topics. *American Journal of Science*, *293*(A), 418.

Westoby, M. (1998). A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and soil*, *199*(2), 213–227. https://doi.org/10.1023/A:1004327224729

Wiens, J. J., & Donoghue, M. J. (2004). Historical biogeography, ecology and species richness. *Trends in ecology & evolution*, *19*(12), 639–644. https://doi.org/10.1016/j.tree.2004.09.011

Wilson, P. G., O'brien, M. M., Heslewood, M. M., & Quinn, C. J. (2005). Relationships within Myrtaceae sensu lato based on a mat K phylogeny. *Plant Systematics and Evolution,* 251(1), 3–19 https://doi.org/10.1007/s00606-004-0162-y

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., et al. (2004). The worldwide leaf economics spectrum. *Nature*, *428*(6985), 821–827. https://doi.org/10.1038/nature02403