Structure Determination of Inactive-State GPCRs with a Universal Nanobody

Michael J. Robertson[1,2], Feng He[1,2], Justin G. Meyerowitz[1,2,3], Alpay B. Seven[1,2], Ouliana

Panova[1,2], Maria-Claudia Peroto[1,2], Tao Che[4,5], Georgios Skiniotis[1,2,6]

[1]Department of Molecular and Cellular Physiology, Stanford University School of

Medicine, Stanford, CA, USA

[2]Department of Structural Biology, Stanford University School of Medicine, Stanford, CA,

USA

[3]Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University

School of Medicine, Stanford, CA 94305, USA.

[4]Department of Anesthesiology, Washington University School of Medicine, St. Louis,

Missouri, 63110, USA.

[5]Center for Clinical Pharmacology, University of Health Sciences & Pharmacy at St. Louis

and Washington University School of Medicine, St. Louis, Missouri, 63110, USA.

[6]Correspondence: yiorgo@stanford.edu

Summary

Cryogenic electron microscopy (cryo-EM) has widened the field of structure-based drug

discovery by allowing for routine determination of membrane protein structures previously

intractable. However, despite representing one of the largest classes of therapeutic

targets, most inactive-state G protein-coupled receptors (GPCRs) have remained

inaccessible for cryo-EM because their small size and membrane-embedded nature

impedes projection alignment for high-resolution map reconstructions. Here we

demonstrate that a camelid single-chain antibody (nanobody) recognizing a grafted

intracellular loop can be used to obtain cryo-EM structures of different inactive-state

GPCRs at resolutions comparable or better than those obtained by X-ray crystallography.

Using this approach, we obtained the structure of human neurotensin 1 receptor (NTSR1)

bound to antagonist SR48692, of μ-opioid receptor (MOR) bound to the clinical

antagonist alvimopan, as well as the structure of the previously uncharacterized

somatostatin receptor 2 (SSTR2) in the apo state; each of these structures yields novel

insights into ligand binding and specificity. We expect this rapid, straightforward approach

to facilitate the broad structural exploration of GPCR inactive states without the need for

extensive engineering and crystallization.

Introduction

The impact of cryo-EM on structure-based drug discovery has been immense,

leading to the characterization of a wide variety of new and valuable membrane protein

drug targets, including ion channels and G protein-coupled receptors (GPCRs)[1]. As a key

requirement for high-resolution cryo-EM map reconstructions, the randomly oriented

particle projections must first be correctly aligned at intermediate-to-low resolution, a step

that often fails for small proteins due to their limited structural features. For relatively small

membrane proteins, the problem is compounded by the presence of a detergent micelle

or lipid nanodisc used for solubilization, which further dampens the contrast of protein

densities. This poses a major bottleneck for cryo-EM structure determination of GCPRs,

which account for roughly 35% of FDA-approved drugs, with approximately half of those

being agonists that activate the receptor and half antagonists or inverse agonists[2] that

engage the inactive-state receptor, thereby blocking endogenous signaling. Active-state

GPCR structures can be determined in complex with heterotrimeric G protein[3], a ~90 kDa

entity that adds significant mass outside the detergent micelle or lipid disc and provides

a source of alignment for the transmembrane receptor. However, there is no such

generalizable tool for cryo-EM structure determination of inactive-state GPCRs,

especially for family A receptors, the largest GCPR subfamily with hundreds of receptors

that are typically ~37-50 kDa and lack sizeable extracellular domains for projection

alignment. Thus, the application of cryo-EM is severely limited for inactive-state GPCRs,

and their structural elucidation mostly relies on extensive engineering and crystallization

trials for X-ray studies. This limitation impacts not only mechanistic investigations of

GPCRs, but also structure-based drug discovery for the countless therapeutically relevant

receptors.

Che *et al.*[4] recently reported a camelid VHH domain antibody, referred to as

nanobody 6 (Nb6), that engages the inactive-state intracellular loop 3 (ICL3) of the κ-

opioid receptor (KOR), even when this loop is grafted onto a wide range of receptors,

including orphan receptors and family B receptors. This tool was initially developed as a

bioluminescence resonance energy transfer (BRET)-based sensor of receptor activation

state, an approach that also allows for cost-effective and robust screening of Nb6-binding

constructs by employing BRET assays. Using the Nb6 approach, we obtained three-

dimensional (3D) map reconstructions of human NTSR1 (hNTSR1) in complex with Nb6

bound to the inverse agonist SR48692[5] at a resolution of 2.4 Å; the mouse MOR bound

to a megabody[6], an enlarged nanobody construct derived from Nb6 (Mb6), in complex

with the antagonist alvimopan[7] at a resolution of 2.8 Å; and the unliganded (apo) structure

of human SSTR2 bound to Nb6 at 3.1 Å resolution. We thus demonstrate that Nb6 is a

general solution for determining high-resolution inactive-state structures by cryo-EM.

Besides alleviating the need for extensive construct engineering and successful

crystallogenesis, the cryo-EM/Nb6 approach comes with two distinct advantages: first, it

does not necessitate the screening and development of receptor-specific nanobodies.

Second, the Nb6 BRET-based sensor allows for rapid and efficient screening of

constructs and conditions instead of time-consuming and expensive cryo-EM-based

screening.

Employing Nb6 for cryo-EM structure determination of GPCRs

Any fiducial marker that is used for cryo-EM structure determination of a target

macromolecule needs to be rigid in its binding and orientation in respect to the structural

target. Furthermore, if the marker is particularly small related to the target, it must be

positioned such that it provides a distinct feature from various angles to allow

unambiguous alignment of the randomly-oriented cryo-EM projections. A crystal structure

of Nb6 bound to KOR revealed an interaction at the base of TM5 and TM6, off-center

from the major axis of the receptor[4]. We hypothesized that this type of interaction would

be an ideal asymmetric fiduciary for cryo-EM particle projection alignment as it would

allow for the determination of the rotational orientation of the receptor around its major

axis, particularly during initial low-resolution alignments.

To evaluate the degree of rigidity of Nb6 when bound to a GPCR, we first

performed all-atom molecular dynamics (MD) simulations using the crystal structure of

the KOR-Nb6 complex. Alignment of the MD trajectories on the receptor atoms and

calculation of the average root mean squared fluctuation (RMSF) for Nb6 atoms over the

final 100 ns of triplicate 500 ns simulations (Fig. 1a) showed that Nb6 is rigidly bound and

characterized by little flexibility, with the exception of the distal loops at the base of the

nanobody.

Encouraged by these computational predictions, we proceeded with employing

Nb6 for *de novo* cryo-EM structure determination of three pharmacologically relevant

family A GPCRs. The first, NTSR1, is broadly associated with neurological and

gastrointestinal processes[8]. Overexpression of NTSR1 in some tumor types promotes

cancer progression and metastasis and is thus associated with poor prognosis[9],

prompting the study and development of NTSR1 inverse agonists for oncology[10]. We also

studied MOR, the target of opioids, which are both the most effective treatment for pain

but also the therapeutic class underlying the ongoing opioid epidemic in the United

States[11]. While current opioid antagonists can treat acute opioid overdose, there is an

urgent need for MOR antagonists with improved pharmacokinetic properties[11,12]. Finally,

SSTR2 plays a pivotal role in the neuroendocrine system by opposing the release of many

hormones, including growth hormone[13]. SSTR agonists are commonly prescribed for

neuroendocrine tumors, while SSTR subtype-selective antagonists are in development

for diabetes, but the lack of structural information for any somatostatin receptor family

member in either active or inactive state has made the development of next-generation

SSTR ligands challenging.

As described in Che *et al.*, we engineered constructs to enable binding of Nb6 to

mouse MOR and human SSTR2 by identifying minimal swaps of ICL3 along with the

cytoplasmic tips of transmembrane helix 5 and 6 (TM5/6) with that of the KOR (Fig. 1c).

In the case of the closely related MOR, the two point mutations M264L and K269$^{6.24}$R

(Ballesteros-Weinstein notation[14]; Fig. 1c) were sufficient to enable Nb6 binding. SSTR2

was able to bind Nb6 with the swap of all 15 residues from S238$^{5.68}$ of TM5 (V256$^{5.68}$ of

KOR) to K252$^{6.31}$ of TM6 (R270$^{6.31}$ of KOR) (Fig. 1c). For human NTSR1 (hNTSR1) we

utilized the same swap from KOR T247$^{5.59}$ to L277$^{6.38}$ as previously described for rat

NTSR1[4] (rNTSR1; Fig. 1c). BRET assays of agonist-induced Nb6 dissociation confirmed

construct functionality, with all three receptors showing dissociation of the nanobody in

response to agonist binding (Fig. 1d-f). In all experimental replicates, the BRET data for

MOR response to DAMGO appears to be biphasic, consistent with prior nanobody-based

MOR activation sensors[15], where the two-phase response is attributed to distinct plasma

membrane and endosomal signaling.

High-resolution cryo-EM structure determination was feasible in all cases. Despite

Nb6 being only 12 kDa, Nb6, or the equivalent region of Mb6, is resolved in all three

maps, providing a critical stable density outside the detergent micelle surrounding

NSTR1, MOR and SSTR2. Notably, the local resolution indications for Nb6 in our

GPCR/Nb6 complex cryo-EM maps showed high correlation with the relative root mean

square fluctuation (RMSF) values for Nb6 calculated from our MD simulations (Fig. 1b).

This observation further demonstrates the strong predictive power of MD simulations in

assessing the behavior of fiducial markers for such work.

Structure of hNTSR1

Obtaining inactive-state crystal structures has proven challenging for NTSR1. The

only available crystal structures were determined in 2021[16]: one with receptor in the apo

Fig 1. | Construct design and evaluation for Nb6-based GPCR structure determination

a, Root mean squared fluctuations (RMSF) in atomic positions averaged over triplicate molecular dynamics simulations for Nb6 bound to KOR. **b,** Local resolution plot of Nb6 from the cryo-EM map of the SSTR2/Nb6 complex. **c,** Overlay of the three KOR ICL3 constructs used to enable Nb6 binding for MOR (two point mutations), SSTR2 (5.68 to 6.31), and hNTSR1 (5.59 to 6.38). Dose-response data representing loss of BRET (net BRET) between Receptor-rLuc and Nb6-mVenus for **d,** NTSR1$_K$ and neurotensin (NT), **e,** MOR$_K$ and DAMGO, and **f,** SSTR2$_K$ and somatostatin 14 (SST14).

state (3.19 Å) and two structures with the receptor bound to the inverse agonists SR48692

(2.64 Å) or SR142948A (2.92 Å), which are small molecules previously in development

for treating cancer. Obtaining these structures required extensive engineering to facilitate

protein crystallization, with that work employing a construct of rat NTSR1 evolved in

bacteria to acquire numerous thermostabilizing mutations and fused in TM7 with a

DARPin[17] protein domain to facilitate crystal packing (rNTSR1-H4)[18]. To assess and

compare the cryo-EM/Nb6 approach with crystallography we obtained the cryo-EM

Fig 2. | Comparison of cryo-EM structure of hNTSR1 and crystal structure of rNTSR1-H4

a, Cryo-EM map of hNTSR1 at 2.4 Å global resolution. The arrow points to the cryo-EM density and model for inverse agonist SR48692, the inset to the left is the map-model fit for TM3 **b,** 2Fo-Fc crystallography map of rNTSR-H4 at 2.6 Å contoured at σ=1.0. The arrow points to the Local 2Fo-Fc map around SR48692 contoured at σ=1.25. **c,** Water molecules resolved in the core of the hNTSR1 receptor and their associated cryo-EM densities. **d,** Overlay of the hNTSR cryoEM structure (protein in green, SR48692 in purple) with the rNSTR-H4 crystal structure (protein in gray, SR48692 in pink) highlighting differences in ECL2. **e,** Overlay of the hNTSR cryoEM structure (protein in green, SR48692 in purple) with the rNSTR-H4 crystal structure (protein in gray, SR48692 in pink) highlighting the residue shift and probable loss of bound water molecule caused by the F7.42V thermostabilizing mutation

structure of hNTSR1/Nb6 complex bound to SR48692 (Fig. 2a, Extended Data Fig. 2).

Notably, the cryo-EM structure was resolved at a global resolution of 2.4 Å, both a higher

nominal resolution than the crystal structure and with significantly better-resolved

features, including the density corresponding to the ligand (Fig. 2a, b insets, Extended

Data Fig. 8a). This resolution range also reveals densities corresponding to extensive

hydration throughout the receptor in the cryo-EM map (Fig. 2c, Extended Data Fig. 3a).

The construct for hNTSR1 was minimally modified beyond the KOR ICL3 swap,

with only N- and C- termini truncations in regions not resolved in typical family A structures

and a single A85$^{1.54}$L mutation to increase expression[19]. While the general ligand pose is

very similar between the two structures, the cryo-EM structure reveals additional

interactions not observed in the rNTSR-H4 crystal structure. First, the remodeling of the

TM7-ECL3 region allows W334 in ECL3 to be resolved in the hNTSR1 structure loosely

capping the top of the hydrophobic chloro-naphthyl and dimethoxy-phenyl moieties of

SR48692 (Fig. 2c). This change may be due to several amino acid sequence differences

at the top of TM7, most prominently P336$^{7.25}$ in hNTSR1 versus T341$^{7.25}$ in rNTSR1-H4,

with the proline inducing a sharp turn between the top of TM7 and ECL3 in hNTSR1. In

addition, the rearrangement of R327$^{6.54}$ in rNTSR-H4 (R322$^{6.54}$ in hNTSR1) appears to

disrupt a water-mediated interaction between R327$^{6.54}$, R328$^{6.55}$ (R323$^{6.55}$ in hNTSR1),

and the carboxylate moiety of SR48692 observed in the hNTSR1 structure (Fig. 2d). This

R327$^{6.54}$ rearrangement likely stems from the F358$^{7.42}$V thermostabilizing mutation in

rNTSR-H4, allowing a downward relaxation of Y324$^{6.51}$ (Y319$^{6.51}$ in hNTSR1) and

R327$^{6.54}$. Notable differences also exist in the TM helices between the two structures;

TM1 is shifted 3.4 Å further outward from the core of the receptor in the crystal structure

(Extended Data Fig. 3b), while the top of TM2 and ECL1 are out of register by one residue.

While this may be due to the differences in experimental constructs, the observed

discrepancy may also result from the fact that even at σ=1 the 2Fo-Fc map of the crystal

structure lacks sidechain information in the ECL1 region, rendering modeling ambiguous

(Extended Data Fig. 3c,d). Overall, the lack of thermostabilizing mutations and DARPin

in TM7 restores a more native-like inactive state as suggested by the conformation of key

amino acids. For example, Y364[7.53] of the hNTSR1 NPxxY motif (Y369[7.53] in rNSTR1)

occupies a position and rotamer more typical of an inactive state family A GPCR

(Extended Data Fig. 3e).

Structure of MOR

In parallel, we obtained a cryo-EM structure of the MOR bound to the non-blood

brain barrier penetrant, MOR selective inverse agonist alvimopan, an FDA-approved drug

for reversing opioid-induced gastrointestinal (GI) symptoms in hospital settings. In the

case of NTSR1/Nb6, vitrifying the sample in a very thin buffer layer on holey gold grids[20]

was necessary, not only for high resolution but also to achieve proper alignment of particle

projections. To assess an alternative approach, we used Nb6 to develop a megabody,

termed 'Mb6', a fusion of a nanobody and a scaffold protein of either 45 or 86 kDa that

can serve as a larger fiducial marker, based on the work of Uchański *et al.*[6]. In that work,

the Mb designs employed for $GABA_A$ and WbaP produced cryo-EM maps that could not

resolve well past the nanobody portion, presumably due to flexibility. We thus opted

instead for using the c7HopQA12 design, which incorporates a twisted linker between the

nanobody and the 45 kDa HopQ scaffold to increase their contact and, ideally, rigidity.

However, our MD simulations suggested that even this design was characterized by

substantial flexibility between the Nb and scaffold (Extended Data Fig. 5a). In agreement

with this result, our cryo-EM 2D class averages of the MOR-Mb6 complex showed that

only the Nb6-receptor portion is well resolved, with the scaffold averaging out (Extended

Data Fig. 4b). Nevertheless, leveraging the ordered Nb6 portion of Mb6 in our cryo-EM

data, we obtained a 2.8 Å cryo-EM map (Fig. 3a, Extended Data Fig. 4) that provided an

unambiguous density for the ligand and receptor (Fig. 3c, Extended Data Fig. 8b). This

level of detail is quite comparable to the 2.8 Å crystal structure of MOR (Fig. 3b, d),

although the cryoEM map further resolves several probable water molecules in the ligand

binding site, including one that bridges interactions between the carboxylate of alvimopan

and the receptor (Extended Data Fig. 5b).

Comparing the binding poses of the MOR-selective alvimopan with that of the

structurally similar ligand JDTic, which is selective for KOR, reveals a striking difference.

Despite their similar scaffolds, the phenol-piperidine moieties of alvimopan and JDTic are

in completely different positions and orientations, nearly flipped in the binding pocket.

Aligning our structure with KOR-JDTic (PDB:4DJH[21]) reveals that JDTic would clash with

Y326$^{7.43}$ and Q124$^{2.60}$ in MOR, while alvimopan would clash with W124 and I294$^{6.55}$ in

KOR (I294$^{6.55}$ of KOR is substituted by a repositioned valine in MOR). In addition to the

steric confines of the binding pocket, charge differences also appear to contribute to the

differences in binding pose between JDTIC and alvimopan. The nitrogen of the

tetrahydroisoquinoline ring system of JDTic is likely also protonated or perhaps

protonated instead of the piperidine, and thus forms salt bridge interactions with D138$^{3.32}$

(D147$^{3.32}$ in MOR), just as the protonated piperidine of alvimopan does in MOR. Of note,

alvimopan shares similarities to fentanyl and its analogues, including a central piperidine

ring. Computational docking of carfentanil[22], a potent MOR agonist, in the active-state

structure of the receptor[3] suggests a pose similar to alvimopan, but with the additional

phenylamine moiety on the piperidine ring positioned to bind into the pocket occupied by

the toggle switch tryptophan W293$^{6.48}$ (Extended Data Fig. 5a). How a MOR ligand

engages this region is often the main determining factor for its agonism versus

Fig 3. | Comparison of cryo-EM structure of alvimopan-bound MOR with crystal structures of MOR/β-FNA and KOR/JDTic **a,** Cryo-EM map of MOR at 2.8 Å global resolution. **b,** 2Fo-Fc crystallography map of MOR at 2.8 Å contoured at σ=2.0. **c,** Cryo-EM density and model for inverse agonist alvimopan. **d,** Local 2Fo-Fc map around β-FNA contoured at σ=1.5. **e,** overlay of the alvimopan pose in the MOR cryo-EM structure (protein in blue, alvimopan in magenta) with the JDTic pose in the KOR crystal structure (protein in grey, JDTic in orange). **f,** Comparison of JDTic and alvimopan chemical structures highlighting similar phenol-piperidine scaffold.

223 antagonism, and we hypothesize that this phenyl moiety of fentanyl and its derivatives is

224 responsible for conversion of the piperidine scaffold to an agonist.

Structure of SSTR2

To test the cryo-EM/Nb6 approach on a receptor without prior structural

characterization we chose SSTR2, a member of the SSTR subfamily for which no

structures have been determined. Despite the absence of a stabilizing inverse agonist or

any other ligand and the lack of an energy filter in the microscope used for cryo-EM data

acquisition (typically beneficial for data quality when imaging small proteins), we were

able to obtain a 3.1 Å map of SSTR2/Nb6 with well-resolved features (Fig. 4a, Extended

Data Fig. 6, Extended Data Fig. 8c). Together with two active-state structures of SSTR2

bound to two different agonists, the endogenous 14-mer SST14 and the synthetic 8-mer

octreotide (detailed in our accompanying manuscript[23]), the inactive-state SSTR2

structure enabled us to characterize several aspects of receptor activation and ligand

recognition. Perhaps the most interesting feature of SSTR2 is the conformational flexibility

of ECL2; while in a well-defined position in the apo state structure, ECL2 is pushed to a

more open conformation when bound to the endogenous agonist SST14, but folds down

over the top of the receptor in the octreotide-bound state (Fig. 4b). As a consequence of

the ECL2 plasticity, different residues are engaged in binding the pan-SSTR agonist

SST14 versus the SSTR2-selective octreotide. This is an important factor to SSTR

subtype selectivity, which is explored further in the accompanying study[23].

Very few structures have been obtained to date for family A GPCRs in the

unliganded, apo state, primarily due to the general need for a stabilizing ligand to obtain

a crystal structure. Experimentally-derived apo structures provide an opportunity to

evaluate predicted structures from the recently published deep-learning based tools,

AlphaFold2[24] and RoseTTAfold[25]. Notably, while both software performed well in

predicting the overall structure of SSTR2, key details of the ligand binding site were in

disagreement with the experimental structure, particularly in residues $D122^{3.32}$, $F92^{2.53}$,

and $Y302^{7.43}$ (Fig. 4c), which are involved in ligand binding and activation, as detailed in

the companion active-state SSTR2 manuscript[23]. Although a more systematic

comparison is needed, this observation underscores both the merits but also the potential

limitations of homology model-based approaches. Differences in side chain positioning,

especially within a ligand binding site, can throw off structure-based drug discovery efforts

and other pursuits in which high-accuracy coordinates are required.

Fig 4. | Cryo-EM structure of the apo SSTR2 and comparison of sodium ion binding site.
a, Cryo-EM map of SSTR2 at 3.1 Å global resolution. **b,** Comparison of the model and cryo-EM maps of apo SSTR2 (teal) and SSTR2 in complex with either SST14 (magenta) or octreotide (purple). **d,** Overlay of SSTR2 (teal), MOR (blue), and NTSR1 (green) cryo-EM structures around the canonical family A sodium ion binding site with the DOR sodium coordination site structure (gray, PDB:4N6H).

Differential sodium accommodation by family A GPCRs

Our near-native inactive-state cryo-EM maps provide the opportunity to examine

the allosteric sodium binding site found deep in the core of many family A receptors. All

three of the receptors studied in this work have experimentally demonstrated negative

allosteric modulation by sodium[26–28]. In both MOR and SSTR2 the identities of the

residues in the sodium binding site are the same as the δ-opioid receptor (DOR), for which

there is a high-resolution crystal structure (1.8 Å) with a well-defined sodium site for

comparison[29] (Fig. 4c). The geometries of the MOR, SSTR2, and DOR sites are very

similar, with the exception of a slight rearrangement of $W^{6.48}$ in the case of both MOR

($W293^{6.48}$) and SSTR2 ($W269^{6.48}$), and $N125^{3.35}$ in SSTR2. In contrast to SSTR2 and

MOR, hNTSR1 has four amino acid differences in the sodium binding site, namely

T155$^{3.39}$ (in place of S$^{3.39}$), S357$^{7.46}$ (in place of N$^{7.46}$), S111$^{2.49}$ (in place of A$^{2.49}$), and

C151$^{3.35}$ (in place of N$^{3.35}$). In the DOR structure, both S135$^{3.39}$ and N131$^{3.35}$ directly

interact with the sodium .

The three maps we determined resolve features consistent with water and/or ions

in the canonical sodium binding region near D2.50, although only NTSR1 has sufficiently

high resolution in this particular region to facilitate unambiguous ion modeling (Extended

Data Fig. 7a-c). The probable ion sites of SSTR2 and MOR match relatively well to those

observed in DOR. In the hNTSR1 map, we observe a density with close proximity to

oxygen atoms of T155$^{3.39}$, S111$^{2.49}$, D112$^{2.50}$ (Extended Data Fig. 7a). Even though

precise identification of water versus ion is challenging in cryo-EM maps, we expect that

this feature corresponds to the sodium, shifted by 1.8 Å compared to the sodium in DOR

(Fig. 4d, Extended Data Fig. 7d, e). As other receptors with resolved sodium ion sites

have cysteine or other hydrophobic residues in place of N$^{3.35}$, S111$^{2.49}$ is likely responsible

for the shift in sodium position in NTSR1. Overall, we find substantial similarity between

the sodium binding sites of inactive state GPCRs from cryo-EM and crystallography.

Discussion

Here, we present inactive-state structures of three family A GPCRs, all obtained

by cryo-EM at resolutions as high as 2.4 Å by employing minimal engineering and a

universal nanobody for projection alignment. By using molecular dynamics simulations to

assess fiducial markers for use in cryo-EM structure determination, we could accurately

predict that Nb6 is sufficiently rigid for cryo-EM particle alignment, while Mb6 is too flexible

for resolution beyond the nanobody portion. Each of the obtained cryo-EM structures

yields new insights into ligand recognition, with resolution that is as good or better than

that of analogous structures produced by x-ray crystallography. In contrast to active state

GPCR-G protein complexes, where the highest-resolution portion of the map is on the

larger G protein due to the center of alignment, the inactive-state Nb6 GPCR structures

display their highest resolution at the receptor core, facilitating accurate modeling of

ligands and their surrounding hydration. This generalizable approach opens this broad

family of drug targets to rapid, crystallization-free structure determination, with

biochemical stability of the receptor and nanobody binding being the only limitations. As

such, we anticipate Nb6 to be a broadly applicable approach to determining inactive state

structures of near-native GPCRs, further propelling drug discovery in this class of

receptors.

Author Contributions

M.J.R. cloned constructs, expressed and purified proteins, processed EM data, built

models, and ran/analyzed molecular dynamics simulations. F.H. cloned constructs and

expressed proteins. J.M. performed BRET assays. A.S. prepared cryo-EM samples and

collected cryo-EM data. M.C. expressed proteins and purified nanobody. O.P. prepared

cryo-EM samples and collected cryo-EM data. T.C. provided constructs for nanobody

expression and BRET assays. M.J.R. and G.S. wrote the manuscript with input from F.H.,

J.M., O.P., and T.C.. G.S. supervised the project.

Data Availability

All data generated or analyzed in this study are included in this article and the

Supplementary Information. The cryo-EM density maps and corresponding coordinates

have been deposited in the Electron Microscopy Data Bank (EMDB) and the Protein Data

Bank (PDB), respectively, under the following accession codes: Raw data has been

deposited in EMPIAR under the following accession codes:

Competing Interests

The authors declare no competing interests.

Acknowledgements

We thank the Kobilka lab for providing plasmids of NTSR1 and MOR. Cryo-EM data

were collected at the Stanford cryo-EM center (cEMc) with support from E. Montabana.

This work was supported, in part, by the Mathers Foundation (G.S.), training grant

T32GM089626 (J.G.M.), and used the Extreme Science and Engineering Discovery

Environment (XSEDE)[30] resource comet-gpu through sdsc-comet allocation TG-

MCB190153 (G.S.), which is supported by National Science Foundation grant number

ACI-1548562.

Methods

Cloning

The cDNA for SSTR2 was obtained from Horizon Discovery and cloned into a pFastBac

vector containing an N-terminal haemagglutinin (HA) signal sequence followed by a FLAG

epitope (DYKDDDD) and a C-terminal C3 protease cleavage site followed by enhanced

green fluorescent protein (eGFP) and a hexahistidine (His6) tag using Gibson cloning.

Constructs for hNTSR1 and MOR were obtained from the lab of Brian Kobilka and have

been described previously[3,19]. Briefly, each is cloned in pFastBac with an N-terminal

haemagglutinin (HA) signal sequence followed by a Flag epitope (DYKDDDD) and a C-

terminal HRV 3C protease cleavage site followed by a hexahistidine (His6) tag. In the

case of hNTSR1, the His6 tag is preceded by enhanced green fluorescent protein (eGFP).

For MOR, two point mutations (M264L and K269R) were introduced with site-directed

mutagenesis; PCR reactions were conducted with Q5 polymerase with the following pair

of primers: 5'-   GGCTCCCGCGAAAAGGACAGGAACCTGCGC   -3';   5'-

CGACAGCAGGCGGACACTCTTGAG

-3'. For hNTSR1, residues between T247$^{5.59}$ to L277$^{6.3}$ of kappa opioid were substituted

with site-directed mutagenesis in three steps, with the following three primer pairs: 5'-

ATCACACGCCTGGTTCTGGCAGTGGTCATCGCCT

-3'   5'-   AATATCATCAGGGTGTACAGGACCGAGATGACCA   -3';   5'-

GGAATCTGCGCCGAATCACACGCCTGGTTCT   -3'   5'-

ACAGATTTCAGCCGCAATATCATCAGGGTGTACAG-3';   5'-

GCCGCTCAGCAGGCGCACAGATTTCAGCCGCA   -3'   5'-

AGCCGCGAAAGGATCGGAATCTGCGCCGAAT -3'. For SSTR2, residues between

S238 and K252 were replaced with those of kappa opioid in two steps with site-directed

mutagenesis with the following primer pair: 5'-

GAGCGGCAGCTCTAAGAGGAAGAAGTCTGAGAA

-3' 5'- AGCAGGCGCACGGACTTCACCTTGATGATAATGAAC

-3'; 5'- CGGAATCTGCGCAAGGTCACCCGAATGGTG

-3', 5'- ATCCTTTTCGCGGCTGCCGCTCAGCA

-3'.

Biolumescence resonance energy transfer (BRET) assays

BRET assays were performed and analyzed as previously described[4] with the following

modifications: HEK-293S cells grown in FreeStyle 293 suspension media (Thermo

Fisher) were transfected at a density of 1 million cells/mL in 2 mL volume using 600 ng

total DNA at 1:1 ratio of Receptor-rLuc:Nb6-mVenus and a DNA:PEI ratio of 1:5,

and incubated in a 24 deep well plate at 220 rpm, 37°C for 48 hours. Cells were harvested

by centrifugation, washed with Hank's Balanced Salt Solution (HBSS) without

Calcium/Magnesium (Gibco), and resuspended in assay buffer (HBSS with 20 mM

HEPES pH 7.45) with 1 μg/mL freshly prepared coelenterazine h (Promega). Cells were

plated in white-walled, white-bottom 96 well plates (Costar) in a volume of 60 μl/well and

60,000 cells/well. Ligands were prepared in drug buffer (assay buffer with 0.1% BSA, 6

mM $CaCl_2$, 6 mM $MgCl_2$), and added at a 1:2 ratio of drug:cell suspension. Ten minutes

after the addition of ligand, plates were read using a SpectraMax iD5 plate reader using

485 nm and 535 nm emission filters with a one-second integration time per well. The

computed BRET ratios (mVenus/RLuc emission) were normalized to ligand-free control

(Net BRET) prior to further analysis.

Expression and Purification of Nb6 and Mb6

Wk6 *E. coli* (ATCC) were transformed via heat shock with Nb6 plasmid[4] and two 5 mL

starter cultures were grown overnight in LB supplemented with 100 μg/ml ampicillin. 2L

of terrific broth were supplemented with 100 μg/ml ampicillin, inoculated with starter

cultures, and grown at 37°C with shaking until OD_{600}=0.7. Expression was then induced

with 1mM IPTG and expression was allowed to proceed at 28°C overnight. Pellets were

harvested via centrifugation and washed once with phosphate-buffered saline before

snap freezing in liquid nitrogen. Pellets containing Nb6 were thawed and resuspended in

50 mM Tris pH 8.0, 0.5 mM EDTA, 20% w/v sucrose (TES) buffer at 15 mL / 1 liter pellet

and supplemented with protease inhibitor cocktail and shaken in an orbital shaker for 1

hour at 4°C. An additional 30 mL per 1 liter pellet of TES diluted 1:4 with water was added

and placed in an orbital shaker for an additional 45 minutes at 4°C. Cell debris was

removed by ultracentrifugation at 100,000xg for 30 minutes. The supernatant was filtered

with a 45 micron filter, supplemented with 20 mM imidazole, and loaded over a gravity Ni-

NTA column at 4°C. The column was washed with 10 CV of a buffer containing 250 mM

NaCl, 50 mM Tris pH 7.5, 10 mM imidazole, and then eluted with buffer containing 250

mM imidazole. Nanobody was concentrated and applied to size exclusion

chromatography with a buffer containing 250 mM NaCl and 50 mM Tris pH 7.5.

Monomeric fractions were pooled, concentrated, supplemented with 10% glycerol, and

snap frozen in liquid nitrogen for later use. Expression and purification of Mb6 was

performed identically to Nb6.

Expression and Purification of Receptor/Nb6 Complexes

All receptors were expressed in Sf9 insect cells (Expression Systems) infected at a

density of 3-4 million cells/ml. At 48 hours post-infection, cells were collected with

centrifugation, washed with phosphate-buffered saline containing 10 μM antagonist

(when used), and pellets were snap frozen in liquid nitrogen for purification. A general

protocol was used for producing receptor/Nb6 complexes (Extended Data Fig. 1), with

some slight deviations. Sf9 pellets containing receptor were lysed in hypotonic buffer

containing 20 mM HEPES pH 7.5, 5 mM $MgCl_2$, protease inhibitor cocktail, benzonase,

10 μM antagonist (when used), and 2 mg/ml iodoacetamide and gently stirred for an hour

at 4°C. Membranes were harvested with ultracentrifugation at 100,000xg, supernatant

was discarded, and membranes were resuspended in solubilization buffer containing 250

mM NaCl, 20 mM HEPES pH 7.5, 1 mM $MgCl_2$, protease inhibitor cocktail, 10 μM

antagonist (when used), and 2 mg/ml iodoacetamide; then drip frozen into liquid nitrogen

and stored at -80°C. Resuspended membranes were thawed and detergent was added

dropwise while stirring at 4°C to a final concentration of 1% LMNG/0.1% CHS/0.1%

Cholate. After 3 hours, insoluble debris was removed with ultracentrifugation at 100,000

g. Solubilized receptor was supplemented with 20 mM imidazole and gravity loaded over

a Ni-NTA resin column. Columns were washed with 10 column volumes of buffer

containing 250 mM NaCl, 20 mM HEPES pH 7.5, 10 μM antagonist (when used), 20 mM

imidazole, and 0.1% LMNG/0.01% CHS and protein was eluted in buffer containing 250

mM NaCl, 20 mM HEPES pH 7.5, 10 μM antagonist (when used), 250 mM imidazole, and

0.01% LMNG/0.001% CHS. Receptor was then supplemented with 5 mM $CaCl_2$ and

loaded onto M1 flag resin; washed with 5 column volumes of 250 mM NaCl, 20 mM

HEPES pH 7.5, 10 μM antagonist (when used), 2 mM CaCl2, and 0.1% LMNG/0.01%

CHS; and eluted with buffer containing 150 mM NaCl, 20 mM HEPES pH 7.5, 10 μM

antagonist (when used), 1 mM EDTA, 0.2 mg/ml FLAG peptide, and 0.1% LMNG/0.01%

CHS. Purified receptor was incubated on ice overnight with HRV 3C protease and purified

Nb6 was added at a 2:1 molar ratio. Receptor/Nb6 complex was concentrated in a 50

kDa molecular weight cutoff spin concentrator and subjected to SEC chromatography

with an ENrich 650 column (Bio-Rad) and a buffer containing 150 mM NaCl, 20 mM

HEPES pH 7.5, 0.001% LMNG, 0.00033% GDN, and 0.0001% CHS. Fractions containing

monomeric receptor/Nb6 complex were pooled for cryo-EM sample preparation. This

protocol was identical for SSTR2 and NTSR1 with the exception of the addition of 100

μM TCEP to the lysis buffer for NTSR1. Preparation of MOR/Mb6 complex was identical

to that of receptor Nb6 complex, with the omission of the HRV-3C cleavage to remove

eGFP as it was not present in the construct.

Cryo-EM Sample Preparation

All samples were prepared on glow-discharged holey gold grids (Quantifoil ultrAufoil

R1.2/1.3), blotted in a FEI Vitrobot Mark IV (Thermo Fisher Scientific) at 4°C and 100%

humidity, and plunge frozen into liquid ethane. Blotting conditions for each sample were

as follows: 3.0 μl of NTSR1/Nb6 complex at 16 mg/ml with an additional 0.05% beta OG;

3.0 μl of MOR/Mb6 complex at 5 mg/ml; and 2.5 μl of SSTR2/Nb6 complex at 16 mg/ml

with an additional 0.05% beta OG.

Cryo-EM Data Collection

All samples were collected on a KRIOS electron microscope at an accelerating voltage

of 300 kV, with an energy filter for NTSR1/Nb6 and MOR/Mb6 and without an energy filter

for SSTR2/Nb6. All data was collected through SerialEM with beam tilt compensation and

recorded on a Gatan K3 direct electron detector. The resulting image stacks have a pixel

size of 0.434 Å for NTSR1/Nb6 and MOR/Mb6 and 0.426 Å for SSTR2/Nb6, all in super

resolution mode. Each NTSR1/Nb6 image stack is composed of 71 frames with an

incident electron dose of 0.85 $e^-/Å^2$ per frame, for a total dose of 61 $e^-/Å^2/s$ per micrograph.

Each MOR/Mb6 image stack is composed of 63 frames with an incident electron dose of

0.98 $e^-/Å^2$ per frame, for a total dose of 61 $e^-/Å^2/s$ per micrograph. Each SSTR2/Nb6

image stack is composed of 55 frames with an incident electron dose of 1.26 $e^-/Å^2$ per

frame, for a total dose of 69 $e^-/Å^2/s$ per micrograph.

Cryo-EM Data Processing

A pictorial depiction of each cryoEM processing workflow can be found in Extended Data

Figs. 2c, 4c, and 6c. Briefly, dose-fractionated image stacks were imported into RELION-

3.1[31] and subjected to beam-induced motion correction and dose weighting with

MotionCor2[32]. Contrast transfer function parameter estimation was performed with

CTFFIND-4.1[33]. Particle selection and extraction was performed with the Laplacian

autopicker function of RELION-3.1 on micrographs with a CTF fit better than 3.5 Å. The

extracted particle stack was imported into CryoSPARC[34] and subjected to multiple rounds

of 2D classification followed by iterative rounds of *ab initio* reconstruction with multiple

classes and heterogeneous refinement to further classify particles. In early 3D iterations,

particles from poor 3D classes were subjected to additional 2D classification and good

particles were 'rescued' for further 3D classification. Once iterative *ab initio* and

heterogenous refinement no longer produced higher resolution reconstruction, particles

were re-imported to RELION-3.1. In the cases of hNTSR1 and MOR CTF fit cutoffs of 3.1

Å and 3.3 Å were applied, respectively. Particles were then subjected to 3D refinement in

RELION-3.1 followed by Bayesian polishing. Polished particles were then imported back

to CryoSPARC for non-uniform refinement, global CTF refinement based on optics group,

and repeated non-uniform and local refinement.

Model Building and Refinement

All initial models were rigid body fit into cryo-EM maps using the ChimeraX[35] software.

For hNTSR1, the cryo-EM model of active state hNTSR1 was used as the initial model

(PDB:6OS9[19]); for MOR the crystal structure of inactive MOR bound to a covalent

antagonist was used as the initial model (PDB:4DKL[36]), and for SSTR2 a homology model

generated from KOR (PDB:6VI4[4]) was used. In all cases KOR/Nb6 complex (PDB:6VI4[4])

was aligned to the receptor to produce the initial model for the nanobody. All structures

were manually refined in COOT[37] with iterative real-space refinement in Phenix[38]. Once

accurate modeling was achieved for the protein components, the GemSpot pipeline[39] was

used for automatic modeling of the ligands alvimopan and SR48692 into MOR and

hNTSR1 respectively. Final refinement was executed in Phenix, followed by model-free

Phenix map modification for NTSR1 and MOR[40]. All maps displayed in this work are those

prior to model-free map modification unless stated otherwise.

Molecular Dynamics Simulations

Simulations of the KOR/Nb6 complex started with the crystal structure PDB:6VI4, with the

pose of JDTic replaced with that of PDB:4DJH[21]. Maestro's protein preparation tool was

used to assign protonation states, optimize hydrogen bonding, and build missing

sidechains and loops. Once the receptor was prepared, the orientation of proteins in

membranes webserver (OPM)[41] was used to align the receptor as it would be in a

membrane in the xy plane and the CHARMM-GUI[42] was used to build the full system with

a 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine POPC/CHS lipid bilayer, TIP3P

water[43], and 150 mM NaCl. PSF and PDB files were generated with the OPLS-AA/M[44]

forcefield in VMD[45] for the system. For Mb6, a homology model was built from PDB:6XVI

with the Prime homology modeling tool[46]. This model was solvated in TIP3P water and

150 mM NaCl in VMD for simulation.

Molecular dynamics simulations were carried out in NAMD[47] with a 2 fs timestep

with SHAKE and SETTLE algorithms[48,49], with a Langevin thermostat and a Nosé-Hoover

Langevin piston barostat at 1 atm with a period of 50 fs and decay of 25 fs. Periodic

boundary conditions were used with nonbonded interactions smoothed starting at 10 Å to

12 Å with long-range interactions treated with particle mesh Ewald (PME). The system

was minimized for 1,500 steps and then slowly heated from 0 to 303.15K in increments

of 20K simulating for 0.4 ns at each increment. For the Kappa/Nb6 simulations, all non-

hydrogen, non-water, and non-ion atoms were restrained with a 1 kcal/mol/ $Å^2$ harmonic

restraint during heating and an additional 10 ns of equilibration. 1 kcal/mol/ $Å^2$ harmonic

restraints were used for an additional 10 ns of equilibration on all non-hydrogen protein

atoms followed by a final 10 ns of restrained equilibration with 1 kcal/mol/ $Å^2$ harmonic

restraint on only Cα atoms. The first 30 ns of unrestrained molecular dynamics were also

considered to be equilibration, with an additional 500 ns of simulation performed. All

simulations were performed in triplicate. For Mb6, the first 30 ns of unrestrained molecular

dynamics were treated as equilibration, and 100 ns of triplicate molecular dynamics

simulations were performed.

Docking

The cryo-EM structure of active MOR (PDB:6DDE)[3] was prepared in Maestro and

carfentanil was docked with Glide XP docking[50].

References

1.   Robertson, M. J., Meyerowitz, J. G. & Skiniotis, G. Drug discovery in the era of cryo-

electron   microscopy.   *Trends   in   Biochemical   Sciences*   (2021).

doi:10.1016/j.tibs.2021.06.008

2.   Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schiöth, H. B. & Gloriam, D. E. Trends

in GPCR drug discovery: New agents, targets and indications. *Nat. Rev. Drug Discov.*

(2017). doi:10.1038/nrd.2017.178

3.   Koehl, A. *et al.* Structure of the μ-opioid receptor-Gi protein complex. *Nature* **558**, 547–552

(2018).

4.   Che, T. *et al.* Nanobody-enabled monitoring of kappa opioid receptor states. *Nat. Commun.*

(2020). doi:10.1038/s41467-020-14889-7

5.   Gully, D. *et al.* Biochemical and pharmacological profile of a potent and selective

nonpeptide antagonist of the neurotensin receptor. *Proc. Natl. Acad. Sci. U. S. A.* (1993).

doi:10.1073/pnas.90.1.65

6.   Uchański, T. *et al.* Megabodies expand the nanobody toolkit for protein structure

determination by single-particle cryo-EM. *Nat. Methods* (2021). doi:10.1038/s41592-020-

01001-6

7.   Zimmerman, D. M. *et al.* Discovery of a Potent, Peripherally Selective trans-3,4-Dimethyl-
4-(3-hydroxyphenyl)piperidine Opioid Antagonist for the Treatment of Gastrointestinal
Motility Disorders. *J. Med. Chem.* (1994). doi:10.1021/jm00041a003

8.   Carraway, R. & Leeman, S. E. Characterization of radioimmunoassayable neurotensin in
the rat. Its differential distribution in the central nervous system, small intestine, and
stomach. *J. Biol. Chem.* (1976). doi:10.1016/s0021-9258(17)32938-1

9.   Christou, N. *et al.* Neurotensin pathway in digestive cancers and clinical applications: an
overview. *Cell Death and Disease* (2020). doi:10.1038/s41419-020-03245-8

10.   Liu, J. *et al.* neurotensin receptor 1 antagonist SR48692 improves response to carboplatin
by enhancing apoptosis and inhibiting drug efflux in ovarian cancer. *Clin. Cancer Res.*
(2017). doi:10.1158/1078-0432.CCR-17-0861

11.   Volkow, N. D. & Blanco, C. The changing opioid crisis: development, challenges and
opportunities. *Molecular Psychiatry* (2021). doi:10.1038/s41380-020-0661-4

12.   Volkow, N. D. & Collins, F. S. The Role of Science in Addressing the Opioid Crisis. *N. Engl.*
J. Med. (2017). doi:10.1056/nejmsr1706626

13.   Günther, T. *et al.* International union of basic and clinical pharmacology. CV. somatostatin
receptors: Structure, function, ligands, and new nomenclature. *Pharmacol. Rev.* (2018).
doi:10.1124/pr.117.015388

14.   Ballesteros, J. A. & Weinstein, H. Integrated methods for the construction of three-
dimensional models and computational probing of structure-function relations in G protein-
coupled receptors. *Methods Neurosci.* (1995). doi:10.1016/S1043-9471(05)80049-7

15.   Stoeber, M. *et al.* A Genetically Encoded Biosensor Reveals Location Bias of Opioid Drug
Action. *Neuron* (2018). doi:10.1016/j.neuron.2018.04.021

16.   Deluigi, M. *et al.* Complexes of the neurotensin receptor 1 with small-molecule ligands
reveal structural determinants of full, partial, and inverse agonism. *Sci. Adv.* (2021).
doi:10.1126/sciadv.abe5504

17.   Mittl, P. R., Ernst, P. & Plückthun, A. Chaperone-assisted structure elucidation with DARPins. *Current Opinion in Structural Biology* (2020). doi:10.1016/j.sbi.2019.12.009

18.   Scott, D. J., Kummer, L., Egloff, P., Bathgate, R. A. D. & Plückthun, A. Improving the apo-state detergent stability of NTS1 with CHESS for pharmacological and structural studies. *Biochim. Biophys. Acta - Biomembr.* (2014). doi:10.1016/j.bbamem.2014.07.015

19.   Kato, H. E. *et al.* Conformational transitions of a neurotensin receptor 1–Gi1 complex. *Nature* **572**, 80–85 (2019).

20.   Russo, C. J. & Passmore, L. A. Ultrastable gold substrates for electron cryomicroscopy. *Science (80-.).* (2014). doi:10.1126/science.1259530

21.   Wu, H. *et al.* Structure of the human κ-opioid receptor in complex with JDTic. *Nature* (2012). doi:10.1038/nature10939

22.   Van Bever, W. F. M., Niemegeers, C. J. E. & Schellekens, K. H. L. P. A. J. N 4 substituted 1 (2 arylethyl) 4 piperidinyl N phenylpropanamides, a novel series of extremely potent analgesics with unusually high safety margin. *Arzneimittel-Forschung/Drug Res.* (1976).

23.   Robertson, M. J., Meyerowitz, J. M., Panova, O., Borrelli, K. W., Skiniotis, G. Plasticity in Ligand Recognition at Somatostatin Receptors. (2021)

24.   Tunyasuvunakool, K. *et al.* Highly accurate protein structure prediction for the human proteome. *Nature* (2021). doi:10.1038/s41586-021-03828-1

25.   Baek, M. *et al.* Accurate prediction of protein structures and interactions using a three-track neural network. *Science (80-.).* (2021). doi:10.1126/science.abj8754

26.   Kong, H., Raynor, K., Yasuda, K., Bell, G. I. & Reisine, T. Mutation of an aspartate at residue 89 in somatostatin receptor subtype 2 prevents Na+ regulation of agonist binding but does not alter receptor-G protein association. *Mol. Pharmacol.* (1993).

27.   Martin, S., Botto, J. M., Vincent, J. P. & Mazella, J. Pivotal role of an aspartate residue in sodium sensitivity and coupling to G proteins of neurotensin receptors. *Mol. Pharmacol.* (1999). doi:10.1124/mol.55.2.210

28. Yabaluri, N. & Medzihradsky, F. Regulation of μ-opioid receptor in neural cells by
extracellular sodium. *J. Neurochem.* (1997). doi:10.1046/j.1471-4159.1997.68031053.x

29. Fenalti, G. DRUG DISCOVERY Molecular control of delta -opioid receptor signalling. *Nat.*
-LONDON- (2014).

30. Towns, J. *et al.* XSEDE: Accelerating scientific discovery. *Comput. Sci. Eng.* (2014).
doi:10.1109/MCSE.2014.80

31. Zivanov, J. *et al.* RELION-3: New tools for automated high-resolution cryo-EM structure
determination. *bioRxiv* (2018). doi:10.1101/421123

32. Zheng, S. Q. *et al.* MotionCor2: Anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nature Methods* **14**, 331–332 (2017).

33. Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).

34. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. CryoSPARC: Algorithms for
rapid unsupervised cryo-EM structure determination. *Nat. Methods* (2017).
doi:10.1038/nmeth.4169

35. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci.* (2021). doi:10.1002/pro.3943

36. Manglik, A. *et al.* Crystal structure of the μ-opioid receptor bound to a morphinan
antagonist. *Nature* (2012). doi:10.1038/nature10954

37. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr. Sect. D Biol. Crystallogr. **66**, 486–501 (2010).

38. Adams, P. D. *et al.* PHENIX: A comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 213–221 (2010).

39. Robertson, M. J., van Zundert, G. C. P., Borrelli, K. & Skiniotis, G. GemSpot: A Pipeline for
Robust Modeling of Ligands into Cryo-EM Maps. *Structure* (2020).
doi:10.1016/j.str.2020.04.018

40.   Terwilliger, T. C., Ludtke, S. J., Read, R. J., Adams, P. D. & Afonine, P. V. Improvement of cryo-EM maps by density modification. *Nat. Methods* (2020). doi:10.1038/s41592-020-0914-9

41.   Lomize, M. A., Pogozheva, I. D., Joo, H., Mosberg, H. I. & Lomize, A. L. OPM database and PPM web server: Resources for positioning of proteins in membranes. *Nucleic Acids Res.* **40**, (2012).

42.   Lee, J. *et al.* CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory Comput.* **12**, 405–413 (2016).

43.   Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935 (1983).

44.   Robertson, M. J., Tirado-Rives, J. & Jorgensen, W. L. Improved Peptide and Protein Torsional Energetics with the OPLS-AA Force Field. *J. Chem. Theory Comput.* **11**, 3499–3509 (2015).

45.   Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J. Mol. Graph.* **14**, 33–38 (1996).

46.   Jacobson, M. P. *et al.* A Hierarchical Approach to All-Atom Protein Loop Prediction. *Proteins Struct. Funct. Genet.* (2004). doi:10.1002/prot.10613

47.   Phillips, J. C. *et al.* Scalable molecular dynamics with NAMD. *Journal of Computational Chemistry* **26**, 1781–1802 (2005).

48.   Ryckaert, J. P., Ciccotti, G. & Berendsen, H. J. C. Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *J. Comput. Phys.* (1977). doi:10.1016/0021-9991(77)90098-5

49.   Miyamoto, S. & Kollman, P. A. Settle: An analytical version of the SHAKE and RATTLE algorithm for rigid water models. *J. Comput. Chem.* (1992). doi:10.1002/jcc.540130805

50.   Friesner, R. A. *et al.* Extra precision glide: Docking and scoring incorporating a model of

649   hydrophobic enclosure for protein-ligand complexes. *J. Med. Chem.* **49**, 6177–6196

650   (2006).

651