SPCS: A Spatial and Pattern Combined Smoothing Method for Spatial Transcriptomic Expression

Yusong Liu[a], Tongxin Wang[b], Ben Duggan[c], Michael Sharpnack[d], Kun Huang[e, f],

Jie Zhang*[g], Xiufen Ye*[a], and Travis S. Johnson*[e,h]

[a] College of Intelligent Systems Science and Engineering, Harbin Engineering University, Harbin, Heilongjiang 150001, China

[b] Department of Computer Science, Indiana University Bloomington, Bloomington, IN 47408, USA

[c] Department of Medicine, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[d] Department of Pathology, University of California San Francisco, San Francisco, CA 94143, USA

[e] Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[f] Regenstrief Institute, Indianapolis, IN 46202, USA

[g] Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[h] Indiana Biosciences Research Institute, Indianapolis, IN 46202, USA

* Indicate corresponding authors.

Corresponding authors:

Travis S. Johnson

Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Tel: +1 (317) 278-5451

E-mail: johnstrs@iu.edu

Xiufen Ye

College of Intelligent Systems Science and Engineering, Harbin Engineering University, Harbin, Heilongjiang 150001, China

Tel: +86 (451) 8251-9410

E-mail: yexiufen@hrbeu.edu.cn

Jie Zhang

Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Tel: +1 (317) 274-2839

E-mail: jizhan@iu.edu

Declarations of Competing Interests:

None

Abstract

High dimensional, localized RNA sequencing is now possible owing to recent developments in spatial transcriptomics (ST). ST is based on highly multiplexed sequence analysis and uses barcodes to match the sequenced reads to their respective tissue locations. ST expression data suffers from high noise and drop-out events; however, smoothing techniques have the promise to improve the data interpretability prior to performing downstream analyses. Single cell RNA sequencing (scRNA-seq) data similarly suffer from these limitations, and smoothing methods developed for scRNA-seq can only utilize associations in transcriptome space (also known as one-factor smoothing methods). Since they do not account for spatial relationships, these one-factor smoothing methods cannot take full advantage ST data. In this study, we present a novel two-factor smoothing technique, Spatial and Pattern Combined Smoothing (SPCS), that employs k-nearest neighbor technique to utilize information from transcriptome and spatial relationships. By performing SPCS on multiple ST slides from pancreatic ductal adenocarcinoma (PDAC), dorsolateral prefrontal cortex (DLPFC), and simulated high-grade serous ovarian cancer (HGSOC) datasets, smoothed ST slides have better separability, partition accuracy, and biological interpretability than the ones smoothed by pre-existing one-factor methods. Source code of SPCS is provided in Github (https://github.com/Usos/SPCS).

Keywords

Spatial transcriptomics, imputation, two-factor expression smoothing, k-nearest neighbors, tissue region partition, pancreatic ductal adenocarcinoma, dorsolateral prefrontal cortex, high-grade serous ovarian cancer

Introduction

Mamallian tissue is highly heterogeneous with phenotypes that depend on their spatial distribution [1, 2]. Until recently, studies of tissue heterogeneity have either sacrificed spatial relationships (e.g. scRNAseq) or produced low-dimensional measurements (e.g. IHC) [3-7]. Novel spatial transcriptomics (ST) techniques allow whole transcriptome profiles to be measured while preserving spatial relationships [8, 9]. These techniques have already been profoundly useful in understanding tumor [10-12] and non-tumor tissue [13-16] heterogeneity. However, improvements in ST library preparation [17], sequencing techniques, and bioinformatic analysis pipelines [18] are still necessary and ongoing, in contrast to more establish scRNA-seq standard practice protocols [19, 20].

The most widely utilized ST technologies are based on highly multiplexed sequence barcoding, which suffers from expression noise and drop-out events [21, 22]. Barcoding-based scRNAseq data suffers from similar limitations, while plate- and in vitro transcription-based techniques such as Smart-seq2 [23] and CEL-seq2 [24], respectively, provide more representative expression profiles per cell at the cost of fewer cells measured per experiment. As a result, a multitude of techniques have been developed to impute the missing expression values and smooth the noise that come directly from the barcode-based non-spatial scRNA-seq. SAVER [25] and MAGIC [26] use sets of correlated genes and relative cell similarity in transcriptome space to impute the drop-out events and eliminate other types of expression errors via machine learning techniques. These methods are termed "one-factor methods" given that they only incorporate expression values. The smoothened expression values give more accurate representations of the true underlying RNA abundances than the raw read counts. ST data has the advantage of providing spatial relationships that can be used in addition to

transcriptomic similarity for smoothing, based on the assumption that nearby cells will

have more similar expression profiles than distant cells.

Here we present a novel two-factor smoothing method, termed Spatial and Pattern

Combined Smoothing (SPCS), specifically designed for ST data that utilizes both the

associations of spatial locations in transcriptome space (expression pattern knowledge)

and in Euclidean space (spatial knowledge). By performing SPCS on multiple ST slides

from pancreatic ductal adenocarcinoma (PDAC), dorsolateral prefrontal cortex (DLPFC),

and simulated high-grade serous ovarian cancer (HGSOC) datasets, smoothed ST slides

have better separability, partition accuracy, and biological interpretability than the ones

smoothed by pre-existing one-factor methods.

Methods
91

Datasets
92

The datasets that we use in this study including two real world ST dataset, PDAC [10] and

DLPFC [27], and a simulating dataset generated from HGSOC single-cell dataset [28].

For the two real world datasets, PDAC include 10 ST slides sourced from traditional ST

platform, while DLPFC is a Visium platform dataset with 12 slides. All the data in these

datasets consist of two different matrices containing gene wise expressions and spatial

coordinates. One matrix consists of the gene expression values for each spatial barcode

hybridized from its corresponding spot on the ST slide. The other matrix contains the

spatial locations in 2D space for each spot's spatial barcode. Using these two matrices,

we can generate a 2D representation of each gene's expression value throughout the

biopsied tissue section. Because these datasets sourced from different ST platform, i.e.

traditional ST platform and new developed Visium platform, we can explore the influence

of smoothing methods more comprehensively. A detailed summary statistics of the data

we used is provided in Supplementary Table S1.

To better explore the ability of smoothing methods to deal with outlier spots, we designed

a simulation experiment based on those used in the BayesSpace study [29]. Simulated

ST data is based on HGSOC single-cell dataset and an immunofluorescence stained

image of an ovarian cancer (OC) biopsy. In the original single-cell analysis of the HGSOC

dataset, all the cells were divided into 15 clusters by the DBSCAN clustering method and

annotated [28]. Considering the limited number of cells, we only used some of the slide.

Ground-truth cluster labels were derived from single-cell level annotation of tumor and

stroma compartments within the image. To make the simulated data reflect biology, we

separated the slide into four clusters: intratumor (including dendritic and fibroblast cells),

stroma (corresponding to macrophage cells) and two tumor clusters (associated with two

different maliganent cell clusters). Detailed information about ground truth clusters is

provided in Supplementary Table S2. To test the ability of different smoothing methods to

find outlier spots, we randomly mixed 5% of other cell types as perturbation for each spatial

cluster. We generated 10 sets of simulating data in the simulation analysis.

Spatial and Pattern Combined Smoothing (SPCS) of Spatial

Transcriptomic Expression

For each spot on an ST slide, there exists not only the gene expression but also their

spatial positions. This means we can improve the quality of the expression values within

each specific spot using the relative similarity to the other spots based on both expression

pattern and spatial location on the ST slide. To achieve this goal, we propose a KNN-

based method, Spatial and Pattern Combined Smoothing (SPCS), to perform the

smoothing and padding. We display the procedure of SPCS method in Figure 1.

Figure 1 placement

In our method, we obtained the smoothed expression of each spot by integrating the

contribution-weighted expression of its pattern and spatial neighbors. Let X_i be a vector

of gene expression values for spot i, smoothed expression X_i' can be calculated by:

$$
132 \quad X_i' = (1-\alpha)X_i + \alpha\left(\beta\frac{\sum_{j\in N_s(i)} c_{ji}^s X_j}{\sum_{k\in N_s(i)} c_{ki}^s} + (1-\beta)\frac{\sum_{j\in N_p(i)} c_{ji}^p X_j}{\sum_{k\in N_p(i)} c_{ki}^p}\right) \quad (1)
$$

In eq. (1), there are two ratio parameters α and β. α is used to regularize the ratio of

original expression to the corrected expression, which avoids the expression of the object

spot becoming over-smoothed. β is used to adjust the ratio of spatial-based to pattern-

based smoothing for different applications. $N_s(i)$ and $N_p(i)$ are the spatial and pattern

neighborhood of the object spot i respectively. The size of neighborhoods can be

determined by parameter τ_s and τ_p which also should be specified in advance. c_{ji}^s and c_{ji}^p

represent the spatial and pattern contributions of a corresponding neighbor spot j to the

object spot i. For traditional ST platform based PDAC dataset, we set $\alpha = 0.6, \beta =$

$0.4, \tau_s = 2$ and $\tau_p = 16$; and for the Visium platform based DLPFC and simulating

datasets, we make $\tau_s = 4$ due to the larger number of spots. We will display the influence

of these parameters on data seprability and discuss the selection of them in the Discussion

section. In addition, SPCS will also fill the missing spots using multiple non-missing spatial

neighbors. The source code of our proposed SPCS method is provided in our Github

repository (https://github.com/Usos/SPCS). In the next section, we introduce the detailed

mathematical definitions of neighborhood, neighborhood contribution, and our missing

spot padding strategy.

Pattern neighborhood

ST data is a type of transcriptomic data that measures gene expression patterns similar to scRNA-seq. Like some scRNA-seq data, where cells can be localized to tissue location of origin, spots which have a similar expression pattern are more likely to belong to the same region in a tissue. Therefore, smoothing the expression of a given spot using other spots with similar gene expression can improve data quality and is similar to one-factor smoothing methods [25, 26] designed for scRNA-seq. A group of the most similar spots based on the expression "pattern" of the spot can be defined as that spot's "pattern neighborhood". Here, we provide the explicit definition of the pattern neighborhood used by the SPCS method.

Definition 1 (Pattern neighborhood): S_p is the gene expression pattern space of an ST slide, $i, j, k \in S_p$ are different spots, d_{ij}^p and d_{ik}^p are pattern distance between spots i, j and i, k respectively. $N_p(i)$ is the τ_p pattern neighborhood of spot i if $|N_p(i)| = \tau_p, \forall j \in N_p(i), \forall k \in S_p - (N_p(i) \cup \{i\})$ s.t. $d_{ij}^p < d_{ik}^p$.

For gene expression data, the overall shapes of gene expression patterns are of greater interest than the individual magnitudes of each feature [30]. Hence, we used Pearson correlation distance to measure pattern distance between different spots. Let ρ_{ij} represent Pearson correlation coefficient (PCC) of coordinate of spots i and j in pattern space, Pearson correlation distance d_{ij} of spots i and j is [31]:

$$d_{ij} = 1 - \rho_{ij} \tag{2}$$

In ST data, some genes are expressed at identical or near-identical levels that lack the variance to establish an accurate pattern neighborhood. Therefore, we used principal component analysis (PCA) [32] to transform the expression of spots into a 10-dimensional

principal component (PC) space before smoothing. These uncorrelated components with

the largest variance from our PCA are considered the gene expression pattern space.

Spatial neighborhood

In contrast to scRNA-seq data, ST data provide the spatial position for each spot in the

slide. Regions in proximity on histopathology slides are more likely to be the same tissue

type. Aside from the pattern associations between spots, we can also use spatial

associations to smooth the expression as a second factor. We named the group of spots

that are spatially near a given spot the "spatial neighborhood" of that spot, which is defined

explicitly below.

Definition 2 (Spatial neighborhood): S represents the set of spatial location indices of an

ST slide, $i, j \in S$, d_{ij}^S is the spatial distance between spot i and j. $N_s(i)$ is the τ_s spatial

neighborhood of spot i, if $\forall j \in N_s(i)$ s.t. $d_{ij}^S \leq \tau_s$.

ST spots are spatially distributed in a checkerboard or honeycomb pattern. Due to the

geometric patterns inherent to ST spot layout, manhattan distance is a suitable metric to

measure the spatial distance between spots inside an ST slide. Thus, we chose

Manhattan distance as the spatial distance to define our spatial neighborhood. Due to the

difference of spatial pattern of spots, the shape of spatial neighborhood could be different

in different platform. Figure 1(b) illustrated the shape of second-order neighborhood of

both traditional ST platform (checkerboard spatial pattern) and Visium platform

(honeycomb spatial pattern).

Contribution of neighbors on smoothing

Different neighbors in the spatial or pattern neighborhood, will have different impacts on

the smoothing for a given spot, which we refer as "contribution". Since the definitions of

spatial and pattern distance are different, we model the corresponding contributions in

different ways. The contributions of both spatial and pattern neighbors are still comparable

since the range of both is $[0,1]$. For spots outside the neighborhood (both spatial and

pattern) of object spot, we assigned their corresponding contribution to 0, which means

they have no contribution to smoothing of the object spot.

In pattern space, to better capture global gene expression patterns, we used Pearson

correlation distance described in eq. (2) as the distance metric, whose range is $[0,2]$. If

the expression of two spots has a negative correlation, the distance based on eq. (2) will

become greater than 1. However, smoothing with negative correlation spots is not

performed since they are dissimilar. Therefore, we set the contribution to 0 if pattern

distance between the object spot and one of its neighbors is greater than 1. We used an

exponential transformation to achieve this goal. For object spot i and its pattern neighbor

j, pattern contribution c_{ji}^p can be defined as:

$$c_{ji}^p = \begin{cases} \exp\left(-\left(\dfrac{d_{ij}^p}{\sigma}\right)^2\right) & d_{ij}^p < 1 \\ 0 & otherwise \end{cases} \qquad (3)$$

209   The exponential function in eq. (3) limits the range of c_{ji}^p to $[0,1]$ and ensures c_{ji}^p decreases

210   as d_{ij}^p increases. σ in this equation is a tuning parameter that controls how the pattern

211   contribution decays with pattern distance. When $d_{ij}^p > \frac{3\sigma}{\sqrt{2}}$, c_{ji}^p will quickly decay to 0 [33,

212   34]. Hence, setting σ to $\frac{\sqrt{2}}{3}$ will remove the effect of negative correlation neighbors and we

213   can simplify the eq. (3) to:

$$c_{ji}^p = e^{-4.5\left(d_{ij}^p\right)^2} \qquad (4)$$

215 For spatial neighbors, we used Manhattan distance, an integer greater than 0, as spatial

216 distance to measure their distance to the object spot. Since the inverse proportional

217 function has a similar decay nature as exponential function, we used the inverse of spatial

218 distance as the spatial contribution, i.e.:

$$c_{ji}^s = \frac{1}{d_{ij}^s}$$

219   (5)

In this case, i is the spot being smoothed and j is the spatial neighbor of spot i.

Padding of missing spots

For ST slides, there are two types of missing values. The first type of missing values are

missing genes, which means the spot itself is located in tissue region, but some specific

gene expression is missing. This is the main form of drop-out events and also frequently

happens in single-cell data, which most of the smoothing methods can handle. The second

type are missing spots, i.e. expression of all genes in the spot is 0. This kind of absent

data is unique in ST data and cannot be padded without spatial position information, which

often indicates the spot has been removed due to a quality problem. Hence, it is necessary

to judge whether the missing spot need to be padded. As shown in Figure 1(c), for each

blank spot in a slide, SPCS will only pad the ones whose pre-determined spatial

neighborhood has more than 50% non-blank spots. This criteria ensures that boundary of

the tissue will not be erroneously expanded. Since there's no expression on missing spots

at all, we estimate the expression of missing spots by their spatial neighbors only. Let i

become a missing spot, its expression X_i can be estimated by:

$$X_i = \frac{\sum_{j \in N_s(i)} c_{ji}^s X_j}{\sum_{k \in N_s(i)} c_{ki}^s}$$

235   (6)

In Eq.(6), $N_s(i)$ is τ_s-order spatial neighborhood of spot i, c_{ji}^S and c_{ki}^S are spatial

contribution of spot j and k to spot i. Spot padding is performed after non-missing spot

smoothing. To evalulate ARI score for padding spots, we assign ground truth labels for

them. The ground truth labels of the padded spots are determined by spatial contribution

weighted major voting of their spatial neighbors and manually inspected by a pathology

resident.

Performance evaluation

To evaluate the effectiveness of our proposed SPCS method, we first performed SPCS

and other one-factor smoothing methods (SAVER and MAGIC) on PDAC, DLPFC and

simulated ST datasets. SAVER and MAGIC are two representative one-factor smoothing

methods using different techniques (i.e., statistical model-based and kNN-based). Then

we partitioned both smoothed and the original unsmoothed real world slides using K-

medoids clustering method [35] and judged how well the clusters were separated after

smoothing by internal evaluation of the unsupervised clustering. In addition, we also

explored how the parameters in SPCS will influence the smoothing. Next, we performed

multiple unsupervised clustering methods, including K-medoids, Louvain [36], mclust [37],

BayesSpace [29] and spaGCN [38], on all the smoothed and unsmoothed slides to find

out how well the clusters match the histopathological labels from the corresponding image

as an external evaluation. Simulating data was also used here to detect outliers for each

smoothing method. As a gene filter, we only kept genes with less than 70% zero expressed

spots in our analysis. For normalization, we performed logarithmic count per million (CPM)

normalization before smoothing using SPCS, SAVER, and MAGIC. The distance metric

used during clustering was Pearson correlation distance as described in eq. (2). For each

dataset, number of clusters is pre-determined by ground truth which is provided in

Supplementary Table S1-S2. PCA was performed prior to clustering and the eigenvectors

with top 20 eigenvalues were selected to reduce the dimensions of expression matrices

and enhance the clustering results.

Internal evaluation

For the internal evaluation, silhouette score was used as the evaluation indicator [39].

Silhouette score, a metric whose range is [-1,1], estimates the average distance between

clusters. A greater silhouette score indicates better cluster separations. For an ST spot i,

let $a(i)$ be the average distance between i and all the other ST spots within the same

cluster and $b(i)$ be the smallest average distance between i and all the ST spots with any

other clusters. The silhouette coefficient $S(i)$ of ST spot i can be expressed by:

270
$$S(i) = \frac{b(i) - a(i)}{\max\{a(i), b(i)\}} \tag{7}$$

The silhouette score of an entire ST slide is the average silhouette coefficient of all spots

in it.

In this work, Pearson correlation based measurment in Eq. (2) was used to calculate the

dissimilarity between ST spots from the imputed ST data. Traditionally, Euclidean distance

is used but here we adopted the dissimilarity metric defined in Eq. (2) since our clustering

method is based on PCC distance. Padded spots are excluded in this analysis. To make

the average silhouette coefficient of clustering under different smoothing methods

comparable, we used the same distance matrix which was based on dimensionality

reduced original unsmoothed ST expression. Hence, in our experiments, compared with

unsmoothed slides, an increase or lack of change in silhouette score on smoothed slides

represents the smoothing method has enhanced the original data distribution while a

decrease indicates the smoothing method has corrupted the original data distribution.

External evaluation

For the external evaluation, we obtained the histopathological labels of the ST spots. The

correspondence between smoothed slides and histopathological labels was evaluated at

both clustering and gene expression level. At the clustering level, the imputed ST data is

clustered using multiple clustering algorithms. Technical details of the clustering methods

is shown in Supplementary Table S4. Next, the concordance was evaluated between the

clusters from the imputed ST data and the labels from the corresponding histopathological

images. Then, the distribution of marker gene expression was compared to the locations

of histopathological labels.

The Adjusted Rand Index (ARI) was used to evaluate the similarity between the clustering

results from the imputed ST data and the histopathological labels [40]. For clusters from

the imputed ST data $\{I_1, I_2, I_3, I_4\}$ and the histopathological categories of ST spots

$\{H_1, H_2, H_3, H_4\}$, we denoted n_{ij} as the number of ST spots that are both in I_i and H_j. The

ARI is defined as:

297
$$ARI = \frac{\sum_{ij} \binom{n_{ij}}{2} - \frac{\sum_i \binom{n_i}{2} \sum_j \binom{n_j}{2}}{\binom{n}{2}}}{\frac{1}{2}\left[\sum_i \binom{n_i}{2} + \sum_j \binom{n_j}{2}\right] - \frac{\sum_i \binom{n_i}{2} \sum_j \binom{n_j}{2}}{\binom{n}{2}}} \tag{8}$$

where n_i is the number of ST spots in I_i and n_j is the number of ST spots in H_j. A higher

ARI value indicates that the imputed ST clusters and the histopathological labels are more

similar. In this analysis, padded spots unable to be clustered with the ground truth will be

treated as an error cluster.

In the marker gene evaluation with the PDAC dataset, two marker genes, *PRSS1* and

TM4SF1, were used to compare their expression distribution spatially. Both genes are

protein-coding genes. *PRSS1* encodes a trypsinogen, which is often highly expressed in

normal pancreatic tissues while *TM4SF1* is a common proto-oncogene and highly

expressed in pancreatic cancer among other malignancies [41, 42]. High expression of

TM4SF1 in the cancerous regions of the PDAC dataset has been detected in previous

research [10]. For the DLPFC dataset, three other marker genes, *MOBP*, *PCP4* and

SNAP25, were used in the analysis. Previous research has reported that these genes can

delineate different cortical layers [27].The expression distribution of these representative

genes can stratify the histopathological regions and can be used to measure the accuracy

of that partition.

Biological analysis

To examine how our algorithm aids in informing the biology of a ST sample, we identified

differentially expressed genes (DEGs) and performed gene ontology enrichment analysis

(GOEA). DEGs were identified in the PDAC slides by comparing two groups of spots with

TM4SF1-high (neoplastic tissue) and low expression (non-neoplastic tissue). To define

the two groups, we first linearly transformed the expression values of *TM4SF1* between

[0,1] by dividing each value by the maximum expression value. The spots on each slide

were split into two groups with one having a transformed expression value less than 0.7

(*TM4SF1* under-expressed) and the other greater (*TM4SF1* over-expressed). The cutoff

0.7 was chosen as it can reflect the boundary of the cancer region accurately.

A Kruskal-Wallis test was performed between the over- and under-expressed groups and

the p-values were adjusted using the Benjamini-Hochberg method to account for the

multiple comparisons [43]. Only genes with an adjusted p-value less than 0.05 were

included. In addition, we used logarithmic foldchange (logFC) to determine the up-

regulated and down-regulated events for genes. Genes with a logFC greater than 1 were

considered up-regulated, and down-regulated if the logFC were less than -1.

GOEA was performed on the DEGs using *g:Profiler* [44]. The significance of enriched

terms were tested by cumulative hypergeometric test and p-values were corrected by

g:SCS method [45]. Only terms with an adjusted p-value below 0.05 were reported. All

data sources offered by *g:Profiler* were used including Gene Ontology (GO) [46],

Reactome [47], KEGG [48], WikiPathways [49], Transfac [50], CORUM [51], Human

Protein Atlas [52], and the Human Phenotype Ontology [53]. Heatmaps for each sample

were then generated to compare the terms found by each smoothing algorithm.

The DLPFC slides were processed slightly differently than the PDAC slides. To find the

differentially expressed genes, one layer of the cortex was compared with the other six

layers. For example, the expression values for all the sports in Layer 1 (L1) were compared

to the expression values for the spots in layers L2, L3, L4, L5, L6, and WM (white matter).

A Kruskal-Wallis test was performed between one layer of the cortex and all other layers,

p-values corrected using the Benjamini-Hochberg method, and logFC calculated. This

resulted in 7 sets of comparisons. The same cuttoff of p-value less than 0.05 and logFC

greater than one were used. The significant DEGs were then compared to a set of DLPFC

layer marker genes published by Zeng et al. [54]. Enrichment analysis was performed

using *g:Profiler* in the same way as the PDAC slides were processed.

Results

In this study, we applied our two-factor smoothing method SPCS and two state-of-the-art

one-factor smoothing methods (MAGIC and SAVER) to smooth ST slides. To compare

the performance of different smoothing methods, we evaluated the quality of generated

clusters after performing K-Medoids unsupervised clustering on both smoothed and

unsmoothed expression. The generated clusters and the distribution of marker gene

expression were compared with pre-labeled histopathological partitions to check if the

smoothed expression more accurately reflected the pathology features of the

corresponding images. In addition, DEGs were identified in different regions with respect

to the marker gene *TM4SF1* expression level and GOEA was performed to reveal the

biological meaning from the smoothed data.

Figure 2 placement

Internal Evaluation

The silhouette score indicates average distance between clusters in a slide. Since we

used the same unsmoothed distance matrix in the different smoothing methods, the

silhouette score reflects how well the smoothing methods kept the original data distribution.

For different smoothing methods, a greater silhouette score than an unsmoothed slide

usually represents better separability of the smoothed expressions compared to the

original data distribution. In contrast, a decreased silhouette score indicates the smoothing

method has changed the original data distribution making it less seperable. After clustering

the spots into corresponding clusters, we calculated the silhouette score of ten PDAC

slides and eight DLPFC slides with seven clusters for each smoothing method in Figure

2(a) and Figure S1. In most of the slides from these two datasets, SPCS and MAGIC got

similar or even greater silhouette score to the unsmoothed slide while SAVER got a

significantly lower score. In addition, SPCS had the greatest average silhouette score over

the other smoothing methods, even slightly higher than the unsmoothed slide. This result

indicates that kNN-based smoothing generally can keep the charactristics of original data

distribution resulting in better data separablity.

Figure 2 (b)-(c) illustrated how the parameters influenced the data separability of SPCS

smoothed PDACA1 slide. For the two ratio parameters α and β, while $\beta \leq 0.4$, silhouette

score were not affected much by the changing of α, but the result became more

sensentive to α once β is large. This result indicated that pattern based smoothing can

help keep the original data distribution. In addition, the results reveal that the data

distribution of spatial neighbors is different from the pattern neighbors. For the

neighborhood size parameters, the results show that τ_s and τ_p, have no significant

influence on data separability. However, while $\tau_s = 2$ and $\tau_p = 4$, silhouette score is

significantly lower than other parameter combinations, which indicates neighborhood size

parameters also should be well tuned according to the dataset to avoid unexpected results.

External Evaluation

In the external evaluation, different smoothing methods were evaluated based on the

consistency of unsupervised clusters with histopathological labels. The result of PDACA1

alone was shown in Figure 3 since the histopathological labels were not available for other

PDAC slides. This slide can be well clustered by K-Medoids clustering even without

smoothing. When examining the results in detail, Figure 3(b) shows that most ST spots in

the cancer cells and desmoplasia region were separated from other clusters. Most ST

spots in the duct epithelium region were also well spearated, with slight mixing of the

interstitium and the normal pancreatic tissue regions. It is worth mentioning that by

including spatial position information, SPCS can pad missing spots, which other one-factor

smoothing method cannot do. Figure 3(c, d) show the influence of different smoothing

methods on marker genes expression. Compared with other smoothing methods, SPCS

generated better marker gene contrast between distinctive histopathological areas due to

two reasons. First, marker gene expressions showed fewer drop-outs and better reflected

expression patterns across different tissue regions with SAVER and SPCS (MAGIC does

not impute the missing expressions). Second and in contrast to SAVER, SPCS imputed

the missing values and kept the spatial distribution of non-missing values stable.

Figure 3 placement

Figure 4 illustrates external evaluation results from 12 slides in DLPFC dataset. Figure

4(a) shows the influence of smoothing methods on different clustering methods. To make

the evaluation comprehensive, we chose three commonly used general clustering

methods (i.e. k-Medoids, Louvain and mclust) and two state-of-the-art clustering methods

developed specifically for ST slides (i.e BayesSpace and SpaGCN). The results revealed

that smoothing achieved a higher ARI score using various clustering methods and SPCS

outperform other one-factor smoothing methods for every clustering method. When

combining SPCS with Louvain or mclust clustering methods on the DLPFC dataset, we

got an ARI score near BayesSpace, indicating that the combination of SPCS with general

clustering methods improves the performance of spatial clustering on ST slides. In addition,

combining SPCS and BayesSpace got the highest ARI score among all the combinations.

Due to the difference of preprocessing steps, the results of SpaGCN clustering were

moderately lower than the original publication [38], which indicates that hyperparameter

settings of SpaGCN is sensitive to preprocessing steps and we hope to optimize this in

the future. Figure 4(b) and Figure S2(a) illustrated clustering ground truth and BayesSpace

clustering results combined with different smoothing methods on slide 151675 and 151673.

Compared with other one-factor smoothing or without smoothing, SPCS can achieve more

clear and accurate boundaries between different regions, which contributes to a higher

ARI score. In addition, smaller spatial neighborhood (τ_s) for SPCS can help to capture

long narrow regions in the slide but may cause over clustering in thicker regions with

similar length and width. Similar to the PDAC dataset, SPCS also helps to obtain a better

marker gene contrast between distinctive cortical layers in DLPFC dataset as shown in

Figure S2(b-d).

<u>Figure 4 placement</u>

We also performed simulation analysis to test the clustering accuracy for different

smoothing methods. Louvain and BayesSpace, the two best general and ST dedicated

clustering methods in the DLPFC experiment, were used in this analysis. The clustering

ground truth is shown in Figure 5(a). For the 10 simulated slides, Figure 5(b) illustrated

distribution of ARI score of each combination of smoothing and clustering method. In

general, since the original single cell dataset was well clustered, every combination of

smoothing and clustering methods got high ARI scores in this experiment. With Louvain

clustering, there was only a small difference in ARI scores on different slides. For different

smoothing methods, SPCS gets a slightly higher average ARI score which indicates that

spatial information helps general clustering methods to cluster ST slides. However,

Louvain seperates the spots into five clusters instead of four, which means Louvain tends

to cluster the spots according to cell types. For the BayesSpace experiments, the ARI

scores were lower than Louvain on average and greater in variance. To better review the

higher variance of BayesSpace, we illustrated BayesSpace clustering results of two

simulating slides in Figure 5(c) and (d). Obviously, BayesSpace tends to merge small

clusters into nearby larger ones which leads to oversmoothing in the simulated dataset,

and smoothing with SAVER and SPCS will aggravate this problem. Overall all methods

performed well on the simulated data (ARI > 0.85) but it is worth noting that SPCS has

higher potential ARI as evaluated by the 75^{th} quantile. This is surprising considering the

other smoothing methods are specifically designed for scRNA-seq data from which these

simulated ST slides are generated.

Figure 5 placement

Biological Analysis

The biological interpretability of the smoothed results was compared between different

smoothing methods. Comparing between *TM4SF1 over- and under-expressed* regions for

the PDAC slides, the significant DEG numbers were displayed in Figure 6(a). There were

no DEGs found in any smoothed or unsmoothed slides of PDACB2 and PDACG because

they lacked *TM4SF1* while a higher number of DEGs were detected by SPCS in six out of

the rest eight ST slides. Correspondingly, the number of enriched GO terms were also

more from SPCS than the other methods, as shown by Figure 6(b), which are further

examined below.

The top ten most significant GO terms found in slide PDACA1 for each algorithm are

shown in Figure 6(c) for the up-regulated DEGs and Figure 6(d) for the down-regulated

DEGs. More than ten terms are shown in the heatmap because most GO terms were not

shared between smoothing methods. Without smoothing the slides, the up-regulated

DEGs identified are related to interleukin-1. Instead, the terms found by SPCS smoothing

are related primarily to cell adhesion, extracellular matrix (ECM) organization and MET

activates PTK2 signaling. The GO terms from all smoothed and unsmoothed up-regulated

slides can be seen in Figure S3. *g:Profiler* could not find any enriched terms for slides

PDACB2, PDACD, and PDACG. Up-regulated enriched terms for all SPCS slides except

PDACB1 appear similar to PDACA1. The results for the other methods had a small

number of terms which do not seem related to PDAC, except for slide PDACF which

contained cell adhesion and ECM terms for the no smoothing and MAGIC.

Figure 6 placement

The enriched GO terms for the down-regulated DEGs (Figure 6(d)) tell a different story.

Without smoothing, GO terms related to digestion as well as infection and autoimmune

related pathways and complement cascade components (C3, C5) were identified. GO

terms related to digestion were seen with SAVER as well. After performing smoothing, we

found many more GO terms related to the ECM, more complement binding, as well as

apoptosis regulation, but the infection and autoimmune-related pathways were absent.

Applying MAGIC and SPCS also helped to find other complement cascade components

such as C1q complex binding. The other slides showed are in Figure S3 and have similar

results to the terms described above.

Figure 6(e) contains terms enriched in the SPCS smoothed data from up-regulated DEGs

that have previously been reported in PDAC [55-58]. These terms, such as cell adhesion,

cadherin binding, PI3K-Akt signaling pathway and focal adhesion, are all significantly

enriched in DEGs from SPCS-smoothed data (albeit not among the top-ten enriched

terms), but were absent from unsmoothed data, reflecting the enhancement of biological

interpretability by performing our novel SPCS smoothing method.

The results of the biological analysis of the DLPFC slides are shown in Figure S4. The

significant DEGs are shown in Figure S4(a) and show a similar number of DEGs for each

algorithm. When comparing the DEGs to those reported by Zeng et al. [54], a similar

number of genes were shared for each smoothing method as shown in Figure S4(b).

These genes include *MOBP* (an oligodendroctye and white matter marker), *PCP4*

(associated with L5 and L6) [27, 54], and *GFAP* (associated with astrocytes and many

neurological disorders) [59]. In addition, comparing with other smoothing methods, SPCS

helped to discover five DEGs (i.e., *CADM4*, *ELOVL5*, *AGR2*, *LGALS3BP* and *SCARB2*)

which the authors have not found reported in the DLPFC. These genes were found in the

white matter layers of the brain. The expression of *CADM4* and *SCARB2* in sample

151673 are shown in Figure S4(c) and Figure S4(d), respectively.

Discussion

Importance of smoothing on ST data

ST data is based on highly multiplexed sequence analysis where barcodes are used to

split the sequenced reads into their respective tissue locations. However, this type of

sequencing suffers from high noise and drop-out events. To keep enough genes to

perform biological analysis, we set a relaxed filtering threshold (less than 70% zero

expressed spots) to filter out non-expressed genes in preprocessing steps. Even with this

rather relaxed threshold, less than 10% of genes in slides of both PDAC and DLPFC

datasets were left, which indicated that drop-out events are highly frequent in ST datasets.

By visualizing the expression distribution of two marker genes, *PRSS1* and *TM4SF1* in

Figure 3, the gene-level drop-out events can be easily seen, as indicated by the black-

colored spots on the slides. For both genes, drop-out events frequently occurred in

interstitium regions of the slide, leading to a failure cluster this area on unsmoothed ST

data. In addition, there's also entire spots missing in multiple ST slides, which can

negatively influence spatial clustering. By performing smoothing methods, the missing and

noisy expression values were controlled to a certain extent so that the partition of regions

and downstream analysis are greatly improved. Therefore, smoothing is an important and

necessary step for analyzing ST data.

SPCS Improves Data Quality

Compared with unsmoothed data, smoothing improves data quality. We have

demonstrated that various smoothing algorithms increase the separability and partition

accuracy of ST spots. Moreover, by performing internal and external evaluations, we

confirm that SPCS-smoothed data show better quality as compared with the two existing

one-factor smoothing methods MAGIC and SAVER. For internal evaluation, from the

results shown in Figure 2, SPCS smoothing method produces greater silhouette scores

than MAGIC and SAVER, which means the ST data smoothed by SPCS has better

separability. In addition, a more similar silhouette score of SPCS smoothed and original

unsmoothed data indicates that SPCS can better preserve the original data distribution

which helps to keep accurate biological analysis results while improving spatial clustering

accuracy.

Our external evaluation verifies the partition accuracy of smoothed data. One main

objective of slide partition is to identify different histopathological regions in the slide.

Hence, we measured the degree of overlap between unsupervised clusters on smoothed

data and histopathological partitions using ARI. Since the spots in the same

histopathological region are usually connected, incorporating spatial knowledge is

expected to boost partition accuracy. Moreover, spatial knowledge can also help to detect

and pad missing spots. Indeed, as expected, results in Figure 3-5 reveal SPCS method

generates a higher ARI score than existing one-factor methods, which means a more

accurate histopathological partition can be acquired by performing two-factor SPCS

method. In addition, it is also clear that SPCS can be used before various clustering

methods to improve clustering accuracy. From the marker gene analysis, SPCS method

recovered the drop-out events and enhanced the expressions of marker genes in the

corresponding regions. This evidence proves that SPCS can improve the accuracy of the

ST spot partitions.

SPCS Enhances Biological Interpretability

Our SPCS method identified many more differentially expressed genes than the other

smoothing methods tested for most of the ST slides. *IL1RN*, *KRT7*, *LAMB3*, *LAMC2*,

NOTCH3, and S100A16 were identified as upregulated DEGs in the PDACA1 slide after

SPCS processing, and each of these genes has been associated with poor survival in

PDAC [56, 63]. *IL1RN* and *LAMB3* were also found using unsmoothed data and *LAMB3*

was detected with MAGIC smoothed data, but the other genes were uniquely detected by

SPCS. A higher number of DEGs associated with GO terms after SPCS processing than

unsmoothed, SAVER, and MAGIC processing. Specifically, only slides PDACE and

PDACF contained up-regulated DEGs and GO for SAVER and MAGIC; although PDACD

was the only slide which had DEGs and no GO terms for the down-regulated genes. The

unsmoothed up-regulated slides PDACA2, PDACB3, PDACD, and PDACE, and down-

regulated slides PDACA3, PDACB3, and PDACD also had DEGs but no GO terms. In

contrast, every SPCS smoothed slides that had DEGs also had GO terms. The SPCS

data demonstrated a *TM4SF1* expression landscape that matched the original

histopathological assignment more accurately than the other smoothing methods.

The GO terms reported in PDACA1 using SPCS related more to pancreatic cancer and

the role of *TM4SF1* than the non-smoothed data and the two one-factor smoothed data.

TM4SF1 has been found to be overexpressed in PDAC and has roles in apoptosis,

proliferation, and cell migration [41]. Previous work found that collagen 1 (GO:0030199)

binds with DDR1 which then interacts with TM4SF1 to activate the focal adhesion kinase

(FAK) [41, 60, 61]. This results in disruption of E-cadherin (GO:0045296) leading to the

Wnt signaling pathway and loss of cell-cell adhesion (GO:0050839) [41, 60]. The FAK

pathway also increases expression of N-cadherin (GO:0045296) which results in migration

564 of cancer cells [41, 60]. TM4SF1 can also activate the *AKT* pathway (KEGG:04151)

565 leading to anti-apoptotic effects and angiogenesis [41, 62]. All of these previously

566 identified GO terms are important factors for PDAC survival and metastasis. They were

567 also identified using SPCS-smoothed data only, which would have been missed using

568 other smoothing methods or with the unsmoothed ST data.

569 The terms found through GOEA were more interpretable in the scenario of PDAC and its

570 pathogenesis when applying SPCS smoothing. Many of the SPCS top ten terms in Figure

571 4(c) and previously reported terms in Figure 4(e) are similar to those found in the literature

572 [55-58] and are consistent between slides. Without applying SPCS, some of the top terms

573 found in Figure 4(d) are involved in typical pancreatic activity such as peptidase regulator

574 activity (GO:0061134), digestion (GO:0007586), and triglyceride lipase activity

575 (GO:0004806) which indicates that important PDAC-pathology related GO terms may be

576 missed when data is not smoothed or not properly smoothed.

577 Identifying the DEGs in different histological regions and checking their associated GO

578 terms can help evaluate the biological interpretability of a smoothed slide. It is worth noting

579 that the gene ontology (i.e. gene set database) and the enrichment tool can yield different

580 results. The simpliest example would be the use of a hypergeometric test to determine

581 enriched gene sets opposed to gene set enrichment analysis which accounts for the

582 significance of DEGs. The hypergeometric test has a longer history of use and is easily

583 interpretable whereas gene set enrichment analysis is a newer approach. Furthermore,

584 the gene sets themselves differ between databases such as KEGG and GO. There could

585 potentially be more enriched KEGG pathways for one sample and more enriched GO

586 terms for another. For these reasons, we used g:Profiler since it incorporates many gene

587 set databases in the analysis and relies on the well-established hypergeometric testing

588 approach.

Using the cortical layers in the DLPFC slides, we found a similar number of DEGs and

shared genes between the smoothing methods. Most of these genes were previously

reported. Due to the enhanced contrast, our proposed SPCS method helped to identify

five DEGs in the white matter while the other methods did not. The Cell Adhesion Molecule

4 (*CADM4*) and scavenger receptor class B Member 2 (*SCARB2*), to the author's

knowledge, have not been identified as white matter markers in the DLPFC. Cell adhesion

molecules (CAMs) like *CADM4* play an important role in myelination by oligodendrocytes.

Higher expression of *CADM4* leads to many short myelin internodes that disrupt the

normal myelination process [64]. *SCARB2* is a lysosomal membrane receptor for the

glucocerebrosidase enzyme. It has been associated with Parkinson's disease and Lewy

Body Disease. Glucocerebrosidase degenerates sphingolipid, which is important for brain

development. There is some evidence that decreases sphingolipids can lead to

demyelination [65, 66]. While the authors do not intend to present *CADM4* and *SCARB2*

as marker genes for white matter, we believe that SPCS can be used to help aid with this

task. The enrichment analysis results for DLPFC are not shown but are similar between

all smoothing methods. This is likely because the number of GO terms found is correlated

with the number of DEGs. It's possible that using a gene markers instead of layers to

identify DEGs could produce different results. Given the generally clearly defined

boundries of the brain, using layer data to get differential gene expression seems more

appropriate.

Determination of SPCS parameters

There are four parameters in SPCS: τ_s, τ_p, α and β. The parameters τ_s and τ_p are

designed to adjust the size of spatial neighborhood and pattern neighborhood respectively.

Including more information while performing smoothing is beneficial for a more robust

result, and increasing the size of neighborhood is a good way to achieve that goal. Blindly

expanding neighborhoods will incorporate some spots that are not similar to the spot being

smoothed; therefore, SPCS uses contribution weighting to reduce this effect, which gives

the size of both spatial and pattern neighborhoods limited influence on data separability

as shown in Figure 2(e). In addition, as shown in Figure 4 (b) and (c), spatial neighborhood

will also influence clustering sensitivity. A smaller spatial neighborhood (τ_s) for SPCS can

help to capture long narrow regions in slides but may cause over clustering in thicker

regions with similar length and width. Therefore, we recommend a modest selection of

these two parameters to balance the tradeoff. For most cases, $\tau_p \leq 16$ and $\tau_s \leq 4$ is

recommended.

The parameters, α and β, are designed to balance the original expression and corrections

from spatial and pattern neighbors, which has a significant effect on smoothing quality.

Due to the pattern similarity between the object spot and its pattern neighbors, corrections

from pattern neighbors tend to enhance the original data distribution features shown by

an increased or stabilized on silhouette score. In contrast, corrections from spatial

neighbors make the object spot expression consistent with its spatial neighbors. This is

beneficial for spatial clustering but may change the original data distribution leading to a

worse silhouette score. To keep the accuracy of spatial clustering and biological analysis

simultaneously, it is important to balance the intensity of correction with the underlying

expression signatures. Results in Figure 2(d) indicate that the influence of smoothing

strength (α) on data separability is heavily reliant on the proportion of spatial and pattern

correction (β). Hence, we recommend setting α between 0.2 to 0.8 and β less than 0.6 in

most cases, and values should be selected carefully according to data distribution. In

addition, as indicating in Figure 5, when combined with spatial clustering methods like

BayesSpace, smaller α and β are recommended to avoid erroroneous merging of small

clusters.

Conclusion

In response to expression noise and drop-out events in barcoding-based sequencing

technologies, smoothing has become an essential data processing step before performing

the downstream analysis on ST data. In this paper, we proposed a novel two-factor ST

data smoothing method, SPCS, which can take full advantage of both the expression

patterns and the spatial patterns contained in ST data. Compared with traditional one-

factor smoothing methods, SPCS improved separability, partition accuracy, and biological

interpretability of ST experiments. SPCS can effectively improve ST data quality for

accurate and meaningful downstream analyses. SPCS is broadly applicable on any

barcoding-based ST technology.

Key Points

•   Due to the common issue of noise and drop-out events in ST data, smoothing has

become a necessary step before downstream analysis on ST data.

•   SPCS is a novel kNN-based two-factor smoothing method that can fully utilize both

expression pattern and spatial knowledge in ST data.

•   Compared with traditional expression pattern knowledge-based one-factor

smoothing methods, SPCS can provide better separability, partition accuracy and

biological interpretability.

Data Availability

658 All of the data used for the analyses in the manuscript are freely available from their

659 original publications.

Funds

661 This work is partially supported by ACS-IRG Grant Mechanism [Grant No. 19-144-34, to

662 T.S.J.], the National Natural Science Foundation of China [Grant No. 41876100, to X.Y.],

663 the State Key Program of National Natural Science Foundation of China [Grant No.

664 61633004, to X.Y.], the Indiana University Precision Health Initiative [to K.H. and J.Z.].

Acknowledgement

666 We special thanks Dr. Edward Zhao and Dr. Raphael Gottardo in Fred Hutchinson Cancer

667 Research Center for providing information and help on simulation experiment design.

Description of Authors

Yusong Liu is a PhD student in the College of Intelligent Systems Science and Engineering,

Harbin Engineering University. His research is focus on machine learning and network

analysis in bioinformatics.

Dr. Tongxin Wang is a recently graduated PhD student from the Department of Computer

Science, Indiana University, who is currently employed as a Research Scientist at

Facebook. His research focus is on deep learning, transfer learning, and adversarial

networks.

Ben Duggan is a medical student at Indiana University School of Medicine. His research

interests include bioinformatics, machine learning, and using computing to improve patient

care.

Dr. Michael Sharpnack is a pathology resident in the Department of Pathology at the
Univeristy of California San Francisco. His research is primarily concerned with novel
immunotherapies, antigen presentation, and bioinformatics methods development as it
applies to lung cancer.

Dr. Kun Huang is a Professor and Chair of the Department of Biostatistics and Health Data
Science, Indiana University School of Medicine, Indiana University School of Medicine
Precision Health Initiative Chair for Genomic Data Science, Director of Data Science and
Informatics for the Precision Health Initiative, Associate Director of Data Science for the
Indiana University Simon Comprehensive Cancer Center, and an Investigator at the
Regenstrief Institute. His main research interests are in medical image analysis, multi-
omics, and machine learning.

Dr. Jie Zhang is an Assistant Professor of Medical and Molecular Genetics who is also a
member of the Center for Computational Biology and Bioinformatics. Her research
interests are applied translational bioinformatics and systems biology for cancer and
neurological disease.

Dr. Xiufen Ye is a professor of College of Intelligent Systems Science and Engineering,
Harbin Engineering University, IEEE Senior Member. Her main research interest includes
image processing, pattern recognition and artificial intelligent.

Dr. Travis S. Johnson is an Assistant Research Professor in the Department of
Biostatistics and Health Data Science, Indiana University School of Medicine. His research
interests include applied machine learning for use with high dimensional omic data as it
applies to cancer and dementia research.

 # Figures

Figure 1 Workflow of our proposed SPCS smoothing method. (a) An example of

smoothing. In this sample, SPOT 1 is the object spot that is going to be smoothed. For ST

slides, preprocessing steps normalize the expressions and filter out genes with zero

expression in most spots. Spatial neighborhood is second order neighborhood of SPOT 1

with nine spots. Spots where all gene expression is 0 are treated as non-tissue reigons

and excluded from spatial neighborhood. Pattern neighborhood consists of the top eight

spots with most similar expression pattern to SPOT1. Both spatial and pattern

contributions are normalized which means total contribution of all neighbors is equal to 1.

Smoothing is performed by integrating parameter- and contribution-weighted expression

of the object spot itself and both it's spatial and pattern neighbors. (b) Shape of second

order spatial neighborhood for ST and Visium platforms. Spatial neighborhood is

determined by Manhattan distance. Spots with same Manhattan distance to object spot

belong to same order of spatial neighborhood. Due to the different geometries of spots,

the shape of spatial neighborhood could be different in different platforms. (c) Padding

strategy of SPCS. A missing spot will not be padded unless there are more than 50% of

unmissing spots inside its spatial neighborhood.

Figure 2 Influence of smoothing on data separability and data distribution. (a) Box plot of average silhouette score of ten PDAC samples and eight 7-layer DLPFC samples without smoothing and with smoothing by different methods (MAGIC, SAVER and SPCS). (b) Influence of parameters α and β on average silhouette score for SPCS smoothed PDACA1 slide when $\tau_p = 16, \tau_s = 2$. (c) Influence of parameters τ_p and τ_s on average silhouette score for SPCS smoothed PDACA1 slide when $\alpha = 0.6, \beta = 0.4$.

Figure 3 Influence of smoothing on clustering accuracy in PDACA1. (a) Original ST slide of PDACA1 and its histopathological partitions. (b) Results of K-Medoids clustering on expression smoothed by different methods. ARI score, which reflects the correlation between clusters and histopathological labels, is marked at bottom-right corner of each figure. Clusters are ordered by their size. Heatmap of smoothed expression of two marker genes, (c) *PRSS1* and (d) *TM4SF1*, are also shown. For demonstration purposes, expressions of genes are linear transformed into range of 0 to 1 as normalization.

Figure 4 Influence of smoothing on clustering accuracy in DLPFC dataset. (a) Box plot of ARI score on various combination of smoothing (No smoothing, MAGIC, SAVER and SPCS) and clustering methods (k-medoids, Louvain, mclust, BayesSpace and SpaGCN) in 12 DLPFC samples. (b) Ground truth label and BayesSpace clustering results of smoothed and unsmoothed sample 151675. Clusters are ordered by their size.

Figure 5 Influence of smoothing on clustering accuracy in HGSOC simulating dataset. (a)
Ground truth labels of a simulated ST slide. (b) Box plot of ARI score on Louvain and
BayesSpace clustered 10 smoothed simulated ST slides. (c) and (d) illustrate BayesSpace
clustering results of two of the simulated ST slides. Clusters are ordered by their size.

Figure 6 Biological analysis of unsmoothed and different methods (MAGIC, SAVER and

SPCS) smoothed PDAC slides. (a) Number of DEGs identified in each slide and (b) the

number of GO terms found from GOEA in each slide shown on a log10 scale. DEGs which

are down-regulated in *TM4SF1* over-expressed region and their corresponding GOEA

terms are marked as slash filled texture. The ten most significantly enriched terms from

each smoothing method found in the (c) up-regulated DEGs and (d) down-regulated DEGs

groups in *TM4SF1* over-expressed region for PDACA1 slide are shown, shaded by -

log10(p-value). (e) GO terms enriched in SPCS smoothed slides from up-regulated DEGs

that have previously been found in PDAC.

Reference

1.   Heindl A, Sestak I, Naidoo K et al. Relevance of Spatial Heterogeneity of Immune
Infiltration for Predicting Risk of Recurrence After Endocrine Therapy of ER+ Breast
Cancer, J Natl Cancer Inst 2018;110.

2.   Valkonen M, Ruusuvuori P, Kartasalo K et al. Analysis of spatial heterogeneity in
normal epithelium and preneoplastic alterations in mouse prostate tumor models, Sci Rep
2017;7:44831.

3.   Yeong   J,   Tan   T,   Chow   ZL   et   al.   Multiplex
immunohistochemistry/immunofluorescence (mIHC/IF) for PD-L1 testing in triple-negative
breast cancer: a translational assay compared with conventional IHC, J Clin Pathol
2020;73:557-562.

4.   Gillett C, Fantl V, Smith R et al. Amplification and overexpression of cyclin D1 in
breast cancer detected by immunohistochemical staining, Cancer Res 1994;54:1812-
1817.

5.   Gorsch SM, Memoli VA, Stukel TA et al. Immunohistochemical staining for
transforming growth factor beta 1 associates with disease progression in human breast
cancer, Cancer Res 1992;52:6949-6952.

6.   Liang H, Li H, Xie Z et al. Quantitative multiplex immunofluorescence analysis
identifies infiltrating PD1(+) CD8(+) and CD8(+) T cells as predictive of response to
neoadjuvant chemotherapy in breast cancer, Thorac Cancer 2020;11:2941-2954.

7.   Sun Z, Nyberg R, Wu Y et al. Developing an enhanced 7-color multiplex IHC
protocol to dissect immune infiltration in human cancers, PLoS One 2021;16:e0247238.

8.   Burgess DJ. Spatial transcriptomics coming of age, Nat Rev Genet 2019;20:317.

9.   Marx V. Method of the Year: spatially resolved transcriptomics, Nat Methods
2021;18:9-14.

10.   Moncada R, Barkley D, Wagner F et al. Integrating microarray-based spatial
transcriptomics and single-cell RNA-seq reveals tissue architecture in pancreatic ductal
adenocarcinomas, Nat Biotechnol 2020;38:333-342.

11.   He B, Bergenstråhle L, Stenbeck L et al. Integrating spatial gene expression and
breast tumour morphology via deep learning, Nat Biomed Eng 2020;4:827-834.

12.   Berglund E, Maaskola J, Schultz N et al. Spatial maps of prostate cancer
transcriptomes reveal an unexplored landscape of heterogeneity, Nat Commun
2018;9:2419.

13.   Chen WT, Lu A, Craessaerts K et al. Spatial Transcriptomics and In Situ
Sequencing to Study Alzheimer's Disease, Cell 2020;182:976-991.e919.

14.   Maynard KR, Collado-Torres L, Weber LM et al. Transcriptome-scale spatial gene

expression in the human dorsolateral prefrontal cortex, Nat Neurosci 2021;24:425-436.

15.   Ortiz C, Navarro JF, Jurek A et al. Molecular atlas of the adult mouse brain, Sci
Adv 2020;6:eabb3446.

16.   Vickovic S, Eraslan G, Salmén F et al. High-definition spatial transcriptomics for in
situ tissue profiling, Nat Methods 2019;16:987-990.

17.   Berglund E, Saarenpää S, Jemt A et al. Automation of Spatial Transcriptomics
library preparation to enable rapid and robust insights into spatial organization of tissues,
BMC Genomics 2020;21:298.

18.   Dries R, Zhu Q, Dong R et al. Giotto: a toolbox for integrative analysis and
visualization of spatial expression data, Genome Biol 2021;22:78.

19.   Haque A, Engel J, Teichmann SA et al. A practical guide to single-cell RNA-
sequencing for biomedical research and clinical applications, Genome Med 2017;9:75.

20.   Linnarsson S, Teichmann SA. Single-cell genomics: coming of age, Genome Biol
2016;17:97.

21.   Ståhl PL, Salmén F, Vickovic S et al. Visualization and analysis of gene expression
in tissue sections by spatial transcriptomics, Science 2016;353:78-82.

22.   Ding J, Adiconis X, Simmons SK et al. Systematic comparison of single-cell and
single-nucleus RNA-sequencing methods, Nat Biotechnol 2020;38:737-746.

23.   Picelli S, Björklund Å K, Faridani OR et al. Smart-seq2 for sensitive full-length
transcriptome profiling in single cells, Nat Methods 2013;10:1096-1098.

24.   Hashimshony T, Senderovich N, Avital G et al. CEL-Seq2: sensitive highly-
multiplexed single-cell RNA-Seq, Genome Biol 2016;17:77.

25.   Huang M, Wang J, Torre E et al. SAVER: gene expression recovery for single-cell
RNA sequencing, Nat Methods 2018;15:539-542.

26.   van Dijk D, Sharma R, Nainys J et al. Recovering Gene Interactions from Single-
Cell Data Using Data Diffusion, Cell 2018;174:716-729.e727.

27.   Maynard KR, Collado-Torres L, Weber LM et al. Transcriptome-scale spatial gene
expression in the human dorsolateral prefrontal cortex, Nature Neuroscience
2021;24:425-436.

28.   Izar B, Tirosh I, Stover EH et al. A single-cell landscape of high-grade serous
ovarian cancer, Nature Medicine 2020;26:1271-1279.

29.   Zhao E, Stone MR, Ren X et al. Spatial transcriptomics at subspot resolution with
BayesSpace, Nature Biotechnology 2021;39:1375-1384.

30.   Jiang DX, Tang C, Zhang AD. Cluster analysis for gene expression data: A survey,
Ieee Transactions on Knowledge and Data Engineering 2004;16:1370-1386.

31.   Berthold MR, Höppner F. On clustering time series using euclidean distance and
pearson correlation, arXiv preprint arXiv:1601.02213 2016.

32.   Wold S, Esbensen K, Geladi P. Principal component analysis, Chemometrics and
intelligent laboratory systems 1987;2:37-52.

33.   Liu Y, Ye X, Zhan X et al. TPQCI: A topology potential-based method to quantify
functional influence of copy number variations, Methods 2021;192:46-56.

34.   Wang Z, Chen Z, Zhao Y et al. A community detection algorithm based on topology
potential and spectral clustering, The Scientific World Journal 2014;2014.

35.   Struyf A, Hubert M, Rousseeuw P. Clustering in an Object-Oriented Environment,
Journal of Statistical Software 1997;1:1 - 30.

36.   Waltman L, van Eck NJ. A smart local moving algorithm for large-scale modularity-
based community detection, The European Physical Journal B 2013;86:471.

37.   Scrucca L, Fop M, Murphy TB et al. mclust 5: Clustering, Classification and Density
Estimation Using Gaussian Finite Mixture Models, R Journal 2016;8:289-317.

38.   Hu J, Li X, Coleman K et al. SpaGCN: Integrating gene expression, spatial location
and histology to identify spatial domains and spatially variable genes by graph
convolutional network, Nature Methods 2021;18:1342-1351.

39.   Shahapure KR, Nicholas C. Cluster Quality Analysis Using Silhouette Score. In:
2020 IEEE 7th International Conference on Data Science and Advanced Analytics (DSAA).
2020, p. 747-748.

40.   Santos JM, Embrechts M. On the use of the adjusted rand index as a metric for
evaluating supervised classification. In: International conference on artificial neural
networks. 2009, p. 175-184. Springer.

41.   Fu F, Yang X, Zheng M et al. Role of Transmembrane 4 L Six Family 1 in the
Development and Progression of Cancer, Front Mol Biosci 2020;7:202.

42.   Liu Q, Guo L, Zhang S et al. PRSS1 mutation: a possible pathomechanism of
pancreatic carcinogenesis and pancreatic cancer, Mol Med 2019;25:44.

43.   Dal Molin A, Baruzzo G, Di Camillo B. Single-Cell RNA-Sequencing: Assessment
of Differential Expression Analysis Methods, Frontiers in Genetics 2017;8.

44.   Raudvere U, Kolberg L, Kuzmin I et al. g: Profiler: a web server for functional
enrichment analysis and conversions of gene lists (2019 update), Nucleic acids research
2019;47:W191-W198.

45.   Reimand J, Kull M, Peterson H et al. g:Profiler--a web-based toolset for functional
profiling of gene lists from large-scale experiments, Nucleic Acids Res 2007;35:W193-200.

46.   Consortium GO. The Gene Ontology resource: enriching a GOld mine, Nucleic
Acids Res 2021;49:D325-d334.

47.   Jassal B, Matthews L, Viteri G et al. The reactome pathway knowledgebase,
Nucleic Acids Res 2020;48:D498-d503.

48.   Kanehisa M, Furumichi M, Sato Y et al. KEGG: integrating viruses and cellular
organisms, Nucleic acids research 2020;49:D545-D551.

49.   Martens M, Ammar A, Riutta A et al. WikiPathways: connecting communities,
Nucleic acids research 2020;49:D613-D621.

50.   Wingender E. The TRANSFAC project as an example of framework technology
that supports the analysis of genomic regulation, Brief Bioinform 2008;9:326-332.

51.   Giurgiu M, Reinhard J, Brauner B et al. CORUM: the comprehensive resource of
mammalian protein complexes-2019, Nucleic Acids Res 2019;47:D559-d563.

52.   Colwill K, Gräslund S. A roadmap to generate renewable protein binders to the
human proteome, Nat Methods 2011;8:551-558.

53.   Köhler S, Gargano M, Matentzoglu N et al. The Human Phenotype Ontology in
2021, Nucleic Acids Res 2021;49:D1207-d1217.

54.   Zeng H, Shen Elaine H, Hohmann John G et al. Large-Scale Cellular-Resolution
Gene Profiling in Human Neocortex Reveals Species-Specific Molecular Signatures, Cell
2012;149:483-496.

55.   Zhou J, Hui X, Mao Y et al. Identification of novel genes associated with a poor
prognosis in pancreatic ductal adenocarcinoma via a bioinformatics analysis, Biosci Rep
2019;39.

56.   Atay S. Integrated transcriptome meta-analysis of pancreatic ductal
adenocarcinoma and matched adjacent pancreatic tissues, PeerJ 2020;8:e10141.

57.   Liu L, Wang S, Cen C et al. Identification of differentially expressed genes in
pancreatic ductal adenocarcinoma and normal pancreatic tissues based on microarray
datasets, Mol Med Rep 2019;20:1901-1914.

58.   Shang M, Zhang L, Chen X et al. Identification of hub genes and regulators
associated with pancreatic ductal adenocarcinoma based on integrated gene expression
profile analysis, Discov Med 2019;28:159-172.

59.   Brenner M. Role of GFAP in CNS injuries, Neuroscience Letters 2014;565:7-13.

60.   Weniger M, Honselmann KC, Liss AS. The Extracellular Matrix and Pancreatic
Cancer: A Complex Relationship, Cancers (Basel) 2018;10.

61.   Yang JC, Zhang Y, He SJ et al. TM4SF1 Promotes Metastasis of Pancreatic
Cancer via Regulating the Expression of DDR1, Sci Rep 2017;7:45895.

62.   Zheng B, Ohuchida K, Cui L et al. TM4SF1 as a prognostic marker of pancreatic
ductal adenocarcinoma is involved in migration and invasion of cancer cells, Int J Oncol
2015;47:490-498.

63. Ye J, Wen J, Ning Y et al. Higher notch expression implies poor survival in
pancreatic ductal adenocarcinoma: A systematic review and meta-analysis,
Pancreatology 2018;18:954-961.

64. Elazar N, Vainshtein A, Golan N et al. Axoglial Adhesion by Cadm4 Regulates
CNS Myelination, Neuron 2019;101:224-231.e225.

65. Alcalay RN, Levy OA, Wolf P et al. SCARB2 variants and glucocerebrosidase
activity in Parkinson's disease, npj Parkinson's Disease 2016;2:16004.

66. Giussani P, Prinetti A, Tringali C. The role of Sphingolipids in myelination and
myelin stability and their involvement in childhood and adult demyelinating disorders,
Journal of Neurochemistry 2021;156:403-414.

909