SPCS: A Spatial and Pattern Combined Smoothing Method for Spatial Transcriptomic Expression

Yusong Liu[a], Tongxin Wang[b], Ben Duggan[c], Kun Huang[d, e], Jie Zhang*[f],

Xiufen Ye*[a], and Travis S. Johnson*[d]

[a] College of Intelligent Systems Science and Engineering, Harbin Engineering University, Harbin, Heilongjiang 150001, China

[b] Department of Computer Science, Indiana University Bloomington, Bloomington, IN 47408, USA

[c] Department of Medicine, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[d] Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[e] Regenstrief Institute, Indianapolis, IN 46202, USA

[f] Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

* Indicate corresponding authors.

Corresponding authors:

Travis S. Johnson

Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Tel: +1 (317) 278-5451

E-mail: johnstrs@iu.edu

Xiufen Ye

College of Intelligent Systems Science and Engineering, Harbin Engineering University, Harbin, Heilongjiang 150001, China

Tel: +86 (451) 8251-9410

E-mail: yexiufen@hrbeu.edu.cn

Jie Zhang

Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Tel: +1 (317) 274-2839

E-mail: jizhan@iu.edu

Declarations of Competing Interests:

None

Abstract

The recently developed spatial transcriptomics (ST) technique has made it possible to view spatial transcriptional heterogeneity in a high throughput manner. It is based on highly multiplexed sequence analysis and uses barcodes to split the sequenced reads into respective tissue locations. However, this type of sequencing technique suffers from high noise and drop-out events in the data, which makes smoothing a necessary step before performing downstream analysis. Traditional smoothing methods used in the similar single cell RNA sequencing (scRNA-seq) data are one-factor methods that can only utilize associations in transcriptome space. Since they do not account for associations in the Euclidean space, i.e. tissue location distances on the ST slide, these one-factor methods cannot take full advantage of all the knowledge in ST data. In this study, we present a novel two-factor smoothing technique, Spatial and Pattern Combined Smoothing (SPCS), that employs k-nearest neighbor technique to utilize associations from transcriptome and Euclidean space from the ST data. By performing SPCS on multiple ST slides from pancreatic ductal adenocarcinoma (PDAC), dorsolateral prefrontal cortex (DLPFC) and simulated high-grade serous ovarian cancer (HGSOC) datasets, smoothed ST slides have better separability, partition accuracy, and biological interpretability than the ones smoothed by pre-existing one-factor methods. Source code of SPCS is provided in Github (https://github.com/Usos/SPCS).

Keywords

Spatial transcriptomics, imputation, two-factor expression smoothing, k-nearest neighbors, tissue region partition, pancreatic ductal adenocarcinoma, dorsolateral prefrontal cortex, high-grade serous ovarian cancer

Introduction

Tumors and organ tissues are highly heterogeneous with different phenotypes distributed throughout the tissue [1, 2]. Until recently, this tissue heterogeneity had to be evaluated using histology image analysis or using scRNA-seq. Histology imaging offers a view on spatial tissue heterogeneity whereas scRNA-seq offers insight into the transcriptional heterogeneity. Methods like immunohistochemistry (IHC) labelled histology slides also allow researchers to view protein in tissue images [3-7]. However, IHC is only capable of capturing protein expression level(s) of one or a few proteins at a time. These technologies lacked the ability to view transcriptional heterogeneity spatially in a high throughput manner.

Fortunately, revolutionary new techniques that can be broadly categorized as spatial transcriptomics (ST) have recently become available that allow transcriptome profiles to be measured for different locations in a tissue [8, 9]. These techniques have already been profoundly useful in understanding tumor heterogeneity [10-12] and heterogeneity of brain tissues [13-16]. In recent years, research on this new ST technology has been prolific -- in 2017 there were only five articles mentioning "spatial transcriptomics" in PubMed (https://pubmed.ncbi.nlm.nih.gov/), but by 2020 there were a total of 70 articles, 14 times as many as just three years earlier. However, ST technology is still largely being studied methodologically to improve the library preparation [17], sequencing techniques, and bioinformatic analysis pipelines [18] used in spatial transcriptomic analysis. In comparison, scRNA-seq analysis has now largely established standard practice protocols [19, 20] such that the data preprocessing and initial bioinformatic analyses are consistent between experiments. Once ST techniques has established the standard practice, like single cell technologies, then they will be widely available for any number of disease and normal

tissues.

The most common form of ST data to date, much like the now widely abundant scRNA-

seq data, is based on highly multiplexed sequence analysis where barcodes are used to

sort the sequenced reads into their respective cells or tissue locations [21]. This type of

sequencing suffers from a wide array of noise and drop-out events in the data which has

been well studied across scRNA-seq methodologies [22]. In barcoding-based single cell

data, a single cell's expression profile on its own is not perfectly representative of the true

underlying RNA abundances from that cell due to the noise introduced during the

sequencing process. There are techniques such as Smart-seq2 [23] and CEL-seq2 [24]

that give more representative expression profiles per cell, but these are much lower

throughput, resulting in many fewer cells. As a result, a multitude of techniques have been

developed to impute the missing expression values and smooth the noise that come

directly from the barcode-based scRNA-seq. These techniques help to smooth the data

by using similar cells to correct for technical errors in the sequencing data. Specifically,

sets of correlated genes and relative cell similarity in transcriptome space is used to impute

the drop-out events and eliminate other types of expression errors via machine learning

techniques. Some of these tools include SAVER [25] and MAGIC [26]. When applied, the

processed expression values give more accurate representations of the true underlying

RNA abundances than the raw read counts.

Because most current ST data are also generated from barcode-based technology, it is

not surprising that ST data also suffer from similar noise and drop-out events. This means

that the distributions of gene expression in the tissue does not always follow the

corresponding phenotypic differences in histopathological images at a high resolution. For

drop-out events, it can be assumed that zero expression values have some probability of

being spuriously missed in the reads simply due to chance. However, ST data, unlike

scRNA-seq, not only has relative similarity in transcriptome space but also in Euclidean

space on the tissue slide. That means locations that are adjacent to one another in the ST

array will likely be similar tissue and as a result have similar transcriptome profiles. In

addition, if two locations in an ST array share a majority of their transcriptome profiles,

then their transcriptional differences can be partially explained by drop-out.

Here we present a novel two-factor ST smoothing method that utilizes both the

associations of spatial locations in transcriptome space (expression pattern knowledge)

and in Euclidean space (spatial knowledge). This is in contrast to the pre-existing one-

factor scRNA-seq techniques that only utilize associations in transcriptome space. We

believe that by using both the spatial location of a given ST spot and its transcriptome

profile we can better smooth the data such that it better represents the true underlying

RNA abundances in the tissue.

119   Methods

120   Datasets

The datasets that we use in this study including two real world ST dataset, PDAC [10] and

DLPFC [27], and a simulating dataset generated from HGSOC single-cell dataset [28].

For the two real world datasets, PDAC include 10 ST slides sourced from traditional ST

platform, while DLPFC is a Visium platform dataset with 12 slides. All the data in these

datasets consist of two different matrices containing gene is expressions and spatial

coordinates. One matrix consists of the gene expression values for each spatial barcode

hybridized from its corresponding spot on the ST slide. The other matrix contains the

spatial locations in 2D space for each spot's spatial barcode. Using these two matrices,

we can generate a 2D representation of each gene's expression value throughout the

biopsied tissue section. Because these datasets sourced from different ST platform, i.e.

traditional ST platform and new developed Visium platform, we can explore the influence

of smoothing methods more comprehensively. A detailed summary statistics of the data

we used is provided in Table S1.

To better explore the ability of smoothing methods on deal with outlier spots, we designed

a simulation experiment referred to BayesSpace [29]. Simulating ST data is based on

HGSOC single-cell dataset and an immunofluorescence stain image associated with

ovarian cancer (OC). In the original single-cell analysis of HGSOC dataset, all the cells

were divided into 15 clusters by DBSCAN clustering method and well annotated [28].

Considering the limitation of number of cells, we used only partial of the slide. Ground-

truth cluster labels were derived from single-cell level annotation of tumor and stroma

compartments within the image. To make simulated data reflect the real nature, we

separated the slide into four clusters, including intratumor, stroma and two tumor clusters.

Among the spatial clusters, intratumor cluster including dendritic cell and fibroblast cell

clusters, stroma cluster corresponded to macrophage cell cluster and tumor clusters

associated to two different malignant cell clusters. Detailed information about ground truth

clusters is provided in Table S2. For the propose of testing the anti-outlier spot ability of

different smoothing methods, we randomly mixed 5% of other types of cells as

perturbation for each spatial cluster. We generated 10 groups of simulating data in the

simulation analysis.

Spatial and Pattern Combined Smoothing (SPCS) of Spatial

Transcriptomic Expression

For each spot on an ST slide, there exists not only the gene expression but also their

spatial positions. This means we can improve the quality of the expression values within

each specific spot using the relative similarity to the other spots based on expression

pattern and spatial location on the ST slide. In addition, spatial positions can also help to

fill the missing spots. To achieve this goal, we put forward a KNN-based method, Spatial

and Pattern Combined Smoothing (SPCS), to perform the smoothing and padding. We

display the procedure of SPCS method in Figure 1.

Figure 1 placement

In our method, we obtained the smoothed expression of each spot by integrating the

contribution-weighted expression of its pattern and spatial neighbors. Let X_i be a vector

of gene expression values for spot i, smoothed expression X_i' can be calculated by:

163
$$X_i' = (1 - \alpha)X_i + \alpha \left(\beta \frac{\sum_{j \in N_s(i)} c_{ji}^s X_j}{\sum_{k \in N_s(i)} c_{ki}^s} + (1 - \beta) \frac{\sum_{j \in N_p(i)} c_{ji}^p X_j}{\sum_{k \in N_p(i)} c_{ki}^p} \right) \tag{1}$$

In eq. (1), there are two ratio parameters α and β. α is used to regularize the ratio of

original expression to the corrected expression, which avoids the expression of object spot

becoming over-smoothed. β is used to adjust the ratio of spatial-based to pattern-based

smoothing for different applications. $N_s(i)$ and $N_p(i)$ are deleted spatial and pattern

neighborhood of object spot i respectively. Unless otherwise stated, the neighborhood in

this paper is deleted neighborhood. The size of neighborhoods can be determined by

parameter τ_s and τ_p which also should be specified in advance. c_{ji}^s and c_{ji}^p are spatial and

pattern smoothing contribution of corresponding neighbor spot j of object spot i. For

traditional ST platform based PDAC dataset, we set $\alpha = 0.6, \beta = 0.4, \tau_s = 2$ and $\tau_p = 16$;

and for the Visium platform based DLPFC and simulating datasets, we make $\tau_s = 4$ due

to the more number of spots. We will display the influence of these parameters on data

seprability and discuss the selection of them in the Discussion section. In addition, SPCS

will also fill the missing spots with multiple non-missing spatial neighbors. Source code of

our proposed SPCS method is provided in Github (https://github.com/Usos/SPCS). In the

next section, we introduce detailed mathematical definitions of neighborhood, contribution

and our missing spots padding strategy.

Pattern neighborhood

ST data is a type of transcriptomic data, which means it includes expression patterns

similar to scRNA-seq. Like some scRNA-seq data, where cells can be localized to tissue

location of origin, spots which have a similar expression pattern are more likely to belong

to the same region in a tissue. Therefore, smoothing the expression of a spot with those

of similarly expressing spots can improve data quality and is common in traditional one-

factor smoothing methods [25, 26]. A group of the most similar spots based on the

expression "pattern" of the spot can be defined as that spot's "pattern neighborhood". Here,

we provide the explicit definition of the pattern neighborhood used by the SPCS method.

Definition 1 (Pattern neighborhood): S_p is gene expression pattern space of a ST slide,

$i, j, k \in S_p$ are different spots, d_{ij}^p and d_{ik}^p are pattern distance between spots i, j and i, k

respectively. $N_p(i)$ is τ_p-deleted pattern neighborhood of spot i if $|N_p(i)| = \tau_p, \forall j \in$

$N_p(i), \forall k \in S_p - (N_p(i) \cup \{i\})$ s.t. $d_{ij}^p < d_{ik}^p$.

For gene expression data, the overall shapes of gene expression patterns are of greater

interest than the individual magnitudes of each feature [30]. Hence, we used Pearson

correlation distance to measure pattern distance between different spots. Let ρ_{ij} represent

Pearson correlation coefficient (PCC) of coordinate of spots i and j in pattern space,

Pearson correlation distance d_{ij} of spots i and j is [31]:

$$198 \qquad\qquad d_{ij} = 1 - \rho_{ij} \qquad\qquad\qquad (2)$$

In ST data, some genes are expressed at identical or near-identical levels, which lack of

variance to establish an accurate pattern neighborhood. Therefore, we used principal

component analysis (PCA) [32], to transform the expression of spots into a 10-dimensional

principal component (PC) space before smoothing, to extract uncorrelated components

with the largest variance. We named this PC space as gene expression pattern space.

Spatial neighborhood

In contrast to scRNA-seq data, ST data can provide spatial position for each spot in the

slide. Regions in proximity on histopathology slides are usually the same tissue type.

Aside from the pattern associations between spots, we can also use spatial associations

to smooth the expression as a second factor. We named the group of spots that are

spatially near a given spot the "spatial neighborhood" of that spot, which is defined

explicitly below.

Definition 2 (Spatial neighborhood): S represents the set of spatial location indices of an

ST slide, $i, j \in S$, d_{ij}^s is the spatial distance between spot i and j. $N_s(i)$ is τ_s-deleted

spatial neighborhood of spot i, if $\forall j \in N_s(i)$ s.t. $d_{ij}^s \leq \tau_s$.

ST spots are spatially distributed in a checkerboard or honeycomb pattern. Manhattan

distance is a suitable metric to measure the spatial distance between spots inside an ST

slide. Thus, we chose Manhattan distance as the spatial distance to define our spatial

neighborhood. Due to the difference of spatial pattern of spots, the shape of spatial

neighborhood could be different in different platform. Figure 1(b) illustrated the shape of

second-order neighborhood of both traditional ST platform (checkerboard spatial pattern)

and Visium platform (honeycomb spatial pattern).

221 Contribution of neighbors on smoothing

Different neighbors in the spatial or pattern neighborhood, will have different impacts on

the smoothing for a given spot, which we refer as "contribution". Since the definitions of

spatial and pattern distance are different, we model the corresponding contributions in

different ways. The contributions of both spatial and pattern neighbors are still comparable

since the range of both is [0,1]. For spots outside the neighborhood (both spatial and

pattern) of object spot, we assigned their corresponding contribution to 0, which means

they have no contribution to smoothing of the object spot.

In pattern space, to better caputure global gene expression patterns, we used Pearson

correlation coefficient distance described in eq. (2) as distance metric, whose range is

[0,2]. If the expression of two spots has a negative correlation, the distance based on eq.

(2) will become greater than 1. However, smoothing with negative correlation spots is not

performed since they are dissimilar. Therefore, we set the contribution to 0 if pattern

distance between the object spot and one of its neighbors is greater than 1. We used an

exponential transformation to achieve this goal. For object spot i and its pattern neighbor

j, pattern contribution c_{ji}^{p} can be defined as:

$$c_{ji}^{p} = \begin{cases} \exp\left(-\left(\dfrac{d_{ij}^{p}}{\sigma}\right)^{2}\right) & d_{ij}^{p} < 1 \\ 0 & otherwise \end{cases} \qquad (3)$$

238 The exponential function in eq. (3) limits the range of c_{ji}^{p} to [0,1] and ensures c_{ji}^{p} decreases

239 as d_{ij}^{p} increases. σ in this equation is a parameter to tune the initial position of exponential

240 function descending. When $d_{ij}^{p} > \frac{3\sigma}{\sqrt{2}}$, c_{ji}^{p} will quickly decay to 0 [33, 34]. Hence, setting σ

241 to $\frac{\sqrt{2}}{3}$ will cease the effect of negative correlation neighbors and we can simplify the eq. (3)

242   as:

$$c_{ji}^{p} = e^{-4.5\left(d_{ij}^{p}\right)^{2}} \tag{4}$$

244   For spatial neighbors, we used Manhattan distance, an integer greater than 0, as spatial

245   distance to measure their distance to object spot. Since the inverse proportional function

246   has a similar decay nature as exponential function, we used the inverse of spatial distance

247   as the spatial contribution, i.e.:

$$c_{ji}^{s} = \frac{1}{d_{ij}^{s}} \tag{5}$$

In this case, i is the spot being smoothed and j is the spatial neighbor of spot i.

Padding of missing spots

For ST slides, there are two types of missing values. The first type of missing values is

gene level missing, which means the spot itself is located in tissue region, but some

specific gene expression is missing. This is the main form of drop-out events and also

frequently happen in single-cell data, which most of the smoothing methods can handle.

The second type is spot level missing, i.e. expression of all genes in the spot is 0. This

kind of missing is unique in ST data and cannot be padded without spatial position

information, which often indicates there's no tissue on the spot or the spot has been

removed due to the quality problem. Hence, it is necessary to judge whether the missing

spot need to be padded. As shown in Figure 1(c), for each blank spot in a slide, SPCS will

only pad the ones whose pre-determined spatial neighborhood has more than 50% non-

blank spots. This criteria ensures that boundary of the tissue will not be errorly expanded.

Since there's no expression on missing spots at all, we estimate the expression of missing

spots by their spatial neighbors only. Let i become a missing spot, its expression X_i can

be estimated by:

$$X_i = \frac{\sum_{j \in N_s(i)} c_{ji}^s X_j}{\sum_{k \in N_s(i)} c_{ki}^s} \tag{6}$$

266   In Eq.(6), $N_s(i)$ is τ_s -order spatial neighborhood of spot i, c_{ji}^s and c_{ki}^s are spatial

267   contribution of spot j and k to spot i. Spot padding is performed after non-missing spot

268   smoothing. To evalulate ARI score for padding spots, we assign ground truth labels for

269   them. The ground truth labels of the padded spots are determined by spatial contribution

270   weighted major voting of their spatial neighbors.

Performance evaluation

To evaluate the effectiveness of our proposed SPCS method, we first performed SPCS

and other one-factor smoothing methods (SAVER and MAGIC) on PDAC, DLPFC and

simulated ST datasets. SAVER and MAGIC are two representative one-factor smoothing

methods on different technical routes (i.e., statistical model-based and kNN-based). Then

we partitioned both smoothed and the original unsmoothed real world slides using K-

medoids clustering method [35] and judged how well the clusters can be separated after

performing multiple smoothing methods by internal evaluation of the unsupervised

clustering. In addition, we also explored how the parameters in SPCS will influence the

smoothing. After all, we performed multiple unsupervised clustering methods, including K-

medoids, Louvain [36], mclust [37], BayesSpace [29] and spaGCN [38], on all the

smoothed and unsmoothed slides to find out how well the clusters match the

histopathological labels from the corresponding image as an external evaluation.

Simulating data also was used here to detect the ability of anti-outliers for each smoothing

method. As a gene filter, we only kept genes with less than 70% zero expressed spots in

our analysis. For normalization, we performed logarithmic count per million (CPM)

normalization before smoothing using SPCS, SAVER, and MAGIC. The distance metric

used during clustering was Pearson correlation distance as described in eq. (2). For each

dataset, number of clusters is pre-determined by ground truth which is provided in Table

S1-S2. PCA was performed prior to clustering, then we selected the eigenvectors with top

20 eigenvalues to reduce the dimensions of expression matrices and enhance the

clustering results.

Internal evaluation

For the internal evaluation, silhouette score was used as the evaluation indicator [39].

Silhouette score, a metric whose range is [-1,1], estimates the average distance between

clusters. A greater silhouette score indicates better cluster separations. For a ST spot i,

let $a(i)$ be the average distance between i and all the other ST spots within the same

cluster and $b(i)$ be the smallest average distance between i and all the ST spots with any

other clusters. The silhouette coefficient $S(i)$ of ST spot i can be expressed by:

300
$$S(i) = \frac{b(i) - a(i)}{\max\{a(i), b(i)\}} \tag{7}$$

Silhouette score of an ST slide is the average silhouette coefficient of all spots in it.

In this piece of work, Pearson correlation based measurment in Eq. (2) was used to

calculate the dissimilarity between ST spots from the imputed ST data. Traditionally,

Euclidean distance is used but here we adopted the dissimilarity metric defined in Eq. (2)

since our clustering method is based on Pearson correlation distance. Padded spots are

excluded in this analysis. To make the average silhouette coefficient of clustering under

different smoothing methods comparable, we used the same distance matrix which was

based on dimensionality reduced original unsmoothed ST expression. Hence, in our

experiments, comparing with unsmoothed slides, an increase or lack of change in

silhouette score on smoothed slides represents the smoothing method has enhanced the

original data distribution while a decrease indicates the smoothing method has changed

the distribution without keeping the originl one.

External evaluation

For the external evaluation, we obtained the histopathological labels of the ST spots. The

correspondence between smoothed slides and histopathological labels was evaluated at

both clustering and gene expression level. At the clustering level, the imputed ST data is

clustered using multiple clustering algorithms. Technical details of the clustering methods

is shown in Table S4. Next, the concordance was evaluated between the clusters from the

imputed ST data and the labels from the corresponding histopathological images.

Furthermore, the distribution of marker gene expression was compared to the locations of

histopathological labels.

The Adjusted Rand Index (ARI) was used to evaluate the similarity between the clustering

results from the imputed ST data and the histopathological labels [40]. For clusters from

the imputed ST data $\{I_1, I_2, I_3, I_4\}$ and the histopathological categories of ST spots

$\{H_1, H_2, H_3, H_4\}$, we denoted n_{ij} as the number of ST spots that are both in I_i and H_j. The

ARI is defined as:

327
$$ARI = \frac{\sum_{ij} \binom{n_{ij}}{2} - \frac{\sum_i \binom{n_i}{2} \sum_j \binom{n_j}{2}}{\binom{n}{2}}}{\frac{1}{2}\left[\sum_i \binom{n_i}{2} + \sum_j \binom{n_j}{2}\right] - \frac{\sum_i \binom{n_i}{2} \sum_j \binom{n_j}{2}}{\binom{n}{2}}} \qquad (8)$$

where n_i is the number of ST spots in I_i and n_j is the number of ST spots in H_j. A higher

ARI value indicates that the imputed ST clusters and the histopathological labels are more

similar. In this analysis, unable to cluster padding spots with ground truth will be treated

as error clustered.

In the marker gene evaluation, For PDAC dataset, two marker genes, *PRSS1* and

TM4SF1, were used to compare their expression distribution spatially. Both genes are

protein-coding genes. *PRSS1* encodes a trypsinogen, which is often highly expressed in

normal pancreatic tissues while *TM4SF1* is a common proto-oncogene and highly

expressed in pancreatic cancer among other malignancies [41, 42]. High expression of

TM4SF1 in the cancerous regions of the PDAC dataset has been detected in previous

research [10]. For DLPFC dataset, three other marker genes, *MOBP*, *PCP4* and *SNAP25*,

is used in the analysis. Previous research has reported that these genes can delineate

different cortical layers [27].The expression distribution of these representative genes can

stratify the histopathological regions and can be used to measure the accuracy of that

partition.

Biological analysis

To examine how our algorithm aids in informing the biology of a ST sample, we identified

differentially expressed genes (DEGs) and performed gene ontology enrichment analysis

(GOEA). DEGs were identified in the PDAC slides by comparing two groups of spots with

TM4SF1-high (neoplastic tissue) and low expression (non-neoplastic tissue). To define

the two groups, we first linearly transformed the expression values of *TM4SF1* between

[0,1] by dividing each value by the maximum expression value. The spots on each slide

were split into two groups with one having a transformed expression value less than 0.7

(*TM4SF1* under-expressed) and the other greater (*TM4SF1* over-expressed). The cutoff

0.7 was chosen as it can reflect the boundary of the cancer region accurately.

A Kruskal-Wallis test was performed between the over- and under-expressed groups and

the p-values were adjusted using the Benjamini-Hochberg method to account for the

multiple comparisons [43]. Only genes with an adjusted p-value less than 0.05 were

included. In addition, we used logarithmic foldchange (logFC) to determine the up-

regulated and down-regulated events for genes. Genes with a logFC greater than 1 were

considered up-regulated, and down-regulated if the logFC were less than -1.

GOEA was performed on the DEGs using *g:Profiler* [44]. The significance of enriched

terms were tested by cumulative hypergeometric test and p-values were corrected by

g:SCS method [45]. Only terms with an adjusted p-value below 0.05 were reported. All

data sources offered by *g:Profiler* were used including Gene Ontology (GO) [46],

Reactome [47], KEGG [48], WikiPathways [49], Transfac [50], CORUM [51], Human

Protein Atlas [52], and the Human Phenotype Ontology [53]. Heatmaps for each sample

were then generated to compare the terms found by each smoothing algorithm.

The DLPFC slides were processed slightly differently than the PDAC slides. To find the

differentially expressed genes, one layer of the cortex was compared with the other six

layers. For example, the expression values for all the sports in Layer 1 (L1) were compared

to the expression values for the spots in layers L2, L3, L4, L5, L6, and WM (white matter).

A Kruskal-Wallis test was performed between one layer of the cortex and all other layers,

p-values corrected using the Benjamini-Hochberg method, and logFC calculated. This

resulted in 7 sets of comparisons. The same cuttoff of p-value less than 0.05 and logFC

greater than one were used. The significant DEGs were then compared to a set of DLPFC

layer marker genes published by Zeng et al. [54]. Enrichment analysis was performed

using *g:Profiler* in the same way as the PDAC slides were processed.

Results

In this study, we applied our two-factor smoothing method SPCS and two state-of-the-art one-factor smoothing methods (MAGIC and SAVER) to smooth ten PDAC ST slides. To compare the performance of different smoothing methods, we evaluated the quality of generated clusters after performed K-Medoids unsupervised clustering on both smoothed and unsmoothed expression. The generated clusters and the distribution of marker gene expressions were compared with pre-labeled histopathological partitions to check if the smoothed expressions are more accurately reflecting the pathology features of the corresponding images. In addition, the DEGs were identified in different regions with respect to the marker gene *TM4SF1* expression level and GOEA was performed to reveal the biological meanings revealed by the smoothing.

Figure 2 placement

Internal Evaluation

The silhouette score indicates average distance between clusters in a slide. Since we used a same unsmoothed distance matrix for each slide smoothed by different smoothing methods, silhouette score in this analysis also can reflect how well the smoothing methods keep the original data distribution. For different smoothing methods, a greater silhouette score than unsmoothed slide usually represents smoothed expressions have better separability based on original data distribution. In the contrast, the decreasion of silhouette score indicates the smoothing method has changed the original data distribution. After clustering the spots into corresponding clusters, we calculated the silhouette score of ten PDAC slides and eight DLPFC slides with seven clusters for each smoothing method in Figure 2(a) and Figure S1. In most of the slides in these two datasets, SPCS and MAGIC

got similar or even greater silhouette score to the unsmoothed slide while SAVER got a

significantly lower score. In addition, SPCS got the greatest average silhouette score than

other smoothing methods, even slightly higher than the unsmoothed slide. This result

indicates that kNN-based smoothing generally can keep the charactristics of original data

distribution better and SPCS can contribute to better data separablity.

Figure 2 (b)-(c) illustrated how the parameters will influence the data separability of SPCS

smoothed PDACA1 slide. For the two ratio parameters α and β, while $\beta \leq 0.4$, silhouette

score will not be affect a lot by the changing of α, but the result will become more

sensentive to α if β is large enough. This result indicating that pattern based smoothing

can help to keep the original data distribution. In addition, the results also reveal a truth

that data distribution nature of spatial neighbors is different from the pattern neighbors.

For the neighborhood size parameters τ_s and τ_p, the results show that they will have no

significant influence on data separability.

External Evaluation

In the external evaluation, different smoothing methods were evaluated based on the

consistency of unsupervised clusters with histopathological labels. The result of PDACA1

was shown in Figure 3 since the histopathological labels were available for PDACA1. This

slide can be well clustered by k-Medoids clustering method even without smoothing. When

examining the results in detail, Figure 3(b) shows that most ST spots in the cancer cells

and desmoplasia region can be separated from other clusters. Most ST spots in the duct

epithelium region can also be well spearated, with slight mixing of the interstitium and the

normal pancreatic tissue regions. It is worth mentioning that by including spatial position

information, SPCS can pad missing spots, which other one-factor smoothing method

cannot do. Figure 3(c, d) show the influence of different smoothing methods on the two

marker genes expression. Compared with other smoothing methods, SPCS generated

better marker gene contrast between distinctive histopathological areas due to two

reasons. 1) By performing smoothing, marker gene expressions showed fewer drop-outs

and better reflected expression patterns across different tissue regions with SAVER and

SPCS (MAGIC does not impute the missing expressions). 2) In contrast to SAVER, SPCS

imputed the missing values and kept the spatial distribution of non-missing values

relatively stable.

Figure 3 placement

Figure 4 illustrates external evaluation results from 12 slides in DLPFC dataset. Figure

4(a) shows the influence of smoothing methods on different clustering methods. To make

the evaluation comprehensive, we chose three commonly used general clustering

methods (i.e. k-Medoids, Louvain and mclust) and two state-of-the-art clustering methods

developed specifically for ST slides (i.e BayesSpace and SpaGCN). The results revealed

that smoothing helps to get a higher ARI score for various clustering methods and SPCS

outperform other one-factor smoothing methods for every clustering method. When

combining SPCS and Louvain or mclust clustering methods, we can get a ARI score near

BayesSpace for DLPFC dataset, which means the combination of SPCS with general

clustering methods is good to performing spatial clustering on ST slides. In addition,

combining of SPCS and BayesSpace got the highest ARI score among all the

combinations. Due to the difference of preprocessing steps, the results of SpaGCN

clustering were moderately lower than the original publication [38], which indicates that

hyperparameter settings of SpaGCN is sensitive to preprocessing steps and we hope to

optimize this in the future. Figure 4(b) and Figure S2(a) illustrated clustering ground truth

and BayesSpace clustering results while combining with different smoothing methods on

slide 151675 and 151673. Compared with other one-factor smoothing or without

smoothing, SPCS can help to get a more clear and accurate boundaries between different

regions, which contribute to a higher ARI score. In addition, smaller spatial neighborhood

(τ_s) for SPCS can help to capture long narrow regions in the slide but may cause over

clustering in thicker regions with similar length and width. Similar to PDAC dataset, SPCS

also helps to obtain a better marker gene contrast between distinctive cortical layers in

DLPFC dataset as shown in Figure S2(b-d).

Figure 4 placement

We also performed simulation analysis to test the clustering accuracy for different

smoothing methods. Louvain and BayesSpace, the two best general and ST dedicated

clustering methods in DLPFC experiment, were used in this analysis. Clustering ground

truth is shown in figure 5(a). For the 10 simulated slides, Figure 5(b) illustrated distribution

of ARI score of each combination of smoothing and clustering method. In general, since

the original single cell dataset was well clustered, every combination of smoothing and

clustering methods got high ARI score in this experiment. On the track of Louvain

clustering, there's only a little difference of ARI score on different slides. For different

smoothing methods, SPCS gets a slightly higher average ARI score which indicates that

spatial information helps general clustering methods to better clustering ST slides.

However, Louvain seperates the spots into five clusters instead four, which means

Louvain tends to cluster the spots according to cell types. For the BayesSpace

experiments, the ARI score was lower than Louvain in average and greater in variance.

To better review the higher variancde of BayesSpace, we illustrated BayesSpace

clustering results of two simulating slides in Figure 5(c) and (d). Obviously, BayesSpace

tends to merge small clusters into nearby larger ones which leads to oversmoothing in the

simulating dataset, and SAVER and SPCS smoothing will aggravate this problem. Overall

all methods performed well on the simulated data (ARI > 0.85) but it is worth noting that

SPCS has higher potential ARI as evaluated by the 75th quantile. This is surprising

considering the other smoothing methods are specifically designed for scRNA-seq data

from which these simulated ST slides are generated.

Figure 5 placement

Biological Analysis

The biological interpretability of the smoothed results was compared between different

smoothing methods. When comparing between *TM4SF1 over- and under-expressed*

regions for the PDAC slides, the significant DEG numbers were displayed in Figure 6(a).

There were no DEGs found in any smoothed or unsmoothed slides of PDACB2 and

PDACG because they lacked *TM4SF1* while a higher number of DEGs was detected by

SPCS in six out of the rest eight ST slides. Correspondingly, the number of enriched GO

terms was also more from SPCS than the other methods, as shown by Figure 6(b), which

are further examined below.

The top ten most significant GO terms found in slide PDACA1 for each algorithm are

shown in Figure 6(c) for the up-regulated DEGs and Figure 6(d) for the down-regulated

DEGs. More than ten terms are shown in the heatmap because most GO terms were not

shared between smoothing methods. Without smoothing the slides, the up-regulated

DEGs identified are related to interleukin-1. Instead, the terms found by SPCS smoothing

are related primarily to cell adhesion, extracellular matrix (ECM) organization and MET

activates PTK2 signaling. The GO terms from all smoothed and unsmoothed up-regulated

slides can be seen in Figure S3. *g:Profiler* could not find any enriched terms for slides

PDACB2, PDACD, and PDACG. Up-regulated enriched terms for all SPCS slides except

PDACB1 appear similar to PDACA1. The results for the other methods had a small

number of terms which do not seem related to PDAC, except for slide PDACF which

contained cell adhesion and ECM terms for the no smoothing and MAGIC.

Figure 6 placement

The enriched GO terms for the down-regulated DEGs (Figure 6(d)) tell a different story.

Without smoothing, GO terms related to digestion as well as infection and autoimmune

related pathways and complement cascade components (C3, C5) were identified. GO

terms related to digestion were seen with SAVER as well. After performing smoothing, we

found many more GO terms related to the ECM, more complement binding, as well as

apoptosis regulation, but the infection and autoimmune-related pathways were absent.

Applying MAGIC and SPCS also helped to find other complement cascade components

such as C1q complex binding. The other slides showed are in Figure S3 and appear

similar results to the terms described above.

Figure 6(e) contains terms enriched in the SPCS smoothed data from up-regulated DEGs

that have previously been reported in PDAC [55-58]. These terms, such as cell adhesion,

Cadherin binding, PI3K-Akt signaling pathway and focal adhesion, are all significantly

enriched in DEGs from SPCS-smoothed data (albeit not among the top-ten enriched

terms), but were absent from unsmoothed data, reflecting the enhancement of biological

interpretability by performing our novel SPCS smoothing method.

The results of the biological analysis of the DLPFC slides are shown in Figure S4. The

significant DEGs are shown in Figure S4(a) and show a similar number of DEGs for each

algorithm. When comparing the DEGs to those reported by Zeng et al. [54], a similar

number of genes were shared for each smoothing method as shown in Figure S4(b).

These genes include *MOBP* (an oligodendroctye and white matter marker), *PCP4*

(associated with L5 and L6) [27, 54], and *GFAP* (associated with astrocytes and many

neurological disorders) [59]. In addition, comparing with other smoothing methods, SPCS

helped to discover five DEGs (i.e., *CADM4*, *ELOVL5*, *AGR2*, *LGALS3BP* and *SCARB2*)

which not previously reported. These genes were all found in the white matter layers of

the brain. The expression of *CADM4* and *SCARB2* in sample 151673 are shown in Figure

S4(c) and Figure S4(d), respectively.

Discussion

Importance of smoothing on ST data

ST data is based on highly multiplexed sequence analysis where barcodes are used to

split the sequenced reads into their respective tissue locations. However, this type of

sequencing suffers from high noise and drop-out events. To keep enough genes to

perform biological analysis, we set a relaxed filtering threshold (less than 70% zero

expressed spots) to filter out non-expressed genes in preprocessing steps. Even with this

rather relaxed threshold, less than 10% of genes in slides of both PDAC and DLPFC

datasets were left, which indicated that drop-out events are highly frequent in ST datasets.

By visualizing the expression distribution of two marker genes, *PRSS1* and *TM4SF1* in

Figure 3, the gene-level drop-out events can be easily seen, as indicated by the black-

colored spots on the slides. For both genes, drop-out events frequently occurred in

interstitium regions of the slide, leading to a failure cluster this area on unsmoothed ST

data. In addition, there's also entire spots missing in multiple ST slides, which can

negatively influence spatial clustering. By performing smoothing methods, the missing and

noisy expression values were controlled to a certain extent so that the partition of regions

and downstream analysis are greatly improved. Therefore, smoothing is an important and

necessary step for analyzing ST data.

SPCS Improves Data Quality

Compared with unsmoothed data, smoothing step improves data quality. We have demonstrated that various smoothing algorithms increase the separability and partition accuracy of ST spots. Moreover, by performing internal and external evaluations, we confirm that SPCS-smoothed data show better quality on both aspects as compared with the two existing one-factor smoothing methods MAGIC and SAVER. For internal evaluation, from the results shown in Figure 2, SPCS smoothing method produces greater silhouette scores than MAGIC and SAVER, which means the ST data smoothed by SPCS has better separability. In addition, a more similar silhouette score of SPCS smoothed and original unsmoothed data indicates that SPCS can better preserve original data distribution of non-missing values, which helps to keep an accurate biological analysis results while improving spatial clustering accuracy.

Our external evaluation verifies the partition accuracy of smoothed data. One main objective of slide partition is to identify different histopathological regions in the slide. Hence, we measured the degree of overlap between unsupervised clusters on smoothed data and histopathological partitions using ARI. Since the spots in the same histopathological region are usually connected, incorporating spatial knowledge is expected to boost partition accuracy. Moreover, spatial knowledge can also help to detect and pad missing spots. Indeed, as expected, results in Figure 3-5 reveal SPCS method generates a higher ARI score than existing one-factor methods, which means a more accurate histopathological partition can be acquired by performing two-factor SPCS method. In addition, it is also cleared that SPCS can be used before various clustering methods to improve clustering accuracy. From the marker gene analysis, SPCS method recovers the drop-out events and enhance the expressions of marker genes in the

567   corresponding regions. This evidence proves that SPCS can improve the accuracy of the

568   ST spot partitions.

SPCS Enhances Biological Interpretability

The GO terms reported in PDACA1 from data using the SPCS smoothing method are

more interpretable in the context of pancreatic cancer and the role of *TM4SF1* in cancer

than non-smoothed data and the two one-factor smoothed data. *TM4SF1* has been found

to be overexpressed in pancreatic cancer and has roles in apoptosis, proliferation, and

cell migration [41]. Previous work found that collagen 1 (GO:0030199) binds with DDR1

which then interacts with TM4SF1 to activate the focal adhesion kinase (FAK) [41, 60, 61].

This results in disruption of E-cadherin (GO:0045296) leading to the Wnt signaling

pathway and loss of cell-cell adhesion (GO:0050839) [41, 60]. The FAK pathway also

increases expression of N-cadherin (GO:0045296) which results in migration of cancer

cells [41, 60]. TM4SF1 can also activate the *AKT* pathway (KEGG:04151) leading to anti-

apoptotic effects and angiogenesis [41, 62]. All of these previously identified GO terms

are important factors for PDAC survival and metastasis. They were also identified using

SPCS-smoothed data only, which would have been missed using other smoothing

methods or with the unsmoothed ST data.

Identifying the DEGs in different histological regions and check their associated GO terms

can also help evaluate the biological interpretability of a smoothed slide. It is worth noting

that the gene ontology (i.e. gene set database) and the enrichment tool can yield different

results. The simpliest example would be the use of a hypergeometric test to determine

enriched gene sets opposed to gene set enrichment analysis which accounts for the

significance of DEGs. The hypergeometric test has a longer history of use and is easily

interpretable whereas gene set enrichment analysis is a newer approach. Furthermore,

the gene sets themselves differ between databases such as KEGG and GO. There could

potentially be more enriched KEGG pathways for one sample and more enriched GO

terms for another. For these reasons, we use g:Profiler since it incorporates many gene

set databases in the analysis and relies on the well-established hypergeometric testing

approach.

In our analyses we chose to use Our SPCS method contributed to identify many more

differentially expressed genes than the other smoothing methods for most of the slides.

Among the DEGs, the up regulation of *IL1RN*, *KRT7*, *LAMB3*, *LAMC2*, *NOTCH3, and*

S100A16, which have previously been reported associate with poor survival in PDAC, was

also detected in the SPCS smoothed PDACA1 slide [56, 63]. *IL1RN* and *LAMB3* were

also found using unsmoothed data and *LAMB3* was detected with MAGIC smoothed data,

but the other genes were uniquely detected by SPCS. Generally, a high number of DEGs

also associate with more GO terms. However, some unsmoothed, SAVER, and MAGIC-

smoothed slides contained DEGs but no GO terms, which may indicate the insufficient

imputing or smoothing without considering the location information. Specifically, only

slides PDACE and PDACF contained up-regulated DEGs and GO for SAVER and MAGIC;

although PDACD was the only slide which had DEGs and no GO terms for the down-

regulated genes. The unsmoothed up-regulated slides PDACA2, PDACB3, PDACD, and

PDACE, and down-regulated slides PDACA3, PDACB3, and PDACD also had DEGs but

no GO terms. In contrast, every SPCS smoothed slides that had DEGs also had GO terms.

The SPCS data demonstrated a *TM4SF1* expression landscape that matched the original

histopathological assignment more accurately than the other smoothing methods. For this

reason, it's possible SPCS found more DEGs because other genes in the slide had more

defined expression after smoothing.

The terms found through GOEA were more interpretable in the scenario of PDAC and its

pathogenesis when applying SPCS smoothing. Many of the SPCS top ten terms in Figure

4(c) and previously reported terms in Figure 4(e) are similar to those found in the literature

[55-58] and are consistent between slides. Without applying SPCS, some of the top terms

found in Figure 4(d) are involved in typical pancreatic activity such as peptidase regulator

activity (GO:0061134), digestion (GO:0007586), and triglyceride lipase activity

(GO:0004806) which indicates that important PDAC-pathology related GO terms may be

missed when data is not smoothed or not properly smoothed.

Using the cortical layers in the DLPFC slides, we found a similar number of DEGs and

shared genes between the smoothing methods. Most of these genes were previously

reported. However, profiting from contrast enhancement, our proposed SPCS method

helped to identify five DEGs in the white matter while the other methods did not. The Cell

Adhesion Molecule 4 (*CADM4*) and scavenger receptor class B Member 2 (*SCARB2*), to

the author's knowledge, have not been identified as white matter markers in the DLPFC.

Cell adhesion molecules (CAMs) like *CADM4* play an important role in myelination by

oligodendrocytes. Higher expression of *CADM4* leads to many short myelin internodes

that disrupt the normal myelination process [64]. *SCARB2* is a lysosomal membrane

receptor for the glucocerebrosidase enzyme. It has been associated with Parkinson's

disease and Lewy Body Disease. Glucocerebrosidase degenerates sphingolipid, which is

important for brain development. There is some evidence that decreases sphingolipids

can lead to demyelination [65, 66]. While the authors do not intend to present *CADM4* and

SCARB2 as marker genes for white matter, we believe that SPCS can be used to help aid

with this task. The enrichment analysis results for DLPFC are not shown but are similar

between all smoothing methods. This is likely because the number of GO terms found is

correlated with the number of DEGs. It's possible that using a gene markers instead of

layers to identify DEGs could produce different results. Given the generally clearly defined

boundries of the brain, using layer data to get differential gene expression seems more

appropriate.

Determination of SPCS parameters

There are four parameters in SPCS: τ_s, τ_p, α and β. These parameters affect the

performance of SPCS algorithm by adjusting neighborhood size and smoothing intensity

and should be tuned according to the quality and distribution of original ST data.

τ_s and τ_p are designed to adjust the size of spatial neighborhood and pattern

neighborhood, respectively. Obviously, including more information while performing

smoothing is beneficial for a more robust result, and increasing the size of neighborhood

is a good way to achieve that goal. However, Considering blindly expanding

neighborhoods will incorporate some spots that are not similar to the spot being smoothed,

SPCS method use contribution weighting to reduce this effect, which gives the size of both

spatial and pattern neighborhoods limited influence on data separability as shown in

Figure 2(e). In addition, as shown in Figure 4 (b) and (c), spatial neighborhood will also

influence clustering sensitivity. A smaller spatial neighborhood (τ_s) for SPCS can help to

capture long narrow regions in slides but may cause over clustering in thicker regions with

similar length and width. Therefore, we recommend a modest selection of these two

parameters to balance the tradeoff. For most cases, $\tau_p \leq 16$ and $\tau_s \leq 4$ is recommended.

The parameters, α and β, are designed to balance the original expression and corrections

from spatial and pattern neighbors, which has a significant effect on smoothing quality.

Due to the pattern similarity between object spot and its pattern neighbors, corrections

from pattern neighbors trend to enhance the original data distribution features, which can

help to increase or stabilize on silhouette score. In the contrast, corrections from spatial

neighbors try to make the object spot expression consistent to its spatial neighbors, which

is beneficial to spatial clustering but may change the original data distribution leading to a

worse silhouette score. To keep the accuracy of spatial clustering and biological analysis

simultaneously, it is important to balance the intensity of correction with the underlying

expression signatures. Results in Figure 2(d) indicate that the influence of smoothing

strength (α) on data separability is heavily reliant on the proportion of spatial and pattern

correction (β). Hence, we recommend setting α between 0.2 to 0.8 and β less than 0.6 in

most cases and should be selected carefully according to data distribution. In addition, as

indicating in Figure 5, when combining with spatial clustering methods like BayesSpace,

smaller α and β are recommended to avoid erroroneous merging of small clusters.

Conclusion

Due to the limitation of barcoding based sequencing technologies which result in noisy

and high drop-out in transcriptomics data, smoothing has become an essential data

processing step before performing the downstream analysis on ST data. In this paper, we

proposed a novel two-factor ST data smoothing method SPCS, which can take full

advantage of both the expression patterns and the spatial patterns contained in ST data.

Compared with traditional one-factor smoothing methods, we have demonstrated that

SPCS can recover the missing or erroneous expression in ST data better and improve the

separability, partition accuracy, and biological interpretability. This reflects that our

proposed SPCS method can effectively improve data quality of ST and contribute to more

accurate and meaningful downstream analysis results. SPCS smoothing is a state-of-the-

art ST smoothing algorithm with implications in numerous diseases where ST data is being

generated.

Key Points

- Due to the common issue of noise and drop-out events in ST data, smoothing has become a necessary step before downstream analysis on ST data.
- SPCS is a novel kNN-based two-factor smoothing method that can fully utilize both expression pattern and spatial knowledge in ST data.
- Compared with traditional expression pattern knowledge-based one-factor smoothing methods, SPCS can provide better separability, partition accuracy and biological interpretability.

Data Availability

All of the data used for the analyses in the manuscript are freely available from their original publications.

Funds

This work is partially supported by ACS-IRG Grant Mechanism (Grant Number: 19-144-34) [to T.S.J.], the National Natural Science Foundation of China [Grant No. 41876100, to X.Y.], the State Key Program of National Natural Science Foundation of China [Grant No. 61633004, to X.Y.], the Indiana University Precision Health Initiative [to K.H. and J.Z.].

Acknowledgement

We special thanks Dr. Edward Zhao and Dr. Raphael Gottardo in Fred Hutchinson Cancer Research Center for providing information and help on simulation experiment design.

Description of Authors

Yusong Liu is a PhD student in College of Intelligent Systems Science and Engineering, Harbin Engineering University. His research is focus on machine learning and network analysis in bioinformatics.

Dr. Tongxin Wang is a recently graduated PhD student from the Department of Computer Science, Indiana University, who is currently employed as a Research Scientist at Facebook. His research focus is on deep learning, transfer learning, and adversarial networks.

Ben Duggan is a medical student at Indiana University School of Medicine. His research interests include bioinformatics, machine learning, and using computing to improve patient care.

Dr. Kun Huang is a Professor and Chair of the Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indiana University School of Medicine Precision Health Initiative Chair for Genomic Data Science, Director of Data Science and Informatics for the Precision Health Initiative, Associate Director of Data Science for the Indiana University Simon Comprehensive Cancer Center, and an Investigator at the Regenstrief Institute. His main research interests are in medical image analysis, multi-omics, and machine learning.

Dr. Jie Zhang is an Assistant Professor of Medical and Molecular Genetics who is also a member of the Center for Computational Biology and Bioinformatics. Her research interests are applied translational bioinformatics and systems biology for cancer and neurological disease.

Dr. Xiufen Ye is a professor of College of Intelligent Systems Science and Engineering,

Harbin Engineering University, IEEE Senior Member. Her main research interest includes

image processing, pattern recognition and artificial intelligent.

Dr. Travis S. Johnson is an Assistant Research Professor in the Department of

Biostatistics and Health Data Science, Indiana University School of Medicine. His research

interests include applied machine learning for use with high dimensional omic data as it

applies to cancer and dementia research.

Figures

Figure 1 Workflow of our proposed SPCS smoothing method. (a) An example of

smoothing. In this sample, SPOT 1 is the object spot that going to be smoothed. For ST

slides, preprocessing steps normalize the expressions and filter out genes with zero

expression in most spots. Spatial neighborhood is second order neighborhood of SPOT 1

with nine spots. Spots where all gene expression is 0 are treated as non-tissue reigons

and excluded from spatial neighborhood. Pattern neighborhood consists of the top eight

spots with most similar expression pattern to SPOT1. Both spatial and pattern

contributions are normalized which means total contribution of all neighbors is equal to 1.

Smoothing is performed by integrating parameter- and contribution-weighted expression

of the object spot itself and both it's spatial and pattern neighbors. (b) Shape of second

order spatial neighborhood for ST and Visium platforms. Spatial neighborhood is

determined by Manhattan distance. Due to the difference of spatial pattern of spots, the

shape of spatial neighborhood could be different in different platforms. (c) Padding

strategy of SPCS. A missing spot will not be padded unless there are more than 50% of

unmissing spots inside its spatial neighborhood.

Figure 2 Influence of smoothing on data separability and data distribution. (a) Box plot of average silhouette score of ten PDAC samples and eight 7-layer DLPFC samples without smoothing and with smoothing by different methods (MAGIC, SAVER and SPCS). (b) Influence of parameters α and β on average silhouette score for SPCS smoothed PDACA1 slide when $\tau_p = 16, \tau_s = 2$. (c) Influence of parameters τ_p and τ_s on average silhouette score for SPCS smoothed PDACA1 slide when $\alpha = 0.6, \beta = 0.4$.

Figure 3 Influence of smoothing on clustering accuracy in PDACA1. (a) Original ST slide of PDACA1 and its histopathological partitions, (b) Results of K-Medoids clustering on expression smoothed by different methods. ARI score, which reflects the correlation between clusters and histopathological labels, is marked at bottom-right corner of each figure. Clusters are ordered by their size. Heatmap of smoothed expressions of two marker genes, (c) *PRSS1* and (d) *TM4SF1*, are also shown. For demonstration purposes, expressions of genes are linear transformed into range of 0 to 1 as normalization.

Figure 4 Influence of smoothing on clustering accuracy in DLPFC dataset. (a) Box plot of ARI score on various combination of smoothing (No smoothing, MAGIC, SAVER and SPCS) and clustering methods (k-medoids, Louvain, mclust, BayesSpace and SpaGCN) in 12 DLPFC samples. (b) Ground truth label and BayesSpace clustering results of smoothed and unsmoothed sample 151675. Clusters are ordered by their size.

Figure 5 Influence of smoothing on clustering accuracy in HGSOC simulating dataset. (a) Ground truth labels of a simulated ST slide. (b) Box plot of ARI score on Louvain and BayesSpace clustered 10 smoothed simulated ST slides. (c) and (d) illustrate BayesSpace clustering results of two of the simulated ST slides. Clusters are ordered by their size.

Figure 6 Biological analysis of unsmoothed and different methods (MAGIC, SAVER and

SPCS) smoothed PDAC slides. (a) Number of DEGs identified in each slide and (b) the

number of GO terms found from GOEA in each slide shown on a log10 scale. DEGs which

are down-regulated in *TM4SF1* over-expressed region and their corresponding GOEA

terms are marked as slash filling texture. The ten most significantly enriched terms from

each smoothing method found in the (c) up-regulated DEGs and (d) down-regulated DEGs

groups in *TM4SF1* over-expressed region for PDACA1 slide are shown, shaded by -

log10(p-value). (e) GO terms enriched in SPCS smoothed slides from up-regulated DEGs

that have previously been found in PDAC.

Reference

1. Heindl A, Sestak I, Naidoo K et al. Relevance of Spatial Heterogeneity of Immune Infiltration for Predicting Risk of Recurrence After Endocrine Therapy of ER+ Breast Cancer, J Natl Cancer Inst 2018;110.

2. Valkonen M, Ruusuvuori P, Kartasalo K et al. Analysis of spatial heterogeneity in normal epithelium and preneoplastic alterations in mouse prostate tumor models, Sci Rep 2017;7:44831.

3. Yeong J, Tan T, Chow ZL et al. Multiplex immunohistochemistry/immunofluorescence (mIHC/IF) for PD-L1 testing in triple-negative breast cancer: a translational assay compared with conventional IHC, J Clin Pathol 2020;73:557-562.

4. Gillett C, Fantl V, Smith R et al. Amplification and overexpression of cyclin D1 in breast cancer detected by immunohistochemical staining, Cancer Res 1994;54:1812-1817.

5. Gorsch SM, Memoli VA, Stukel TA et al. Immunohistochemical staining for transforming growth factor beta 1 associates with disease progression in human breast cancer, Cancer Res 1992;52:6949-6952.

6. Liang H, Li H, Xie Z et al. Quantitative multiplex immunofluorescence analysis identifies infiltrating PD1(+) CD8(+) and CD8(+) T cells as predictive of response to neoadjuvant chemotherapy in breast cancer, Thorac Cancer 2020;11:2941-2954.

7. Sun Z, Nyberg R, Wu Y et al. Developing an enhanced 7-color multiplex IHC protocol to dissect immune infiltration in human cancers, PLoS One 2021;16:e0247238.

8. Burgess DJ. Spatial transcriptomics coming of age, Nat Rev Genet 2019;20:317.

9. Marx V. Method of the Year: spatially resolved transcriptomics, Nat Methods 2021;18:9-14.

10. Moncada R, Barkley D, Wagner F et al. Integrating microarray-based spatial transcriptomics and single-cell RNA-seq reveals tissue architecture in pancreatic ductal adenocarcinomas, Nat Biotechnol 2020;38:333-342.

11. He B, Bergenståhle L, Stenbeck L et al. Integrating spatial gene expression and breast tumour morphology via deep learning, Nat Biomed Eng 2020;4:827-834.

12. Berglund E, Maaskola J, Schultz N et al. Spatial maps of prostate cancer transcriptomes reveal an unexplored landscape of heterogeneity, Nat Commun 2018;9:2419.

13. Chen WT, Lu A, Craessaerts K et al. Spatial Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease, Cell 2020;182:976-991.e919.

14. Maynard KR, Collado-Torres L, Weber LM et al. Transcriptome-scale spatial gene

expression in the human dorsolateral prefrontal cortex, Nat Neurosci 2021;24:425-436.

15.   Ortiz C, Navarro JF, Jurek A et al. Molecular atlas of the adult mouse brain, Sci
Adv 2020;6:eabb3446.

16.   Vickovic S, Eraslan G, Salmén F et al. High-definition spatial transcriptomics for in
situ tissue profiling, Nat Methods 2019;16:987-990.

17.   Berglund E, Saarenpää S, Jemt A et al. Automation of Spatial Transcriptomics
library preparation to enable rapid and robust insights into spatial organization of tissues,
BMC Genomics 2020;21:298.

18.   Dries R, Zhu Q, Dong R et al. Giotto: a toolbox for integrative analysis and
visualization of spatial expression data, Genome Biol 2021;22:78.

19.   Haque A, Engel J, Teichmann SA et al. A practical guide to single-cell RNA-
sequencing for biomedical research and clinical applications, Genome Med 2017;9:75.

20.   Linnarsson S, Teichmann SA. Single-cell genomics: coming of age, Genome Biol
2016;17:97.

21.   Ståhl PL, Salmén F, Vickovic S et al. Visualization and analysis of gene expression
in tissue sections by spatial transcriptomics, Science 2016;353:78-82.

22.   Ding J, Adiconis X, Simmons SK et al. Systematic comparison of single-cell and
single-nucleus RNA-sequencing methods, Nat Biotechnol 2020;38:737-746.

23.   Picelli S, Björklund Å K, Faridani OR et al. Smart-seq2 for sensitive full-length
transcriptome profiling in single cells, Nat Methods 2013;10:1096-1098.

24.   Hashimshony T, Senderovich N, Avital G et al. CEL-Seq2: sensitive highly-
multiplexed single-cell RNA-Seq, Genome Biol 2016;17:77.

25.   Huang M, Wang J, Torre E et al. SAVER: gene expression recovery for single-cell
RNA sequencing, Nat Methods 2018;15:539-542.

26.   van Dijk D, Sharma R, Nainys J et al. Recovering Gene Interactions from Single-
Cell Data Using Data Diffusion, Cell 2018;174:716-729.e727.

27.   Maynard KR, Collado-Torres L, Weber LM et al. Transcriptome-scale spatial gene
expression   in   the   human   dorsolateral   prefrontal   cortex,   Nature   Neuroscience
2021;24:425-436.

28.   Izar B, Tirosh I, Stover EH et al. A single-cell landscape of high-grade serous
ovarian cancer, Nature Medicine 2020;26:1271-1279.

29.   Zhao E, Stone MR, Ren X et al. Spatial transcriptomics at subspot resolution with
BayesSpace, Nature Biotechnology 2021;39:1375-1384.

30.   Jiang DX, Tang C, Zhang AD. Cluster analysis for gene expression data: A survey,
Ieee Transactions on Knowledge and Data Engineering 2004;16:1370-1386.

31.   Berthold MR, Höppner F. On clustering time series using euclidean distance and
pearson correlation, arXiv preprint arXiv:1601.02213 2016.

32.   Wold S, Esbensen K, Geladi P. Principal component analysis, Chemometrics and
intelligent laboratory systems 1987;2:37-52.

33.   Liu Y, Ye X, Zhan X et al. TPQCI: A topology potential-based method to quantify
functional influence of copy number variations, Methods 2021;192:46-56.

34.   Wang Z, Chen Z, Zhao Y et al. A community detection algorithm based on topology
potential and spectral clustering, The Scientific World Journal 2014;2014.

35.   Struyf A, Hubert M, Rousseeuw P. Clustering in an Object-Oriented Environment,
Journal of Statistical Software 1997;1:1 - 30.

36.   Waltman L, van Eck NJ. A smart local moving algorithm for large-scale modularity-
based community detection, The European Physical Journal B 2013;86:471.

37.   Scrucca L, Fop M, Murphy TB et al. mclust 5: Clustering, Classification and Density
Estimation Using Gaussian Finite Mixture Models, R Journal 2016;8:289-317.

38.   Hu J, Li X, Coleman K et al. SpaGCN: Integrating gene expression, spatial location
and histology to identify spatial domains and spatially variable genes by graph
convolutional network, Nature Methods 2021;18:1342-1351.

39.   Shahapure KR, Nicholas C. Cluster Quality Analysis Using Silhouette Score. In:
2020 IEEE 7th International Conference on Data Science and Advanced Analytics (DSAA).
2020, p. 747-748.

40.   Santos JM, Embrechts M. On the use of the adjusted rand index as a metric for
evaluating supervised classification. In: International conference on artificial neural
networks. 2009, p. 175-184. Springer.

41.   Fu F, Yang X, Zheng M et al. Role of Transmembrane 4 L Six Family 1 in the
Development and Progression of Cancer, Front Mol Biosci 2020;7:202.

42.   Liu Q, Guo L, Zhang S et al. PRSS1 mutation: a possible pathomechanism of
pancreatic carcinogenesis and pancreatic cancer, Mol Med 2019;25:44.

43.   Dal Molin A, Baruzzo G, Di Camillo B. Single-Cell RNA-Sequencing: Assessment
of Differential Expression Analysis Methods, Frontiers in Genetics 2017;8.

44.   Raudvere U, Kolberg L, Kuzmin I et al. g: Profiler: a web server for functional
enrichment analysis and conversions of gene lists (2019 update), Nucleic acids research
2019;47:W191-W198.

45.   Reimand J, Kull M, Peterson H et al. g:Profiler--a web-based toolset for functional
profiling of gene lists from large-scale experiments, Nucleic Acids Res 2007;35:W193-200.

46.   Consortium GO. The Gene Ontology resource: enriching a GOld mine, Nucleic
Acids Res 2021;49:D325-d334.

47. Jassal B, Matthews L, Viteri G et al. The reactome pathway knowledgebase,
Nucleic Acids Res 2020;48:D498-d503.

48. Kanehisa M, Furumichi M, Sato Y et al. KEGG: integrating viruses and cellular
organisms, Nucleic acids research 2020;49:D545-D551.

49. Martens M, Ammar A, Riutta A et al. WikiPathways: connecting communities,
Nucleic acids research 2020;49:D613-D621.

50. Wingender E. The TRANSFAC project as an example of framework technology
that supports the analysis of genomic regulation, Brief Bioinform 2008;9:326-332.

51. Giurgiu M, Reinhard J, Brauner B et al. CORUM: the comprehensive resource of
mammalian protein complexes-2019, Nucleic Acids Res 2019;47:D559-d563.

52. Colwill K, Gräslund S. A roadmap to generate renewable protein binders to the
human proteome, Nat Methods 2011;8:551-558.

53. Köhler S, Gargano M, Matentzoglu N et al. The Human Phenotype Ontology in
2021, Nucleic Acids Res 2021;49:D1207-d1217.

54. Zeng H, Shen Elaine H, Hohmann John G et al. Large-Scale Cellular-Resolution
Gene Profiling in Human Neocortex Reveals Species-Specific Molecular Signatures, Cell
2012;149:483-496.

55. Zhou J, Hui X, Mao Y et al. Identification of novel genes associated with a poor
prognosis in pancreatic ductal adenocarcinoma via a bioinformatics analysis, Biosci Rep
2019;39.

56. Atay S. Integrated transcriptome meta-analysis of pancreatic ductal
adenocarcinoma and matched adjacent pancreatic tissues, PeerJ 2020;8:e10141.

57. Liu L, Wang S, Cen C et al. Identification of differentially expressed genes in
pancreatic ductal adenocarcinoma and normal pancreatic tissues based on microarray
datasets, Mol Med Rep 2019;20:1901-1914.

58. Shang M, Zhang L, Chen X et al. Identification of hub genes and regulators
associated with pancreatic ductal adenocarcinoma based on integrated gene expression
profile analysis, Discov Med 2019;28:159-172.

59. Brenner M. Role of GFAP in CNS injuries, Neuroscience Letters 2014;565:7-13.

60. Weniger M, Honselmann KC, Liss AS. The Extracellular Matrix and Pancreatic
Cancer: A Complex Relationship, Cancers (Basel) 2018;10.

61. Yang JC, Zhang Y, He SJ et al. TM4SF1 Promotes Metastasis of Pancreatic
Cancer via Regulating the Expression of DDR1, Sci Rep 2017;7:45895.

62. Zheng B, Ohuchida K, Cui L et al. TM4SF1 as a prognostic marker of pancreatic
ductal adenocarcinoma is involved in migration and invasion of cancer cells, Int J Oncol
2015;47:490-498.

63. Ye J, Wen J, Ning Y et al. Higher notch expression implies poor survival in
pancreatic ductal adenocarcinoma: A systematic review and meta-analysis,
Pancreatology 2018;18:954-961.

64. Elazar N, Vainshtein A, Golan N et al. Axoglial Adhesion by Cadm4 Regulates
CNS Myelination, Neuron 2019;101:224-231.e225.

65. Alcalay RN, Levy OA, Wolf P et al. SCARB2 variants and glucocerebrosidase
activity in Parkinson's disease, npj Parkinson's Disease 2016;2:16004.

66. Giussani P, Prinetti A, Tringali C. The role of Sphingolipids in myelination and
myelin stability and their involvement in childhood and adult demyelinating disorders,
Journal of Neurochemistry 2021;156:403-414.