Trehalose-6-phosphate links singlet oxygen-induced signalling with metabolic signalling in *Chlamydomonas reinhardtii*

Waeil Al Youssef[1], Regina Feil[2], Maureen Saint-Sorny[3], Xenie Johnson[3], John E. Lunn[2], Bernhard Grimm[1], and Pawel Brzezowski[1*]

Affiliation:

[1]Humboldt-Universität zu Berlin, Institut für Biologie/Pflanzenphysiologie, 10115 Berlin, Germany

[2]Max Planck Institute of Molecular Plant Physiology, 14476 Potsdam-Golm, Germany

[3]Aix Marseille Université, CNRS, CEA, Institut de Biosciences et Biotechnologies Aix-Marseille, Photosynthesis and Environment team, CEA Cadarache, 13108 Saint-Paul-lez-Durance, France.

*e-mail: pawel.brzezowski@hu-berlin.de

Abstract

Singlet oxygen ($^1O_2$) induces retrograde signalling in chloroplasts. Using a novel mutant screen, we identified a mutation in the *TREHALOSE-6-PHOSPHATE PHOSPHATASE 1* (*T6PP1*) gene that results in accumulation of trehalose 6-phosphate, a reprogramming of cell metabolism, and impairment of $^1O_2$-induced retrograde signalling in *Chlamydomonas reinhardtii*. From transcriptomic analysis and metabolite profiling, we conclude that accumulation or deficiency of certain metabolites directly affect $^1O_2$-signalling. $^1O_2$-inducible *GLUTATHIONE PEROXIDASE 5* (*GPX5*) gene expression is suppressed by increased content of fumarate, an intermediate in the tricarboxylic acid cycle (TCA cycle) in mitochondria and dicarboxylate metabolism in the cytosol, while it is promoted by another TCA cycle intermediate, aconitate. Furthermore, genes encoding known essential components of chloroplast-to-nucleus $^1O_2$-signalling show decreased transcript levels in a *t6pp1* mutant, which can be rescued by exogenous application of aconitate. We demonstrate that chloroplast retrograde signalling involving $^1O_2$ depends on mitochondrial and cytosolic processes and that the metabolic status of the cell determines the response to $^1O_2$.

Introduction

Nearly forty years ago, tetrapyrrole biosynthesis (TBS) intermediates were proposed to be involved in chloroplast retrograde (organelle-to-nucleus) signalling. This conclusion was based on the observation that accumulation of chlorophyll precursors negatively affects transcript level of light-harvesting chlorophyll a/b-binding (LHCB) protein[1]. Later studies on chloroplast retrograde signalling involved mutants[2], treatment with a carotenoid biosynthesis inhibitor, norflurazon[3], or TBS inhibitors, such as α,α-dipyridyl [1] or thujaplicin[4]. Experiments with norflurazon led to the discovery of the *genomes uncoupled* (*gun*) mutants with a common phenotype of chloroplast status-independent expression of photosynthesis-associated nuclear genes[3].

It is noteworthy that treatment with inhibitors causing carotenoid deficiencies results in generation of reactive oxygen species (ROS) and photooxidative damage to chloroplasts. Furthermore, the end-products of TBS, haem and chlorophyll, as well as many of their intermediates (Supplementary Fig. 1) are light-absorbing and redox-reactive molecules, capable to generate ROS. Singlet oxygen ($^1O_2$) can be produced through interaction of ground (triplet)-state oxygen ($^3O_2$) with triplet-state chlorophyll or TBS intermediates, e.g. protoporphyrin IX (Proto), which are excited by light[5]. ROS are also metabolic products of other cellular processes in plants, such as photosynthesis and respiration.

Although $^1O_2$ is not considered to be the most reactive oxygen species, it is thought to be the major ROS involved in photooxidative damage[6]. However, it was shown that production of $^1O_2$ in the chloroplast induces stress responses that do not result exclusively from physicochemical damage, but also rely on signal transduction triggered by ROS[7]. Additionally, studies on the *fluorescent* (*flu*) mutant of *Arabidopsis thaliana*, which is impaired in regulation of TBS and accumulates protochlorophyllide (Supplementary Fig. 1), demonstrated that $^1O_2$-retrograde signalling is integrated into a broader signalling network and affects not only stress responses but also metabolism[8].

However, $^1O_2$ has a short half-life (about 200 ns) in the cell[9]. As a result, the distance that it may move was calculated to be approximately 10 nm, based on predicted diffusion rates[10,11]. Its diffusion range is also limited due to its high reactivity with membrane lipids. Therefore, $^1O_2$ could play a specific role as an activator of a stress response only if it is detected close to its source, which strongly suggests that other components mediating the $^1O_2$ signals should exist. Alternatively, altered metabolite contents triggered by $^1O_2$ may also mediate $^1O_2$-retrograde signalling, or as proposed by Pfannschmidt [12], a certain metabolic signature may be required to trigger changes in nuclear gene expression.

Despite the profound effect of $^1O_2$ on the chloroplast redox state and its apparent
involvement in altering nuclear gene expression, little is known about the components
involved in $^1O_2$-dependent retrograde signalling. The protein factors identified so far include
EXECUTER1 (EX1) and EX2 in the *flu* mutant of *A. thaliana*[13], PSBP DOMAIN-
CONTAINING PROTEIN 2 (PSBP2[14] and SINGLET OXYGEN ACCLIMATION (SAK1[15]
in *Chlamydomonas reinhardtii*, or METHYLENE BLUE SENSITIVITY (MBS1[16] shown to
be involved in $^1O_2$-signalling in both *A. thaliana* and *C. reinhardtii*.

In our search for tetrapyrrole-signalling components, we hypothesized that $^1O_2$
generated via the photosensitizing activity of Proto, rather than porphyrin accumulation itself,
triggers signalling cascades that alter nuclear gene expression. Therefore, we used a *C.*
reinhardtii mutant *chlD-1*[17] that accumulates Proto due to a dysfunctional Mg-chelatase
(MgCh, Supplementary Fig.1). The GENOMES UNCOUPLED 4 (GUN4) protein involved
in MgCh function (Supplementary Fig. 1) and signalling degrades upon Proto
accumulation[18]. However, *chlD-1* with introduced over-expression of GUN4, *chlD-1/GUN4*,
showed higher GUN4 content than *chlD-1*, but retained the chlorophyll-free phenotype[18].
Additionally, *chlD-1/GUN4* demonstrated higher expression of *GPX5* and higher tolerance to
$^1O_2$ than *chlD-1*[18]. Thus, to minimize the GUN4-deficient phenotype and to maintain high
and stable $^1O_2$-inducibility of *GPX5* expression, *chlD-1/GUN4* was used as a background
strain instead of *chlD-1* in the present study to monitor $^1O_2$-signalling. For this purpose,
expression of the *GLUTATHIONE PEROXIDASE 5* (*GPX5*) gene, which is specifically
induced by $^1O_2$ in *C. reinhardtii*[19,20] was used as a marker gene. The $^1O_2$-inducible promoter
region of *GPX5* fused to the promoterless *ARYLSULFATASE 2* gene (*ARS2*) was shown to be
an effective reporter to study *GPX5* expression[14,20,21].

Mutagenesis of the strain carrying the *GPX5-ARS2* reporter (in a *chlD-1/GUN4*
mutant background), followed by a screening for decreased reporter gene expression
identified a mutant impaired in $^1O_2$-signalling. The lack of the retrograde response to $^1O_2$ in
this mutant was verified to decrease expression of *GPX5*, as well as *SAK1*, *MBS*, and *PSBP2*.
The causal mutation was found in a gene annotated as *TREHALOSE 6-PHOSPHATE*
PHOSPHATASE (hereafter *T6PP1*). Besides clear impairment in $^1O_2$-signalling, mutation in
T6PP1 also resulted in accumulation of trehalose 6-phosphate (Tre6P). Therefore, T6PP1 is
the first confirmed phosphatase acting on Tre6P in *C. reinhardtii* and in fact the first enzyme
with a confirmed function in trehalose metabolism in this organism. Our data indicate that
accumulation of Tre6P causes changes to the expression of genes involved in several
metabolic pathways. However, Tre6P is an intermediate rather than the primary cause of the

impaired $^1O_2$-signalling. Here using a combination of genetics, gene expression and

metabolic phenotypes, we reveal a complex interaction between chloroplast, mitochondria

and cytosol in $^1O_2$ signal transmission to the nucleus.

Results

Forward genetics screen to identify components of the $^1O_2$-signalling

To generate a mutant impaired in $^1O_2$-dependent signalling, we first created a reporter strain

in a known $^1O_2$-generating mutant, which was subsequently subjected to mutagenesis. The

GPX5-ARS2 reporter construct (Supplementary Fig. 2) was introduced into the genome of

chlD-1/GUN4 to express the ARS2 protein in a $^1O_2$-dependent manner, and the transformants

were tested for the enzymatic activity of ARS2. The reporter strain, which showed the lowest

ARS2 activity in darkness and the highest activity in light was named *signalling Reporter*

(*sigRep*; Fig. 1a) and was used in further applications. Expression kinetics of the cytosolic

(*GPX5$_{cyt}$*) and chloroplastic (*GPX5$_{cp}$*) version of *GPX5*[22]; Supplementary Fig. 2), as well as

the *GPX5-ARS2* construct were examined by quantitative Real-Time PCR (qRT-PCR) upon

transfer from dark to light. As expected, higher induction of *GPX5$_{cp}$* and *GPX5$_{cyt}$* were

observed upon light illumination in *sigRep* compared to WT (Fig. 1b). Induced *GPX5-ARS2*

expression in *sigRep* in the light (Fig. 1c) confirmed the results obtained in the ARS activity

assay (Fig. 1a).

Subsequently, a random mutagenesis was performed on *sigRep* using bleomycin

resistance cassette (ble^R) as an insert, followed by screening to isolate strains with lower or

undetectable ARS2 activity in the light compared to *sigRep*, and thus, possibly disrupted $^1O_2$-

dependent signalling. Screen of 804 transformants allowed us to isolate nine, which were

named *genomes uncoupled Singlet Oxygen Signalling* (*gunSOS*). The *gunSOS1* mutant (Fig.

1d) was selected for further analysis. Higher sensitivity to light was recorded in *gunSOS1*

compared to the *sigRep* background strain (Fig. 1e), in agreement with a higher sensitivity or

impaired acclimation to $^1O_2$. Mutant impaired in $^1O_2$-signalling showed similar Proto

accumulation in light compared to *sigRep* and *chlD-1/GUN4* (Supplementary Fig. 3), which

eliminated the possibility that spurious mutations in *gunSOS1* resulted in reduced content of

this photosensitizer.

The ble^R insertion in *gunSOS1* was located in the first exon of the gene annotated as

TREHALOSE 6-PHOSPHATE PHOSPHATASE 1 (here *T6PP1*; locus Cre12.g497750) in the

JGI portal (Department of Energy Joint Genome Institute, https://phytozome-

next.jgi.doe.gov, v13, genome v5.6; Fig. 1f). The *T6PP1* transcript abundance was

determined in *gunSOS1* and compared to *sigRep* by qRT-PCR using primers annealing to the

coding sequence upstream of the insertion site. At 2 h after transfer from dark to light, *T6PP1*

mRNA content increased 2.5-fold in *sigRep* compared to WT, while it was virtually absent in

gunSOS1 (Fig. 1g). Expression of $GPX5_{cyt}$ and $GPX5_{cp}$ was nearly 10 and 8 times lower in

gunSOS1 compared to *sigRep*, respectively (Fig. 1h). The transcript abundance for *GPX5-*

ARS2 was 7 times lower in *gunSOS1* than in *sigRep* (Fig. 1i), and explains undetectable

ARS2 activity in the $^1O_2$-signalling mutant (Fig. 1d).

C. reinhardtii T6PP1 is a functional Tre6P phosphatase

While *gunSOS1* is clearly impaired in $^1O_2$-signalling, it was necessary to determine the

primary phenotype caused by the mutation in *T6PP1*. The content of Tre6P in *sigRep* and

gunSOS1 was determined by anion-exchange high performance liquid chromatography

coupled to tandem mass spectrometry (LC-MS/MS). Tre6P accumulation increased in

gunSOS1 upon exposure to light in a time-dependent manner, while it decreased in *sigRep*

(Fig. 2a). Accumulation of Tre6P in *gunSOS1* indicates that *C. reinhardtii* T6PP1 is a

functional phosphatase dephosphorylating Tre6P to trehalose. Subsequently, *T6PP1*

expression was found to be induced in the light in *sigRep*, but not in the WT, indicating its

inducibility by photooxidative stress rather than light (Fig. 2b). Mutation of the *T6PP1* gene

would be expected to lower Tre6P phosphatase activity and dephosphorylation of Tre6P to

trehalose, but we also observed accumulation of trehalose in *gunSOS1* (Fig. 2c). This

apparent discrepancy can be explained by the existence of another enzyme in *C. reinhardtii*

with possible Tre6P phosphatase activity, in combination with higher levels of Tre6P (see

Discussion section).

Rescue of the T6PP1 deficiency in *gunSOS1* was performed with the isolated genomic

DNA fragment carrying the wild-type *T6PP1* gene. Several independent transformants

showed rescued $^1O_2$-signalling, which was indicated by the *GPX5-ARS2* expression

determined in the ARS assay, and data from three representative strains are shown in

Supplementary Fig. 4a. The analysed rescued strains also showed increased $GPX5_{cp}$ transcript

content compared to *gunSOS1* (Supplementary Fig. 4b). A peptide-specific antibody for

T6PP1 was produced to compare the protein content in *gunSOS1* to strains showing $^1O_2$-

dependent signalling. A faint immune signal of approximately 43 kDa was detected in

gunSOS1 and *chlD-1*, but two narrow bands were seen in *chlD-1/GUN4*, *sigRep* and the

T6PP1-rescued strains (Fig. 2d). WT and *gunSOS1* showed an additional unspecific

immunoreaction with protein of approximately 53 kDa. As expected, GPX5 content was

lower in *gunSOS1* compared to *sigRep* (Fig. 2d), which correlated with qRT-PCR results on *GPX5* transcript levels (Fig. 1 h). Strains with rescued $^1O_2$-signalling, *T6PP1-R1, -R2, and -R3*, showed increased content of GPX5 (Fig. 2d) and higher tolerance to light compared to *gunSOS1* (Fig. 2e).

The *gunSOS1* mutant shows altered metabolism compared to *sigRep*

In *A. thaliana* and other angiosperms, Tre6P functions as a sucrose signal and homeostatic regulator of sucrose metabolism[23,24], which links plant growth and development to the availability of sucrose[25], reviewed in[26]. To determine the effect of accumulated Tre6P on metabolism in *C. reinhardtii*, we performed comparative metabolite profiling of *gunSOS1* and *sigRep* (Supplementary Table 1), which revealed a relatively wide range of variation. Aside from Tre6P and trehalose, 11 out of 27 analysed metabolites showed significantly increased content in *gunSOS1* compared to *sigRep* in the light (Figure 3a). By means of the MetaboAnalyst 5.0 portal (https://www.metaboanalyst.ca) these metabolites were assigned to the metabolic processes in the cell. A hypergeometric test indicated the most affected metabolic pathways, i.e. the TCA cycle, pyruvate metabolism, and starch and sucrose metabolism (Fig. 3b, Supplementary Table 2). Eight metabolites showed significantly decreased content in *gunSOS1* compared to *sigRep* (Fig. 4a), with the highest impact (MetaboAnalyst 5.0) on fructose and mannose metabolism, amino sugar and nucleotide sugar metabolism, and starch and sucrose metabolism (Fig. 4b, Supplementary Table 3). Metabolites, which did not show a significant change in *gunSOS1* compared to *sigRep* are presented in Supplementary Fig. 5.

Starch content was determined in *gunSOS1* and compared to *sigRep*, because of changes in several intermediates of starch and sucrose metabolism (Fig. 3b and Fig. 4b). Indeed, *gunSOS1* accumulated only 60% of the starch content of *sigRep* when the strains were grown in acetate-supplemented media (TAP, Fig. 4c). Both *sigRep* and *gunSOS1* are obligate heterotrophs. *SigRep* contained less starch after transfer from TAP and cultivation in TP for 48 h, whereas the starch content of *gunSOS1* was essentially unchanged (Fig. 4c), indicating that there was net degradation of starch in *sigRep* but not in *gunSOS1*.Starch formation is strongly induced during nitrogen (N) deprivation[27]. The *sigRep* cells did not increase starch reserves after 3 days in N-deficient medium, suggesting that starch accumulation had already reached its maximum under non-stress conditions. The starch content in *gunSOS1* deprived of N did increase by 50% and was similar to the content in *sigRep* (Fig. 4c). We conclude that *sigRep* accumulates high levels of starch as a

consequence of the *chlD-1/GUN4* genetic background and that the accumulation of Tre6P

represses this response in *gunSOS1* leading to a lower accumulation of starch. The *gunSOS1*

cells are still capable of responding to the N-deprivation stress and accumulating starch

despite disturbed metabolism. Starch degradation on the other hand, is compromised in

gunSOS1, which could indicate a general impairment of catabolic processes or a defect in

sensing carbon limitation. The map-overview of the selected metabolic pathways with

depicted metabolites that showed different content, or were not changed, in *gunSOS1* relative

to *sigRep* is presented in Figure 5.

Increased Tre6P content alters metabolism in *C. reinhardtii* via alteration of the

transcriptome

To determine the extent to which chloroplast retrograde signalling is affected in *gunSOS1*, we

performed comparative RNA-sequencing (RNA-seq) of the transcriptomes of *gunSOS1* and

sigRep upon transfer from dark to light. The analysis showed that 8120 genes were expressed

in both strains, but 1377 and 1087 additional genes were uniquely expressed in *gunSOS1* and

sigRep, respectively (Fig. 6a, detailed listing can be found in Supplementary Data Set 1).

Furthermore, differential gene expression analysis showed that the transcript contents of 1606

genes and 1706 genes were lower or higher, respectively, in *gunSOS1* compared to *sigRep*

(Supplementary Data Set 2).

Based on the KEGG ontology (http://www.kegg.jp/) several genes associated with

metabolic pathways were down- (Fig. 6b) or up-regulated (Fig. 6c) in *gunSOS1* compared to

sigRep (Supplementary Data Set 3). Given the strong overlap between differentially

expressed genes and corresponding pathway intermediates, we hypothesized that the altered

metabolism can be directly explained by altered gene expression. To decipher a complex

correlation between altered signalling, gene expression and metabolism, accumulation or

deficiency in metabolites from given pathway(s) was correlated with altered gene expression

in *gunSOS1* compared to *sigRep*. Fumarate and aconitate will be described in more detail, as

representative examples of the metabolites showing respectively increased or decreased

content in *gunSOS1* compared to *sigRep*.

KEGG enrichment of the RNA-seq data indicated increased expression of the gene

encoding FUMARATE HYDRATASE CLASS II (FUM2, Enzyme Commission (EC)

Number 4.2.1.2, Cre01.g020223) in *gunSOS1* compared to *sigRep*. FUM is responsible for

reversible stereospecific interconversion of malate to fumarate. However, specific isoforms of

this protein act in defined pathways and may favour one direction over the other depending

on the subcellular localisation and environment[28]. The mitochondrial isoform catalyses

hydration of fumarate to L-malate in the TCA cycle, while the cytosolic form catalyses

dehydration of L-malate to fumarate[29,30]. Two isoforms exist in *C. reinhardtii*, FUM1

encoded in locus Cre06.g254400 and FUM2. Based on the predicted subcellular localisation

(PredAlgo; http://lobosphaera.ibpc.fr/cgi-bin/predalgodb2.perl?page=main), FUM1 localizes

to the mitochondria, while FUM2 is assigned to the "other" compartment and is likely to be a

cytosolic protein. Pathway (KEGG) diagrams for the TCA cycle and pyruvate metabolism

can be found in Supplementary Fig. 6, showing integration of metabolite, RNA-seq and qRT-

PCR data. Comparison of the transcriptomes of *gunSOS1* and *sigRep* also indicated increased

expression of the *FUMARYLACETOACETASE* gene (locus *Cre17.g732802*, EC 3.7.1.2)

involved in tyrosine catabolism. FUMARYLACETOACETASE catalyses hydrolysis of 4-

fumarylacetoacetate to acetoacetate and fumarate (Supplementary Fig. 7). Increased

expression of genes encoding FUM2 and FUMARYLACETOACETASE could explain

accumulation of fumarate in *gunSOS1* compared to *sigRep*.

Aconitate deficiency in *gunSOS1* compared to *sigRep* can be explained by

transcriptional downregulation of *ACONITATE HYDRATASE* (*ACH1*, Cre01.g042750,

EC 4.2.1.3). ACH1 interconverts citrate and isocitrate, via cis-aconitate, in the TCA cycle

(Supplementary Fig. 8) and is also involved in glyoxylate and dicarboxylate metabolism. At

equilibrium, the reactants of the ACH1 reaction are present in the following ratio: 91%

citrate, 6% isocitrate and 3% aconitate. With the smallest pool of the three tricarboxylic

acids, fluctuations in the level of aconitate are expected to be more pronounced compared to

the other metabolites involved, especially citrate, whose content was similar in *gunSOS1* and

sigRep in the light (Supplementary Fig. 5). Taken together, our data indicate that altered

fumarate and aconitate content in *gunSOS1* compared to *sigRep* can be explained by altered

expression of genes encoding enzymes involved in key metabolic processes affecting those

metabolites.

266 Impaired $^1O_2$-signalling in *gunSOS1* correlates with decreased expression of *PSBP2*,

267 *MBS*, and *SAK1*

268 The PSBP2[14], MBS[16], and SAK1[15] proteins are required for $^1O_2$-induced chloroplast

269 retrograde signalling in *C. reinhardtii*. Therefore, we determined the transcript levels of

270 *PSBP2*, *MBS*, and *SAK1*, which were 8-, 9-, and nearly 41-fold lower, respectively, in

271 *gunSOS1* compared to *sigRep* (Fig. 7a). Decreased expression of *SAK1* in *gunSOS1* prompted

272 us to compare transcript levels of selected $^1O_2$-responsive genes between *gunSOS1* and *sak1*,

which lacks acclimation response to $^1O_2$[15]. SAK1 is a key regulator of the gene expression response and its involvement in $^1O_2$-signalling was determined following treatment with rose bengal[15], which produces $^1O_2$ in the light. Nevertheless, based on our qRT-PCR analysis, all tested genes had the same reduced inducibility in *gunSOS1* compared to *sigRep* (Fig. 7b), as it was observed in *sak1* compared to WT following $^1O_2$-exposure[15]. Among the genes that were down-regulated in both *gunSOS1* (Fig. 7b) and *sak1*[15] were two *CYCLOPROPANE FATTY ACID SYNTHASES*, *CFA1* and *CFA2* (*CPLD27*), involved in lipid and sterol metabolism, as well as the gene encoding SOUL1 heme-binding protein (Fig. 7b). It is noteworthy that attenuated expression of both *CFA1* and *CFA2*, as well as *SOUL1*, was also observed in studies on a *gpx5* mutant[31]. Furthermore, in agreement with the *sak1* phenotype presented in Wakao, et al. [15], *gunSOS1* also showed decreases in transcript content of *PEPTIDE METHIONINE SULFOXIDE REDUCTASE* (*MSRA3*), *ALCOHOL DEHYDROGENASE* (*ADH7*), *RETINALDEHYDE BINDING PROTEIN-RELATED* (*RABPR2*), and Cre01.g007300 encoding N-ACETYLPHOSPHATIDYLETHANOLAMINE-HYDROLYZING PHOSPHOLIPASE D (Fig. 7b). Thus, there is an overlap between the phenotypes of *gunSOS1* and *sak1*[15] with respect to their attenuated expression of genes induced during elevated $^1O_2$ (Fig. 7b).

Altered metabolite content affects $^1O_2$-signalling

We hypothesized that at least some of the metabolites showing increased or decreased content in *gunSOS1* compared to *sigRep*, attenuate or propagate $^1O_2$-signalling, respectively. We selected fumarate and aconitate as representatives of either group. To test our hypothesis, metabolites were added separately to the liquid cultures at empirically determined sublethal concentrations, followed by incubation in the light to induce $^1O_2$ generation by Proto. Subsequently, the $GPX5_{cyt}$ and $GPX5_{cp}$ transcript levels were used as readouts for the $^1O_2$ signalling efficiency in *gunSOS1* and *sigRep*.

Exogenous application of fumarate significantly ($P < 0.001$, a detailed report from two-way ANOVA analyses is presented in Supplementary Tables 4 and 5) attenuated expression of both $GPX5_{cyt}$ and $GPX5_{cp}$ in a concentration-dependent manner in *sigRep* (Fig. 8a). As expected, fumarate did not affect *GPX5* expression in *gunSOS1*. Application of 20 μM aconitate significantly ($P < 0.0001$) increased expression of *GPX5* in *gunSOS1* and even further in *sigRep* compared to untreated cells ($P < 0.0001$), but 100 μM had a quenching effect on the *GPX5* expression in *sigRep* (Fig. 8b, Supplementary Tables 6 and 7). This indicates that aconitate up to a certain threshold concentration promotes $^1O_2$ signalling.

Subsequently, we determined the expression of *SAK1* upon feeding *gunSOS1* and
sigRep with aconitate. Exogenous application of aconitate at 20 μM increased *SAK1*
transcript abundance in *gunSOS1* more than 120 times compared to the untreated *gunSOS1*
control, and exceeded values observed in *sigRep* subjected to the same treatment by a factor
of 5 (Fig. 8c). Due to the increased *SAK1* expression in *gunSOS1* upon feeding with
aconitate, we also determined the expression of *PSBP2* and *MBS* in the same conditions. The
PSBP2 response to aconitate was similar to *SAK1* in terms of the increased expression (Fig.
8c), but with less pronounced dependence on the metabolite concentration compared to
SAK1. Although 20 μM aconitate increased *PSBP2* expression in *gunSOS1* by a factor of 48
compared to the untreated control, it was still 2.7 times lower compared to *sigRep* subjected
to the same treatment (Fig. 8c). Similarly to *SAK1* and *PSBP2*, 20 μM aconitate also
increased expression of *MBS* in *gunSOS1*, although values did not exceed those observed in
sigRep subjected to the same treatment (Fig. 8c). Statistical analyses indicated that the effect
of 20 μM aconitate on *SAK1*, *PSBP2* and *MBS* expression in *sigRep* and *gunSOS1*,
interaction between mutants and aconitate, as well as the mutant-dependent expression of
these genes were always significant ($P < 0.001$, Supplementary Tables 8-10).
323

Specificity of the $^1O_2$-signalling pathway(s) attenuated in *gunSOS1*

The profound effect of accumulating Tre6P on the transcriptome (Fig. 6a-c) and consequently
the metabolism of *gunSOS1* (Fig. 3 and Fig. 4) may have been indicative of a general
photooxidative stress response being impaired in *gunSOS1*. To determine the specificity of
this response, we compared *gunSOS1* and *sigRep* in terms of the expression of selected genes
that are known to be induced by various reactive species or conditions causing
photooxidative stress other than $^1O_2$.

Upon H_2O_2 or organic *tert*-butyl hydroperoxide (*t*-BOOH) treatment, Blaby, et al. [32]
observed an increase in the expression of the *MSD3* gene, encoding plastid-localised Mn
SUPEROXIDE DISMUTASE 3. In our studies, we did not observe an increase in *MSD3*
transcript in *sigRep* compared to WT (Fig. 9a), which shows that *MSD3* expression is not
inducible by $^1O_2$ produced by Proto. However, based on qRT-PCR analysis, a 9-fold increase
was observed in *gunSOS1* compared to *sigRep* (Fig. 9a, see also RNA-seq in Supplementary
Data Set 2). Upon H_2O_2 or *t*-BOOH treatment, Urzica, et al. [33] also observed induced
expression of genes involved in the glutathione-ascorbate system, *GDP-L-GALACTOSE*
PHOSPHORYLASE (*VTC2*) and *DEHYDROSASCORBATE REDUCTASE* (*DHAR1*). Based
on our qRT-PCR results, *VTC2* transcript content was not changed in *sigRep* compared to

WT, but an increase was observed in *gunSOS1* (Fig. 9b; Supplementary Data Set 2). *DHAR1* expression was also stimulated in *gunSOS1* compared to *sigRep* (Fig. 9c; Supplementary Data Set 2).

Similarly to *sak1* following treatment with rose bengal[15], we observed increased expression of *GLUTATHIONE S-TRANSFERASE* (*GSTS1*) in *gunSOS1* compared to *sigRep* (Fig. 9d, Supplementary Data Set 2). However, in another study increased expression of *GSTS1* was shown after treatment with acrolein, which suggests its transcriptional induction by reactive electrophile species (RES)[34]. Acrolein was also shown to induce *FSD1* encoding Fe SUPEROXIDE DISMUTASE (FeSOD[34], which was expressed both in *sigRep* and *gunSOS1* (Supplementary Data Set 1), and no significant difference could be observed in DEG analysis (Supplementary Data Set 2) or qRT-PCR (Fig. 9e). It can be concluded that, despite impaired $^1O_2$-signalling, *gunSOS1* retained the ability to respond to photooxidative stress caused by other ROS such as H_2O_2, organic peroxides, or RES.

Discussion

ROS are formed as a by-product of biological redox reactions[35] mostly in the mitochondria or chloroplasts[36,37]. Although excess ROS production can cause oxidative damage to cell components, ROS or the oxidation products play an important role in the signal transduction processes.

Different retrograde signalling pathways have been proposed to involve also TBS-intermediates in plants and green algae (reviewed in[38,39]). While involvement of Mg-porphyrins in chloroplast retrograde signalling is now excluded[40,41], in the present study we have demonstrated that $^1O_2$ produced by the photosensitizing activity of Proto in the light, but not Proto itself, triggers signalling cascades that alter nuclear gene expression. Using strains that endogenously accumulate Proto, we identified several mutants that are impaired in $^1O_2$-signalling. Endogenous accumulation of Proto in mutants is advantageous for studying $^1O_2$-signalling, because it eliminates the need for exogenous application of TBS or carotenoid biosynthesis inhibitors, as well as photosensitizers, such as rose bengal or neutral red (Fischer et al., 2004) to induce $^1O_2$ generation. The lack of chlorophyll provides another advantage, because such mutants do not have functional photosynthetic electron transport, so that $^1O_2$ production in photosynthesis is avoided. The accumulating Proto is thus the dominant source of generated $^1O_2$.

The $^1O_2$-induced signalling phenotype in the *gunSOS1* mutant was due to a lesion in the *T6PP1* gene. We demonstrated that T6PP1 is a functional phosphatase capable of

dephosphorylating Tre6P. *T6PP1* is the only representative of the classical plant *TPP* gene family in *C. reinhardtii*, contrasting with the large *TPP* gene families in angiosperms[42]. In addition, *C. reinhardtii* has one representative of the class I *TREHALOSE-6-PHOSPHATE SYNTHASE* (*TPS*) family (Cre16.g662350, hereafter *T6PS1*), and two members of the class II *TPS* family (Cre06.g278221, here *T6PSP1* and Cre16.g686200, here *T6PSP2*). Both class I and class II TPS proteins have glucosyltransferase and TPP-like domains, but only the class I TPS proteins have demonstrated TPS activity[43-45]. T6PP1 belongs to the haloacid dehalogenase superfamily of proteins and contains the characteristic DXDX(T/V) active site motif – $_{107}$DYDGT$_{112}$ – in which the initial Asp residue forms a phospho-acyl intermediate during catalysis[46]. T6PSP1 also contains the complete active site motif ($_{592}$DYDGT$_{597}$), so we cannot exclude the possibility that the T6PSP1 protein in *C. reinhardtii* has TPP activity, which could explain the increased content of trehalose in *gunSOS1* compared to *sigRep*. Although the intracellular localizations of these proteins have not been determined experimentally in *C. reinhardtii*, PredAlgo analysis indicated possible localization of T6PP1 and T6PSP1 in mitochondria, while T6PS1 and T6PSP2 are likely to be cytosolic proteins, because they were not assigned to any specific organelle.

Correlation between *T6PP1* (Fig. 2b) and *GPX5* (Fig. 1b, c) expression indicates that low Tre6P content is required for efficient $^1O_2$-signalling. However, the apparent negative effect of Tre6P on $^1O_2$-signalling is not direct and involves a complex metabolic reprogramming (Fig. 10). Study on the contrasting phenotypes between *A. thaliana* overexpressing bacterial TPS, TPP, or TREHALOSE PHOSPHATE HYDROLASE (TPH) pointed to Tre6P, rather than trehalose, playing a signalling function[25]. Tre6P was shown to be highly correlated with sucrose, leading to the proposal that it functions as a signal of sucrose status[23]. Tre6P was also shown to inhibit starch degradation in *A. thaliana*[47,48], which is also true in Tre6P-acumulating *gunSOS1* mutant of *C. reinhardtii*.

It was demonstrated that Tre6P acts as an inhibitor of SUCROSE-NON-FERMENTING (SNF)-RELATED PROTEIN KINASE1 (SnRK1) in developing tissues and that this is dependent on a so-far unidentified protein factor[49-51]. Tre6P also inhibits the activation of SnRK1 by SnRK1-activating kinases/geminivirus Rep interacting kinases[52]. SnRK1 belongs to the AMPK-SNF1-SnRK family of protein kinases, which is represented in all eukaryotes[53]. In plants, SnRK1 plays a central role in energy and metabolic homeostasis, and is activated during energy deficient conditions caused by stresses like nutrient starvation, pathogen attack, or ROS[54]. Baena-González, et al. [55] established approximately 1000 genes as markers of SnRK1 in *A. thaliana*, which indicates an extensive SnRK1-dependent

transcriptional reprogramming. In *C. reinhardtii*, involvement of various SnRKs in responses

to stress was observed during sulphur[56,57] and nitrogen deprivation[58], or cold stress[59].

Genome-wide analysis revealed the existence of 21 genes as potential orthologues of the

plant SnRK α, β and γ/βγ subunits in *C. reinhardtii*. It was suggested that the proteins

encoded by these genes play the same role in cell survival and stress response in *C.*

reinhardtii as SnRKs in land plants[58].

In analogy to *A. thaliana*, altered SnRKs activity was shown to cause metabolic

remodelling also in algae[59]. In our study, the majority of changes to gene expression observed

in *gunSOS1*, relative to *sigRep*, originates from accumulation of Tre6P and, based on

previous studies, this leads to inhibition of one or more SnRKs (Fig. 10). Our analysis

showed that altered gene expression in *gunSOS1* relative to *sigRep* (Supplementary Fig. 6-8)

is directly responsible for altered fumarate and aconitate content in *gunSOS1* compared to

sigRep. Thus, Tre6P accumulating in *gunSOS1* is not directly involved in $^1O_2$-signalling but

controls other processes in the cell which affect $^1O_2$-signalling more directly (Fig. 10).

We demonstrated that accumulation of fumarate is responsible for attenuating $^1O_2$-

signalling (Fig. 8a, Fig. 10). Fumarate is a well-recognized oncometabolite in mammalian

cells[60], associated with development of tumours by competitive inhibition of 2-oxoglutarate-

dependent oxygenases, including HYPOXIA INDUCIBLE FACTOR (HIF) hydroxylases,

leading to stabilisation of HIF and activation of oncogenic HIF-dependent pathways[61]. In

fact, accumulation of fumarate in human cells was linked to an aggressive variant of

hereditary kidney cancer[62]. In mice, fumarate was also shown to directly modify some

proteins by succination of cysteine residues to form 2-succinocysteine derivatives[63].

Succination of three cysteines crucial for iron-sulphur cluster binding was identified in

mitochondrial ACONITASE2 (ACO2) in a *fumarate hydratase 1* knockout (Fh1KO) mouse

embryonic fibroblast (MEF) cell line[63]. Analysis of tryptic peptides derived from ACO2 in

Fh1KO MEFs, indicated succination of Cys385. Another tryptic peptide of ACO2 was

identified as a mixture of two isomers in Fh1KO MEFs, which showed succination at Cys451

or Cys448[63]. Succination of these cysteine residues in ACO2 leads to its inhibition[63].

Equivalent cysteine residues are conserved in *C. reinhardtii* ACH1, which explains the

aconitate deficiency in *gunSOS1* (Fig. 4a).

The *gunSOS1* mutant showed decreased levels of aconitate compared to *sigRep*, while

exogenous application of this metabolite to *gunSOS1* restored $^1O_2$-signalling. Mammalian

ACO2 is thought to be a mitochondrial oxidative stress sensor, which requires an active

$[Fe_4S_4]^{2+}$ cluster[64]. It is, therefore, proposed that accumulation of fumarate deactivates ACH1

and reduces the aconitate content in *gunSOS1*, thereby impairing oxidative stress sensing and

$^1O_2$-signalling. Diminished expression of *ACH1* in *gunSOS1* relative to *sigRep*

(Supplementary Fig. 8a, b) may be an additional factor adding to the aconitate depletion in

gunSOS1.

Results obtained in our study indicate that there is a metabolic signature conditioning

$^1O_2$-signalling. A mechanism in which the chloroplast controls nuclear gene expression via

metabolite levels was postulated before[12,65]. It was shown that altered chloroplast redox

conditions result in changes in the metabolome in *A. thaliana*, reflected also by the

reallocation of energy resources[66]. Importantly, although our study shows that the metabolic

configuration of the cell is essential for $^1O_2$-signalling, this does not exclude the involvement

of protein components in this signalling network (Fig. 10). On the contrary, our findings

support the direct involvement of proteins in $^1O_2$-signalling, but we demonstrated that

expression of *PSBP2*[14], *MBS*[16], and *SAK1*[15] depends on the metabolic status (Fig. 7a and Fig.

8a, b). The overlap between the *gunSOS1* and the SAK1-dependent transcriptomes (Fig. 7b)

as well as other defined ROS or RES-dependent signalling pathways being functional (Fig.

9a-e) indicate specificity of the impaired $^1O_2$-signalling in *gunSOS1*. Our results indicate that

fumarate and aconitate affect the expression of *PSBP*, *MBS*, or *SAK1*, which subsequently

convey information about the metabolic status of the cell to the nucleus and trigger specific

responses (Fig. 10). It is not clear yet how the specificity of the signalling is achieved,

because the mechanisms by which metabolites affect the expression of these proteins, as well

as downstream components of these signalling pathways, remain unknown. Additionally, the

protein components involved in $^1O_2$-signalling may be subject to additional regulatory

mechanisms, such as phosphorylation, as proposed for SAK1 by Wakao, et al. [15].

We have shown that retrograde signalling triggered by $^1O_2$, which is mostly generated

in the chloroplast, strictly depends on the mitochondrial metabolic status. PROLYL-tRNA

SYNTHETASE (PRORS1) in *A. thaliana*, a component of the organellar gene expression

machinery indicated that mitochondria may contribute to chloroplast retrograde signalling[67].

PRORS1 is targeted both to the plastid and to mitochondria, but downregulation of specific

photosynthesis-associated nuclear genes was observed only in a plastidial *prpl11* and

mitochondrial *mrpl11* double mutant, but not in *prpl11* or *mrpl11* single mutants[67]. Another

study showed that the ROS-dependent signals from chloroplast and mitochondria are

integrated at the RADICAL-INDUCED CELL DEATH 1 (RCD1) protein located in the

nucleus[68]. RCD1 suppresses transcription factors ANAC013 and ANAC017, which mediate

the ROS-signal from mitochondria, while chloroplast regulation of RCD1 takes place through

3'-phosphoadenosine 5'-phosphate. It was proposed that RCD1 may function at the
intersection of mitochondrial and chloroplastic retrograde signalling pathways[68] .

The retrograde signalling pathways identified so far assume the existence of separate
signalling routes. However, chloroplast retrograde signalling involves a network of different
but inter-connected mechanisms, and observation of specific signalling pathways within the
network depends on the experimental conditions. Pharmacological or genetic interventions
that disrupt specific pathways can be used to characterize alternative pathways. In the
manuscript presented here, we report that the metabolic status of the cell, reflecting levels of
Tre6P, and components of the mitochondrial TCA cycle are integral factors in the $^1O_2$-
dependent retrograde signalling network in *C. reinhardtii*.

Methods

***Chlamydomonas reinhardtii* cultures and genetic manipulations**

All strains were cultivated heterotrophically in Tris-acetate-phosphate (TAP) medium in the
dark. Additionally, starch measurements were performed in TAP, Tris-phosphate (TP)
medium without acetate, or nitrogen-depleted (TAP-N) medium. Experiments were
performed in TAP medium in the dark or upon a shift to 20 μmol photons m^{-2} s^{-1} light. Three
strains used in this study were described elsewhere, *chlD-1* in von Gromoff, et al. [17], *chlD-*
1/GUN4 in Brzezowski, et al. [18], and wild type (4A+) in Dent, et al. [69].

To generate the *GPX5-ARS2* construct (Supplementary Fig. 2), the *GPX5* regulatory
region (*GPX5* 5' RR) was amplified by PCR using forward and reverse primers carrying
XhoI and EcoRV restrictions sites, respectively (Supplementary Table 11**)**. The obtained
fragment was subcloned and ligated between the XhoI/EcoRV sites of the pSL18 vector[70],
replacing the existing promoter region of the gene encoding the PSAD and positioning *GPX5*
5'RR in reverse orientation with respect to the paromomycin resistance cassette (paro[R,71]. The
DNA fragment carrying paro[R] and *GPX5* 5' RR was excised by restriction digestion with
KpnI and EcoRV and ligated into the corresponding restriction sites of pJD54[72], which
carries a copy of the promoterless version of the gene encoding ARYLSULFATASE 2
(ARS2), which allowed *ARS2* expression to be controlled by the *GPX5* 5'RR (Supplementary
Fig. 2). The *GPX5-ARS2* construct was verified by sequencing.

The *sigRep* strain was produced in the *chlD-1/GUN4* background by transformation
with a *GPX5-ARS2* construct. Transformant selection was performed on TAP agar plates with
10 μg mL^{-1} paromomycin in dark, followed by a screen for *GPX5-ARS2* inducibility by $^1O_2$.
The *sigRep* strain showed low ARS2 activity in the dark and high activity in the light (Fig.

1a) and was selected for further applications. The *gunSOS1* mutant was generated by random insertional mutagenesis performed on *sigRep* using a bleomycin resistance cassette (ble^R) isolated from the pMS188 vector[73] using NotI and KpnI. Following mutagenesis, selection was performed on TAP agar plates containing 15 µg mL^{-1} zeocin in the dark. More than 800 obtained colonies were screened for decreased or not detectable ARS2 activity in the light and nine mutants that showed impaired inducibility of *GPX5-ARS* expression in light were selected for further analysis. Restriction enzyme site-directed amplification PCR[74] allowed us to identify the genomic DNA flanking the ble^R insertion sites in five out of the nine mutants. Only analysis of the *gunSOS1* mutant is presented here. Rescue of *gunSOS1* (*t6pp1*) was conducted with 5242 bp fragment of genomic DNA carrying 3241 bp *T6PP1* amplified by PCR (see Supplementary Table 11 for primers) using bacterial artificial chromosome PTQ5987 (Clemson University Genomics Institute, Clemson, SC, USA) as a template. The amplified DNA fragment included 1500 bp upstream and 501 bp downstream of the annotated *T6PP1*. Wild-type *T6PP1* was introduced into *gunSOS1* by co-transformation with a spectinomycin resistance cassette isolated from the pALM32 vector[75] with AleI and KpnI endonucleases. Transformant selection was performed on TAP agar plates supplemented with 100 µg mL^{-1} spectinomycin in the dark. All genetic transformations were performed by electroporation.

Arylsulfatase activity assay

To assess the level of *GPX5-ARS2* expression in transformed cells, enzymatic assays for arylsulfatase activity[76] were performed essentially as described before[14]. Cells were spotted onto agar-solidified TAP medium plates and cultured in either light or dark conditions. Arylsulfatase is expected to be secreted into the medium if *GPX5-ARS2* is expressed. After removal of the cells, plates were flooded with detection solution containing 0.1 mg mL^{-1} 2-naphthyl sulfate (potassium salt; Santa Cruz Biotechnology, Inc., Dallas, TX, USA) as a chromogenic substrate coupled with 1 mg mL^{-1} tetrazotized-o-dianisidine chloride (Fast Blue B salt, Santa Cruz Biotechnology, Inc., Dallas, TX, USA). Following 1 h incubation, purple spots appearing on the agar plates identified expressed ARS2.

Analysis of protoporphyrin IX content

The Proto content was analysed by High Pressure Liquid Chromatography (HPLC) essentially as described in Czarnecki, et al. [77], with modified sample preparation for *C. reinhardtii*[18]. In short, cultures were grown in the dark and transferred to 20 µmol photons m$^-$

545   ^{2} s^{-1} light for 2 h. Samples containing 1.2×10^{8} cells were centrifuged at 3000 x g for 5 min at

546   4°C and the pellets were snap-frozen in liquid N$_2$. Proto was extracted in cold (-20°C)

547   acetone/0.1 M NH$_4$OH (9/1, v/v) in a three-step cycle of resuspension and centrifugation.

548   Proto analysis was performed using a Nova-Pak C18 column (Waters, 3.9×150 mm, 4 μm,

549   at 20 °C). The results were normalized to pmol/10^{6} cells.

RNA isolation, qRT-PCR, and RNA-seq

The total RNA was isolated, after a shift from dark to 20 μmol photons m^{-2} s^{-1} light for 2 h,

using TRIzol Reagent (Thermo Fisher Scientific, Waltham, MA, USA), according to the

manufacturer's protocol. RNA quality was assessed by electrophoresis on a 1% (w/v) agarose

gel, while quantity was determined using a Nanodrop 2000 (Thermo Fischer Scientific,

Waltham, MA). Aliquots of 2 μg RNA were treated with DNase and RiboLock RNase

inhibitor (Thermo Fisher Scientific, Waltham, MA, USA) and used to synthesize cDNA with

RevertAid Reverse Transcriptase (Thermo Fisher Scientific, Waltham, MA, USA) and

oligo(dT)$_{18}$ primer. Transcript analysis by qRT-PCR were performed using 2× ChamQ

Universal SYBR qPCR Master Mix (Viazyme Biotech Co., Ltd., Nanjing, China) and a

CFX96-C1000 96-well plate thermocycler (Bio-Rad, Hercules, CA, USA), the 18S rRNA

was used as a reference gene[22]. All primers were designed using Primer3Plus

(http://primer3plus.com), and are listed in Supplementary Table 11.

RNA-sequencing was performed on total RNA samples isolated from biological

triplicates of *sigRep* and *gunSOS1* after the shift from dark to 20 μmol photons m^{-2} s^{-1} light

for 2 h. Library preparation, sequencing, and analysis services were commercially provided

by Novogene Europe (Novogene (UK) Company Ltd., Cambridge, UK) and performed using

an Illumina NovaSeq 6000 platform operated in 150 bp pair end sequencing mode, with a

sequencing depth of 20 million reads per sample. Six gigabases of sequencing data per library

was filtered for high-quality reads, which were mapped to the *C. reinhardtii* v5.5

(Department of Energy JGI). Estimated expression was obtained in fragments per kilobase of

transcript sequence per millions base pairs sequenced (FPKM). Biological replicates were

averaged to obtain sample expression estimate. Differential expression analysis between two

mutants (three biological replicates per mutant) was performed using DESeq2.v1.20.0, which

provides statistical routines for determining differential expression using a model based on

the negative binomial distribution. The resulting p values were adjusted using the Benjamini

and Hochberg's approach for controlling the False Discovery Rate (FDR). Genes with a

log$_2$FoldChange ≥ 2 and adjusted p value (padj) < 0.05 found by DESeq2.v1.20.0 were

defined as differentially expressed (DEGs). Gene Ontology (GO) analysis and Kyoto
Encyclopedia of Genes and Genomes (KEGG; www.genome.jp/kegg/pathway.html) pathway
analysis were conducted to identify DEGs at the biologically functional level using
clusterProfiler R package. GO terms and KEGG pathways with a padj < 0.05 were
considered to be significantly enriched.

Metabolite analysis
Cultures were grown in TAP in the dark until they reached mid-log phase of 3 x 10^6 cells mL$^-$
^{1}. Samples were always normalized to contain 6 x 10^7 cells, all centrifugation steps were
carried at 3000 x g at 4°C for 5 min, followed by snap-freezing in liquid N_2. Samples from
dark conditions were collected immediately, while the remainder of each culture was
transferred to 20 μmol photons m^{-2} s^{-1} light. Subsequent sampling took place at 15-min
intervals, up to 1 h. Metabolites were extracted using chloroform-methanol as described by
Lunn et al. (2006). Tre6P, other phosphorylated intermediates and organic acids were
measured by anion-exchange HPLC coupled to tandem mass spectrometry (LC-MS/MS) as
described in Lunn et al. (2006) with modifications as described in[78]. Sugars and sugar
alcohols were measured by LC-MS/MS as described in[79]. Obtained results were normalized
to pmol/10^6 cells.

For starch analysis, cultures were grown in TAP in the dark until they reached mid-
log phase of 3 x 10^6 cells mL^{-1}. For starch determination cultures were either transferred in
TAP to the light (20 μmol photons m^{-2} s^{-1}) for 2 h before harvesting, or centrifuged cells were
resuspended in Tris-phosphate (TP) medium (without acetate), keeping the same cell
concentration, and transferred back to the dark for 24 h. Subsequently, cultures were
transferred to 20 μmol photons m^{-2} s^{-1} for 2 h before harvesting. A similar protocol was
applied for starch analysis in TAP devoid of N (TAP-N), except that cells were cultivated in
TAP-N for 3 days in the dark, followed by exposure to 20 μmol photons m^{-2} s^{-1} for 2 h. For
each sampling, 5 x 10^6 cells were centrifuged and lyophilized. Starch content was determined
by enzymatic degradation and glucose quantification following the protocol of DuBois, et al.
[80].

Protein extraction and immunoblot analysis
Cultures were grown in the dark, followed by a transfer to 20 μmol photons m^{-2} s^{-1} for 3 h.
Cells were pelleted by centrifugation and total proteins were extracted in 400 μL buffer
containing: 56 mM Na_2CO_3, 56 mM DTT, 2% (w/v) SDS, 12% (w/v) sucrose and 2 mM

EDTA, pH 8.0. Proteins were quantified using Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA) and separated by SDS-PAGE on a 12% polyacrylamide gel, followed by transfer by electroblotting to nitrocellulose membrane (GE Healthcare, Chicago, IL, USA). T6PP1 was detected using a rabbit crude antiserum (dilution 1:500) raised by inoculation with a T6PP1-specific peptide (VEWSKSDSNGWRAKPC) against *C. reinhardtii* T6PP1 (calculated MW 42 kDa). GPX5 was detected with a commercially available antibody (AS15 2882, dilution 1:1000) obtained from Agrisera (Vännäs, Sweden). CHLI1 antibody (PHY5510S, dilution 1:1000) was purchased from PhytoAB (San Jose, CA, USA). For application, all antibodies were diluted in CrossDown buffer (AppliChem, AppliChem GmbH, Darmstadt, Germany). The secondary antibody (AS09 602, goat anti-rabbit IgG, dilution 1:10,000) conjugated to horseradish peroxidase was obtained from Agrisera. The immunoblotting signals were detected using a CCD camera (Intas Biopharmaceuticals, Ahmedabad, India) after application of enhanced chemiluminescence detection kit (Clarity™ Western ECL Substrate; Bio-Rad, Hercules, CA, USA.

Chemical treatments of *C. reinhardtii* cells

Fumarate (sodium fumarate dibasic) and aconitate (*cis*-aconitic acid) were obtained from Sigma-Aldrich (Sigma-Aldrich Chemie GmbH, Taufkirchen, Germany), dissolved in H_2O to a stock solution of 100 mM and added individually to the mid-log phase (3×10^6 cells mL^{-1}) cultures to final concentrations of 20, 50, and 100 μM, followed by exposure to light for 2 h.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

1   Johanningmeier, U. & Howell, S. H. Regulation of light-harvesting chlorophyll-binding protein mRNA accumulation in *Chlamydomonas reinhardii*. Possible involvement of chlorophyll synthesis precursors. *J. Biol. Chem.* **259**, 3541-3549 (1984).
2   Oelmüller, R. Photooxidative destruction of chloroplasts and its effect on nuclear gene expression and extraplastidic enzyme levels. *Photochem. Photobiol.* **49**, 229-239, doi:10.1111/j.1751-1097.1989.tb04101.x (1989).
3   Susek, R. E., Ausubel, F. M. & Chory, J. Signal transduction mutants of Arabidopsis uncouple nuclear CAB and RBCS gene expression from chloroplast development. *Cell* **74**, 787-799 (1993).

4   Oster, U., Brunner, H. & Rudiger, W. The greening process in cress seedlings .5. Possible
interference of chlorophyll precursors, accumulated after thujaplicin treatment, with light-
regulated expression of Lhc genes. *J. Photoch. Photobio. B* **36**, 255-261, doi:10.1016/s1011-
1344(96)07388-5 (1996).
5   Lydon, J. & Duke, S. O. Porphyrin synthesis is required for photobleaching activity of the
para-nitrosubstituted diphenyl ether herbicides. *Pestic. Biochem. Physiol.* **31**, 74-83,
doi:10.1016/0048-3575(88)90031-4 (1988).
6   Triantaphylides, C. *et al.* Singlet oxygen is the major reactive oxygen species involved in
photooxidative damage to plants. *Plant Physiol.* **148**, 960-968, doi:10.1104/pp.108.125690
(2008).
7   op den Camp, R. G. L. *et al.* Rapid induction of distinct stress responses after the release of
singlet oxygen in Arabidopsis. *Plant Cell* **15**, 2320-2332, doi:10.1105/tpc.014662 (2003).
8   Baruah, A., Simkova, K., Apel, K. & Laloi, C. Arabidopsis mutants reveal multiple singlet
oxygen signaling pathways involved in stress response and development. *Plant Mol. Biol.* **70**,
547-563, doi:10.1007/s11103-009-9491-0 (2009).
9   Gorman, A. A. & Rodgers, M. A. J. Current perspectives of singlet oxygen detection in
biological environments. *J. Photoch. Photobio. B* **14**, 159-176 (1992).
10 Beck, C. F. Signaling pathways from the chloroplast to the nucleus. *Planta* **222**, 743-756,
doi:10.1007/s00425-005-0021-2 (2005).
11 Sies, H. & Menck, C. F. M. Singlet oxygen induced DNA damage. *Mutat. Res.* **275**, 367-375
(1992).
12 Pfannschmidt, T. Plastidial retrograde signalling - a true "plastid factor" or just metabolite
signatures? *Trends Plant Sci.* **15**, 427-435, doi:10.1016/j.tplants.2010.05.009 (2010).
13 Kim, C. *et al.* Chloroplasts of Arabidopsis are the source and a primary target of a plant-
specific programmed cell death signaling pathway. *Plant Cell* **24**, 3026-3039,
doi:10.1105/tpc.112.100479 (2012).
14 Brzezowski, P., Wilson, K. E. & Gray, G. R. The PSBP2 protein of Chlamydomonas
reinhardtii is required for singlet oxygen-dependent signaling. *Planta* **236**, 1289-1303,
doi:10.1007/s00425-012-1683-1 (2012).
15 Wakao, S. *et al.* Phosphoprotein SAK1 is a regulator of acclimation to singlet oxygen in
Chlamydomonas reinhardtii. *eLife* **3**, e02286, doi:10.7554/eLife.02286 (2014).
16 Shao, N., Duan, G. Y. & Bock, R. A mediator of singlet oxygen responses in
Chlamydomonas reinhardtii and Arabidopsis identified by a luciferase-based genetic screen in
algal cells. *Plant Cell* **25**, 4209-4226, doi:10.1105/tpc.113.117390 (2013).
17 von Gromoff, E. D., Alawady, A., Meinecke, L., Grimm, B. & Beck, C. F. Heme, a plastid-
derived regulator of nuclear gene expression in *Chlamydomonas*. *Plant Cell* **20**, 552-567,
doi:10.1105/tpc.107.054650 (2008).
18 Brzezowski, P. *et al.* The GUN4 protein plays a regulatory role in tetrapyrrole biosynthesis
and chloroplast-to-nucleus signalling in *Chlamydomonas reinhardtii*. *Plant J.* **79**, 285-298,
doi:10.1111/tpj.12560 (2014).
19 Fischer, B. B., Krieger-Liszkay, A. & Eggen, R. I. L. Photosensitizers neutral red (Type I)
and rose bengal (Type II) cause light-dependent toxicity in Chlamydomonas reinhardtii and
induce the Gpxh gene via increased singlet oxygen formation. *Environ. Sci. Technol.* **38**,
6307-6313, doi:10.1021/es049673y (2004).
20 Leisinger, U. *et al.* The glutathione peroxidase homologous gene from Chlamydomonas
reinhardtii is transcriptionally up-regulated by singlet oxygen. *Plant Mol. Biol.* **46**, 395-408
(2001).
21 Fischer, B. B., Il Eggen, R. & Niyogi, K. K. Characterization of singlet oxygen-accumulating
mutants isolated in a screen for altered oxidative stress response in Chlamydomonas
reinhardtii. *BMC Plant Biol.* **10**, doi:279 10.1186/1471-2229-10-279 (2010).
22 Fischer, B. B. *et al.* Function and regulation of the glutathione peroxidase homologous gene
GPXH/GPX5 in Chlamydomonas reinhardtii. *Plant Mol. Biol.* **71**, 569-583,
doi:10.1007/s11103-009-9540-8 (2009).

23 Lunn, J. E. *et al.* Sugar-induced increases in trehalose 6-phosphate are correlated with redox
activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in
Arabidopsis thaliana. *Biochem. J.* **397**, 139-148, doi:10.1042/bj20060083 (2006).
24 Yadav, U. P. *et al.* The sucrose-trehalose 6-phosphate (Tre6P) nexus: specificity and
mechanisms of sucrose signalling by Tre6P. *J. Exp. Bot.* **65**, 1051-1068,
doi:10.1093/jxb/ert457 (2014).
25 Schluepmann, H., Pellny, T., van Dijken, A., Smeekens, S. & Paul, M. Trehalose 6-phosphate
is indispensable for carbohydrate utilization and growth in Arabidopsis thaliana. *Proc Natl*
Acad Sci U S A **100**, 6849-6854, doi:10.1073/pnas.1132018100 (2003).
26 Fichtner, F. & Lunn, J. E. The Role of Trehalose 6-Phosphate (Tre6P) in Plant Metabolism
and Development. *Annu. Rev. Plant Biol.* **72**, 737-760, doi:10.1146/annurev-arplant-050718-
095929 (2021).
27 Miller, R. *et al.* Changes in transcript abundance in Chlamydomonas reinhardtii following
nitrogen deprivation predict diversion of metabolism. *Plant Physiol.* **154**, 1737-1752,
doi:10.1104/pp.110.165159 (2010).
28 Ajalla Aleixo, M. A., Rangel, V. L., Rustiguel, J. K., de Pádua, R. A. P. & Nonato, M. C.
Structural, biochemical and biophysical characterization of recombinant human fumarate
hydratase. *FEBS J.* **286**, 1925-1940, doi:10.1111/febs.14782 (2019).
29 Yogev, O. *et al.* Fumarase: a mitochondrial metabolic enzyme and a cytosolic/nuclear
component of the DNA damage response. *PLoS Biol.* **8**, e1000328,
doi:10.1371/journal.pbio.1000328 (2010).
30 Jiang, Y. *et al.* Local generation of fumarate promotes DNA repair through inhibition of
histone H3 demethylation. *Nat. Cell Biol.* **17**, 1158-1168, doi:10.1038/ncb3209 (2015).
31 Ma, X. *et al.* Transcriptomic and Physiological Responses to Oxidative Stress in a
Chlamydomonas reinhardtii Glutathione Peroxidase Mutant. *Genes* **11**,
doi:10.3390/genes11040463 (2020).
32 Blaby, I. K. *et al.* Genome-wide analysis on Chlamydomonas reinhardtii reveals the impact of
hydrogen peroxide on protein stress responses and overlap with other stress transcriptomes.
Plant J. **84**, 974-988, doi:10.1111/tpj.13053 (2015).
33 Urzica, E. I. *et al.* Impact of oxidative stress on ascorbate biosynthesis in Chlamydomonas via
regulation of the VTC2 gene encoding a GDP-L-galactose phosphorylase. *J. Biol. Chem.* **287**,
14234-14245, doi:10.1074/jbc.M112.341982 (2012).
34 Roach, T., Stöggl, W., Baur, T. & Kranner, I. Distress and eustress of reactive electrophiles
and relevance to light stress acclimation via stimulation of thiol/disulphide-based redox
defences. *Free radical biology & medicine* **122**, 65-73,
doi:10.1016/j.freeradbiomed.2018.03.030 (2018).
35 Arora, A., Sairam, R. K. & Srivastava, G. C. Oxidative stress and antioxidative system in
plants. *Curr. Sci.* **82**, 1227-1238 (2002).
36 Moller, I. M. Plant mitochondria and oxidative stress: Electron transport, NADPH turnover,
and metabolism of reactive oxygen species. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**,
561-591 (2001).
37 Asada, K. Production and scavenging of reactive oxygen species in chloroplasts and their
functions. *Plant Physiol.* **141**, 391-396, doi:10.1104/pp.106.082040 (2006).
38 Brzezowski, P., Richter, A. S. & Grimm, B. Regulation and function of tetrapyrrole
biosynthesis in plants and algae. *Biochimica Et Biophysica Acta-Bioenergetics* **1847**, 968-
985, doi:10.1016/j.bbabio.2015.05.007 (2015).
39 Larkin, R. M. Tetrapyrrole Signaling in Plants. *Front. Plant Sci.* **7**,
doi:10.3389/fpls.2016.01586 (2016).
40 Moulin, M., McCormac, A. C., Terry, M. J. & Smith, A. G. Tetrapyrrole profiling in
Arabidopsis seedlings reveals that retrograde plastid nuclear signaling is not due to Mg-
protoporphyrin IX accumulation. *Proc. Natl. Acad. Sci. USA* **105**, 15178-15183,
doi:10.1073/pnas.0803054105 (2008).
41 Mochizuki, N., Tanaka, R., Tanaka, A., Masuda, T. & Nagatani, A. The steady-state level of
Mg-protoporphyrin IX is not a determinant of plastid-to-nucleus signaling in Arabidopsis.
Proc. Natl. Acad. Sci. USA **105**, 15184-15189, doi:10.1073/pnas.0803245105 (2008).

42 Vandesteene, L. *et al.* Expansive evolution of the trehalose-6-phosphate phosphatase gene
family in Arabidopsis. *Plant Physiol.* **160**, 884-896, doi:10.1104/pp.112.201400 (2012).
43 Ramon, M. *et al.* Extensive expression regulation and lack of heterologous enzymatic activity
of the Class II trehalose metabolism proteins from Arabidopsis thaliana. *Plant Cell Environ.*
32, 1015-1032, doi:10.1111/j.1365-3040.2009.01985.x (2009).
44 Vandesteene, L., Ramon, M., Le Roy, K., Van Dijck, P. & Rolland, F. A single active
trehalose-6-P synthase (TPS) and a family of putative regulatory TPS-like proteins in
Arabidopsis. *Mol. Plant* **3**, 406-419, doi:10.1093/mp/ssp114 (2010).
45 Delorge, I., Figueroa, C. M., Feil, R., Lunn, J. E. & Van Dijck, P. Trehalose-6-phosphate
synthase 1 is not the only active TPS in Arabidopsis thaliana. *Biochem. J.* **466**, 283-290,
doi:10.1042/bj20141322 (2015).
46 Collet, J. F., Stroobant, V., Pirard, M., Delpierre, G. & Van Schaftingen, E. A new class of
phosphotransferases phosphorylated on an aspartate residue in an amino-terminal
DXDX(T/V) motif. *J. Biol. Chem.* **273**, 14107-14112, doi:10.1074/jbc.273.23.14107 (1998).
47 Martins, M. C. *et al.* Feedback inhibition of starch degradation in Arabidopsis leaves
mediated by trehalose 6-phosphate. *Plant Physiol.* **163**, 1142-1163,
doi:10.1104/pp.113.226787 (2013).
48 Dos Anjos, L. *et al.* Feedback regulation by trehalose 6-phosphate slows down starch
mobilization below the rate that would exhaust starch reserves at dawn in Arabidopsis leaves.
Plant direct **2**, e00078, doi:10.1002/pld3.78 (2018).
49 Zhang, Y. *et al.* Inhibition of SNF1-related protein kinase1 activity and regulation of
metabolic pathways by trehalose-6-phosphate. *Plant Physiol.* **149**, 1860-1871,
doi:10.1104/pp.108.133934 (2009).
50 Debast, S. *et al.* Altering trehalose-6-phosphate content in transgenic potato tubers affects
tuber growth and alters responsiveness to hormones during sprouting. *Plant Physiol.* **156**,
1754-1771, doi:10.1104/pp.111.179903 (2011).
51 Nunes, C. *et al.* Inhibition of SnRK1 by metabolites: tissue-dependent effects and cooperative
inhibition by glucose 1-phosphate in combination with trehalose 6-phosphate. *Plant Physiol.*
Biochem. **63**, 89-98, doi:10.1016/j.plaphy.2012.11.011 (2013).
52 Zhai, Z. *et al.* Trehalose 6-Phosphate Positively Regulates Fatty Acid Synthesis by Stabilizing
WRINKLED1. *Plant Cell* **30**, 2616-2627, doi:10.1105/tpc.18.00521 (2018).
53 Baena-González, E. & Lunn, J. E. SnRK1 and trehalose 6-phosphate - two ancient pathways
converge to regulate plant metabolism and growth. *Curr. Opin. Plant Biol.* **55**, 52-59,
doi:10.1016/j.pbi.2020.01.010 (2020).
54 Depaepe, T. *et al.* At the Crossroads of Survival and Death: The Reactive Oxygen Species-
Ethylene-Sugar Triad and the Unfolded Protein Response. *Trends Plant Sci.* **26**, 338-351,
doi:10.1016/j.tplants.2020.12.007 (2021).
55 Baena-González, E., Rolland, F., Thevelein, J. M. & Sheen, J. A central integrator of
transcription networks in plant stress and energy signalling. *Nature* **448**, 938-942,
doi:10.1038/nature06069 (2007).
56 Gonzalez-Ballester, D., Pollock, S. V., Pootakham, W. & Grossman, A. R. The central role of
a SNRK2 kinase in sulfur deprivation responses. *Plant Physiol.* **147**, 216-227,
doi:10.1104/pp.108.116137 (2008).
57 González-Ballester, D. *et al.* RNA-seq analysis of sulfur-deprived Chlamydomonas cells
reveals aspects of acclimation critical for cell survival. *Plant Cell* **22**, 2058-2084,
doi:10.1105/tpc.109.071167 (2010).
58 Colina, F. *et al.* Genome-wide identification and characterization of CKIN/SnRK gene family
in Chlamydomonas reinhardtii. *Sci. Rep.* **9**, 350, doi:10.1038/s41598-018-35625-8 (2019).
59 Valledor, L., Furuhashi, T., Hanak, A. M. & Weckwerth, W. Systemic cold stress adaptation
of Chlamydomonas reinhardtii. *Mol. Cell. Proteomics* **12**, 2032-2047,
doi:10.1074/mcp.M112.026765 (2013).
60 Yang, M., Soga, T., Pollard, P. J. & Adam, J. The emerging role of fumarate as an
oncometabolite. *Frontiers in oncology* **2**, 85, doi:10.3389/fonc.2012.00085 (2012).

61 O'Flaherty, L. *et al.* Dysregulation of hypoxia pathways in fumarate hydratase-deficient cells
is independent of defective mitochondrial metabolism. *Hum. Mol. Genet.* **19**, 3844-3851,
doi:10.1093/hmg/ddq305 (2010).
62 Shanmugasundaram, K. *et al.* The oncometabolite fumarate promotes pseudohypoxia through
noncanonical activation of NF-κB signaling. *J. Biol. Chem.* **289**, 24691-24699,
doi:10.1074/jbc.M114.568162 (2014).
63 Ternette, N. *et al.* Inhibition of mitochondrial aconitase by succination in fumarate hydratase
deficiency. *Cell reports* **3**, 689-700, doi:10.1016/j.celrep.2013.02.013 (2013).
64 Lloyd, S. J., Lauble, H., Prasad, G. S. & Stout, C. D. The mechanism of aconitase: 1.8 A
resolution crystal structure of the S642a:citrate complex. *Protein Sci.* **8**, 2655-2662,
doi:10.1110/ps.8.12.2655 (1999).
65 Kleine, T., Voigt, C. & Leister, D. Plastid signalling to the nucleus: messengers still lost in
the mists? *Trends Genet.* **25**, 185-190, doi:10.1016/j.tig.2009.02.004 (2009).
66 Bräutigam, K. *et al.* Dynamic plastid redox signals integrate gene expression and metabolism
to induce distinct metabolic states in photosynthetic acclimation in Arabidopsis. *Plant Cell*
21, 2715-2732, doi:10.1105/tpc.108.062018 (2009).
67 Pesaresi, P. *et al.* Nuclear photosynthetic gene expression is synergistically modulated by
rates of protein synthesis in chloroplasts and mitochondria. *Plant Cell* **18**, 970-991,
doi:10.1105/tpc.105.039073 (2006).
68 Shapiguzov, A. *et al.* Arabidopsis RCD1 coordinates chloroplast and mitochondrial functions
through interaction with ANAC transcription factors. *eLife* **8**, doi:10.7554/eLife.43284
(2019).
69 Dent, R. M., Haglund, C. M., Chin, B. L., Kobayashi, M. C. & Niyogi, K. K. Functional
genomics of eukaryotic photosynthesis using insertional mutagenesis of Chlamydomonas
reinhardtii. *Plant Physiol.* **137**, 545-556, doi:10.1104/pp.104.055244 (2005).
70 Depege, N., Bellafiore, S. & Rochaix, J. D. Role of chloroplast protein kinase Stt7 in LHCII
phosphorylation and state transition in Chlamydomonas. *Science* **299**, 1572-1575 (2003).
71 Sizova, I., Fuhrmann, M. & Hegemann, P. A Streptomyces rimosus aphVIII gene coding for a
new type phosphotransferase provides stable antibiotic resistance to Chlamydomonas
reinhardtii. *Gene* **277**, 221-229 (2001).
72 Davies, J. P., Weeks, D. P. & Grossman, A. R. Expression of the arylsulfatase gene from the
β2-tubulin promoter in Chlamydomonas reinhardtii. *Nucleic Acids Res.* **20**, 2959-2965
(1992).
73 Schroda, M., Beck, C. F. & Vallon, O. Sequence elements within an HSP70 promoter
counteract transcriptional transgene silencing in Chlamydomonas. *Plant J.* **31**, 445-455,
doi:10.1046/j.1365-313X.2002.01371.x (2002).
74 Gonzalez-Ballester, D., de Montaigu, A., Galvan, A. & Fernandez, E. Restriction enzyme
site-directed amplification PCR: A tool to identify regions flanking a marker DNA. *Anal.*
Biochem. **340**, 330-335, doi:10.1016/j.ab.2005.01.031 (2005).
75 Meslet-Cladière, L. & Vallon, O. Novel shuttle markers for nuclear transformation of the
green alga Chlamydomonas reinhardtii. *Eukaryot. Cell* **10**, 1670-1678, doi:10.1128/ec.05043-
11 (2011).
76 Ohresser, M., Matagne, R. F. & Loppes, R. Expression of the arylsulphatase reporter gene
under the control of the nit1 promoter in Chlamydomonas reinhardtii. *Curr. Genet.* **31**, 264-
271 (1997).
77 Czarnecki, O., Peter, E. & Grimm, B. in *Chloroplast research in Arabidopsis: Methods and*
Protocols, Vol II Vol. 775   357-385 (2011).
78 Figueroa, C. M. *et al.* Trehalose 6-phosphate coordinates organic and amino acid metabolism
with carbon availability. *Plant J.* **85**, 410-423, doi:10.1111/tpj.13114 (2016).
79 Fichtner, F. *et al.* Trehalose 6-phosphate is involved in triggering axillary bud outgrowth in
garden pea (Pisum sativum L.). *Plant J.* **92**, 611-623, doi:10.1111/tpj.13705 (2017).
80 DuBois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A. & Smith, F. Colorimetric Method
for Determination of Sugars and Related Substances. *Analytical chemistry* **28**, 350-356,
doi:10.1021/ac60111a017 (1956).

81   Johnson, X. & Alric, J. Central carbon metabolism and electron transport in Chlamydomonas
reinhardtii: metabolic constraints for carbon partitioning between oil and starch. *Eukaryot.*
Cell **12**, 776-793, doi:10.1128/ec.00318-12 (2013).
82   Findinier, J. *et al.* The Chlamydomonas mex1 mutant shows impaired starch mobilization
without maltose accumulation. *J. Exp. Bot.* **68**, 5177-5189, doi:10.1093/jxb/erx343 (2017).

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (DFG-GR 936/20-2 to B.G.), by the Agence Nationale de la Recherche (ChloroPaths: ANR-14-CE05-0041-01 to X.J.), and by the Max Planck Society (R.F. and J.E.L.).

Author contributions

P.B. designed the research. P.B. and W.A.Y. performed most of the experiments. R.F. and J.E.L. performed the LC-MS/MS measurements. M.S.-S. and X.J. performed the starch quantification. P.B., B.G., and J.E.L. analysed the data. P.B. wrote the manuscript. P.B., B.G., J.E.L., and X.J. revised the manuscript. All authors discussed the results and commented upon the manuscript.

Competing interests

The authors declare no competing financial interests.

Additional information

Supplementary information accompanies this manuscript:

Supplementary Figure 1. Schematic representation of tetrapyrrole biosynthesis pathway.

Supplementary Figure 2. Schematic representation of the *GPX5-ARS2* reporter gene construct.

Supplementary Figure 3. Analysis of Proto content in *gunSOS1*.

Supplementary Figure 4. Rescue of the $^1O_2$-signalling in *gunSOS1* by introduction of the wild-type *T6PP1*.

Supplementary Figure 5. Metabolites content in *gunSOS1* compared to *sigRep*.

Supplementary Figure 6. Accumulation of fumarate in *gunSOS1* compared to *sigRep* could be explained by increased expression of *FUMARATE HYDRATASE 2*.

Supplementary Figure 7. Accumulation of fumarate in *gunSOS1* compared to *sigRep* could be explained by increased expression of *FUMARYLACETOACETASE*.

Supplementary Figure 8. Deficiency in aconitate in *gunSOS1* compared to *sigRep* could be
explained by decreased expression of *ACONITATE HYDRATASE 1*.
Supplementary Table 1. Metabolites analyzed in *gunSOS1* compared to *sigRep* (LC-
MS/MS).
Supplementary Table 2. Impact of accumulating metabolites on cell metabolism.
Supplementary Table 3. Impact of deficient metabolites on cell metabolism.
Supplementary Table 4. Statistical analyses for $GPX5_{cyt}$ expression following treatment
with fumarate, presented in Figure 7a.
Supplementary Table 5. Statistical analyses for $GPX5_{cp}$ expression following treatment with
fumarate, presented in Figure 7a.
Supplementary Table 6. Statistical analyses for $GPX5_{cyt}$ expression following treatment
with aconitate, presented in Figure 7b.
Supplementary Table 7. Statistical analyses for $GPX5_{cp}$ expression following treatment with
aconitate, presented in Figure 7b.
Supplementary Table 8. Statistical analyses for *SAK1* expression following treatment with
aconitate, presented in Figure 7c.
Supplementary Table 9. Statistical analyses for *PSBP2* expression following treatment with
aconitate, presented in Figure 7c.
Supplementary Table 10. Statistical analyses for *MBS* expression following treatment with
aconitate, presented in Figure 7c.
Supplementary Table 11. Primers used in this study.
Supplementary Data Set 1. Co-expression analysis
Supplementary Data Set 2. DEG analysis
Supplementary Data Set 3. KEGG analysis
922

Figure 1. Generation and initial characterization of the mutant impaired in $^1O_2$-
signalling. a Arylsulfatase (ARS2) assay for selection of the strain expressing *GPX5-ARS2*
reporter gene in a $^1O_2$-inducible manner. The mutant selected for further applications was
named *signalling Reporter* (*sigRep*). WT and *chlD-1/GUN4* do not carry *GPX5-ARS2* and
were used as negative controls in ARS2 assay. **b** Expression kinetics of cytosolic (*GPX5$_{cyt}$*)
and chloroplast *GPX5* (*GPX5$_{cp}$*) in *sigRep* compared to WT upon exposure to light; the
values were normalized to WT in dark. **c** Expression of *GPX5-ARS2* in *sigRep* positively
correlates with time of exposure to light. WT was used as a negative control. **d** ARS2 activity
assay following mutagenesis of *sigRep*. Screen was performed to identify mutants not
expressing *GPX5-ARS2* compared to *sigRep* in the same conditions. The mutant selected for
further applications was *named genomes uncoupled Singlet Oxygen Signalling 1* (*gunSOS1*).
WT and *chlD-1/GUN4* were used as negative controls. **e** Strains with impaired $^1O_2$-signalling
demonstrated increased sensitivity to light compared to *sigRep* and *chlD-1/GUN4*. Higher
light tolerance of *chlD-1/GUN4* and *sigRep* compared to *chlD-1* is also visible. **f** *T6PP1*
(Cre12.g497750) gene model. The insertion (ble^R) was identified in the first exon of 3241 bp
T6PP1. **g** Expression of *T6PP1* in *gunSOS1* compared to *sigRep*. **h** Expression of *GPX5$_{cyt}$*
and *GPX5$_{cp}$* in *gunSOS1* compared to *sigRep*. **i** Expression of *GPX5-ARS2* in *gunSOS1*
compared to *sigRep*. WT was used as a negative control. Transcript analyses were performed
by qRT-PCR on biological triplicates, values were calculated either as a normalized fold
expression ($2^{-\Delta\Delta Ct}$; WT=1) or relative expression ($2^{\Delta Ct}$). For b and c, horizontal bars represent
the calculated mean ($n = 3$), vertical error bars represent calculated ±SD; significant
differences were calculated using two-tailed Student's *t*-test and are indicated by asterisks
(non-significant not shown), $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. For
g-i, error bars indicate calculated ±SD; one-way ANOVA, pair-wise comparison with the
Tukey's post-hoc test (non-significant not shown), $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and
$****P < 0.0001$.

Figure 2. *C. reinhardtii* **T6PP1 is a functional phosphatase induced during photooxidative stress. a** Kinetics of Tre6P accumulation in *gunSOS1* compared to *sigRep* upon transfer from dark to light. **b** Kinetics of *T6PP1* expression upon transfer from dark to light in *gunSOS1* compared to WT indicate inducibility by photooxidative stress rather than light; values were calculated as a normalized fold expression ($2^{-\Delta\Delta Ct}$; WT=1). **c** Higher content of trehalose in *gunSOS1* compared to *sigRep* upon transfer from dark to light may be resulting from the function of T6PS/T6PP1. **d** Immuno-blot analysis of T6PP1 (calculated MW of 42 kDa) and GPX5 content in *gunSOS1* compared to *sigRep* and strains rescued with wild-type copy of *T6PP1*. The CHLI1 content was used as a loading control. The unspecific immunoreaction is indicated by an asterisk. **e** Light sensitivity examination showed that the strains with rescued $^1O_2$-signalling have an increased tolerance to light compared to *gunSOS1*. Experiments in a, b and c were performed in biological triplicates (*n* = 3) horizontal bars represent the calculated mean, vertical error bars represent ±SD; significant differences were calculated using two-tailed Student's *t*-test and are indicated by asterisks (non-significant not shown), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.

Figure 3. Metabolites with significantly increased content in *gunSOS1* compared to *sigRep* in the light. a LC-MS/MS analysis; fumaric acid (fumarate), oxoglutaric acid (2-OG), myo-Inositol, ADP-glucose (ADPGlc), galactose 1-phosphate (Gal1P), fructose 1-phosphate (Fru1P), alpha-D-Glucose 1,6-bisphosphate (Glc1, 6BP), glycerol 3-phosphate (Gly3P), phosphoenolpyruvic acid (PEP), L-Malic acid (malate). Measurements were performed in biological triplicates and are presented as mean (n = 3), horizontal bars represent the calculated mean, vertical error bars represent calculated ±SD; significant differences were calculated using two-tailed Student's t-test and are indicated by asterisks (non-significant not shown), *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001. **b** Impact of accumulating metabolites on cell metabolism analysed using MetaboAnalyst 5.0 (https://www.metaboanalyst.ca), sorted by the P value; 1. Tricarboxylic acid cycle (TCA cycle; P = 4.41E-05), 2. Pyruvate metabolism (P = 0.00136), 3. Starch and sucrose metabolism (P = 0.00136), 4. Glyoxylate and dicarboxylate metabolism (P = 0.00429), 5. Arginine biosynthesis (P = 0.01473), 6. Alanine, aspartate and glutamate metabolism (P = 0.02019), 7. Glycerolipid metabolism (P = 0.02019), 8. Carbon fixation in photosynthetic organisms (P = 0.02217), 9. Galactose metabolism (P = 0.03566), 10. Amino sugar and nucleotide sugar metabolism (P = 0.07616), 11. Fructose and mannose metabolism (P = 0.13874), 12. Tyrosine metabolism (P = 0.1872), 13. Phenylalanine, tyrosine and tryptophan biosynthesis (P = 0.22414), 14. Inositol phosphate metabolism (P = 0.24202), 15. Glycolysis / Gluconeogenesis (P = 0.24202), 16. Phosphatidylinositol signaling system (P = 0.25082), 17. Glycine, serine and threonine metabolism (P = 0.27665), 18. Glycerophospholipid metabolism (P = 0.34157); detailed results are presented in Supplementary Table 2.

Figure 4. Metabolites with significantly decreased content in *gunSOS1* compared to *sigRep* in the light. a LC-MS/MS analysis; cis-Aconitic acid (aconitate), mannose 6-phosphate (Man6P), Glucose 6-phosphate (Glc6P), D-fructose (frictose), Fructose 6-phosphate (Fru6P), Glucose 1-phosphate (Glc1P), Fructose 1,6-bisphosphate (Fru1, 6BP), sorbitol. Measurements were performed in biological triplicates and are presented as mean ($n = 3$), horizontal bars represent the calculated mean, vertical error bars represent calculated ±SD; significant differences were calculated using two-tailed Student's *t*-test and are indicated by asterisks (non-significant not shown), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$. **b** Impact of deficient metabolites on cell metabolism analysed using MetaboAnalyst 5.0 (https://www.metaboanalyst.ca), sorted by the *P* value; 1. Fructose and mannose metabolism ($P = 6.23E-05$), 2. Amino sugar and nucleotide sugar metabolism ($P = 8.99E-05$), 3. Starch and sucrose metabolism ($P = 0.000282$), 4. Glycolysis / Gluconeogenesis ($P = 0.01094$), 5. Galactose metabolism ($P = 0.013767$), 6. Pentose and glucuronate interconversions ($P = 0.081139$), 7. Pentose phosphate pathway ($P = 0.1195$), 8. Tricarboxylic acid cycle (TCA cycle; $P = 0.13197$), 9. Glycerolipid metabolism ($P = 0.13197$), 10. Carbon fixation in photosynthetic organisms ($P = 0.13815$), 11. Inositol phosphate metabolism ($P = 0.15646$), 12. Glyoxylate and dicarboxylate metabolism ($P = 0.19785$); detailed results are presented in Supplementary Table 3. **c** Starch accumulation in *gunSOS1* compared to *sigRep*. Measurements were performed in biological triplicates ($n = 3$); median is shown as a center line; upper and lower quartiles are shown as dotted lines. Significant differences were calculated using two-tailed Student's *t*-test and are indicated by asterisks (non-significant not shown), *$P < 0.05$, **$P < 0.01$.

1050
1051

Figure 5. The overview of the selected metabolic pathways in *gunSOS1* compared to *sigRep*. The map was constructed based on the measured metabolites presented in Figures 3, 4, and Supplementary Figure 5. Metabolites increased in *gunSOS1* compared to *sigRep*: fumaric acid (fumarate), oxoglutaric acid (2-OG), myo-Inositol, ADP-glucose (ADPGlc), galactose 1-phosphate (Gal1P), fructose 1-phosphate (Fru1P), alpha-D-Glucose 1,6-bisphosphate (Glc1, 6BP), glycerol 3-phosphate (Gly3P), phosphoenolpyruvic acid (PEP), L-Malic acid (malate). Metabolites decreased in *gunSOS1* compared to *sigRep*: cis-Aconitic acid (aconitate), mannose 6-phosphate (Man6P), Glucose 6-phosphate (Glc6P), D-fructose (frictose), Fructose 6-phosphate (Fru6P), Glucose 1-phosphate (Glc1P), Fructose 1,6-bisphosphate (Fru1, 6BP), and sorbitol. Metabolites that did not show significant difference in gunSOS1 relative to sigRep: D-Glucose (glucose), D-Maltose (maltose), maltotriose, citric acid (citrate), isocitric acid (isocitrate), pyruvic acid (pyruvate), succinic acid (succinate), 3-phosphoglyceric acid (3PGA). The colour-coding: blue = increased, red = decreased, black = unchanged, grey = not measured or [a]below the limit of detection. Significant differences were calculated using two-tailed Student's *t*-test. The maximal significant differences from any time point in light (Fig. 4 and 5) are indicated by asterisks, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. The metabolic map was constructed based on the review by[81] and[82].

1073
Figure 6. Transcriptome analysis based on RNA-seq. a Venn diagram showing number of
genes expressed in *gunSOS1* and *sigRep* (8120), or uniquely expressed in *gunSOS1* (1377) or
in *sigRep* (1087). **b** Pathway enrichment analysis (KEGG, http://www.kegg.jp/) based on the
differentially expressed genes (RNA-seq); genes down-regulated in *gunSOS1* compared to
sigRep. **c** KEGG analysis showing genes up-regulated in *gunSOS1* compared to *sigRep.*
Effect on metabolic processes is sorted by the *p* value. Asterisk indicates the core
(significant) enrichment.

1081

Figure 7. The qRT-PCR analysis of gene expression in *gunSOS1* compared to *sigRep*. a
Expression of genes encoding proteinaceous components previously associated with the
chloroplast retrograde signalling involving $^1O_2$, *PSBP2*[14], *SAK1*[15], and *MBS*[16]. **b** transcript
levels of the selected genes showing attenuated expression in *gunSOS1* compared to *sigRep*
in the context of the SAK1-transcriptome[15], genes description in the text. Experiments were
performed in biological replications; results are presented as normalized fold expression ($2^{-\Delta\Delta Ct}$, WT=1); the error bars represent calculated ±SD. Significant differences were calculated
using one-way ANOVA, pair-wise comparison with the Tukey's post-hoc test (non-
significant not shown), $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$.

Figure 8. Effect of selected metabolites applied exogenously on $^1O_2$-signalling inducing $GPX5_{cyt}$ or $GPX5_{cp}$ expression. a Fumarate (Fum) significantly ($P < 0.0001$) decreased $GPX5$ expression in *sigRep* in concentration-dependent manner. Interaction between mutants and fumarate, as well as the mutant-dependent expression of $GPX5$ were also significant, $P < 0.0001$ and $P < 0.001$, respectively. Fumarate had no effect on $GPX5_{cyt}$ expression in *gunSOS1* or WT, but attenuating effect was observed for $GPX5_{cp}$ expression in WT ($P \leq 0.0001$). Detailed reports from statistical analyses are presented in Supplementary Tables 4 and 5. **b** Aconitate (Acon) at 20 µM concentration rescues $GPX5$ expression in *gunSOS1* and increases $GPX5$ expression in *sigRep*. Statistical analysis showed that 20 µM aconitate significantly ($P \leq 0.0001$) affect $GPX5_{cyt}$ expression in *gunSOS1* and *sigRep*, but 100 µM concentration ($P = 0.0001$) decreased this expression in *sigRep*. An increase in $GPX5_{cp}$ expression ($P < 0.05$) was also observed in WT upon treatment with 50 µM aconitate (Supplementary Table 6 and 7). **c** Transcripts content for proteins necessary for the $^1O_2$-signalling, *SAK1*, *PSBP2*, and *MBS* significantly ($P < 0.01$) increased upon treatment with 20 µM aconitate. An increase in *PSBP2* and *MBS* expression was also observed in WT treated with 50 µM aconitate. Detailed statistical analyses are presented in Supplementary Tables 8-10. The qRT-PCR experiments were performed in biological replications ($n = 3$); results are presented as normalized fold expression ($2^{-\Delta\Delta Ct}$, untreated WT=1); the error bars represent calculated ±SD. Significant differences were calculated using two-way ANOVA with Dunnett's multiple comparison test and are indicated by asterisks (non-significant changes are not shown on the graph for clarity), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.

1137
Figure 9. Expression of genes, which induction was previously associated with H_2O_2 or organic peroxides, but not with $^1O_2$. a *Mn superoxide dismutase 3* (*MSD3*). **b** *ASCORBATE PEROXIDASE 1* (*APX1*). **c** *Mono-functional catalase* (*CAT1*). **d** *GDP-L-galactose Phosphorylase* (*VTC2*). **e** *monodehydroascorbate reductase* (*MDAR1*). **f** *dehydrosascorbate reductase* (*DHAR1*) **g** *GLUTATHIONE REDUCTASE* (*GSHR1*). **h** *γ-glutamylcysteine synthetase* (*GSH1*). **i** *Glutathione S-transferase* (*GSTS1*). **j** *Fe superoxide dismutase* (*FSD1*). Experiments were performed in biological replications (*n* = 3); results are presented as normalized fold expression ($2^{-\Delta\Delta Ct}$, WT=1); the error bars represent calculated ±SD. Significant differences were calculated using one-way ANOVA, pair-wise comparison with the Tukey's post-hoc test (non-significant not shown), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.

1149

1150

1151
1152
1153
1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

Figure 10. Model of the chloroplast $^1O_2$-induced retrograde signalling depending on the
metabolite content. Lack of functional Mg-chelatase (MgCh) leads to accumulation of
protoporphyrin IX (Proto), which in light generates $^1O_2$. Lack of functional T6PP1 results in
accumulation of trehalose 6-phosphate (Tre6P), which inhibits SnRK. Blocked SnRK
signalling pathway leads to the inadequate transcriptomic response to stress and disturbed
metabolites content. Fumarate accumulated due to the altered expression of genes involved in
pyruvate metabolism, tyrosine metabolism and the TCA cycle. Accumulation of fumarate and
disturbed TCA cycle lead to deficiency in aconitate. Fumarate attenuate $^1O_2$-signalling
through yet unknown factor (marked with a question mark). The proteinaceous components,
PSBP2, MBS, and SAK1, previously demonstrated to be required for $^1O_2$-signalling are
shown. Aconitate is promoting expression of *PSBP2*, *MBS*, and *SAK1* in *gunSOS1* mutant.
Mitochondria are marked by an asterisk.

1178