1 **MTG16 (CBFA2T3) regulates colonic epithelial differentiation, colitis, and tumorigenesis**
2 **by repressing E protein transcription factors**
3

4 Rachel E. Brown[1,2], Justin Jacobse[3,4,5], Shruti A. Anant[6], Koral M. Blunt[5], Bob Chen[7,8], Paige N.
5 Vega[8,9], Chase T. Jones[5], Jennifer M. Pilat[1], Frank Revetta[3], Aidan H. Gorby[5], Kristy R.
6 Stengel[10], Yash A. Choksi[1,5,11,12], Kimmo Palin[13,14,15], M. Blanca Piazuelo[5], M. Kay Washington[3],
7 Ken S. Lau[8,9,16], Jeremy A. Goettel[1,3,5,12], Scott W. Hiebert[10,16], Sarah P. Short[1,5,12], and
8 Christopher S. Williams[1,2,5,11,12]
9
[1] Program in Cancer Biology, Vanderbilt University School of Medicine, Nashville, Tennessee,
37232, USA
[2] Medical Scientist Training Program, Vanderbilt University School of Medicine, Nashville,
Tennessee, 37232, USA
[3] Department of Pathology, Microbiology, and Immunology, Vanderbilt University, Nashville, TN,
USA
[4] Willem-Alexander Children's Hospital, Leiden University Medical Center, Leiden, The
Netherlands
[5] Department of Medicine, Division of Gastroenterology, Hepatology, and Nutrition, Vanderbilt
University Medical Center, Nashville, Tennessee, 37232, USA
[6] Vanderbilt University, Nashville, Tennessee, 37232, USA
[7] Program in Chemical and Physical Biology, Vanderbilt University School of Medicine,
Nashville, Tennessee, 37232, USA
[8] Epithelial Biology Center, Vanderbilt University Medical Center, Nashville, Tennessee, 37232,
USA
[9] Department of Cell and Developmental Biology, Vanderbilt University School of Medicine,
Nashville, Tennessee, 37232, USA
[10] Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN 37232,
USA
[11] Veterans Affairs Tennessee Valley Health Care System, Nashville, Tennessee, 37232, USA
[12] Center for Mucosal Inflammation and Cancer, Vanderbilt University Medical Center, Nashville,
Tennessee, 37232, USA
[13] Department of Medical and Clinical Genetics, University of Helsinki, Helsinki, Finland
[14] Applied Tumor Genomics Research Program, Research Programs Unit, University of
Helsinki, Helsinki, Finland
[15] iCAN Digital Precision Cancer Medicine Flagship, University of Helsinki, Helsinki, Finland
[16] Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville,
Tennessee, 37232, USA
38
39 **Corresponding author**
40 Christopher S. Williams; Address: 1075 Light Hall/MRBIV, 2215-B Garland Ave., Nashville, TN
41 37232; Tel: 615-322-3642; Fax: 615-343-6229; Email: christopher.s.williams@vumc.org

42
43 **Conflict-of-interest statement**
44 The authors have declared that no conflict of interest exists.

Abstract

Aberrant epithelial differentiation and regeneration contribute to colon pathologies including inflammatory bowel disease (IBD) and colitis-associated cancer (CAC). MTG16 (CBFA2T3) is a transcriptional corepressor expressed in the colonic epithelium. MTG16 deficiency in mice exacerbates colitis and increases tumor burden in CAC, though the underlying mechanisms remain unclear. Here, we identified MTG16 as a central mediator of epithelial differentiation, promoting goblet and restraining enteroendocrine cell development in homeostasis and enabling regeneration following dextran sulfate sodium (DSS)-induced colitis. Transcriptomic analyses implicated increased E box-binding transcription factor (E protein) activity in MTG16-deficient colon crypts. Using a novel mouse model with a point mutation that disrupts MTG16:E protein complex formation ($Mtg16^{P209T}$), we established that MTG16 exerts control over colonic epithelial differentiation and regeneration by repressing E protein-mediated transcription. Mimicking murine colitis, $MTG16$ expression was increased in biopsies from patients with active IBD compared to unaffected controls. Finally, uncoupling MTG16:E protein interactions only partially phenocopied the enhanced tumorigenicity of $Mtg16^{-/-}$ colon in the azoxymethane(AOM)/DSS-induced model of CAC, indicating that MTG16 protects from tumorigenesis through additional mechanisms. Collectively, our results demonstrate that MTG16, via its repression of E protein targets, is a key regulator of cell fate decisions during colon homeostasis, colitis, and cancer.

Graphical Abstract

Introduction

The colonic epithelium is a complex, self-renewing tissue comprised of specialized cell types with diverse functions (1). Stem cells at the base of the colon crypt divide and differentiate into absorptive and secretory cells. Conversely, colonic epithelial regeneration in response to injury occurs through de-differentiation of committed absorptive and secretory lineage cells (2–4). Secretory lineage dysregulation is implicated in inflammatory bowel disease (IBD), diabetes, and even behavioral changes (5–9), and aberrant regenerative programs may lead to uncontrolled cell proliferation and dysplasia (10). Thus, these processes must be tightly controlled by coordinated networks of transcriptional activators and repressors (1).

Myeloid translocation gene 16 (MTG16, also known as CBFA2T3) is a transcriptional corepressor that regulates cell fate decisions by bridging transcription factors and chromatin modifiers to repress transcription of target loci (11–14). Repression targets of MTG16 vary depending on the expression gradients of other components of the repression complex (14, 15). MTG16 represses stem cell genes and pan-secretory genes in the homeostatic small intestine (SI) (16). We have previously shown that MTG16 deficiency leads to increased injury in dextran sulfate sodium (DSS)-induced colitis (17) and tumor burden in azoxymethane (AOM)/DSS-induced inflammatory carcinogenesis and that these were epithelial-specific phenotypes (18). However, the precise mechanisms by which MTG16 controls colonic homeostasis, regeneration, and tumorigenesis remain unknown.

One family of transcription factors repressed by MTG16 are basic helix-loop-helix (bHLH) transcription factors (14, 19–22). Ubiquitously expressed (class I) and tissue-specific (class II) bHLH transcription factors dimerize to control stem cell dynamics, lineage allocation, and differentiation (23–25). Class I bHLH transcription factors are also known as E proteins because they bind to consensus Ephrussi box (E box) DNA sequences and include E12 and

E47 (splice variants of *E2A*), HEB, and E2-2 (22, 23). However, the functions of E proteins have

mainly been described outside of the colon (26–32).

In this study, we defined the topology of *Mtg16* expression in the colon and discovered

new functions of MTG16 in the colonic epithelium. We used a novel MTG16:E protein

uncoupling point mutant mouse model to demonstrate that the mechanism driving these

phenotypes was repression of E protein transcription factors by MTG16. We tested the

functional impact of this regulatory relationship in murine models of IBD and colitis-associated

cancer (CAC) and correlated our observations with patient data. Overall, we demonstrate novel,

context-specific roles for MTG16 in colonic epithelial lineage allocation and protection from

colitis and tumorigenesis that depend on its repression of E protein-mediated transcription.

Results

MTG16 is expressed in goblet cells in human and murine colon. To understand the
topography of _MTG16_ expression in the human colon, we queried single-cell RNA-seq (scRNA-
seq) datasets generated from 2 independent cohorts of normal human colon samples (10).
Interestingly, we observed that _MTG16_ was predominantly expressed in goblet cells (**Fig. 1A,
Fig. S1A**). To understand if this expression pattern was evolutionarily conserved, we queried
our inDrop scRNA-seq generated from WT murine colon crypts (33, 34) and similarly found that
Mtg16 was expressed in goblet and enteroendocrine clusters (**Fig. 1B, Fig. S1B**). RNAscope
for _Mtg16_ and _Muc2_, the predominant mucin expressed by colonic goblet cells (35),
demonstrated strong co-localization in the murine colonic epithelium (**Fig. 1C**). Collectively,
these data suggested a role for MTG16 in the colonic secretory lineage.

Loss of MTG16 distorts colonic secretory cell differentiation. We next assessed secretory
lineage allocation in the homeostatic colon. $Mtg16^{-/-}$ mice had fewer goblet cells but,
interestingly, more synaptophysin-positive (SYP[+]) enteroendocrine cells per colon crypt (**Fig.
1D-E**). Tuft cells were decreased in the colons of $Mtg16^{-/-}$ mice (**Fig. S1C**). In agreement with
prior studies (16, 36), $Mtg16^{-/-}$ mice had fewer goblet cells, but no increase in enteroendocrine
cells, in the SI (**Fig. S2**). Together, these data implicated a colon-specific role for MTG16 in
orchestrating cell type allocation within the secretory lineage.

Prior studies have demonstrated subtle differences in secretory cell differentiation along
the length of the colon(35, 37–40). Thus, we stratified our human scRNA-seq datasets by
proximal and distal colon and examined _MTG16_ expression. _MTG16_ was consistently
expressed in goblet cells, but its expression pattern in other secretory cell types varied (**Fig.
S3**). We next queried the _Tabula Muris_ (41), a murine colon scRNA-seq dataset that specifies
whether certain clusters originate from the proximal or distal colon, for _Mtg16_ expression. In the
proximal colon, _Mtg16_ was enriched in all goblet cell clusters (**Fig. S4A**). In the distal colon,

Mtg16 was enriched in *Lgr5*⁺ amplifying, undifferentiated cells and most goblet cells, except

those located at the top of the crypt (**Fig. S4B**). Bulk RNA-seq of proximal and distal WT colon

revealed higher *Mtg16* expression in the proximal colon, potentially reflecting its higher goblet

cell density (**Fig. S4C**). We confirmed this *Mtg16* expression pattern by RNAscope (**Fig. S5**).

Due to these differences, we performed RNA-seq of colon epithelial isolates from the

proximal and distal colon separately. In both colon segments, goblet-associated genes were

downregulated, and enteroendocrine-associated genes were upregulated compared to WT (**Fig.**

1F-G). However, goblet cell-associated gene downregulation was more pronounced in the

proximal colon (**Fig. 1F**), while enteroendocrine cell-associated gene upregulation was more

striking in the distal colon (**Fig. 1G**). Closer inspection of the distal *Mtg16*⁻/⁻ colon revealed

marked upregulation of class II bHLH transcription factors required for enteroendocrine cell

differentiation (*Neurog3, Neurod1*) and enteroendocrine cell hormone products (*Chga, Gcg, Sct*)

(**Fig. 2A**). Due to this novel phenotype, we chose to further investigate the distal colon using

Gene Set Enrichment Analysis (GSEA) with custom gene sets derived from the literature

(described in **Table S3**). GSEA of distal colon RNA-seq demonstrated significant enrichment of

all stages of enteroendocrine cells during differentiation except the earliest progenitor cells in

which expression of *Neurog3*, which is required for enteroendocrine cell differentiation (42–45),

has not yet been induced (44) (**Fig. 2B**). Concurrently, the goblet cell gene signature was

significantly de-enriched in the absence of MTG16 (**Fig. 2B**).

Notably, *Mtg16*⁻/⁻ vs. WT RNA-seq did not show significant differences in the stem cell

genes *Lgr5* and *Ascl2*, distinct from *Mtg16*⁻/⁻ SI (16). Additionally, GSEA using gene sets

representing stem cells and other early progenitors (**Table S3**) did not demonstrate enrichment

of any gene sets representing *Lgr5*⁺ crypt base columnar stem cells (CBCs) (**Fig. 2C**, **Fig. S6A**).

We did observe enrichment of gene sets representing *Bmi1*⁺ "label-retaining" and *mTert*⁺ stem

cells (**Fig. 2C**), recently shown to be enteroendocrine cells expressing chromogranin A (*Chga*)

(46). While CHGA⁺ cells were increased overall, they were not localized to the crypt base

(including the +4/+5 cell position) (**Fig. 2D**). Altogether, these data indicate that MTG16

promotes goblet cell differentiation, at the expense of enteroendocrine cell differentiation, during

colonic secretory lineage allocation.

MTG16-driven goblet cell differentiation is dependent on its repression of E protein-

mediated transcription. We next sought to illuminate potential mechanisms by which MTG16

promotes the goblet lineage and restricts the enteroendocrine lineage. We and others have

previously shown that MTG16 negatively regulates major transcription factor effectors of the

WNT and Notch pathways in the hematopoietic system (15, 47) and SI (16). We thus performed

GSEA for WNT and Notch activation, as well as other pathways important in colonic epithelial

homeostasis and differentiation (1). Surprisingly, GSEA did not identify global canonical WNT or

Notch pathway upregulation in $Mtg16^{-/-}$ colon crypts (**Fig. S6B**), despite upregulation of the

Notch ligand *Dll3* (**Fig. 1G, Fig. 2A**). Additionally, several WNT pathway inhibitors (*Wif1*, *Cxxc4*)

and the Notch pathway inhibitor *Neurl1a* were significantly upregulated (**Fig. 2A**). Thus, it

appeared unlikely that the effects of MTG16 on colonic differentiation were due to interaction

with, and thus repression of, canonical WNT and Notch effectors.

We then investigated whether target genes of known MTG16-interacting factors,

including E proteins (14, 19–22), were dysregulated in $Mtg16^{-/-}$ colon crypts. As stated

previously, E proteins coordinate lineage-specific differentiation by interacting with cell type-

specific class II bHLH transcription factors (23). Furthermore, E proteins such as E2A and HEB

are known to be expressed in SI crypts and adenomas (48). Here, we observed enrichment of

all E protein gene signatures by GSEA in $Mtg16^{-/-}$ colon crypts (**Fig. 2E**, **Table S3**). Gene sets

associated with upregulation of Id1 and Id2, which counteract E protein activity and increase

stemness in colon cancer (23, 49–52), were not enriched (**Fig. 2E**). These data demonstrated a

strong induction of E protein-mediated transcription in the absence of repression by MTG16.

Thus, we hypothesized that MTG16 drives goblet cell differentiation from secretory progenitors

by inhibiting E protein-mediated transcription of key enteroendocrine transcription factors.

We next tested this hypothesis by selectively disrupting the MTG16:E protein interface *in*

vivo. To do so, we leveraged a CRISPR-generated mouse with a point mutation resulting in a

proline-to-threonine substitution in MTG16 (*Mtg16*[P209T]) that impairs MTG16:E2A and

MTG16:HEB binding and, consequently, MTG16-mediated repression of E protein

transcriptional activity (20). Similar to *Mtg16*[-/-] mice, homozygous *Mtg16*[P209T] mutant (*Mtg16*[T/T])

mice had fewer periodic acid-Schiff-positive (PAS[+]) goblet cells and more CHGA[+]

enteroendocrine cells per crypt compared to littermate-matched WT controls, though tuft cell

frequency per crypt was unchanged (**Fig. 3A-C**). We next performed bulk RNA-seq on proximal

and distal colon crypt isolates. In both colon segments, goblet cell genes were downregulated

compared to WT, while enteroendocrine cell genes were upregulated compared to WT, in a

pattern resembling that of *Mtg16*[-/-] colon (**Fig. 3D-E**). Notably, the distal colon exhibited

significant upregulation of *Neurog3* (**Fig. 3E**). Additionally, GSEA of the overall enteroendocrine

signature, key enteroendocrine progenitor genes, *Bmi1*[+] and *mTert*[+] stem cell signatures, and E

protein signatures all largely phenocopied the *Mtg16*[-/-] distal colon (**Fig. 3F-H, Table S3**). As

expected, the gene sets representing E2A and HEB transcriptional targets were also enriched

(**Fig. 3H**).

We next compared RNA-seq results from *Mtg16*[-/-] vs. WT and *Mtg16*[T/T] vs. WT distal

colon crypts and observed 22 genes upregulated in both datasets (**Fig. 4A, Table 1**). Many of

these genes (e.g. *Neurog3, Insm1, Insrr, Pex5l, Syt13, Syngap1, Kcnh6, and Agt*) were

enteroendocrine-associated (**Table 1**). A publicly available MTG16 ChIP-seq dataset from SI

crypts (16) exhibited significant MTG16 occupancy proximal to 11 of the 22 genes (**Table 1**),

including a large peak at an E box-rich site upstream of *Neurog3* (**Fig. 4B**), suggesting that

these genes are repressed by MTG16. Finally, *Neurog3*[+] enteroendocrine progenitor cells were

markedly increased in *Mtg16*[-/-] and *Mtg16*[T/T] colon crypts by RNAscope (**Fig. 4C-D**). Altogether,

we discovered an MTG16:E protein-dependent mechanism of colonic secretory lineage

allocation, specifically inhibition of NEUROG3-mediated commitment to the enteroendocrine cell

lineage.

***MTG16* is upregulated in IBD patients with active disease.** There has been considerable

interest in the role of secretory cell lineage misallocation and dysfunction in the pathogenesis of

IBD, which includes ulcerative colitis (UC) and Crohn's disease and is defined by continuous

epithelial injury and regeneration (8, 35, 53–57). Additionally, several large RNA-seq studies of

human UC biopsies have recently been published, including the pediatric PROTECT study (58,

59) and mucosal biopsies from adult UC patients with active disease, UC patients in remission,

and healthy control patients (60). We found that *MTG16* was upregulated in UC patients in both

datasets (**Fig. 5A-B**). *MTG16* expression decreased with remission, although remained

elevated compared to unaffected control patients (**Fig. 5B**). Lastly, a microarray dataset from

pediatric colon biopsies (61) demonstrated upregulation of *MTG16* in both UC and Crohn's

colitis (**Fig. 5C**). Collectively, these data show that increased *MTG16* expression is correlated

with active injury and regeneration in IBD.

MTG16 contributes to colon crypt regeneration following DSS-induced colitis. As stated

previously, MTG16 deficiency increases disease severity in DSS-induced colitis (17), a mouse

model of IBD (schematic in **Fig. 5D**) that recapitulates histologic features of human UC (62). We

first assessed *Mtg16* expression in DSS-induced colitis to determine whether it would

recapitulate the increased *MTG16* expression in human IBD. Because DSS treatment largely

affects the distal colon (62), we compared distal WT colon crypt epithelial isolates post-DSS-

induced injury and regeneration to WT distal colon crypts at baseline. Indeed, *Mtg16* expression

was significantly increased in regenerating colon crypts (**Fig. 5E**), similar to human IBD. Thus,

we posited that MTG16 expression may be critical for epithelial regeneration in colitis.

The cellular sources of intestinal repair have been a major focus of the field over the last

several decades. Recently, Murata *et al.* demonstrated that any CBC progeny cell can de-

differentiate and repopulate ablated CBCs by inducing expression of *Ascl2* (2). After CBC

ablation, 238 significantly upregulated genes defined the de-differentiating ASCL2$^+$ cells

compared to resting CBCs. Interestingly, *Mtg16* was one of these few significantly upregulated

genes (log$_2$[fold change] = 2.05, p_{adj} = 0.004) (2). Furthermore, ASCL2 ChIP-seq data in the de-

differentiating population (2) yielded a peak at a second transcription start site of *Mtg16*,

suggesting that ASCL2 induces *Mtg16* expression during colonic epithelial regeneration (**Fig.**

5F).

To confirm that the ASCL2-driven regeneration program was relevant to crypt

regeneration following DSS-induced injury, we constructed a gene set consisting of the 238

genes upregulated in *Ascl2*$^+$ regenerating cells (2) (**Table S3**), performed GSEA, and observed

significant enrichment of this signature in WT distal colon crypts after DSS-induced regeneration

(**Fig. 5G**). Conversely, this regeneration signature was significantly de-enriched in *Mtg16*$^{-/-}$ colon

crypts following DSS-induced injury, despite enrichment of other gene signatures associated

with intestinal epithelial regeneration (**Fig. 5H-I**, **Table S3**). Altogether, these data implicate

MTG16 as a central component of ASCL2-driven regeneration and repair in the colon. Similar to

our findings in colon homeostasis, E protein transcriptional signatures were enriched in post-

DSS *Mtg16*$^{-/-}$ colon by GSEA (**Fig. 5J-K**), leading us to hypothesize that the contribution of

MTG16 to colon crypt regeneration occurs via repression of E protein-mediated transcription.

MTG16-mediated colonic epithelial regeneration is dependent on its repression of E

proteins. To test whether MTG16-dependent regeneration occurs via an E protein-dependent

mechanism, we next analyzed colonic epithelial regeneration following DSS-induced injury (**Fig.**

6A) in *Mtg16*$^{T/T}$ vs. WT mice. Like *Mtg16*$^{-/-}$ mice, *Mtg16*$^{T/T}$ mice developed worse colitis and poor

regeneration measured by increased weight loss (**Fig. 6B**), decreased colon length (**Fig. 6C**),

and increased histologic injury-regeneration score (**Fig. 6D, Table S2**) compared to WT.

Mtg16$^{T/T}$ mice also exhibited increased extent of injury along the length of the *Mtg16*$^{T/T}$ colon

(**Fig. 6E**). Next, we injected mice with 5-Ethynyl-2'-deoxyuridine (EdU) 1 h prior to sacrifice

following DSS injury and regeneration and found that *Mtg16*$^{T/T}$ mice had fewer S-phase cells in

injury-adjacent crypts (**Fig. 6F**), indicating a proliferation defect in the crypts that actively

regenerate the ulcerated colonic epithelium following DSS-induced colitis (63). Finally, like

Mtg16$^{-/-}$ mice, RNA-seq of *Mtg16*$^{T/T}$ distal colon crypt epithelial isolates following DSS injury and

regeneration was de-enriched for ASCL2-mediated de-differentiation despite enrichment of

other gene signatures associated with regeneration (**Fig. 6G-H, Table S3**). These data showed

that MTG16 repression of E protein-mediated transcription is required for ASCL2-mediated

colonic epithelial regeneration.

***MTG16* expression is decreased in dysplasia.** Because *MTG16* expression was upregulated

in IBD, we wondered whether *MTG16* expression would be increased in IBD-driven colon

cancer (CAC) compared to sporadic CRC. We previously observed decreased *MTG16*

expression in sporadic CRC and CAC compared to normal colon tissue (18), but had not directly

compared expression levels of *MTG16* between these different neoplastic origins. To do so, we

first confirmed our previous findings in sporadic CRC using DESeq2 on raw counts generated

by Rahman *et al.* (64) from The Cancer Genome Atlas (TCGA) (65) (**Fig. 7A**). We then

analyzed *MTG16* expression in bulk RNA-seq of CRC and CAC tumors (66) and found no

significant difference in *MTG16* expression (**Fig. 7B**), indicating that *MTG16* expression is

reduced in colon dysplasia regardless of whether it is driven by IBD.

MTG16-mediated protection from CAC is partially dependent on its repression of E

protein transcription factors. We previously showed that *Mtg16*$^{-/-}$ mice have increased

mortality, tumor burden, tumor grade, and tumor cell proliferation and apoptosis in AOM/DSS-

induced CAC (18). We next tested whether the mechanism driving these phenotypes was E

protein-dependent by treating *Mtg16$^{T/T}$* mice with AOM/DSS (**Fig. 7C**). Llike *Mtg16$^{-/-}$* mice,

Mtg16$^{T/T}$ mice did exhibit increased histologic injury, tumor multiplicity, and tumor size compared

to WT (**Fig. 7D-G**). Interestingly, *Mtg16$^{T/T}$* mice had more tumors and larger tumors in the

middle and proximal colon than their WT counterparts (**Fig. 7L-K**). This was consistent with a

greater extent of DSS-induced injury (**Fig. 6E**). These data suggest that this phenotype could be

driven by increased injury in the proximal colon.

However, unlike *Mtg16$^{-/-}$* mice, *Mtg16$^{T/T}$* mice did not have significantly more weight loss

(**Fig. 7J**), colon shortening (**Fig. 7K**), mortality (**Fig. 7L**), or tumor dysplasia (**Fig. 7M**) compared

to WT. Additionally, further characterization of *Mtg16$^{T/T}$* tumors (**Fig. S7**) demonstrated no

difference in proliferation (**Fig. S7A**). We did, however, observe a subtle increase in apoptotic

epithelial cells per tumor (**Fig. S7B**), similar to our prior findings in *Mtg16$^{-/-}$* tumors. These

phenotypes were recapitulated in tumoroids derived from distal, but not proximal, WT and

Mtg16$^{T/T}$ tumors (**Fig. S8**). Overall, these data indicate that uncoupling MTG16:E protein

complexes recapitulates some, but not all, *Mtg16$^{-/-}$* phenotypes, suggesting that MTG16 protects

from tumorigenesis through additional mechanisms.

Discussion

In this study, we investigated the mechanism underlying context-specific functions of the

transcriptional corepressor MTG16. We identify previously unappreciated roles for MTG16 as a

critical regulator of colonic secretory cell differentiation and active colon regeneration. To define

the mechanism underlying these phenotypes, we utilized a novel mouse model with a single

point mutation in MTG16, $Mtg16^{P209T}$, that disrupts MTG16 binding to the E proteins E2A and

HEB(20). $Mtg16^{T/T}$ mice homozygous for this MTG16:E protein uncoupling mutation largely

recapitulated $Mtg16^{-/-}$ lineage allocation and regeneration phenotypes, although the effect of

MTG16 loss in dysplasia appears to only partially be due to increased E protein activity. Thus,

these data indicate a new role for MTG16:E protein complexes in the colonic epithelium.

First, we demonstrated a novel role for MTG16 in colonic epithelial homeostasis. MTG16

has been shown to repress stem, goblet, and enteroendocrine cell genes to drive enterocyte

differentiation in the SI (16). However, in accordance with numerous studies demonstrating

important differences between SI and colon biology (2, 35, 67–70), $Mtg16^{-/-}$ colon did not exhibit

increased stem cell transcriptional signatures or an overall decrease in secretory genes.

Instead, we observed specific upregulation of the enteroendocrine lineage, occurring at least in

part due to unrepressed expression of $Neurog3$, the class II bHLH transcription factor required

for enteroendocrine cell differentiation (42–45). This could be due to the fact that MTG16

repression targets, and its effects on them, vary in different tissues and cell types due to

expression gradients of its binding partners (14). Another potential explanation for the difference

between SI and colonic phenotypes is that $Neurog3^{+}$ progenitor cells in the SI may still

differentiate into goblet and Paneth cells but are fully committed to the enteroendocrine lineage

in the colon (45, 68).

Enteroendocrine cell lineage commitment is dependent on a cascade of type II bHLH

transcription factors including NEUROG3 and its downstream target NEUROD1 (71). E proteins

are known to induce transcription of *Neurog1-3* in neurogenesis (30, 32), and NEUROD1

heterodimerizes with E proteins in other systems (72). Although E proteins have been detected

in SI stem cells and adenomas (48), E protein function in colon homeostasis remained

unknown. Here, we mechanistically probed the extent to which MTG16:E protein interactions

drive MTG16-associated lineage allocation functions in the colon using a novel mouse model

with a single amino acid substitution in *Mtg16* (*Mtg16*[P209T]) (20). Strikingly, *Mtg16*[T/T] mice

recapitulated secretory cell phenotypes of *Mtg16*[-/-] mice including decreased goblet cells,

increased enteroendocrine cells, and increased *Neurog3* tone. One potential non-E protein-

mediated pathway that was increased (FDR q-value = 0.11) in *Mtg16*[-/-], but not *Mtg16*[T/T] colon

crypts, is the non-canonical, β-catenin-independent WNT-planar cell polarity (PCP) pathway.

This could be responsible for the slight difference in magnitude between *Mtg16*[-/-] and *Mtg16*[T/T]

phenotypes, as the WNT-PCP pathway may enable enteroendocrine cells to differentiate

directly from "unipotentially-primed" intestinal stem cells[99].

An area of considerable interest in gastrointestinal biology is the relationship between

epithelial differentiation and regeneration (4). Although it was long thought that dedicated *Bmi1*[+]

or *mTert*[+] "reserve" stem cells at the +4/+5 position replenished the stem cell compartment in

intestinal epithelial regeneration, recent data indicate that these cells are enteroendocrine cells

that can de-differentiate into stem cells following injury (46). Recently, "committed" progenitors

from both the secretory and absorptive lineages were demonstrated to share this ability through

induction of ASCL2 (2). In this study, *Ascl2*[+] de-differentiating cells were isolated from the

colonic epithelium following CBC ablation (2). Interestingly, these cells were not enriched for *Clu*

or fetal epithelial genes (2), previously thought to enhance intestinal epithelial response to injury

(4, 73, 74). *Mtg16* was one of the few genes upregulated in these regenerating cells, and ChIP-

seq demonstrated an ASCL2 peak at a second TSS in the *Mtg16* locus (2). Demonstrating

relevance to DSS-induced colon regeneration, both *Mtg16* and the *Ascl2*[+] regenerating cell

signature were upregulated in regenerating WT colon epithelium compared to baseline. *MTG16*

was also upregulated in multiple IBD patient cohorts and even decreased with remission, during

which the colon epithelium is closer to homeostatic renewal. Finally, RNA-seq and GSEA of

$Mtg16^{-/-}$ and $Mtg16^{T/T}$ colonic epithelial isolates following DSS-induced injury demonstrated

enrichment of "revival" stem cell signatures and fetal epithelial signatures, but de-enrichment of

the $Ascl2^+$ regenerating cell signature. These data indicate an active, E protein-mediated role for

MTG16 in colon regeneration and provide a potential explanation for defective colonic epithelial

regeneration in $Mtg16^{-/-}$ mice despite an apparent baseline enrichment of $Bmi1^+$ and $mTert^+$

"reserve" stem cells. Future work is necessary to determine the E protein targets repressed by

MTG16 in $Ascl2^+$ de-differentiating cells. One possibility is that MTG16 represses $Neurog3$ in

these de-differentiating cells to prevent them from reversing course and differentiating toward

the enteroendocrine lineage.

Finally, we investigated the role of MTG16:E protein complexes in inflammatory

carcinogenesis. We previously showed that $Mtg16^{-/-}$ mice develop greater tumor burden than

their WT counterparts with AOM/DSS treatment (18). Although E proteins maintain lineage

commitment in homeostasis, they have been shown to be both anti-tumorigenic and pro-

tumorigenic in colon cancer (75, 76). Although we did observe increased histologic injury, tumor

burden, and tumor size in $Mtg16^{T/T}$ mice, $Mtg16^{T/T}$ tumors did not recapitulate the much higher

mortality or grade of dysplasia observed in $Mtg16^{-/-}$ tumors (18). Thus, disabling one known

function of MTG16 was not sufficient to fully replicate the $Mtg16^{-/-}$ phenotype. Indeed, our group

recently showed that MTG16-dependent effects on CAC development is dependent on Kaiso

(ZBTB33) (77). Future studies are necessary to determine additional MTG16 binding partners

and repression targets responsible for suppressing carcinogenesis.

In conclusion, we discovered key new functions of MTG16 in colonic secretory lineage

allocation, regeneration following colitis, and colitis-associated tumorigenesis, dependent on its

repression of E proteins. Confirming translational relevance, we determined that $MTG16$ is

upregulated in patients with active IBD, reduced with restitution, and decreased in dysplasia.

Thus, *MTG16* may be a candidate biomarker for disease activity or a target to modulate

differentiation and regeneration.

Methods

Animal models. *Mtg16⁻/⁻* mice were previously generated and validated (78). *Mtg16^{P209T}* mutant mice were CRISPR-generated as recently described (20). Both models are on a predominantly C57BL/6 background. Littermate-matched WT and *Mtg16⁻/⁻* and WT and homozygous *Mtg16^{P209T}* mutant (*Mtg16^{T/T}*) mice were bred using heterozygous x heterozygous breeding schemes, housed in the same facility with a 12:12 light-dark cycle, provided with standard rodent chow (#5L0D, LabDiet) *ad libitum*. Approximately equal numbers of 8-12-wk-old male and female mice were used.

scRNA-seq. InDrop scRNA-seq and analysis were performed on 2 independent (DIS, discovery and VAL, validation) cohorts of human colon samples as previously described (10). These data are publicly available through the Human Tumor Atlas Network (HTAN) (https://data.humantumoratlas.org/) (79). scRNA-seq of colon crypts isolated from WT mice was performed as previously described (33, 34, 80). These data (#GSE145830, #GSE114044) are available in the Gene Expression Omnibus (GEO) (81, 82).

RNAscope. RNAscope, high-resolution RNA *in situ* hybridization (83), was performed on FFPE sections using the RNAscope Multiplex Fluorescent V2 Assay (#323100, ACDBio). Slides were boiled in Target Retrieval Reagents for 15 min and treated with Protease IV for 30 min. Probes were specific for Mm-*Cbfa2t3* (#43601-C2, ACDBio), Mm-*Muc2* (#315451, ACDBio), and Mm-*Neurog3* (#422401, ACDBio). 1:750 TSA Cy3 (#NEL704A001KT, PerkinElmer) and TSA Cy5 (#NEL75A001KT, Perkin Elmer) were used for probe visualization.

Chromogenic and immunofluorescent IHC. SI and colon were dissected, Swiss-rolled, and fixed in 10% neutral-buffered formalin (NBF) for 24 h at RT. formalin (NBF) for 24 h at RT. The Vanderbilt Translational Pathology Shared Resource (TPSR) paraffin-embedded the tissue and performed H&E and PAS staining. Chromogenic and immunofluorescent IHC were performed

as previously described (77, 84). Briefly, 5-µm FFPE sections were deparaffinized, rehydrated in

graded ethanol concentrations, and permeabilized using TBS-T (Tris-buffered saline with 0.05%

Tween-20) followed by antibody-specific antigen retrieval (**Table S1**). Next, for chromogenic

IHC, endogenous peroxidases were quenched in 0.03% H_2O_2 with sodium azide for 5 min.

Slides were then incubated in primary antibody (**Table S1**) O/N at RT. Detection was performed

by 30 min. incubation with Dako Envision+ System HRP-Labeled Polymers (#K4003, Agilent)

followed by a 5-min incubation with DAB. For immunofluorescent staining, slides were blocked

in 5% normal goat serum (#01-620-1, ThermoFisher), incubated in primary antibody (**Table S1**)

O/N at 4 °C, washed in TBS-T, and incubated in the appropriate secondary antibodies (#A-

11011, #A-11077, #A-21131, Invitrogen) for 2 h at RT. EdU was visualized using the Click-iT

EdU Cell Proliferation Kit for Imaging (#C10037, ThermoFisher). Nuclear counterstaining was

performed using ProLong Gold Antifade Mountant with DAPI (#P36931, ThermoFisher).

Slide imaging and quantification. Slides were imaged using a Nikon Eclipse E800 microscope

with NIS-Elements Basic Research Software or by the Vanderbilt University Medical Center

Digital Histology Shared Resource (DHSR). For chromogenic IHC, whole slides were imaged by

the DHSR using a Leica SCN400 Slide Scanner (Leica Biosystems). For fluorescent staining,

whole slides were imaged by the DHSR using an Aperio Versa 200 automated slide scanner

(Leica Biosystems) and visualized using Aperio ImageScope (v12.4.3) (Leica Biosystems).

Quantification was performed by an observer blinded to genotype by counting positive cells in

well-oriented crypts or crypt-villus units.

Colon crypt isolation. Murine colon was harvested and splayed longitudinally. The proximal (5

cm from cecum) and distal (5 cm from anus) colon were washed in PBS without calcium or

magnesium and minced. Tissue fragments were incubated in chelation buffer (2 mM EDTA in

PBS without calcium or magnesium) with nutation for 90 min at 4 °C and washed with PBS

followed by 2-min cycles of gentle shaking in shaking buffer (43.3 mM sucrose and 54.9 mM

sorbitol in PBS without calcium or magnesium) to free colon crypts. Intact crypts were collected

by centrifugation at 150 $x\ g$, 4 °C for 12 min for RNA isolation and bulk RNA-seq as described

below.

DSS-induced colitis. Bedding was mixed between cages to normalize microbiome 2 wk prior to

experiments. Water deprivation caps were placed to accustom mice to drinking from water

bottles 1 wk prior to experiments. Water bottles were filled with 2% (w/v) DSS (#DS1004, Gojira

Fine Chemicals) for 5 d and then replaced with water for 4 d of recovery prior to sacrifice. At

sacrifice, colons were dissected, Swiss-rolled, fixed in 10% NBF, and processed by the

Vanderbilt TPSR. H&Es were scored for histologic injury and regeneration by a gastrointestinal

pathologist blinded to genotype (MKW) (scoring system described in **Table S2**). Distal colon

crypts were collected as described above for RNA isolation and RNA-seq as described below.

For EdU experiments, mice were intraperitoneally injected with 50 mg/kg EdU (#NE08701,

Biosynth Carbosynth) in PBS 1 h prior to sacrifice. Mice were weighed daily throughout all

experiments.

Inflammatory carcinogenesis. Mice were prepared as described above and injected *i.p.* with

10 mg/kg AOM followed by three 4-day cycles of 3% (w/v) DSS (#DB001, TdB Labs), each

followed by 16 d of recovery. Mice were weighed daily. At sacrifice, colons were dissected,

flushed with PBS, splayed longitudinally, and imaged using a Nikon SMZ1270 dissecting scope.

Tumors were identified by vascular, non-dimpled appearance from these images in conjunction

with direct visualization and measured macroscopically in 2 perpendicular dimensions using

calipers. Tumor volume was calculated using the previously validated formula $W^2{*}L/2$, where

width (W) was the shorter caliper measurement, and length (L) was the longer caliper

measurement (85). Tumoroids were generated from a subset of tumors (see **Supplemental**

Methods). Histologic injury and regeneration were assessed by MKW (scoring system in **Table**

S2). Grade of dysplasia was evaluated as previously described (86) by a gastrointestinal

pathologist blinded to genotype (MBP).

RNA isolation, bulk RNA-seq, and analysis. Murine colon crypts were collected as described

above followed by resuspension in TRIzol Reagent (#15596018, Invitrogen) and

homogenization by 5x shearing through a 25G needle. Murine tumors were mechanically

homogenized in TRIzol using an electronic homogenizer. Samples were centrifuged for 10 min

at 16,200 x *g*, 4 °C to remove insoluble material. Cleared homogenate was chloroform-

extracted. RNA was isolated from the aqueous layer using the RNeasy Mini Kit (#74106,

Qiagen) with on-column DNase treatment (#79254, Qiagen). Library preparation and paired-end

150 bp RNA-seq using the Illumina NovaSeq6000 were performed by Vanderbilt Technologies

for Advanced Genomics. Raw reads in FASTQ format were trimmed with fastp (v0.20.0) with

default parameters (87). Quantification was performed using Salmon (v1.3.0) (88) against a

decoy transcriptome generated from *Mus musculus* GENCODE (v21) (89). Further analysis was

performed in R (v3.6.3) as described previously (90). Briefly, quantification files were imported

with tximeta (v1.4.5) (91). Genes with counts ≤ 1 were omitted. Differential expression analysis

(DEA) was performed on raw transcript counts using DESeq2 (v1.26.0) (92) and annotated with

AnnotationDbi (v1.46.1) (93). GSEA (94) was performed on DESeq2-normalized transcript

counts using the GSEA software (v4.1.0) with 1,000 gene permutations, gene set

randomization, and gene sets found in the Molecular Signatures Database (MSigDB) (v7.2) (95)

or derived from the literature (described in **Table S3**). RNA-seq of human CRC and CAC

samples collected from 9 regional hospitals in Finland was performed and analyzed as

described previously (96) (**Supplemental Methods**).

In silico **analyses of publicly available datasets.** Raw and processed ChIP-seq data from

Baulies *et al.* (16) (#GSE124186) and Murata *et al.* (2) (#GSE130822) were retrieved from the

GEO (81). Data were visualized and E box (5'-CATNNG) sequences were annotated using the

Broad Institute Integrated Genomics Viewer (v2.9.1) (97). Metadata and SRR files from the

PROTECT study (58, 59) (#GSE109142) were downloaded using the NCBI SRA Toolkit and

processed as described above for murine samples, with reads quantified against a decoy-aware

transcriptome generated from the GENCODE human reference genome (v29) (89). DEA was

performed on raw counts using DESeq2 (v1.26.0) (92). Pairwise DEA was also performed on

raw counts from RNA-seq of colon biopsies from adults with UC (60) (#GSE128682) and raw

counts generated from the TCGA (65) by Rahman *et al*. (64) (#GSE62944) using DESeq2

(v1.33.1) (92). Microarray data from IBD patients (61) (#GSE9686) were analyzed using limma

in Geo2R (82). The *Tabula Muris* (41) was queried for *Mtg16* expression using their web

interface (https://tabula-muris.ds.czbiohub.org/).

Statistics. Statistical analyses of human CAC and sporadic colorectal cancer (CRC) samples

were performed using limma (v3.34.9) (98). Statistical analyses of all other bulk RNA-seq were

performed using DESeq2 (92). Statistical analyses of microarray data were performed using

limma in Geo2R (82). $p_{adj} < 0.05$ was considered significant. Statistical analyses for GSEA were

performed using the GSEA program (v4.1.0). FDR (q-value) < 0.05 was considered significant.

All other statistical analyses were performed in GraphPad Prism (v9.0.1) as indicated in each

figure legend. Data are displayed as arithmetic mean ± SEM unless otherwise noted. $p < 0.05$

was considered significant.

Study approval. All animal experiments were carried out in accordance with protocols

approved by the Vanderbilt IACUC. Human CRC and CAC studies (66) were approved by the

Ethics Committee of the Hospital District of Helsinki and the Finnish Institute for Health and

Welfare.

Author Contributions

REB designed experiments, performed experiments, and analyzed data. SAA, KMK, CTJ, JMP, and AHG performed experiments and analyzed data. JJ and REB designed and performed bulk RNA-seq data analysis, respectively. KSL, BC, and PNV designed, collected, and analyzed human and murine colon single cell RNA-seq. FR validated antibodies for and performed immunohistochemical staining. MBP and MKW performed histological analyses and provided expertise in gastrointestinal pathology. JJ, RB, and KP analyzed RNA-seq from human tumors. CSW, SPS, SWH, KSL, JAG, KRS, and YAC provided expertise regarding experimental models and experimental design. REB wrote the manuscript. All authors edited and approved the manuscript.

Acknowledgments

This work was supported by the National Institutes of Health (F30DK120149 to REB, F31DK127687 to PNV, F31CA232272 to JMP, R01DK103831 to KSL, U01CA215798 to KSL, R03DK123489 to JAG, R01CA178030 to SWH, K01DK123495 to SPS, F32DK108492 to SPS, R01DK099204 to CSW, P30DK058404 to the Vanderbilt Digestive Disease Research Center [DDRC], P50CA236733 to the Vanderbilt-Ingram Cancer Center [VICC] Spore in Gastrointestinal Cancer, P30CA068485 and U24DK059637 to the Vanderbilt Translational Pathology Shared Resource [TPSR], and T32GM00734 to the Vanderbilt Medical Scientist Training Program), the U.S. Department of Veterans Affairs Office of Medical Research (1I01BX001426 to CSW and IK2BX004648 to YAC), and the Crohn's and Colitis Foundation (623541 to CSW and 662877 to SPS). JJ was supported by the Royal Netherlands Academy of Arts and Sciences (Academy Ter Meulen Grant) and the Prince Bernhard Cultural Foundation (Cultural Foundation Grant). KP was supported by grants from Academy of Finland (Finnish Center of Excellence Program 2018–2025, 312041), the iCAN Digital Precision Cancer Medicine Flagship (320185), and the Sigrid Jusélius Foundation.

We thank Barbara Fingleton, Stephen Brandt, Matthew Stier, and other members of the Williams and Hiebert Labs for their critical insights into this manuscript. We would also like to thank the TPSR, Digital Histology Shared Resource (DHSR), Vanderbilt Technologies for Advanced Genomics (VANTAGE), and Advanced Computing Center for Research and Education (ACCRE) for aid with histology, imaging, RNA-seq, and data storage/processing, respectively. Additional computational resources were provided by the ELIXIR node, hosted at the CSC–IT Center for Science, Finland.

References

1. Beumer J, Clevers H. Cell fate specification and differentiation in the adult mammalian
intestine. *Nat Rev Mol Cell Bio* 2021;22(1):39–53.

2. Murata K et al. Ascl2-Dependent Cell Dedifferentiation Drives Regeneration of Ablated
Intestinal Stem Cells. *Cell Stem Cell* 2020;26(3):377-390.e6.

3. Jadhav U et al. Dynamic Reorganization of Chromatin Accessibility Signatures during
Dedifferentiation of Secretory Precursors into Lgr5+ Intestinal Stem Cells. *Cell Stem Cell*
2017;21(1):65-77.e5.

4. Liu Y, Chen Y-G. Intestinal epithelial plasticity and regeneration via cell dedifferentiation. *Cell*
Regen 2020;9(1):14.

5. Worthington JJ, Reimann F, Gribble FM. Enteroendocrine cells-sensory sentinels of the
intestinal environment and orchestrators of mucosal immunity. *Mucosal Immunol* 2018;11(1):3–
20.

6. Skibicka KP, Dickson SL. Enteroendocrine hormones—central effects on behavior. *Curr Opin*
Pharmacol 2013;13(6):977–982.

7. Yamane S, Inagaki N. Control of intestinal stem cell fate: A novel approach to treating
diabetes. *J Diabetes Invest* 2016;7(2):166–168.

8. Parikh K et al. Colonic epithelial cell diversity in health and inflammatory bowel disease.
Nature 2019;567(7746):49–55.

9. Aliluev A et al. Diet-induced alteration of intestinal stem cell function underlies obesity and
prediabetes in mice. *Nat Metabolism* 2021;3(9):1202–1216.

10. Chen B et al. Human colorectal pre-cancer atlas identifies distinct molecular programs
underlying two major subclasses of pre-malignant tumors. *Biorxiv* 2021;2021.01.11.426044.

11. Gelmetti V et al. Aberrant recruitment of the nuclear receptor corepressor-histone
deacetylase complex by the acute myeloid leukemia fusion partner ETO. *Mol Cell Biol*
1998;18(12):7185--91.

12. Cai Y et al. Eto2/MTG16 and MTGR1 are heteromeric corepressors of the TAL1/SCL
transcription factor in murine erythroid progenitors. *Biochem Bioph Res Co* 2009;390(2):295--
301.

13. Wang J, Hoshino T, Redner RL, Kajigaya S, Liu JM. ETO, fusion partner in t(8;21) acute
myeloid leukemia, represses transcription by interaction with the human N-CoR/mSin3/HDAC1
complex. *Proc National Acad Sci* 1998;95(18):10860--10865.

14. Steinauer N, Guo C, Zhang J. Emerging Roles of MTG16 in Cell-Fate Control of
Hematopoietic Stem Cells and Cancer. *Stem Cells Int* 2017;2017:1--12.

15. Moore AC et al. Myeloid Translocation Gene Family Members Associate with T-Cell Factors
(TCFs) and Influence TCF-Dependent Transcription. *Mol Cell Biol* 2007;28(3):977–987.

16. Baulies A et al. The Transcription co-Repressors MTG8 and MTG16 Regulate Exit of
Intestinal Stem Cells From Their Niche and Differentiation into Enterocyte vs Secretory
Lineages. *Gastroenterology* 2020;159(4):1328-1341.e3.

17. Williams CS et al. MTG16 contributes to colonic epithelial integrity in experimental colitis..
Gut 2013;62(10):1446--55.

18. McDonough EM et al. MTG16 is a tumor suppressor in colitis-associated carcinoma.. *Jci*
Insight 2017;2(16). doi:10.1172/jci.insight.78210

19. Hunt A, Fischer M, Engel ME, Hiebert SW. Mtg16/Eto2 contributes to murine T-cell
development. *Mol Cell Biol* 2011;31(13):2544--51.

20. Acharya P et al. Mtg16-dependent repression of E protein activity is required for early
lymphopoiesis. *Biorxiv* 2020;2020.07.24.220525.

21. Ghosh HS et al. ETO family protein Mtg16 regulates the balance of dendritic cell subsets by
repressing Id2.. *J Exp Medicine* 2014;211(8):1623--35.

22. Ghosh HS, Liu K, Hiebert S, Reizis B. Eto2/MTG16 Regulates E-Protein Activity and Subset
Specification in Dendritic Cell Development. *Blood* 2012;120(21):1229–1229.

23. Wang L-H, Baker NE. E Proteins and ID Proteins: Helix-Loop-Helix Partners in Development
and Disease. *Dev Cell* 2015;35(3):269--280.

24. Slattery C, Ryan MP, McMorrow T. E2A proteins: Regulators of cell phenotype in normal
physiology and disease. *Int J Biochem Cell Biology* 2008;40(8):1431–1436.

25. Peng V et al. E proteins orchestrate dynamic transcriptional cascades implicated in the
suppression of the differentiation of group 2 innate lymphoid cells. *J Biol Chem*
2020;295(44):14866–14877.

26. Zhuang Y, Cheng P, Weintraub H. B-lymphocyte development is regulated by the combined
dosage of three basic helix-loop-helix genes, E2A, E2-2, and HEB.. *Mol Cell Biol*
1996;16(6):2898–2905.

27. Cisse B et al. Transcription Factor E2-2 Is an Essential and Specific Regulator of
Plasmacytoid Dendritic Cell Development. *Cell* 2008;135(1):37–48.

28. Beck K, Peak MM, Ota T, Nemazee D, Murre C. Distinct roles for E12 and E47 in B cell
specification and the sequential rearrangement of immunoglobulin light chain loci. *J Exp Med*
2009;206(10):2271–2284.

29. Bhattacharya A, Baker NE. A Network of Broadly Expressed HLH Genes Regulates Tissue-
Specific Cell Fates. *Cell* 2011;147(4):881–892.

30. Dréau GL et al. E proteins sharpen neurogenesis by modulating proneural bHLH
transcription factors' activity in an E-box-dependent manner. *Elife* 2018;7:e37267.

31. Bouderlique T et al. The Concerted Action of E2-2 and HEB Is Critical for Early Lymphoid
Specification. *Front Immunol* 2019;10:455.

32. Rao C, Malaguti M, Mason JO, Lowell S. The transcription factor E2A drives neural
differentiation in pluripotent cells. *Development* 2020;147(12):dev.184093.

33. Banerjee A et al. Succinate Produced by Intestinal Microbes Promotes Specification of Tuft
Cells to Suppress Ileal Inflammation. *Gastroenterology* 2020;159(6):2101-2115.e5.

34. Herring CA et al. Unsupervised Trajectory Analysis of Single-Cell RNA-Seq and Imaging
Data Reveals Alternative Tuft Cell Origins in the Gut. *Cell Syst* 2018;6(1):37-51.e9.

35. Nyström EEL et al. An intercrypt subpopulation of goblet cells is essential for colonic mucus
barrier function. *Science* 2021;372(6539):eabb1590.

36. Poindexter SV et al. Transcriptional corepressor MTG16 regulates small intestinal crypt
proliferation and crypt regeneration after radiation-induced injury. *Am J Physiol-gastr L*
2015;308(6):G562--71.

37. Gunawardene AR, Corfe BM, Staton CA. Classification and functions of enteroendocrine
cells of the lower gastrointestinal tract. *Int J Exp Pathol* 2011;92(4):219–231.

38. Fothergill LJ, Furness JB. Diversity of enteroendocrine cells investigated at cellular and
subcellular levels: the need for a new classification scheme. *Histochem Cell Biol*
2018;150(6):693–702.

39. Akiyama S et al. CCN3 Expression Marks a Sulfomucin-nonproducing Unique Subset of
Colonic Goblet Cells in Mice. *Acta Histochem Cytoc* 2017;50(6):17027.

40. Billing LJ et al. Single cell transcriptomic profiling of large intestinal enteroendocrine cells in
mice – Identification of selective stimuli for insulin-like peptide-5 and glucagon-like peptide-1 co-
expressing cells. *Mol Metab* 2019;29:158–169.

41. Schaum N et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris.
Nature 2018;562(7727):367–372.

42. Jenny M et al. Neurogenin3 is differentially required for endocrine cell fate specification in
the intestinal and gastric epithelium. *Embo J* 2002;21(23):6338–6347.

43. López-Díaz L et al. Intestinal Neurogenin 3 directs differentiation of a bipotential secretory
progenitor to endocrine cell rather than goblet cell fate. *Dev Biol* 2007;309(2):298–305.

44. Gehart H et al. Identification of Enteroendocrine Regulators by Real-Time Single-Cell
Differentiation Mapping. *Cell* 2019;176(5):1158-1173.e16.

45. Li HJ, Ray SK, Kucukural A, Gradwohl G, Leiter AB. Reduced Neurog3 Gene Dosage Shifts
Enteroendocrine Progenitor Towards Goblet Cell Lineage in the Mouse Intestine. *Cell Mol*
Gastroenterology Hepatology [published online ahead of print: 2020];
doi:10.1016/j.jcmgh.2020.08.006

46. Yan KS et al. Intestinal Enteroendocrine Lineage Cells Possess Homeostatic and Injury-
Inducible Stem Cell Activity. *Cell Stem Cell* 2017;21(1):78-90.e6.

47. Engel ME, Nguyen HN, Mariotti J, Hunt A, Hiebert SW. Myeloid translocation gene 16
(MTG16) interacts with Notch transcription complex components to integrate Notch signaling in
hematopoietic cell fate specification.. *Mol Cell Biol* 2010;30(7):1852--63.

48. Flier LG van der et al. Transcription Factor Achaete Scute-Like 2 Controls Intestinal Stem
Cell Fate. *Cell* 2009;136(5):903–912.

49. Wice BM, Gordon JI. Forced Expression of Id-1 in the Adult Mouse Small Intestinal
Epithelium Is Associated with Development of Adenomas*. *J Biol Chem* 1998;273(39):25310–
25319.

50. O'Brien CA et al. ID1 and ID3 Regulate the Self-Renewal Capacity of Human Colon Cancer-
Initiating Cells through p21. *Cancer Cell* 2012;21(6):777–792.

51. Biyajima K et al. Id2 deletion attenuates Apc-deficient ileal tumor formation. *Biol Open*
2015;4(8):993–1001.

52. Rockman SP et al. Id2 Is a Target of the β-Catenin/T Cell Factor Pathway in Colon
Carcinoma*. *J Biol Chem* 2001;276(48):45113–45119.

53. Yu Y, Yang W, Li Y, Cong Y. Enteroendocrine Cells: Sensing Gut Microbiota and Regulating
Inflammatory Bowel Diseases. *Inflamm Bowel Dis* 2019;26(1):11–20.

54. Khaloian S et al. Mitochondrial impairment drives intestinal stem cell transition into
dysfunctional Paneth cells predicting Crohn's disease recurrence. *Gut* 2020;69(11):1939–1951.

55. Gersemann M et al. Differences in goblet cell differentiation between Crohn's disease and
ulcerative colitis. *Differentiation* 2009;77(1):84–94.

56. Sünderhauf A et al. Loss of mucosal p32/gC1qR/HABP1 triggers energy deficiency and
impairs goblet cell differentiation in ulcerative colitis. *Cell Mol Gastroenterology Hepatology*
[published online ahead of print: 2021]; doi:10.1016/j.jcmgh.2021.01.017

57. Treveil A et al. Regulatory network analysis of Paneth cell and goblet cell enriched gut
organoids using transcriptomics approaches. *Mol Omics* 2019;16(1):39–58.

58. Hyams JS et al. Clinical and biological predictors of response to standardised paediatric
colitis therapy (PROTECT): a multicentre inception cohort study. *Lancet*
2019;393(10182):1708–1720.

59. Haberman Y et al. Ulcerative colitis mucosal transcriptomes reveal mitochondriopathy and
personalized mechanisms underlying disease severity and treatment response. *Nat Commun*
2019;10(1):38.

60. Fenton CG, Taman H, Florholmen J, Sørbye SW, Paulssen RH. Transcriptional Signatures
That Define Ulcerative Colitis in Remission. *Inflamm Bowel Dis* 2020;27(1):94–105.

61. Carey R et al. Activation of an IL-6:STAT3-dependent transcriptome in pediatric-onset
inflammatory bowel disease. *Inflamm Bowel Dis* 2008;14(4):446–457.

62. Perse M, Cerar A. Dextran Sodium Sulphate Colitis Mouse Model: Traps and Tricks. *J
Biomed Biotechnol* 2012;2012:1--13.

63. Dieleman et al. Chronic experimental colitis induced by dextran sulphate sodium (DSS) is
characterized by Th1 and Th2 cytokines. *Clin Exp Immunol* 1998;114(3):385–391.

64. Rahman M et al. Alternative preprocessing of RNA-Sequencing data in The Cancer
Genome Atlas leads to improved analysis results. *Bioinformatics* 2015;31(22):3666–3672.

65. Chang K et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*
2013;45(10):1113–1120.

66. Rajamäki K et al. Genetic and epigenetic characteristics of inflammatory bowel disease
associated colorectal cancer. *Gastroenterology* [published online ahead of print: 2021];
doi:10.1053/j.gastro.2021.04.042

67. Andrews N et al. An unsupervised method for physical cell interaction profiling of complex
tissues. *Nat Methods* 2021;18(8):912–920.

68. Schonhoff SE, Giel-Moloney M, Leiter AB. Neurogenin 3-expressing progenitor cells in the
gastrointestinal tract differentiate into both endocrine and non-endocrine cell types. *Dev Biol*
2004;270(2):443–454.

69. Wang Y et al. Single-cell transcriptome analysis reveals differential nutrient absorption
functions in human intestine. *J Exp Med* 2019;217(2):e20191130.

70. Beumer J et al. High-Resolution mRNA and Secretome Atlas of Human Enteroendocrine
Cells. *Cell* 2020;181(6):1291-1306.e19.

71. Li HJ, Ray SK, Singh NK, Johnston B, Leiter AB. Basic helix-loop-helix transcription factors
and enteroendocrine cell differentiation. *Diabetes Obes Metabolism* 2011;13(s1):5–12.

72. Longo A, Guanga GP, Rose RB. Crystal Structure of E47−NeuroD1/Beta2 bHLH
Domain−DNA Complex: Heterodimer Selectivity and DNA Recognition † , ‡. *Biochemistry-us*
2008;47(1):218–229.

73. Yui S et al. YAP/TAZ-Dependent Reprogramming of Colonic Epithelium Links ECM
Remodeling to Tissue Regeneration. *Cell Stem Cell* 2018;22(1):35-49.e7.

74. Qu M et al. Establishment of intestinal organoid cultures modeling injury-associated
epithelial regeneration. *Cell Res* 2021;1–13.

75. Lee C-C et al. TCF12 Protein Functions as Transcriptional Repressor of E-cadherin, and Its
Overexpression Is Correlated with Metastasis of Colorectal Cancer*. *J Biol Chem*
2012;287(4):2798–2809.

76. Zhao H et al. E2A suppresses invasion and migration by targeting YAP in colorectal cancer
cells. *J Transl Med* 2013;11(1):317.

77. Short SP et al. Kaiso is required for MTG16-dependent effects on colitis-associated
carcinoma. *Oncogene* 2019;38(25):5091–5106.

78. Chyla BJ et al. Deletion of Mtg16, a Target of t(16;21), Alters Hematopoietic Progenitor Cell
Proliferation and Lineage Allocation ▿ †. *Mol Cell Biol* 2008;28(20):6234–6247.

79. Rozenblatt-Rosen O et al. The Human Tumor Atlas Network: Charting Tumor Transitions
across Space and Time at Single-Cell Resolution. *Cell* 2020;181(2):236–249.

80. Liu Q et al. Quantitative assessment of cell population diversity in single-cell landscapes.
Plos Biol 2018;16(10):e2006687.

81. Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* 2002;30(1):207–210.

82. Barrett T et al. NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids*
Res 2013;41(D1):D991–D995.

83. Wang F et al. Technical Advance RNAscope A Novel in Situ RNA Analysis Platform for
Formalin-Fixed, Paraffin-Embedded Tissues. *J Mol Diagnostics* 2012;14:22--29.

84. Short SP et al. p120-Catenin is an obligate haploinsufficient tumor suppressor in intestinal
neoplasia. *J Clin Invest* 2017;127(12):4462–4476.

85. Faustino-Rocha A et al. Estimation of rat mammary tumor volume using caliper and
ultrasonography measurements. *Lab Animal* 2013;42(6):217–224.

86. Boivin GP et al. Pathology of mouse models of intestinal cancer: Consensus report and
recommendations. *Gastroenterology* 2003;124(3):762–777.

87. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 2018;34(17):i884–i890.

88. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon: fast and bias-aware
quantification of transcript expression using dual-phase inference. *Nat Methods*
2017;14(4):417–419.

89. Frankish A et al. GENCODE reference annotation for the human and mouse genomes.
Nucleic Acids Res 2018;47(D1):gky955-.

90. RNA-seq workflow: gene-level exploratory analysis and differential expression
[Internet]https://www.bioconductor.org/packages/devel/workflows/vignettes/rnaseqGene/inst/doc
/rnaseqGene.html. cited February 5, 2021

91. Love MI et al. Tximeta: Reference sequence checksums for provenance identification in
RNA-seq. *Plos Comput Biol* 2020;16(2):e1007664.

92. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* 2014;15(12):550.

93. H P, M C, S F, N L. Bioconductor - AnnotationDbi [Internet]. *AnnotationDbi: Manipulation of*
SQLite-based annotations in Bioconductor. R package version 1.54.1.
2021;https://bioconductor.org/packages/release/bioc/html/AnnotationDbi. cited

94. Subramanian A et al. Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *P Natl Acad Sci Usa* 2005;102(43):15545–15550.

95. Liberzon A et al. Molecular signatures database (MSigDB) 3.0. *Bioinformatics*
2011;27(12):1739–1740.

96. Rajamäki K et al. Genetic and epigenetic characteristics of inflammatory bowel disease
associated colorectal cancer. *Gastroenterology* [published online ahead of print: 2021];
doi:10.1053/j.gastro.2021.04.042

97. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Brief Bioinform* 2013;14(2):178–192.

98. Ritchie ME et al. limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* 2015;43(7):e47–e47.

Tables

Genes up in distal colon	$Mtg16^{-/-}$ vs. WT (log$_2$FC)	$Mtg16^{T/T}$ vs. WT (log$_2$FC)	ChIP-seq peak (nearest TSS)?
Agt	1.55	1.29	No
Ces1c	2.77	2.85	No
Dbn1	2.07	0.95	No
Dll3	2.73	2.25	Yes
Dmpk	1.80	1.26	Yes
Gfra3	1.07	0.89	No
Gnao1	1.42	1.20	No
Guca2b	1.14	0.79	Yes
Id4	1.92	1.19	No
Insm1	1.49	0.65	No
Insrr	4.30	3.22	Yes
Kcnh6	1.94	1.17	Yes
Neurog3	**2.35**	**1.09**	**Yes**
Ooep	2.83	2.66	No
Pex5l	2.65	2.02	No
Prss23	2.52	1.59	Yes
Rbp4	1.66	0.88	No
Spon2	2.84	1.93	No
Stxbp1	1.04	0.87	Yes
Syngap1	3.02	3.16	No
Syt13	1.97	0.83	No
Tff3	0.72	0.92	Yes

747 **Table 1. Genes upregulated in both *Mtg16$^{-/-}$* vs. WT and *Mtg16$^{T/T}$* vs. WT distal colon RNA-seq.**
748 **Log2FC.** The log$_2$(fold change) (log$_2$FC) of each gene in each dataset (*Mtg16$^{-/-}$* vs. WT distal colon
749 and *Mtg16$^{T/T}$* vs. WT distal colon) and whether they have a TSS nearest to an MTG16 ChIP-seq
750 peak in a dataset generated by Baulies *et al.* (16) is shown.

Figures

Figure 1. MTG16 is expressed in a subset of colonic secretory cells, and its deficiency alters lineage allocation. (A) *MTG16* expression in cell populations generated from inDrop scRNA-seq of normal human colon biopsies (discovery cohort, n = 35 samples, 30,374 cells) (validation cohort in Fig. S1A). **(B)** *Mtg16* expression in inDrop scRNA-seq of WT mouse colon (n = 3 mice, 3,653 cells). Color gradient represents the average *Mtg16* expression level in each cell population. Dot size represents the percentage of cells in each population expressing *Mtg16.* UMAP plots are in Fig. S1B. **(C)** RNAscope *in situ* hybridization of *Mtg16* and *Muc2* in WT mouse colon. Scale bar = 50 μm. Goblet cells are outlined in the insets at right. **(D-E)** WT and *Mtg16⁻/⁻* mouse colon (n = 10 WT, 9 *Mtg16⁻/⁻*) stained for **(D)** goblet cells by periodic acid-Schiff (PAS) stain and **(E)** enteroendocrine cells by IHC for synaptophysin (SYP). Representative images at left. SYP⁺ cells are indicated with white arrowheads. Scale bars = 100 μm. *$p < 0.05$, ****$p < 0.0001$ by Mann-Whitney test. **(F-G)** Volcano plots demonstrating differentially expressed genes in **(F)** proximal (n = 2 WT, 2 *Mtg16⁻/⁻*) and **(G)** distal (n = 4 WT, 4 *Mtg16⁻/⁻*) colonic epithelial isolates by RNA-seq. Horizontal dashed line indicates $p_{adj} < 0.05$ by DESeq2. Vertical dashed lines indicate fold change = 1.5. Dot colors indicate goblet-associated (blue), enteroendocrine (EE)-associated (red), and secretory-associated (black) genes.

768
Figure 2. MTG16 deficiency is associated with increased enteroendocrine cell markers,
reserve stem cell gene signatures, and E protein target upregulation in the distal colon. (A)
Unsupervised hierarchical clustering of differentially expressed genes in *Mtg16*[-/-] vs. WT (n = 4)
distal colonic epithelial isolates by RNA-seq. Gene expression in individual samples is displayed by
Z-score (normalized count). Genes of interest are labeled at right. **(B)** GSEA of distal colon RNA-seq
using gene sets representing epithelial cell types (described in Table S3). EE, enteroendocrine.
Individual GSEA plots of enteroendocrine and goblet cell signatures are displayed at right. **(C)** GSEA
of distal colon RNA-seq using gene sets representing stem and other progenitor cell populations
(Table S3). **(D)** Staining and quantification of CHGA[+] cells/crypt (top) stratified by location along the
crypt axis (bottom). Scale bar = 50 μm. n = 10 WT, 9 *Mtg16*[-/-]. *$p < 0.05$, **$p < 0.01$ by Mann-Whitney
test. **(E)** GSEA of distal colon RNA-seq using MSigDB (95) gene sets representing E protein and Id
protein transcriptional signatures (Table S3). Individual GSEA plot of the E47 target gene signature
at right. **(B-C, E)** NES, normalized enrichment score (ES). Tag % is defined as the percentage of
gene hits before (for positive ES) or after (for negative ES) the peak in the running ES, indicating the
percentage of genes contributing to the ES. FDR q-value < 0.05 is considered significant.

784
Figure 3. MTG16-driven colonic epithelial lineage allocation is dependent on its repression of
the E protein transcription factors E2A and HEB. (A) WT and *Mtg16^T/T^* (T/T) mouse colon (n = 11
WT, 10 *Mtg16^T/T^*) stained for **(A)** goblet cells by periodic acid-Schiff (PAS) stain, **(B)** enteroendocrine
cells by IHC for synaptophysin (SYP), and **(C)** tuft cells by IHC for doublecortin-like kinase 1
(DCLK1). Representative images at left. Scale bars = 100 μm. *p < 0.05, **p < 0.01 by Mann-
Whitney test. **(D-E)** Volcano plots demonstrating differentially expressed genes in **(D)** proximal (n = 2
WT, 2 *Mtg16^T/T^*) and **(E)** distal (n = 3 WT, 4 *Mtg16^T/T^*) colonic epithelial isolates by RNA-seq.
Horizontal dashed line indicates p_{adj} < 0.05 by DESeq2. Vertical dashed lines indicate fold change =
1.5. Dot colors indicate goblet-associated (blue), enteroendocrine (EE)-associated (red), and
secretory-associated (black) genes. Dots outlined in orange represent genes that are upregulated or
downregulated in the same direction in the corresponding *Mtg16^-/-^* colon segment ("Dysreg. in *16^-/-^*").
(F-H) GSEA of distal colon RNA-seq using gene sets representing **(F-G)** epithelial cell types and **(H)**
E protein transcriptional signatures (Table S3). EE, enteroendocrine. NES, normalized enrichment
score (ES). Tag % is defined as the percentage of gene hits before (for positive ES) or after (for
negative ES) the peak in the running ES, indicating the percentage of genes contributing to the ES.
FDR q-value < 0.05 is considered significant.

801
Figure 4. MTG16 represses E protein-mediated transcription of *Neurog3*. (A) Venn diagram of
upregulated ($p_{adj} < 0.05$) genes from bulk RNA-seq of WT vs. *Mtg16$^{-/-}$* (orange) and WT vs *Mtg16$^{T/T}$*
(T/T) (purple) distal colonic crypt isolates (n = 3-4) and genes with transcription start sites (TSS)
nearest to MTG16 peaks in a ChIP-seq dataset generated by Baulies *et al.* (16) (green). Table 1
provides more information about the common upregulated genes. **(B)** MTG16 occupancy of an E
box (5'-CATNNG)-rich region 5' to *Neurog3* in ChIP-seq by Baulies *et al.* (16). Scale: 0-2. **(C-D)**
Staining and quantification of *Neurog3$^+$* enteroendocrine progenitor cells by RNAscope in **(C)** WT vs.
Mtg16$^{-/-}$ colon (n = 9-10) and **(D)** WT vs. *Mtg16$^{T/T}$* colon (n = 4-5). *$p < 0.05$, **$p < 0.01$ by Mann-
Whitney test.

811
Figure 5. *Mtg16* is upregulated in experimental colitis and patients with IBD and is required for ASCL2-mediated colonic epithelial regeneration. (A) *MTG16* expression from an RNA-seq dataset generated from rectal biopsies of treatment-naïve pediatric UC patients (n = 206) vs. unaffected controls (n = 20) in the PROTECT study (58). Transcript counts were normalized by DESeq2. ****p_{adj} < 0.0001 by DESeq2. **(B)** *MTG16* expression in an RNA-seq dataset from adult ulcerative colitis (UC) patients with active disease (n = 14), UC patients in remission (n = 14), and unaffected control patients (n = 16) (60). Transcript counts were normalized by DESeq2. **p_{adj} < 0.01 and ****p_{adj} < 0.0001 by pairwise DESeq2. **(C)** *MTG16* expression in a microarray dataset comparing UC (n = 5), Crohn's colitis (n = 20), and unaffected (n = 8) patient biopsies (61). **p_{adj} < 0.01 and ****p_{adj} < 0.0001 by limma in GEO2R. Normalized probe intensity is plotted for data visualization. **(A-C)** The lines within each box represent the mean, the bounds of the boxes represent 25th to 75th %-iles, and the bounds of the whiskers represent the range of the data. All data points are shown. **(D)** Schematic of the dextran sulfate sodium (DSS) injury-regeneration model in which mice are treated with 2% for 5 d followed by 4 d of recovery. **(E)** Volcano plot demonstrating significantly downregulated (blue) and upregulated (red) genes in distal colon crypt isolates from WT mice following DSS-induced injury and regeneration compared to baseline (n = 3). Horizontal

dashed line indicates $p_{adj} < 0.05$ by DESeq2. Vertical dashed lines indicate fold change = 2. **(F)**
ASCL2 occupancy near an *Mtg16* transcription start site (TSS 2) in a ChIP-seq dataset of FACS-
sorted *Ascl2*+ regenerating cells compared to *Lgr5*+ homeostatic crypt-base columnar stem cells
(CBCs) generated by Murata *et al.* (2). Scale: 0-800. **(G)** GSEA plot demonstrating significant
enrichment of the *Ascl2*+ de-differentiating cell signature (Table S3) in in distal colon crypt isolates
from WT mice following DSS-induced colitis vs. WT mice at baseline (n = 3). **(H-K)** GSEA performed
on RNA-seq of *Mtg16*-/- vs. WT distal colon crypt isolates following DSS injury-regeneration (n = 3
WT, 4 *Mtg16*-/-) using **(H-I)** epithelial regeneration-associated gene sets and **(J-K)** gene sets
representing E protein transcriptional signatures (described in Table S3). **(G-K)** NES, normalized
enrichment score (ES). Tag % is defined as the percentage of gene hits before (for positive ES) or
after (for negative ES) the peak in the running ES, indicating the percentage of genes contributing to
the ES. FDR q-value < 0.05 is considered significant.

840
Figure 6. MTG16-mediated colonic epithelial regeneration is dependent on its repression of E
protein transcription factor-mediated transcription. (A) Schematic of the dextran sulfate sodium
(DSS) injury-regeneration model. **(B-D)** Colitis severity in *Mtg16^{T/T}* (T/T) vs. WT mice evaluated
using **(B)** weight loss, **(C)** colon length, and **(D)** histologic injury (scoring system described in Table
S2). **(D)** Representative images at left. Scale bar = 200 μm. **(E)** Percent of colon with histologic
findings, estimated from the distal end, as a measure of proximal extent of disease. **(B-E)** Data are
combined from 2 independent experiments (n = 15 WT, 16 *Mtg16^{T/T}*). **(F)** EdU+ cells in injury-
adjacent, regenerating crypts in colons from mice injected with EdU 1 h prior to sacrifice on day 9 (n
= 30-50 crypts from n = 5 mice). Dashed white line indicates ulcerated epithelium. Scale bar = 100
μm. **(B-F)** *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by **(B)** 2-way ANOVA followed by
Sidak's multiple comparison tests or **(C-F)** Mann-Whitney test. **(G-H)** GSEA of distal *Mtg16^{T/T}* vs. WT
colon crypt isolates following DSS injury-regeneration (n = 3) (gene sets described in Table S3).
NES, normalized enrichment score (ES). Tag % is defined as the percentage of gene hits before (for
positive ES) or after (for negative ES) the peak in the running ES, indicating the percentage of genes
contributing to the ES. FDR q-value < 0.05 is considered significant.

856
Figure 7. MTG16-mediated protection from tumorigenesis is partially dependent on
repression of E protein activity. (A) *MTG16* expression in sporadic colorectal cancer (CRC) (n =
482) compared to normal colon tissue (n = 41) by DESeq2 of TCGA raw counts generated by
Rahman *et al.* (64). The lines within each box represent the mean, the bounds of the boxes
represent the 25th to 75th %-iles, and the bounds of the whiskers represent the range of the data. All
individual data points are shown. *p_{adj} < 0.05 by DESeq2. **(B)** *MTG16* expression in sporadic CRC (n
= 47) and colitis-associated cancer (CAC) (n = 17) tumors. *MTG16* expression was normalized by
DESeq2 and limma batch correction. The box extends from the lower to upper quartile values of the
data, with a line at the median. The lower whisker is at the lowest datum above Q1 - 1.5*(Q3-Q1),
and the upper whisker at the highest datum below Q3 + 1.5*(Q3-Q1), where Q1 and Q3 are the first
and third quartiles. **(C)** Schematic of the azoxymethane/dextran sulfate sodium (AOM/DSS) model in
which mice are injected *i.p.* with AOM followed by 3 cycles of DSS-induced injury and recovery prior
to sacrifice. **(D-G)** CAC severity in AOM/DSS-treated *Mtg16^{T/T}* (T/T) vs. WT mice assessed by **(D)**
histologic injury-regeneration score (scoring system described in Table S2), **(E-F)** tumor multiplicity,
and **(G)** average tumor volume per mouse. **(H)** Average tumor position normalized to colon length
(distance from anus/colon length*100%) per mouse. **(J)** Weight loss compared using repeated
measures 2-way ANOVA (n = 26 WT, 25 *Mtg16^{T/T}*, representative of 2 independent experiments).

(K) Colon length measured at sacrifice. **(L)** Survival curves compared using Log-rank (Mantel-Cox)
test. **(O)** Percent of tumors in each grade of dysplasia, evaluated by a pathologist blinded to
genotype and compared by X^2 test. **(D-F, G-I, K-M)** Data are pooled from 2 independent
experiments with n = 35 WT, 26 $Mtg16^{T/T}$ remaining at sacrifice. *$p < 0.05$ by **(D, F-H, K)** Mann-
Whitney test or **(I)** least-squares regression.

A Human colon

Absorptive colonocytes
Crypt top colonocytes
Enteroendocrine
Goblet
Stem
Transit amplifying
Tuft

MTG16 expression

B Mouse colon

Enteroendocrine
Tuft
Goblet
Stem
Absorptive colonocytes

Average expression level

% cells

Mtg16 expression

C

Merge

Mtg16   *Muc2*

DAPI   DAPI

D

Periodic Acid-Schiff

WT   *Mtg16⁻/⁻*

PAS⁺ cells/crypt/mouse

WT   *16⁻/⁻*

E

Synaptophysin

WT   *Mtg16⁻/⁻*

SYP cells/crypt/mouse

WT   *16⁻/⁻*

F *Mtg16⁻/⁻* vs. WT - Proximal colon

- Goblet-assoc.
- EE-assoc.
- Sec.-assoc.

$-\log_{10}(p_{adj})$

Ces1c
Reg4
Muc1
Spdef
Mmp7   *Atoh1*   *Tff3*
Best2   *Stxbp1*
B3galt2   *Sst*   *Pyy*
Tac1
Kif1a

$\log_2(\text{fold change})$

G *Mtg16⁻/⁻* vs. WT - Distal colon

$-\log_{10}(p_{adj})$

Chgb
Tph1   *Chga*
Nk2-2   *Neurod1*   *Syngap1*
Pax6   *Neurog3*
Insm1   *Pax4*   *Asxl3*
Atoh1   *Lmx1a*   *Sct*   *Ces1c*
Clca1   *Tff3*   *Dll3*
B3galt2   *Stxbp1*
Kif1a   *Myt1*

$\log_2(\text{fold change})$

A Periodic Acid-Schiff — WT / T/T

PAS⁺ cells/crypt/mouse; WT vs T/T (*)

B DAPI / E-cadherin / CHGA — WT / T/T

CHGA⁺ cells/crypt/mouse; WT vs T/T (**)

C DCLK1 — WT / T/T

DCLK1⁺ cells/crypt/mouse; WT vs T/T (ns)

D T/T vs. WT - Proximal colon

Legend:
- Goblet-assoc. (blue)
- EE-assoc. (red)
- Sec.-assoc. (black)
- Dysreg. in $16^{-/-}$ (orange open)

Axes: $-\log_{10}(p_{adj})$ vs $\log_2$(fold change)

Labeled genes: *Muc1*, *Clca4b*, *B3galt2*, *Mmp7*, *Plet1*, *Pla2g2a*, *Defa24*, *Ces1d*, *Ang4*, *Tff3*, *Chga*, *Chgb*, *Clca1*, *Dll1*, *Kif1a*, *Agt*, *Etv5*, *Reg4*

E T/T vs. WT - Distal colon

Axes: $-\log_{10}(p_{adj})$ vs $\log_2$(fold change)

Labeled genes: *Ces1c*, *Agt*, *Syngap1*, *Stxbp1*, *Insm1*, *Tff3*, *Syt13*, *Neurog3*, *Reg3g*, *Reg3b*, *Plet1*, *Klb*, *Dll3*, *Dlk1*

F Differentiated cells / EE progenitors

Differentiated cells:
- EE
- Enterochromaffin
- L-cell
- Absorptive
- Tuft
- Goblet
- BEST4/OTOP2⁺ colonocyte

EE progenitors:
- Early EE
- Early/Intermediate EE
- Intermediate EE
- Intermediate/Late EE
- Late EE
- *Neurog3*-EGFP⁺/⁺

Axis: NES; FDR q-val.; Tag %

G Stem cells and progenitors

- *mTert*-GFP⁺
- *Bmi1*-GFP⁺
- FVR^Low
- *Rbpj*-DBZ Sec-Pro
- Cd166^Hi
- *MKI67*^Hi
- *Lrig1*-GFP⁺
- *Lgr5*⁺ ISC
- *Lgr5*⁺ ISC (colon)

Axis: NES; FDR q-val.; Tag %

H E protein and Id protein targets

- E12_Q6
- E47_02
- HEB_Q6
- E47_01
- E2A_Q2
- E2-2 (TCF4_Q5)
- ID2_TARGETS
- ID1_TARGETS

Axis: NES; FDR q-val.; Tag %

A

Genes upregulated in distal colon

115 | 33 | ChIP-seq peaks (nearest genes)

5893

13 | 9

3 | 5

Mtg16⁻/⁻

T/T

B

Input_1 0-2

Input_2

MTG16_1

MTG16_2

5'-CATNNG (E box)

Neurog3 →

C

DAPI *Neurog3*

WT

Mtg16⁻/⁻

Neurog3⁺ cells/crypt/mouse

WT | 16⁻/⁻

**

D

DAPI *Neurog3*

WT

T/T

Neurog3⁺ cells/crypt/mouse

WT | T/T

*

A 2% DSS | Recovery | ▼Sac
Day 0   5   9

B
% Initial weight
   0 1 2 3 4 5 6 7 8 9
   —DSS— —Water—
        Days
○ WT
● T/T
* *** **** **** ****

C
Colon length (cm)
**

D
Hist. injury score (0-24)
WT T/T
****

E
% Colon affected
WT T/T
**

F
EdU
DAPI
EdU⁺ cells/injury-adjacent crypt
60
40
20
0
WT T/T
*

G
Mtg16^{T/T} DSS colitis vs. WT DSS colitis
Intestinal epithelial regeneration signatures

DSS-induced colitis regenerating epithelium
Composite injury-regeneration signature
Fetal epithelial gene signature
scRNA-seq cluster SSC2c "revival" cells
Clu⁺ "revival" cell signature
Ascl2⁺ de-differentiating cell signature

NES -3 -2 -1 0 1 2 3

FDR q-val.
1.00
0.75
0.50
0.25
0.00

Tag %
30
40
50
60

H
Enrichment score
0.6
0.5
0.4
0.3
0.2
0.1
0.0
DSS-induced colitis regenerative epithelium (Wang *et al.* 2019)
NES: 2.58
FDR: 0.00
T/T colitis   WT colitis

Enrichment score
0.0
-0.1
-0.2
-0.3
-0.4
-0.5
Ascl2 de-differentiating cell signature (Murata *et al.* 2020)
NES: -2.69
FDR: 0.00
T/T colitis   WT colitis

A

MTG16 expression (normalized counts)

B

MTG16 expression (normalized counts)

C

AOM *i.p.*

| DSS cycle 1 | Recovery cycle 1 | DSS cycle 2 | Recovery cycle 2 | DSS cycle 3 | Recovery cycle 3 |

Day   0   3   7   23   27   43   47   63

Sac

D

Histologic injury score (0-24)

E

WT

T/T

F

Gross tumor multiplicity (tumors/mouse)

G

Gross tumor volume (mm³/mouse)

H

Avg. tumor position/mouse norm. to colon length (%)

I

Gross tumor volume (mm³)

Distance from anus normalized to colon length (%)

J

% Initial weight

Days

K

Colon length (cm)

L

% Survival

Days

M

% of tumors

Tumor grade
Invasive adenocarcinoma
Intramucosal adenocarcinoma
High grade dysplasia
Low grade dysplasia