Integrative Modelling of Signalling Network Dynamics Identifies Cell Type-selective Therapeutic Strategies for FGFR4-driven Cancers

Sung-Young Shin[1,2,*], Nicole J Chew[1,2,*], Milad Ghomlaghi[1,2], Anderly C Chüeh[1,2], Yunhui Jeong[1,2], Lan K. Nguyen[1,2,#] and Roger J Daly[1,2,#]

[1]Cancer Program, Biomedicine Discovery Institute, Monash University, Melbourne, VIC 3800, Australia.
[2] Department of Biochemistry and Molecular Biology, Monash University, Melbourne, VIC 3800, Australia.

* These authors contributed equally to the work
# Correspondence: roger.daly@monash.edu; lan.k.nguyen@monash.edu

Abstract

Oncogenic FGFR4 signalling represents a potential therapeutic target in various cancer types, including triple negative breast cancer (TNBC) and hepatocellular carcinoma (HCC). However, resistance to FGFR4 single-agent therapy remains a major challenge, emphasizing the need for effective combinatorial treatments. Our study sought to develop a comprehensive computational model of FGFR4 signalling and provide network-level insights into resistance mechanisms driven by signalling dynamics. Our integrated approach, combining computational network modelling with experimental validation, uncovered potent AKT reactivation following FGFR4 targeting in the TNBC cell line MDA-MB-453. By systematically simulating the model to analyse the effects of co-targeting specific network nodes, we were able to predict, and subsequently confirm through experimental validation, the strong synergy of co-targeting FGFR4 and AKT or specific ErbB kinases, but not PI3K. Incorporating protein expression data from hundreds of cancer cell lines, we then adapted our model to diverse cellular contexts. This revealed that while AKT rebound is common, it is not a general phenomenon. ERK reactivation, for example, occurs in certain cell types, including the FGFR4-driven HCC cell line Hep3B, where there is a synergistic effect of co-targeting FGFR4 and MEK, but not AKT. In summary, our study offers key insights into drug-induced network remodelling and the role of protein expression heterogeneity in targeted therapy responses. We underscore the utility of computational network modelling for designing cell type-selective combination therapies and enhancing precision cancer treatment.

Running title - Computational modelling of the FGFR4 signalling network

Significance: This study underscores the potential of computational predictive modelling in deciphering mechanisms of cancer cell resistance to targeted therapies and in designing more effective, cancer type-specific combination treatments.

Keywords: cancer cell resistance, combination therapy, network rewiring, signal transduction, computational mechanistic modelling, cancer type-specific treatments.

Introduction

Aberrant signalling by specific members of the FGFR family, comprising FGFR1-4, occurs in a variety of human cancers, reflecting FGFR gene alterations and/or enhanced ligand expression (1). This has led to the development of selective small molecule drugs targeting these receptors, including Erdafitinib (an inhibitor of FGFR1-4), which was recently FDA-approved for patients with metastatic urothelial carcinoma (2). While initial interest focused on FGFR1-3, oncogenic roles for FGFR4 in a variety of cancers have now become evident (3). In breast cancer, enhanced FGFR4 expression and mutation is associated with metastatic progression, particularly in the lobular subtype (4), and high FGFR4 expression positively correlates with endocrine resistance (4,5). FGFR4 also drives phenotypic switching of luminal A breast cancers to a HER2-enriched gene expression phenotype (6), and is overexpressed in approximately one third of triple negative breast cancers (TNBCs) (7). A second cancer where FGFR4 is strongly implicated is hepatocellular carcinoma (HCC). Here, FGFR4 is activated as a consequence of overexpression of its ligand, FGF19, which is a 'driver' oncogene on the chromosome 11q13.3 amplicon (8,9). These oncogenic roles for FGFR4 have led to the development of BLU9931, BLU-554 (fisogatinib) and H3B-6527, small molecule drugs that selectively target FGFR4 and exhibit promising pre-clinical activity in FGF19-driven HCC (9-11). Both fisogatinib and H3B-6527 are currently under evaluation in clinical trials for patients with advanced HCC (e.g. NCT04194801 and NCT02834780).

While targeted therapies directed towards specific oncogenic protein kinases have greatly improved clinical management of particular cancers (12), intrinsic and acquired drug resistance remain a major problem (13). One mechanism that can limit the efficacy of targeted therapy is dynamic, adaptive rewiring of the cellular signalling network in response to treatment (1). For example, treatment with AKT or MEK inhibitors can lead to increased expression and/or activation of a suite of receptor tyrosine kinases (RTKs) that dampens the cellular response to the drug (14,15). Importantly, characterization of network adaptation in response to drug treatment can inform the rational design of combination therapies that exhibit improved efficacy (14).

Despite these advances, the complexity of kinase signalling networks presents a major roadblock to understanding mechanisms of adaptive resistance to targeted therapy (1). This complexity reflects the presence of features such as pathway crosstalk and both positive and negative feedback loops, and results in non-intuitive behaviour that makes logical prediction of signalling and biological outputs challenging. A powerful approach to address this issue is to integrate mathematical network modelling with experimental analysis in order to generate and validate predictive models. This can provide fundamental insights into the wiring of signalling networks and how this generates specific signalling behaviours and biological outputs (16,17). In addition, it can be exploited to identify novel therapeutic strategies, as exemplified by accurate prediction of synergistic drug combinations by establishing a quantitative mechanistic model of the EGFR-PYK2-MET signalling network in TNBC (18).

In this study, we developed a comprehensive mechanistic model of the FGFR4 signalling network and used an integrative approach of computational modelling and experimental studies to analyse how the network responds to anti-FGFR4 inhibition across different cancer types and to identify combinatorial approaches that circumvent drug resistance. Overall, our work establishes an integrative framework for network-level analysis of drug-induced signalling adaptation that can be applied not only to FGFR4 inhibition but also other kinase-directed targeted therapies.

RESULTS

FGFR4 inhibition rewires signalling networks in MDA-MB-453 cells leading to upregulation of AKT activity

To determine the effects of the selective and irreversible FGFR4 inhibitor BLU9931 on FGFR4 downstream signalling and cell proliferation in TNBC, the MDA-MB-453 cell line that exhibits an activating mutation in FGFR4 was utilised (19). The MDA-MB-468 cell line with no detected FGFRs was used as a negative control (20). MDA-MB-453 cells were treated with different concentrations of BLU9931 for 1 h. Total FGFR4 expression was not affected by BLU9931, but the treatment decreased pFGFR at concentrations of 3nM and higher (Fig. 1A), and this effect was paralleled by a significant decrease in downstream signalling proteins pFRS2, pERK and pAKT (Fig. 1A, S1). MDA-MB-453 cell proliferation was significantly decreased by BLU9931 at concentrations of 10nM or higher, starting from day 3 after treatment (Fig. 1B). However, BLU9931 did not have any effect on MDA-MB-468 cell proliferation (Fig. 1C).

To understand how FGFR4 inhibition temporally influences downstream signalling, MDA-MB-453 cells were treated with 10nM of BLU9931 for 1, 4, 8 and 24 h (Fig. 1D). The levels of pFRS2 and pERK were initially decreased, followed by some signal recovery at 24 h (Fig. 1D). However, pAKT levels decreased after 1 h of drug treatment but the signal recovered at 4 h and then increased at 8 h and 24 h relative to the vehicle control (Fig. 1D). To identify if the bounce-back of pAKT is independent of the inhibitor used, MDA-MB-453 cells were treated with another FGFR4 inhibitor, H3B-6527, a specific FGFR4 inhibitor currently undergoing clinical trials for solid tumours (NCT02834780, NCT03424577). This resulted in the same pAKT rebound effect at the same time points (Fig. 1D-F). This indicates a general rewiring phenomenon following FGFR4 inhibition in MDA-MB-453 cells.

A new mechanistic model of the integrated FGFR4 signalling network

To achieve a holistic and systematic understanding of drug-induced network rewiring, we developed a novel mechanistic mathematical model of the integrated FGFR signalling network. (Fig. 2A, S2). The model scope and assumptions along with supporting evidence are described in detail in Supplementary Text S1. The model was formulated using ordinary differential equations

(ODEs), which describe the time-dependent rates of change of each species in the network based
on the underlying biochemical reactions and their kinetics (see Materials and Methods). Detailed
description of the reaction rates, kinetic laws used, and full set of ODEs are provided in
Supplementary Tables S1-2.

To render our model context-specific and predictive, we first trained it using combined time- and
dose-dependent signalling datasets from MDA-MB-453 cells, measured by Western blot (Fig.
1A,D). This was achieved by employing genetic algorithm-based optimisation procedures
(Supplementary Text S1), which involved estimating the model parameters to ensure that model
simulations fit the experimental data. Next, we conducted model identifiability analysis (21-24) to
test if unique best-fitted parameter values could be obtained (see Materials & Methods and
Supplementary Text S1). The analysis revealed partial identifiability (Fig. S3-4); i.e. at least one
parameter remained non-uniquely identified, which aligns with the complexity of our model
(24,25). This implies existence of multiple optimal parameter sets that fit the data similarly. We
iterated the optimisation process 300 times using a genetic algorithm, yielding 50 equally well-
fitted parameter sets (Supplementary Data S2-4). These sets were collectively used for simulations,
which displayed strong concordance with the quantified experimental data (Fig. 2B). This
confirms that the calibrated model, hereafter referred to as *model-1*, effectively reproduces the data
both qualitatively and quantitatively.

Concomitant inhibition of AKT, but not PI3K, eliminates pAKT rebound and AKT co-
targeting enhances inhibition of MDA-MB-453 cell proliferation

To determine whether the rebound in AKT activation following FGFR4 inhibition is dosage-
dependent, we utilised model-1 to simulate the temporal response of pAKT to escalating doses of
BLU9931. Interestingly, the model predicted that high doses of BLU9931 induced an even
stronger rebound of pAKT than low doses, with the effect being more pronounced at later time
points (Fig. 3A). Given that PI3K is a direct upstream kinase of AKT signalling, we asked if co-
targeting FGFR4 with a PI3K inhibitor could eliminate the rebound in pAKT. Model simulations
showed that while combined FGFR and PI3K inhibition significantly suppressed pAKT levels, it
did not completely eliminate the rebound pattern of pAKT (Fig. 3B).

To experimentally validate these model predictions, we first selected specific concentrations of the
PI3Kα-specific inhibitor BYL719 to combine with BLU9931 (Fig. S5). Next, MDA-MB-453 cells
were treated with BLU9931 and BYL719 alone or in combination for 1 and 24 h (Fig. 3C). The
highest dose of BLU9931 and BYL719 decreased pAKT at 1 h, as did all the combination
treatments (Fig. 3C). However, consistent with our model prediction, the higher dose of BLU9931
triggered a stronger rebound of pAKT at 24 h (Fig. 3C). Moreover, while the BLU9931 + BYL719
combination treatment suppressed the pAKT signal recovery, it did not completely eliminate the
rebound, as predicted by our model.

Model simulations predicted that co-inhibition of AKT would strongly suppress the
phosphorylation of AKT, as expected (Fig. 3D). To validate this experimentally, MDA-MB-453
cells were treated with BLU9931 and AKT inhibitor MK2206 alone or in combination for 4 h and
24 h (Fig. 3E, Fig. S6). The BLU9931 + MK2206 combination treatment potently suppressed
pAKT at 24 hr, in agreement with our model prediction (Fig. 3D). MK2206 treatment alone also
significantly suppressed pAKT to a greater extent than BLU9931 alone at 4 hr (Fig. 3E). In line
with these findings, treatment with MK2206 significantly decreased the proliferation of MDA-
MB-453 cells (Fig. 3F). Moreover, combining BLU9931 with MK2206 was significantly more
effective in blocking cell proliferation than either agent alone (Fig. 3F). This work was further
developed using the FGFR4 inhibitor H3B-6527. MDA-MB-453 cells were treated with different
dosages of H3B-6527 and MK-2206 alone or in combination. Consistent with the BLU9931 data,
the combined H3B-6527 + MK-2206 treatments caused synergistic effects on cell viability,
confirming that the effect is independent of the FGFR4 inhibitor used (Fig. 3G). Drug synergism
was quantified using the Bliss Independence and Loewe methods (Fig. S7A-B).
178
The prediction that inhibiting AKT is more effective than inhibiting PI3K in eliminating the pAKT
rebound, despite the same target inhibition levels and PI3K being upstream of AKT, was intriguing
(Fig. S8A). To gain further insight into this observation, we first performed model-based
sensitivity analysis and identified network links that control the differential effects of PI3Ki versus
AKTi on FGFR4i-induced pAKT reactivation. The sensitivity analysis (Fig. S8B) and time-course
simulations (Fig. S9) demonstrated that the links related to negative feedback regulation, including
the CBL-, SPRY2-, and PTP-mediated negative feedback loops, enhanced the differential effect
of AKTi and PI3Ki on pAKT rebound (red line, Fig S8C). Conversely, the links associated with
GAB proteins and positive feedback reduced the differential effects (blue line, Fig S8C). Next, we
analysed the correlation between the magnitude of pAKT rebound and the differential effects of
AKTi and PI3Ki. For this, we introduced the rebound (RB) score to describe the early response of
pAKT to FGFR4i, followed by a rebound at a later time (Fig. S8D). We found that a stronger
pAKT rebound exaggerates the differential effect between AKTi and PI3Ki on suppression of
pAKT, as shown in Fig. S8E. This indicates that when the FGFR4-induced pAKT rebound is more
pronounced, AKT is a more suitable co-target of FGFR4 than PI3K.
194
195 Collectively, our integrative studies indicated that the efficacy of FGFR4 inhibitors in this TNBC
196 cell line is limited by the rebound of AKT activity, which can be more effectively inhibited by co-
197 targeting AKT rather than PI3K.
198
199 **The role of specific RTKs in modulating FGFR4i sensitivity**
200
To further investigate the dynamic signalling remodelling following FGFR4 inhibition, we
enhanced our mathematical model by including additional time-course and dose-response
signalling data from combined FGFR and PI3K inhibition experiments (Fig. S10). These datasets,
previously excluded from model-1, helped recalibrate the model. The refined model-2 accurately

replicated all training data using a new suite of 50 best-fitted parameter sets (Fig. S11; model
identifiability analysis results in Figs S12-13).

Next, we extended our model simulation readouts beyond pAKT to include phosphorylated IRS,
IGF1R and ErbBs in response to FGFR4i, as these proteins are regulated by multiple negative
feedback loops (Fig. 2A). Model simulations predicted strong upregulation of pIGF1R, pIRS and
pErbB at 24 hr following BLU9931 treatment at both low and high doses (Fig. 4A). Interestingly,
higher BLU9931 concentrations triggered stronger upregulation of these proteins (Fig S14), as
seen previously for pAKT (Figs. 3A-B). These predictions suggest that FGFR4 inhibition
remodelled the RTK network in TNBC cells, causing pAKT rebound and compromising the
inhibitory effect of FGFR4i on cell proliferation. To test this hypothesis and the model predictions,
we assessed the expression and phosphorylation of specific RTKs in MDA-MB-453 cells at 1 hr
and 24 hr after BLU9931 treatment. Consistent with the model predictions, we observed significant
upregulation of pIGF1R, pIRS, and pErbB2-3 at 24 hr compared to the control or 1 hr treatment
(Fig. 4B).

In addition to investigating the short-term remodelling of the FGFR4 network in response to
BLU9931, we also interrogated its long-term effects. To this end, we exposed MDA-MB-453 cells
to BLU9931 for at least 3 months to allow isolation of resistant cells (see Material and Methods).
Long-term drug-treated cells showed no significant growth inhibition at 10 nM BLU9931
compared to parental cells (Fig. 4C), while 100 nM of the drug reduced growth, although this
potency was stronger in parental cells (Fig. 4D). To investigate the mechanism underlying long-
term resistance to BLU9931, we sequenced the FGFR4 kinase domain and characterized the
expression profile of FGFR4 signalling network components in parental and long-term BLU9931-
treated cells. FGFR4 gatekeeper mutations (26) were not detected (Fig S15), and similar to parental
cells treated for 24 h, long-term BLU9931-treated cells showed a marked upregulation of pAKT,
pIGF1R, pIRS, and pErbB2-3 (Fig. 4B). Furthermore, levels of phosphorylated GSK3α/β and
pY1248 ErbB2 were higher in long-term BLU9931-treated cells than in parental cells treated for
24 hrs. While increased activation of particular RTKs induced by 24-hour BLU9931 treatment is
maintained in the long-term drug-refractory cells, additional changes likely underpin long-term
resistance to BLU9931. These data suggest potential targets in addition to AKT for combination
therapies with FGFR4 inhibition, which will be addressed next.

Model prediction and validation of synergistic drug combinations co-targeting FGFR4

Our computational model provides a powerful framework for the systematic prediction of nodes
that could be co-targeted with FGFR4 to overcome adaptive resistance to anti-FGFR4
monotherapy. To achieve this, we further refined model-2 by incorporating the new experimental
data from Fig. 4B and S16 for model recalibration, evolving it to become *model-3*. The refined
model reproduces all the training data well (Fig. S17), and displays improved identifiability
compared to the previous models (Figs. S18-20). Using model-3, we conducted in silico

simulations to evaluate the efficacy of 19 possible pair-wise drug combinations co-targeting
FGFR4 and 19 other network nodes (Fig. 5A). Highest Single Agent (HSA) and the coefficient of
drug interaction (CDI) were computed to evaluate the theoretical performance of each combination
in suppressing cell proliferation relative to single-drug treatments (Fig. 5A and S21A, also see
Materials and Methods). By sorting these synergy scores, we were able to rank and prioritise the
drug combinations based on their predicted synergistic effects (Fig. 5A and S21A).
252
Our model predicted synergistic drug combinations for co-targeting FGFR4, with dual FGFR4 and
AKT inhibition showing the most synergistic effect (Fig. 5A). In contrast, co-targeting FGFR4
and MEK was predicted to be non-synergistic. Interestingly, co-targeting ErbB receptors was
predicted to be synergistic (Fig. 5A). Our data demonstrating upregulation of ErbBs and IGF1R in
response to FGFR4 inhibition (Fig. 4) are consistent with our model's prediction that ErbBs would
be a synergistic FGFR4 co-target despite being in a different signalling branch than the IGF1R-
PI3K-AKT cascade. To validate this prediction, we treated long-term BLU9931-maintained cells
with Lapatinib, which predominantly targets ErbB2 in MDA-MB-453 cells since EGFR/ErbB1
expression is low (27), and selected 100 nM as the most appropriate dose for subsequent
experiments (Fig. 5B). Lapatinib treatment alone had no effect on parental or resistant cells (Fig.
5C-D) but the combination treatment of BLU9931 and Lapatinib induced cell death in both
parental and resistant cells (Fig. 5C-D). Note that at 10 nM BLU9931, the combination treatment
with Lapatinib restored growth inhibition only in cells maintained long-term (Fig. S21B-D). To
further confirm the model prediction of non-synergistic combination, we conducted an experiment
where MDA-MB-453 cells were treated with different doses of H3B-6527 and MEK inhibitor
trametinib, either alone or in combination, and their effects on cell viability were assessed (Fig
5E). Interestingly, unlike the synergistic effects observed with the combined H3B-6527+MK-2206
treatments (Fig. 3G), the combination of H3B-6527+trametinib failed to exhibit any synergistic
effects on cell viability inhibition for almost all dose combinations (Fig. 5E and S22).
272
273 Together, these results indicate that long-term BLU9931-maintained cells are more dependent on
274 ErbB2, consistent with the enhanced ErbB2 Y1248 phosphorylation (Fig 4B), and demonstrate
275 that co-targeting ErbB2 synergistically enhances the efficacy of FGFR4 inhibition in MDA-MB-
276 453 TNBC cells.
277
278 **Analysis of network connections that drive pAKT and pErbB rebounds**
279
Our analyses identified RTKs such as ErbBs and IGF1R as potential mediators of pAKT rebound
following FGFR4i treatment. However, the network features that drive this rebound remain
unclear. To address this issue, we conducted a sensitivity analysis using model-3 where we
systematically blocked network links and observed the resulting changes in pAKT rebound
following BLU9931 treatment (Fig. 6A). While a visual assessment indicated strong and varied
changes to the pAKT dynamics depending on the perturbed model parameters (Fig. 6A), to assess

these changes more accurately we computed the rebound metric introduced earlier (*RB*, Fig. S8D)
for each of the perturbed and control conditions (Fig. 6B).

We found that inhibiting certain parameters decreased the RB score, indicating a positive role for
the corresponding network links in mediating pAKT rebound (Fig. 6B). In contrast, for almost a
quarter of the remaining parameters, blocking them had the opposite effect, while the remaining
three-quarters did not have a detectable impact on the RB score (Fig. 6B). By mapping these
parameters onto the network interaction diagram, we found that the magnitude of the pAKT RB
score is under the positive control of the negative feedback loops mediated by CBL and PTP, with
the PTP-to-ErbB feedback loop having the most significant impact (Fig. 6B-C). As these feedback
loops act primarily upon the RTK receptors, including ErbB and IGF1R, the findings align with
our prior experimental results showing the upregulation of these RTKs following FGFR4
inhibition (Fig. 4). On the other hand, we found that network links that had a negative impact on
the pAKT RB score tended to be positioned along the core RTK signalling axes, such as ligand-
to-IGFR1/FGFR4, which promote the activation of the RTKs and downstream signals (Fig. 6C).

To further investigate the mechanism of ErbB signal reactivation since this is likely associated
with the pAKT rebound, we conducted a similar sensitivity analysis on the dynamics of pErbB.
The result revealed that, similar to pAKT rebound, the links related to PTP- and CBL-mediated
negative feedback positively regulated the ErbB RB score (Fig. 6D-F). However, we observed that
the ERK-to-ErbB negative feedback had an impact only on ErbB reactivation, indicating that these
links may be more specific to the ErbB signalling pathway (Fig. 6F). Together, these systematic
analyses identify key network features that drive AKT and ErbB signal reactivation, providing
important insights into the mechanism underlying the signalling dynamic remodelling following
FGFR4 inhibition.

Diverse pAKT/pERK response profiles following FGFR4 inhibition across cancer cell lines

In our previous work (28), we showed that cell-to-cell variability in protein expression levels is a
major source of heterogeneity in signalling response. To explore the hypothesis that FGFR4i-
induced signalling dynamics are dictated by the protein expression profiles across specific cancer
cell types, we obtained protein expression data from the Cancer Cell Line Encyclopedia (CCLE)
consortium for 350 different cancer cell lines representing diverse tissue origins (29). We utilised
this data to customize our MDA-MB-453 based model-3, generating 350 models that reflect the
specific protein expression profile of each cell type (see Fig. S23A for detailed workflow). By
simulating the temporal response of pAKT and pERK after FGFR4 inhibition using these cell type-
specific models, we identified a remarkable heterogeneity in the response of these outputs, ranging
from no significant change to strong rebound to straight increase patterns for pAKT (Fig. 7A),
while pERK predominantly displayed rebound or monotonic decrease patterns (Fig. 7B).

We defined three composite patterns (REB: rebound; INC/NOC: increase or no significant change;
and DEC: decrease) and used them to classify the cell lines into four major subgroups (namely
groups 1-4: REB-REB; INC/NOC-REB; REB-DEC; and INC/NOC-DEC) based on their distinct
response profiles of pAKT/pERK respectively (Fig. 7C and Fig. S23B). Our simulations revealed
that FGFR4 inhibition induced pERK rebound in about two-thirds of the cell lines (66%),
indicating that reactivation of ERK represents a salient response feature post FGFR4i treatment in
a subset of cell types that likely blunts the treatment effect. Although pAKT displayed rebound in
about 71% of the cell lines, our simulations suggest that FGFR4 inhibition either did not
significantly affect or induced a straight increase of pAKT in the remaining cell lines (29%).
Notably, only 5% of the cell lines exhibited concomitant pAKT increase/no change and pERK
decrease, while 42% exhibited rebound in both pAKT and pERK (Fig. 7C).
337
Next, to examine whether the modulation of network components could potentially trigger a
transition between different response groups, we employed the MDA-MB-453 model (group 3)
and systematically altered the expression levels of each component. The perturbation analysis
indicated that downregulation of FGFR4, FRS2, Grb2, or Ras, or upregulation of PDK1, AKT,
PIP2 or PI3K transitioned pAKT from a rebound state to an increase or no change state, resulting
in a group 3 → 4 transition (Fig. S24).
344
345   Together, these studies highlight the heterogeneity and adaptability of signalling responses to
346   FGFR4-targeted therapy in cancer cells. These findings underscore the necessity of considering
347   the network context when analysing drug-induced network remodelling and carry implications for
348   the development of personalized cancer therapies.
349
350   **Alternative network rewiring following selective FGFR4 inhibition in Hep3B cells**
351
352   Our model-based analysis predicted distinct cancer cell subgroups with different response patterns
353   for pAKT-pERK, which can potentially limit the effect of FGFR4 inhibition. The MDA-MB-453
354   cell line represents group 3 (Fig. 1D, F), and to validate the model prediction of group 2, we
355   identified Hep3B, an HCC cell line with active FGFR4 signalling due to FGF19 gene amplification
356   (9), as a member of this group.
357
Western blot analysis of Hep3B cells treated with H3B-6527 revealed a significant decrease in
phosphorylated FRS2 and ERK after 1 hr (Fig. 7D, E). However, we also observed an early and
temporally increasing rebound of pERK levels, despite the durable suppression of pFRS2 (Fig. 7
E). Notably, pAKT showed no significant change after H3B-6527 treatment in these cells (Fig. 7E
and Fig. S25). These dynamics are consistent with the predicted behaviour of group 2, confirming
the validity our model-based approach for identifying cell type-specific response patterns to anti-
FGFR4 treatment. For the Hep3B model, downregulation of ERK, MEK, Raf, or Ras, or
upregulation of AKT transitioned pERK from a rebound state to a decrease, triggering a group 2
→ 4 transition (Fig. S26).

The H3B-6527 treatment-induced rebound of pERK suggests that co-targeting ERK may enhance
the sensitivity of Hep3B cells to H3B-6527. To test this hypothesis, we first determined the
appropriate concentration of trametinib, an inhibitor that targets ERK's direct upstream kinase
MEK, to be used in combination treatments. Treatment with 5nM or higher of trametinib led to
decreased pERK in Hep3B cells (Fig. S27). We then tested the effects of combined H3B-6527 and
trametinib in comparison to single-drug treatments. While H3B-6527 and trametinib alone
decreased Hep3B cell proliferation (Fig. 7F), their combination resulted in strong synergistic
effects (Fig. 7F, S28A-B). Interestingly, when we treated the cells with H3B-6527 and the AKT
inhibitor MK-2206, we did not observe strong synergistic effects between them (Fig. 7G and
S28C-D). This observation is consistent with the fact that pAKT was not significantly affected by
H3B-6527 treatment (Fig. 7G). These results indicate that the efficacy of H3B-6527 is improved
by blocking the ERK signal bounce-back and highlights the combination of H3B-6527 and
trametinib as a promising therapeutic approach for HCC with this network behaviour.

Discussion

In this study we develop a comprehensive computational model for FGFR4 signalling and shed
new light on the mechanisms utilised by cancer cells to counter FGFR4 inhibition. Furthermore,
we identify heterogeneity in tumour cell responses to FGFR4 inhibition and based on this,
highlight potential context-specific combined drug strategies for enhanced personalised treatment
outcomes.

Our new computational model of the FGFR4 signalling network integrates key signalling cascades
downstream of FGFR4 and is rigorously calibrated using diverse time-course and dose-response
data sets, employing ordinary differential equation formalism. This represents a significant step
forward from early models, which either failed to capture intracellular signalling (30-34) or
neglected key downstream pathways (35,36). Some recent studies have adopted logic-based
Boolean modelling (37); however, these cannot capture the complex temporal aspects of drug-
induced signalling dynamics. Therefore, our model fills these critical gaps by providing a more
detailed and comprehensive platform for the analysis and prediction of these dynamics.

The integration of our computational modelling and experimental analyses has shed light on the
network rewiring occurring in breast cancer cells to resist FGFR4 inhibition. In MDA-MB-453
TNBC cells we determined that selective inhibition of FGFR4 leads to significant reactivation of
the pro-survival protein AKT. However, an interesting predicted and validated phenomenon was
that targeting AKT more effectively eliminated AKT rebound than inhibiting PI3K. Sensitivity
analysis showed that blocking negative feedback mechanisms, such as CBL-, SPRY2-, and PTP-
mediated loops, mitigated the differential effect of AKT and PI3K inhibitors on pAKT rebound.
Further analysis revealed a correlation between the strength of pAKT rebound and the differential
impact of AKT and PI3K inhibitors. Stronger pAKT rebound widens this differential effect, while

its absence narrows it, indicating a dominant role for feedback loops like CBL and PTP in
controlling the contrasting impacts of AKT and PI3K inhibitors on pAKT response. Although
experimental validation of these predictions is pending, our computational findings have
illuminated otherwise non-intuitive observations.
412
The focus of most research has been on combining FGFR4 inhibition with radio or
chemotherapeutics, leaving the area of combinatorial targeting of FGFR4 relatively understudied
(38,39). To expedite the identification of novel combination treatments that target FGFR4 we
predicted the impact of pair-wise drug combinations in a systematic and unbiased manner. This
highlighted ErbB receptors as FGFR4 co-targets in MDA-MB-453 cells, consistent with the
activation of alternative RTKs as an escape mechanism previously reported in FGFR-resistance
models (40,41). For example, FGFR3-dependent bladder cancer cell lines developed resistance to
the pan-FGFR inhibitor BGJ398 by switching receptor signalling to ErbB2 or ErbB3 (40).
421
Given these insights, it is important to note that the mechanisms underlying adaptive resistance
can vary significantly depending on the context, even when the same drug is used (42). This
variability is driven by differences in protein expression and mutational profiles among different
cancer cell types, which influence feedback strength and crosstalk, ultimately dictating drug
response (18,28). To address this, we incorporated cell type-specific protein expression data to
generate hundreds of cell type-specific computational models, which led to the identification of
distinct response groups that exhibit unique behaviours after FGFR4 inhibition. We identified and
experimentally validated the HCC cell line Hep3B as a cell type exhibiting pERK rebound with
no significant change in pAKT. Importantly, co-targeting FGFR4 and MEK, upstream of ERK, in
Hep3B cells significantly and synergistically enhanced the inhibition of cell proliferation,
highlighting this approach as a promising precision treatment strategy for HCC. The unique
clustering of response dynamics uncovered here underscore the need to consider network context
in drug resistance analysis, and has potential implications in clinical settings, especially in the
context of personalized cancer therapies.
436
We also demonstrated that manipulation of network interference points could shift response
behaviours, such as transitioning from rebound to progressive increase in pAKT activation. This
illustrates how cancer cells can exploit the plasticity of signalling networks to evolve from drug-
sensitive to drug-resistant states. Furthermore, this new knowledge can assist in designing new
therapeutic strategies to counteract drug-induced reactivation of pro-growth signalling by
transforming pAKT/pERK (re)activation into a reduction in activity. These findings highlight the
effectiveness of using computational modelling in identifying nodal alterations that manipulate
dynamic signalling behaviour. Adding to the novelty of our study, we introduced innovative
metrics that enable systematic and quantitative detection and characterization of drug-induced
adaptive rebound in signalling proteins, including the rebound (RB) metric. Not only does this
offer a more precise definition of adaptive signalling rebound, but it also enables systematic
comparison across various conditions. Indeed, the utilization of these metrics in our study has

facilitated model sensitivity analysis of potential mechanisms underlying the rebound of AKT and
ErbB activities following FGFR4 inhibition, which would have been challenging otherwise.

In conclusion, by taking an integrative approach, we have uncovered the mechanisms that drive
FGFR4 network rewiring and discovered potential combinatorial targeted therapies that deliver
improved efficacy for FGFR4-driven TNBC and HCC cancers. Our finding that co-targeting
FGFR4 and ErbB2 was particularly effective suggests that this targeted treatment strategy could
extend to other breast cancer subtypes with poor prognosis, including luminal B and HER2 breast
cancers. Furthermore, the computational models and integrative systems-based approach utilised
in this study lay a robust foundation for future investigations into network rewiring and adaptive
resistance related to targeting other RTKs.

Materials and Methods

Cell lines, cell culture and reagents
The TNBC cell line, MDA-MB-453 and HCC cell line, Hep3B were purchased from ATCC.
MDA-MB-453 was cultured in RPMI-1640 (Gibco) supplemented with 10% (v/v) FBS
(Moregate), 10 μg/mL Actrapid penfill insulin (Clifford Hallam Healthcare) and 20 mM HEPES
(Gibco). Hep3B was cultured in EMEM (USbio) supplemented with 10% (v/v) FBS and 1 mM
sodium pyruvate (Gibco).

Inhibitors and treatment
The following inhibitors were purchased from Selleckchem: FGFR4 inhibitor BLU9931 and H3B-
6527, ErbB family inhibitor Lapatinib, PI3Kα inhibitor BYL719, AKT inhibitor MK2206 and
MEK inhibitor Trametinib. All inhibitors were reconstituted in DMSO.
For inhibitor treatment, cells were seeded into culture plates with an 80% end point confluence for
all cell lines. After 24 h, cells were treated for the indicated times with the specific inhibitor or
DMSO as vehicle control.

Generation of FGFR4 inhibitor resistant cells
MDA-MB-453 cells were seeded into 10 cm plates at a density of 500 cells. After 48 h, cells were
treated with DMSO (as vehicle control) or FGFR4 inhibitor BLU9931 for at least 3 months.
Culture medium was replaced twice a week. When cells formed colonies visible to the naked eye,
small pieces of sterile filter paper soaked in trypsin were used to detach cells which were collected
for further maintenance in media containing DMSO or FGFR4 inhibitor. These cells were termed
long-term BLU9931 MDA-MB-453 cells in this study.

Immunoblotting
Protein lysates were prepared in RIPA buffer and subjected to Western blotting as described
previously (43).

The following antibodies were purchased from Cell Signaling Technology: pan-phosFGFR
(Y653/654) (3471), AKT (4685), ERK (4695), pAKT (S473) (4058), pERK (T202, Y204) (4370),
pFRS2 (Y436) (3861), ErbB2 (2165), pErbB2 (Y1248) (2247), ErbB3 (4754), pErbB3 (Y1328)
(8017), IGF-1R (9750) and GSK-3α/β (5676). The following antibodies were purchased from
Santa Cruz Biotechnology: FGFR4 (sc-136988) and β-actin (sc-69879). The α-tubulin (T5168)
and FRS2 (05-502) antibodies were purchased from Sigma-Aldrich. The pIRS-1 (Y612) (44-
816G) and pIGF-1R (Y1162/1163) (44-804G) were purchased from Biosource. An IRS-1 (6248)
antibody was purchased from Upstate and a pGSK-3α/β (Y216/279) (ab4797-50) was purchased
from Abcam.
498
499 **Cell viability assays**
500 Trypsinized cells were stained with Trypan blue (EVS-1000, NanoTek), then transferred to an
501 EVE cell counting slide (EVS-1000, NanoTek) and counted with the EVE automatic cell counter
502 (EVE-MC-DEMO, NanoTek) according to the manufacturer's protocol. MTS proliferation assays
503 were undertaken as previously described (43).
504
RNA isolation, RT-PCR and Sanger sequencing
Total RNA was isolated from parental MDA-MB-453 and long-term BLU9931 MDA-MB-453
cells using a RNeasy mini kit (Qiagen) following the manufacturer's protocol. RNA was quantified
using a Nanodrop ND-1000 (NanoDrop Technologies). RNAs were reverse transcribed using a
high-capacity cDNA reverse transcription kit (Thermoscientific). Subsequently, cDNA was
amplified by PCR to identify gatekeeper mutations in the FGFR4 kinase domain using forward (F)
primers and reverse (R) primers (Table S3). The PCR products were resolved by gel
electrophoresis, and the bands at the predicted product size were excised and purified with a gel
and PCR clean-up system (Promega). Sanger sequencing was completed by the Micromon facility
at Monash University. Reactions were repeated on three biological replicates.
515
516 **Statistical analysis**
517 Quantification of western blots by densitometry was performed using ImageLab version 5.2.1
518 (Bio-Rad) and statistical t-tests were performed using GraphPad Prism 8 and Microsoft-Excel.
519
Computational modelling
The FGFR4-centered model was formulated using ordinary differential equations (ODEs). The
rate equations and full set of ODEs are given in Supplementary Tables S1-2. The model
construction and calibration processes were implemented in MATLAB (The MathWorks. Inc.
2023a) and the IQM toolbox (http://www.intiquan.com/intiquan-tools/) was used to compile the
IQM file for a MEX file which makes the simulation faster. To facilitate model exchange, an
exchangeable Systems Biology Markup Language (SBML) file of the model is provided as
Supplementary Data S1-3. The code for the modelling has been deposited to Github and can be
accessed at https://github.com/NguyenLabNetworkModeling/FGFR4-Signaling-Network-
Model.git.

Model calibration

The adequacy and specificity of a mathematical model is generally justified by its ability to recapitulate experimental data. This is achieved through model calibration (also referred to as model training or fitting) that involves estimation of the unmeasured model parameters so that simulations using these parameter values (called best-fitted parameter sets) could recapitulate the training data (Fig. S30-32). Here, parameter estimation was done by minimizing an objective function J that quantifies the discrepancies between model simulations and the corresponding experimental measurements. A genetic algorithm was used to optimize the objective function (44-46), utilising the Global Optimization Toolbox and the function *ga* in MATLAB. Model calibration was carried out on a virtual machine consisting of 32 Intel Xeon 2.2 GHz processors running in parallel. A more detailed description of the model calibration process is given in Supplementary Text S1. As a result, we obtained 50 best-fitted parameter sets that were collectively used for model simulations (Fig. S32); these are given in Supplementary Data S2-4.

Model-based simulation and computation of drug synergy

Because the synergy between two drugs can depend on the specific doses at which they are combined, we used IC50 value of individual drugs as a reference concentration for the simulation (see Supplementary Information for details). Using the simulated dose-response matrix, we evaluated in silico the efficacy and possible synergism of 19 possible combinatorial strategies co-targeting FGFR4 and each of 19 network components. Drug synergy was computed based on the coefficient of drug interaction (CDI) and highest single agent (HSA) metric (47-49): CDI = $R_{12}/(R_1 \times R_2)$, where R_{12} is a normalized biological response (e.g., cell viability) by the combined treatment of drug 1 and 2, and R_1 and R_2 are the response by the single drug treatment, respectively. Note $R = 1$ (and 0) indicates no drug response (complete inhibition, respectively). CDI <1, $= 1$ or >1 indicates that the drugs are synergistic, additive or antagonistic, respectively. According to the HSA model, the synergistic effect is equal to the greater effect of individual drugs (49). Thus, HSA $= E_{12} - \max(E_1, E_2)$. E_{12} is a normalized drug effect by the combined treatment of drug 1 and 2, and E_1 and E_2 are the response by the single drug treatment, respectively. Note $E = 0$ (and 1) indicates no drug effect (maximum effect, respectively). For the readability and easy comparison between CDI and HAS, we used a log2-transformed CDI metric (-log2(CDI)): Thus, for the both metrics the score > 0, $= 0$ or < 0 indicates that the drugs are synergistic, additive or antagonistic, respectively. In addition, we employed Bliss and Loewe models and SynergyFinder+, an open-source package (50), which can be accessible at www.synergyfinderplus.org, to quantify experimental data for drug synergy.

To comparatively assess the effect of the single-drug and combination treatments, we introduced a theoretical cell viability (ICV) function, defined as the aggregate of the activated levels of the major pro-growth signalling nodes as follows:

$$ICV = \sum_i W_i \cdot R_i$$

where R_i is the level of activated pro-growth signalling proteins (pAKT, pERK, pS6K) and W_i is the corresponding protein's contributing weight to cell viability (see Supplementary Text 1 for more detail).

Data availability

The data generated in this study are available within the article and its supplementary data files, and raw data can be provided upon request from the corresponding authors.

Acknowledgements and Funding

L.K.N was supported by a Research Fellowship from Victorian Cancer Agency (MCRF18026). L.K.N's laboratory was supported for this work by an Investigator Initiated Research Scheme grant from National Breast Cancer Foundation (IIRS-20–094) and a Venture Grant from Cancer Council Victoria. R.J.D's laboratory was supported for this work by a Venture Grant from Cancer Council Victoria.

Author Contributions

Conception and design: R.J.D, L.K.N.

Development of methodology: SY.S., N.J.C., M.G., L.K.N, R.J.D.

Acquisition of data: N.J.C., A.C.C., SY.S., M.G., YH.J.

Modelling and simulation: SY.S., M.G.

Analysis and interpretation of data: SY.S., N.J.C., M.G., A.C.C., YH.J., L.K.N, R.J.D.

Writing of the manuscript: SY.S., N.J.C., L.K.N, R.J.D.

Study supervision: R.J.D, L.K.N.

All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no conflict of interest.

References

1. Babina IS, Turner NC. Advances and challenges in targeting FGFR signalling in cancer. Nature reviews Cancer **2017**;17:318-32
2. Loriot Y, Necchi A, Park SH, Garcia-Donas J, Huddart R, Burgess E, *et al.* Erdafitinib in Locally Advanced or Metastatic Urothelial Carcinoma. N Engl J Med **2019**;381:338-48
3. Levine KM, Ding K, Chen L, Oesterreich S. FGFR4: A promising therapeutic target for breast cancer and other solid tumors. Pharmacology & Therapeutics **2020**;214:107590
4. Levine KM, Priedigkeit N, Basudan A, Tasdemir N, Sikora MJ, Sokol ES, *et al.* FGFR4 overexpression and hotspot mutations in metastatic ER+ breast cancer are enriched in the lobular subtype. npj Breast Cancer **2019**;5:19
5. Meijer D, Sieuwerts AM, Look MP, van Agthoven T, Foekens JA, Dorssers LC. Fibroblast growth factor receptor 4 predicts failure on tamoxifen therapy in patients with recurrent breast cancer. Endocr Relat Cancer **2008**;15:101-11
6. Garcia-Recio S, Thennavan A, East MP, Parker JS, Cejalvo JM, Garay JP, *et al.* FGFR4 regulates tumor subtype differentiation in luminal breast cancer and metastatic disease. J Clin Invest **2020**;130:4871-87
7. Chew NJ, Lim Kam Sian TCC, Nguyen EV, Shin SY, Yang J, Hui MN, *et al.* Evaluation of FGFR targeting in breast cancer through interrogation of patient-derived models. Breast Cancer Res **2021**;23:82
8. Sawey ET, Chanrion M, Cai C, Wu G, Zhang J, Zender L, *et al.* Identification of a therapeutic strategy targeting amplified FGF19 in liver cancer by Oncogenomic screening. Cancer Cell **2011**;19:347-58
9. Joshi JJ, Coffey H, Corcoran E, Tsai J, Huang CL, Ichikawa K, *et al.* H3B-6527 Is a Potent and Selective Inhibitor of FGFR4 in FGF19-Driven Hepatocellular Carcinoma. Cancer Res **2017**;77:6999-7013
10. Hagel M, Miduturu C, Sheets M, Rubin N, Weng W, Stransky N, *et al.* First Selective Small Molecule Inhibitor of FGFR4 for the Treatment of Hepatocellular Carcinomas with an Activated FGFR4 Signaling Pathway. Cancer Discov **2015**;5:424-37
11. Kim RD, Sarker D, Meyer T, Yau T, Macarulla T, Park J-W, *et al.* First-in-Human Phase I Study of Fisogatinib (BLU-554) Validates Aberrant FGF19 Signaling as a Driver Event in Hepatocellular Carcinoma. Cancer Discovery **2019**;9:1696-707
12. Fleuren ED, Zhang L, Wu J, Daly RJ. The kinome 'at large' in cancer. Nature reviews Cancer **2016**;16:83-98
13. Saraon P, Pathmanathan S, Snider J, Lyakisheva A, Wong V, Stagljar I. Receptor tyrosine kinases and cancer: oncogenic mechanisms and therapeutic approaches. Oncogene **2021**;40:4079-93
14. Chandarlapaty S, Sawai A, Scaltriti M, Rodrik-Outmezguine V, Grbovic-Huezo O, Serra V, *et al.* AKT inhibition relieves feedback suppression of receptor tyrosine kinase expression and activity. Cancer Cell **2011**;19:58-71
15. Duncan JS, Whittle MC, Nakamura K, Abell AN, Midland AA, Zawistowski JS, *et al.* Dynamic reprogramming of the kinome in response to targeted MEK inhibition in triple-negative breast cancer. Cell **2012**;149:307-21
16. Shin SY, Kim T, Lee HS, Kang JH, Lee JY, Cho KH, *et al.* The switching role of beta-adrenergic receptor signalling in cell survival or death decision of cardiomyocytes. Nature communications **2014**;5:5777

17. Nakakuki T, Birtwistle MR, Saeki Y, Yumoto N, Ide K, Nagashima T*, et al.* Ligand-
specific c-Fos expression emerges from the spatiotemporal control of ErbB network
dynamics. Cell **2010**;141:884-96
18. Shin SY, Muller AK, Verma N, Lev S, Nguyen LK. Systems modelling of the EGFR-
PYK2-c-Met interaction network predicts and prioritizes synergistic drug combinations
for triple-negative breast cancer. PLoS Comput Biol **2018**;14:e1006192
19. Roidl A, Foo P, Wong W, Mann C, Bechtold S, Berger HJ*, et al.* The FGFR4 Y367C
mutant is a dominant oncogene in MDA-MB453 breast cancer cells. Oncogene
2010;29:1543-52
20. Chew NJ, Nguyen EV, Su SP, Novy K, Chan HC, Nguyen LK*, et al.* FGFR3 signaling
and function in triple negative breast cancer. Cell Commun Signal **2020**;18:13
21. Maiwald T, Hass H, Steiert B, Vanlier J, Engesser R, Raue A*, et al.* Driving the Model to
Its Limit: Profile Likelihood Based Model Reduction. PLOS ONE **2016**;11:e0162366
22. Rateitschak K, Winter F, Lange F, Jaster R, Wolkenhauer O. Parameter Identifiability
and Sensitivity Analysis Predict Targets for Enhancement of STAT1 Activity in
Pancreatic Cancer and Stellate Cells. PLOS Computational Biology **2012**;8:e1002815
23. Raue A, Kreutz C, Maiwald T, Bachmann J, Schilling M, Klingmüller U*, et al.* Structural
and practical identifiability analysis of partially observed dynamical models by exploiting
the profile likelihood. Bioinformatics **2009**;25:1923-9
24. Villaverde AF, Barreiro A, Papachristodoulou A. Structural Identifiability of Dynamic
Systems Biology Models. PLOS Computational Biology **2016**;12:e1005153
25. Kearney AL, Norris DM, Ghomlaghi M, Kin Lok Wong M, Humphrey SJ, Carroll L*, et
al.* Akt phosphorylates insulin receptor substrate to limit PI3K-mediated PIP3 synthesis.
eLife **2021**;10:e66942
26. Hatlen MA, Schmidt-Kittler O, Sherwin CA, Rozsahegyi E, Rubin N, Sheets MP*, et al.*
Acquired On-Target Clinical Resistance Validates FGFR4 as a Driver of Hepatocellular
Carcinoma. Cancer Discovery **2019**;9:1686-95
27. Jeong H, Kim J, Lee Y, Seo JH, Hong SR, Kim A. Neuregulin-1 induces cancer stem cell
characteristics in breast cancer cell lines. Oncol Rep **2014**;32:1218-24
28. Norris D, Yang P, Shin SY, Kearney AL, Kim HJ, Geddes T*, et al.* Signaling
Heterogeneity is Defined by Pathway Architecture and Intercellular Variability in Protein
Expression. iScience **2021**;24:102118
29. Nusinow DP, Szpyt J, Ghandi M, Rose CM, McDonald ER, Kalocsay M*, et al.*
Quantitative Proteomics of the Cancer Cell Line Encyclopedia. Cell **2020**;180:387-
402.e16
30. Fannon M, Forsten KE, Nugent MA. Potentiation and Inhibition of bFGF Binding by
Heparin: A Model for Regulation of Cellular Response. Biochemistry **2000**;39:1434-45
31. Filion RJ, Popel AS. A reaction-diffusion model of basic fibroblast growth factor
interactions with cell surface receptors. Ann Biomed Eng **2004**;32:645-63
32. Forsten KE, Fannon M, Nugent MA. Potential mechanisms for the regulation of growth
factor binding by heparin. J Theor Biol **2000**;205:215-30
33. Gopalakrishnan M, Forsten-Williams K, Täuber UC. Ligand-induced coupling versus
receptor pre-association: cellular automaton simulations of FGF-2 binding. J Theor Biol
2004;227:239-51
34. Ibrahimi OA, Zhang F, Hrstka SC, Mohammadi M, Linhardt RJ. Kinetic model for FGF,
FGFR, and proteoglycan signal transduction complex assembly. Biochemistry
2004;43:4724-30

35.   Yamada S, Taketomi T, Yoshimura A. Model analysis of difference between EGF
692   pathway and FGF pathway. Biochemical and Biophysical Research Communications
693   **2004**;314:1113-20
694   36.   Kanodia J, Chai D, Vollmer J, Kim J, Raue A, Finn G, *et al.* Deciphering the mechanism
695   behind Fibroblast Growth Factor (FGF) induced biphasic signal-response profiles. Cell
696   Communication and Signaling **2014**;12:34
697   37.   Lee J, Choi SR, Cho KH. Network Dynamics Caused by Genomic Alteration Determine
698   the Therapeutic Response to FGFR Inhibitors for Lung Cancer. Biomolecules **2022**;12
699   38.   Turkington R, Longley D, Allen W, Stevenson L, McLaughlin K, Dunne P, *et al.*
700   Fibroblast growth factor receptor 4 (FGFR4): a targetable regulator of drug resistance in
701   colorectal cancer. Cell death & disease **2014**;5:e1046-e
702   39.   Ahmed MA, Selzer E, Dörr W, Jomrich G, Harpain F, Silberhumer GR, *et al.* Correction:
703   Fibroblast growth factor receptor 4 induced resistance to radiation therapy in colorectal
704   cancer. Oncotarget **2019**;10:5385
705   40.   Wang J, Mikse O, Liao RG, Li Y, Tan L, Janne PA, *et al.* Ligand-associated ERBB2/3
706   activation confers acquired resistance to FGFR inhibition in FGFR3-dependent cancer
707   cells. Oncogene **2015**;34:2167-77
708   41.   Herrera-Abreu MT, Pearson A, Campbell J, Shnyder SD, Knowles MA, Ashworth A, *et*
709   *al.* Parallel RNA interference screens identify EGFR activation as an escape mechanism
710   in FGFR3-mutant cancer. Cancer discovery **2013**;3:1058-71
711   42.   Cunningham DL, Sarhan AR, Creese AJ, Larkins KPB, Zhao H, Ferguson HR, *et al.*
712   Differential responses to kinase inhibition in FGFR2-addicted triple negative breast
713   cancer cells: a quantitative phosphoproteomics study. Sci Rep **2020**;10:7950
714   43.   Chew NJ, Nguyen EV, Su S-P, Novy K, Chan HC, Nguyen LK, *et al.* FGFR3 signaling
715   and function in triple negative breast cancer. Cell Communication and Signaling
716   **2020**;18:13
717   44.   Reali F, Priami C, Marchetti L. Optimization Algorithms for Computational Systems
718   Biology. Frontiers in Applied Mathematics and Statistics **2017**;3
719   45.   Man KF, Tang KS, Kwong S. Genetic algorithms: concepts and applications [in
720   engineering design]. IEEE Transactions on Industrial Electronics **1996**;43:519-34
721   46.   Shin SY, Kim T, Lee HS, Kang JH, Lee JY, Cho KH, *et al.* The switching role of β-
722   adrenergic receptor signalling in cell survival or death decision of cardiomyocytes.
723   Nature communications **2014**;5:5777
724   47.   Li X, Lin Z, Zhang B, Guo L, Liu S, Li H, *et al.* β-elemene sensitizes hepatocellular
725   carcinoma cells to oxaliplatin by preventing oxaliplatin-induced degradation of copper
726   transporter 1. Scientific Reports **2016**;6:21010
727   48.   Liu F, Shang Y, Chen S-z. Chloroquine potentiates the anti-cancer effect of lidamycin on
728   non-small cell lung cancer cells in vitro. Acta Pharmacol Sin **2014**;35:645-52
729   49.   Yadav B, Wennerberg K, Aittokallio T, Tang J. Searching for Drug Synergy in Complex
730   Dose-Response Landscapes Using an Interaction Potency Model. Comput Struct
731   Biotechnol J **2015**;13:504-13
732   50.   Zheng S, Wang W, Aldahdooh J, Malyutina A, Shadbahr T, Tanoli Z, *et al.*
733   SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination
734   Screening Datasets. Genomics Proteomics Bioinformatics **2022**;20:587-96

736 **FIGURES**

737
738

Figure 1. Effect of the FGFR4 inhibitor BLU9931 on FGFR4 downstream signalling
pathways and proliferation in the MDA-MB-453 cell line. (**A**) Dose dependent effect of
BLU9931 on expression and activation of downstream signalling proteins in MDA-MB-453 cells,
1 h post-treatment with the indicated doses. Effect of BLU9931 on proliferation in the FGFR4-
positive MDA-MB-453 (**B**) and FGFR4-negative control MDA-MB-468 cell lines (**C**). Cell
proliferation was determined by direct cell counting. Error bars: mean ± standard error of three
biological replicates. ** indicates p-value of < 0.01 and *** < 0.001, comparing individual
BLU9931 concentrations to the DMSO vehicle control. U indicates untreated control, D indicates
DMSO vehicle control. (**D**) Time course analysis of BLU9931 treatment. Expression and
activation of downstream signalling proteins 1, 4, 8 and 24 h post-treatment with 10 nM of
BLU9931. (**E**) Dose dependent effect of H3B-6527 on expression and activation of downstream
signalling proteins in MDA-MB-453 cells, 1 h post-treatment with the indicated doses. (**F**) Time
course analysis of H3B-6527 treatment. Expression and activation of downstream signalling
proteins 1, 4, 8 and 24 h post-treatment with 10 nM of H3B-6527. Representative of three
biological replicates. Note that the pFGFR signals were detected using a pan-phospho-FGFR
antibody.

Figure 2. Development and calibration of a mathematical model of the FGFR4 signalling

network. (A) A simplified schematic diagram depicting the network interactions included in the

FGFR4 model, consisting of major RTKs, converging downstream signalling pathways and key

feedback and crosstalk mechanisms. A detailed model reaction diagram is provided in Figure S2.

The full model reaction rates and ODEs are given in Supplemental Table S1-2. **(B)** Model fitting

to experimental data. Comparison of model simulation using best-fitted parameter sets (blue lines,

error bars: mean ± standard error, n=50) against experimental time-course and dose-response data

(red lines, error bars: mean ± standard error, n≥2) demonstrate good agreement between simulation

and data. Data used for model calibration were quantified from Figures 1A, D.

768
Figure 3. Co-inhibition of FGFR4 and AKT, but not PI3K, eliminates pAKT rebound and
suppresses MDA-MB-453 cell proliferation. (A-B) Model prediction of phosphorylated AKT
(pAKT) dynamics in response to FGFR4 inhibitor BLU9931 and PI3Kα inhibitor BYL719 in
single or combination treatment. Error bars: mean ± standard error of 50 best-fitted parameter sets.

We used the IC99 (FGFR4i (high)) and IC90 concentration (FGFR4i (low)) to inhibit FGFR4. For
the inhibition of PI3K we used the IC90 concentration. **(C)** Dose dependent effect of BLU9931 in
combination with BYL719 on AKT phosphorylation in MDA-MB-453 cells. pAKT was
characterized 1 h and 24 h post-treatment with the indicated treatments. U indicates untreated
control, D indicates DMSO vehicle control (represenative of 3 independent experiments). **(D)**
Model prediction of pAKT dynamics in response to FGFR4 and AKT inhibitors in single or
combination treatments. We used the IC99 (FGFR4i(high)) and IC90 concentration
(FGFR4i(low)) to inhibit FGFR4. For the inhibition of AKT we used the IC90 concentration. Error
bars: mean ± standard error of 50 best-fitted parameter sets. **(E)** Dose dependent effect of
BLU9931 in combination with AKT inhibitor MK-2206 on pAKT in MDA-MB-453 cells. pAKT
was characterized 1 h and 24 h post-treatment with the indicated treatments. U indicates untreated
control, D indicates DMSO vehicle control (represenative of 3 independent experiments). **(F)**
MDA-MB-453 cells were subjected to single inhibitor treatment or a combination of BLU9931
and MK-2206. Cell proliferation was determined by MTS assay. Error bars: mean ± standard error
of four biological replicates. ** indicates p-value of < 0.01. **(G)** Combined drug effect of FGFR4
inhibitor H3B-6537 and AKT inhibitor MK-2206 on cell proliferation in the MDA-MB-453 cell
line. Cell proliferation was determined by MTS assay. Error bars: mean ± standard error of three
biological replicates.
791

Figure 4. Identification of upregulated RTKs in parental and BLU9931-resistant MDA-MB-453 cells. **(A)** Model prediction of the dynamic responses of specific signalling proteins, predicting they are temporally upregulated following BLU9931 treatment. Error bars: mean ± standard error of 50 best-fitted parameter sets. **(B)** Expression and activation of RTKs and FGFR4 downstream signalling proteins at 1 and 24 h post-treatment with the indicated BLU9931 concentrations in parental MDA-MB-453 cells, compared with long-term (LT) BLU9931-resistant MDA-MB-453 cells. The latter were harvested 24 h post-seeding. **(C-D)** The proliferation of parental and long-term BLU9931-resistant MDA-MB-453 cells treated with DMSO vehicle control, 10 nM (C) or 100 nM BLU9931 (D). Cell proliferation was determined by MTS assay. Error bars: mean ± standard error of six biological replicates. ** indicates p-value of < 0.01, *** < 0.001.

803
804
Figure 5. Model prediction of synergistic drug combinations and experimental validation.
(A) In silico prediction of the effect of 19 possible pair-wise drug combinations co-targeting
FGFR4 and various network components, assessed using drug synergy scores of HSA. The value
that is >1, =1, <1 indicate synergistic, additive or antagonistic effects, respectively; and higher
HSA scores indicate stronger synergism (Materials and Methods). Error bars: mean ± standard
error of 50 best-fitted parameter sets. **(B)** Dose dependent effect of Lapatinib on the expression
and phosphorylation of ErbB2 in long-term 100 nM BLU9931-resistant MDA-MB-453 cells. **(C-**

D) Proliferation of parental and long-term 100 nM BLU9931-resistant MDA-MB-453 cells

subjected to single inhibitor or combination treatment of BLU9931 (BLU) and Lapatinib (Lap) at

the indicated doses. Cell proliferation was determined by MTS assay. **(E)** Combined drug effect

of FGFR4 inhibitor H3B-6537 and MEK inhibitor trametinib on cell proliferation of the MDA-

MB-453 cell line. Cell proliferation was determined by MTS assay. Error bars: mean $\pm$ standard

error of three biological replicates. * indicates p-value of <0.05, ** < 0.01, *** < 0.001.

Figure 6. Model-based sensitivity analysis and identification of network architecture controlling pAKT and pErbB rebound. (**A, D**) A 3D plot (waterfall plot) showing in silico simulation of the temporal phosphorylated AKT and ErbB dynamics in response to FGFR4 inhibition. The dynamics of the intact (non-perturbed) model is highlighted in white as a control. The pAKT and pErbB time profiles were averaged across 50 best-fitted parameter sets. (**B, E**) Changes in the RB score of BLU9931-induced pAKT and pErbB response were measured by a sensitivity score, relative to the control (no perturbation) condition. If the sensitivity score is less than 1, it indicates that the parameters promote the reactivation of pAKT and ErbB. If the score is greater than 1, it indicates that the parameters inhibit it. (**C, F**) Simplified network models and key mechanisms that play a critical role in inducing the reactivation of pAKT and pErbB.

830
831
Figure 7. Analysis of drug-induced pAKT/pERK response profiles across a large panel of
cancer cell types. (A, B) Model prediction of pAKT and pERK time-course profiles in response
to FGFR4 inhibition in 350 cancer cell lines using cell-type specific models generated by
incorporating protein expression data from CCLE. **(C)** Classification of the cell lines into
subgroups based on the response patterns of pAKT and pERK. INC: increase, DEC: decrease,
REB: rebound, NOC: no change. **(D)** Dose dependent effect of H3B-6527 on expression and
activation of FGFR4 downstream signalling proteins in Hep3B cells 1 h post-treatment with the
indicated doses. **(E)** Time course analysis of treatment using the FGFR4 inhibitor H3B-6527 on
FGFR4 downstream signalling pathways in the Hep3B cell line. Expression and activation of
downstream signalling proteins were characterized 1, 4, 8 and 24 h post-treatment with 50 nM

H3B-6527. U indicates untreated control, D indicates DMSO vehicle control. Representative of
three biological replicates. Combined drug effect of FGFR4 inhibitor H3B-6537 and MEK
inhibitor Trametinib (**F**) or AKT inhibitor MK-2206 (**G**) on cell proliferation in the Hep3B cell
line. Error bars: mean ± standard error of three biological replicates. Cell proliferation was
determined by MTS assay.