Selecting species for restoration in foundational assemblages

Joshua S. Madin[1]*, Michael McWilliam[1], Kate Quigley[2], Line K. Bay[2], David Bellwood[3], Thomas Bridge[4], Christopher Doropoulos[5], Leanne Fernandes[6], Peter Harrison[7], Andrew S. Hoey[3], Peter J. Mumby[8], Zoe T. Richards[9], Cynthia Riginos[8], David J. Suggett[10], Madeleine J. H. van Oppen[2,11]

1. Hawai□i Institute of Marine Biology, University of Hawai□i at Manoa, Kāneʻohe, Hawai□i, USA
2. Australian Institute of Marine Science, Townsville, Queensland, Australia
3. ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland, Australia
4. Museum of Tropical Queensland, Townsville, Queensland, Australia
5. CSIRO Oceans & Atmosphere, Brisbane, Queensland, Australia
6. Great Barrier Reef Marine Park Authority, Townsville, Queensland, Australia
7. Marine Ecology Research Centre at Southern Cross University, New South Wales, Australia
8. School of Biological Sciences, The University of Queensland, St. Lucia, Queensland, Australia
9. Coral Conservation and Research Group, Trace and Environmental DNA Laboratory, School of Molecular and Life Sciences, Curtin University, Bentley, Western Australia, Australia
10. University of Technology Sydney, Climate Change Cluster, Sydney, New South Wales, Australia
11. School of BioSciences, The University of Melbourne, Parkville, Victoria, Australia

*jmadin@hawaii.edu

Abstract

1. Humans have long sought to restore species but little attention has been directed at how best to do so for rich assemblages of foundation species that support ecosystems, like rainforests and coral reefs that are increasingly threatened by environmental change.

2. We developed a two-part triage process for selecting optimized sets of species for restoration. We demonstrated this process using phenotypic traits and ecological characteristics for reef building corals found along the east coast of Australia. Without clear linkages between phenotypic traits and ecosystem functions, the first part of the triage hedges against function loss by ensuring an even spread of life history traits. The second part hedges against future species losses by weighting species based on characteristics that are known to increase their ecological persistence to current environmental pressures—abundance, species range and thermal bleaching tolerance—as well as their amenability to restoration methods.

3. We identified sets of ecologically persistent and restorable species most likely to protect against functional loss by examining marginal returns in occupancy of phenotypic trait space per restored species.

4. We also compared sets of species with those from the southern-most accretional reef as well as a coral restoration program to demonstrate how trait space occupancy is likely to protect against local loss of ecosystem function.

5. *Synthesis and applications*. A quantitative approach to selecting sets of foundational species for restoration can inform decisions about ecosystem protection to guide and optimize future restoration efforts. The approach addresses the need to insure against unpredictable losses of ecosystem functions by investing in a wide range of phenotypes. Furthermore, the flexibility of the approach enables the functional goals of restoration to vary depending on environmental context, stakeholder values, and the spatial and temporal scales at which meaningful impacts can be achieved.

Keywords

Restoration; foundation species; triage; phenotypic traits; ecosystem function; hermatypic corals

Introduction

The rate and extent of environmental change experienced by contemporary ecosystems have resulted in major deviations from their historical state (Hobbs et al., 2011). Conservation alone may therefore no longer suffice to preserve biodiversity and ecosystem functions, and restoration is often considered a now required addition. The objective of ecosystem restoration is, through human intervention, to recover a disturbed or degraded ecosystem as far as possible towards some previous state. Interventions in the coastal and marine realm can be direct, such as propagation and field deployment of habitat builders through seeds (e.g., seagrass in Virginia – Orth et al., 2020), propagules (e.g., oysters in South Australia - Vanderklift et al., 2020), early recruits (e.g., kelp - Fredriksen et al., 2020; coral - Randall et al., 2020) or parts of adult tissues (e.g., coral gardening - Rinkevich 2014). Indirect interventions, such as physical stabilization of degraded reef structures are also possible (Ceccarelli et al., 2020).

To date, the augmentation or reintroduction of one or few species has been the most common approach, such as the restoration of the endangered Caribbean coral species *Acropora cervicornis* and *A. palmata* (Ladd et al., 2019), the reintroduction of the greater bilby in some parts of Australia (Lott et al., 2020), the gray wolf across parts of Europe and North America (Ripple et al 2014), and assisted colonization of the Tasmanian Devil to the Australian mainland (Brainard 2020). However, climate change is now affecting many assemblages of foundation species in most if not all the world's ecosystems—forests, kelp forest, and coral reefs—leading to a necessary broadening of focus of restoration activities to encompass more species and their contributions to ecosystem functioning (Brudvig & Mabry 2008; Ladouceur & Shackelford 2021).

Prioritizing sets of species for the restoration of biodiverse ecosystems is a challenging task, and different approaches for decision making have been used. For instance, some approaches focus on the roles that species play in providing particular ecosystem goods or services, including carbon storage in rainforests (Strassburg et al., 2020) or mangrove forests (Adame et al., 2014), or reef accretion on coral reefs for coastal protection (Bellwood et al., 2019). Another often used focus is on keystone species: species that maintain the organization, stability, and function of

their communities, and have disproportionately large, inimitable impacts on their ecosystems

(Hale et al., 2018). Alternatively, weedy pioneer species may quickly restore habitat functions

such as such as providing shelter or stabilizing substratum, such as the emphasis on fast growing

acroporids in coral gardening initiatives (Bostrom-Einarsson et al., 2020). However, the

objective selection of species based on defined ecological, functional and logistical criteria are

rare (Suding et al., 2004; Lamb 2018). Some examples exist for forest restoration (Meli et al.,

2013), and some have used linkages between phenotypic traits and ecosystem functions to select

species (Giannini et al., 2017; Rayome et al., 2019).

An overarching challenge is that restoration initiatives need to anticipate future ecosystem states

which are expected to be very different due to the escalating impacts of climate change (Rogers

et al., 2015; Gaitán☐Espitia & Hobday 2021). Faced with complex ecosystems, multiple threats

to biodiversity and limited funding, conservation practitioners must prioritize investment into

different management options, including restoration actions, and difficult decisions must be

made about which sets of species to allocate resources to (Game et al., 2018). Strategic decisions

must be taken about supporting those most likely to do better to improve future persistence and

resilience, and those that will struggle and potentially push them through a period of elevated and

prolonged stress; especially those that are already closer to their existing physiological limits,

like reef-building corals. Protecting habitat-forming species such as corals is imperative for

securing the ecological functions and socio-economic services they provide such as reef

building, habitat and food provisioning for commercially important species, primary production,

nutrient recycling, natural products, and social, cultural and recreational opportunities.

In this paper, we developed a two-part triage process for selecting sets of species for the

restoration or maintenance of foundational assemblages, for which we use reef-building corals as

an example. The first part centered on the restoration of ecosystem functions provided by

foundational assemblages, such as habitat engineering (Ellison et al., 2005) (Fig. 1). However,

rather than targeting specific ecosystem functions (Fig. 1a), we propose to minimize loss of

function by maximizing phenotypic variation (Fig. 1b) for several reasons. First, mechanistic

linkages between phenotypic traits of species and functions are poorly understood, especially for

coral reefs (Bellwood et al., 2019). Second, the species that support specific functions are likely

to differ depending on the situation (e.g., disturbance history) and site-specific conditions. Third,

maximizing life history variation minimizes the risk of wholesale species loss, because no

species is at a selective optimum in all situations and environments (Stearns 1992). Fourth,

important ecosystem functions, such as habitat engineering and ecological succession, tend to be

supported by a broad range of life history combinations (e.g., builders, fillers and cementers

[Goreau 1963]; weedy, competitive and stress-tolerant [Darling et al., 2012]). Finally, a bet

hedging approach also acts to increase phylogenetic diversity because many life history traits are

evolutionarily conserved (Westoby et al., 2002).

The second part of the triage process is to select species based on characteristics that make them

better equipped to avoid depletion and resist or recover from large-scale events like fires or

marine heatwaves (Fig. 1c). For example, species with small range sizes and small local

populations generally have a higher extinction risk (Staude et al., 2020), while species with

higher local abundances tend to bounce back faster following disturbance (Halford et al., 2004).

Often there are synergistic relationships—for example, extinction risk tends to be greater for

geographically limited and locally rare species (double jeopardy; Brown 1984)—but not always

(Hughes et al., 2014). Meanwhile, some species are more tolerant to disturbances and gradual

changes that are expected to become more frequent or more intense in the future through rapid

adaptation. Suitability of species to restoration can also be considered as a characteristic of

ecological persistence; albeit one requiring human intervention (Suggett et al., 2019). For

example, restoration may be facilitated by the use of species that can be easily propagated in the

laboratory via sexual reproduction, grown in nurseries and outplanted, or manipulated in the field

(Rinkevich 2020; Randall et al., 2020). Furthermore, some phenotypic traits and ecological

characteristics make some species better candidates for restoration than others. For instance, it is

easier to generate coral fragments from branching species than from massive species, therefore

most restoration efforts on coral reefs have historically focused on branching corals (Bostrom-

Einarsson et al., 2020). Similarly, restoring local areas with species that have large geographical

ranges will improve connectivity (Hock et al., 2017).

The overall aim was to develop a quantitative approach to select a set of *n* species for restoration,

recognizing that the goals of restoration can vary enormously across systems. The approach we

developed aims to maximize the probability of maintaining phylogenetic diversity and ecosystem

function, and therefore protect against a range of sensitivities to future stressors. We demonstrate

this approach with reef building corals found on the east coast of Australia, including the Great

Barrier Reef, a region that has declined severely over the past few decades (De'ath et al., 2012;

Hughes et al., 2017). Here, coral reef restoration activities are already underway to address the

local-scale depletion of coral populations (Boström-Einarsson et al., 2020; Howlett et al., in

review) and larger scale restoration interventions are in the research and development phase

(Anthony et al. 2020). Although the data and results are only illustrative at this stage, and further

scrutiny is required before making formal restoration decisions at appropriate and manageable

scales (McAfee et al., 2021), it provides a quantitative, reproducible and adaptable basis for

selecting species for restoration projects and restoration research.

Materials and methods

*Spectrum of life histories—*The first part of the process required collation of quantitative

phenotypic traits for species that capture as many life history trade-offs as possible (Gallagher et

al., 2020), such as acquisition-conservation and propagule size-number (Westoby et al., 2002).

For our demonstration, we use a dataset for 396 species found along the east coast of Australia

from McWilliam et al. (2018) with the following traits: growth rate, corallite width,

rugosity/branch spacing, surface area per unit volume, colony height, maximum colony

size/diameter, and skeletal density. The trait data enabled us to capture important dimensions of

species life history, ranging from fast to slow growth (Darling et al., 2012), fragile to robust

morphologies (Zawada et al., 2019), and small to large colonies that drives up colony fecundity

(Alvarez-Noriega et al., 2016). The trait space was calculated using a principal components

analysis (PCA) of the seven traits and is presented in Fig. 2a for the 396 species.

We also viewed the trait space through the lens of one ecosystem function, reef building, by

adopting Goreau's (1963) classification of reef species into builders, fillers and cementers; an

approach that has been supported by modern synthesis (González-Barrios & Alverez Filip 2018).

Within the McWilliam et al. (2018) trait space, builders were classified as species with the

highest values of size, height, and volume; fillers with largest values of size and rugosity; and

cementers with the largest sizes and smallest height values (Fig. 2a, red vectors). Ecosystem

functions such as reef building and habitat construction requires a broad range of life histories—

ranging from slow growing, potentially large builders (i.e., late successional species) to dense

skeleton, encrusting cementers to fast growing, morphological complex fillers (i.e., weedy or

early successional species). Other functions tend to require specific trait combinations (Fig. 2b).

For example, coral species with higher surface area to volume ratios, rugosities and growth rates

tend to generate habitat complexity that is reportedly important as fish habitats (Graham & Nash

2013) and have a high value for human uses, such as tourism and recreation, because they are

generally considered aesthetically appealing (Marshall et al., 2019).

While there are many definitions of phenotypic trait diversity (Villéger et al., 2008), our goal

under a hedging strategy was to evenly capture the largest area of trait space with the fewest

species, and therefore to ensure a spread of species along important trait dimensions. This goal

was accomplished by iteratively removing the species closest to other species in the two-

dimensional area defined by PC1 and PC2 (which captured about 70% total trait variation) until

a given number of species n remained. Selection could also happen at higher dimensions to

capture higher levels of variation. We tried two approaches to measure the proximity of species

using R (R Core Team 2021): (1) the areas of Voronoi cells using the *voronoi.mosaic* function in

the *tripack* package (Renka et al., 2020), and (2) the nearest neighbor distances using the *nndist*

function in the *spatstat* package and only considering the single closest species (i.e., k=1;

Baddeley et al., 2015). Proximity values were normalized at each iteration by dividing by the

maximum distance or area, depending on the approach. A grid-based approach was used to

assess trait diversity throughout the study, whereby a grid of a given resolution was

superimposed onto the trait space (e.g., a 5 by 5 cell grid is shown in Fig. 3a and a 10 by 10 cell

grid in Fig. S2). Trait diversity was the proportion of possible grid cells with at least one species;

redundancy was the mean number of species in occupied possible grid cells.

*Ecological persistence—*For the second part of the triage process we focused on three

characteristics (ecological abundance, geographic range size and thermal bleaching

susceptibility) broadly defined as factors contributing to ecological persistence of reef building

coral species. We acknowledge that these characteristics will depend on current taxonomic

designations that are currently being revised (Cowman et al., 2020). We used typical abundance
of species data from Veron (2000) and geographic distribution data from Hughes et al. (2013)
downloaded for the 396 species from the Coral Trait Database (Madin et al., 2016). Ecological
abundance was categorized by Veron (2000) as common, uncommon and rare, which we
normalized as 1, 0.5 and 0.25, respectively. Geographic extent was normalized by dividing the
range size of each species by the maximum range size for a species. Normalizing puts
characteristics on the same scale (i.e., between 0 and 1) (Fig. S1). Weightings can also be applied
to characteristics of ecological persistence to augment or diminish their importance, but we did
not do so here.

Thermal bleaching susceptibility is an increasingly relevant characteristic for coral ecological
persistence, but it is context dependent, highly variable, and poorly understood. Nonetheless, we
use the Coral Bleaching Index (BI) from Swain et al. (2016) to demonstrate how this variable
might be included in the triage analysis. BI is a value between 0 and 100, where higher values
correspond with more thermally vulnerable species. Therefore, we normalized BI by dividing by
100 and subtracting the result from 1 (i.e., species with values closer to 1 are more resistant to
bleaching based on Swain et al. [2016]). BI values were available at the species level for 212 of
the species, and therefore genus level BIs were used for the remainder of the analysis in order to
retain all 396 species.

Restoration of reef corals is a relatively new field (Hein et al., 2021), and so there is little long-
term knowledge of what makes species more or less amenable to the restoration process. A meta-
analysis of coral restoration studies ranked the use of coral growth forms in restoration projects
(Boström-Einarsson et al., 2020). While this ranking likely reflects a historical focus on coral
gardening (i.e., fragmentation) as well as specific situations, such as the demise of branching
Acropora species in the Caribbean, we nonetheless utilize this ranking as an index of species
amenability to restoration. Species growth form was downloaded from the Coral Trait Database
and species were ranked from 1 to 6: columnar (1), tabular (2), encrusting (3), foliose (4),
massive (5), and branching (6), which includes corymbose and digitate. This ranking was
normalized by dividing values by six.

Pairwise associations between PC1, PC2 and normalized ecological persistence variables for
species are shown in Fig. S1. Weighting species by characteristics was done by multiplying
normalized values. Species with values closer to 1—i.e., large ranges, common, and resistant to
bleaching—were considered ecologically persistent. To redirect focus of species selection to
vulnerable species, normalized variables were subtracted from 1 before proceeding. For example,
we also explored the triage process for species that were wide ranging and common, but
susceptible to bleaching.
251
Triage assessment—We compared triage results with two contrasting reef assemblages. The first
was the southern-most accretional reef assemblage along the east coast of Australia, Lord Howe
Island, built by approximately 50 coral species; however, we only include the 15 most common
species for our analysis, having greater than 5% mean cover (Table S1). Contrasts were made by
comparing trait combinations in the Lord Howe assemblage with those generated by the triage
process with the same numbers of species (n=15). The second set of species are those used for
out-planting for fragments by the Coral Nurture Program on the Great Barrier Reef throughout
the first two years of its planting activity across six northern GBR high value tourism reef sites
(~20,000 out-plants; August 2018 - April 2020; Howlett et al., in review). The program is run by
tourism operators, and so tends to focus on abundant, fast growing branching species with
recognised aesthetic values to the tourism industry. Contrasts were made by comparing trait
combinations of the 39 Coral Nurture Program species with those generated by the triage process
for n=39 species.
265
266   **Results**
267
268   The nearest neighbor distance method produced the most even spread of species in the trait space
269   (Fig. 2c, points) because the Voronoi area method could not calculate areas for peripheral points
270   (Fig. 2c, crosses) that were inadvertently retained during the iterative species removal process.
271
Selecting species based solely on maximizing species spread in trait space (i.e., via nearest
neighbor distances) resulted in levels of occupancy in the trait space that were significantly
greater than random (Fig. 3a, yellow line sitting largely above the shaded 95% confidence band).

Conversely, selecting species based solely on ecological persistence led to both the lowest
occupancy in the trait space (i.e., spread of morphological and life history traits) and the highest
levels of redundancy (Figs. 3a and 3b green curves, respectively). These levels of occupancy and
redundancy were not significantly different to randomized species selection (black curves in
Figs. 3a and 3b, respectively, where the green "persistence" curves are captured by the shaded
95% confidence bands). Selecting species based on both ecological persistence and trait diversity
resulted in some loss of trait diversity, but with levels of occupancy much greater than ecological
persistence alone (Fig. 3, red curves). Marginal returns in terms of occupancy per species was
high (approximately 5% per species) for the combined triage scenario (i.e., red curve, Fig. 3a) up
until $n=11$ species, medium (approximately 2% per species) between from $n=11$ to $n=28$, and
low (<0.3% per species) above $n=28$. The general patterns shown in Fig. 3 were robust to the cell
size of the grid used to calculate occupancy (Fig. S2); with the proviso that more species are
required to maintain specified levels of occupancy and redundancy for finer grids. For example,
the region of medium marginal returns occurred between $n=29$ and $n=55$ for the 10 by 10 cell
grid (Fig. S2A, red curve).
290
Based on marginal returns, the optimal number of species selected occurred somewhere between
11 and 28. Fig. 4 shows triage values when selecting $n=20$ species at different stages of the
process for two scenarios: (1) considering ecologically persistent species (large geographic
range, ecologically abundant and resistant to bleaching) (Fig. 4a) and (2) considering large
geographic ranges and ecologically abundant species that are susceptible to bleaching (Fig. 4b).
While there is some overlap among the triage stages, Fig. 4 demonstrated how focal species can
change throughout the selection process. It also shows the selection outcomes of switching
species that are better ecologically, but are difficult to restore based on our criteria, while
simultaneously retaining an even spread of species in the trait space. Fig. 2d contrasts triage for
20 species with and without consideration of trait diversity, illustrating that, while cementers are
captured either way, because they tend to be ecologically persistent species, while builders and
fillers are not captured to the same extent if focused solely on ecological persistence
characteristics.
304

The 15 species of Lord Howe Island assemblage showed lower trait diversity and higher
redundancy than would be produced by the triage process (Fig. 3, asterisks). Nonetheless, these
species occupied three highly distinctive trait combinations that largely overlap with the three
functional groups originally put forward by Goreau (1963) (Fig. 2e). The 39 species of the Coral
Nurture Program showed trait diversity markedly lower than expected by randomly selecting
species (Fig. 3a), resulting in high levels of redundancy (Fig. 3b). Indeed, the Lord Howe Island
assemblage showed similar levels of occupancy as the Coral Nurture Program with less than half
the species (Fig. 3a). Fig. 2f illustrates that these out-planted species are the fast growing, high
surface area to volume ratio species, with the exception of *Galaxea fascicularis* (Fig. 3e, red
arrow); they also only tend to capture Goreau's filler species category.

Discussion

The triage process developed here identified coral species for restoration projects based on both
ecologically beneficial characteristics and diversity of life history trait values. Selection based on
ecological persistence is important for hedging against future species loss, while phenotypic trait
diversity is important for hedging against both species loss and local ecosystem function loss.
The importance of individual species in ecosystem functions are poorly understood, particularly
for reef corals, and therefore our tactic was to prioritize an even spread of species across the trait
space rather than prioritizing particular phenotypic trait values or targeting regions in the trait
space (e.g., Fig. 2b). Even spread across the trait space (measured as gridded occupancy)
increased with the number of species selected; however, marginal returns declined at relatively
low numbers of species (~20 species, Fig. 3a). However, this depended on the resolution at
which the trait space was gridded (Fig. S2). How finely spread species should be across trait
space for preserving particular ecosystem functions remains an open question. Meanwhile, the
number of species that can be selected for a restoration project will ultimately depend upon
project goals, as well as resource and logistical constraints. The flexible triage process developed
here can serve as a framework for such decisions.

The species lists shown in Fig. 4 is an illustration of outputs that include one potential planning
process, hence the anonymous labelling, as they are based on a relatively narrow set of

ecological characteristic and phenotypic trait data. Future studies may build on our initial analyses by including a broad range of traits (rather than primarily morphological traits presented here) and imputing trait values for species with missing trait data (Fig. 1). For instance, certain *Acropora* species are underrepresented in Fig. 4a due to their high bleaching sensitivity. We used the colony growth form ranking by Bostrom-Einarsson et al. (2020) as the suitability for restoration trait. This trait is relevant for coral gardening approaches that fragment, grow and then out-plant adult colonies (Rinkevich 2014). However, for prioritization of species in restoration initiatives based on propagation of sexually produced coral stock (dela Cruz and Harrison 2020), we recommend traits other than growth form to be used for defining species' suitability for restoration, such as the mode of reproduction (large amounts of sexually produced offspring are easier to obtain from broadcast spawning as compared to brooding species; Doropoulos et al., 2019), high early life survivorship, and fecundity (species with higher fecundity can provide larger numbers of sexually produced offspring; Alvarez-Noriega et al., 2016). An obstacle for selecting species for restoration based on sexual reproduction is that the knowledge of husbandry is limited for many spawning coral species, and it will always be challenging to obtain high abundances of sexually produced offspring from brooding species (Randall et al., 2020). However, the triage process can be used to highlight where husbandry effort should be directed. For example, Fig. 4 contrasts the best set of species for restoration based on ecological persistence and trait diversity (middle columns in panels a and b); however, this list changes when also considering which of these species are amenable to fragmentation and gardening approaches (last columns). Ideally, coral husbandry and out-planting research should focus on reducing this mismatch (e.g., Baria☐Rodriguez et al., 2019).

Our understanding of the species and traits that drive reef functions and services is still emerging, placing severe limitations on our capacity to select foundation species for restoration. Many reviews have dealt with the importance of ecosystem functions and services on coral reefs (Harborne et al., 2006, Brandl et al., 2019; Woodhead et al., 2019), including in the context for reef restoration practices (Hein et al., 2021). However, for corals, the experimental and observational evidence linking species to functions is still limited (Brandl et al., 2019), perhaps because of the long timescales of reef-building. For example, recent restoration work from the Florida Keys achieved large increases in the cover of *Acropora cervicornis* but little benefit to

broader ecological functions (Ladd et al., 2019). Furthermore, in their blueprint for protecting

coral reef functions, Bellwood et al (2019) suggest a hierarchical approach, recognizing that all

functions are not equal, and that priority functions are likely to be context specific and should be

defined in specific restoration objectives. Selecting species based on functions is therefore likely

to vary across systems depending on the specific environment, and the values of the local

stakeholders. Our triage approach addresses both these limitations by hedging against loss of

range of function by capturing a wide range of morphological and life history traits, while

simultaneously enhancing the probability that these traits persist in a range of environments.

Given the uncertainty inherent to all restoration efforts, we suggest that better knowledge of

functions and a clear vision of the goals of restoration is likely to enhance rather than supersede

this triage approach.

Restoration efforts should explicitly consider the capacity of species to persist in future

environments. An increasingly visible pattern is that there are species that are likely to persist in

future predicted environments ('winners') and those that are likely to decline in future

environments ('losers') (Adam et al., 2021). Whether to protect the hardy or the vulnerable is a

widespread debate throughout conservation biology. For example, when selecting protected

areas, the decision to protect areas of imminent threat or resistant areas generating the best

returns is highly context-dependent (Sacre et al., 2019). Similarly, whether to select winners or

losers for restoration is likely to depend on the severity of the threat to the ecosystem, and the

traits and functions that are most critical to ecological integrity. Indeed, a species-oriented focus

on restoration is likely to favor rare or depleted species with low persistence, while ecosystem-

based restoration is likely to favor foundation species that are dominant and therefore drive

ecosystem functions. The triage approach developed here is specifically designed so that both

these viewpoints can be incorporated.

This paper's goal was to develop and present a simple and flexible process for decision making

around target species for restoration using real and interpolated data. However, the data and

results presented here should not be used for decisions without further consideration,

consultation and analysis. For example, a multi-criteria decision-making approach is one

pathway (Gouezo et al., 2021). Expert elicitation by coral reef scientists is needed but

inadequate, and this process should also include other stakeholders (managers, tourism
operators) and pivotally, First Nations people. Moreover, the value of restoration as a
management intervention per se must be evaluated against or complement alternative tools, such
as protected areas, fish aggregation devices, and artificial structures, and the outcomes of such
assessments vary from one ecosystem function to another (Rogers et al 2015). The ultimate goals
of restoration, and particularly the ecosystem functions being targeted, should be based on the
scale at which restoration has the greatest impact. Moreover, the success of coral restoration
hinges upon addressing local and global actions that facilitate natural coral growth and
reproduction, such as regulating climate, overharvesting and land-use.
407

408 **Authors contributions**
409

410 All authors conceived the idea during a working group meeting organized by MJHvO and KQ.
411 JSM and MM developed the idea, gathered data and ran analyses. JSM, MM, KQ and MJHvO
412 wrote the first draft. All authors critically revised drafts and added intellectual content.
413

414 **Acknowledgements**
415

416 The workshop was funded by the Australian Research Council Laureate Fellowship
417 FL180100036 to MJHvO. CD and LKB acknowledges the Reef Restoration and Adaptation
418 program. ZR acknowledges Australian Research Council LP160101508.

420 **Conflict of Interest**
421

422 The authors declare that they have no known competing financial interests or personal
423 relationships that could have appeared to influence the work reported in this paper.
424

425 **Data availability statement**
426

All data and code are available at https://github.com/jmadinlab/species_choice

References

Adam, A.A., Garcia, R.A., Galaiduk, R., Tomlinson, S., Radford, B., Thomas, L., & Richards,
Z.T. (2021). Diminishing potential for tropical reefs to function as coral diversity
strongholds under climate change conditions. *Diversity & Distributions*, 27, 2245-2261.

Adame, M.F., Teutli, C., Santini, N.S., Caamal, J.P., Zaldívar-Jiménez, A., Hernández, R., &
Herrera-Silveira, J.A. (2014). Root Biomass and Production of Mangroves Surrounding a
Karstic Oligotrophic Coastal Lagoon. *Wetlands*, 34, 479–488.

Álvarez-Noriega, M., Baird, A.H., Dornelas, M., Madin, J.S., Cumbo, V.R., & Connolly, S.R.
(2016). Fecundity and the demographic strategies of coral morphologies. *Ecology*, 97,
3485–3493.

Anthony, K.R.N., Helmstedt, K.J., Bay, L.K., Fidelman, P., Hussey, K.E., Lundgren, P., Mead,
D., McLeod, I.M., Mumby, P.J., Newlands, M., Schaffelke, B., Wilson, K.A., Hardisty,
P.E. (2020) Interventions to help coral reefs under global change—A complex decision
challenge. PLOS ONE, 15, e0236399.

Baddeley, A., Rubak, E., & Turner, R. (2015). *Spatial Point Patterns: Methodology and
Applications with R*. London: Chapman and Hall/CRC Press.

Baria☐Rodriguez, M.V., dela Cruz, D.W., Dizon, R.M., Yap, H.T., & Villanueva R.D. (2019).
Performance and cost☐effectiveness of sexually produced Acropora granulosa juveniles
compared with asexually generated coral fragments in restoring degraded reef areas.
Aquatic Conservation: Marine and Freshwater Ecosystems, 29, 891-900.

Bellwood, D.R., Pratchett, M.S., Morrison, T.H., Gurney, G.G., Hughes, T.P., Álvarez-Romero,
J.G., Day, J.C., Grantham, R., Grech, A., Hoey, A.S., Jones, G.P., Pandolfi, J.M., Tebbett,
S.B., Techera, E., Weeks, R., & Cumming, G.S. (2019). Coral reef conservation in the
Anthropocene: Confronting spatial mismatches and prioritizing functions. *Biological
Conservation*, 236, 604–615.

Boström-Einarsson, L., Babcock, R.C., Bayraktarov, E., Ceccarelli D., Cook, N., Ferse, S.C.A.,
Hancock, B., Harrison, P., Hein, M., Shaver, E., Smith, A., Suggett, D., Stewart-Sinclair,
P.J., Vardi, T., McLeod, I.M. (2020). Coral restoration – A systematic review of current
methods, successes, failures and future directions. *PLoS ONE*, 15,.e0226631.

Brainard, J. (2020). Tasmanian devil reintroduced. *Science*, 370,268-268.

Brandl, S.J., Rasher, D.B., Côté, I.M., Casey, J.M., Darling, E.S., Lefcheck, J.S., Duffy, J.E.
(2019). Coral reef ecosystem functioning: eight core processes and the role of biodiversity.
Front Ecol Environ, 17, 445–454.

Brown, J.H. (1984). On the Relationship between Abundance and Distribution of Species. The
American Naturalist, 124, 255–279.

Brudvig, L.A., Mabry, C.M. (2008). Trait-Based Filtering of the Regional Species Pool to Guide
Understory Plant Reintroductions in Midwestern Oak Savannas, U.S.A. *Restoration*
Ecology, 16, 290-304.

Ceccarelli, D.M., McLeod, I.M., Boström-Einarsson, L., Bryan, S.E., Chartrand, K.M., Emslie,
M.J., Gibbs, M.T., Gonzalez Rivero, M., Hein, M.Y., Heyward, A., Kenyon, T.M., Lewis,
B.M., Mattocks, N., Newlands, M., Schläppy, M.-L., Suggett, D.J., Bay, L.K. (2020).
Substrate stabilisation and small structures in coral restoration: State of knowledge, and
considerations for management and implementation. *PLOS ONE*, 15, e0240846.

Cowman, P.F., Quattrini, A.M., Bridge, T.C.L., Watkins-Colwell, G.J., Fadli, N., Grinblat, M.,
Roberts, T.E., McFadden, C.S., Miller, D.J., Baird, A.H. (2020). An enhanced target-
enrichment bait set for Hexacorallia provides phylogenomic resolution of the staghorn
corals (Acroporidae) and close relatives. Molecular Phylogenetics and Evolution, 153,
106944.

Darling, E.S., Alvarez-Filip, L., Oliver, T.A., McClanahan, T.R., Côté, I.M. (2012). Evaluating
life-history strategies of reef corals from species traits. *Ecology Letters*, 15,1378–1386.

De'ath G, Fabricius KE, Sweatman H, Puotinen M (2012) The 27-year decline of coral cover on
the Great Barrier Reef and its causes. Proc Natl Acad Sci USA 109:17995-17999

dela Cruz DW, Harrison PL (2020) Enhancing coral recruitment through assisted mass
settlement of cultured coral larvae. PLOS ONE 15:e0242847

Doropoulos C, Vons F, Elzinga J, ter Hofstede R, Salee K, van Koningsveld M, Babcock RC
(2019) Testing Industrial-Scale Coral Restoration Techniques: Harvesting and Culturing
Wild Coral-Spawn Slicks. Front Mar Sci 6:658

Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, Ford CR, Foster DR, Kloeppel BD,
Knoepp JD, Lovett GM, Mohan J, Orwig DA, Rodenhouse NL, Sobczak WV, Stinson KA,
Stone JK, Swan CM, Thompson J, Von Holle B, Webster JR (2005) Loss of foundation

489 species: consequences for the structure and dynamics of forested ecosystems. Frontiers in
490  Ecology and the Environment 3:479–486.

491 Fredriksen S, Filbee-Dexter K, Norderhaug KM, Steen H, Bodvin T, Coleman MA, Moy F,
492  Wernberg T (2020) Green gravel: a novel restoration tool to combat kelp forest decline.
493  Scientific reports 10:3983

494 Gaitán⬜Espitia JD, Hobday AJ (2021) Evolutionary principles and genetic considerations for
495  guiding conservation interventions under climate change. Glob Change Biol 27:475–488

496 Gallagher RV, Falster DS, Maitner BS, Salguero-Gómez R, Vandvik V, Pearse WD, Schneider
497  FD, Kattge J, Poelen JH, Madin JS, Ankenbrand MJ, Penone C, Feng X, Adams VM,
498  Alroy J, Andrew SC, Balk MA, Bland LM, Boyle BL, Bravo-Avila CH, et al. (2020) Open
499  Science principles for accelerating trait-based science across the Tree of Life. Nature
500  Ecology & Evolution 4:294–303

501 Game ET, Tallis H, Olander L, Alexander SM, Busch J, Cartwright N, Kalies EL, Masuda YJ,
502  Mupepele A-C, Qiu J, Rooney A, Sills E, Sutherland WJ (2018) Cross-discipline evidence
503  principles for sustainability policy. Nat Sustain 1:452–454

504 Giannini TC, Giulietti AM, Harley RM, Viana PL, Jaffe R, Alves R, Pinto CE, Mota NFO,
505  Caldeira CF, Imperatriz-Fonseca VL, Furtini AE, Siqueira JO (2017) Selecting plant
506  species for practical restoration of degraded lands using a multiple-trait approach. Austral
507  Ecology 42:510–521

508 González-Barrios FJ, Álvarez-Filip L (2018) A framework for measuring coral species-specific
509  contribution to reef functioning in the Caribbean. Ecological Indicators 95:877–886

510 Goreau TF (1963) Calcium carbonate deposition by coralline algae and corals in relation to their
511  roles as reef⬜builders. Comparative Biology of Calcified Tissue, Volume 109, Issue1 pp.
512  127-167.

513 Gouezo M, Fabricius K, Harrison P, Golbuu Y, Doropoulos C (2021) Optimizing coral reef
514  recovery with context-specific management actions at prioritized reefs. Journal of
515  Environmental Management 295:113209

516 Graham NAJ, Nash KL (2013) The importance of structural complexity in coral reef ecosystems.
517  Coral Reefs 32:315–326

518 Hale SL, Koprowski JL (2018) Ecosystem-level effects of keystone species reintroduction: a
519  literature review: Effects of keystone species reintroduction. Restor Ecol 26:439–445

520   Halford A, Cheal AJ, Ryan D, Williams DMcB (2004) Resilience to large-scale disturbance in
521   coral and fish assemblages on the Great Barrier Reef. Ecology 85:1892–1905

522   Harborne AR, Mumby PJ, Micheli F, Perry CT, Dahlgren CP, Holmes KE, Brumbaugh DR
523   (2006) The Functional Value of Caribbean Coral Reef, Seagrass and Mangrove Habitats to
524   Ecosystem Processes. Advances in Marine Biology. Elsevier, pp 57–189

525   Hein MY, Vardi T, Shaver EC, Pioch S, Boström-Einarsson L, Ahmed M, Grimsditch G,
526   McLeod IM (2021) Perspectives on the Use of Coral Reef Restoration as a Strategy to
527   Support and Improve Reef Ecosystem Services. Front Mar Sci 8:618303

528   Hobbs RJ, Hallett LM, Ehrlich PR, Mooney HA (2011) Intervention Ecology: Applying
529   Ecological Science in the Twenty-first Century. BioScience 61:442–450

530   Hock K, Wolff NH, Ortiz JC, Condie SA, Anthony KRN, Blackwell PG, Mumby PJ (2017)
531   Connectivity and systemic resilience of the Great Barrier Reef. PLoS Biol 15:e2003355

532   Howlett L, et al. (in review) Adoption of coral propagation and out-planting via the tourism
533   industry to advance site stewardship on the northern Great Barrier Reef.

534   Hughes TP, Bellwood DR, Connolly SR, Cornell HV, Karlson RH (2014) Double Jeopardy and
535   Global Extinction Risk in Corals and Reef Fishes. Current Biology 24:2946–2951

536   Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH,
537   Babcock RC, Beger M, Bellwood DR, Berkelmans R (2017) Global warming and recurrent
538   mass bleaching of corals. Nature 543:373-377

539   Hughes, T. P., Connolly, S. R., Keith, S. A. (2013) Geographic ranges of reef corals (Cnidaria:
540   Anthozoa: Scleractinia) in the Indo-Pacific. Ecology 94, 1659

541   Ladd MC, Burkepile DE, Shantz AA (2019) Near‐term impacts of coral restoration on target
542   species, coral reef community structure, and ecological processes. Restor Ecol 27:1166–
543   1176

544   Ladouceur E, Shackelford N (2021) The power of data synthesis to shape the future of the
545   restoration community and capacity. Restor Ecol 29:

546   Lamb D (2018) Undertaking large-scale forest restoration to generate ecosystem services:
547   Landscape restoration and ecosystem services. Restor Ecol 26:657–666

548   Lott MJ, Wright BR, Kemp LF, Johnson RN, Hogg CJ (2020) Genetic Management of Captive
549   and Reintroduced Bilby Populations. J Wildl Manage 84:20-32

Madin JS, Anderson KD, Andreasen MH, Bridge TCL, Cairns SD, Connolly SR, Darling ES,
Diaz M, Falster DS, Franklin EC, Gates RD, Hoogenboom MO, Huang D, Keith SA,
Kosnik MA, Kuo C-Y, Lough JM, Lovelock CE, Luiz O, Martinelli J, et al. (2016) The
Coral Trait Database, a curated database of trait information for coral species from the
global oceans. Scientific Data 3:160017

Marshall N, Marshall P, Curnock M, Pert P, Smith A, Visperas B (2019) Identifying indicators
of aesthetics in the Great Barrier Reef for the purposes of management. PLoS ONE
14:e0210196

McAfee D, Costanza R, Connell SD (2021) Valuing marine restoration beyond the 'too small
and too expensive.' Trends in Ecology & Evolution 36:968–971

McWilliam M, Hoogenboom MO, Baird AH, Kuo C-Y, Madin JS, Hughes TP (2018)
Biogeographical disparity in the functional diversity and redundancy of corals. Proceedings
of the National Academy of Sciences 115:3084–3089

Meli P, Martínez-Ramos M, Rey-Benayas JM (2013) Selecting Species for Passive and Active
Riparian Restoration in Southern Mexico: Selecting Species for Riparian Restoration.
Restor Ecol 21:163–165

Orth RJ, Lefcheck JS, McGlathery KS, Aoki L, Luckenbach MW, Moore KA, Oreska MPJ,
Snyder R, Wilcox DJ, Lusk B (2020) Restoration of seagrass habitat leads to rapid
recovery of coastal ecosystem services. Science Advances 6:eabc6434

R Core Team (2021). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Randall CJ, Negri AP, Quigley KM, Foster T, Ricardo GF, Webster NS, Bay LK, Harrison PL,
Babcock RC, Heyward AJ (2020) Sexual production of corals for reef restoration in the
Anthropocene. Mar Ecol Prog Ser 635:203-232

Rayome D, DiManno N, Ostertag R, Cordell S, Fung B, Vizzone A, Pante P, Tate R (2019)
Restoring Ecosystem Services Tool (REST): a Computer Program for Selecting Species for
Restoration Projects Using a Functional-Trait Approach. 58

Renka RJ, Gebhardt A, Eglen S, Zuyev S, White D (2020) tripack: Triangulation of Irregularly
Spaced Data. R package version 1.3-9.1

Rinkevich B (2014) Rebuilding coral reefs: does active reef restoration lead to sustainable reefs?
Current Opinion in Environmental Sustainability 7:28-36

Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, Berger J,
Elmhagen B, Letnic M, Nelson MP, Schmitz OJ, Smith DW, Wallach AD, Wirsing AJ
(2014) Status and Ecological Effects of the World's Largest Carnivores. Science
343:1241484

Rogers A, Harborne AR, Brown CJ, Bozec Y-M, Castro C, Chollett I, Hock K, Knowland CA,
Marshell A, Ortiz JC, Razak T, Roff G, Samper-Villarreal J, Saunders MI, Wolff NH,
Mumby PJ (2015) Anticipative management for coral reef ecosystem services in the 21st
century. Glob Change Biol 21:504–514

Sacre E, Bode M, Weeks R, Pressey RL (2019) The context dependence of frontier versus
wilderness conservation priorities. Conservation Letters 12:

Staude IR, Navarro LM, Pereira HM (2020) Range size predicts the risk of local extinction from
habitat loss. Global Ecol Biogeogr 29:16–25

Stearns, S. (1992). *The Evolution of Life Histories*. Oxford ; New York: Oxford University Press.

Strassburg BBN, Iribarrem A, Beyer HL, Cordeiro CL, Crouzeilles R, Jakovac CC, Braga
Junqueira A, Lacerda E, Latawiec AE, Balmford A, Brooks TM, Butchart SHM, Chazdon
RL, Erb K-H, Brancalion P, Buchanan G, Cooper D, Díaz S, Donald PF, Kapos V, Leclère
D, Miles L, Obersteiner M, Plutzar C, de M. Scaramuzza CA, Scarano FR, Visconti P
(2020) Global priority areas for ecosystem restoration. Nature 586:724–729

Suding KN, Gross KL, Houseman GR (2004) Alternative states and positive feedbacks in
restoration ecology. Trends in Ecology & Evolution 19:46–53

Suggett DJ, Camp EF, Edmondson J, Boström☐Einarsson L, Ramler V, Lohr K, Patterson JT
(2019) Optimizing return☐on☐effort for coral nursery and outplanting practices to aid
restoration of the Great Barrier Reef. Restor Ecol 27:683–693

Swain TD, Vega☐Perkins JB, Oestreich WK, Triebold C, DuBois E, Henss J, Baird A, Siple M,
Backman V, Marcelino L (2016) Coral bleaching response index: a new tool to standardize
and compare susceptibility to thermal bleaching. Glob Change Biol 22:2475–2488

Vanderklift MA, Doropoulos C, Gorman D, Leal I, Minne AJP, Statton J, Steven ADL,
Wernberg T (2020) Using Propagules to Restore Coastal Marine Ecosystems. Front Mar
Sci 7:724

Veron, J. E. N. (2000) Corals of the World. Australian Institute of Marine Science and CCR Qld
Pty Ltd.

Villéger S, Mason NWH, Mouillot D (2008) New multidimensional functional diversity indices
for a multifaceted framework in functional ecology. Ecology 89:2290–2301
Westoby M, Falster DS, Moles AT, Vesk PA, Wright IJ (2002) Plant Ecological Strategies:
Some Leading Dimensions of Variation Between Species. Annual Review of Ecology and
Systematics 33:125–159
Woodhead, A.J., Hicks, C.C., Norström, A.V., Williams, G.J., Graham, N.A.J. (2019) Coral reef
ecosystem services in the Anthropocene. Funct Ecol, 1365-2435.13331.
Zawada, K.J.A., Dornelas, M., Madin, J.S. (2019) Quantifying coral morphology. Coral Reefs
38,1281–1292.

Figure 1. Selecting sets of species for restoration with a two-part triage approach. The orange panel shows ecological functions that are important or threatened. If mechanistic linkages between life history traits and functions are established, then triage could focus on specific high-priority functions (left blue panel). Alternatively, a bet hedging approach that maximizes the diversity of functional and life history traits increases the likelihood that multiple functions are performed across a range of environments (right blue panel). Next, the species supporting functions are narrowed down further to include only those with characteristics that enhance or diminish their chances of persisting (green panel), resulting in different lists of species (yellow panel).

Figure 2. Phenotypic trait space is represented as the first two principal component axes for 396 east coast Australia coral species. (**a**) The trait loadings: growth rate (GR), corallite width (CW), rugosity/branch spacing (R), surface area per unit volume (SAV), colony height (CH), maximum colony size/diameter (MCS), and skeletal density (SD). Red vectors are Goreau's (1963) categories of essential reef builders. (**b**) An overlay illustrating where several functions approximately lie in trait space. Some functions require species broadly across the space, whereas others only a limited region of the space. (**c**) Selections of evenly spread species (n=20) calculated using nearest neighbor distances (points) and Voronoi cell areas (crosses and dashed line). (**d**) Selections based on ecological persistence characteristics only (crosses and grey shading) and then with the addition of trait diversity weighting (points) demonstrating the two-part triage. (**e**) The 15 common Lord Howe Island species (crosses) and 15 species selected using the triage approach (points). (**f**) The 39 Coral Nurture Program species (crosses) and 39 species selected using the triage approach (points). The red arrow shows an outlier species, *Galaxea fascicularis*.

Figure 3. (**a**) Proportion occupancy and (**b**) redundancy in trait space as a function of number of

species n using a 5 by 5 cell grid (inset in panel **a**). Regions of high, medium and low marginal

returns delineated with dotted vertical lines. Included are symbols that show trait diversity and

redundancy for common species at Lord Howe Island (*n*=15) and for the Coral Nurture Program

(*n*=39). The grey shaded region shows 95% CIs for randomized species selection.

Figure 4. Species selected (*n*=20) at different stages of the triage process when focusing on (**a**)
ecological persistent species in terms of range size and local abundance, bleaching resistance,
and suitability for restoration, and (**b**) species that have a large range size and are ecologically
abundant, vulnerable to bleaching, and suitable for restoration. Values (and heat colors)
correspond with a species triage score at successive stages. Asterisks (*) denote species found at
Lord Howe Island; pluses (+) denote Coral Nurture Program species from Howlett et al. (in
review).