Transcriptomics analysis reveals intracellular mutual regulation of coral-zooxanthella holobionts

Yunchi Zhu[1], Xin Liao[2], Tingyu Han[1], J.-Y. Chen[3], Chunpeng He[1*] and Zuhong Lu[1*]

[1] State Key Laboratory of Bioelectronics, Southeast University, Nanjing, Jiangsu, China
[2] Guangxi Key Lab of Mangrove Conservation and Utilization, Guangxi Mangrove Research Center, Beihai, Guangxi, China
[3] Nanjing Institute of Paleontology and Geology, Nanjing, China

* Corresponding author(s):
E-mail: cphe@seu.edu.cn; zhlu@seu.edu.cn

Abstract

Corals should make excellent models for cross-kingdom regulation research because of their natural animal-photobiont holobiont composition, yet a lack of studies and experimental data restricts their use. Here we integrate new full-length transcriptomes and small RNAs of four common reef-building corals with the published *Symbiodinium C1* genome to gain deeper insight into mutual gene regulation in coral-zooxanthella holobionts. We show that zooxanthellae secrete miRNA to downregulate rejection from host coral cells, and that a potential correlation exists between miRNA diversity and physiological activity. Convergence of these holobionts' biological functions in different species is also revealed, which implies the low gene impact on bottom ecological niche organisms. This work provides evidence for the early origin of cross-kingdom regulation as a mechanism of self-defense autotrophs can use against heterotrophs, sheds more light on coral-zooxanthella holobionts, and contributes valuable data for further coral research.

Introduction

miRNA is a type of small, single-stranded non-coding RNA found in plants, animals, and some viruses, and that functions in RNA silencing and post-transcriptional regulation of gene expression (1). miRNA is recognized to be involved in various regulation pathways, such as cross-cell regulation, cross-organ regulation, and even cross-generation regulation, however, whether cross-kingdom regulation occurs has been controversial. In 2012, Zhang and colleagues first proposed this concept after they identified stable plant miRNAs in several human and animal organs (2). They argued that intaken plant miRNAs could remain active and exert biological effects in animal tissues. A great deal of subsequent research has supported their views, for example, (3) revealed a correlation between fruit juice intake and fruit miRNA concentration in human serum; one honeysuckle miRNA (4) and two rapeseed miRNAs (5) were observed to be effectively absorbed by mice; (6) integrated ddPCR and transcriptome analyses to explore silkworms' absorption of mulberry miRNAs; etc. Some "avid" supporters of the idea of cross-kingdom miRNA activity advanced the medicinal value of plant miRNA against IAV (4), tumors (7), and even SARS-CoV-2 (8). Nevertheless, ever since the concept was put forward, the doubts continue to be voiced (9, 10). The proposers and supporters of cross-kingdom regulation have been working on explaining miRNA absorption mechanisms in detail in order to refute these objections (11), while not all research of interorganismal miRNA exchange is focused on food-intake-derived regulation, such as (12) studied the role of dodder miRNA in the parasitic process.

Most studies on cross-kingdom miRNA activity study humans or other complex animals as experimental subjects and look into whether plant miRNAs can affect animals' physiological functions. From our perspective, these models may be too complex in terms of behavior and physiology to support accurate conclusions about the origins or even the existence of such small molecules. Instead, we suggest a focus on more basal species exhibiting less complex behavior to explore the origins and details of the dominant regulation mechanisms of miRNA, which could lay the theoretical and practical foundation for studies of more complex organisms. We recommend coral as a potential animal model for cross-kingdom regulation research. Known co-evolutionary patterns exist for coral microbial communities and coral phylogeny. Corals obtain the majority of their energy and nutrients from photosynthetic unicellular dinoflagellates of the genus *Symbiodinium*, commonly known as symbiotic zooxanthellae. Zooxanthellae living within corals' tissues not only supply their hosts with the products of photosynthesis but also aid in coral calcification and waste removal. Coral-zooxanthella holobionts are naturally formed "animal-plant mixtures," that offer the advantages of simple behavior and ease of observation relative to more complex animal or

plant models. It is not difficult to assume that cross-kingdom regulation, regarded by its proposers as evidence of long-term co-evolution between photosynthetic organisms and animals, may play an important role in these holobionts.

However, in contrast to more common model organisms, there is a lack of public omics data from reef-building corals, not commensurate with their important status in the marine ecosystem and scientists' sharp focus on them. At the end of 2020, only 24 coral genomes at a medium assembly level are available on NCBI, while transcriptomes and proteomes are even more lacking. Many other well-known databases, such as miRBase (13) and KEGG (14), contain very little or even no coral data. Various factors, including geographic location, technical limitations, and experimental challenges, contribute to the limited number of coral studies serving as data sources, resulting in a "data gap." This lack of data deprives bioinformaticians of the opportunity to conduct dry lab work on corals, adding to a vicious circle exerting further negative effects on coral research. In-depth research underpinned by high quality bio-data is in urgent demand to break this cycle.

In order to explore intracellular mutual regulation of coral-zooxanthella holobionts, we sequenced the full-length transcriptomes and small RNAs of four common and frequently dominant reef-building corals, including *Acropora muricata*, *Montipora foliosa*, *Montipora capricornis* and *Pocillopora verrucosa*. For each species, we also performed Illumina sequencing to quantify gene expression. Every full-length transcript was annotated with Nr, Nt, Pfam, KOG, Swiss-Prot, GO and KEGG to distinguish coral and zooxanthella genes. A public *Symbiodinium C1* genome was employed to identify zooxanthella miRNAs, whose targeting sites were predicted later. Finally, highly expressed coral genes, zooxanthella genes and miRNA-target genes were selected for enrichment analysis. To a certain extent, zooxanthellae become organelles when they are inside coral cells, limiting the expression of most of their genes. Some of the miRNAs secreted by zooxanthellae tend to target immune-related genes in corals, suggesting that cross-kingdom regulation be a way for the zooxanthellae to prevent rejection by their hosts. Results also demonstrate that coral genes turn out to be functionally similar despite structural differences among coral species. These results suggest that the experimental subject selection for a coral model organism may be flexible, undoubtedly a benefit for researchers. This work sheds light into the coral-zooxanthellae holobiont, an ancient association in which an early example of cross-kingdom regulation is found. We also curate our data in a full-service online database to support further research (http://coral.bmeonline.cn).

Results

All raw sequencing data are available on NCBI (**Table 1**). 28 transcriptomes have been sequenced in total. For each species, the biological replicate numbers of full-length SMRT transcriptome sequencing, small RNA sequencing and Illumina RNA-seq sequencing are designated as 1:3:3 in order to ensure accuracy.

Table 1. Raw sequencing data on NCBI SRA. There are 28 transcriptomes in total. Each sample represents one biological replicate.

Species	Full-length transcript	Small RNA	Illumina RNA-seq
Acropora muricata	SRR9613488	SRR13442147	SRR12904786
		SRR13442146	SRR12904785
		SRR13442145	SRR12904784
Montipora foliosa	SRR9129316	SRR13442144	SRR12904783
		SRR13442143	SRR12904782

		SRR13442142	SRR12904781
		SRR13442141	SRR12904780
Montipora capricornis	SRR9129315	SRR13442150	SRR12904792
		SRR13442149	SRR12904791
		SRR13442152	SRR12904794
Pocillopora verrucosa	SRR9129314	SRR13442151	SRR12904793
		SRR13442148	SRR12904787

Transcriptome sequencing and data processing

The SMRT-sequencing technology was performed with the PacBio Sequel II platform to acquire offline polymer read bases of full-length transcriptomes using SMRTlink v. 7.0 software. The offline polymer read bases of *A. muricata*, *M. foliosa*, *M. capricornis* and *P. verrucosa* samples are 12.6G, 20.16G, 16G and 14.97G respectively (**File S1** for more details). The subreads, CCSs, FLNCs, consensus sequences, corrected consensus reads and unigenes are shown in **Table 2**, which also covers all information revealed in subsequent analyses.

Table 2. Sequencing data statistics of four full-length coral transcriptomes.

Sample name	*A. muricata*	*M. foliosa*	*M. capricornis*	*P. verrucosa*
Subread base (G)	12.37	19.82	15.62	14.61
Subread (number)	5,034,975	7,847,499	7,535,422	7,357,796
Average subread length (Nt)	2,456	2,526	2,073	1,986
CCS (number)	358,971	403,628	354,221	317,878
FLNC (number)	264,098	291,254	248,892	236,923
Consensus read (number)	22,925	23,067	18,354	20,696
Corrected consensus (number)	22,925	23,067	18,354	20,696
Unigene (number)	14,926	14,040	11,410	12,982

The full-length transcriptomes were annotated with the aid of Nr, Nt, Pfam, KOG, Swiss-Prot, GO and KEGG databases, and related unigene statistics are shown in **Fig 1.** In function-related databases, over 70% of unigenes get GO and KEGG annotations, paving the way for functional enrichment analysis. In protein-related databases, more than 96% of the unigenes of investigated corals are annotated in Nr, the basic protein primary sequence database. Such wide coverage indicates that Nr could be utilized for gene-ID mapping among different corals. As illustrated in **Fig S1**, Nr annotations map most genes of the four corals to the cnidarians *A. digitifera*, *E. pallida* and *N. vectensis,* with more than 85% overlap. Acceptable as it is, this result still reflects the lack of coral data for those annotations rarely correspond with the actual species in question.

Fig 1. Statistics of gene functional annotation. Photos of investigated corals are listed in the upper left corner.

According to Nr annotation, there are 43 *Symbiodinium* genes in *A. muricata*, 159 in *M. foliosa*, 45 in *M. capricornis,* and 129 in *P. verrucosa*, less than 0.07% of the total gene number. These numbers should be described as significantly few, although errors in annotation databases could account for this phenomenon to some extent. This result is similar to the condition of several well-researched organelles, such as mitochondria and chloroplasts, that are hypothesized to have become "intracellular slaves" during the "war" between archaebacteria and proteobacteria one billion years ago, working all the time but hardly expressing any genes of their own. It is not unreasonable to wonder whether zooxanthellae have undergone an analogous situation in the holobionts.

The RNA-seq sequencing was performed on the Illumina HiSeq X Ten platform. The total read bases are 22.1G (7.8G + 7.2G + 7.1G) from *A. muricata*, 29G (9.9G + 8.4G + 10.7G) from *M. foliosa*, 23.2G (7G + 7.1G + 9.1G) from *M. capricornis* and 23.1G (6.9G + 7.5G + 8.7G) from *P. verrucosa*.

Small RNAs were sequenced with the Illumina HiSeq X Ten platform as well. The raw read bases are 4.412G (1.548G + 1.359G + 1.505G) from *A. muricata*, 3.865G (1.275G + 1.053G + 1.537G) from *M. foliosa*, 3.295G (1.177G + 0.893G + 1.225G) from *M. capricornis* and 6.235G (2.67G + 1.494G + 2.071G) from *P. verrucosa* (**File S2** for more details). The raw read, clean read, unique sRNA, rRNA read, tRNA read, snRNA read, snoRNA read, known coral miRNA, novel coral miRNA and novel Symbiodinium miRNA are shown in **Table 3**.

Table 3. Sequencing data statistics of four coral small RNAs. For each species, miRNA identification results of three biological replicates were combined.

Sample name	A. muricata	M. foliosa	M. capricornis	P. verrucosa
	30,957,494	25,501,875	23,546,531	53,396,078
Raw read (number)	27,178,173	21,065,650	17,864,735	29,888,758
	30,090,650	30,736,399	24,502,555	41,410,154

Clean read (number)	28,243,807	23,557,745	22,858,723	22,623,421
	24,810,283	19,871,339	16,535,193	18,441,529
	27,669,070	29,809,596	23,400,051	16,787,161
Unique sRNA (number)	592,845	2,010,281	3,467,649	188,202
	314,918	1,567,677	1,976,367	154,649
	4,113,033	3,117,977	2,396,498	111,338
rRNA, tRNA, snRNA and snoRNA read (number)	1,740,899	307,766	233,107	28,462
	1,451,405	507,083	193,556	56,179
	324,321	313,366	609,494	14,949
known coral miRNA (number)	26	17	15	4
novel coral miRNA (number)	155	87	115	35
novel *Symbiodinium* miRNA (number)	7	9	4	6

436 coral miRNAs and 11 novel *Symbiodinium* miRNAs are identified, the detailed information
of which, such as loci, sequence and precursor structure, have been added to corresponding
miRNA datasets named CoralmiR (http://coral.bmeonline.cn/miR/) and CoralSym
(http://coral.bmeonline.cn/sym/). Comparing the two, CoralmiR is large yet flawed in that
there are inevitably several false positive results in it due to the de novo analysis process, while
CoralSym is small but precise because CoralSym miRNAs are supported by published
zooxanthellae genomes, enhancing their credibility.

153
Gene expression analysis

Illumina RNA-seqs were mapped to full-length transcripts using RSEM for gene expression
quantification and differential expression pattern identification. The results of gene
expression analysis are demonstrated in **Fig 2.** Gene numbers of differential expression
patterns are illustrated in **Fig 2.a,** where patterns representing differentially expressed genes
in single species occupy the top three spots, while no Pattern01 gene co-expressed among all
species is identified. Combined with the gene distribution revealed in **Fig 2.b,** the unique genes
account for the majority of total genes in quantity and expression for each species, suggesting
that gene structural differences among different corals may be significant. **Fig S2** displays the
distribution of gene expression and quantity in specific sub-sets, and these supplemental
results are consistent with the above findings for total gene expression.

Fig 2. Results of gene expression analysis. AM, MF, MC, PV are short for *A. muricata*, *M. foliosa*, *M. capricornis,* and *P. verrucosa*. In the enrichment plots, each species is indicated by a different colour and statistical significance is indicated by transparency. (a) Statistics of gene differential expression patterns among the four coral species. The vertical axis is the pattern name and the horizontal axis is the number of genes. Each pattern is illustrated with a contrasting color band listed beside the vertical axis. The top three patterns in quantity are Pattern08, Pattern02 and Pattern05, which represent genes expressed differentially in *A. muricata*, *P. verrucosa* and *M. foliosa*, while all Pattern01 genes identified as co-expressed among species fail to meet the statistical significance criteria (PPDE>0.95). (b) Venn graph of gene distribution among four species. Text is in the form of "*zooxanthellae gene number | holobiont gene number*." (c) KEGG enrichment of *Symbiodinium* genes. Pathways related to

photosynthesis are put on the plot. Nitrogen metabolism is outstanding in both quantity and statistical significance, while few other pathways exhibit enrichment. (d) GO enrichment of highly expressed coral-zooxanthella holobiont genes. Significantly enriched terms are "translation" in Biological Process, "ribosome" in Cellular Component and "structural constituent of ribosome" in Molecular Function, presenting the features of housekeeping genes.

However, **Fig 2.c** and **Fig 2.d** reveal gene functional similarities among the corals in spite of their structural differences. Among the zooxanthella genes, as **Fig 2.c** indicates, nitrogen metabolism is the common and dominant pathway. Considering the large number of pathways involving photosynthesis, zooxanthella gene functions seem fairly narrow, which is reminiscent of mitochondria, in that they obtain most of the proteins they need from their host cells rather than producing them on their own. This finding may be conducive to the hypothesis that zooxanthellae have effectively degenerated into coral organelles. Highly expressed coral genes are significantly enriched in the basic cellular functions, as shown in **Fig 2.d.** The top GO terms are "translation" in Biological Processes, "ribosome" in Cellular Components and "structural constituent of ribosome" in other words, the most highly expressed genes in the corals studied are housekeeping genes. Based on the enrichment results above, genes of coral-zooxanthella holobionts from the same habitat are believed to play analogous roles that are not significantly affected by species-specific structural differences, which may help to simplify the selection of a particular coral as an experimental model subject.

Table 4 lists the results of a hypothetical test of zooxanthella gene expression against total gene expression. Most zooxanthella genes appear to be limited in expression, hinting at suppression by host coral cells, while those from *M. foliosa* present abnormally high levels of expression in all three biological replicates. These differences provide a potential entry point for exploring cross-kingdom regulation mechanisms in coral-zooxanthella holobionts.

Table 4. Hypothetical test of zooxanthella gene expression against total gene expression. T-test method was utilized. $H_0: Exp_{zoox} \leq \overline{Exp}$; $H_1: Exp_{zoox} > \overline{Exp}$

Species	P_{rep1}	P_{rep2}	P_{rep3}
A. muricata	0.2066	0.0558	0.1367
M. foliosa	0.0007	0.0012	0.0014
M. capricornis	0.991	0.9934	0.988
P. verrucosa	0.3783	0.5881	0.4764

miRNA-target analysis

Eleven *Symbiodinium* miRNAs are identified from the total small RNA, as shown in **Fig 3**, numbered as smg-miR-1 to 11. Their expression profiles are supplemented in **Table S2**. Four miRNAs, smg-miR-1, 2, 4, and 7, exist in all investigated corals, and their expression levels are relatively high. It is as expected that, in intracellular environments, photobiont miRNA can maintain a certain concentration far beyond concentrations observed in traditional cross-kingdom research. This phenomenon may illustrate the advantage of coral as an observer-friendly model organism to some extent.

Fig 3. Results of miRNA-targets analysis. AM, MF, MC, PV are short for *A. muricata*, *M. foliosa*, *M. capricornis*, and *P. verrucosa*. GO.BP, GO.CC, GO.MF are short for biological process, cellular component and molecular function in gene ontology. KEGG.CP, KEGG.EIP, KEGG.GIP, KEGG.OS are short for cellular processes, environmental information processing, genetic Information processing and organismal systems in KEGG. Enrichment plots are generated with categories marked by color and statistical significance marked by transparency. Left: identified *Symbiodinium* miRNAs in investigated corals. The vertical axis indicates the miRNA name and the horizontal axis is the species name. Results of miRDP2 are marked with dark green while expanded results of mirdeep2 quantifier are marked with light green. Upper right: enrichment results of smg-miR-3,6,8,9,10 targets in *M. foliosa*. Notable terms include a series of GO terms related to cytoskeleton as well as cell migration, cell adhesion, and cell proliferation, including "regulation of Rho protein signal transduction," "DNA replication," and "microtubule-based movement" in biological process, "myosin complex" and "chromosome centromeric region" in cellular component and "Rho guanyl-nucleotide exchange factor activity," "Ran GTPase binding," "motor activity," and "chromatin binding" in molecular function. Lower right: enrichment results of smg-miR-11 targets in *A. muricata*. Enriched pathways are immune-related pathways including NOD-like receptor signaling pathway, IL-17 signaling pathway, RIG-I-like signaling pathway and Toll-like receptor signaling pathway.

As noted in **Fig 3**, *M. foliosa* has a unique miRNA group consisting of smg-miR-3, 6, 8, 9 and 10. In the results of enrichment analysis of their target genes, a series of GO terms related to the cytoskeleton as well as to cell migration, cell adhesion, and cell proliferation emerge, among which Rho-related terms are notable. Rho GTPases belonging to the Ras superfamily of small

GTP-binding protein serve as "molecular switches" in various signaling pathways, mainly by
exerting effects on the cytoskeleton (15). They are widely distributed in immune-related cells
and are frequently observed to participate in immune regulation. They usually cycle between
an active GTP-bound state and an inactive GDP-bound state, triggering immune responses
when bound to GTP until they are hydrolyzed into the GDP-bound state. Three proteins are
recognized to regulate this cycle, among which Rho guanyl-nucleotide exchange factor (Rho-
GEF) contributes to the activation of Rho GTPases, playing the role of positive regulator. From
this it is assumed that zooxanthellae secrete these miRNAs to downregulate the activity of
positive regulators in immune pathways. These miRNAs may also help remodel the
cytoskeleton of corals to allow for the entry of zooxanthellae, in a similar way to how some
parasites affect their hosts (16, 17).

Rho-related biological processes and molecular functions are also enriched in the target genes
of smg-miR-1 in *M. capricornis* and smg-miR-7 in *P. verrucoses* shown in **Table S1**. For *A.*
muricata, there is only one unique *Symbiodinium* miRNA, smg-miR-11. As illustrated in **Fig 3**,
several cytoskeleton-related terms appear in GO enrichment results, while KEGG enrichment
results directly demonstrate pathways in immune system. Though KEGG annotation may not
be exactly terminologically accurate for corals, in that most KEGG data sources are from
humans and other dominant model organisms, the results can be considered as further
evidence of immune-related cross-kingdom regulation in coral-zooxanthella holobionts. In
addition, autophagy-related genes are identified as these miRNAs' targets as well, which may
strengthen the credibility of the assumptions outlined above in view of many observations
that Rho GTPases upregulate autophagosome fusion and transport. **Fig 4** displays a predicted
pathway by which zooxanthellae block autophagosome production via miRNA-silencing the
upstream Rho "switch," nevertheless the possibility that zooxanthellae miRNAs directly target
autophagy-related genes cannot yet be ruled out.

259

Fig 4. Predicted downregulating rejection pathway in coral-zooxanthella holobionts. Zooxanthellae
secrete miRNAs targeting and silencing coral Rho-GEF genes to prevent the activation of Rho GTPases
to block threatening autophagosome production in the coral.

In contrast to the similarities among genes in different species of corals examined in this study,
zooxanthella miRNAs present functional differences among species. For instance, smg-miR-11
exists in *P. verrucosa*, while few significant functions are enriched in its target genes. smg-miR-
1, 2, 4 and 7 are observed to be co-expressed among all coral samples, while their target genes

are not functionally centralized. It seems that immune downregulation is the rare common
role they play in the investigated corals.

No coral miRNA targeting sites are identified in the genes of zooxanthellae. It is thus inferred
that miRNA silencing is not the main way corals regulate zooxanthellae, yet real matching
criterion of *Symbiodinium* targeting sites may not be as strict as that applied in this research.

Discussion

273
274

Role of miRNAs in coral-zooxanthella holobiont

Our work provides evidence for an early origin of cross-kingdom regulation as a self-defense
mechanism of autotrophs against heterotrophs. It is proposed that miRNAs target and silence
immune-related coral genes to reduce rejection, helping zooxanthellae to move beyond
provoking an antigenic response in potential coral hosts to becoming components of the
holobiont. Conversely, no coral miRNAs are found to target zooxanthella genes, indicating that
coral does not utilize miRNA as its main way to regulate zooxanthellae.

In view of the differences in gene function and expression levels revealed in **Fig 2.c** and **Table
4**, it could be further assumed that miRNA multiplicity may be conducive to function diversity
for zooxanthellae. For example, among the investigated corals, the unique miRNA group in *M.
foliosa*'s zooxanthellae is rather highly expressed and has diverse functions, while the miRNA
in *M. capricornis*'s zooxanthellae, none of which is unique, are expressed at the lowest level.
The group targeting effect of miRNA may reduce the inhibition of zooxanthellae by host coral
cells so as to permit greater flexibility in the physiological activity of zooxanthellae. Records in
Coral Trait (18) indicate that *M. foliosa* may be able to inhabit mesophotic coral reefs (19)
more than 30m deep (https://coraltraits.org/species/1014/traits/92), where few symbiont
zooxanthellae would be able to photosynthesize. The mutual dependence between coral and
zooxanthellae in this holobiont under these conditions would be much lower, not only
conforming to our finding but also supporting our hypothesis.

Coral model and "ecological niche effect"

From the experimental point of view, when conducting studies of miRNA interactions between
photobiont and animals, corals as model organisms have advantages in observer-friendliness
and simplicity of behavior. For one thing, the single-cell structure of zooxanthellae narrows
the observation range down to the intracellular environment and reduces miRNAs lost by
transport and degradation, helping to maintain a relatively higher photobiont miRNA
concentration for easier identification. For another, traits and behaviors of corals have been
well-researched, while the internal cross-kingdom regulation occurring inside corals appears
to be to be unidirectional from zooxanthellae to coral according to our research, thus it will
not be too difficult to record the impact of zooxanthellae on corals under appropriate
experimental conditions.

The coral-zooxanthella holobionts investigated in this study appear to have simple behavior.
The majority of highly expressed coral genes are devoted to basic biological processes, such
as the construction of ribosomes, while most exogenous zooxanthellae work in specific
pathways under various limitations. Though our work has not identified any miRNA-based
coral-to-zooxanthellae regulations, corals' inhibitory effects on zooxanthellae are certainly
observed. It is not unreasonable to predict that similar intracellular "dystopias" exist in a
variety of ancient autotroph-heterotroph holobionts in which mutual regulations between
symbiotic organisms vary in mechanism and are imbalanced in intensity.

It is further suggested that the closer the organism is to the bottom ecological niche, the
greater the non-gene impact on it. Though coral reefs are often made up of a community
exhibiting significant species diversity, our research shows that corals from the same habitat
share many physiological commonalities. The "ecological niche effect" hypothesis is expected
to support the feasibility of this coral model and provide guidelines for specimen collection for
further research.

320
Limitation and prospect

This work is typical omics-based research, mining information from large-scale bio-data, thus
it is impossible to avoid systematic errors coming from bioinformatics tools and databases,
especially when taking the "coral data gap" into account. Insufficient or incorrect data about
coral in bio-databases affecting annotations, tags and reference sequences have emerged
more than once during this research. However, the largest limitation to this research project
actually comes from the lack of data about zooxanthellae. There exist various sub-clades of
Symbiodinium in one coral, the genome differences among which may be as huge as that
between humans and *Pan troglodytes*. Although the negative impact might be sharply reduced
by the "ecological niche effect" mentioned above, using one clade-C1 genome to represent all
clade-C genomes is destined to cause omissions. It has been observed that Clade D
zooxanthellae also occupy a certain percentage of the *P. verrucosa* holobiont (**Video S1**),
implying that a great deal of information is yet to be mined. Besides, the mixed *Symbiodinium*
samples extracted from coral are still hard to isolate and culture in vitro currently. Acquiring
genomes of specific *Symbiodinium* species precisely from coral-zooxanthellae holobionts can
be described as the most essential challenge related to cross-kingdom regulation research.

In summary, our work integrates new full-length transcriptomes and small RNAs of four
common and frequently dominant reef-building corals with a public *Symbiodinium C1* genome
to explore coral-zooxanthella holobionts, shedding light on intracellular mutual regulation
between these ancient heterotrophs and autotrophs. It reveals the miRNA-based anti-
rejection mechanism used by zooxanthellae, proposes the concept of coral as a model
organism for cross-kingdom regulation along with ecological niche effect, and offers a
foundation for further cross-kingdom research.

We have added all our data, including gene annotation and miRNA information, to the
CoralBioinfo database (http://coral.bmeonline.cn), in a gesture to build a high-quality multi-
omics platform for further coral research. Multiple web services have been developed for it,
most of which turn out to be more user-friendly to bioinformaticians familiar with "dry lab"
skills. Data contribution requests for joint efforts are always welcome.

Materials and Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Chunpeng He (cphe@seu.edu.cn), or by Zuhong Lu (zhlu@seu.edu.cn).

Materials availability

This study did not generate any new reagents.

Data availability

Raw data from our own full-length transcriptome sequencing and small RNA sequencing are available from NCBI (PRJNA544778).

Though we sincerely recommend acquiring data via our CoralBioinfo database (http://coral.bmeonline.cn) introduced in this paper, we have already uploaded necessary supplementary data to better serve researchers.

(https://seunic-my.sharepoint.cn/:f:/g/personal/230218818_seu_edu_cn/Eo4QciTXwQRBkYqcKEacEN8BRCKffoV-299v2HFILGF6lQ?e=dlQJx9)

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All coral samples were collected and processed in accordance with local laws for invertebrate protection and approved by the Ethics Committee of Institutional Animal Care and Use Committee of Nanjing Medical University (protocol code IACUC-1910003 and date of approval is 10 October 2019).

The species in the study were collected from the Xisha Islands in the South China Sea (latitude 15°40'–17°10' north, longitude 111°–113° east).

The coral samples were cultured in our laboratory coral tank with conditions conforming to their habitat environment. All the species were raised in a RedSea® tank (redsea575, Red Sea Aquatics Ltd) at 26°C and 1.025 salinity (Red Sea Aquatics Ltd). The physical conditions of the coral culture system are as follows: three coral lamps (AI®, Red Sea Aquatics Ltd), a protein skimmer (regal250s, Reef Octopus), a water chiller (tk1000, TECO Ltd), two wave devices (VorTechTM MP40, EcoTech Marine Ltd), and a calcium reactor (Calreact 200, Reef Octopus), etc.

METHOD DETAILS

RNA extraction

All RNA extraction procedures follow the manufacturers' instructions. The total RNA was isolated with TRIzol LS Reagent (Thermo Fisher Scientific, 10296028) and treated with DNase I (Thermo Fisher Scientific, 18068015). The high quality mRNA was isolated with a FastTrack MAG Maxi mRNA Isolation Kit (Thermo Fisher Scientific, K1580-02). Samples were separated from healthy *Acropora muricata*, *Montipora foliosa*, *Montipora capricornis* and *Pocillopora*

verrucosa to ensure that enough high quality RNA—more than 10 µg—could be obtained for
a cDNA transcriptome library.

Full-length transcriptome sequencing

Before establishing a cDNA transcriptome library, the quality of total RNA had to be tested.
Agarose gel electrophoresis was used to analyze the degree of degradation of RNA and
whether it was contaminated. A Nanodrop nucleic acid quantifier was used to detect the purity
of RNA (OD260/280 ratio), a Qubit RNA assay was used to quantify the RNA concentration
accurately, and an Agilent 2200 TapeStation was used to accurately detect the integrity of the
RNA.

The Clontech SMARTer® PCR cDNA Synthesis Kit (Clontech Laboratories, 634926) and the
BluePippin Size Selection System protocol, as described by Pacific Biosciences (PN 100-092-
800-03) were using to prepare the Iso-Seq library according to the Isoform Sequencing
protocol (Iso-Seq).

The RNA-seq library was prepared according to the standard library construction for the
Illumina HiSeq X Ten, using three biological replicates for each species of coral.

We used the PacBio Sequel II platform with SMRT (single molecular real time) sequencing
technology and SMRTlink v7.0 software (minLength 50; maxLength 15,000; minPasses 1) to
process sequencing samples (20). After polymer read bases were performed, the
subreads.bam files were obtained by removing the joint and the original offline data with
lengths of under 50 bp. The CCSs (circular consensus sequences) were obtained by putting the
subreads.bam file through the CCS algorithm, which is self-correcting for the single molecule
multiple sequencing sequence. Consequently, the full-length-non-chimera (FLNC) and nFL
(non-full-length, non-chimera) sequences were identified by detecting whether CCSs
contained 5'-primer, 3'-primer and poly-A. FLNC sequences of the same transcript were
clustered with a hierarchical n * log (n) algorithm to obtain consensus sequences. The
corrected consensus reads were polished from consensus sequences (Arrow polishing) using
LoRDEC (21) v0.7 software using the RNA-seq data sequenced with the Illumina HiSeq X Ten
platform. Using CD-HIT (22) software (-c 0.95 -T 6 -G 0 -aL 0.00 -aS 0.99), all redundancies were
removed in corrected consensus reads to acquire final full-length transcripts and unigenes for
subsequent bioinformatics analyses.

Small RNA sequencing

A total amount of 3 µg of RNA per sample was used as input material for the small RNA library.
Sequencing libraries were generated using the NEBNext® Multiplex Small RNA Library Prep Set
for Illumina® (NEB, USA) following the manufacturer's recommendations, and index codes
were added to attribute sequences to each sample. Briefly, the NEB 3' SR Adapter was directly
and specifically ligated to the 3' end of miRNA, siRNA and piRNA. After the 3' ligation reaction,
the SR RT Primer was hybridized to the excess of 3' SR Adapter that remained free after the 3'
ligation reaction, transforming the single-stranded DNA adapter into a double-stranded DNA
molecule. This step is important to prevent adapter-dimer formation, and dsDNAs are also not
substrates for ligation mediated by T4 RNA Ligase 1, so they do not ligate to the 5' SR Adapter
in the subsequent ligation step. Next, the 5' ends adapter was ligated to the 5' ends of miRNAs,
siRNA and piRNA. Then the first strand of cDNA was synthesized using M-MuLV Reverse
Transcriptase (RNase H–). PCR amplification was performed using LongAmp Taq 2X Master
Mix, SR Primer for Illumina and index (X) primer. PCR products were purified on an 8%
polyacrylamide gel (100V, 80 min). DNA fragments corresponding to 140~160 bp (the length
of small noncoding RNA plus the 3' and 5' adapters) were recovered and dissolved in 8 µL

elution buffer. Finally, library quality was assessed with the Agilent Bioanalyzer 2100 system
using DNA High Sensitivity Chips.

The clustering of the index-coded samples was performed on a cBot Cluster Generation
System using a TruSeq SR Cluster Kit v3-cBot-HS (Illumia) according to the manufacturer's
instructions. After cluster generation, the library preparations were sequenced on an Illumina
HiSeq X Ten platform and 50bp single-end reads were generated.

Clean data for downstream analysis were obtained by removing reads with low quality, poly-
N|A|T|G|C, and 5' adapter contaminants. Reads without the 3' adapter or the insert tag were
cleaned as well.

Gene function annotation

Gene function was annotated using the following databases: Nr (NCBI non-redundant protein
sequences) (23), Nt (NCBI non-redundant nucleotide sequences), Pfam (Protein family) (24),
KOG (Clusters of Orthologous Groups of proteins) (25), Swiss-Prot (A manually annotated and
reviewed protein sequence database) (26), GO (Gene Ontology) (27) and KEGG (Kyoto
Encyclopedia of Genes and Genomes) (14). We use BLAST 2.7.1+ in NCBI setting the e-value
'1e-5' in Nt database analysis, Diamond v0.8.36 BLASTX software, setting the e-value to '1e-5'
in Nr, KOG, Swiss-Prot and KEGG databases analysis, and HMMER 3.1 package for Pfam
database analysis.

Gene expression analysis

455 We used RSEM (28) to align Illumina reads to full-length transcripts and to quantify gene
456 expression. Transcript names were converted to Nr IDs to build the multi-coral expression
457 profile. EBSeq (29) was finally called by RSEM to identify differential expression patterns.

miRNA identification

After quality control and length filtering (18–35 nt) of raw small RNA sequences, known miRNA
alignment and novel miRNA prediction were conducted. For known miRNA alignment,
miRBase was used as the reference, while mirdeep2 (30) and srna-tools-cli (http://srna-
workbench.cmp.uea.ac.uk) were employed to identify miRNAs and draw their secondary
structures. For novel miRNA prediction, small RNA tags were mapped to RepeatMasker (31)
and Rfam (32) to remove tags originating from protein-coding genes, repeat sequences, rRNA,
tRNA, snRNA, and snoRNA. Next, miREvo (33) and mirdeep2 were integrated to predict novel
miRNA among the remaining unannotated tags. All miRNA targeting sites on zooxanthella
genes were predicted with psRobot (34) (no mismatches allowed).

miRDP2 (35) was employed to identify *Symbiodinium* miRNAs in each coral separately after
building the bowtie index of the *Symbiodinium C1* genome (36) downloaded from
Reefgenomics (37). Initial outputs were corrected by genome annotation in which sequences
with loci hitting CDS were removed. Then mirdeep2 using filtered sequences as inputs was run
on all raw data to quantify miRNA expression as well to eliminate false negatives. All miRNA
targeting sites were finally predicted with miRanda (38) (-sc 145 -en -10 -scale 4 -strict).

Enrichment analysis

475 For each coral, the top 500 most highly expressed genes (that each entered the top 500 among
476 three biological replicates more than once), and the zooxanthellae genes annotated by Nr and
477 predicted target genes were all selected for GO/KEGG enrichment using clusterProfiler (39)
478 (pvalueCutoff = 0.05, pAdjustMethod = 'BH', qvalueCutoff = 0.2).

Website development

Website development

CoralBioinfo (http://coral.bmeonline.cn) was developed under the LEMP (Linux, Nginx, MySQL
and PHP) stack. Web functions were implemented with PHP 7.2, while Nginx works as the web
server. Most services have been built into Docker images, which are beneficial for cross-
platform deployment and extended development.

All data are stored in MySQL tables with indexes to improve query efficiency. In addition,
sequence data have been formatted into NCBI-BLAST local databases for alignment and
download services.

ADDITIONAL RESOURCES

Detailed database structure manuals along with user tutorials are all available at
http://coral.bmeonline.cn/help.php.

API documents are available at http://coral.bmeonline.cn/api/.

Supplemental information

Fig S1. Annotation of four full-length coral transcriptomes with Nr database. Horizontal axis is species (txid) and vertical axis is unigene number. The top three species with the highest number of annotated unigenes are *A. digitifera*, *E. pallida* and *N. vectensis*, which account for 92.8% in *A. muricata*, 89.8% in *M. foliosa*, 90.5% in *M. capricornis* and 85.9% in *P. verrucosa*.

Fig S2. Supplemental results of gene expression analysis. (a) Differential zooxanthella gene expression patterns among the four coral species studied. The horizontal axis is the pattern name and the vertical axis is the gene number. The top three patterns in quantity are Pattern05, Pattern02 and Pattern08, which represent differentially expressed genes in *M. foliosa*, *P. verrucosa* and *A. muricata*, while all Pattern01 genes identified as being co-expressed among species fail to meet the statistical significance criteria (PPDE>0.95). (b) Venn graph of top 500 most highly expressed genes among four species.

File S1. Supplementary processing details of full-length transcriptomes.

File S2. Supplementary processing details of small RNAs.

Table S1. Complete results of miRNA-target gene enrichment analysis. GeneIDs are CSG/T IDs in CoralSym (http://coral.bmeonline.cn/sym/).

Table S2. Expression profiles of *Symbiodinium* miRNAs. AM_x, MF_x, MC_x, PV_x are short for the biological replicates of *A. muricata*, *M. foliosa*, *M. capricornis,* and *P. verrucosa*. The expression level is quantified in RPM.

Video S1. Demo of searching CoralTrait++ via CoralAPI. CoralTrait++ is a manually-built simplified version of the Coral Trait database. It consists of 12 quantitative traits and six descriptive traits with all units standardized, and more than 45,000 records. It was revealed in this demo that clade D zooxanthellae also hold a certain percentage of the traits seen in the *Pocillopora verrucosa* holobiont.

Data S1. Unigene sequences of *A. muricata*, *M. foliosa*, *M. capricornis,* and *P. verrucosa*. BLAST service is available at http://coral.bmeonline.cn/blast_unigene/.

Data S2. Gene annotation of four corals from Nr, Nt, Pfam, KOG, Swiss-Prot, GO and KEGG databases. All gene information is searchable at http://coral.bmeonline.cn/gene/.

Data S3. CoralSym dataset including information of 11 novel *Symbiodinium* miRNAs and their targeting sites. The format manual is available at http://coral.bmeonline.cn/sym/help.php #database-content. Exact searching of targeting information is accessible at /sym/ search.php.

References

1. Bartel DP. Metazoan MicroRNAs. Cell. 2018 Mar 22;173(1):20-51. doi: 10.1016/j.cell.2018.03.006.

2. Zhang L, Hou D, Chen X, Li D, Zhu L, Zhang Y, Li J, Bian Z, Liang X, Cai X, Yin Y, Wang C, Zhang T, Zhu D, Zhang D, Xu J, Chen Q, Ba Y, Liu J, Wang Q, Chen J, Wang J, Wang M, Zhang Q, Zhang J, Zen K, Zhang CY. Exogenous plant MIR168a specifically targets mammalian LDLRAP1: evidence of cross-kingdom regulation by microRNA. Cell Res. 2012 Jan;22(1):107-26. doi: 10.1038/cr.2011.158.

3. Liang H, Zhang S, Fu Z, Wang Y, Wang N, Liu Y, Zhao C, Wu J, Hu Y, Zhang J, Chen X, Zen K, Zhang CY. Effective detection and quantification of dietetically absorbed plant microRNAs in human plasma. J Nutr Biochem. 2015 May;26(5):505-12. doi: 10.1016/j.jnutbio.2014.12.002.

4. Zhou Z, Li X, Liu J, Dong L, Chen Q, Liu J, Kong H, Zhang Q, Qi X, Hou D, Zhang L, Zhang G, Liu Y, Zhang Y, Li J, Wang J, Chen X, Wang H, Zhang J, Chen H, Zen K, Zhang CY. Honeysuckle-encoded atypical microRNA2911 directly targets influenza A viruses. Cell Res. 2015 Jan;25(1):39-49. doi: 10.1038/cr.2014.130.

5. Chen X, Dai GH, Ren ZM, Tong YL, Yang F, Zhu YQ. Identification of Dietetically Absorbed Rapeseed (Brassica campestris L.) Bee Pollen MicroRNAs in Serum of Mice. Biomed Res Int. 2016;2016:5413849. doi: 10.1155/2016/5413849.

6. Jia L, Zhang D, Xiang Z, He N. Nonfunctional ingestion of plant miRNAs in silkworm revealed by digital droplet PCR and transcriptome analysis. Sci Rep. 2015 Jul 21;5:12290. doi: 10.1038/srep12290.

7. Chin AR, Fong MY, Somlo G, Wu J, Swiderski P, Wu X, Wang SE. Cross-kingdom inhibition of breast cancer growth by plant miR159. Cell Res. 2016 Feb;26(2):217-28. doi: 10.1038/cr.2016.13.

8. Zhou LK, Zhou Z, Jiang XM, Zheng Y, Chen X, Fu Z, Xiao G, Zhang CY, Zhang LK, Yi Y. Absorbed plant MIR2911 in honeysuckle decoction inhibits SARS-CoV-2 replication and accelerates the negative conversion of infected patients. Cell Discov. 2020 Aug 5;6:54. doi: 10.1038/s41421-020-00197-3.

9. Dickinson B, Zhang Y, Petrick JS, Heck G, Ivashuta S, Marshall WS. Lack of detectable oral bioavailability of plant microRNAs after feeding in mice. Nat Biotechnol. 2013 Nov;31(11):965-7. doi: 10.1038/nbt.2737.

10. Campbell K. Do the microRNAs we eat affect gene expression? Nature. 2020;582(7812):S10. doi:10.1038/d41586-020-01767-x

11. Chen Q, Zhang F, Dong L, Wu H, Xu J, Li H, Wang J, Zhou Z, Liu C, Wang Y, Liu Y, Lu L, Wang C, Liu M, Chen X, Wang C, Zhang C, Li D, Zen K, Wang F, Zhang Q, Zhang CY. SIDT1-dependent absorption in the stomach mediates host uptake of dietary and orally administered microRNAs. Cell Res. 2021 Mar;31(3):247-258. doi: 10.1038/s41422-020-0389-3.

12. Shahid S, Kim G, Johnson NR, Wafula E, Wang F, Coruh C, Bernal-Galeano V, Phifer T, dePamphilis CW, Westwood JH, Axtell MJ. MicroRNAs from the parasitic plant Cuscuta campestris target host messenger RNAs. Nature. 2018 Jan 3;553(7686):82-85. doi: 10.1038/nature25027.

13. Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. miRBase: tools for microRNA genomics. Nucleic Acids Res. 2008 Jan;36(Database issue):D154-8. doi: 10.1093/nar/gkm952.

14. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res. 2017 Jan 4;45(D1):D353-D361. doi: 10.1093/nar/gkw1092.

15. Etienne-Manneville S, Hall A. Rho GTPases in cell biology. Nature. 2002 Dec 12;420(6916):629-35. doi: 10.1038/nature01148.

16. Gonzalez V, Combe A, David V, Malmquist NA, Delorme V, Leroy C, Blazquez S, Ménard
R, Tardieux I. Host cell entry by apicomplexa parasites requires actin polymerization in
the host cell. Cell Host Microbe. 2009 Mar 19;5(3):259-72. doi:
10.1016/j.chom.2009.01.011.
17. Walker ME, Hjort EE, Smith SS, Tripathi A, Hornick JE, Hinchcliffe EH, Archer W, Hager
KM. Toxoplasma gondii actively remodels the microtubule network in host cells.
Microbes Infect. 2008 Nov-Dec;10(14-15):1440-9. doi: 10.1016/j.micinf.2008.08.014.
18. Madin JS, Anderson KD, Andreasen MH, Bridge TC, Cairns SD, Connolly SR, Darling ES,
Diaz M, Falster DS, Franklin EC, Gates RD, Harmer A, Hoogenboom MO, Huang D, Keith
SA, Kosnik MA, Kuo CY, Lough JM, Lovelock CE, Luiz O, Martinelli J, Mizerek T, Pandolfi
JM, Pochon X, Pratchett MS, Putnam HM, Roberts TE, Stat M, Wallace CC, Widman E,
Baird AH. The Coral Trait Database, a curated database of trait information for coral
species from the global oceans. Sci Data. 2016 Mar 29;3:160017. doi:
10.1038/sdata.2016.17.
19. Loya, Yossi, Kimberly A. Puglise, and Tom CL Bridge, eds. Mesophotic coral ecosystems.
Vol. 12. Springer, 2019. doi:10.1007/978-3-319-92735-0.
20. Chin CS, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, Dunn C, O'Malley
R, Figueroa-Balderas R, Morales-Cruz A, Cramer GR, Delledonne M, Luo C, Ecker JR,
Cantu D, Rank DR, Schatz MC. Phased diploid genome assembly with single-molecule
real-time sequencing. Nat Methods. 2016 Dec;13(12):1050-1054. doi:
10.1038/nmeth.4035.
21. Salmela L, Rivals E. LoRDEC: accurate and efficient long read error correction.
Bioinformatics. 2014 Dec 15;30(24):3506-14. doi: 10.1093/bioinformatics/btu538.
22. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics. 2012 Dec 1;28(23):3150-2. doi:
10.1093/bioinformatics/bts565.
23. Pruitt KD, Tatusova T, Klimke W, Maglott DR. NCBI Reference Sequences: current status,
policy and new initiatives. Nucleic Acids Res. 2009 Jan;37(Database issue):D32-6. doi:
10.1093/nar/gkn721.
24. El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M, Richardson
LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan D, Tosatto SCE,
Finn RD. The Pfam protein families database in 2019. Nucleic Acids Res. 2019 Jan
8;47(D1):D427-D432. doi: 10.1093/nar/gky995.
25. Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM,
Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan
S, Wolf YI, Yin JJ, Natale DA. The COG database: an updated version includes eukaryotes.
BMC Bioinformatics. 2003 Sep 11;4:41. doi: 10.1186/1471-2105-4-41.
26. Bairoch A, Apweiler R. The SWISS-PROT protein sequence database and its supplement
TrEMBL in 2000. Nucleic Acids Res. 2000 Jan 1;28(1):45-8. doi: 10.1093/nar/28.1.45.
27. The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing
strong. Nucleic Acids Res. 2019 Jan 8;47(D1):D330-D338. doi: 10.1093/nar/gky1055.
28. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 2011 Aug 4;12:323. doi:
10.1186/1471-2105-12-323.
29. Leng N, Dawson JA, Thomson JA, Ruotti V, Rissman AI, Smits BM, Haag JD, Gould MN,
Stewart RM, Kendziorski C. EBSeq: an empirical Bayes hierarchical model for inference
in RNA-seq experiments. Bioinformatics. 2013 Apr 15;29(8):1035-43. doi:
10.1093/bioinformatics/btt087.
30. Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. miRDeep2 accurately
identifies known and hundreds of novel microRNA genes in seven animal clades.
Nucleic Acids Res. 2012 Jan;40(1):37-52. doi: 10.1093/nar/gkr688.

31. Chen N. Using Repeat Masker to identify repetitive elements in genomic sequences.
Current protocols in bioinformatics. 2004;5(1):4-10.
32. Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, Bateman A,
Finn RD, Petrov AI. Rfam 13.0: shifting to a genome-centric resource for non-coding
RNA families. Nucleic Acids Res. 2018 Jan 4;46(D1):D335-D342. doi:
10.1093/nar/gkx1038.
33. Wen M, Shen Y, Shi S, Tang T. miREvo: an integrative microRNA evolutionary analysis
platform for next-generation sequencing experiments. BMC Bioinformatics. 2012 Jun
21;13:140. doi: 10.1186/1471-2105-13-140.
34. Wu HJ, Ma YK, Chen T, Wang M, Wang XJ. PsRobot: a web-based plant small RNA meta-
analysis toolbox. Nucleic Acids Res. 2012 Jul;40(Web Server issue):W22-8. doi:
10.1093/nar/gks554.
35. Kuang Z, Wang Y, Li L, Yang X. miRDeep-P2: accurate and fast analysis of the microRNA
transcriptome in plants. Bioinformatics. 2019 Jul 15;35(14):2521-2522. doi:
10.1093/bioinformatics/bty972.
36. Liu H, Stephens TG, González-Pech RA, Beltran VH, Lapeyre B, Bongaerts P, Cooke I,
Aranda M, Bourne DG, Forêt S, Miller DJ, van Oppen MJH, Voolstra CR, Ragan MA, Chan
CX. Symbiodinium genomes reveal adaptive evolution of functions related to coral-
dinoflagellate symbiosis. Commun Biol. 2018 Jul 17;1:95. doi: 10.1038/s42003-018-
0098-3.
37. Liew YJ, Aranda M, Voolstra CR. Reefgenomics.Org - a repository for marine genomics
data. Database (Oxford). 2016 Dec 26;2016:baw152. doi: 10.1093/database/baw152.
38. Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. MicroRNA targets in
Drosophila. Genome Biol. 2003;5(1):R1. doi: 10.1186/gb-2003-5-1-r1.
39. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological
themes among gene clusters. OMICS. 2012 May;16(5):284-7. doi:
10.1089/omi.2011.0118.

Acknowledgments

We are grateful for Dr. Xiaojun Xia's technical support in our production environment.

Author Contributions

YZ: experiment, database, writing and editing. TH: data uploading. ZL & JC: reviewing. CH: supervision. XL: project approval. All authors contributed to the article and approved the submitted version.

Declaration of Interests

All authors declare that no conflict of interest exists.