1

Banp regulates DNA damage response and chromosome segregation during the cell cycle in zebrafish retina

Swathy Babu, Yuki Takeuchi, and Ichiro Masai[#]

Developmental Neurobiology Unit, Okinawa Institute of Science and Technology Graduate University (OIST)

1919-1 Tancha, Onna, Okinawa, Japan

Correspondence should be addressed: masai@oist.jp

Keywords: Banp, DNA damage, chromosome segregation, cell cycle, cenpt, ncapg, tp53

Word Count for Introduction/Results/Discussion: 6068

Abstract

Btg3-associated nuclear protein (Banp) was originally identified as a nuclear matrix-associated protein and it functions as a tumor suppressor. At molecular level, Banp regulates transcription of metabolic genes via a CGCG-containing motif called the Banp motif. However, its physiological roles in embryonic development are unknown. Here we report that Banp is indispensable for DNA damage response and chromosome segregation during mitosis. Zebrafish *banp* mutants show mitotic arrest and apoptosis in developing retina. We found that DNA replication stress and tp53-dependent DNA damage responses were activated to induce apoptosis in *banp* mutants, suggesting that Banp is required for integrity of DNA replication and DNA damage repair. Furthermore, in *banp* mutants, chromosome segregation was not smoothly processed from prometaphase to anaphase, leading to a prolonged M-phase. Our RNA- and ATAC-sequencing identified 31 candidates for direct Banp target genes that carry the Banp motif. Interestingly, two chromosome segregation regulators, *cenpt* and *ncapg*, are included in this list. Thus, Banp directly regulates transcription of *cenpt* and *ncapg* to promote chromosome segregation during mitosis. Our findings provide the first *in vivo* evidence that Banp is required for cell-cycle progression and cell survival by regulating DNA damage responses and chromosome segregation during mitosis.

(197 words)

Introduction

The vertebrate retina is composed of six major classes of neurons and one type of glia, namely retinal ganglion cells (RGCs), three types of interneurons (amacrine cells, bipolar cells, and horizontal cells), two types of photoreceptors (rods and cones), and Müller glia These retinal cells are assembled to form the neural circuit responsible for phototransduction and visual processing (Dowling, 2012). During development, retinal progenitor cells are multipotent and give rise to all seven retinal cell types in response to extrinsic and intrinsic cues (Boije et al., 2014). In addition, retinal neurogenesis is spatially and temporally regulated among vertebrates. In zebrafish, neurogenesis starts in the ventro-nasal retina at 25 h post-fertilization (hpf) and then spreads into the entire neural retina by 72 hpf (Hu and Easter, 1999; Masai et al., 2000). In general, early-born post-mitotic cells differentiate into RGCs and cone photoreceptors, followed by amacrine cells, bipolar cells, and horizontal cells, and late-born cells differentiate into rod photoreceptors and Müller glia, indicating stage-dependent retinal histogenesis.

Another important characteristic of retinal development is that retinal neurogenesis is linked to cell-cycle regulation. In zebrafish, cell-cycle length of retinal progenitor cells is around 32–49 h during 6 to 25 hpf, but shortened to 8–10 h once retinal neurogenesis starts at 25 hpf (Li et al., 2000), suggesting that activation of cell-cycle progression is associated with neurogenesis. It is generally accepted that active cell proliferation may increase DNA replication errors, which induces DNA replication stress and subsequent DNA damage (Berti et al., 2020; Blackford and Jackson, 2017; Waterman et al., 2020). Thus, DNA damage response is essential to ensure genomic integrity (Murga et al., 2009). Otherwise, retinal progenitor cells undergo cell-cycle arrest or apoptosis. Consistently, in zebrafish, retinal apoptosis occurs in the absence of DNA damage response regulators such as DNA damage-binding protein 1 (DDB1) (Hu et al., 2015), Fanconi anemia group D2 (Fancd2) (Liu et al., 2003), and Ku80 (Xrcc80) (Bladen et al., 2005). We also showed that retinal apoptosis occurs in zebrafish *prim1* mutants, in which DNA replication stress is chronically elevated. In this mutant, a central regulator of the DNA replication checkpoint, Ataxia telangiectasia-mutated (ATM) and Rad3-related (ATR), is activated, which subsequently activates DNA damage response regulators, ATM and Chk2, leading to tp53-mediated apoptosis. Thus, DNA replication checkpoint and DNA damage response function as safeguards to

ensure integrity of retinal cell proliferation and neurogenesis (Yamaguchi et al., 2008).

However, it remains to be seen how the DNA damage response pathway is integrated

into retinal cell proliferation and neurogenesis.

BTG3-associated nuclear protein (Banp) (also known as Scaffold/matrix

attachment region binding protein 1 (SMAR1)) was identified as a nuclear protein that

binds to the matrix-associated region (MAR) of the β-locus of murine T cell receptors

(Birot et al., 2000; Chattopadhyay et al., 2000). MARs are often found near gene

promoters (Scheuermann and Garrard, 1999) and constitute a periodic loop domain,

through which chromosomal DNA is topologically organized inside the nucleus

(Blasquez et al., 1989; Hart and Laemmli, 1998). Very recently, it was reported that

Banp binds to the nucleotide sequence TCTCGCGAGA, called the "Banp motif", which

is enriched near the transcription initiation site of CpG island promoters, and regulates

gene transcription in a DNA methylation-dependent manner. Banp promotes

transcription of essential metabolic genes in pluripotent stem and terminal differentiated

neuronal cells in humans and mice (Grand et al., 2021). Furthermore, Banp functions as

a tumor suppressor (Kaul et al., 2003; Kaul-Ghanekar et al., 2005). In humans, the *banp*

gene is located on chromosome 16q24, the genomic region of which often suffers from

loss-of-heterozygosity in various tumors (Tsuda et al., 1994; Wang et al., 1999). Banp

interacts with tumor suppressor tp53 to modulate its functions (Kaul et al., 2003;

Pavithra et al., 2009; Sinha et al., 2012; Sinha et al., 2010). Overexpression of Banp

suppresses cancer growth (Bhagat et al., 2018; Liu et al., 2014; Taye et al., 2018).

However, these cellular functions of Banp were proposed mostly from *in vitro* studies,

and genetic deletion of *banp* was reportedly unsuccessful (Grand et al., 2021), whereas

banp knockout mice show embryonic lethality (Chemmannur et al., 2015), which makes

in vivo analysis difficult. Thus, physiological functions of Banp are still unknown.

In this study, we identified zebrafish *banp* mutants, which show mitotic arrest

and apoptosis in the developing retina. DNA replication checkpoint and tp53-dependent

DNA damage response are activated to induce apoptosis in *banp* mutant retinas. Thus,

Banp is required for integrity of DNA replication and DNA damage repair. Furthermore,

live imaging revealed that chromosome segregation fails to proceed smoothly during

mitosis in *banp* mutants, resulting in a prolonged M-phase. Our RNA-sequencing and

ATAC-sequencing enabled us to identify 31 candidates for direct Banp target genes that

carry the Banp motif near their promoters. Interestingly, two chromosome segregation
regulators, *cenpt* and *ncapg*, are included in this list. Thus, Banp directly regulates the
transcription of *cenpt* and *ncapg* to promote chromosome segregation during mitosis.
Taken together, these data suggest that Banp is required for cell-cycle progression and
cell survival in zebrafish retinas by regulating DNA damage response and chromosome
segregation during mitosis.
109

Results

Zebrafish *rw337* mutants show mitotic cell accumulation and cell death in the retina after 2.5 dpf

We performed large-scale mutagenesis using zebrafish and identified a zebrafish mutant, *rw337*. *rw337* mutants have smaller eyes and a smaller optic tectum at 4 days post-fertilization (dpf) (Fig. 1A). Around 6–7 dpf, *rw337* mutants die, indicating a lethal mutation. Next, we examined retinal phenotypes (Fig. 1B). At 1.5 dpf, the retina consists of retinal progenitor cells and early differentiating neurons. There was no difference in retinal morphology between *rw337* mutants and wild-type siblings. However, at 2 dpf, mitotic-like round cells had accumulated in the apical region of the retina in *rw337* mutants, indicating mitotic arrest. At 2.5 dpf, in wild-type retinas, three nuclear layers, the RGC layer, the inner nuclear layer (INL) and the outer nuclear layer (ONL), and two plexiform layers, the inner plexiform layer (IPL) and the outer plexiform layer (OPL), had formed. In *rw337* mutants, only the RGC layer was observed; however, many pyknotic nuclei were observed in the outer retinal region. At 3 dpf, the RGC layer, IPL and INL appeared in *rw337* mutants. However, pyknotic nuclei were densely packed in the outer region of *rw337* mutant retinas. At 4 dpf, most dead cell debris had decreased in *rw337* mutants. The RGC layer, IPL, and INL were formed; however, no ONL was observed, despite very small, patchy photoreceptor-like cells. At 3 and 4 dpf, the ciliary marginal zone (CMZ), where retinal stem and progenitor cells are located, was observed in *rw337* mutants. Thus, RGCs differentiate, but mitotic arrest and cell death occur after 2.5 dpf in *rw337* mutants.

Next, we examined neuronal differentiation in *rw337* mutants at 4 dpf. The transgene *Tg[ath5:GFP]* drives GFP expression in RGCs (Fig. 1-figure supplement 1A) (Masai et al., 2000). Pax6 is strongly expressed in amacrine cells and weakly in RGCs (Fig. 1-figure supplement 1C) (Macdonald and Wilson, 1997). Prox1 is expressed in bipolar cells and horizontal cells (Fig. 1-figure supplement 1E) (Jusuf and Harris, 2009). zpr1 antibody labels double cone photoreceptors (Fig. 1-figure supplement 1G) (Larison and Bremiller, 1990). Anti-Gluta

mine synthetase (GS) antibody labels Müller cells (Fig. 1-figure supplement 1I) (Peterson et al., 2001). We observed that all these markers are expressed in *rw337*

mutants (Fig. 1-figure supplement 1B, 1D, 1F, 1H, 1J). However, the numbers of strong

Pax6+ cells and GS+ cells were decreased in *rw337* mutants (Fig. 1-figure supplement

1D, 1J), suggesting that amacrine cells and Müller cells were reduced. Prox1+ cells

were misplaced and less numerous in *rw337* mutants, suggesting that horizontal and

bipolar cells are reduced (Fig. 1-figure supplement 1F). zpr1+ cells were not detected,

except for a small patch of zpr1+ cells in the ventral retina, where photoreceptor

differentiation initially occurs, in *rw337* mutants (Fig. 1-figure supplement 1H),

suggesting that most cone photoreceptors fail to differentiate. Anti-PCNA antibody

labels retinal stem and progenitor cells in the CMZ (Fig. 1-figure supplement 1K),

which was not affected in *rw337* mutants (Fig. 1-figure supplement 1L). Phalloidin

labeling shows IPL and OPL in wild-type retinas (Fig. 1-figure supplement 1M). *rw337*

mutants showed IPL, but not OPL (Fig. 1-figure supplement 1N), which is consistent

with zpr1 labeling (Fig. 1-figure supplement 1H). Thus, retinal stem and progenitor

cells are maintained. RGCs do differentiate; however, amacrine cells, horizontal cells,

bipolar cells, and Müller cells are decreased in number and photoreceptors fail to

differentiate in *rw337* mutants.

***rw337* mutant gene encodes Banp**

We mapped the *rw337* mutation to zebrafish chromosome 25 (Fig. 1C). Two

polymorphic markers, 070702B and zC93F2D, restricted the genomic region

encompassing *rw337* mutation, in which only four candidate genes: *BTG3-*

associated nuclear protein (*banp*); *programmed cell death 2-like* (*pdcd2l*);

carboxylesterase 2b (*ces2b*); *protein disulfide isomerase family A, member 3* (*pdia3*),

were annotated. We cloned all these cDNAs from mRNA prepared from *rw337*

homozygous mutants, sequenced them, and found that a part of the intron sequence

between exons 4 and 5 (intron 4/5) failed to be spliced out of the *banp* cDNA (Fig. 1D).

A nucleotide T at the exon-intron junction of exon 4 is converted to G, which results in

an alteration of the donor splice site of exon 4 from GT to GG. This mutation leads to

the abnormal insertion of a 52-bp intron sequence after exon 4, resulting in a premature

stop codon in *banp* cDNA (Fig. 1D). The *banp* gene encodes a protein containing a

coiled-coil domain, a BEN domain and a DNA-binding domain, which includes a

MAR-binding domain (Fig. 1E). Overall, zebrafish Banp protein shows high similarity

to those of humans and mice, especially in the BEN domain (Fig. 1-figure supplement

2). This mutation is predicted to generate nonfunctional, truncated Banp protein in

rw337 mutants (Fig. 1E). There was no mutation in the other three genes in *rw337*

mutants, suggesting that the *rw337* mutant gene encodes Banp.

To confirm that the loss of Banp causes retinal phenotypes in *rw337* mutants, we

carried out three sets of experiments. First, we examined another nonsense mutation

allele, *banp^{sa12976}*, which has a premature stop codon in exon 4 (Fig. 1-figure

supplement 3A). Homozygous *banp^{sa12976}* mutant embryos and trans-heterozygous

banp^{rw337}; *banp^{sa12976}* mutants phenocopied *rw337* mutants (Fig. 1-figure supplement

3B). Second, we inhibited Banp using ATG-morpholino antisense oligos against Banp

(banp MO). We confirmed MO specificity, because banp MO inhibited translation of

mRNA encoding C-terminal EGFP-tagged wild-type Banp (banp(wt)-EGFP), whereas

the control, 5-mismatch banp MO (banp 5misMO) did not (Fig. 1-figure supplement 4A

and 4B). *banp* morphants phenocopied *rw337* mutants (Fig. 1F). Third, we

overexpressed wild-type Banp in *banp^{rw337}* mutants. We injected mRNA encoding

Banp(wt)-EGFP into wild-type embryos at the one-cell stage. In the developing retinas,

Banp(wt)-EGFP protein was localized in the nuclei during interphase, whereas it spread

into the whole cell region during mitosis (Fig. 1-figure supplement 4C and 4D),

confirming nuclear localization of zebrafish Banp. We expressed Banp(wt)-EGFP, N-

terminal EGFP tagged wild-type (EGFP-Banp(wt)) and the *rw337* mutant form of Banp

(EGFP-Banp(rw337)) in wild-type retinas. Only EGFP-Banp(rw337) was not stably

maintained (Fig. 1-figure supplement 4E), suggesting that the mutant form of Banp

protein is degraded, probably by nonsense mutation decay. Finally, overexpression of

Banp(wt)-EGFP rescued retinal phenotypes in *rw337* mutants at 2 dpf, but

overexpression of EGFP-Banp(rw337) did not (Fig. 1G). Taken together, retinal

phenotypes of *rw337* mutants are caused by the loss of Banp.

***banp* mRNA is prominently expressed in the developing CNS of zebrafish**

We examined *banp* mRNA expression by *in-situ* hybridization (Fig. 1-figure

supplement 5A). *banp* mRNA shows ubiquitous expression from the one-cell stage to

the 21-somite stage, suggesting maternal and zygotic expression. mRNA was restricted

to the brain, including the retina at 30 and 48 hpf. Frontal sections show that mRNA is

expressed in neural retina at 24 hpf, but restricted in the CMZ at 48 hpf, suggesting that
retinal progenitor cells express *banp* mRNA (Fig. 1-figure supplement 5B). In addition
to the brain and retinas, *banp* mRNA expression was observed in fin and notochord at
48 hpf and in neuromasts of the lateral line after 72 hpf. *banp* mRNA expression was
not detected in *banp^{rw337}* homozygous mutants at 3 dpf (Fig. 1-figure supplement 5C),
suggesting nonsense mutation-mediated decay. Thus, *banp* mRNA is prominently
expressed in the developing brain in zebrafish.

banp mutants show cell cycle arrest followed by cell death of retinal progenitor
cells

Although retinal progenitor cells produce RGCs in *banp^{rw337}* mutant retinas, the number
of other neurons, especially photoreceptors, was decreased at 4 dpf (Fig. 1-figure
supplement 1). Indeed, mitotic-like round cells and pyknotic nuclei were accumulated in
the outer region of *banp^{rw337}* mutant retinas from 2 to 3 dpf (Fig. 1B), suggesting that
mitotic arrest and cell death eliminate retinal neurons. Accordingly, we performed TdT-
mediated dUTP Nick End Labeling (TUNEL), which visualizes apoptosis. TUNEL+
cells were rarely observed in wild-type retinas throughout the stages from 29 to 101 hpf,
as well as in *banp^{rw337}* mutant retinas at 24 hpf (Fig. 2A). However, in *banp^{rw337}* mutants,
TUNEL+ cells appeared in the neural retina at 53 hpf, and became more localized to the
outer retina at 77hpf (Fig. 2A) and to the interface between the central retina and the
CMZ at 77 and 101 hpf (Fig. 2A). After 53 hpf, the number of TUNEL+ cells was
significantly higher in *banp^{rw337}* mutants than in wild-type siblings (Fig. 2C). Thus,
retinal cells undergo apoptosis in *banp^{rw337}* mutants. Next, we examined mitosis by
labeling with anti-phosphorylated (Ser-10) Histone H3 (pH3) antibody, which labels
mitotic cells in pro-metaphase (Prigent and Dimitrov, 2003) (Fig. 2B). The number of
pH3+ cells did not differ between wild-type and *banp^{rw337}* mutant retinas at 24 hpf.
However, pH3+ cells were markedly increased and located at the apical surface of the
neural retina in *banp^{rw337}* mutant retinas at 53 hpf (Fig. 2B), suggesting mitotic arrest. At
77 and 101 hpf, pH3+ cells disappeared from the central retina, but still were
accumulated in the CMZ in *banp^{rw337}* mutants (Fig. 2B). The number of pH3+ cells in
banp^{rw337} mutant retinas increased transiently at 53 hpf, and then decreased, but was

still significantly higher than that of wild-type sibling retinas after 77 hpf because of

accumulation of pH3+ cells at the CMZ (Fig. 2D).

To clarify how mitotic arrest and apoptosis are induced during retinal cell

proliferation and neurogenesis in $banp^{rw337}$ mutants, we examined BrdU incorporation

and ath5:EGFP expression at 48 hpf (Fig. 2-figure supplement 1A and 1B). The

percentage of BrdU+ area to total retinal area did not differ significantly between wild-

type siblings and $banp^{rw337}$ mutants (Fig. 2-figure supplement 1C), whereas the

percentage of ath5:EGFP+ area to the total retinal area was slightly, but significantly

lower in $banp^{rw337}$ mutants than in wild-type siblings (Fig. 2-figure supplement 1C),

suggesting that retinal neurogenesis is slightly reduced in $banp^{rw337}$ mutants at 48 hpf,

probably due to mitotic arrest or apoptosis.

Next, we examined the spatial position of mitotic arrest and apoptosis at 48 hpf

by labeling with anti-pH3 and anti-activated caspase 3 antibodies (Fig. 2-figure

supplement 1D and 1E). Both caspase 3+ and pH3+ cells were increased in $banp^{rw337}$

mutants (Fig. 2-figure supplement 1F). Furthermore, caspase 3+ cells were mostly

located in the intermediate zone of the neural retina along the apico-basal axis, whereas

pH3+ cells were localized at the apical surface of the neural retina (Fig. 2-figure

supplement 1G), suggesting that apoptosis occurs mostly in retinal progenitor cells

undergoing G1 or S phase, or differentiating neurons. Next, we examined whether

apoptosis occurs in early differentiating retinal neurons, using double labeling with anti-

activated caspase 3 antibody and ath5:EGFP (Fig. 2-figure supplement 1H). The

percentage of ath5:EGFP- and caspase 3-double-positive cells to the total number of

caspase 3+ cells was 9.44±13.1% in $banp$ mutants at 48 hpf and higher than that of

wild-type siblings (0.00±0.00%), although the difference was not significant (Fig. 2-

figure supplement 1I). Next, we examined whether apoptosis occurs in mature retinal

neurons. HuC/D is expressed in maturing neurons (Park et al., 2000). At 48 hpf, anti-

HuC/D antibody labeled RGCs in zebrafish (Fig. 2-figure supplement 1J). The fraction

of caspase 3- and HuC/D-double-positive cells in the total caspase 3+ cells was almost

zero in both wild-type and $banp^{rw337}$ mutants (Fig. 2-figure supplement 1K), suggesting

that apoptosis does not occur in maturing retinal neurons. Taken together, in $banp^{rw337}$

mutants, retinal progenitor cells undergo mitotic arrest at 48 hpf. On the other hand,

apoptosis is induced in retinal progenitor cells at G1 or S-phase or in newly

differentiating retinal neurons, although apoptosis in newly differentiating retinal

neurons is minor (Fig. 2-figure supplement 1L). Since mitotic arrest was transiently

observed at 53 hpf (Fig. 2B), but apoptosis continues in the central retina of *banp*[rw337]

mutants after 53 hpf (Fig. 2A), it is likely that apoptosis occurs following mitotic arrest

(Fig. 2E and 2F).

tp53 pathway mediates retinal apoptosis in *banp* mutants

To understand molecular mechanisms of mitotic arrest and apoptosis in *banp*[rw337]

mutants, we identified differentially expressed mRNAs by RNA-seq analysis. We

prepared mRNA from 48-hpf wild-type siblings and *banp*[rw337] mutants. Illumina

sequencing generated more than 50 million paired-end reads from cDNA libraries.

27,771 protein-coding genes were detected. Among them, 258 genes were upregulated,

and 81 genes were downregulated in *banp*[rw337] mutants (Fig. 3A). The top 15

upregulated genes were enriched regulators of tp53 dependent-DNA damage response

(Fig. 3B). Quantitative RT-PCR revealed that four tp53 target genes: *p21(cdkn1a),*

puma (bbc3), *ccng1*, and *mdm2* are significantly upregulated in *banp*[rw337] mutants (Fig.

3C). Western blotting showed that full length tp53 (FL tp53) protein is stabilized and

more than 11 fold higher in *banp*[rw337] mutants than in wild type (Fig. 3D and 3E). We

previously showed that the ATR/ATM-Chk2-tp53 DNA damage pathway induces

retinal apoptosis in zebrafish (Yamaguchi et al., 2008). Indeed, mRNA expression of

ATR and ATM was upregulated in *banp*[rw337] mutants (Fig. 3F). So, we examined

whether retinal apoptosis depends on tp53 in *banp*[rw337] mutants. Injection of tp53

antisense morpholino (tp53 MO) completely inhibited retinal apoptosis in *banp*[rw337]

mutants at 48 hpf (Fig. 3G and 3H). However, retinal apoptosis reappeared in *banp*[rw337]

mutants injected with tp53 MO at 72 hpf (Fig. 3G), although there was no statistical

difference in apoptotic cell number between wild-type and *banp*[rw337] mutant retinas at

72 hpf (Fig. 3H). Thus, retinal apoptosis in *banp*[rw337] mutants depends on tp53, although

it is possible that some tp53-independent mechanism induces retinal apoptosis after 48

hpf.

Transcription of Δ113-tp53 is preferentially elevated in *banp* mutants

RNA-seq analysis showed upregulation of tp53 mRNA (Fig. 3B); however, surprisingly,

quantitative RT-PCR showed that the mRNA level of FL tp53 is not significantly

different between $banp^{rw337}$ mutants and wild-type siblings at 48 hpf (Fig. 4A). There

are several isoforms of tp53 protein (Joruiz et al., 2020; Joruiz and Bourdon, 2016).

Importantly, in zebrafish, the isoform Δ113tp53 is generated from the alternative

transcription start site in intron 4. Δ113tp53 preferentially activates the transcription of

target genes related to cell-cycle arrest to promote DNA repair and cell survival,

whereas FL tp53 preferentially activates transcription of target apoptotic genes to

eliminate severely DNA-damaged cells (Gong et al., 2015; Gong et al., 2020).

Differential expression of tp53 isoforms is thought to regulate the balance between cell

survival and apoptosis in response to DNA damage. Indeed, quantitative RT-PCR

confirmed that only Δ113tp53 was upregulated in $banp^{rw337}$ mutants (Fig. 4A). ATAC-

seq revealed that a genomic region in intron 4 of tp53 is specifically open in $banp^{rw337}$

mutants but not in wild-type siblings (Fig. 4B, ATAC-seq). Consistently, transcription

of the Δ113tp53 mRNA was specifically activated in $banp^{rw337}$ mutants, but not in wild-

type siblings (Fig. 4B, RNA-seq). Thus, the internal promoter at intron 4 (Chen et al.,

2009; Chen et al., 2005) is specifically activated in $banp^{rw337}$ mutants (Fig. 4B). These

data raise the possibility that the lack of Banp increases DNA replication stress and

promotes an accumulation of DNA damage, resulting in activation of ATR and ATM,

which stabilize FL tp53, such that FL tp53 triggers Δ113tp53 transcription in $banp^{rw337}$

mutants (Fig. 4G).

DNA double-strand breaks accumulate in *banp* mutant retinas

To examine whether DNA damage is increased due to loss of Banp functions, we

labeled 48-hpf retinas with anti-γ-H2AX antibody, which marks DNA double-strand

breaks (Rogakou et al., 1998; Shroff et al., 2004). The number of γ-H2AX+ cells

increased in both $banp^{rw337}$ mutant (Fig. 4C, 4F) and *banp* morphant retinas (Fig. 4-

figure supplement 1A, 1B), confirming an accumulation of DNA damage in the absence

of Banp. Next, to determine how DNA damage is induced in $banp^{rw337}$ mutants, we

examined the fraction of S-phase cells, mitotic cells, and post-mitotic neurons in γ-

H2AX+ retinal cells in $banp^{rw337}$ mutants at 48 hpf. First, wild-type and $banp^{rw337}$

mutant retinas carrying the transgene *Tg[ath5:EGFP]* were labeled with anti-BrdU and

anti-γ-H2AX antibodies (Fig. 4-figure supplement 1C). The fraction of γ-H2AX+ cells

increased in *banp^{rw337}* mutant retinas, whereas the fraction was almost zero in wild-type

sibling retinas (Fig. 4-figure supplement 1D). Next, we compared the fraction of BrdU+

cells between the total retinal cells and γ-H2AX+ retinal cells in *banp^{rw337}* mutants (Fig.

4-figure supplement 1E). The BrdU+ fraction was significantly higher in γ-H2AX+

cells than in the total retinal cells (Fig. 4-figure supplement 1F), indicating that S-phase

fraction is 4.3 fold higher in DNA damaged retinal cells than in the total retinal cells, in

banp^{rw337} mutants. Thus, DNA double-strand breaks are abnormally induced or fail to be

repaired in retinal progenitor cells undergoing S phase. Next, we examined the fraction

of ath5:EGFP+ cells (Fig. 4-figure supplement 1G). There was no significant difference

in ath5:EGFP+ cell fraction between the total retinal cells and γ-H2AX+ retinal cells in

banp^{rw337} mutants (Fig. 4-figure supplement 1H). Finally, we determined the fraction of

mitotic cells by labeling with anti-pH3 and anti-γ-H2AX antibodies in *banp^{rw337}* mutant

retinas at 48 hpf (Fig. 4-figure supplement 1I). There is no significant difference in

pH3+ cell fraction between the total retinal cells and γ-H2AX+ retinal cells (Fig. 4-

figure supplement 1J). Although DNA damage may occur in post mitotic neurons and

mitotic cells in *banp^{rw337}* mutants, it is likely that most DNA damage is introduced in S-

phase of retinal progenitor cells in the absence of Banp.

It is generally accepted that tp53 promotes DNA damage repair or eliminates

DNA-damaged cells via apoptosis, so we expected that tp53 knockdown may enhance

accumulation of DNA damage in *banp^{rw337}* mutants. However, compared with standard-

MO injection control (n=120.8±60.6/section), the injection of FL tp53-MO significantly

reduced   the   number   of   γ-H2AX+   cells   in   *banp^{rw337}*   mutant   retinas

(n=55.9±15.9/section) at 48 hpf (Fig. 4C, 4F). The injection of Δ113tp53-MO mildly

reduced the number of γ-H2AX+ cells (n=85.4±15.2/section); however, the difference

was not significant (Fig. 4C, 4F). It is likely that tp53 or Δ113tp53 does not repair DNA

damage in *banp^{rw337}* mutants, but rather it may enhance DNA damage in *banp^{rw337}*

mutants. Thus, Banp normally suppresses DNA replication stress upstream of tp53, or

promotes DNA damage repair in concert with tp53 (Fig. 4G).

In addition, we investigated whether retinal apoptosis and mitotic arrest in

banp^{rw337} mutants depend on FL tp53-MO or Δ113tp53-MO. Compared with Standard-

MO injection control (n=174.7±30.5/section), retinal apoptosis was completely

inhibited by FL tp53-MO in *banp^rw337* mutant retinas (n=0.50±0.58/section) at 48 hpf

(Fig. 4C, 4D). However, inhibition of apoptosis in *banp^rw337* mutant retinas by

Δ113tp53-MO was significant, but ineffective (n=58.00±11.0/section) (Fig. 4C, 4D).

This result is consistent with previous reports proposing that Δ113tp53 preferentially

activates transcription of target genes related to cell-cycle arrest rather than apoptosis

(Gong et al., 2015; Gong et al., 2020). However, compared with Standard-MO injection

controls (n=376.6±74.8/section), the number of pH3+ cells in *banp^rw337* mutants was

significantly decreased by FL tp53-MO (n=262.4±68.3/section) (Fig. 4C, 4E). However,

the number of pH3+ cells was still higher in *banp^rw337* mutant retinas injected with FL

tp53-MO than in wild-type control retinas injected with FL tp53-MO. The injection of

Δ113tp53-MO did not significantly rescue mitotic arrest in *banp^rw337* mutants

(n=303.8±46.6/section) (Fig. 4C, 4F). These data suggest that mitotic arrest in *banp^rw337*

mutants is mostly mediated by a tp53-independent mechanism.

378   Banp is required for the expression of chromatin segregation genes

Mitotic arrest in *banp^rw337* mutants is independent of tp53. RNA-seq analysis revealed

that two chromosomal segregation regulators, *cenpt* and *ncapg*, are downregulated in

banp^rw337 mutants at 48 hpf (Fig. 3B). Cenpt and Ncapg are essential for chromatin

segregation (Hung et al., 2017; Seipold et al., 2009; Zhang et al., 2018). Quantitative

RT-PCR confirmed their downregulation in *banp^rw337* mutants and *banp* morphants (Fig.

5A, 5C). Furthermore, ATAC-seq revealed that chromatin accessibility is decreased

near the transcription start site (TSS) of both genes in *banp^rw337* mutants, leading to

reduction of transcription activity (Fig. 5B, 5D). Thus, Banp is required for

transcriptional activation of *cenpt* and *ncapg* genes. Interestingly, two independent chip-

sequencing studies using Banp antibody revealed a specific binding site within 100 bp

of the TSS of *cenpt* and *ncapg* genes in mice and humans (Grand et al., 2021; Mathai et

al., 2016), suggesting that the requirement for Banp in *cenpt* and *ncapg* transcription is

conserved in zebrafish, mice, and humans. Furthermore, using motif discovery software

(HOMER), we discovered that TCTCGCGAGA, which was reported to be the Banp

motif (Grand et al., 2021), is the most enriched motif in the closed chromatin region of

banp^rw337 mutants (Fig. 5E, Fig. 5-figure supplement 1A). Indeed, among 81 down-

regulated genes in *banp^rw337* mutants (Fig. 3A), we identified the Banp motif near the

TSS of 31 genes (Fig. 5F and Supplementary table 1), 26 % of which was cell-cycle

regulators (Fig. 5G, 5H) including *cenpt* and *ncapg* (Fig. 5I and 5J). However, we

could not find the Banp motif in 258 upregulated genes. Furthermore, the Banp motif

occurs upstream of the TSS of both *cenpt* and *ncapg* in mice and humans (Fig. 5-figure

supplement 1B, 1C). Their Banp motif position is consistent with the previous report,

that the Banp motif is not located at a constant distance from TSSs, but rather between

20 and 70 nucleotides upstream on both strands, and that the frequency of the Banp

motif occurrence upstream of TSSs may correlate to a transcription rate of target genes

(Mahpour et al., 2018) (Supplementary table 1). Taken together, it is likely that Banp

directly regulates transcription of *cenpt* and *ncapg* via the Banp motif.

Chromosomal segregation is abnormal in *banp* mutants

Knockdown of Cenpt and Ncapg causes defects in chromosomal segregation during

mitosis in zebrafish (Hung et al., 2017; Seipold et al., 2009). So, we conducted time-

lapse imaging of mitosis in *banp*^{rw337} mutant retinas for 1-2 h from 48 to 54 hpf, using a

transgenic line *Tg[h2afv:GFP; EF1α:mCherry-zGem]*, which visualizes cell-cycle

phases (Mochizuki et al., 2014). In wild-type retinas injected with control banp

5misMO, chromatin starts to be condensed in prophase, kinetochores and a mitotic

spindle appear in prometaphase, chromosomes are centrally aligned and start to be

pulled toward the opposite poles by mitotic spindles in metaphase, sister chromatids

move toward the opposite poles in anaphase, and chromosomes arrive at the opposite

poles in telophase, whereupon cytokinesis starts (Fig. 6A, Video 1). The duration from

prophase to telophase was 22.9±5.8 minutes (Fig. 6B). On the other hand, mitosis did

not proceed smoothly in *banp* morphant retinas, especially from prophase to anaphase

(Fig. 6A, Video 2). It took longer from chromosomal condensation in prophase to

chromosomal alignment at the division plane in anaphase in *banp* morphant retinas

(t=15 to t=46 in banp-MO, Fig. 6A). Aligned chromosomes were often dispersed

during chromosomal segregation by the mitotic spindle (t=46 to t=64 in banp-MO, Fig.

6A), although all 28 cell divisions eventually completed cytokinesis in *banp* morphants.

The duration from prophase to telophase was 49.2± 17.0 minutes and significantly

longer in *banp* morphants (Fig. 6B). Thus, chromosomal segregation is abnormal during

mitosis in the absence of Banp. Furthermore, we labeled 48-hpf retinas with anti-α-

tubulin and anti-pH3 antibodies to visualize the mitotic spindle and prometaphase

chromosomes, respectively. In *banp^{rw337}* mutants, mitotic spindles are irregularly

attached to less condensed and misaligned chromosome (Fig. 6C). The percentage of

cells with abnormal mitosis was significantly higher in *banp^{rw337}* mutant retinas than

those of wild-type siblings (Fig. 6D). These data suggest that the interaction between

chromosomes and mitotic spindles is compromised in the absence of Banp, probably

due to kinetochore attachment defects caused by lack of Cenpt and Ncapg. Thus, Banp

regulates chromosomal segregation during mitosis.

Taken together, Banp plays essential roles in cell-cycle regulation in the

zebrafish developing retina. During the S phase, Banp suppresses DNA replication

stress and promotes DNA damage repair in concert with tp53-mediated DNA damage

response. In the M phase, Banp promotes mitosis by activating the transcription of two

chromosome segregation regulators, *cenpt* and *ncapg*, most likely via their Banp motifs

(Fig. 7).

Discussion

Banp was initially identified as a nuclear matrix-associated protein and reported to function as a tumor suppressor (Malonia et al., 2011). Very recently, it was reported that Banp binds to the Banp motif enriched near the TSS of CpG island promoters and regulates transcription of metabolic genes (Grand et al., 2021). However, physiological functions of Banp during embryonic development remain to be discovered. In this study, we identified zebrafish *banp* mutants, in which retinal progenitor cells show mitotic arrest and subsequent apoptosis during embryonic development. RNA-seq analysis revealed that expression of tp53-dependent DNA damage response regulators is upregulated in *banp* mutants. In a non-stress condition, tp53 is degraded through ubiquitination by an E3-ubiqutin ligase, Mdm2, and subsequent actions of proteasomes, keeping its protein level very low (Oliner et al., 1993). On the other hand, under stress, such as DNA replication stalling or DNA double-strand breaks, ATR and ATM are activated and phosphorylate the N-terminal serine of tp53 (Saito et al., 2002; Tibbetts et al., 1999). This phosphorylation reduces the binding affinity of Mdm2 for tp53, resulting in stabilization of tp53 protein (Yadahalli et al., 2019). Indeed, the tp53 protein level is increased in *banp* mutants, and mRNA expression of ATR and ATM is also elevated. tp53 knockdown effectively suppresses retinal apoptosis in *banp* mutants. These data suggest that DNA replication stress is elevated in *banp* mutants, resulting in activation of tp53-dependent DNA damage response, which subsequently induces apoptosis of retinal progenitor cells.

It is generally accepted that tp53 has dual, but opposing roles: cell survival versus cell death in response to DNA replication stress and DNA damage (Chen, 2016). In the initial phase of the DNA damage response, tp53 binds to its alternative promoter located in intron 4 and activates transcription of an N-terminal truncated isoform Δ113-tp53, which preferentially activates transcription of target genes related to cell-cycle arrest for promoting DNA repair and cell survival (Chen et al., 2009). However, if DNA damage fails to be repaired, but accumulates chronically, FL tp53, in turn, activates transcription of apoptotic genes for eliminating DNA-damaged cells. Previous *in vitro* studies of human *banp* suggest that Banp physically interacts with tp53 to promote transcription of cell-cycle arrest genes and suppresses transcription of apoptotic genes, leading to promotion of its cell survival role (Jalota et al., 2005; Sinha et al., 2010).

Indeed, our quantitative RT-PCR showed that transcription of Δ113-tp53 is significantly
elevated in *banp* mutants, whereas the mRNA level of FL tp53 is very low. RNA-seq
and ATAC-seq also confirmed a higher level of transcription and chromatin
accessibility of Δ113tp53 in *banp* mutants. These data suggest that Δ113tp53-mediated
transcription is selectively activated in *banp* mutants. Thus, it is important to clarify the
role of tp53, including Δ113tp53 in mitotic arrest in *banp* mutants, because pH3+ cells
are markedly accumulated in *banp* mutant retinas prior to apoptosis. However, tp53-MO
mildly inhibited accumulation of pH3+ cells in *banp* mutants. There is no significant
difference in pH3+ cell accumulation between tp53-MO and Δ113tp53-MO, or between
wild-type siblings and Δ113tp53-MO. So, these data suggest that the contribution of
tp53 and Δ113tp53 to mitotic arrest in *banp* mutants seems to be minor, and that a tp53-
independent pathway is responsible for mitotic arrest phenotypes. Our live imaging of
mitosis in *banp* morphant retinas reveals that mitosis is significantly longer due to
chromosome segregation defects; however, importantly, almost all mitoses are
eventually completed in *banp* mutants, suggesting that accumulation of pH3+ cells in
banp mutants is transient and depends mostly on prolonged mitosis, which is caused by
chromosome segregation defects. At present, we have not observed S-phase arrest in
banp mutants, because the fraction of BrdU+ cells in the total retinal region is not
increased in *banp* mutants, although mRNA expression of a cdk inhibitor, *cdkn1a* (*p21*),
is higher in *banp* mutants than in wild-type siblings. Taken together, we conclude that
tp53 mainly mediates retinal apoptosis, and that its role in cell-cycle arrest is minor in
banp mutants.

Labeling with anti-γ-H2AX antibody revealed that DNA damage is increased in
retinal progenitor cells in *banp* mutants, suggesting that DNA damage is abnormally
introduced or fails to be repaired in the absence of Banp. Importantly, the level of DNA
damage in *banp* mutant retinas is not increased by either tp53-MO and Δ113tp53-MO,
suggesting that tp53 does not repair DNA damage without Banp. There are two possible
functions of Banp in DNA damage regulation. Banp may maintain the integrity of DNA
replication. Banp functions as a transcription factor to regulate expression of essential
metabolic genes in pluripotent stem cells and differentiating neurons (Grand et al.,
2021). Therefore, Banp may promote transcription of DNA replication regulators. Our
RNA-seq data identified 81 downregulated genes in *banp* mutants, compared to wild-

type siblings. Among them, 31 genes have a Banp motif near their TSSs and contain

eight cell-cycle regulatory genes. It will be interesting to examine whether these Banp

targets are involved in the activation of DNA replication stress in *banp* mutants.

Another possibility is that Banp is required for DNA repair. A previous study using

human *in vitro* culture cells revealed that Banp promotes HDAC6-mediated

deacetylation of a DNA end joining protein Ku70, which is required for non-

homologous end-joining repair (Chaudhary et al., 2014). Since Ku70 is recruited to the

double-strand damage site after γ-H2AX accumulation, Ku70 is an interesting candidate.

Since knockdown of Ku70 along with radiation-induced DNA damage markedly

induces apoptosis in zebrafish embryos (Bladen et al., 2007), it will be interesting to

investigate whether this scenario is the case in zebrafish.

In addition to the role of Banp in DNA replication stress and DNA repair

mechanisms, mRNA expression of two important regulators of chromosome

segregation, *ncapg* and *cenpt*, is significantly downregulated in *banp* mutants. ATAC-

seq and RNA-seq confirmed that chromatin accessibility is decreased near the TSSs of

these genes in *banp* mutants, suppressing transcription of these mRNAs. Our *de novo*

motif enrichment study identified the Banp motif sequence, TCTCGCGAGA, as the

most enriched motif of closed chromatin regions of downregulated genes. Thus, it is

highly likely that Banp directly binds to the Banp motifs of both *ncapg* and *cenpt*, opens

their chromatin, and promotes their transcription. Consistently, as reported in zebrafish

ncapg and *cenpt* knockdown embryos (Hung et al., 2017; Seipold et al., 2009),

chromosome segregation is compromised in mitosis of retinal progenitor cells in *banp*

mutants, suggesting that Banp is an essential regulator of mitosis. Since a Banp motif is

found in the promoter region of *ncapg* and *cenpt* in mice and humans, and Chip

sequencing with Banp antibody validated TSS peaks of both *ncapg* and *cenpt* genes in

humans and mice (Grand et al., 2021; Mathai et al., 2016), it will be interesting to

confirm whether Banp regulates chromosome segregation during mitosis in mammals.

Lastly, we identified 26 candidates of Banp direct target genes, including *ncapg*

and *cenpt,* which carry the Banp motifs near their TSSs, and show markedly decreased

chromatin accessibility and transcription in the absence of Banp. This gene list will

provide a useful starting point to investigate the Banp-mediated transcriptional network.

It was reported that the binding of Banp to the Banp motif is dependent on DNA

methylation of the Banp motifs (Grand et al., 2021), suggesting that an epigenetic mechanism regulates Banp-mediated transcription. Thus, zebrafish *banp* mutants provide an excellent new model in which to study how epigenetic factors regulate cellular processes for embryonic development via the Banp motifs of CpG island promoter genes. In addition, Banp has been proposed as a tumor suppressor (Malonia et al., 2011). It is generally accepted that DNA methylation of essential genes is altered in cancer cells (Baylin et al., 2001), which is expected to modify a pattern of gene transcription through Banp motifs (Grand et al., 2021). It will be interesting to investigate how the Banp-mediated transcription profile is changed in cancer models of zebrafish. Future research on zebrafish *banp* mutants will provide more in-depth understanding of cancer growth and metastasis, as well as new insights into therapy development of tumors in human.

Material and Methods

Zebrafish strains and maintenance

Zebrafish (*Danio rerio*) were maintained using standard procedures (Westerfield, 1993).

RIKEN Wako (RW) was used as a wild-type strain for mutagenesis in which *banp*[rw337]

was generated. Mapping of the *rw337* mutation and analysis of retinal phenotypes of

banp[rw337] mutants were carried out in the genetic background of WIK and Okinawa

wild-type (oki), respectively. Two *banp* mutant alleles, *banp*[rw337] and *banp*[sa12976] were

used. *banp*[sa12976] was obtained from the Zebrafish International Resource Center (ZIRC,

Eugene, OR, USA). Zebrafish transgenic lines, *Tg[h2afv:GFP]*[kca6/kca66] (Pauls et al.,

2001), *Tg[EF1α:mCherry-zGem]*[oki011] (Mochizuki et al., 2014) and *Tg[EF1α:mCherry-*

CAAX][oki049] (Mochizuki et al., 2017) were used to visualize chromatin, S/G2/M phases

and plasma membranes, respectively.

Ethics statement

All zebrafish experiments were carried out in accordance with the Okinawa Institute of

Science and Technology Graduate School's Animal Care and Use Program, which is

based on the National Research Council of the National Academy's Guide for the Care

and Use of Laboratory Animals and has been accredited by the Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC International).

The OIST Institutional Animal Care and Use Committee approved all experimental

protocols.

Zebrafish mutagenesis, mapping and cloning of the *banp* mutant gene

Mutagenesis, mapping, and cloning were performed as previously described (Masai et

al., 2003). The polymorphic markers, 070702B and zC93F2D, shown below were

designed and used for restricting the genomic region covering the *banp*[rw337] mutation.

070702B forward: 5'-ACTTCTTATCAGGGCTGTGC-3'

070702B reverse: 5'-TCAGTCAAGAGCAGTGAGAG-3'

zC93F2D forward: 5'-TGGGATCTCTTTAAGTGAGTGAG-3'

zC93F2D reverse: 5'-TCCAACTATGTGGGTCAAACC-3'

Genotyping of *banp*^{rw337} mutant embryos

banp^{rw337} mutant phenotypes were difficult to distinguish until 2 dpf. Identification of homozygous mutant embryos at earlier time points was performed using PCR amplification of genomic regions using specific primers that enabled restriction digestion of only mutated sequence using MboII. Primers used, forward: 5'-CGATGTTGATATCCATCAGTCAGGCGATC-3'; reverse primer: 5'-GGTGCTGGTGTATAAATCACATGACCTATGGTCCTCTT-3'.

Acridine Orange staining

Larvae were incubated at 28°C, in 5 mg/mL Acridine Orange (AO) in E3 embryonic medium (5mM NaCl, 0.17mM KCl, 0.33mM $CaCl_2$, 0.33mM $MgSO_4$) for 30 min, protected from light. Embryos were rinsed in E3 medium and viewed with an epifluorescence microscope (SteREO Discovery, Zeiss) to identify live mutant embryos with dead cell accumulation.

In situ **hybridization**

Whole-mount *in situ* hybridization of zebrafish embryos was performed as previously described (Xu et al., 1994). To prepare RNA probes, full-length cDNAs of zebrafish *banp* gene were amplified with PCR and sub-cloned into pBluescript II SK(+) vector (Stratagene/Agilent Technologies). Plasmids were digested with an appropriate restriction enzyme and used for DNA templates to make antisense or sense RNA probes. Digoxigenin-labeled RNA probes were generated by *in vitro* transcription using a DIG RNA Labeling Kit (Roche) and purified using Micro Bio-Spin® Chromatography Columns (BioRad). *In situ* hybridization signals in whole-mount embryos were detected using anti-DIG alkaline phosphatase (AP)-conjugated antibody (Roche) with an AP substrate, BM-Purple (Roche) and imaged using SteREO Discovery.V12 (ZEISS) microscope.

Rescue experiment of *banp* mutants with overexpression of wild-type Banp

The full-length coding region of the *banp* gene was amplified with PCR. To overexpress wild-type Banp protein, a DNA construct encoding zebrafish wild-type

Banp with an EGFP tag at the N-terminus or C-terminus, EGFP-banp(wt) or banp(wt)-

EGFP, was prepared. A DNA construct encoding an N-terminal EGFP-tagged *rw337*

mutant form of Banp, EGFP-Banp(rw337), was prepared as a negative control. These

DNA constructs were subcloned into the pCS2 expression vector. Capped mRNA was

generated in a standard *in vitro* transcription reaction using an SP6 polymerase

(mMESSAGE mMACHINE™ SP6 Transcription Kit, Invitrogen). We performed

mRNA injections in one-cell-stage embryos obtained from pair-wise crosses of

banp^{rw337} heterozygous mutant male and female fish. At 2 dpf, embryos were used to

determine their genotype with PCR and to identify homozygous mutants. Plastic

sections were obtained to evaluate retinal phenotypes of *banp^{rw337}* mutant embryos

injected with mRNA encoding banp(wt)-EGFP, or EGFP-banp(rw337). We also

evaluated protein stability and nuclear localization of wild-type and *rw337* mutant form

of Banp by live confocal imaging of wild-type retinas expressing banp(wt)-EGFP,

EGFP-banp(wt), or EGFP-banp(rw337) at 24 hpf.

Morpholino injection

ATG morpholino antisense oligos for *banp* (banp MO) and its 5 mismatch *banp* MO

(banp 5misMO) were designed as 5'-CCACTAAATCTTGCTCTGACATCAT-3' and

5'-CCtCaAAATgTTcCTCTcACATCAT-3', respectively. Morpholino antisense oligos

for tp53 (tp53 MO) (5'-GCGCCATTGCTTTGCAAGAATTG-3') (Langheinrich et al.,

2002), standard MO (STD MO) (5'-CCTCTTACCTCAGTTACAATTTATA-3') were

used. To knock down Δ113tp53, morpholino antisense oligos for Δ113tp53 (Δ113tp53

MO; also referred to as MO6-tp53 in zebrafish database ZFIN) (5'-

GCAAGTTTTTGCCAGCTGACAGAAG-3') was used (Chen et al., 2009). All MOs

were injected into one-cell-stage eggs at 1 mM.

Histology

Plastic sectioning, immunostaining of cryosections, paraffin sectioning, and BrdU

incorporation were performed as described previously (Imai et al., 2010), except that we

injected BrdU solution into the yolk of embryos at 48 hpf instead of soaking embryos in

10mM BrdU solution. Anti- pH3 antibody (Cell signalling technology, 6G3, 9706;

Sigma-Aldrich, 06-570), anti-Pax6 antibody (Covance; PRB-278P), zpr1 antibody

(ZIRC, Eugene, Oregon), anti-Prox1 antibody (GeneTex, GTX128354), anti-glutamine

synthetase (GS) (Millipore, MAB302), anti-BrdU antibody (abcam, Ab6326), anti-

PCNA (Sigma-Aldrich, P8825) were used at 1:500, 1:500, 1:100, 1:500, 1:100, 1:200,

1:200 dilutions, respectively. Nuclear staining was performed using SYTOX Green

(Molecular Probes). Filamentous actin (F-actin) was stained using 0.1 mM rhodamine-

conjugated phalloidin (Molecular Probes). Terminal deoxynucleotide transferase-

mediated dUTP nick-end labeling (TUNEL) was performed using the In Situ Cell Death

Detection Kit (Roche).

Whole-mount immunostaining with anti-α-tubulin antibody (Sigma-Aldrich;

#T5168) to label the mitotic spindle was performed as follows. Homozygous mutant

embryos were selected by AO staining, and fixed in 2% trichloroacetic acid (TCA) at

room temperature (RT) for 3 h. Then embryos were washed three times in 1X PBST

(PBS, 0.1 % tritonx100), and incubated in 0.2% trypsin for 4 min at 4^{o}C to increase

antibody permeabilization through the skin. Then, embryos were rinsed thrice with PBS.

After embryos were re-fixed with 4% PFA for 5 min at 4^{o}C, they were incubated in

blocking solution (10% Goat serum in PBST) for 1 h at RT and transferred to 1%

blocking solution with diluted anti-α-tubulin antibody at 1:1000 dilution for 2 days at

$4°$C. Embryos were washed with PBST four times for 15 min and then incubated with

fluorescent conjugated anti-mouse antibody in 1% blocking solution in the dark for 2

days at $4°$C. Next, all samples were washed in PBST three times for 15 min followed by

glycerol storage at 4^{o}C until imaging. Samples were imaged with a confocal microscope

LSM 880 (Zeiss) followed by image analysis using IMARIS software (ver.9.1.2

Bitplane). Whole-mount immunostaining for anti-pH3, anti-γ-H2AX (GeneTex,

GTX127342) (Sidi et al., 2008), anti-active caspase3 (BD Pharmingen, Clone C92-

605), anti-HuC/D (Thermo Fisher, A-21271) (Sorrells et al., 2013) were performed as

described previously (Imai et al., 2010). Images were scanned using an LSM710

confocal laser scanning microscope (Zeiss). Number of fluorescent-positive cells and

percentage of fluorescence-positive area relative to the region of interest was calculated

using original scanning images with IMARIS (Bitplane) and Zen (Zeiss) or Image-J

(NIH) software, respectively.

RT-PCR

Total RNA was extracted from embryo heads at the indicated developmental stages
using a Sepasol RNA I Super (Nacalai, Japan). cDNA synthesis was carried out from
500ng of RNA using Toyobo ReverseTra Ace® qPCR RT master mix with gDNA
remover according to the manufacturer's instructions. Expression profiles of the genes
listed below were analyzed with the specified primers using Luna® Universal qPCR
Master Mix (NEB). Quantitative RT-PCR was performed using StepOnePlus (Applied
Biosystems). The relative expression was calculated using ΔΔCt method with the
reference gene ef1α. Primers used in qRT-PCR analysis; mdm2 (Wilkins et al., 2013),
p21 (Stiff et al., 2016), FL tp53 and Δ113 tp53 (Chen et al., 2016), ef1α (McCurley and
Callard, 2008).

Gene	RT PCR-Forward primer	RT PCR-Reverse primer
ccng1	ccctggagattgaggatcag	cacacaaaccaggtctccaa
mdm2	caggaggaggagaagcagtg	agggaaaagctgtccgactt
p21(cdkn1a)	aagcgcaaacagaccaacat	tcagctactggccggattt
FL tp53	tggagaggaggtcggcaaaatcaa	gactgcgggaacctgagcctaaat
Δ113 tp53	atatcctggcgaacatttggaggg	cctcctggtcttgtaatgtcac
Puma (bbc3)	ctgaggaggaccccacact	tctccagttctgccagtgc
cenpt	tcatgaggagattgtggaagatg	ggtgagctctgcgagttatt
ncapg	ctgatgtgagggagcctattt	gagtctgtttggcctccatta
atm	cctcaaggctgtggagaact	aggggattttctttacaccactc
atr	aggaacccaatctgccagt	gatgtccagtgccagctctc
ef1α	cttctcaggctgactgtgc	ccgctagcattaccctcc

Western blotting

Total 21 embryos were divided into 3 groups (each n=7) for 48 hpf wild-type and
banp[rw337] mutants, respectively. Heads were dissected in ice-cold E3 medium. Seven
heads removed from each group were homogenized together in sample buffer (125 mM
NaCl, 50 mM Tris-HCl pH 7.4, 0.5 mM EDTA, 1% Triton X-100, 1X Protease
inhibitor) and used as one sample. Appropriate volumes of samples were diluted to
provide equal protein amounts and used for SDS-PAGE (BIO-RAD, Mini-PROTEAN

TGX Gels). Western blot analysis was performed to 3 samples for wild-type and

banp[rw337] mutants, respectively, according to standard protocols using anti-tp53

antibody (GeneTex, GTX128135) at a 1:1000 dilution. β-actin protein level was

evaluated by western blot analysis of rehybridized membranes with anti-β-actin

antibody (Sigma, A5441) at a 1:5000 dilution, and used to normalize tp53 protein level

for each sample. For secondary antibodies, horseradish peroxidase-conjugated anti-

rabbit or anti-mouse IgG antibodies (Amersham, NA931 and NA934) were used.

Enzyme activity was detected using Immunostar® LD (Wako) and luminescence was

imaged and quantified using iBright 1500 (Thermo Fisher Scientific).

Live imaging

Banp-MO was injected into one-cell-stage eggs of *Tg[h2afv:GFP; EF1α: mCherry-*

zGem] transgenic line. Banp-5misMO was used as a control. Two dpf embryos were

anesthetized using 5% (v/v) Tricane (3-amino benzoic acid-ethylester) and mounted

laterally using 1% low-melting agarose on a depression slide to image retinas without

left or right preference. Time-lapse images were obtained every 1.1 min using a

confocal laser scanning microscope LSM710 (Zeiss) with 40X0.8/1.0W Plane

Apochromat lens. Data were evaluated using IMARIS software (ver.9.1.2 Bitplane) to

calculate the time taken to complete each cell division.

RNA sequencing

Embryo heads were homogenized using a Polytron 1200E homogenizer

(KINEMATICA) with TRIzol reagent (Invitrogen). RNA was isolated using a Direct-

zol RNA Miniprep Kit (ZYMO RESEARCH) from three replicates of *banp*[rw337] mutant

and wild-type embryos at 48 hpf. Each sample contained a pool of 7 embryo heads.

RNA isolation was followed by library preparation for sequencing using an NEBNext®

Ultra™ II Directional RNA Library Prep Kit for Illumina® according to the

manufacturer's instructions. 151-base paired-end sequencing was performed using a

NovaSeq6000 SP. The quality of sequenced reads was evaluated with FastQC (Andrews,

2010) and adaptor trimming and quality filtering were done with fastp (Chen et al.,

2018). Cleaned reads were then mapped to the zebrafish reference genome (GRCz11)

using hisat2.1.0 (Kim et al., 2019) and mapped reads were counted with FeatureCounts

from the Subread package (Liao et al., 2014) using annotation from Ensembl

(Danio_rerio.GRCz11.95.gtf). Differentially expressed gene (DEG) analysis was

performed using the EdgeR package (Robinson and Oshlack, 2010). Genes with false

discovery rates (FDR) < 0.01 were considered significant DEGs. The EnhancedVolcano

package (Blighe et al., 2018) was used to create a volcano plot with $Log_2FoldChange$

(wildtype – mutant) values and FDR calculated with EdgeR. Differences in alternative

splicing between mutant and wild type was tested using the DEXseq package (Anders et

al., 2012). Adjusted p-value < 0.01 was considered as significant.

ATAC sequencing

48-hpf embryos were used for ATAC sequencing. We used the Kaestner Lab ATAC seq

protocol (Ackermann and Kaestner, 2019) until library preparation, followed by

sequencing using a NovaSeq6000 SP. We generated reads for three independent

$banp^{rw337}$ mutant samples, as well as three independent $banp^{rw337}$ wild-type sibling

samples. Each sample contained a pool of 3 embryo heads. A quality check of reads was

performed using FastQC. Reads from each sample were adaptor trimmed and aligned

using fastp and Bowtie2 to the zebrafish reference genome (GRCz11) (Langmead and

Salzberg, 2012) respectively. For quality control, aligned reads were filtered for low-

quality reads, non-unique alignments, and PCR duplicates using samtools (Li et al.,

2009; Yan et al., 2020). Moreover, alignment files were peak-called using Genrich, with

removal of mitochondrial reads (Gaspar, 2019). Differences in differentially enriched

peaks in wild-type and mutant embryos were identified using the DiffBind R package

(Stark and Brown, 2021). The obtained peaks were annotated using the annotatePeaks

tool of HOMER software (Heinz et al., 2010). The findMotifsGenome tool of HOMER

(Hypergeometric Optimization of Motif EnRichment) was used to find enriched motifs

in differentially expressed peaks. To compare coverage tracks across genes of interest in

$banp^{rw337}$ mutants and wild-type fish in RNA-seq and ATAC-seq data, distributions of

mapped reads in each replicate were averaged and plotted with SparK (Kurtenbach and

Harbour, 2019).

Statistics

Significance of differences between groups (wild type vs mutant) was tested using

GraphPad Prism (GraphPad Software Version 9.1.2). The number of samples used per

experiment is represented within figures or figure legends. Tests used to calculate

statistical significance are mentioned in figure legends. All data are presented as means

$\pm$ SDs. A p value of 0.05 was considered statistically significant. In the histogram, p*<

0.05, p**< 0.01, p***< 0.001, p****< 0.0001, or ns (not significant) is indicated.

Acknowledgements

We thank four former lab members, Masahiro Yamaguchi, Noriko Tonou-Fujimori,

Yukihiro Yoshimura, Eri Oguri for supporting the initial phase of mapping and cloning

experiments, and three technical staff members, Jeff Liner, Mamoru Fujiwara, and

Tetsuya Harakuni for supporting mutant fish maintenance, immunohistochemistry, and

DNA construction. We thank SQC, Research support division, Okinawa Institute of

Science and Technology Graduate University for assistance in sequencing experiments.

We thank Steven D. Aird for editing the manuscript.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: SB, YT, IM; Methodology: SB, YT, IM; Software: SB, YT, IM;

Validation: SB, YT, IM; Formal analysis: SB, YT, IM; Investigation: SB, YT, IM;

Resources: SB, YT, IM; Data curation: SB, YT, IM; Writing - original draft: SB, IM;

Writing - review & editing: SB, IM; Visualization: SB, IM; Supervision: IM; Project

administration: SB, IM; Funding acquisition: IM.

Funding

This work was supported by a grant from the Okinawa Institute of Science and

Technology Graduate University to IM.

Data availability

Raw RNA-seq and ATAC-seq datasets of *banp*[rw337] mutant and wild-type siblings are

available at DDBJ Sequence Read Archive (DRA012572).

References

References

Ackermann, A., and L. Kaestner. 2019. **UPDATED** ATAC-seq Protocol.

Anders, S., A. Reyes, and W. Huber. 2012. Detecting differential usage of exons from
RNA-seq data. *Genome Research*. 22:2008-2017.

Andrews, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.

Baylin, S.B., M. Esteller, M.R. Rountree, K.E. Bachman, K. Schuebel, and J.G. Herman.
2001. Aberrant patterns of DNA methylation, chromatin formation and gene
expression in cancer. *Hum Mol Genet*. 10:687-692.

Berti, M., D. Cortez, and M. Lopes. 2020. The plasticity of DNA replication forks in
response to clinically relevant genotoxic stress. *Nat Rev Mol Cell Bio*. 21:633-
651.

Bhagat, P.N., S.H. Jadhav, S. Chattopadhyay, and K.M. Paknikar. 2018. Carbon
nanospheres mediated nuclear delivery of SMAR1 protein (DNA binding
domain) controls breast tumor in mice model. *Nanomedicine (Lond)*. 13:353-
372.

Birot, A., L. Duret, L. Bartholin, B. Santalucia, I. Tigaud, J. Magaud, and J. Rouault. 2000.
Identification and molecular analysis of BANP. *Gene*. 253:189-196.

Blackford, A.N., and S.P. Jackson. 2017. ATM, ATR, and DNA-PK: The Trinity at the
Heart of the DNA Damage Response. *Molecular Cell*. 66:801-817.

Bladen, C.L., W.K. Lam, W.S. Dynan, and D.J. Kozlowski. 2005. DNA damage response
and Ku80 function in the vertebrate embryo. *Nucleic Acids Res*. 33:3002-3010.

Bladen, C.L., S. Navarre, W.S. Dynan, and D.J. Kozlowski. 2007. Expression of the Ku70
subunit (XRCC6) and protection from low dose ionizing radiation during
zebrafish embryogenesis. *Neurosci Lett*. 422:97-102.

Blasquez, V.C., A.O. Sperry, P.N. Cockerill, and W.T. Garrard. 1989. Protein - DNA
Interactions at Chromosomal Loop Attachment Sites. *Genome*. 31:503-509.

Blighe, K., S. Rana, and M. Lewis. 2018. EnhancedVolcano: Publication-ready volcano
plots with enhanced colouring and labeling.

Boije, H., R.B. MacDonald, and W.A. Harris. 2014. Reconciling competence and
transcriptional hierarchies with stochasticity in retinal lineages. *Curr Opin
Neurobiol*. 27:68-74.

Chattopadhyay, S., R. Kaul, A. Charest, D. Housman, and J.Z. Chen. 2000. SMAR1, a
novel, alternatively spliced gene product, binds the scaffold/matrix-associated
region at the T cell receptor beta locus. *Genomics*. 68:93-96.

Chaudhary, N., K.K. Nakka, P.L. Chavali, J. Bhat, S. Chatterjee, and S. Chattopadhyay.
2014. SMAR1 coordinates HDAC6-induced deacetylation of Ku70 and dictates
cell fate upon irradiation. *Cell Death Dis*. 5:e1447.

Chemmannur, S.V., A.J. Badhwar, B. Mirlekar, S.K. Malonia, M. Gupta, N. Wadhwa, R.
Bopanna, U. Mabalirajan, S. Majumdar, B. Ghosh, and S. Chattopadhyay. 2015.
Nuclear matrix binding protein SMAR1 regulates T-cell differentiation and
allergic airway disease. *Mucosal Immunol*. 8:1201-1211.

Chen, J. 2016. The Cell-Cycle Arrest and Apoptotic Functions of p53 in Tumor Initiation
and Progression. *Cold Spring Harb Perspect Med*. 6:a026104.

Chen, J., S.M. Ng, C. Chang, Z. Zhang, J.C. Bourdon, D.P. Lane, and J. Peng. 2009. p53
isoform delta113p53 is a p53 target gene that antagonizes p53 apoptotic
activity via BclxL activation in zebrafish. *Genes Dev*. 23:278-290.
Chen, J., H. Ruan, S.M. Ng, C. Gao, H.M. Soo, W. Wu, Z. Zhang, Z. Wen, D.P. Lane, and J.
Peng. 2005. Loss of function of def selectively up-regulates Delta113p53
expression to arrest expansion growth of digestive organs in zebrafish. *Genes*
Dev. 19:2900-2911.
Chen, S.F., Y.Q. Zhou, Y.R. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*. 34:884-890.
Chen, Y.C., S. Semenova, S. Rozov, M. Sundvik, J.L. Bonkowsky, and P. Panula. 2016. A
Novel Developmental Role for Dopaminergic Signaling to Specify Hypothalamic
Neurotransmitter Identity. *Journal of Biological Chemistry*. 291:21880-21892.
Dowling, J.E. 2012. The Retina, An Approachable Part of the Brain. Harvard University
press.
Gaspar, J.M. 2019. ATAC-seq Guidelines.
Gong, L., H. Gong, X. Pan, C. Chang, Z. Ou, S. Ye, L. Yin, L. Yang, T. Tao, Z. Zhang, C. Liu,
D.P. Lane, J. Peng, and J. Chen. 2015. p53 isoform Delta113p53/Delta133p53
promotes DNA double-strand break repair to protect cell from death and
senescence in response to DNA damage. *Cell Res*. 25:351-369.
Gong, L., X. Pan, G.K. Abali, J.B. Little, and Z.M. Yuan. 2020. Functional interplay
between p53 and Delta 133p53 in adaptive stress response. *Cell Death and*
Differentiation. 27:1618-1632.
Grand, R.S., L. Burger, C. Grawe, A.K. Michael, L. Isbel, D. Hess, L. Hoerner, V.
Iesmantavicius, S. Durdu, M. Pregnolato, A.R. Krebs, S.A. Smallwood, N. Thoma,
M. Vermeulen, and D. Schubeler. 2021. BANP opens chromatin and activates
CpG-island-regulated genes. *Nature*.
Hart, C.M., and U.K. Laemmli. 1998. Facilitation of chromatin dynamics by SARs.
Current Opinion in Genetics & Development. 8:519-525.
Heinz, S., C. Benner, N. Spann, E. Bertolino, Y.C. Lin, P. Laslo, J.X. Cheng, C. Murre, H.
Singh, and C.K. Glass. 2010. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage
and B cell identities. *Mol Cell*. 38:576-589.
Hu, M., and S.S. Easter. 1999. Retinal neurogenesis: The formation of the initial central
patch of postmitotic cells. *Dev Biol*. 207:309-321.
Hu, Z., J. Holzschuh, and W. Driever. 2015. Loss of DDB1 Leads to Transcriptional p53
Pathway Activation in Proliferating Cells, Cell Cycle Deregulation, and Apoptosis
in Zebrafish Embryos. *PLoS One*. 10:e0134299.
Hung, C.Y., B. Volkmar, J.D. Baker, J.W. Bauer, E. Gussoni, S. Hainzl, A. Klausegger, J.
Lorenzo, I. Mihalek, O. Rittinger, M. Tekin, J.E. Dallman, and O.A. Bodamer.
2017. A defect in the inner kinetochore protein CENPT causes a new syndrome
of severe growth failure. *PLoS One*. 12:e0189324.
Imai, F., A. Yoshizawa, N. Fujimori-Tonou, K. Kawakami, and I. Masai. 2010. The
ubiquitin proteasome system is required for cell proliferation of the lens

epithelium and for differentiation of lens fiber cells in zebrafish. *Development*.
137:3257-3268.

Jalota, A., K. Singh, L. Pavithra, R. Kaul-Ghanekar, S. Jameel, and S. Chattopadhyay.
2005. Tumor suppressor SMAR1 activates and stabilizes p53 through its
arginine-serine-rich motif. *J Biol Chem*. 280:16019-16029.

Joruiz, S.M., J.A. Beck, I. Horikawa, and C.C. Harris. 2020. The Delta 133p53 Isoforms,
Tuners of the p53 Pathway. *Cancers*. 12.

Joruiz, S.M., and J.C. Bourdon. 2016. p53 Isoforms: Key Regulators of the Cell Fate
Decision. *Cold Spring Harb Perspect Med*. 6.

Jusuf, P.R., and W.A. Harris. 2009. Ptf1a is expressed transiently in all types of
amacrine cells in the embryonic zebrafish retina. *Neural Dev*. 4.

Kaul, R., S. Mukherjee, F. Ahmed, M.K. Bhat, R. Chhipa, S. Galande, and S.
Chattopadhyay. 2003. Direct interaction with and activation of p53 by SMAR1
retards cell-cycle progression at G2/M phase and delays tumor growth in mice.
Int J Cancer. 103:606-615.

Kaul-Ghanekar, R., S. Majumdar, A. Jalota, N. Gulati, N. Dubey, B. Saha, and S.
Chattopadhyay. 2005. Abnormal V(D)J recombination of T cell receptor beta
locus in SMAR1 transgenic mice. *J Biol Chem*. 280:9450-9459.

Kim, D., J.M. Paggi, C. Park, C. Bennett, and S.L. Salzberg. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*.
37:907-+.

Kurtenbach, S., and J.W. Harbour. 2019. SparK: A Publication-quality NGS Visualization
Tool. *bioRxiv*.

Langheinrich, U., E. Hennen, G. Stott, and G. Vacun. 2002. Zebrafish as a model
organism for the identification and characterization of drugs and genes
affecting p53 signaling. *Curr Biol*. 12:2023-2028.

Langmead, B., and S.L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods. 9:357-U354.

Larison, K.D., and R. Bremiller. 1990. Early Onset of Phenotype and Cell Patterning in
the Embryonic Zebrafish Retina. *Development*. 109:567-&.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.
Durbin, and G.P.D. Proc. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinformatics*. 25:2078-2079.

Li, Z., M. Hu, M.J. Ochocinska, N.M. Joseph, and S.S. Easter, Jr. 2000. Modulation of cell
proliferation in the embryonic retina of zebrafish (Danio rerio). *Dev Dyn*.
219:391-401.

Liao, Y., G.K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics*.
30:923-930.

Liu, H.C., F. Ma, Y. Shen, Y.Q. Hu, and S. Pan. 2014. Overexpression of SMAR1 Enhances
Radiosensitivity in Human Breast Cancer Cell Line MCF7 via Activation of p53
Signaling Pathway. *Oncol Res*. 22:293-300.

Liu, T.X., N.G. Howlett, M. Deng, D.M. Langenau, K. Hsu, J. Rhodes, J.P. Kanki, A.D.
D'Andrea, and A.T. Look. 2003. Knockdown of zebrafish Fancd2 causes
developmental abnormalities via p53-dependent apoptosis. *Dev Cell*. 5:903-914.
Macdonald, R., and S.W. Wilson. 1997. Distribution of Pax6 protein during eye
development suggests discrete roles in proliferative and differentiated visual
cells. *Dev Genes Evol*. 206:363-369.
Mahpour, A., B.S. Scruggs, D. Smiraglia, T. Ouchi, and I.H. Gelman. 2018. A methyl-
sensitive element induces bidirectional transcription in TATA-less CpG island-
associated promoters. *Plos One*. 13.
Malonia, S.K., S. Sinha, P. Lakshminarasimhan, K. Singh, A. Jalota-Badhwar, S. Rampalli,
R. Kaul-Ghanekar, and S. Chattopadhyay. 2011. Gene regulation by SMAR1:
Role in cellular homeostasis and cancer. *Biochim Biophys Acta*. 1815:1-12.
Masai, I., Z. Lele, M. Yamaguchi, A. Komori, A. Nakata, Y. Nishiwaki, H. Wada, H. Tanaka,
Y. Nojima, M. Hammerschmidt, S.W. Wilson, and H. Okamoto. 2003. N-
cadherin mediates retinal lamination, maintenance of forebrain compartments
and patterning of retinal neurites. *Development*. 130:2479-2494.
Masai, I., D.L. Stemple, H. Okamoto, and S.W. Wilson. 2000. Midline signals regulate
retinal neurogenesis in zebrafish. *Neuron*. 27:251-263.
Mathai, J., S.P. Mittal, A. Alam, P. Ranade, D. Mogare, S. Patel, S. Saxena, S. Ghorai, A.P.
Kulkarni, and S. Chattopadhyay. 2016. SMAR1 binds to T(C/G) repeat and
inhibits tumor progression by regulating miR-371-373 cluster. *Sci Rep*. 6:33779.
McCurley, A.T., and G.V. Callard. 2008. Characterization of housekeeping genes in
zebrafish: male-female differences and effects of tissue type, developmental
stage and chemical treatment. *Bmc Molecular Biology*. 9.
Mochizuki, T., Y.J. Luo, H.F. Tsai, A. Hagiwara, and I. Masai. 2017. Cell division and
cadherin-mediated adhesion regulate lens epithelial cell movement in zebrafish.
Development. 144:708-719.
Mochizuki, T., S. Suzuki, and I. Masai. 2014. Spatial pattern of cell geometry and cell-
division orientation in zebrafish lens epithelium. *Biol Open*. 3:982-994.
Murga, M., S. Bunting, M.F. Montana, R. Soria, F. Mulero, M. Canamero, Y. Lee, P.J.
McKinnon, A. Nussenzweig, and O. Fernandez-Capetillo. 2009. A mouse model
of ATR-Seckel shows embryonic replicative stress and accelerated aging. *Nat
Genet*. 41:891-U846.
Oliner, J.D., J.A. Pietenpol, S. Thiagalingam, J. Gyuris, K.W. Kinzler, and B. Vogelstein.
1993. Oncoprotein MDM2 conceals the activation domain of tumour
suppressor p53. *Nature*. 362:857-860.
Park, H.C., S.K. Hong, H.S. Kim, S.H. Kim, E.J. Yoon, C.H. Kim, N. Miki, and T.L. Huh. 2000.
Structural comparison of zebrafish elav/Hu and their differential expressions
during neurogenesis. *Neuroscience Letters*. 279:81-84.
Pauls, S., B. Geldmacher-Voss, and J.A. Campos-Ortega. 2001. A zebrafish histone
variant H2A.F/Z and a transgenic H2A.F/Z:GFP fusion protein for in vivo studies
of embryonic development. *Dev Genes Evol*. 211:603-610.

Pavithra, L., S. Mukherjee, K. Sreenath, S. Kar, K. Sakaguchi, S. Roy, and S.
974   Chattopadhyay. 2009. SMAR1 forms a ternary complex with p53-MDM2 and
975   negatively regulates p53-mediated transcription. *J Mol Biol*. 388:691-702.
976   Peterson, R.E., J.M. Fadool, J. McClintock, and P.J. Linser. 2001. Muller cell
977   differentiation in the zebrafish neural retina: Evidence of distinct early and late
978   stages in cell maturation. *Journal of Comparative Neurology*. 429:530-540.
979   Prigent, C., and S. Dimitrov. 2003. Phosphorylation of serine 10 in histone H3, what
980   for? *J Cell Sci*. 116:3677-3685.
981   Robinson, M.D., and A. Oshlack. 2010. A scaling normalization method for differential
982   expression analysis of RNA-seq data. *Genome Biol*. 11.
983   Rogakou, E.P., D.R. Pilch, A.H. Orr, V.S. Ivanova, and W.M. Bonner. 1998. DNA double-
984   stranded breaks induce histone H2AX phosphorylation on serine 139. *Journal of*
985   *Biological Chemistry*. 273:5858-5868.
986   Saito, S., A.A. Goodarzi, Y. Higashimoto, Y. Noda, S.P. Lees-Miller, E. Appella, and C.W.
987   Anderson. 2002. ATM mediates phosphorylation at multiple p53 sites, including
988   Ser(46), in response to ionizing radiation. *J Biol Chem*. 277:12491-12494.
989   Scheuermann, R.H., and W.T. Garrard. 1999. MARs of antigen receptor and co-
990   receptor genes. *Crit Rev Eukar Gene*. 9:295-310.
991   Seipold, S., F.C. Priller, P. Goldsmith, W.A. Harris, H. Baier, and S. Abdelilah-Seyfried.
992   2009. Non-SMC condensin I complex proteins control chromosome segregation
993   and survival of proliferating cells in the zebrafish neural retina. *BMC Dev Biol*.
994   9:40.
995   Shroff, R., A. Arbel-Eden, D. Pilch, G. Ira, W.M. Bonner, J.H. Petrini, J.E. Haber, and M.
996   Lichten. 2004. Distribution and dynamics of chromatin modification induced by
997   a defined DNA double-strand break. *Curr Biol*. 14:1703-1711.
998   Sidi, S., T. Sanda, R.D. Kennedy, A.T. Hagen, C.A. Jette, R. Hoffmans, J. Pascual, S.
999   Imamura, S. Kishi, J.F. Amatruda, J.P. Kanki, D.R. Green, A.A. D'Andrea, and A.T.
1000   Look. 2008. Chk1 suppresses a caspase-2 apoptotic response to DNA damage
1001   that bypasses p53, Bcl-2, and caspase-3. *Cell*. 133:864-877.
1002   Simossis, V.A., and J. Heringa. 2005. PRALINE: a multiple sequence alignment toolbox
1003   that integrates homology-extended and secondary structure information.
1004   *Nucleic Acids Research*. 33:W289-W294.
1005   Sinha, S., S.K. Malonia, S.P. Mittal, J. Mathai, J.K. Pal, and S. Chattopadhyay. 2012.
1006   Chromatin remodelling protein SMAR1 inhibits p53 dependent transactivation
1007   by regulating acetyl transferase p300. *Int J Biochem Cell Biol*. 44:46-52.
1008   Sinha, S., S.K. Malonia, S.P. Mittal, K. Singh, S. Kadreppa, R. Kamat, R. Mukhopadhyaya,
1009   J.K. Pal, and S. Chattopadhyay. 2010. Coordinated regulation of p53 apoptotic
1010   targets BAX and PUMA by SMAR1 through an identical MAR element. *EMBO J*.
1011   29:830-842.
1012   Sorrells, S., C. Toruno, R.A. Stewart, and C. Jette. 2013. Analysis of Apoptosis in
1013   Zebrafish Embryos by Whole-mount Immunofluorescence to Detect Activated
1014   Caspase 3. *Jove-J Vis Exp*.
1015   Stark, R., and G. Brown. 2021. DiffBind: Differential binding analysis of ChIP- Seq peak
1016   data.

Stiff, T., T.C. Tena, M. O'Driscoll, P.A. Jeggo, and M. Philipp. 2016. ATR promotes cilia
signalling: links to developmental impacts. *Hum Mol Genet*. 25:1574-1587.

Taye, N., A. Alam, S. Ghorai, D.G. Chatterji, A. Parulekar, D. Mogare, S. Singh, P.
Sengupta, S. Chatterjee, M.K. Bhat, M.K. Santra, P.B. Salunkhe, S.K. Finston, and
S. Chattopadhyay. 2018. SMAR1 inhibits Wnt/beta-catenin signaling and
prevents colorectal cancer progression. *Oncotarget*. 9:21322-21336.

Tibbetts, R.S., K.M. Brumbaugh, J.M. Williams, J.N. Sarkaria, W.A. Cliby, S.Y. Shieh, Y.
Taya, C. Prives, and R.T. Abraham. 1999. A role for ATR in the DNA damage-
induced phosphorylation of p53. *Genes Dev*. 13:152-157.

Tsuda, H., D.F. Callen, T. Fukutomi, Y. Nakamura, and S. Hirohashi. 1994. Allele loss on
chromosome 16q24.2-qter occurs frequently in breast cancers irrespectively of
differences in phenotype and extent of spread. *Cancer Res*. 54:513-517.

Wang, X., L. Gleich, Z.P. Pavelic, Y.Q. Li, N. Gale, S. Hunt, J.L. Gluckman, and P.J.
Stambrook. 1999. Cervical metastases of head and neck squamous cell
carcinoma correlate with loss of heterozygosity on chromosome 16q. *Int J
Oncol*. 14:557-561.

Waterman, D.P., J.E. Haber, and M.B. Smolka. 2020. Checkpoint Responses to DNA
Double-Strand Breaks. *Annu Rev Biochem*. 89:103-133.

Westerfield, M. 1993. The zebrafish book : a guide for the laboratory use of zebrafish
(Brachydanio rerio).

Wilkins, B.J., K. Lorent, R.P. Matthews, and M. Pack. 2013. p53-Mediated Biliary
Defects Caused by Knockdown of cirh1a, the Zebrafish Homolog of the Gene
Responsible for North American Indian Childhood Cirrhosis. *Plos One*. 8.

Xu, Q., N. Holder, R. Patient, and S.W. Wilson. 1994. Spatially regulated expression of
three receptor tyrosine kinase genes during gastrulation in the zebrafish.
Development. 120:287-299.

Yadahalli, S., J.L. Neira, C.M. Johnson, Y.S. Tan, P.J.E. Rowling, A. Chattopadhyay, C.S.
Verma, and L.S. Itzhaki. 2019. Kinetic and thermodynamic effects of
phosphorylation on p53 binding to MDM2. *Sci Rep*. 9:693.

Yamaguchi, M., N. Fujimori-Tonou, Y. Yoshimura, T. Kishi, H. Okamoto, and I. Masai.
2008. Mutation of DNA primase causes extensive apoptosis of retinal neurons
through the activation of DNA damage checkpoint and tumor suppressor p53.
Development. 135:1247-1257.

Yan, F., D.R. Powell, D.J. Curtis, and N.C. Wong. 2020. From reads to insight: a
hitchhiker's guide to ATAC-seq data analysis. *Genome Biol*. 21.

Zhang, Q., R. Su, C. Shan, C. Gao, and P. Wu. 2018. Non-SMC Condensin I Complex,
Subunit G (NCAPG) is a Novel Mitotic Gene Required for Hepatocellular Cancer
Cell Proliferation and Migration. *Oncol Res*. 26:269-276.

1055

Figure Legends

Figure 1. Zebrafish *banp^rw337* mutants show mitotic cell accumulation and cell death in the developing retina.

(A) Morphology of wild-type and *rw337* mutant embryos at 4 dpf. *rw337* mutant embryos show smaller eyes compared to wild type at 4 dpf (red arrow). The black arrow shows the cloudy tectum in *rw337* mutant embryos, representing cell death. Scale bars: 100 μm.

(B) Plastic sections of wild-type and *rw337* mutant retinas during development. Higher magnification images of green squares in the left columns are shown in the right columns. Retinal neuroepithelium is normal in *rw337* mutants until 2 dpf. However, many round cells reminiscent of mitotic cells are accumulated in the apical region of *rw337* mutant retinas at 2 dpf (open green arrowheads), suggesting mitotic arrest of retinal progenitor cells. In wild-type retinas, three nuclear layers and two plexiform layers are form at 2.5 dpf. However, only RGCs appear near the lens and the outer region was occupied by pyknotic nuclei, indicating dead cells in *rw337* mutants. At 3 dpf, INL is formed, but the outer region is still occupied by pyknotic nuclei in *rw337* mutants. At 4 dpf, pyknotic nuclei are reduced in *rw337* mutants; however, ONL and OPL are absent despite the small number of photoreceptor-like columnar cells in putative INL (PR, asterisks). At 3 and 4 dpf, the CMZ is maintained in *rw337* mutants (green double asterisks).

(C) The *rw337* mutation is mapped in the genomic region flanked by two polymorphic markers, 070702B and zC93F2D, on chromosome 25. This genomic region contains four genes including *banp*.

(D) *banp* cDNA consists of 13 exons. In *rw337* mutants, T at the donor splice site of the intron between exons 4 and 5 (intron4/5) is converted to G, leading to the insertion of 52 nucleotides (red region) from intron 4/5. A new stop codon appears within this insertion.

(E) Wild-type Banp protein has a coiled-coil domain, a BEN domain and a DNA-binding domain. An NLS domain, an RS-rich domain, and MAR-binding domain are shown. A truncated protein missing BEN and DNA-binding domains is predicted in *rw337* mutants.

(F) Retinal sections of wild-type, wild-type embryos injected with Banp-5misMO and

banp MO. *banp* morphant retinas have similar phenotypes to *banp^rw337* mutants,

with increased cell death and loss of the ONL. Yellow open arrowheads show

pyknotic nuclei.

(G) Retinal section of wild-type retinas, *banp^rw337* mutant retinas, and *banp^rw337* mutant

retinas overexpressing Banp(wt)-EGFP and EGFP-Banp(rw337). Overexpression of

Banp(wt)-EGFP inhibits cell death and recovers retinal lamination in *banp^rw337*

mutants, whereas overexpression of EGFP-Banp(rw337) does not inhibit retinal

defects in *banp^rw337* mutants. Yellow open arrowheads indicate pyknotic cells in

banp^rw337 mutant retinas and *banp^rw337* mutant retinas overexpressing EGFP-

Banp(rw337).

Scale bars: 20 μm for (B, F, G).

RGC, retinal ganglion cell; AC, amacrine cell, BP, bipolar cell; HC, horizontal cell; PR,

photoreceptor; OPL, outer plexiform layer; IPL, inner plexiform layer; INL, inner

nuclear layer; CMZ, ciliary marginal zone.

Figure Supplement 1. Retinal cell differentiation in *banp^rw337* mutants at 4 dpf.

Figure Supplement 2. Amino acid conservation of Banp protein between zebrafish,

mice and humans.

Figure Supplement 3. *banp^sa12976* homozygous mutants phenocopy *rw337* mutants.

Figure Supplement 4. Confirmation of specificity of banp MO and nuclear localization

of zebrafish Banp protein.

Figure Supplement 5. Spatio-temporal pattern of zebrafish *banp* mRNA expression.

1109

Figure 2. *banp* mutant retinas show mitotic arrest and apoptosis.

(A) TUNEL of wild-type (upper) and *banp^rw337* mutant (lower) retinas. In *banp^rw337*

mutants, apoptotic cells start to be observed in the neural retinas at 53 hpf, and

increase in the apical region of retinas (white arrowheads) and at the interface

between the CMZ and the central retina (yellow open arrowheads) at 77 and 101

hpf.

(B) Labeling of wild-type (upper) and *banp^rw337* mutant (lower) retinas with anti-pH3

antibody. Mitotic cells are accumulated at the apical surface of *banp^rw337* mutant

retinas at 53 hpf (white open arrowheads) and restricted in the apical region of the

CMZ (yellow open arrow) at 77 and 101 hpf.

(C) The number of TUNEL-positive cells per retina. Apoptosis is significantly

increased in *banp^{rw337}* mutants after 53 hpf. Unpaired t-test, two-tailed, Mean ± SD.

[n=3, p**< 0.01]

(D) The number of pH3-positive cells per retina. The number of mitotic cells is

significantly increased in *banp^{rw337}* mutants at 53 hpf. The number is decreased, but

still significantly higher in *banp^{rw337}* mutants than in wild-type siblings after 77 hpf.

Unpaired t-test, two-tailed, Mean ± SD. [n=3, p**< 0.01]

(E) Schematic diagram of wild-type retinas from 48 to 77 hpf. Neuronal differentiation

proceeds from the central to the peripheral retina. At 48 hpf, the proliferating area is

gradually restricted in the CMZ; however, there are still retinal progenitor cells that

proliferate in the central retina. Daughter cells start to exit from the cell cycle to

produce differentiated neurons (light green color). At 77 hpf, most progenitor cells

exit from the cell cycle, except at the CMZ. BrdU-labeled cells and pH3-positive

cells are indicated in blue and green, respectively.

(F) Schematic diagram of *banp^{rw337}* mutant retinas from 48 to 77 hpf. At 48 hpf, mitotic

cells are significantly accumulated at the apical surface of the neural retina and

apoptosis starts in the intermediate region of the central retina. At 77 hpf, the

apoptotic area is more restricted to the apical region of the retina and at the interface

between the central retina and the CMZ. Mitotic cells accumulate in the apical

region of the CMZ at 77 hpf. TUNEL-positive cells are indicated in magenta.

Scale bars: 20 μm for (A, B).

Source data 1. Data for Figure 2CD.

Figure Supplement 1. Apoptosis is induced in retinal progenitor cells and newly

differentiating neurons in *banp^{rw337}* mutants.

1144

1145 **Figure 3. tp53-mediated DNA damage response is activated in the absence of Banp.**

1146 (A) A volcano plot indicating gene differential expression of 48 hpf wild-type versus

1147   *banp^{rw337}* mutant embryos. Red dots show genes that were upregulated ≥2-fold in

1148   mutants, while blue dots represent genes downregulated ≥ 2-fold.

1149 (B) Heat map of the top 15 differentially expressed genes.

(C) Relative mRNA expression levels for *bbc3*, *ccng1*, *mdm2*, *cdkn1a* in wild type and *banp^{rw337}* mutants. Quantitative RT-PCR confirmed the up-regulation of p53 pathway genes in *banp^{rw337}* mutants at 48 hpf. Unpaired t-test (two-tailed), Mean ± SD. [n=3, p*< 0.05, p**< 0.01, p***< 0.001, p****< 0.0001].

(D) Western blotting of wild-type and *banp^{rw337}* mutants with anti-tp53 antibody. Full-length tp53 protein is stabilized in *banp^{rw337}* mutants, compared to wild type. β-actin protein is an endogenous control and used for normalization of tp53 protein level.

(E) Quantification of tp53 protein level relative to a control protein, β-actin, in wild-type and *banp^{rw337}* mutants. The tp53 protein level is significantly higher in *banp^{rw337}* mutants than in wild type at 48 hpf. Unpaired t-test (two-tailed), Mean ± SD. [n=3, p*< 0.05].

(F) Relative mRNA expression levels for *atm* and *atr* in wild type and *banp^{rw337}* mutants. Both mRNAs are significantly up-regulated in *banp^{rw337}* mutants at 48 hpf. Unpaired t-test (two-tailed) [n=3, p*< 0.05].

(G) TUNEL of wild-type and *banp^{rw337}* mutant retinas injected with STD MO and tp53 MO at 48 hpf and 72 hpf. Scale bars: 20 μm.

(H) Histogram of the number of TUNEL-positive cells per retinal section in wild-type and *banp^{rw337}* mutant retinas injected with STD MO and tp53 MO at 48 hpf and 72 hpf. Two way ANOVA, Turkey's multiple comparisons test, Mean ± SD. [n=3 for STD-MO at 48 hpf and tp53-MO at 72 hpf; n=4 for tp53-MO at 48 hpf and STD-MO at 72 hpf, p****< 0.0001].

Source data 1. Data for Figure 3CEFH.

Source data 2. Data for Figure 3D.

Figure 4. Banp is required for the integrity of DNA replication and DNA damage repair.

(A) Quantitative RT-PCR-based validation of FL tp53 and Δ113tp53 mRNA expression at 48 hpf. Δ113tp53 mRNA expression is markedly higher in *banp^{rw337}* mutants than in wild-type siblings, whereas FL tp53 mRNA is not significantly different between mutants and wild-type siblings. Unpaired t-test (two-tailed), Mean ± SD. [n=3, p**< 0.01, ns (not significant)].

(B) ATAC-seq- and RNA-seq-based validation of chromatin accessibility (upper panel)

and transcription level (bottom panel) of the genomic region covering both tp53

and Δ113tp53 transcription units. Data of wild-type and *banp^{rw337}* mutants are

indicated in blue and red color, respectively. Light blue indicates overlapping level.

The red box in ATAC-seq shows open chromatin in *banp^{rw337}* mutants in intron 4,

signifying differential transcription of Δ113tp53 using an internal promoter.

(C) TUNEL (top), anti-pH3 antibody labeling (middle), and anti-γ-H2AX antibody

labeling (bottom) of wild-type and *banp^{rw337}* mutant retinas injected with STD MO,

tp53 MO, and Δ113tp53 MO. tp53 MO-mediated knockdown completely rescues

apoptosis in *banp^{rw337}* mutant retinas at 48 hpf. However, Δ113tp53 MO-mediated

knockdown mildly inhibits apoptosis in *banp^{rw337}* mutant retinas. Neither tp53 MO

nor Δ113tp53 MO effectively rescued mitotic arrest and accumulation of γ-H2AX-

positive cells (white arrowheads) in *banp^{rw337}* mutant retinas. Scale bars: 20 μm.

(D–F) Histogram of the number of TUNEL-positive cells (D), pH3-positive cells (E),

and γ-H2AX-positive cells (F) per retinal section in wild-type and *banp^{rw337}* mutants

injected with STD-MO, tp53 MO, and Δ113tp53 MO. Two way ANOVA with Tukey's

multiple comparisons test, Mean ± SD. [n= in graph, p*< 0.05, p**< 0.01, p***< 0.001,

p****< 0.0001, ns (not significant)].

(G) Possible role of Banp in tp53-mediated the DNA damage response pathway. DNA

replication stress and subsequent DNA damage activate ATR and ATM, respectively,

which subsequently inhibit Mdm2-mediated tp53 degradation. Stabilized tp53 mainly

promotes transcription of the Δ113tp53 isoform, which activates cell-cycle arrest genes

and promotes DNA damage repair. Chronic activation of tp53 activates apoptotic genes

to induce apoptosis. Banp may normally suppress DNA replication stress and also

promote DNA damage repair.

Source data 1. Data for Figure 4ADEF.

Figure supplement 1. DNA damage is induced in retinal progenitor cells in *banp*

knockdown.

1209

Figure 5. Banp promotes transcription of chromosome segregation regulators,

cenpt **and** *ncapg***.**

(A) Quantitative RT-PCR of *ncapg* expression at 48 hpf. *ncapg* mRNA expression is

down-regulated in *banp^{rw337}* mutants and *banp* morphants. Unpaired t-test (two-

tailed), Mean ± SD. [n=3, p**< 0.01].

(B) ATAC-seq and RNA-seq analysis of *ncapg* mRNA transcription. In *banp^{rw337}*

mutants, chromatin access is reduced at the TSS of *ncapg* (ATAC-seq) and

transcript expression is markedly decreased (RNA seq).

(C) Quantitative RT-PCR of *cenpt* expression at 48 hpf. *cenpt* mRNA expression is

down-regulated in *banp^{rw337}* mutants and *banp* morphants. Unpaired t-test (two-

tailed), Mean ± SD. [n=3, p**< 0.01].

(D) ATAC-seq and RNA-seq analysis on *cenpt* mRNA transcription. In *banp^{rw337}*

mutants, chromatin access is reduced at TSS of *cenpt* (ATAC-seq) and transcript

expression is markedly decreased (RNA seq).

(E) The motif enriched in genomic regions in which chromatin accessibility is reduced

in *banp^{rw337}* mutants. This sequence shows a high similarity to the Banp motif.

(F) A schematic diagram representing the flow of RNA-sequencing combined with

ATAC-sequencing to find candidates for zebrafish Banp direct target genes.

HOMER was used to scan for Banp motifs in downregulated candidate genes, from

RNA- and ATAC-sequencing in *banp^{rw337}* mutants. RNA sequencing yielded 31

candidates, whereas ATAC sequencing yielded 72, with 26 target genes consistent

in both. These 26 candidate genes are the most likely to be target genes of Banp.

(G) STRING interactome analysis showing the interaction of Banp motif-containing

genes downregulated in RNA sequencing. Black arrows specify genes related to

cell-cycle regulation.

(H) Banp target candidate genes related to cell-cycle regulation. Gene ontology (GO)

analysis was applied to 31 Banp motif-containing genes downregulated in RNA

sequencing. 26% of these genes are engaged in the cell-cycle regulation. Others

participate in a variety of non-categorized pathways.

(I) Banp motif upstream to the 5'-UTR of *ncapg*.

(J) Banp motif upstream to the 5'-UTR of *cenpt.*

Source data 1. Data for Figure 5AC.

Figure supplement 1. Motifs enriched in the closed and open chromatin area in

banp^{rw337} mutants.

1244

Figure 6. Banp is required for chromosomal segregation during mitosis.

(A) Time-lapse images of mitosis of retinal progenitor cells injected with banp 5misMO

(upper) and banp MO (lower) at 2 dpf using *Tg [h2afv: GFP; EF1α: mCherry-*

zGem] transgenic zebrafish. In retinas injected with banp 5misMO, mitosis proceeds

sequentially from prophase (t=5~10 min), metaphase (t=16 min), anaphase (t=21

min), and telophase (t=24 min). On the other hand, in *banp* morphant retinas, the

duration of mitosis from prophase to telophase was 67 min, significantly longer than

that of the control MO (24 min). Furthermore, a prophase-like chromosome

arrangement appeared abnormally (t=59) after several metaphase-like alignments

(t=36 and 46), followed by anaphase (t=64) and telophase. Scale bar: 5μm.

(B) Violin plot of the duration of mitosis from prophase to telophase in retinas injected

with banp 5misMO and banp MO. Mitosis in *banp* morphants is significantly longer

than that of banp 5misMO injected embryos. Mann Whitney test, Mean ± SD. [n=3,

p****< 0.0001]

(C) Confocal images of metaphase cells in zebrafish retinas at 48 hpf. The mitotic

spindle is labeled with anti-α-tubulin antibody, whereas metaphase chromosomes

are labeled with anti-pH3 antibody. The mitotic spindle is correctly attached to

metaphase chromosomes in wild-type retinal cells (left panel). On the other hand,

mitotic cells in *banp*^rw337^ mutants exhibit inefficient mitotic spindle attachment and

poorly orientated metaphase chromosomes (right panels). Dotted circles indicate the

outline of metaphase cells. Scale bar: 2μm.

(D) Percentage of cells with normal and abnormal spindle attachment in wild-type and

banp^rw337^ mutant embryos. Two way ANOVA with Šídák's multiple comparisons

test, Mean ± SD. [n=3, p****< 0.0001].

Source data 1. Data for Figure 6BD.

1270

1271 **Figure 7. A possible model of Banp functions in DNA damage repair and**

1272 **chromosome segregation during mitosis.**

1273 Our study reveals two novel roles of Banp in cell-cycle regulation of retinal progenitor

1274 cells. First, Banp suppresses DNA replication stress and promotes DNA damage repair.

In the absence of Banp, a DNA replication stress sensor, ATR, is activated, leading to accumulation of DNA double-strand breaks, which activate ATM and the tp53-mediated DNA damage response pathway. tp53 activates the apoptotic pathway and promotes transcription of Δ113tp53, both of which promote cell-cycle arrest and DNA repair. Second, Banp is required for chromosome segregation during mitosis by promoting transcription of two mitotic regulators, *cenpt* and *ncapg*. Since a Banp motif is found near the TSSs of these genes, it is likely that *cenpt* and *ncapg* are direct targets of Banp.

 Supplementary Figure Legends

 Figure 1-figure supplement 1. Retinal cell differentiation in *banp*^{rw337} mutants at 4

 dpf.

(A, B) ath5:EGFP expression in wild-type (A) and *rw337* mutant (B) retinas at 4 dpf.

ath5:EGFP is strongly expressed in RGCs and weakly expressed in amacrine cells and

photoreceptor precursors. In *rw337* mutants, ath5:EGFP expression in RGCs, amacrine

cells and photoreceptor precursors is detected.

(C, D) Wild-type (C) and *rw337* mutant (D) retinas labeled with anti-Pax6 antibody at 4

dpf. Pax6 is strongly expressed in amacrine cells and weakly expressed in RGCs. In

rw337 mutants, amacrine cells seem to be decreased in number.

(E, F) Wild-type (E) and *rw337* mutant (F) retinas labeled with anti-Prox1 antibody at 4

dpf. Prox1is expressed in bipolar cells and horizontal cells. In *rw337* mutants, both

bipolar-like round nuclei and horizontal-like flat nuclei are observed, but horizontal-like

cells are very few.

(G, H) Wild-type (G) and *rw337* mutant (H) retinas labeled with zpr1 antibody at 4 dpf.

zpr1 antibody labels double-cone-type photoreceptors in zebrafish. In *rw337* mutants,

only a small cluster of zpr1-positive cells is observed in the ventro-nasal retina,

suggesting that a majority of photoreceptors fail to differentiate in *rw337* mutants.

(I, J) Wild-type (I) and *rw337* mutant (J) retinas labeled with anti-GS antibody at 4 dpf.

Anti-GS antibody labels Müller cells. In *rw337* mutants, Müller cells are decreased in

number.

(K, L) Wild-type (K) and *rw337* mutant (L) retinas labeled with anti-PCNA antibody at

4 dpf. Anti-PCNA antibody labels retinal progenitor cells in the CMZ at 4 dpf. In *rw337*

mutants, retinal progenitor cells are maintained in the CMZ.

(M, N) Phalloidin labeling of wild-type (M) and *rw337* mutant (N) retinas. Phalloidin

visualizes retinal synaptic layers, the IPL and OPL. In *rw337* mutants, the IPL is

detected, but not fully formed on the ventral side. Furthermore, there is no OPL, which

is consistent with the lack of photoreceptors in *rw337* mutants. Scale bars: 20μm.

RGC, retinal ganglion cell, AC, amacrine cell; BP, bipolar cell; HC, horizontal cell; PR,

photoreceptor; MG, Müller cell; RPC, retinal progenitor cell; OPL, outer plexiform

layer; IPL, inner plexiform layer.

1316

Figure 1-figure supplement 2. Amino acid conservation of Banp protein between zebrafish, mice and humans.

1319 Alignment of Banp protein of humans (upper, ENSP00000376902), mice (middle, ENSMUSP00000026) and zebrafish (bottom, ENSDARP00000023) using a multiple sequence alignment program, PRALINE (Simossis and Heringa, 2005). Amino acid conservation is indicated by scoring and color coding from the least conserved (0, blue) to the most conserved (10, red). The BEN domain is indicated.

1324

Figure 1-figure supplement 3. *banp^{sa12976}* homozygous mutants phenocopy *rw337* mutants.

1327 (A) Another *banp* mutant allele, *sa12976*, carries a nonsense mutation in exon 4.

1328 (B) Comparison of phenotypes between *banp* mutant alleles. *sa12976* homozygous *banp* mutant alleles and trans-heterozygous *rw337/sa12976* mutants show similar phenotypes to *rw337*: small eyes (left, red arrowheads) and retinal phenotypes: apoptosis and no ONL (right, open red arrowheads). Scale bars: 100 μm (left panel). Scale bars: 20 μm (right panel).

1333

Figure 1-figure supplement 4. Confirmation of specificity of banp MO and nuclear localization of zebrafish Banp protein.

1336 (A) Experimental design to confirm the specificity of banp MO and banp 5misMO (control MO). Co-injection of MO and RNA encoding Banp(wt)-EGFP to one-cell-stage eggs enables us to examine EGFP expression in gastrula stage-embryos at 6 hpf.

1340 (B) banp MO effectively inhibits translation of Banp(wt)-EGFP protein in zebrafish embryos at 6 hpf and the retina at 24 hpf. On the other hand, banp 5misMO did not inhibit the translation of Banp(wt)-EGFP protein, suggesting that banp MO specifically inhibits Banp protein translation. Yellow dotted line indicates blastoderm cells covering yolk, in which GFP fluorescence was detected only in 5mismatch morphant embryos. Scale bars: 20μm.

(C) Time-lapse confocal scanning of mitosis of *Tg[EF1α:mCherry-CAAX]* transgenic wild-type retinal cells expressing Banp(wt)-EGFP. EGFP fluorescence is indicated in green, whereas mCherry-CAAX fluorescence labels plasma membrane and is indicated in magenta. White arrowheads indicate time-lapse change of a retinal progenitor cell undergoing mitosis. Banp(wt)-EGFP is localized in the nucleus in the interphase, whereas it spreads into the whole cell region during mitosis. Scale bars: 4μm.

(D) Histogram of intensity of Banp(wt)-EGFP during cell division shown in (A). Signals decline from mitosis to cytokinesis.

(E) Expression of Banp(wt)-EGFF, EGFP-Banp(wt) and EGFP-Banp(rw337) in zebrafish retina at 24 hpf. Both N-terminal and C-terminal EGFP-tagged wild-type Banp show nuclear localization, indicating that nuclear localization is conserved in zebrafish Banp. However, EGFP-Banp(rw337) is unstable and not maintained, suggesting that EGFP-Banp(rw337) is degraded, probably by the nonsense mutation decay mechanism. Scale bar: 10μm.

Source data 1. Data for Figure 1-figure supplement 4D.

Figure 1-figure supplement 5. Spatio-temporal pattern of zebrafish *banp* mRNA expression.

(A) Whole-mount *in situ* hybridization of zebrafish *banp* mRNA probe. *banp* mRNA is expressed in the 4-cell stage, suggesting maternal expression. *banp* mRNA is ubiquitously expressed during development from the sphere stage until the 21-somite stage. mRNA expression is restricted in the brain including the retina at 30 hpf. At 48 hpf, mRNA is prominently expressed in the brain, fins, and notochord. At 72 hpf and later, mRNA expression appears in neuromasts of the lateral lines (red open arrowheads). br, brain; re, retina; LL, neuromasts of the lateral line; n, notochord. Scale bars: 200 μm

(B) Frontal sections of zebrafish head labeled with *banp* RNA probe at 24 hpf and 48 hpf. *banp* mRNA is expressed in the entire neural retina at 24 hpf and in the CMZ at 48 hpf, suggesting the expression of *banp* mRNA in retinal progenitor cells. OT, optic tectum; CMZ, ciliary marginal zone. Scale bars: 50 μm

1377 (C) Whole-mount *in situ* hybridization of *banp* mRNA expression in wild-type and

1378 *banp^{rw337}* mutant embryos at 72 hpf. *banp* mRNA expression is absent in *banp^{rw337}*

1379 mutants, suggesting non-sense mutation decay of mRNA. Scale bars: 200 μm

1380

Figure 2-figure supplement 1. Apoptosis is induced in retinal progenitor cells and

newly differentiating neurons in *banp^{rw337}* mutants.

(A) BrdU labeling of wild-type and *banp^{rw337}* mutant retinas at 48 hpf.

(B) ath5:EGFP expression of wild-type and *banp^{rw337}* mutant retinas at 48 hpf.

(C) Percentage of BrdU-positive area or ath5:EGFP-positive area relative to the total

retinal area. There is no significant difference in BrdU-positive area between wild-

type and *banp^{rw337}* mutant retinas. The fraction of ath5:EGFP-positive area is

slightly, but significantly reduced in *banp^{rw337}* mutants compared with wild-type

siblings. Unpaired t test (two tailed), Mean ± SD. [n=6, $p^*< 0.05$, ns (not

significant)].

(D) Labeling of wild-type and *banp^{rw337}* mutant retinas with anti-activated caspase 3

antibody at 48 hpf.

(E) Labeling of wild-type and *banp^{rw337}* mutant retinas with anti-pH3 antibody at 48 hpf.

(F) The number of caspase 3-positive or pH3-positive cells per section of retina. The

number of caspase 3-positive cells and pH3-positive cells is significantly higher in

banp^{rw337} mutants than in wild-type siblings. Unpaired t-test (two tailed), Mean ±

SD. [n= 3, $p^*< 0.05$, $p^{**}< 0.01$].

(G) The relative position of caspase 3-positive cells or pH3-positive cells along the

apico-basal axis of the neural retina: Positional value from 0 to 1 corresponds to the

range from apical to basal position, respectively. In wild-type retinas (upper panel),

pH3-positive cells are located in the apical region of the neural retina (peak of

positional value is 0.2), whereas only one caspase 3-positive cell is located in the

intermediate position of the neural retina (peak of positional value is 0.4). On the

other hand, in *banp^{rw337}* mutant retina, pH3-positive cells are located in the apical

region of the neural retina (peak of positional value is 0.1), whereas caspase3-

positive cells are increased in number and most of them are located in the

intermediate zone (peak of positional value is 0.3). Although pH3- and caspase3-

positive positional ranges are slightly overlapped in *banp^{rw337}* mutant retinas, these

data suggest that apoptosis occurs in S phase of retinal progenitor cells or newly

differentiating neurons migrating toward the basal region in *banp*^{rw337} mutants.

(H) Double labeling of wild-type and *banp*^{rw337} mutant retinas with ath5:EGFP

transgene and anti-caspase 3 antibody.

(I) Percentage of ath5:EGFP-positive area relative to the total retinal area (left) and

percentage of ath5:EGFP- and caspase 3-double positive cells relative to the total

number of caspase 3-positive cells (right). Mann Whitney test, Mean ± SD. [n=7,

p**< 0.01].

(J) Double labeling of wild-type and *banp*^{rw337} mutant retinas with anti-HuC/D

antibody and anti-caspase 3 antibody.

(K) Percentage of HuC/D-positive area relative to the total retinal area (left) and

percentage of HuC/D- and caspase 3-double positive cells relative to the total

number of caspase 3-positive cells (right). Mann Whitney test, Mean ± SD. [n=7 for

wild-type, n=6 for *banp*^{rw337} mutants, p*< 0.05].

(L) A possible model on how apoptosis is induced during retinal progenitor cell

proliferation and retinal neurogenesis in *banp*^{rw337} mutants.

Scale bar: 20μm for (A, B, D, E, H, J)

Source data 1. Data for Figure 2-figure supplement 1CFGIK

1427

Figure 4-figure supplement 1. DNA damage is induced in retinal progenitor cells in

***banp* knockdown.**

(A) Labeling of 48 hpf wild-type retinas injected with STD MO and banp MO with anti-

γ-H2AX antibody. γ-H2AX-positive cells are increased in *banp* morphant retinas.

Scale bars: 20μm.

(B) The number of γ-H2AX-positive cells in 48 hpf wild-type retinas injected with STD

MO and banp MO. The number of γ-H2AX-positive cells is significantly higher in

banp morphant retinas than in wild-type retinas injected with STD MO. Mann

Whitney test, Mean ± SD. [n=5, p**< 0.01].

(C) Labeling of 48 hpf wild-type and *banp*^{rw337} mutant retinas with anti-BrdU antibody

(red), anti-γ-H2AX antibody (blue), and ath5:EGFP transgene (green). Two

independent *banp*^{rw337} mutant retinas are shown. Scale bars: 20 μm.

(D) The fraction of γ-H2AX-positive cells in the total retinal area in wild-type and

banp[rw337] mutants. Mann Whitney test, Mean ± SD. [n=6, p <**0.01].

(E) Double labeling of retinal CMZ in *banp*[rw337] mutants with anti-BrdU antibody (red),

anti-γ-H2AX antibody (blue). Higher magnification images of square in *banp*[rw337]

mutant retinas shown in (C).

(F) The fraction of BrdU-positive cells in the total retinal cells and γ-H2AX-positive

cells in *banp*[rw337] mutants. Mean of difference between the total retinal cells and γ-

H2AX-positive cells is shown in the right side. The fraction is significantly higher

in γ-H2AX-positive cells than in the total retinal cells. Paired t-test. [n=6, p

<***0.001]. Correlation plot (F') indicates significantly effective pairing [p< *0.05].

(G) Double labeling of retinal CMZ in *banp*[rw337] mutants with ath5:EGFP transgene

(green), anti-γ-H2AX antibody (blue). Higher magnification images of square in

banp[rw337] mutant retinas shown in (C).

(H) The fraction of ath5:EGFP-positive cells in the total retinal cells and γ-H2AX-

positive cells in *banp*[rw337] mutants. Mean of difference between the total retinal cells

and γ-H2AX-positive cells is shown in the right side. There is no significant

difference between the total retinal cells and γ-H2AX-positive cells. Paired t-test.

[n=6, ns]. Correlation plot (H') indicates no significant pairing. [ns].

(I) Labeling of 48 hpf wild-type and *banp*[rw337] mutant retinas with anti-pH3 antibody

(magenta), anti-γ-H2AX antibody (green). Top and bottom right-side panels of

banp[rw337] mutant retinas indicate original fluorescent images and their Imaris

surface-rendered images, respectively. Scale bars: 20 μm.

(J) The fraction of pH3-positive cells in the total retinal cells and γ-H2AX-positive

cells in *banp*[rw337] mutants. Mean of difference between the total retinal cells and γ-

H2AX-positive cells is shown in the right side. There is no significant difference

between the total retinal cells and γ-H2AX-positive cells. Paired t-test [n=10, ns].

Correlation plot (J') indicates no significant pairing. [ns].

Source data 1. Data for Figure 4-figure supplement 1BDFHJ.

1468

Figure 5-figure supplement 1. Motifs enriched in the closed and open chromatin

area in *banp*[rw337] mutants.

(A) List of top6 motifs enriched in genomic regions of decreased (downregulated) and

increased (upregulated) accessibility in *banp^rw337* mutants. We found that the Banp

motif (TCTCGCGAGA) is the most enriched sequence in regions with reduced

chromatin accessibility in *banp^rw337* mutants (large arrowhead).

(B, C) A putative Banp motif with a conserved CGCG element upstream of the 5' UTR

of *ncapg* (B) and *cenpt* (C) in zebrafish, mice and humans.

1477

Video File Legends

Video 1. Time-lapse movie of mitosis of wild-type retinal progenitor cells.

Tg [h2afv: GFP; EF1α: mCherry-zGem] transgenic embryos were injected with banp

5misMO and mitosis of retinal progenitor cells was scanned with confocal LSM.

Mitosis proceeds sequentially from prophase (t=5~10 min), metaphase (t=16 min),

anaphase (t=21 min) and telophase (t=24 min). Green indicates GFP fluorescence,

representing chromatin. On the other hand, magenta indicates fluorescence from

mCherry, representing S/G2/M phase cells.

1486

Video 2. Time-lapse movie of mitosis of *banp* morphant retinal progenitor cells.

Tg [h2afv: GFP; EF1α: mCherry-zGem] transgenic embryos were injected with banp

MO and mitosis of retinal progenitor cells was scanned with confocal LSM. Mitosis did

not procced smoothly, and it took longer from chromosome condensation in prophase

(t=15 min) to chromosome alignment at the division plane in anaphase (t=46 min).

Aligned chromosomes were often dispersed during chromosome segregation by mitotic

spindle (t=46~64 min). Green indicates fluorescence from GFP, representing chromatin.

On the other hand, magenta indicates fluorescence from mCherry, representing S/G2/M

phase cells.

1496

1497

Supplementary Information

Supplementary Figures

Figure 1-figure supplement 1. Retinal cell differentiation in $banp^{rw337}$ mutants at 4 dpf.

Figure 1-figure supplement 2. Amino acid conservation of Banp protein between

zebrafish, mice and humans.

Figure 1-figure supplement 3. $banp^{sa12976}$ homozygous mutants phenocopy $rw337$

mutants.

Figure 1-figure supplement 4. Confirmation of specificity of banp MO and nuclear

localization of zebrafish Banp protein.

Figure 1-figure supplement 5. Spatio-temporal pattern of zebrafish $banp$ mRNA

expression.

Figure 2-figure supplement 1. Apoptosis is induced in retinal progenitor cells and

newly differentiating neurons in $banp^{rw337}$ mutants.

Figure 4-figure supplement 1. DNA damage is induced in retinal progenitor cells in

$banp$ knockdown.

Figure 5-figure supplement 1. Motifs enriched in the closed and open chromatin area

in $banp^{rw337}$ mutants.

1515

1516 **Supplementary Table file**

1517 **Supplementary Table 1.**

1518

Source data files

Figure 2-Source data 1. Data for Figure 2CD.

Figure 3-Source data 1. Data for Figure 3CEFH.

Figure 3-Source data 2. Data for Figure 3D.

Figure 4-Source data 1. Data for Figure 4ADEF.

Figure 5-Source data 1. Data for Figure 5AC.

Figure 6-Source data 1. Data for Figure 6BD.

Figure 1-figure supplement 4-Source data 1. Data for Figure 1-figure supplement 4D.

Figure 2-figure supplement 1-Source data 1. Data for Figure 2-figure supplement

1CFGIK

Figure 4-figure supplement 1-Source data 1. Data for Figure 4-figure supplement

1BDFHJ.

1531

1532 **Video files**

1533 **Video 1.** Time-lapse movie of mitosis of wild-type retinal progenitor cells.

1534 **Video 2.** Time-lapse movie of mitosis of *banp* morphant retinal progenitor cells.

1535

1536

1537

1538

1539

1540

1541

1542

1543

1544

1545

Appendix

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (Danio Rerio)	*banp^{rw337}*	This paper		
strain, strain background (Danio Rerio)	*banp^{sa12976}*	ZIRC	ZFIN: ZDB-ALT-130411-2647	
strain, strain background (Danio Rerio)	*Tg[EF1α:mCherry-zGem]^{oki011}*	PMID: 25260917	ZFIN: ZDB-ALT-150128-2	
strain, strain background (Danio Rerio)	*Tg[EF1α:mCherry-CAAX]^{oki049}*	PMID: 28196805	ZFIN: ZDB-TGCONSTRCT-181026-1	
strain, strain background (Danio Rerio)	*Tg[h2afv:GFP]$^{kca6/kca66}$*	PMID: 11819118	ZFIN: ZDB-ALT-020918-4, ZDB-ALT-020717-2	
strain, strain background (Danio	*Tg[ath5:EGFP]^{rw021}*	PMID: 15728672	ZFIN: ZDB-ALT-050627-2	
antibody	pH3 (Mouse ,monoclonal)	Cell signaling technology	6G3, 9706	1:500 (IHC)
antibody	pH3 (Rabbit, polyclonal)	Sigma-Aldrich (Merk)	06-570	1:500 (IHC)

antibody	Caspase3 (Rabbit, monoclonal)	BD Pharmingen	Clone C92-605	1:200 (IHC)
antibody	Pax6 (Rabbit, polyclonal)	Covance	PRB-278P	1:500 (IHC)
antibody	zpr1 (Mouse, monoclonal)	ZIRC, Eugene, Oregon	ZFIN: ZDB-ATB-081002-43	1:100 (IHC)
antibody	glutamine synthetase (GS) (Mouse, monoclonal)	Millipore	MAB302, clone GS-6	1:100 (IHC)
antibody	PCNA (Mouse, monoclonal)	Sigma-Aldrich	Clone PC10,P8825	1:200 (IHC)
antibody	Prox1 (Rabbit, polyclonal)	Gene Tex	GTX128354	1:500 (IHC)
antibody	Alpha tubulin (Mouse, monoclonal)	Sigma-Aldrich	T5168, clone B512	1:1000 (IHC)
antibody	γ-H2AX (Rabbit, polyclonal)	Gene Tex	GTX127342	1:500 (IHC)
antibody	BrdU (Rat, monoclonal)	abcam	Ab6326	1:200 (IHC)
antibody	HuC/D (Mouse, monoclonal)	Thermo Fisher	A-21271	1:200 (IHC)

antibody	β-actin (Mouse, monoclonal)	Sigma-Aldrich	A5441	1:5000 (western blot)
antibody	tp53 (Rabbit, polyclonal)	Gene Tex	GTX128135	1:1000 (western blot)
antibody	Secondary antibodies, horseradish peroxidase-conjugated (Mouse)	Amersham	NA931	1:5000 (western blot)
antibody	secondary antibodies, horseradish peroxidase-conjugated (Rabbit)	Amersham	NA934	1:5000 (western blot)
recombinant DNA reagent	pBluescript II SK(+) (vector)	Stratagene/Agilent Technologies		*in vitro* transcription (In situ hybridization probe synthesis)
recombinant DNA reagent	pCS2 (expression vector)	PMID: **7926732**		*in vitro* transcription (Capped mRNA synthesis)
sequence-based reagent (mutation mapping)	Polymorphic marker to link with banp mutation locus	This paper	PCR primers	70702B forward: 5'-ACTTCTTATCAGGGCTGTGC-3' 070702B reverse: 5'-TCAGTCAAGAGCAGTGAGAG-3'
sequence-based reagent (mutation mapping)	Polymorphic markers to link banp mutation locus	This paper	PCR primers	zC93F2D forward: 5'-TGGGATCTCTTTAAGTGAGTGAG-3' zC93F2D reverse:

				5'-TCCAACTATGTGGG TCAAACC-3'
sequence-based reagent (Morpholino)	banp MO	This paper	Morpholino antisense oligos	5'-CCACTAAATCTT GCTCTGACATCA T-3'
sequence-based reagent (Morpholino)	banp 5misMO	This paper	Morpholino antisense oligos	5'-CCtCaAAATgTTc CTCTcACATCAT-3'
sequence-based reagent (Morpholino)	tp53 MO	PMID: 12477391	Morpholino antisense oligos	5'-GCGCCATTGCTT TGCAAGAATTG-3'
sequence-based reagent (Morpholino)	Δ113 tp53 MO	PMID: 19204115	Morpholino antisense oligos	5'-GCAAGTTTTTGC CAGCTGACAGAA G-3'
sequence-based reagent (Morpholino)	STD MO	PMID: 30322969	Morpholino antisense oligos	5'-CCTCTTACCTCAG TTACAATTTATA-3'
sequence-based reagent (primer)	ccng1	This paper	Primers for qRT-PCR	**Forward primer**: ccctggagattgaggatc ag **Reverse primer:** cacacaaaccaggtctc caa
sequence-based reagent (primer)	mdm2	PMID: 24147052	Primers for qRT-PCR	**Forward primer**: caggaggaggagaagca gtg **Reverse primer:** agggaaaagctgtcc gactt

sequence-based reagent (primer)	p21(cdkn1a)	PMID: 26908596	Primers for qRT-PCR	**Forward primer**: aagcgcaaacagaccaacat **Reverse primer**: tcagctactggccggattt
sequence-based reagent (primer)	FL tp53	PMID: 27539857	Primers for qRT-PCR	**Forward primer**: tggagaggaggtcggcaaaatcaa **Reverse primer**: gactgcgggaacctgagcctaaat
sequence-based reagent (primer)	Δ113 tp53	PMID: 27539857	Primers for qRT-PCR	**Forward primer**: atatcctggcgaacatttggaggg **Reverse primer**: cctcctggtcttgtaatgtcac
sequence-based reagent (primer)	Puma (bbc3)	This paper	Primers for qRT-PCR	**Forward primer**: ctgaggaggacccacact **Reverse primer**: tctccagttctgccagtgc
sequence-based reagent (primer)	cenpt	This paper	Primers for qRT-PCR	**Forward primer**: tcatgaggagattgtggaagatg **Reverse primer**: ggtgagctctgcgagttatt
sequence-based reagent (primer)	ncapg	This paper	Primers for qRT-PCR	**Forward primer**: ctgatgtgagggagcctattt **Reverse primer**: gagtctgtttggcctccatta

sequence-based reagent (primer)	atm	This paper	Primers for qRT-PCR	**Forward primer:** cctcaaggctgtggagaact **Reverse primer:** aggggattttctttacaccactc
sequence-based reagent (primer)	atr	This paper	Primers for qRT-PCR	**Forward primer:** aggaacccaatctgccagt **Reverse primer:** gatgtccagtgccagctctc
sequence-based reagent (primer)	ef1α	PMID: 19014500	Primers for qRT-PCR	**Forward primer:** cttctcaggctgactgtgc **Reverse primer:** ccgctagcattaccctcc
sequence-based reagent (sequencing primer)	Genotyping primers for *banp*[sa12976]	This paper	Primers	**Forward primer:** TGTTGATATCCATCAGTCAG **Reverse primer:** GGTGTATAAATCACATGACC
sequence-based reagent (primer)	Genotyping primers for *banp*[rw337]	This paper	Primers	**forward:** 5'-CGATGTTGATATCCATCAGTCAGGCGATC-3'; **reverse primer:** 5'-GGTGCTGGTGTATAAATCACATGACCTATGGTCCTCTT-3'.
commercial assay or kit	DIG RNA Labelling Kit	Roche	11175025 910	In situ hybridization

commercial assay or kit	mMESSAGE mMACHINE™ SP6 Transcription Kit	Invitrogen	AM1340	Capped mRNA synthesis
commercial assay or kit	In Situ Cell Death Detection Kit, TMR red	Roche	12156792910	apoptosis
commercial assay or kit	ReverTra Ace® aPCR master mix with gDNA remover	Toyobo	FSQ-301	cDNA synthesis
commercial assay or kit	Luna® Universal qPCR Master Mix	NEB	M3003L	qRT-PCR
commercial assay or kit	Direct-zol RNA Miniprep Kit	ZYMO RESEARCH	R2050	RNA-sequencing
commercial assay or kit	NEBNext® Ultra™ II Directional RNA Library Prep Kit	NEB	E7760	Illumina RNA-sequencing
chemical compound, drug	Acridine orange	WALDECK (CHROMA)	1B-307	Live cell death detection
software	IMARIS	Bitplane	ver.9.1.2	http://www.bitplane.com/imaris; RRID: SCR_007370
software	Image J	NIH	Version 2.1.0/1.53c	Percentage area calculation
software	Zen	Zeiss	LSM 710	Percentage area calculation
software	GraphPad Prism	GraphPad Software	Version 9.1.2	https://www.graphpad.com/scientific-software/prism/
other	SYTOX Green	Molecular Probes	S34862	Nuclear stain 1:1000 (IHC)
other	rhodamine-conjugated	Molecular Probes	R415	Filamentous actin (F-actin) stain

	phalloidin			1: 40 (IHC)
other	Restore™ PLUS Western Blot Stripping Buffer	Thermo Scientific	46430	Remove high-affinity antibodies from membranes (western blot)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7