1

Banp regulates DNA damage response and chromosome segregation during the cell cycle in zebrafish retina

Swathy Babu, Yuki Takeuchi, and Ichiro Masai[#]

Developmental Neurobiology Unit, Okinawa Institute of Science and Technology
Graduate University (OIST)
1919-1 Tancha, Onna, Okinawa, Japan

Correspondence should be addressed: masai@oist.jp

Keywords: Banp, DNA damage, chromosome segregation, cell cycle, cenpt, ncapg,
tp53

Word Count for Introduction/Results/Discussion: 7,162

Abstract

Btg3-associated nuclear protein (Banp) was originally identified as a nuclear matrix-associated region (MAR)-binding protein and it functions as a tumor suppressor. At the molecular level, Banp regulates transcription of metabolic genes via a CGCG-containing motif called the Banp motif. However, its physiological roles in embryonic development are unknown. Here we report that Banp is indispensable for the DNA damage response and chromosome segregation during mitosis. Zebrafish *banp* mutants show mitotic cell accumulation and apoptosis in developing retina. We found that DNA replication stress and tp53-dependent DNA damage responses were activated to induce apoptosis in *banp* mutants, suggesting that Banp is required for regulation of DNA replication and DNA damage repair. Furthermore, consistent with mitotic cell accumulation, chromosome segregation was not smoothly processed from prometaphase to anaphase in *banp* morphants, leading to a prolonged M-phase. Our RNA- and ATAC-sequencing identified 31 candidates for direct Banp target genes that carry the Banp motif. Interestingly, a DNA replication fork regulator, *wrnip1,* and two chromosome segregation regulators, *cenpt* and *ncapg*, are included in this list. Thus, Banp directly regulates transcription of *wrnip1* for recovery from DNA replication stress, and *cenpt* and *ncapg* for chromosome segregation during mitosis. Our findings provide the first *in vivo* evidence that Banp is required for cell-cycle progression and cell survival by regulating DNA damage responses and chromosome segregation during mitosis.

(221 words)

Introduction

BTG3-associated nuclear protein (BANP) (also known as Scaffold/matrix attachment region binding protein 1 (SMAR1)) was identified as a nuclear protein that binds to the matrix-associated region (MAR) of the β-locus of murine T cell receptors (Birot et al., 2000; Chattopadhyay et al., 2000). MARs are often found near gene promoters (Scheuermann and Garrard, 1999) and constitute a periodic loop domain, through which chromosomal DNA is topologically organized inside the nucleus (Blasquez et al., 1989; Hart and Laemmli, 1998). Furthermore, BANP functions as a tumor suppressor (Kaul et al., 2003; Kaul-Ghanekar et al., 2005). In humans, the *BANP* gene is located on chromosome 16q24, the genomic region of which often suffers from loss-of-heterozygosity in cervical and breast cancers (Tsuda et al., 1994; Wang et al., 1999). BANP interacts with tumor suppressor tp53 to modulate its functions (Kaul et al., 2003; Pavithra et al., 2009; Sinha et al., 2012; Sinha et al., 2010). Overexpression of BANP suppresses cancer growth (Bhagat et al., 2018; Liu et al., 2014; Taye et al., 2018). However, these proposed cellular functions of BANP were based mostly on *in vitro* studies, and genetic deletion of *Banp* was reportedly unsuccessful (Grand et al., 2021), whereas *Banp* knockout mice show embryonic lethality (Chemmannur et al., 2015), which makes *in vivo* analysis difficult. Thus, physiological functions of BANP, especially in cell-cycle regulation and tumor suppression, are still unknown. Very recently, it was reported that BANP binds to the nucleotide sequence TCTCGCGAGA, called the "Banp motif", which is enriched near the transcription initiation site of CpG island promoters, a genomic region in which it is difficult to replicate and repair DNA, and which regulates gene transcription in a DNA methylation-dependent manner. BANP promotes transcription of essential metabolic genes in pluripotent stem cells and terminal differentiated neuronal cells in humans and mice (Grand et al., 2021), indicating that BANP functions as a transcription factor at the molecular level. These findings raise the question, "How is the BANP-mediated transcriptional network involved in cell-cycle regulation and tumor suppression?"

The vertebrate retina is composed of six major classes of neurons and one type of glia [retinal ganglion cells (RGCs), three types of interneurons (amacrine cells, bipolar cells, and horizontal cells), two types of photoreceptors (rods and cones), and Müller glia]. These retinal cells are assembled to form the neural circuit responsible for

phototransduction and visual processing (Dowling, 2012). During development, retinal progenitor cells are multipotent and give rise to all seven retinal cell types in response to extrinsic and intrinsic cues (Boije et al., 2014). An important characteristic of retinal development is that retinal neurogenesis is linked to cell-cycle regulation. In zebrafish, cell-cycle length of retinal progenitor cells is around 32–49 h from 6 to 25 hpf, but shortened to 8–10 h once retinal neurogenesis starts at 25 hpf (Li et al., 2000), suggesting that activation of cell-cycle progression is associated with neurogenesis. It is generally accepted that active cell proliferation may increase DNA replication errors, which induces DNA replication stress and subsequent DNA damage (Berti et al., 2020; Blackford and Jackson, 2017a; Waterman et al., 2020). Thus, factors involved in DNA replication checkpoint and DNA damage response are essential to ensure genomic integrity (Murga et al., 2009). Otherwise, retinal progenitor cells accumulate genetic defects leading to apoptosis. Consistently, in zebrafish, retinal apoptosis occurs in the absence of DNA damage response regulators such as DNA damage-binding protein 1 (DDB1) (Yildirim et al., 2015), Fanconi anemia group D2 (Fancd2) (Liu et al., 2003), and Ku80 (Xrcc80) (Bladen et al., 2005). We also showed that retinal apoptosis occurs in zebrafish *prim1* mutants, in which DNA replication stress is chronically elevated. In this mutant, a central regulator of the DNA replication checkpoint, Ataxia telangiectasia-mutated and Rad3-related (ATR), is activated, which subsequently activates DNA damage response regulators, Ataxia telangiectasia-mutated (ATM) and Chk2, leading to tp53-mediated apoptosis. Thus, DNA replication checkpoint and DNA damage response function as safeguards to ensure integrity of retinal cell proliferation and neurogenesis (Yamaguchi et al., 2008). Therefore, developing retina is a highly proliferating tissue, in which a spatiotemporal pattern of neurogenesis is tightly coordinated by cell-cycle regulation. So, vertebrate retina provides a great model for studying how cell-cycle regulation, including DNA damage response, ensures neurogenesis and subsequent cell differentiation.

In this study, we identified zebrafish *banp* mutants, which show mitotic cell accumulation and apoptosis in the developing retina. DNA replication checkpoint and tp53-dependent DNA damage response are activated to induce apoptosis in *banp* mutant retinas. Thus, Banp is required for regulation of DNA replication and DNA damage repair. Furthermore, consistent with mitotic cell accumulation in *banp* mutants, our live

imaging revealed that chromosome segregation fails to proceed smoothly during mitosis in *banp* morphants, resulting in a prolonged M-phase. Our RNA-sequencing and ATAC-sequencing enabled us to identify 31 candidates for direct Banp target genes that carry the Banp motif near their promoters. Interestingly, both zebrafish and human genes encoding two chromosome segregation regulators, *Cenpt* and *Ncapg*, and a DNA replication fork regulator, *Werner helicase interacting protein 1* (*Wrnip1*) are included in this list. Thus, Banp regulates transcription of *cenpt* and *ncapg* to promote chromosome segregation during mitosis, whereas Banp maintains transcription of *wrnip1* for recovery from DNA replication stress during S phase. Taken together, these data suggest that Banp is required for cell-cycle progression and cell survival in zebrafish retinas by regulating DNA damage response and chromosome segregation during mitosis.

Results

Zebrafish *rw337* mutants show mitotic cell accumulation and cell death in the retina after 2.5 dpf

We performed large-scale mutagenesis using zebrafish and identified a zebrafish mutant, *rw337*. *rw337* mutants have smaller eyes and a smaller optic tectum at 4 days post-fertilization (dpf) (Fig. 1A). Around 6–7 dpf, *rw337* mutants die, indicating that this mutation is lethal. Next, we examined retinal phenotypes (Fig. 1B). At 1.5 dpf, the retina consists of retinal progenitor cells and early differentiating neurons. There was no difference in retinal morphology between *rw337* mutants and wild-type siblings. However, at 2 dpf, mitotic-like round cells had accumulated in the apical region of the retina in *rw337* mutants, suggesting mitotic arrest or prolonged duration of mitosis. At 2.5 dpf, in wild-type retinas, three nuclear layers, the RGC layer, the inner nuclear layer (INL) and the outer nuclear layer (ONL), and two plexiform layers, the inner plexiform layer (IPL) and the outer plexiform layer (OPL), had formed. In *rw337* mutants, only the RGC layer was observed; however, many pyknotic nuclei were observed in the outer retinal region. At 3 dpf, the RGC layer, IPL and INL appeared in *rw337* mutants. However, pyknotic nuclei were densely packed in the outer region of *rw337* mutant retinas. At 4 dpf, most dead cell debris had decreased in *rw337* mutants. The RGC layer, IPL, and INL were formed; however, no ONL was observed, despite very small, patchy photoreceptor-like cells. At 3 and 4 dpf, the ciliary marginal zone (CMZ), where retinal stem and progenitor cells are located, was observed in *rw337* mutants. Thus, RGCs differentiate, but cell death occurs after 2.5 dpf in *rw337* mutants.

Next, we examined neuronal differentiation in *rw337* mutants at 4 dpf. The transgene *Tg[ath5:EGFP]* drives EGFP expression strongly in RGCs and weakly in amacrine cells and photoreceptors (Fig. 1-figure supplement 1A) (Masai et al., 2000; Masai et al., 2005). Pax6 is strongly expressed in amacrine cells and weakly in RGCs (Fig. 1-figure supplement 1C) (Macdonald and Wilson, 1997). Prox1 is expressed in bipolar cells and horizontal cells (Fig. 1-figure supplement 1E) (Jusuf and Harris, 2009). zpr1 antibody labels double cone photoreceptors (Fig. 1-figure supplement 1G) (Larison and Bremiller, 1990). Anti-glutamine synthetase (GS) antibody labels Müller cells (Fig. 1-figure supplement 1I) (Peterson et al., 2001). We observed that all these markers are expressed in *rw337* mutants (Fig. 1-figure supplement 1B, 1D, 1F, 1H, 1J). However,

the numbers of strong Pax6+ cells and GS+ cells were decreased in *rw337* mutants (Fig.

1-figure supplement 1D, 1J), suggesting that amacrine cells and Müller cells were

reduced. Prox1+ cells were misplaced and less numerous in *rw337* mutants, suggesting

that horizontal and bipolar cells are also reduced (Fig. 1-figure supplement 1F). zpr1+

cells were not detected, except for a small patch of zpr1+ cells in the ventral retina,

where photoreceptor differentiation initially occurs in *rw337* mutants (Fig. 1-figure

supplement 1H), suggesting that most cone photoreceptors fail to differentiate. Anti-

PCNA antibody labels retinal stem and progenitor cells in the CMZ (Fig. 1-figure

supplement 1K), which was not affected in *rw337* mutants (Fig. 1-figure supplement

1L). Phalloidin labeling shows IPL and OPL in wild-type retinas (Fig. 1-figure

supplement 1M). *rw337* mutants showed IPL, but not OPL (Fig. 1-figure supplement

1N), which is consistent with zpr1 labeling (Fig. 1-figure supplement 1H). Thus, retinal

stem and progenitor cells are maintained. RGCs do differentiate; however, amacrine

cells, horizontal cells, bipolar cells, and Müller cells are decreased in number and

photoreceptors fail to differentiate in *rw337* mutants.

***rw337* mutant gene encodes Banp**

We mapped the *rw337* mutation to zebrafish chromosome 25 (Fig. 1C). Two

polymorphic markers, 070702B and zC93F2D, restricted the genomic region

encompassing *rw337* mutation, in which only four candidate genes: *BTG3-*

associated nuclear protein (*banp*); *programmed cell death 2-like* (*pdcd2l*);

carboxylesterase 2b (*ces2b*); *protein disulfide isomerase family A, member 3* (*pdia3*),

were annotated. We cloned all these cDNAs from mRNA prepared from *rw337*

homozygous mutants, sequenced them, and found that part of the intron sequence

between exons 4 and 5 (intron 4/5) failed to be spliced out of the *banp* cDNA (Fig. 1D).

A nucleotide T at the exon-intron junction of exon 4 is converted to G, which alters the

donor splice site of exon 4 from GT to GG. This mutation leads to abnormal insertion of

a 52-bp intron sequence after exon 4, resulting in a premature stop codon in *banp* cDNA

(Fig. 1D). The *banp* gene encodes a protein containing a coiled-coil domain, a BEN

domain, which was identified as a novel domain found in chromatin associated factors

and DNA viral proteins and proposed to mediate protein-DNA and protein-protein

interactions during chromatin organization and transcription (Abhiman et al., 2008), and

a DNA-binding domain, which includes a MAR-binding domain (Fig. 1E). Overall,

zebrafish Banp protein shows high similarity to those of humans and mice, especially in

the BEN domain (Fig. 1-figure supplement 2). This mutation is predicted to generate

nonfunctional, truncated Banp protein in *rw337* mutants (Fig. 1E). There was no

mutation in the other three genes in *rw337* mutants, suggesting that the *rw337* mutant

gene encodes Banp.

To confirm that the loss of Banp causes retinal phenotypes in *rw337* mutants, we

carried out three sets of experiments. First, we examined another nonsense mutation

allele, *banp*^{sa12976}, which has a premature stop codon in exon 4 (Fig. 1-figure

supplement 3A). Homozygous *banp*^{sa12976} mutant embryos and trans-heterozygous

banp^{rw337}; *banp*^{sa12976} mutants phenocopied *rw337* mutants (Fig. 1-figure supplement

3B). Second, we inhibited Banp using ATG-morpholino antisense oligos against Banp

(banp MO). We confirmed MO specificity, because banp MO inhibited translation of

mRNA encoding C-terminal EGFP-tagged wild-type Banp (banp(wt)-EGFP), whereas

the control, 5-mismatch banp MO (banp 5misMO) did not (Fig. 1-figure supplement 4A

and 4B). *banp* morphants phenocopied *rw337* mutants (Fig. 1F). Third, we

overexpressed wild-type Banp in *banp*^{rw337} mutants. We injected mRNA encoding

Banp(wt)-EGFP into wild-type embryos at the one-cell stage. In developing retinas,

Banp(wt)-EGFP protein was localized in the nuclei during interphase, whereas it spread

into the whole cell region during mitosis (Fig. 1-figure supplement 4C and 4D),

confirming nuclear localization of zebrafish Banp. We expressed Banp(wt)-EGFP, N-

terminal EGFP tagged wild-type (EGFP-Banp(wt)) and the *rw337* mutant form of Banp

(EGFP-Banp(rw337)) in wild-type retinas. Only EGFP-Banp(rw337) was not stably

maintained (Fig. 1-figure supplement 4E). Finally, overexpression of Banp(wt)-EGFP

rescued retinal phenotypes in *rw337* mutants at 2 dpf, but overexpression of EGFP-

Banp(rw337) did not (Fig. 1G and Fig. 1-figure supplement 5). Taken together, retinal

phenotypes of *rw337* mutants are caused by the loss of Banp.

banp mRNA is prominently expressed in the developing CNS of zebrafish

We examined *banp* mRNA expression by *in-situ* hybridization (Fig. 1-figure

supplement 6A). *banp* mRNA shows ubiquitous expression from the one-cell stage to

the 21-somite stage, suggesting maternal and zygotic expression. mRNA was restricted

to the brain, including the retina at 30 and 48 hpf. Frontal sections show that mRNA is

expressed in neural retina at 24 hpf, but restricted in the CMZ at 48 hpf, suggesting that

retinal progenitor cells express *banp* mRNA (Fig. 1-figure supplement 6B). In addition

to brain and retina, *banp* mRNA expression was observed in fin and notochord at 48 hpf

and in neuromasts of the lateral line after 72 hpf. *banp* mRNA expression was not

detected in *banp^{rw337}* homozygous mutants at 3 dpf (Fig. 1-figure supplement 6C),

suggesting mRNA decay. Thus, *banp* mRNA is prominently expressed in developing

brain in zebrafish.

***banp* mutants show mitotic defects and cell death of retinal progenitor cells**

Although retinal progenitor cells produce RGCs in *banp^{rw337}* mutant retinas, the number

of other neurons, especially photoreceptors, was decreased at 4 dpf (Fig. 1-figure

supplement 1). Indeed, mitotic-like round cells and pyknotic nuclei were accumulated in

the outer region of *banp^{rw337}*mutant retinas from 2 to 3 dpf (Fig. 1B), suggesting that

mitotic defects and cell death reduce the number of retinal neurons. Thus, we conducted

double labeling of wild-type and *banp^{rw337}*mutant retinas with TdT-mediated dUTP

Nick End Labeling (TUNEL), which labels blunt ends of DNA double-strand breaks

(DSBs) to visualize apoptosis, and with anti-phosphorylated (Ser-10) Histone H3 (pH3)

antibody, which labels mitotic cells in prometaphase, metaphase and early anaphase

(Prigent and Dimitrov, 2003) (Fig. 2A). Then, to evaluate TUNEL and pH3 signals

more precisely, we used only fluorescent channel for TUNEL (Fig. 2B) or anti-pH3

antibody labeling (Fig. 2B'), with nuclear staining with Sytoxgreen. TUNEL+ cells

were rarely observed in wild-type retinas throughout stages from 29 to 101 hpf, as well

as in *banp^{rw337}*mutant retinas at 24 hpf (Fig. 2B). However, in*banp^{rw337}*mutants,

TUNEL+ cells appeared in the neural retina at 53 hpf, and became more localized to the

outer retina at 77hpf (Fig. 2B) and to the interface between the central retina and the

CMZ at 77 and 101 hpf (Fig. 2B). After 53 hpf, the number of TUNEL+ cells was

significantly higher in *banp^{rw337}* mutants than in wild-type siblings (Fig. 2C). Thus,

retinal cells undergo apoptosis in *banp^{rw337}*mutants. Next, we examined mitosis by

labeling with anti-pH3 antibody (Fig. 2B'). The number of pH3+ cells did not differ

between wild-type and *banp^{rw337}*mutant retinas at 24 hpf. However, pH3+ cells were

markedly increased and located at the apical surface of the neural retina in

*banp^{rw337}*mutant retinas at 53 hpf (Fig. 2B'), confirming mitotic cell accumulation. At

77 and 101 hpf, pH3+ cells disappeared from the central retina, but still were

accumulated in the CMZ in *banp^{rw337}* mutants (Fig. 2B'). The number of pH3+ cells in

banp^{rw337} mutant retinas increased transiently at 53 hpf, and then decreased, but was

still significantly higher than that of wild-type sibling retinas after 77 hpf because of

accumulation of pH3+ cells at the CMZ (Fig. 2D).

To clarify how mitotic cell accumulation and apoptosis are induced during

retinal cell proliferation and neurogenesis in *banp^{rw337}* mutants, we examined BrdU

incorporation and ath5:EGFP expression at 48 hpf (Fig. 2-figure supplement 1A and

1B). The percentage of BrdU+ area to total retinal area did not differ significantly

between wild-type siblings and *banp^{rw337}* mutants (Fig. 2-figure supplement 1C),

whereas the percentage of ath5:EGFP+ area to the total retinal area was slightly, but

significantly lower in *banp^{rw337}* mutants than in wild-type siblings (Fig. 2-figure

supplement 1C), suggesting that retinal neurogenesis is slightly reduced in *banp^{rw337}*

mutants at 48 hpf, probably due to mitotic defects or apoptosis.

Next, we examined the spatial position of mitotic cells and apoptosis at 48 hpf

by labeling with anti-pH3 and anti-activated caspase 3 antibodies (Fig. 2-figure

supplement 1D and 1E). Both caspase 3+ and pH3+ cells were increased in *banp^{rw337}*

mutants (Fig. 2-figure supplement 1F). Furthermore, caspase 3+ cells were mostly

located in the intermediate zone of the neural retina along the apico-basal axis, whereas

pH3+ cells were localized at the apical surface of the neural retina (Fig. 2-figure

supplement 1G), suggesting that apoptosis occurs mostly in retinal progenitor cells

undergoing G1, S, or G2 phase, or differentiating neurons. Next, we examined whether

apoptosis occurs in early differentiating retinal neurons, using double labeling with anti-

activated caspase 3 antibody and ath5:EGFP (Fig. 2-figure supplement 1H). The

percentage of ath5:EGFP- and caspase 3-double-positive cells to the total number of

caspase 3+ cells was 9.44±13.1% in *banp* mutants at 48 hpf and higher than that of

wild-type siblings (0.00±0.00%), although the difference was not significant (Fig. 2-

figure supplement 1I). Next, we examined whether apoptosis occurs in mature retinal

neurons. HuC/D is expressed in maturing neurons (Park et al., 2000). At 48 hpf, anti-

HuC/D antibody labeled RGCs in zebrafish (Fig. 2-figure supplement 1J). The fraction

of caspase 3- and HuC/D-double-positive cells among total caspase 3+ cells was almost

zero in both wild-type and *banp^rw337* mutants (Fig. 2-figure supplement 1K), suggesting

that apoptosis does not occur in maturing retinal neurons. Taken together, in *banp^rw337*

mutants, retinal progenitor cells primarily show mitotic defects at 48 hpf. On the other

hand, apoptosis is likely induced in retinal progenitor cells at interphase or in newly

differentiating retinal neurons, although apoptosis in newly differentiating retinal

neurons is minor (Fig. 2-figure supplement 1L). Since mitotic cell accumulation was

transiently observed in the central retina at 53 hpf (Fig. 2B'), and apoptosis was not

overlapped with this mitotic cell accumulation area (Fig. 2-figure supplement G), it is

less likely that mitotic cells undergo permanent mitotic arrest leading to apoptosis.

Rather, mitotic cell accumulation seems to be caused by prolonged mitosis. Thus,

banp^rw337 mutants show mitotic cell accumulation and apoptosis in retinal progenitor

cells during interphase and immature neurons (Fig. 2E and 2F).

tp53 pathway mediates retinal apoptosis in *banp* mutants

To understand the molecular mechanism of mitotic defects and apoptosis in *banp^rw337*

mutants, we identified differentially expressed mRNAs using RNA-seq analysis. We

prepared mRNA from 48-hpf wild-type siblings and *banp^rw337* mutants. Illumina

sequencing generated more than 50 million paired-end reads from cDNA libraries.

27,771 protein-coding genes were detected. Among them, 258 genes were upregulated,

and 81 genes were downregulated in *banp^rw337* mutants (Fig. 3A). The top 15

upregulated genes contained a significant fraction of tp53 dependent-DNA damage

response regulators (Fig. 3B). Quantitative RT-PCR revealed that four tp53 target

genes: *p21(cdkn1a), puma (bbc3), ccng1*, and *mdm2* are significantly upregulated in

banp^rw337 mutants (Fig. 3C). Western blotting showed that full length tp53 (FL tp53)

protein is stabilized and more than 11-fold higher in *banp^rw337* mutants than in wild type

(Fig. 3D and 3E). We previously showed that the ATR/ATM-Chk2-tp53 DNA damage

pathway induces retinal apoptosis in zebrafish (Yamaguchi et al., 2008). Indeed, the top

15 upregulated genes contain exo5, a critical regulator for ATR-dependent replication

fork restart (Hambarde et al., 2021) (Fig. 3B). mRNA expression of ATR and ATM was

upregulated in *banp^rw337* mutants (Fig. 3F). So, we examined whether retinal apoptosis

depends on tp53 in *banp^rw337* mutants. Injection of tp53 antisense morpholino (tp53

MO) completely inhibited retinal apoptosis in *banp^rw337* mutants at 48 hpf (Fig. 3G and

3H).   However, retinal apoptosis reappeared in those tp53 MO-injected *banp^rw337*

mutants at 72 hpf (Fig. 3G), although there was no statistical difference in apoptotic cell

number between wild-type and *banp^rw337* mutant retinas at 72 hpf (Fig. 3H). Thus,

retinal apoptosis in *banp^rw337* mutants depends on tp53, although it is possible that some

tp53-independent mechanism induces retinal apoptosis after 48 hpf.

Transcription of Δ113-tp53 is preferentially elevated in *banp* mutants

RNA-seq analysis showed upregulation of *tp53* mRNA (Fig. 3B); however, surprisingly,

quantitative RT-PCR showed that the mRNA level of FL tp53 is not significantly

different between *banp^rw337* mutants and wild-type siblings at 48 hpf (Fig. 4A). There

are several isoforms of tp53 protein (Joruiz et al., 2020; Joruiz and Bourdon, 2016).

Importantly, in zebrafish, the isoform Δ113tp53 is generated from the alternative

transcription start site in intron 4. It was reported that FL tp53 initially activates

transcription of Δ113tp53 after tp53 protein is stabilized in response to cellular stress

(Chen et al., 2009). Δ113tp53 preferentially activates transcription of target genes

related to cell-cycle arrest to promote DNA repair and cell survival, whereas FL tp53

preferentially activates transcription of target apoptotic genes to eliminate severely

DNA-damaged cells (Gong et al., 2015; Gong et al., 2020). Differential expression of

tp53 isoforms is thought to regulate the balance between cell survival and apoptosis in

response to DNA damage. Indeed, quantitative RT-PCR confirmed that only Δ113tp53

was upregulated in *banp^rw337* mutants (Fig. 4A). ATAC-seq revealed that a genomic

region in intron 4 of tp53 is specifically open in *banp^rw337* mutants, but not in wild-type

siblings (Fig. 4B, ATAC-seq). Consistently, transcription of the Δ113tp53 mRNA was

specifically activated in *banp^rw337* mutants, but not in wild-type siblings (Fig. 4B, RNA-

seq). Thus, the internal promoter at intron 4 (Chen et al., 2009; Chen et al., 2005) is

specifically activated in *banp^rw337* mutants (Fig. 4B). These data raise the possibility that

the lack of Banp increases DNA replication stress and promotes an accumulation of

DNA damage, resulting in activation of ATR and ATM, which stabilize FL tp53, such

that FL tp53 triggers Δ113tp53 transcription in *banp^rw337* mutants (Fig. 4G).

DNA damage accumulates during replication in *banp* mutant retinas

To examine whether replicative DNA damage is increased due to loss of Banp functions,

we labeled 48-hpf retinas with anti-γ-H2AX antibody. In response to DSBs, ATM

rapidly phosphorylates histone H2A isoform H2AX at a C-terminal serine-139, which is

called γ-H2AX (Blackford and Jackson, 2017b; Rogakou et al., 1998a; Rogakou et al.,

1998b; Shroff et al., 2004). Furthermore, ATR phosphorylates H2AX at stalled

replication forks, and later, when single-ended DSBs form at persistent stalled forks,

ATM and DNA-PKs act jointly to further propagate γ-H2AX (Sirbu et al., 2011; Ward

and Chen, 2001). Thus, γ-H2AX is a global marker of stalled replication forks and

DSBs. The number of γ-H2AX+ cells increased in both $banp^{rw337}$ mutant (Fig. 4C, 4F)

and *banp* morphant retinas (Fig. 4-figure supplement 1A, 1B), confirming an

accumulation of DNA damage in the absence of Banp. Next, to determine how DNA

damage is induced in $banp^{rw337}$ mutants, we examined the fraction of S-phase cells,

mitotic cells, and post-mitotic neurons in γ-H2AX+ retinal cells in $banp^{rw337}$ mutants at

48 hpf. First, wild-type and $banp^{rw337}$ mutant retinas carrying the transgene

Tg[ath5:EGFP] were labeled with anti-BrdU and anti-γ-H2AX antibodies (Fig. 4-figure

supplement 1C). The fraction of γ-H2AX+ cells increased in $banp^{rw337}$ mutant retinas,

whereas the fraction was almost zero in wild-type sibling retinas (Fig. 4-figure

supplement 1D). Next, we compared the fraction of BrdU+ cells between the total

retinal cells and γ-H2AX+ retinal cells in $banp^{rw337}$ mutants (Fig. 4-figure supplement

1E). The BrdU+ fraction was significantly higher in γ-H2AX+ cells than in total retinal

cells (Fig. 4-figure supplement 1F), indicating that S-phase fraction is 4.3-fold higher in

DNA damaged retinal cells than in total retinal cells, in $banp^{rw337}$ mutants. Thus,

replicative DNA damage is abnormally increased in retinal progenitor cells undergoing

S phase. Next, we examined the fraction of ath5:EGFP+ cells (Fig. 4-figure supplement

1G). There was no significant difference in the ath5:EGFP+ cell fraction between the

total retinal cells and γ-H2AX+ retinal cells in $banp^{rw337}$ mutants (Fig. 4-figure

supplement 1H). Finally, we determined the fraction of mitotic cells by labeling with

anti-pH3 and anti-γ-H2AX antibodies in $banp^{rw337}$ mutant retinas at 48 hpf (Fig. 4-

figure supplement 1I). There is no significant difference in the pH3+ cell fraction

between the total retinal cells and γ-H2AX+ retinal cells (Fig. 4-figure supplement 1J).

Despite of no statistical significance, interestingly, averaged fraction of pH3+ cells in γ-

H2AX+ retinal cells was lower than that of total retinal cells. Since $banp^{rw337}$ mutants

show increase in γ-H2AX + cells, pH3 + cells and apoptotic cells at 48hpf, it is likely

that DNA damaged cells are eliminated by apoptosis in S phase prior to mitosis. This

also suggests that M phase defects is independent of replicative DNA damage

introduced in S phase. Although DNA damage may occur in post-mitotic neurons and

mitotic cells in $banp^{rw337}$ mutants, it is likely that most DNA damage is introduced in S-

phase of retinal progenitor cells due to replication fork stalling in the absence of Banp.

It is generally accepted that tp53 promotes DNA damage repair or eliminates

DNA-damaged cells via apoptosis, so we expected that tp53 knockdown might enhance

accumulation of DNA damage in $banp^{rw337}$ mutants. However, compared with standard-

MO injection control (n=120.8±60.6/section), the injection of FL tp53-MO significantly

reduced the number of γ-H2AX+ cells in $banp^{rw337}$ mutant retinas

(n=55.9±15.9/section) at 48 hpf (Fig. 4C, 4F). Injection of Δ113tp53-MO mildly

reduced the number of γ-H2AX+ cells (n=85.4±15.2/section); however, the difference

was not significant (Fig. 4C, 4F). It is likely that neither tp53 nor Δ113tp53 repairs

DNA damage in $banp^{rw337}$ mutants. Thus, under normal conditions, Banp probably

suppresses DNA replication stress upstream of tp53 by promoting recovery from stalled

forks or repair of DSBs (Fig. 4G).

We already showed that, compared with Standard-MO injection controls

(n=174.7±30.5/section), retinal apoptosis was completely inhibited by FL tp53-MO in

$banp^{rw337}$ mutant retinas (n=0.50±0.58/section) at 48 hpf (Fig. 3G, 3H; also shown in

Fig. 4C, 4D). Next, we investigated whether retinal apoptosis in $banp^{rw337}$ mutants

depend on Δ113tp53-MO, and compared inhibition levels of retinal apoptosis between

FL tp53-MO and Δ113tp53-MO. However, inhibition of apoptosis in $banp^{rw337}$ mutant

retinas by Δ113tp53-MO was significant, but ineffective (n=58.00±11.0/section) (Fig.

4C, 4D). This result is consistent with previous reports proposing that Δ113tp53

preferentially activates transcription of target genes related to cell-cycle arrest rather

than apoptosis (Gong et al., 2015; Gong et al., 2020).

Furthermore, we investigated whether mitotic cell accumulation in $banp^{rw337}$

mutants depend on FL tp53-MO or Δ113tp53-MO. Compared with Standard-MO

injection controls (n=376.6±74.8/section), the number of pH3+ cells in $banp^{rw337}$

mutants was significantly decreased by FL tp53-MO (n=262.4±68.3/section) (Fig. 4C,

4E). However, the number of pH3+ cells was still higher in $banp^{rw337}$ mutant retinas

injected with FL tp53-MO than in wild-type control retinas injected with FL tp53-MO,

indicating that there is a substantial tp53-independent fraction. Furthermore, the

injection of Δ113tp53-MO did not significantly rescue mitotic phenotypes in *banp^{rw337}*

mutants (n=303.8±46.6/section) (Fig. 4C, 4F). These data suggest that there is a tp53-

independent mechanism that causes mitotic cell accumulation in *banp^{rw337}* mutants.

Banp regulates chromatin segregation during mitosis by promoting transcription

of *cenpt* and *ncapg* genes

A substantial fraction of accumulated mitotic cells in *banp^{rw337}* mutants is independent

of tp53 (Fig. 4E). RNA-seq analysis revealed that two chromosomal segregation

regulators, *cenpt* and *ncapg*, are downregulated in *banp^{rw337}* mutants at 48 hpf (Fig. 3B).

Cenpt and Ncapg are essential for chromatin segregation (Hung et al., 2017; Seipold et

al., 2009; Zhang et al., 2018). Using *in situ* hybridization, we confirmed that their

mRNAs are expressed in retinal progenitor cells and that expression of both mRNAs is

markedly decreased in *banp* mutants at 48 hpf (Fig. 5-figure supplement 1).

Quantitative RT-PCR also confirmed their downregulation in *banp^{rw337}* mutants and

banp morphants (Fig. 5A, 5C). Furthermore, ATAC-seq revealed that chromatin

accessibility is decreased near the transcription start site (TSS) of both genes in

banp^{rw337} mutants, leading to reduction of transcription activity (Fig. 5B, 5D). Thus,

Banp is required for transcriptional activation of *cenpt* and *ncapg* genes. Interestingly,

two independent chip-sequencing studies using BANP antibody revealed a specific

binding site within 100 bp of the TSS of *Cenpt* and *Ncapg* genes in mice and humans

(Grand et al., 2021; Mathai et al., 2016), suggesting that the requirement for Banp in

cenpt and *ncapg* transcription is conserved in zebrafish, mice, and humans.

Knockdown of Cenpt and Ncapg causes defects in chromosomal segregation

during mitosis in zebrafish (Hung et al., 2017; Seipold et al., 2009). To examine

whether *banp* mutants show similar chromosome segregation defects, we conducted

time-lapse imaging of mitosis in *banp^{rw337}* mutant retinas for 1-2 h from 48 to 54 hpf,

using a transgenic line *Tg[h2afv:GFP; EF1α:mCherry-zGem]*, which visualizes cell-

cycle phases (Mochizuki et al., 2014). In wild-type retinas injected with control banp

5misMO, chromatin starts to be condensed in prophase. Then, kinetochores and a

mitotic spindle appear in prometaphase, chromosomes are centrally aligned and start to

be pulled toward opposite poles by mitotic spindles in metaphase, sister chromatids

move toward the opposite poles in anaphase, and chromosomes arrive at opposite poles

in telophase, whereupon cytokinesis starts (Fig. 5E, Video 1). The duration from

prophase to telophase was 22.9±5.8 minutes (Fig. 5F). On the other hand, mitosis did

not proceed smoothly in *banp* morphant retinas, especially from prophase to anaphase

(Fig. 5E, Video 2). It took longer from chromosomal condensation in prophase to

chromosomal alignment at the division plane in anaphase in *banp* morphant retinas

(t=15 to t=46 in banp-MO, Fig. 5E). Aligned chromosomes were often dispersed during

chromosomal segregation by the mitotic spindle (t=46 to t=64 in banp-MO, Fig. 5E),

although all 28 cell divisions eventually completed cytokinesis in *banp* morphants. The

duration from prophase to telophase was 49.2± 17.0 minutes and significantly longer in

banp morphants (Fig. 5F). Thus, chromosomal segregation is abnormal during mitosis

in the absence of Banp. Furthermore, we labeled 48-hpf retinas with anti-α-tubulin and

anti-pH3 antibodies to visualize the mitotic spindle and prometaphase chromosomes,

respectively. In *banp*^{rw337} mutants, mitotic spindles are irregularly attached to less

condensed and misaligned chromosomes (Fig. 5G). The percentage of cells with

abnormal mitosis was significantly higher in *banp*^{rw337} mutant retinas than those of wild-

type siblings (Fig. 5H). These data suggest that the interaction between chromosomes

and mitotic spindles is compromised in the absence of Banp, probably due to

kinetochore attachment defects caused by lack of Cenpt and Ncapg. Thus, Banp

regulates chromosomal segregation during mitosis.

Banp regulates multiple gene targets as a transcription activator through the Banp

motif during embryonic development

Recently, it was reported that BANP binds to the nucleotide sequence TCTCGCGAGA,

called the "Banp motif" and regulates transcription of essential metabolic genes in

humans and mice (Grand et al., 2021). To identify direct transcriptional targets of Banp

in zebrafish, we examined whether genes whose mRNA level or chromatin accessibility

is decreased in *banp* mutants carry Banp motif in their promoters. So, we first searched

for enriched motifs in *banp*^{rw337} mutants using a motif discovery software called

Hypergeometric Optimization of Motif EnRichment (HOMER) with ATAC-seq data.

Consistent with a previous report on human and mouse BANP (Grand et al., 2021), our

analysis revealed that TCTCGCGAGA, is the most enriched motif in the closed

chromatin region of *banp^{rw337}* mutants (Fig. 6A and Fig. 6-figure supplement 1A).

Indeed, among 81 down-regulated genes in *banp^{rw337}* mutants (Fig. 3A), we identified

the Banp motif near TSS of 31 genes (Fig. 6B, C and Supplementary File 1). However,

we could not find the Banp motif among 258 upregulated genes (Fig. 6-figure

supplement 1A). These data suggest that Banp is a transcription activator of multiple

gene targets containing Banp motifs. The position of the Banp motif is consistent with

the previous report, that it is not located at a constant distance from TSSs, but rather

between 20 and 70 nucleotides upstream on both strands. We also observed multiple

Banp motifs in some genes (Supplementary File 1). It is likely that the frequency of

Banp motifs upstream of TSSs may correlate with a transcription rate of target genes

(Mahpour et al., 2018).

Next, we examined profiles of these direct-target genes. Among the transcription

targets of Banp that are downregulated in *banp^{rw337}* mutants, 26 percent were involved

in regulating cell cycle (Fig. 6D), including *ncapg* (Fig. 6E and Fig. 6-figure

supplement 1B), *cenpt* (Fig. 6F and Fig. 6-figure supplement 1C), and *wrnip1* (Fig. 6G

and Fig. 6-figure supplement 2A). These data again confirmed that Banp profoundly

regulates chromosome segregation during mitosis by directly activating transcription of

ncapg and *cenpt* genes. In addition, *wrnip1* is interesting, because it protects stalled

replication forks from MRE11-mediated degradation and promotes fork restart after

replication stress (Leuzzi et al., 2016a; Leuzzi et al., 2016b). We confirmed that *wrnip1*

mRNA is normally expressed in retinal progenitor cells in zebrafish and that its

expression is reduced in *banp^{rw337}* mutants at 48 hpf (Fig. 6-figure supplement 2B).

Quantitative RT-PCR analysis revealed a significant reduction of *wrnip1* mRNA

expression in both *banp* mutants and morphants (Fig. 6-figure supplement 2C).

Consistently, RNA-seq data showed downregulation of *wrnip1* (Fig. 6-figure

supplement 2D), although ATAC-seq data showed comparable expression in wild-type

and mutants. These data suggest that Banp is required to maintain *wrnip1* mRNA

expression in retinal progenitor cells in zebrafish. Loss of *wrnip1* is expected to cause

genomic instability due to excessive fork degradation (Leuzzi et al., 2016b). So, the

reduction of *wrnip1* mRNA expression may explain the significant accumulation of γ-

H2AX+ retinal cells in *banp^{rw337}* mutants (Fig. 4-figure supplement 1C, D). Taken

together, these data suggest that Banp regulates transcription of multiple target genes,

including these cell-cycle regulators.

In summary, Banp serves essential functions in cell-cycle regulation in the zebrafish

developing retina (Fig. 7A) . During S phase, Banp suppresses DNA replication stress

by activating transcription of a DNA replication stress regulator, *wrnip1*, and promotes

DNA damage repair in concert with tp53-mediated DNA damage response. In M phase,

Banp promotes mitosis by activating transcription of two chromosome segregation

regulators, *cenpt* and *ncapg*, most likely via their Banp motifs. Thus, Banp regulates

progression in S phase and M phase through transcriptional control of distinct sets of

cell-cycle regulators. In *banp* mutants, stalled DNA replication and subsequent DSBs

are accumulated in S phase, resulting in increase in apoptosis. In M phase, *banp* mutants

show chromosome segregation defects, which prolongs mitosis but eventually complete

it. So, DNA damage mostly induced in S phase is likely to be inherited through mitosis

by daughter cells, which reenter the cell-cycle or become post-mitotic neurons. Neurons

with a high level of DNA damage may undergo apoptosis. Thus, S phase defects and M

phase defects influence each other along cell-cycle progression of retinal progenitor

cells in the absence of Banp (Fig. 7B).

Discussion

BANP was initially identified as a nuclear MAR-binding protein and reported to function as a tumor suppressor (Malonia et al., 2011). Very recently, it was reported that BANP binds to the Banp motif enriched near the TSS of CpG island promoters and regulates transcription of metabolic genes (Grand et al., 2021). However, physiological functions of Banp remained to be discovered. In this study, we identified zebrafish *banp* mutants, in which retinal progenitor cells show mitotic defects and apoptosis during embryonic development. RNA-seq analysis revealed that expression of tp53-dependent DNA damage response regulators is upregulated in *banp* mutants. In non-stress conditions, tp53 is degraded through ubiquitination by an E3-ubiqutin ligase, Mdm2, and subsequent actions of proteasomes, keeping its protein level very low (Oliner et al., 1993). On the other hand, under stress, such as DNA replication stalling or DSBs, ATR and ATM are activated and phosphorylate the N-terminal serine of tp53 (Banin et al., 1998; Canman et al., 1998; Siliciano et al., 1997; Tibbetts et al., 1999). This phosphorylation reduces the binding affinity of Mdm2 for tp53, resulting in stabilization of tp53 protein (Shieh et al., 1997; Yadahalli et al., 2019). Indeed, the tp53 protein level is increased, and tp53 knockdown effectively suppresses retinal apoptosis in *banp* mutants. Furthermore, the number of γ-H2AX+ cells is significantly increased in *banp* mutants, suggesting elevation of replicative DNA damage. These data suggest that DNA replication stress is elevated in *banp* mutants, resulting in activation of tp53-dependent DNA damage response, which subsequently induces apoptosis of retinal progenitor cells.

It is generally accepted that tp53 has dual, but opposing roles: cell survival versus cell death in response to DNA replication stress and DNA damage (Chen, 2016; Lees et al., 2021). In the initial phase of the DNA damage response, tp53 binds to its alternative promoter located in intron 4 and activates transcription of an N-terminal truncated isoform, Δ113-tp53, which preferentially activates transcription of target genes related to cell-cycle arrest to promote DNA repair and cell survival (Chen et al., 2009). However, if DNA damage fails to be repaired, but accumulates chronically, FL tp53, in turn, activates transcription of apoptotic genes to eliminate DNA-damaged cells. Previous *in vitro* studies of human *BANP* suggest that BANP interacts with tp53 to promote transcription of cell-cycle arrest genes and to suppress transcription of apoptotic genes, leading to promotion of its cell survival role (Sinha et al., 2010).

Indeed, our quantitative RT-PCR showed that transcription of Δ113-tp53 is significantly

elevated in *banp* mutants, whereas the mRNA level of FL tp53 is very low. RNA-seq

and ATAC-seq also confirmed a higher level of transcription and chromatin

accessibility of Δ113tp53 in *banp* mutants. These data suggest that Δ113tp53-mediated

transcription is selectively activated in *banp* mutants. Thus, it is important to clarify the

role of tp53, including Δ113tp53 in mitotic cell accumulation in *banp* mutants, because

pH3+ cells are markedly accumulated in *banp* mutant retinas prior to apoptosis.

However, tp53-MO mildly inhibited accumulation of pH3+ cells in *banp* mutants.

There is no significant difference in pH3+ cell accumulation between tp53-MO and

Δ113tp53-MO, or between wild-type siblings and Δ113tp53-MO. So, these data suggest

that the contribution of tp53 to mitotic cell accumulation in *banp* mutants is limited, and

that a tp53-independent pathway is responsible for mitotic phenotypes. Our live

imaging of mitosis in *banp* morphant retinas reveals that mitosis is significantly longer

due to chromosome segregation defects; however, importantly, almost all mitoses are

eventually completed in *banp* mutants, suggesting that accumulation of pH3+ cells in

banp mutants is transient and depends mostly on prolonged mitosis, which is caused by

chromosome segregation defects. At present, we have not observed S-phase arrest in

banp mutants, because the fraction of BrdU+ cells in the total retinal region is not

increased in *banp* mutants, although it is still possible that *banp* mutant retinal cells are

partially arrested in G1 phase, because mRNA expression of a cdk inhibitor, *cdkn1a*

(*p21*), is higher in *banp* mutants than in wild-type siblings. Taken together, we conclude

that tp53 mainly mediates retinal apoptosis, and that its role in mitotic defects is limited

in *banp* mutants.

Labeling with anti-γ-H2AX antibody revealed that replicative DNA damage is

increased in retinal progenitor cells in *banp* mutants, suggesting that DNA lesions that

cause stalled DNA replication are abnormally induced, or stalled DNA replication and

DSBs fail to be repaired in the absence of Banp. Importantly, the number of γ-H2AX+

cells in *banp* mutant retinas was not increased by either tp53-MO and Δ113tp53-MO,

suggesting that tp53 does not repair replicative DNA damage caused by loss of Banp

functions. There are two possible functions of Banp in DNA damage regulation. Banp

may maintain DNA replication integrity. BANP functions as a transcription factor to

regulate expression of essential metabolic genes in pluripotent stem cells and

differentiating neurons (Grand et al., 2021). Banp may promote transcription of DNA

replication regulators. Our RNA-seq data identified 81 downregulated genes in *banp*

mutants, compared to wild-type siblings. Among them, 31 genes have a Banp motif

near their TSSs and contain eight cell-cycle regulatory genes. Among these eight cell-

cycle regulators, interestingly, we identified *wrnip1*. WRNIP1 protects stalled

replication forks, especially fork reversal, from MRE11- and SLX4-mediated

degradation and promotes fork restart by cooperating with BRCA2 to stabilize RAD51

on single-stranded DNA of stalled forks (Leuzzi et al., 2016a; Leuzzi et al., 2016b;

Porebski et al., 2019). In addition, *wrnip1* regulates another fork protection mechanism

called damage bypass, consisting of translesion synthesis (TLS) (Yoshimura et al.,

2019) and template switching (TS) (Jimenez-Martin et al., 2020). WRNIP1 also

functions in interstrand crosslink repair (ICR) by promoting the recruitment of

FANCD2/FANCI complex at the damaged site (Socha et al., 2020). Indeed, *wrnip1*

mRNA expression is markedly reduced in *banp* mutants, suggesting that Banp is

required for maintenance of *wrnip1* transcription. In this case, loss of Wrnip1 may cause

defects in stalled fork stabilization and ICR repair, leading to genomic instability

(Kondratick et al., 2021; Nickoloff et al., 2021; Scully et al., 2021). It will be interesting

to examine whether knockdown of zebrafish *wrnip1* causes similar defects in DNA

replication stress and apoptosis to *banp* mutants. Another possibility is that Banp

regulates DNA repair independent of its role as a transcription factor. A previous study

using human *in vitro* culture cells revealed that BANP promotes HDAC6-mediated

deacetylation of a DNA end-joining protein, Ku70, which is required for non-

homologous end-joining repair (Chaudhary et al., 2014). Since Ku70 is recruited to

DSB sites after γ-H2AX accumulation, Ku70 is an interesting candidate. Since

knockdown of Ku70 along with radiation-induced DNA damage markedly induces

apoptosis in zebrafish embryos (Bladen et al., 2007), it will be interesting to investigate

whether this scenario occurs in zebrafish.

In addition to the role of Banp in DNA replication stress and DNA repair

mechanisms, mRNA expression of two important regulators of chromosome

segregation, *ncapg* and *cenpt*, is significantly downregulated in *banp* mutants. ATAC-

seq and RNA-seq confirmed that chromatin accessibility is decreased near the TSSs of

these genes in *banp* mutants, suppressing transcription of these mRNAs. Our *de novo*

motif enrichment study identified the Banp motif sequence, TCTCGCGAGA, as the
most enriched motif of closed chromatin regions of downregulated genes. Thus, it is
highly likely that Banp directly binds to the Banp motifs of both *ncapg* and *cenpt*, opens
their chromatin, and promotes their transcription. Consistently, as reported in zebrafish
ncapg and *cenpt* knockdown embryos (Hung et al., 2017; Seipold et al., 2009),
chromosome segregation is compromised in mitosis of retinal progenitor cells in *banp*
mutants, suggesting that Banp is an essential regulator of mitosis. Since a Banp motif is
found in promoter regions of *Ncapg* and *Cenpt* in mice and humans, and Chip
sequencing with BANP antibody validated TSS peaks of both *Ncapg* and *Cenpt* genes
in mice and humans (Grand et al., 2021; Mathai et al., 2016), it will be interesting to
confirm whether BANP regulates chromosome segregation during mitosis in mammals.

Lastly, we identified 26 candidate Banp direct-target genes, including cell-cycle
regulators such as *ncapg, cenpt* and *wrnip1,* which carry Banp motifs near their TSSs,
and show markedly decreased chromatin accessibility and transcription in the absence
of Banp. This gene list will provide a useful starting point to investigate the Banp-
mediated transcriptional network. It was reported that binding of BANP to Banp motifs
is dependent on DNA methylation of Banp motifs (Grand et al., 2021), suggesting that
an epigenetic mechanism regulates Banp-mediated transcription. Thus, zebrafish *banp*
mutants provide an excellent new model in which to study how epigenetic factors
regulate cellular processes for embryonic development via Banp motifs of CpG island
promoter genes. In addition, BANP has been proposed as a tumor suppressor (Malonia
et al., 2011). It is generally accepted that DNA methylation of essential genes is altered
in cancer cells (Baylin et al., 2001), which is expected to modify the pattern of gene
transcription through Banp motifs (Grand et al., 2021). It will be interesting to
investigate how the Banp-mediated transcription profile is changed in cancer models of
zebrafish. Future research on zebrafish *banp* mutants will provide more in-depth
understanding of cancer growth and metastasis, as well as new insights into
development of therapeutics for human tumors.

645
646
647

Material and Methods

Zebrafish strains and maintenance

Zebrafish (*Danio rerio*) were maintained using standard procedures (Westerfield, 1993).

RIKEN Wako (RW) was used as a wild-type strain for mutagenesis in which *banp*[rw337]

was generated. Mapping of the *rw337* mutation and analysis of retinal phenotypes of

banp[rw337] mutants were carried out in the genetic background of WIK and Okinawa

wild-type (oki), respectively. Two *banp* mutant alleles, *banp*[rw337] and *banp*[sa12976] were

used. *banp*[sa12976] was obtained from the Zebrafish International Resource Center (ZIRC,

Eugene, OR, USA). Zebrafish transgenic lines, *Tg[h2afv:GFP]*[kca6/kca66] (Pauls et al.,

2001), *Tg[EF1α:mCherry-zGem]*[oki011] (Mochizuki et al., 2014), *Tg[ath5:EGFP]*[rw021]

(Masai et al., 2005) and *Tg[EF1α:mCherry-CAAX]*[oki049] (Mochizuki et al., 2017) were

used to visualize chromatin, S/G2/M phases, RGCs/amacrine cells/photoreceptors, and

plasma membranes, respectively.

Ethics statement

Zebrafish (*Danio rerio*) were maintained on a 14:10 hour light: dark cycle at 28°C.

Collected embryos were cultured in E3 embryo medium (5 mM NaCl, 0.17 mM KCl,

0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$) containing 0.003% 1-phenyl-2-thiouera (PTU) to

prevent pigmentation and 0.01% methylene blue to prevent fungal growth. All

experiments were performed on zebrafish embryos between 36 hpf and 4 dpf prior to

sexual differentiation. Therefore, sexes of the embryos could not be determined.

All zebrafish experiments were carried out in accordance with the Okinawa Institute of

Science and Technology Graduate School's Animal Care and Use Program, which is

based on the National Research Council of the National Academy's Guide for the Care

and Use of Laboratory Animals and has been accredited by the Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC International).

The OIST Institutional Animal Care and Use Committee approved all experimental

protocols (Protocol no. 2019-267, 2019-268, 2019-269, 2019-270).

Zebrafish mutagenesis, mapping and cloning of the *banp* mutant gene

Mutagenesis, mapping, and cloning were performed as previously described (Masai et

al., 2003). The polymorphic markers, 070702B and zC93F2D, shown below were

designed and used for restricting the genomic region covering the *banp*[rw337] mutation.

070702B forward: 5'-ACTTCTTATCAGGGCTGTGC-3'

070702B reverse: 5'-TCAGTCAAGAGCAGTGAGAG-3'

zC93F2D forward: 5'-TGGGATCTCTTTAAGTGAGTGAG-3'

zC93F2D reverse: 5'-TCCAACTATGTGGGTCAAACC-3'

Genotyping of *banp*[rw337] mutant embryos

banp[rw337] mutant phenotypes are difficult to distinguish until 2 dpf. Identification of

homozygous mutant embryos at earlier time points was performed using PCR

amplification of genomic regions using specific primers that enabled restriction

digestion of only mutated sequence using MboII. Primers used, forward: 5'-

CGATGTTGATATCCATCAGTCAGGCGATC-3'; reverse primer: 5'-

GGTGCTGGTGTATAAATCACATGACCTATGGTCCTCTT-3'.

Acridine Orange staining

53-hpf embryos were incubated at 28°C, in 5 mg/mL Acridine Orange (AO) in E3

embryonic medium (5mM NaCl, 0.17mM KCl, 0.33mM $CaCl_2$, 0.33mM $MgSO_4$) for

30 min, protected from light. Embryos were rinsed in E3 medium and used for detection

of AO+ signals. Retinas of live embryos were scanned from the lateral view with a

confocal laser scanning microscope (LSM) (LSM710, Zeiss). After scanning of wild-

type embryos, *banp* mutants, and *banp* mutants injected with mRNA encoding

banp(wt)-EGFP and EGFP-banp(rw337), these scanned embryos were used for DNA

extraction and genotyping. Five samples were selected from each group and used for

statistical analysis.

In situ **hybridization**

Whole-mount *in situ* hybridization of zebrafish embryos was performed as previously

described (Xu et al., 1994). To prepare RNA probes, full-length cDNAs of the zebrafish

banp gene were amplified with Polymerase Chain Reaction (PCR) and sub-cloned into

pBluescript II SK(+) vectors (Stratagene/Agilent Technologies). cDNA fragments were

amplified from mRNA with PCR using the primers for *cenpt*, *ncapg* and *wrnip1*, and

subcloned into pCRII TOPO vector. All the primer sequences of *banp*, *cenpt*, *ncapg*,

and *wrnip1* for PCR amplification are provided in Key Resources Table. Plasmids were

digested with an appropriate restriction enzyme to make DNA templates, which were

used to make antisense or sense RNA probes. Digoxigenin-labeled RNA probes were

generated by *in vitro* transcription using a DIG RNA Labeling Kit (Roche) and purified

using Micro Bio-Spin® Chromatography Columns (BioRad). *In situ* hybridization

signals in whole-mount embryos were detected using anti-DIG alkaline phosphatase

(AP)-conjugated antibody (Roche) with an AP substrate, BM-Purple (Roche) and

imaged using an SteREO Discovery.V12 (ZEISS) microscope.

Rescue experiment with *banp* mutants overexpressing wild-type Banp

The full-length coding region of the *banp* gene was amplified with PCR. To

overexpress wild-type Banp protein, a DNA construct encoding zebrafish wild-type

Banp with an EGFP tag at the N-terminus or C-terminus, EGFP-banp(wt) or banp(wt)-

EGFP, was prepared. A DNA construct encoding an N-terminal EGFP-tagged *rw337*

mutant form of Banp, EGFP-Banp(rw337), was prepared as a negative control. These

DNA constructs were subcloned into the pCS2 expression vector. Capped mRNA was

generated in a standard *in vitro* transcription reaction using an SP6 polymerase

(mMESSAGE mMACHINE™ SP6 Transcription Kit, Invitrogen). We performed

mRNA injections in one-cell-stage embryos obtained from pair-wise crosses of

banp^{rw337} heterozygous mutant male and female fish. At 2 dpf, embryos were used to

determine their genotype with PCR and to identify homozygous mutants. Plastic

sections were obtained to evaluate retinal phenotypes of *banp*^{rw337} mutant embryos

injected with mRNA encoding banp(wt)-EGFP, or EGFP-banp(rw337). Furthermore,

we conducted AO labeling of wild-type embryos, *banp*^{rw337} mutants, *banp*^{rw337} mutants

expressing banp(wt)-EGFP, and *banp*^{rw337} mutants expressing EGFP-banp(rw337). The

fractional AO+ area of the total area of retinal confocal sections was measured using the

LSM software, ZEN (Zeiss), to evaluate the rescue level. We also evaluated protein

stability and nuclear localization of wild-type and *rw337* mutant form of Banp by live

confocal imaging of wild-type retinas expressing banp(wt)-EGFP, EGFP-banp(wt), or

EGFP-banp(rw337) at 24 hpf.

Morpholino injection

ATG morpholino antisense oligos for *banp* (banp MO) and 5 mismatch *banp* MO (banp

5misMO) were designed as 5'-CCACTAAATCTTGCTCTGACATCAT-3' and 5'-

CCtCaAAATgTTcCTCTcACATCAT-3', respectively. Morpholino antisense oligos for

tp53 (tp53 MO) (5'-GCGCCATTGCTTTGCAAGAATTG-3') (Langheinrich et al.,

2002), standard MO (STD MO) (5'-CCTCTTACCTCAGTTACAATTTATA-3') were

used. To knock down Δ113tp53, morpholino antisense oligos for Δ113tp53 (Δ113tp53

MO; also referred to as MO6-tp53 in the zebrafish database ZFIN) (5'-

GCAAGTTTTTGCCAGCTGACAGAAG-3') was used (Chen et al., 2009). All MOs

were injected into one-cell-stage eggs at 1 mM.

Histology

Plastic sectioning, immunostaining of cryosections, paraffin sectioning, and BrdU

incorporation were performed as described previously (Imai et al., 2010), except that we

injected BrdU solution into the yolk of embryos at 48 hpf instead of soaking embryos in

10mM BrdU solution. Anti-pH3 antibody (Cell signalling technology, 6G3, 9706;

Sigma-Aldrich, 06-570), anti-Pax6 antibody (Covance; PRB-278P), zpr1 antibody

(ZIRC, Eugene, Oregon), anti-Prox1 antibody (GeneTex, GTX128354), anti-glutamine

synthetase (GS) (Millipore, MAB302), anti-BrdU antibody (abcam, Ab6326), anti-

PCNA (Sigma-Aldrich, P8825) were used at 1:500, 1:500, 1:100, 1:500, 1:100, 1:200,

1:200 dilutions, respectively. Nuclear staining was performed using SYTOX Green

(Molecular Probes). Filamentous actin (F-actin) was stained using 0.1 mM rhodamine-

conjugated phalloidin (Molecular Probes). TUNEL was performed using the In Situ

Cell Death Detection Kit (Roche).

Whole-mount immunostaining with anti-α-tubulin antibody (Sigma-Aldrich;

#T5168) to label the mitotic spindle was performed as follows. Homozygous mutant

embryos were selected by AO staining, and fixed in 2% trichloroacetic acid (TCA) at

room temperature (RT) for 3 h. Then embryos were washed three times in 1X PBST

(PBS, 0.1% Triton X-100), incubated in 0.2% trypsin for 4 min at 4^{o}C to increase

antibody permeabilization through the skin, and rinsed three times with PBS. After

embryos were re-fixed with 4% PFA for 5 min at 4°C, they were incubated in blocking

solution (10% Goat serum in PBST) for 1 h at room temperature and transferred to 1%

blocking solution with diluted anti-α-tubulin antibody at a 1:1000 dilution for 2 days at

4°C. Embryos were washed with PBST four times for 15 min and then incubated with

fluorescent conjugated anti-mouse antibody in 1% blocking solution in the dark for 2

days at 4°C. Next, all samples were washed in PBST three times for 15 min followed by

70% glycerol storage at 4°C until imaging. Samples were imaged with a confocal

microscope (LSM 880, Zeiss) followed by image analysis using IMARIS software

(ver.9.1.2 Bitplane). Whole-mount immunostaining for anti-pH3 and anti-γ-H2AX

(GeneTex, GTX127342) antibodies were performed as described previously (Sidi et al.,

2008), and for anti-active caspase3 (BD Pharmingen, Clone C92-605) and anti-HuC/D

(Thermo Fisher, A-21271) antibodies as described (Sorrells et al., 2013), except using

goat serum as a blocking reagents. Images were scanned using an LSM710 confocal

laser scanning microscope (Zeiss). The number of fluorescent-positive cells and

percentage of fluorescence-positive area relative to the region of interest was calculated

using original scanning images with IMARIS (Bitplane) and Zen (Zeiss) or Image-J

(NIH) software, respectively.

793 TUNEL of wild-type sibling and *banp^rw337* mutant retinas injected with STD MO,

794 FL tp53 MO, and Δ113tp53 MO

We performed three MO experiments (STD-MO, FL tp53-MO, Δ113tp53-MO injection

to wild-type sibling and *banp*^rw337 mutant embryos) at 48 hpf and two MO experiments

(STD-MO, FL tp53-MO injection to wild-type sibling and *banp*^rw337 mutant embryos) at

72 hpf. After genotyping of all the injected embryos, wild-type and *banp*^rw337 mutant

embryos were selected for TUNEL. First, STD-MO and FL tp53-MO data at 48 and 72

hpf were used for statistical analysis to compare inhibition levels of retinal apoptosis

between STD-MO and FL tp53-MO (Fig. 3H). Next, we used STD-MO, tp53-MO, and

Δ113-tp53-MO data at 48 hpf for statistical analysis to compare inhibition levels of

retinal apoptosis between STD-MO, tp53-MO, and Δ113-tp53-MO (Fig. 4D). Thus,

STD-MO and tp53-MO data at 48 hpf are shared with these two statistical analyses. All

the MO injections were done using embryos produced by *banp* heterozygous embryos,

which were siblings outcorssed from the same *banp* heterozygous parent, in order to
make genetic variation minimum. All the MOs were used at the same concentration.

RT-PCR

Total RNA was extracted from embryo heads at the indicated developmental stages
using a Sepasol RNA I Super (Nacalai, Japan). cDNA synthesis was carried out from
500 ng of RNA using Toyobo ReverTra Ace® qPCR RT master mix with gDNA
remover, according to the manufacturer's instructions. Expression profiles of the genes
listed below were analyzed with the specified primers using Luna® Universal qPCR
Master Mix (NEB). Quantitative RT-PCR was performed using StepOnePlus (Applied
Biosystems). Relative expression was calculated using the ΔΔCt method with the
reference gene *ef1α*. Primers used in qRT-PCR analysis; *mdm2* (Wilkins et al., 2013),
p21(cdkn1a) (Stiff et al., 2016), *FL tp53* and *Δ113 tp53* (Chen et al., 2016), *ef1α*
(McCurley and Callard, 2008). All the primer sequences of *ccng1*, *mdm2*, *p21 (cdkn1a)*,
FL tp53, *Δ113 tp53*, *puma (bbc3)*, *cenpt*, *ncapg*, *atm*, *atr*, *wrnip1*, and *ef1α* are provided
in Key Resources Table.

Western blotting

In total, 21 embryos were divided into 3 groups (each n=7) for 48 hpf wild-type and
banp[rw337] mutants, respectively. Heads were dissected in ice-cold E3 medium. Seven
heads removed from each group were homogenized together in sample buffer (125 mM
NaCl, 50 mM Tris-HCl pH 7.4, 0.5 mM EDTA, 1% Triton X-100, 1X Protease
inhibitor) and used as one sample. Appropriate volumes of samples were diluted to
provide equal protein amounts and used for SDS-PAGE (BIO-RAD, Mini-PROTEAN
TGX Gels). Western blot analysis was performed on 3 samples for wild-type and
banp[rw337] mutants, respectively, according to standard protocols using anti-tp53
antibody (GeneTex, GTX128135) at a 1:1000 dilution. β-actin protein level was
evaluated by western blot analysis of rehybridized membranes with anti-β-actin
antibody (Sigma, A5441) at a 1:5000 dilution, and used to normalize tp53 protein level
for each sample. For secondary antibodies, horseradish peroxidase-conjugated anti-
rabbit or anti-mouse IgG antibodies (Amersham, NA931 and NA934) were used.

Enzyme activity was detected using Immunostar® LD (Wako) and luminescence was

imaged and quantified using iBright 1500 (Thermo Fisher Scientific).

Live imaging

Banp-MO was injected into one-cell-stage eggs of *Tg[h2afv:GFP; EF1α: mCherry-*

zGem] transgenic line. Banp-5misMO was used as a control. Two-dpf embryos were

anesthetized using 5% (v/v) Tricane (3-amino benzoic acid-ethylester) and mounted

laterally using 1% low-melting agarose on a depression slide to image retinas without

left or right preference. Time-lapse images were obtained every 1.1 min using a

confocal laser scanning microscope (LSM710, Zeiss) with a 40X0.8/1.0W Plane

Apochromat lens. Data were evaluated using IMARIS software (ver.9.1.2 Bitplane) to

calculate the time taken to complete each cell division.

RNA sequencing

Embryo heads were homogenized using a Polytron 1200E homogenizer

(KINEMATICA) with TRIzol reagent (Invitrogen). RNA was isolated using a Direct-

zol RNA Miniprep Kit (ZYMO RESEARCH) from three replicates of *banp^{rw337}* mutant

and wild-type embryos at 48 hpf. Each sample contained a pool of 7 embryo heads.

RNA isolation was followed by library preparation for sequencing using an NEBNext®

Ultra™ II Directional RNA Library Prep Kit for Illumina® according to the

manufacturer's instructions. 151-base paired-end sequencing was performed using a

NovaSeq6000 SP. The quality of sequenced reads was evaluated with FastQC (Andrews,

2010) and adaptor trimming and quality filtering were done with fastp (Chen et al.,

2018). Cleaned reads were then mapped to the zebrafish reference genome (GRCz11)

using hisat2.1.0 (Kim et al., 2019) and mapped reads were counted with FeatureCounts

from the Subread package (Liao et al., 2014) using annotation from Ensembl

(Danio_rerio.GRCz11.95.gtf). Differentially expressed gene (DEG) analysis was

performed using the EdgeR package (Robinson and Oshlack, 2010). Genes with false

discovery rates (FDR) < 0.01 were considered significant DEGs. The EnhancedVolcano

package (Blighe et al., 2018) was used to create a volcano plot with $Log_2FoldChange$

(wildtype – mutant) values and FDR calculated with EdgeR. Differences in alternative

splicing between mutant and wild type were tested using the DEXseq package (Anders
et al., 2012). Adjusted p-values < 0.01 were considered as significant.
870

ATAC sequencing
48-hpf embryos were used for ATAC sequencing. We used the Kaestner Lab ATAC seq
protocol (Ackermann and Kaestner, 2019) until library preparation, followed by
sequencing using a NovaSeq6000 SP. We generated reads for three independent
$banp^{rw337}$ mutant samples, as well as three independent $banp^{rw337}$ wild-type sibling
samples. Each sample contained a pool of 3 embryo heads. A quality check of reads was
performed using FastQC. Reads from each sample were adaptor trimmed and aligned
using fastp and Bowtie2 to the zebrafish reference genome (GRCz11) (Langmead and
Salzberg, 2012) respectively. For quality control, aligned reads were filtered for low-
quality reads, non-unique alignments, and PCR duplicates using samtools (Li et al.,
2009; Yan et al., 2020). Moreover, alignment files were peak-called using Genrich, with
removal of mitochondrial reads (Gaspar, 2019). Differences in differentially enriched
peaks in wild-type and mutant embryos were identified using the DiffBind R package
(Stark and Brown, 2021). The obtained peaks were annotated using the annotatePeaks
tool of HOMER software (Heinz et al., 2010). The findMotifsGenome tool of HOMER
was used to find enriched motifs in differentially expressed peaks. To compare coverage
tracks across genes of interest in $banp^{rw337}$ mutants and wild-type fish in RNA-seq and
ATAC-seq data, distributions of mapped reads in each replicate were averaged and
plotted with SparK (Kurtenbach and Harbour, 2019).
890

Statistics
Significant differences between groups (wild type vs mutant) were tested using
GraphPad Prism (GraphPad Software Version 9.1.2). The number of samples used per
experiment is presented in figures or figure legends. Tests used to calculate statistical
significance are mentioned in figure legends. All data are presented as means ± SDs. A
p value of 0.05 was considered statistically significant. In the histogram, p*< 0.05, p**<
0.01, p***< 0.001, p****< 0.0001, and ns (not significant) are indicated.
898
899

Acknowledgements

We thank four former lab members, Masahiro Yamaguchi, Noriko Tonou-Fujimori, Yukihiro Yoshimura, Eri Oguri for supporting the initial phase of mapping and cloning experiments, and three technical staff members, Jeff Liner, Mamoru Fujiwara, and Tetsuya Harakuni for supporting mutant fish maintenance, immunohistochemistry, and DNA construction. We thank SQC, Research support division, Okinawa Institute of Science and Technology Graduate University for assistance in sequencing experiments. We thank Steven D. Aird for editing the manuscript.

Competing interests

The authors declare no competing or financial interests.

Funding

This work was supported by a grant from the Okinawa Institute of Science and Technology Graduate University to IM.

Data availability

Raw RNA-seq and ATAC-seq datasets of *banp^{rw337}* mutant and wild-type siblings are available at DDBJ Sequence Read Archive (DRA012572).

References

Abhiman, S., L.M. Iyer, and L. Aravind. 2008. BEN: a novel domain in chromatin factors
and DNA viral proteins. *Bioinformatics*. 24:458-461.
Ackermann, A., and L. Kaestner. 2019. **UPDATED** ATAC-seq Protocol.
Anders, S., A. Reyes, and W. Huber. 2012. Detecting differential usage of exons from
RNA-seq data. *Genome Research*. 22:2008-2017.
Andrews, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.
Banin, S., L. Moyal, S. Shieh, Y. Taya, C.W. Anderson, L. Chessa, N.I. Smorodinsky, C.
Prives, Y. Reiss, Y. Shiloh, and Y. Ziv. 1998. Enhanced phosphorylation of p53 by
ATM in response to DNA damage. *Science*. 281:1674-1677.
Baylin, S.B., M. Esteller, M.R. Rountree, K.E. Bachman, K. Schuebel, and J.G. Herman.
2001. Aberrant patterns of DNA methylation, chromatin formation and gene
expression in cancer. *Hum Mol Genet*. 10:687-692.
Berti, M., D. Cortez, and M. Lopes. 2020. The plasticity of DNA replication forks in
response to clinically relevant genotoxic stress. *Nat Rev Mol Cell Bio*. 21:633-
651.
Bhagat, P.N., S.H. Jadhav, S. Chattopadhyay, and K.M. Paknikar. 2018. Carbon
nanospheres mediated nuclear delivery of SMAR1 protein (DNA binding
domain) controls breast tumor in mice model. *Nanomedicine (Lond)*. 13:353-
372.
Birot, A., L. Duret, L. Bartholin, B. Santalucia, I. Tigaud, J. Magaud, and J. Rouault. 2000.
Identification and molecular analysis of BANP. *Gene*. 253:189-196.
Blackford, A.N., and S.P. Jackson. 2017a. ATM, ATR, and DNA-PK: The Trinity at the
Heart of the DNA Damage Response. *Molecular Cell*. 66:801-817.
Blackford, A.N., and S.P. Jackson. 2017b. ATM, ATR, and DNA-PK: The Trinity at the
Heart of the DNA Damage Response. *Mol Cell*. 66:801-817.
Bladen, C.L., W.K. Lam, W.S. Dynan, and D.J. Kozlowski. 2005. DNA damage response
and Ku80 function in the vertebrate embryo. *Nucleic Acids Res*. 33:3002-3010.
Bladen, C.L., S. Navarre, W.S. Dynan, and D.J. Kozlowski. 2007. Expression of the Ku70
subunit (XRCC6) and protection from low dose ionizing radiation during
zebrafish embryogenesis. *Neurosci Lett*. 422:97-102.
Blasquez, V.C., A.O. Sperry, P.N. Cockerill, and W.T. Garrard. 1989. Protein - DNA
Interactions at Chromosomal Loop Attachment Sites. *Genome*. 31:503-509.
Blighe, K., S. Rana, and M. Lewis. 2018. EnhancedVolcano: Publication-ready volcano
plots with enhanced colouring and labeling.
Boije, H., R.B. MacDonald, and W.A. Harris. 2014. Reconciling competence and
transcriptional hierarchies with stochasticity in retinal lineages. *Curr Opin
Neurobiol*. 27:68-74.
Canman, C.E., D.S. Lim, K.A. Cimprich, Y. Taya, K. Tamai, K. Sakaguchi, E. Appella, M.B.
Kastan, and J.D. Siliciano. 1998. Activation of the ATM kinase by ionizing
radiation and phosphorylation of p53. *Science*. 281:1677-1679.
Chattopadhyay, S., R. Kaul, A. Charest, D. Housman, and J.Z. Chen. 2000. SMAR1, a
novel, alternatively spliced gene product, binds the scaffold/matrix-associated
region at the T cell receptor beta locus. *Genomics*. 68:93-96.

Chaudhary, N., K.K. Nakka, P.L. Chavali, J. Bhat, S. Chatterjee, and S. Chattopadhyay.
2014. SMAR1 coordinates HDAC6-induced deacetylation of Ku70 and dictates
cell fate upon irradiation. *Cell Death Dis*. 5:e1447.
Chemmannur, S.V., A.J. Badhwar, B. Mirlekar, S.K. Malonia, M. Gupta, N. Wadhwa, R.
Bopanna, U. Mabalirajan, S. Majumdar, B. Ghosh, and S. Chattopadhyay. 2015.
Nuclear matrix binding protein SMAR1 regulates T-cell differentiation and
allergic airway disease. *Mucosal Immunol*. 8:1201-1211.
Chen, J. 2016. The Cell-Cycle Arrest and Apoptotic Functions of p53 in Tumor Initiation
and Progression. *Cold Spring Harb Perspect Med*. 6:a026104.
Chen, J., S.M. Ng, C. Chang, Z. Zhang, J.C. Bourdon, D.P. Lane, and J. Peng. 2009. p53
isoform delta113p53 is a p53 target gene that antagonizes p53 apoptotic
activity via BclxL activation in zebrafish. *Genes Dev*. 23:278-290.
Chen, J., H. Ruan, S.M. Ng, C. Gao, H.M. Soo, W. Wu, Z. Zhang, Z. Wen, D.P. Lane, and J.
Peng. 2005. Loss of function of def selectively up-regulates Delta113p53
expression to arrest expansion growth of digestive organs in zebrafish. *Genes
Dev*. 19:2900-2911.
Chen, S.F., Y.Q. Zhou, Y.R. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*. 34:884-890.
Chen, Y.C., S. Semenova, S. Rozov, M. Sundvik, J.L. Bonkowsky, and P. Panula. 2016. A
Novel Developmental Role for Dopaminergic Signaling to Specify Hypothalamic
Neurotransmitter Identity. *Journal of Biological Chemistry*. 291:21880-21892.
Dowling, J.E. 2012. The Retina, An Approachable Part of the Brain. Harvard University
press.
Gaspar, J.M. 2019. ATAC-seq Guidelines.
Gong, L., H. Gong, X. Pan, C. Chang, Z. Ou, S. Ye, L. Yin, L. Yang, T. Tao, Z. Zhang, C. Liu,
D.P. Lane, J. Peng, and J. Chen. 2015. p53 isoform Delta113p53/Delta133p53
promotes DNA double-strand break repair to protect cell from death and
senescence in response to DNA damage. *Cell Res*. 25:351-369.
Gong, L., X. Pan, G.K. Abali, J.B. Little, and Z.M. Yuan. 2020. Functional interplay
between p53 and Delta 133p53 in adaptive stress response. *Cell Death and
Differentiation*. 27:1618-1632.
Grand, R.S., L. Burger, C. Grawe, A.K. Michael, L. Isbel, D. Hess, L. Hoerner, V.
Iesmantavicius, S. Durdu, M. Pregnolato, A.R. Krebs, S.A. Smallwood, N. Thoma,
M. Vermeulen, and D. Schubeler. 2021. BANP opens chromatin and activates
CpG-island-regulated genes. *Nature*.
Hambarde, S., C.L. Tsai, R.K. Pandita, A. Bacolla, A. Maitra, V. Charaka, C.R. Hunt, R.
Kumar, O. Limbo, R. Le Meur, W.J. Chazin, S.E. Tsutakawa, P. Russell, K.
Schlacher, T.K. Pandita, and J.A. Tainer. 2021. EXO5-DNA structure and BLM
interactions direct DNA resection critical for ATR-dependent replication restart.
Mol Cell. 81:2989-3006 e2989.
Hart, C.M., and U.K. Laemmli. 1998. Facilitation of chromatin dynamics by SARs.
Current Opinion in Genetics & Development. 8:519-525.
Heinz, S., C. Benner, N. Spann, E. Bertolino, Y.C. Lin, P. Laslo, J.X. Cheng, C. Murre, H.
Singh, and C.K. Glass. 2010. Simple combinations of lineage-determining

transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*. 38:576-589.

Hung, C.Y., B. Volkmar, J.D. Baker, J.W. Bauer, E. Gussoni, S. Hainzl, A. Klausegger, J. Lorenzo, I. Mihalek, O. Rittinger, M. Tekin, J.E. Dallman, and O.A. Bodamer. 2017. A defect in the inner kinetochore protein CENPT causes a new syndrome of severe growth failure. *PLoS One*. 12:e0189324.

Imai, F., A. Yoshizawa, N. Fujimori-Tonou, K. Kawakami, and I. Masai. 2010. The ubiquitin proteasome system is required for cell proliferation of the lens epithelium and for differentiation of lens fiber cells in zebrafish. *Development*. 137:3257-3268.

Jimenez-Martin, A., I. Saugar, C.R. Joseph, A. Mayer, C.P. Lehmann, B. Szakal, D. Branzei, and J.A. Tercero. 2020. The Mgs1/WRNIP1 ATPase is required to prevent a recombination salvage pathway at damaged replication forks. *Sci Adv*. 6:eaaz3327.

Joruiz, S.M., J.A. Beck, I. Horikawa, and C.C. Harris. 2020. The Delta 133p53 Isoforms, Tuners of the p53 Pathway. *Cancers*. 12.

Joruiz, S.M., and J.C. Bourdon. 2016. p53 Isoforms: Key Regulators of the Cell Fate Decision. *Cold Spring Harb Perspect Med*. 6.

Jusuf, P.R., and W.A. Harris. 2009. Ptf1a is expressed transiently in all types of amacrine cells in the embryonic zebrafish retina. *Neural Dev*. 4.

Kaul, R., S. Mukherjee, F. Ahmed, M.K. Bhat, R. Chhipa, S. Galande, and S. Chattopadhyay. 2003. Direct interaction with and activation of p53 by SMAR1 retards cell-cycle progression at G2/M phase and delays tumor growth in mice. *Int J Cancer*. 103:606-615.

Kaul-Ghanekar, R., S. Majumdar, A. Jalota, N. Gulati, N. Dubey, B. Saha, and S. Chattopadhyay. 2005. Abnormal V(D)J recombination of T cell receptor beta locus in SMAR1 transgenic mice. *J Biol Chem*. 280:9450-9459.

Kim, D., J.M. Paggi, C. Park, C. Bennett, and S.L. Salzberg. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*. 37:907-+.

Kondratick, C.M., M.T. Washington, and M. Spies. 2021. Making Choices: DNA Replication Fork Recovery Mechanisms. *Semin Cell Dev Biol*. 113:27-37.

Kurtenbach, S., and J.W. Harbour. 2019. SparK: A Publication-quality NGS Visualization Tool. *bioRxiv*.

Langheinrich, U., E. Hennen, G. Stott, and G. Vacun. 2002. Zebrafish as a model organism for the identification and characterization of drugs and genes affecting p53 signaling. *Curr Biol*. 12:2023-2028.

Langmead, B., and S.L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*. 9:357-U354.

Larison, K.D., and R. Bremiller. 1990. Early Onset of Phenotype and Cell Patterning in the Embryonic Zebrafish Retina. *Development*. 109:567-&.

Lees, A., T. Sessler, and S. McDade. 2021. Dying to Survive-The p53 Paradox. *Cancers (Basel)*. 13.

Leuzzi, G., V. Marabitti, P. Pichierri, and A. Franchitto. 2016a. WRNIP1 protects stalled
forks from degradation and promotes fork restart after replication stress.
EMBO J. 35:1437-1451.
Leuzzi, G., V. Marabitti, P. Pichierri, and A. Franchitto. 2016b. WRNIP1: A new guardian
of genome integrity at stalled replication forks. *Mol Cell Oncol*. 3:e1215777.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.
Durbin, and G.P.D. Proc. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinformatics*. 25:2078-2079.
Li, Z., M. Hu, M.J. Ochocinska, N.M. Joseph, and S.S. Easter, Jr. 2000. Modulation of cell
proliferation in the embryonic retina of zebrafish (Danio rerio). *Dev Dyn*.
219:391-401.
Liao, Y., G.K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics*.
30:923-930.
Liu, H.C., F. Ma, Y. Shen, Y.Q. Hu, and S. Pan. 2014. Overexpression of SMAR1 Enhances
Radiosensitivity in Human Breast Cancer Cell Line MCF7 via Activation of p53
Signaling Pathway. *Oncol Res*. 22:293-300.
Liu, T.X., N.G. Howlett, M. Deng, D.M. Langenau, K. Hsu, J. Rhodes, J.P. Kanki, A.D.
D'Andrea, and A.T. Look. 2003. Knockdown of zebrafish Fancd2 causes
developmental abnormalities via p53-dependent apoptosis. *Dev Cell*. 5:903-914.
Macdonald, R., and S.W. Wilson. 1997. Distribution of Pax6 protein during eye
development suggests discrete roles in proliferative and differentiated visual
cells. *Dev Genes Evol*. 206:363-369.
Mahpour, A., B.S. Scruggs, D. Smiraglia, T. Ouchi, and I.H. Gelman. 2018. A methyl-
sensitive element induces bidirectional transcription in TATA-less CpG island-
associated promoters. *Plos One*. 13.
Malonia, S.K., S. Sinha, P. Lakshminarasimhan, K. Singh, A. Jalota-Badhwar, S. Rampalli,
R. Kaul-Ghanekar, and S. Chattopadhyay. 2011. Gene regulation by SMAR1:
Role in cellular homeostasis and cancer. *Biochim Biophys Acta*. 1815:1-12.
Masai, I., Z. Lele, M. Yamaguchi, A. Komori, A. Nakata, Y. Nishiwaki, H. Wada, H. Tanaka,
Y. Nojima, M. Hammerschmidt, S.W. Wilson, and H. Okamoto. 2003. N-
cadherin mediates retinal lamination, maintenance of forebrain compartments
and patterning of retinal neurites. *Development*. 130:2479-2494.
Masai, I., D.L. Stemple, H. Okamoto, and S.W. Wilson. 2000. Midline signals regulate
retinal neurogenesis in zebrafish. *Neuron*. 27:251-263.
Masai, I., M. Yamaguchi, N. Tonou-Fujimori, A. Komori, and H. Okamoto. 2005. The
hedgehog-PKA pathway regulates two distinct steps of the differentiation of
retinal ganglion cells: the cell-cycle exit of retinoblasts and their neuronal
maturation. *Development*. 132:1539-1553.
Mathai, J., S.P. Mittal, A. Alam, P. Ranade, D. Mogare, S. Patel, S. Saxena, S. Ghorai, A.P.
Kulkarni, and S. Chattopadhyay. 2016. SMAR1 binds to T(C/G) repeat and
inhibits tumor progression by regulating miR-371-373 cluster. *Sci Rep*. 6:33779.

McCurley, A.T., and G.V. Callard. 2008. Characterization of housekeeping genes in
zebrafish: male-female differences and effects of tissue type, developmental
stage and chemical treatment. *Bmc Molecular Biology*. 9.

Mochizuki, T., Y.J. Luo, H.F. Tsai, A. Hagiwara, and I. Masai. 2017. Cell division and
cadherin-mediated adhesion regulate lens epithelial cell movement in zebrafish.
Development. 144:708-719.

Mochizuki, T., S. Suzuki, and I. Masai. 2014. Spatial pattern of cell geometry and cell-
division orientation in zebrafish lens epithelium. *Biol Open*. 3:982-994.

Murga, M., S. Bunting, M.F. Montana, R. Soria, F. Mulero, M. Canamero, Y. Lee, P.J.
McKinnon, A. Nussenzweig, and O. Fernandez-Capetillo. 2009. A mouse model
of ATR-Seckel shows embryonic replicative stress and accelerated aging. *Nat
Genet*. 41:891-U846.

Nickoloff, J.A., N. Sharma, L. Taylor, S.J. Allen, and R. Hromas. 2021. The Safe Path at
the Fork: Ensuring Replication-Associated DNA Double-Strand Breaks are
Repaired by Homologous Recombination. *Front Genet*. 12:748033.

Oliner, J.D., J.A. Pietenpol, S. Thiagalingam, J. Gyuris, K.W. Kinzler, and B. Vogelstein.
1993. Oncoprotein MDM2 conceals the activation domain of tumour
suppressor p53. *Nature*. 362:857-860.

Park, H.C., S.K. Hong, H.S. Kim, S.H. Kim, E.J. Yoon, C.H. Kim, N. Miki, and T.L. Huh. 2000.
Structural comparison of zebrafish elav/Hu and their differential expressions
during neurogenesis. *Neuroscience Letters*. 279:81-84.

Pauls, S., B. Geldmacher-Voss, and J.A. Campos-Ortega. 2001. A zebrafish histone
variant H2A.F/Z and a transgenic H2A.F/Z:GFP fusion protein for in vivo studies
of embryonic development. *Dev Genes Evol*. 211:603-610.

Pavithra, L., S. Mukherjee, K. Sreenath, S. Kar, K. Sakaguchi, S. Roy, and S.
Chattopadhyay. 2009. SMAR1 forms a ternary complex with p53-MDM2 and
negatively regulates p53-mediated transcription. *J Mol Biol*. 388:691-702.

Peterson, R.E., J.M. Fadool, J. McClintock, and P.J. Linser. 2001. Muller cell
differentiation in the zebrafish neural retina: Evidence of distinct early and late
stages in cell maturation. *Journal of Comparative Neurology*. 429:530-540.

Porebski, B., S. Wild, S. Kummer, S. Scaglione, P.L. Gaillard, and K. Gari. 2019. WRNIP1
Protects Reversed DNA Replication Forks from SLX4-Dependent Nucleolytic
Cleavage. *iScience*. 21:31-41.

Prigent, C., and S. Dimitrov. 2003. Phosphorylation of serine 10 in histone H3, what
for? *J Cell Sci*. 116:3677-3685.

Robinson, M.D., and A. Oshlack. 2010. A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome Biol*. 11.

Rogakou, E.P., D.R. Pilch, A.H. Orr, V.S. Ivanova, and W.M. Bonner. 1998a. DNA double-
stranded breaks induce histone H2AX phosphorylation on serine 139. *Journal of
Biological Chemistry*. 273:5858-5868.

Rogakou, E.P., D.R. Pilch, A.H. Orr, V.S. Ivanova, and W.M. Bonner. 1998b. DNA double-
stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol
Chem*. 273:5858-5868.

Scheuermann, R.H., and W.T. Garrard. 1999. MARs of antigen receptor and co-
receptor genes. *Crit Rev Eukar Gene*. 9:295-310.
Scully, R., R. Elango, A. Panday, and N.A. Willis. 2021. Recombination and restart at
blocked replication forks. *Curr Opin Genet Dev*. 71:154-162.
Seipold, S., F.C. Priller, P. Goldsmith, W.A. Harris, H. Baier, and S. Abdelilah-Seyfried.
2009. Non-SMC condensin I complex proteins control chromosome segregation
and survival of proliferating cells in the zebrafish neural retina. *BMC Dev Biol*.
9:40.
Shieh, S.Y., M. Ikeda, Y. Taya, and C. Prives. 1997. DNA damage-induced
phosphorylation of p53 alleviates inhibition by MDM2. *Cell*. 91:325-334.
Shroff, R., A. Arbel-Eden, D. Pilch, G. Ira, W.M. Bonner, J.H. Petrini, J.E. Haber, and M.
Lichten. 2004. Distribution and dynamics of chromatin modification induced by
a defined DNA double-strand break. *Curr Biol*. 14:1703-1711.
Sidi, S., T. Sanda, R.D. Kennedy, A.T. Hagen, C.A. Jette, R. Hoffmans, J. Pascual, S.
Imamura, S. Kishi, J.F. Amatruda, J.P. Kanki, D.R. Green, A.A. D'Andrea, and A.T.
Look. 2008. Chk1 suppresses a caspase-2 apoptotic response to DNA damage
that bypasses p53, Bcl-2, and caspase-3. *Cell*. 133:864-877.
Siliciano, J.D., C.E. Canman, Y. Taya, K. Sakaguchi, E. Appella, and M.B. Kastan. 1997.
DNA damage induces phosphorylation of the amino terminus of p53. *Genes*
Dev. 11:3471-3481.
Simossis, V.A., and J. Heringa. 2005. PRALINE: a multiple sequence alignment toolbox
that integrates homology-extended and secondary structure information.
Nucleic Acids Research. 33:W289-W294.
Sinha, S., S.K. Malonia, S.P. Mittal, J. Mathai, J.K. Pal, and S. Chattopadhyay. 2012.
Chromatin remodelling protein SMAR1 inhibits p53 dependent transactivation
by regulating acetyl transferase p300. *Int J Biochem Cell Biol*. 44:46-52.
Sinha, S., S.K. Malonia, S.P. Mittal, K. Singh, S. Kadreppa, R. Kamat, R. Mukhopadhyaya,
J.K. Pal, and S. Chattopadhyay. 2010. Coordinated regulation of p53 apoptotic
targets BAX and PUMA by SMAR1 through an identical MAR element. *EMBO J*.
29:830-842.
Sirbu, B.M., F.B. Couch, J.T. Feigerle, S. Bhaskara, S.W. Hiebert, and D. Cortez. 2011.
Analysis of protein dynamics at active, stalled, and collapsed replication forks.
Genes Dev. 25:1320-1327.
Socha, A., D. Yang, A. Bulsiewicz, K. Yaprianto, M. Kupculak, C.C. Liang, A.
Hadjicharalambous, R. Wu, S.P. Gygi, and M.A. Cohn. 2020. WRNIP1 Is
Recruited to DNA Interstrand Crosslinks and Promotes Repair. *Cell Rep*.
32:107850.
Sorrells, S., C. Toruno, R.A. Stewart, and C. Jette. 2013. Analysis of Apoptosis in
Zebrafish Embryos by Whole-mount Immunofluorescence to Detect Activated
Caspase 3. *Jove-J Vis Exp*.
Stark, R., and G. Brown. 2021. DiffBind: Differential binding analysis of ChIP- Seq peak
data.
Stiff, T., T.C. Tena, M. O'Driscoll, P.A. Jeggo, and M. Philipp. 2016. ATR promotes cilia
signalling: links to developmental impacts. *Hum Mol Genet*. 25:1574-1587.

Taye, N., A. Alam, S. Ghorai, D.G. Chatterji, A. Parulekar, D. Mogare, S. Singh, P.
Sengupta, S. Chatterjee, M.K. Bhat, M.K. Santra, P.B. Salunkhe, S.K. Finston, and
S. Chattopadhyay. 2018. SMAR1 inhibits Wnt/beta-catenin signaling and
prevents colorectal cancer progression. *Oncotarget*. 9:21322-21336.
Tibbetts, R.S., K.M. Brumbaugh, J.M. Williams, J.N. Sarkaria, W.A. Cliby, S.Y. Shieh, Y.
Taya, C. Prives, and R.T. Abraham. 1999. A role for ATR in the DNA damage-
induced phosphorylation of p53. *Genes Dev*. 13:152-157.
Tsuda, H., D.F. Callen, T. Fukutomi, Y. Nakamura, and S. Hirohashi. 1994. Allele loss on
chromosome 16q24.2-qter occurs frequently in breast cancers irrespectively of
differences in phenotype and extent of spread. *Cancer Res*. 54:513-517.
Wang, X., L. Gleich, Z.P. Pavelic, Y.Q. Li, N. Gale, S. Hunt, J.L. Gluckman, and P.J.
Stambrook. 1999. Cervical metastases of head and neck squamous cell
carcinoma correlate with loss of heterozygosity on chromosome 16q. *Int J*
Oncol. 14:557-561.
Ward, I.M., and J. Chen. 2001. Histone H2AX is phosphorylated in an ATR-dependent
manner in response to replicational stress. *J Biol Chem*. 276:47759-47762.
Waterman, D.P., J.E. Haber, and M.B. Smolka. 2020. Checkpoint Responses to DNA
Double-Strand Breaks. *Annu Rev Biochem*. 89:103-133.
Westerfield, M. 1993. The zebrafish book : a guide for the laboratory use of zebrafish
(Brachydanio rerio).
Wilkins, B.J., K. Lorent, R.P. Matthews, and M. Pack. 2013. p53-Mediated Biliary
Defects Caused by Knockdown of cirh1a, the Zebrafish Homolog of the Gene
Responsible for North American Indian Childhood Cirrhosis. *Plos One*. 8.
Xu, Q., N. Holder, R. Patient, and S.W. Wilson. 1994. Spatially regulated expression of
three receptor tyrosine kinase genes during gastrulation in the zebrafish.
Development. 120:287-299.
Yadahalli, S., J.L. Neira, C.M. Johnson, Y.S. Tan, P.J.E. Rowling, A. Chattopadhyay, C.S.
Verma, and L.S. Itzhaki. 2019. Kinetic and thermodynamic effects of
phosphorylation on p53 binding to MDM2. *Sci Rep*. 9:693.
Yamaguchi, M., N. Fujimori-Tonou, Y. Yoshimura, T. Kishi, H. Okamoto, and I. Masai.
2008. Mutation of DNA primase causes extensive apoptosis of retinal neurons
through the activation of DNA damage checkpoint and tumor suppressor p53.
Development. 135:1247-1257.
Yan, F., D.R. Powell, D.J. Curtis, and N.C. Wong. 2020. From reads to insight: a
hitchhiker's guide to ATAC-seq data analysis. *Genome Biol*. 21.
Yildirim, C., D.Y. Vogel, M.R. Hollander, J.M. Baggen, R.D. Fontijn, S. Nieuwenhuis, A.
Haverkamp, M.R. de Vries, P.H. Quax, J.J. Garcia-Vallejo, A.M. van der Laan, C.D.
Dijkstra, T.C. van der Pouw Kraan, N. van Royen, and A.J. Horrevoets. 2015.
Galectin-2 induces a proinflammatory, anti-arteriogenic phenotype in
monocytes and macrophages. *PLoS One*. 10:e0124347.
Yoshimura, A., M. Oikawa, H. Jinbo, Y. Hasegawa, T. Enomoto, and M. Seki. 2019.
WRNIP1 Controls the Amount of PrimPol. *Biol Pharm Bull*. 42:764-769.

Zhang, Q., R. Su, C. Shan, C. Gao, and P. Wu. 2018. Non-SMC Condensin I Complex,
Subunit G (NCAPG) is a Novel Mitotic Gene Required for Hepatocellular Cancer
Cell Proliferation and Migration. *Oncol Res*. 26:269-276.
1229

 Figure Legends

 Figure 1. Zebrafish *banp*^{rw337} mutants show mitotic cell accumulation and cell

 death in the developing retina.

(A) Morphology of wild-type and *rw337* mutant embryos at 4 dpf. *rw337* mutant

embryos have smaller eyes compared to wild type at 4 dpf (red arrow). The black

arrow shows the cloudy tectum in *rw337* mutant embryos, representing cell death.

Scale bars: 100 μm.

(B) Plastic sections of wild-type and *rw337* mutant retinas during development. Higher

magnification images of green squares in the left columns are shown in the right

columns. Retinal neuroepithelium is normal in *rw337* mutants until 2 dpf. However,

many round cells reminiscent of mitotic cells are accumulated in the apical region

of *rw337* mutant retinas at 2 dpf (open green arrowheads), suggesting mitotic

defects in retinal progenitor cells. In wild-type retinas, three nuclear layers (RGCL,

INL, and ONL) and two plexiform layers (IPL, and OPL) are formed at 2.5 dpf.

INL consists of amacrine cells (ACs), bipolar cells (BPs), and horizontal cells

(HCs). ONL consists of two types of photoreceptors (PRs), rods and cones.

However, only RGCs appear near the lens and the outer region was occupied by

pyknotic nuclei, indicating dead cells in *rw337* mutants. At 3 dpf, INL is formed,

but the outer region is still occupied by pyknotic nuclei in *rw337* mutants. At 4 dpf,

pyknotic nuclei are reduced in *rw337* mutants; however, ONL and OPL are absent

despite the small number of photoreceptor-like columnar cells in putative INL (PR,

asterisks). At 3 and 4 dpf, the CMZ is maintained in *rw337* mutants (green double

asterisks).

(C) The *rw337* mutation is mapped in the genomic region flanked by two polymorphic

markers, 070702B and zC93F2D, on chromosome 25. This genomic region contains

four genes, including *banp*.

(D) *banp* cDNA consists of 13 exons. In *rw337* mutants, T at the donor splice site of the

intron between exons 4 and 5 (intron4/5) is converted to G, leading to the insertion

of 52 nucleotides (red region) from intron 4/5. A new stop codon appears within

this insertion.

(E) Wild-type Banp protein has a coiled-coil domain, a BEN domain and a DNA-

binding domain. An NLS domain, an RS-rich domain, and MAR-binding domain

are shown. A truncated protein missing BEN and DNA-binding domains is

predicted in *rw337* mutants.

(F) Retinal sections of wild-type, wild-type embryos injected with Banp-5misMO, and

banp MO. *banp* morphant retinas have similar phenotypes to *banp^{rw337}* mutants,

with increased cell death and loss of the ONL. Yellow open arrowheads show

pyknotic nuclei.

(G) Retinal sections of wild-type retinas, *banp^{rw337}* mutant retinas, and *banp^{rw337}* mutant

retinas overexpressing Banp(wt)-EGFP and EGFP-Banp(rw337). Overexpression of

Banp(wt)-EGFP inhibits cell death and recovers retinal lamination in *banp^{rw337}*

mutants, whereas overexpression of EGFP-Banp(rw337) does not inhibit retinal

defects in *banp^{rw337}* mutants. Yellow open arrowheads indicate pyknotic cells in

banp^{rw337} mutant retinas and *banp^{rw337}* mutant retinas overexpressing EGFP-

Banp(rw337).

Scale bars: 20 μm for (B, F, G).

RGC, retinal ganglion cell; AC, amacrine cell, BP, bipolar cell; HC, horizontal cell; PR,

photoreceptor; OPL, outer plexiform layer; IPL, inner plexiform layer; INL, inner

nuclear layer; RGCL, retinal ganglion cell layer; CMZ, ciliary marginal zone.

Figure Supplement 1. Retinal cell differentiation in *banp^{rw337}* mutants at 4 dpf.

Figure Supplement 2. Amino acid conservation of Banp protein between zebrafish,

mice and humans.

Figure Supplement 3. *banp^{sa12976}* homozygous mutants phenocopy *rw337* mutants.

Figure Supplement 4. Confirmation of specificity of banp MO and nuclear localization

of zebrafish Banp protein.

Figure Supplement 5. Overexpression of Banp(wt)-EGFP reduces cell death in

banp^{rw337} mutants.

Figure Supplement 6. Spatio-temporal pattern of zebrafish *banp* mRNA expression.

1288

Figure 2. *banp* mutant retinas show mitotic cell accumulation and apoptosis.

(A) Double labeling of wild-type (upper) and *banp^{rw337}* mutant (lower) retinas with

TUNEL (red) and anti-pH3 antibody (blue). Nucleus was counterstained with

Sytoxgreen (green).

(B) Only TUNEL and nuclear fluorescent channel of images shown in Fig. 2A. TUNEL
and nuclear staining are indicated in magenta and green, respectively. In $banp^{rw337}$
mutants, apoptotic cells start to be observed in neural retinas at 53 hpf, and increase
in the apical region of retinas (white arrowheads) and at the interface between the
CMZ and the central retina (yellow open arrowheads) at 77 and 101 hpf.

(B') Only pH3 and nuclear fluorescent channel of images shown in Fig. 2A. pH3
signals and nuclear staining are indicated in magenta and green, respectively.
Mitotic cells are accumulated at the apical surface of $banp^{rw337}$ mutant retinas at 53
hpf (white open arrowheads) and restricted in the apical region of the CMZ (yellow
open arrow) at 77 and 101 hpf.

(C) The number of TUNEL-positive cells per retina. Apoptosis is significantly
increased in $banp^{rw337}$ mutants after 53 hpf. Unpaired t-test, two-tailed, Mean ± SD.
[n=3, p**< 0.01]

(D) The number of pH3-positive cells per retina. The number of mitotic cells is
significantly increased in $banp^{rw337}$ mutants at 53 hpf. The number is decreased, but
still significantly higher in $banp^{rw337}$ mutants than in wild-type siblings after 77 hpf.
Unpaired t-test, two-tailed, Mean ± SD. [n=3, p**< 0.01]

(E) Schematic diagram of wild-type retinas. Retinal progenitor cells undergo cell-cycle
progression in which the nuclei move along the apico-basal axis depending on cell-
cycle phase (the most basal location in S phase and the most apical location in M
phase), called interkinetic nuclear migration. Mitosis occurs in the apical surface of
the neural retina. Daughter cells reenter the cell cycle as retinal progenitor cells or
start to differentiate into postmitotic neurons, which migrate into the basal region
and become mature neurons.

(F) Schematic diagram of $banp^{rw337}$ mutant retinas. A fraction of interphase progenitor
cells (G1, S, or G2 phase cells) undergo apoptosis after 48hpf. Mitotic cells are
accumulated at the apical surface, which may be caused by prolonged mitotic
duration. A fraction of immature differentiating neurons also undergo apoptosis.
Thus, $banp^{rw337}$ mutant retinal cells show multiple defects in cell-cycle progression
of retinal progenitor cells and retinal neurogenesis.

Scale bars: 20 μm for (A, B, B').

Source data 1. Data for Figure 2CD.

Figure Supplement 1. Apoptosis is induced in retinal progenitor cells and newly
differentiating neurons in *banp*^{rw337} mutants.

1327

Figure 3. tp53-mediated DNA damage response is activated in the absence of Banp.

(A) A volcano plot indicating gene differential expression of 48 hpf wild-type versus
banp^{rw337} mutant embryos. Red dots show genes that were upregulated ≥2-fold in
mutants, while blue dots represent genes downregulated ≥ 2-fold.

(B) Heat map of the top 15 differentially expressed genes.

(C) Relative mRNA expression levels for *bbc3*, *ccng1*, *mdm2*, *cdkn1a* in wild type and
banp^{rw337} mutants. Quantitative RT-PCR confirmed the up-regulation of p53
pathway genes in *banp*^{rw337} mutants at 48 hpf. Unpaired t-test (two-tailed), Mean ±
SD. [n=3, p*< 0.05, p**< 0.01, p***< 0.001, p**** < 0.0001].

(D) Western blotting of wild-type and *banp*^{rw337} mutants with anti-tp53 antibody. Full-
length tp53 protein is stabilized in *banp*^{rw337} mutants, compared to wild type. β-actin
protein is an endogenous control and used for normalization of tp53 protein level.

(E) Quantification of tp53 protein level relative to a control protein, β-actin, in wild-
type and *banp*^{rw337} mutants. The tp53 protein level is significantly higher in
banp^{rw337} mutants than in wild type at 48 hpf. Unpaired t-test (two-tailed), Mean ±
SD. [n=3, p*< 0.05].

(F) Relative mRNA expression levels for *atm* and *atr* in wild type and *banp*^{rw337}
mutants. Both mRNAs are significantly up-regulated in *banp*^{rw337} mutants at 48 hpf.
Unpaired t-test (two-tailed) [n=3, p*< 0.05].

(G) TUNEL of wild-type and *banp*^{rw337} mutant retinas injected with STD MO and tp53
MO at 48 hpf and 72 hpf. Scale bars: 20 μm.

(H) Histogram of the number of TUNEL-positive cells per retinal section in wild-type
and *banp*^{rw337} mutant retinas injected with STD MO and tp53 MO at 48 hpf and 72
hpf. Two-way ANOVA, Tukey's multiple comparisons test, Mean ± SD. [n=3 for
STD-MO at 48 hpf and tp53-MO at 72 hpf; n=4 for tp53-MO at 48 hpf and STD-
MO at 72 hpf, p**** < 0.0001].

Source data 1. Data for Figure 3CEFH.
Source data 2. Data for Figure 3D.

 Figure 4. Banp is required for integrity of DNA replication and DNA damage

 repair.

 (A) Quantitative RT-PCR-based validation of FL tp53 and Δ113tp53 mRNA

    expression at 48 hpf. Δ113tp53 mRNA expression is markedly higher in *banp*^{rw337}

    mutants than in wild-type siblings, whereas FL tp53 mRNA is not significantly

    different between mutants and wild-type siblings. Unpaired t-test (two-tailed),

    Mean ± SD. [n=3, p**< 0.01, ns (not significant)].

 (B) ATAC-seq- and RNA-seq-based validation of chromatin accessibility (upper panel)

    and transcription level (bottom panel) of the genomic region covering both tp53

    and Δ113tp53 transcription units. Data of wild-type and *banp*^{rw337} mutants are

    indicated in blue and red, respectively. Light blue indicates overlapping level. The

    red box in ATAC-seq shows open chromatin in *banp*^{rw337} mutants in intron 4,

    signifying differential transcription of Δ113tp53 using an internal promoter.

 (C) TUNEL (top), anti-pH3 antibody labeling (middle), and anti-γ-H2AX antibody

    labeling (bottom) of wild-type and *banp*^{rw337} mutant retinas injected with STD MO,

    tp53 MO, and Δ113tp53 MO. TUNEL data of 48 hpf wild-type sibling and

    banp^{rw337} mutant retinas injected with STD MO and tp53 MO are shared with those

    shown in Fig. 3G. tp53 MO-mediated knockdown completely rescues apoptosis in

    banp^{rw337} mutant retinas at 48 hpf. However, Δ113tp53 MO-mediated knockdown

    mildly inhibits apoptosis in *banp*^{rw337} mutant retinas. Neither tp53 MO nor

    Δ113tp53 MO effectively rescued mitotic cell accumulation and accumulation of γ-

    H2AX-positive cells (white arrowheads) in *banp*^{rw337} mutant retinas. Scale bars: 20

    μm.

 (D–F) Histogram of the number of TUNEL-positive cells (D), pH3-positive cells (E),

 and γ-H2AX-positive cells (F) per retinal section in wild-type and *banp*^{rw337} mutants

 injected with STD-MO, tp53 MO, and Δ113tp53 MO. TUNEL data of 48 hpf wild-type

 sibling and *banp*^{rw337} mutant retinas injected with STD MO and tp53 MO are shared

 with those shown in Fig. 3H. Two-way ANOVA with Tukey's multiple comparisons

 test, Mean ± SD. [n= in graph, p*< 0.05, p**< 0.01, p***< 0.001, p****< 0.0001, ns

 (not significant)].

(G) Possible role of Banp in tp53-mediated the DNA damage response pathway. DNA
replication stress activates ATR, which subsequently promotes DNA replication fork
recovery. Failure of stalled fork recovery causes DSBs, which activate ATM-dependent
DNA damage repair. ATR and ATM inhibit Mdm2-mediated tp53 degradation.
Stabilized tp53 initially promotes transcription of the Δ113tp53 isoform, which
activates cell-cycle arrest genes and promotes DNA damage repair. Chronic activation
of tp53 promotes transcription of apoptotic genes to induce apoptosis. Banp may
normally suppress DNA replication stress by promoting DNA replication fork recovery
or DNA damage repair. Banp may suppress genotoxic stress-mediated activation of
DNA replication stress and DSB formation.
Source data 1. Data for Figure 4ADEF.
Figure supplement 1. DNA damage is induced in retinal progenitor cells in *banp*
knockdown.
1400
Figure 5. Banp promotes transcription of chromosome segregation regulators,
***cenpt* and *ncapg*.**
(A) Quantitative RT-PCR of *ncapg* expression at 48 hpf. *ncapg* mRNA expression is
down-regulated in *banp^{rw337}* mutants and *banp* morphants. Unpaired t-test (two-
tailed), Mean ± SD. [n=3, p**< 0.01].
(B) ATAC-seq and RNA-seq analysis of *ncapg* mRNA transcription.
In *banp^{rw337}* mutants, chromatin access is reduced at the TSS of *ncapg* (ATAC-seq)
and transcript expression is markedly decreased (RNA seq).
(C) Quantitative RT-PCR of *cenpt* expression at 48 hpf. *cenpt* mRNA expression is
down-regulated in *banp^{rw337}* mutants and *banp* morphants. Unpaired t-test (two-
tailed), Mean ± SD. [n=3, p**< 0.01].
(D) ATAC-seq and RNA-seq analysis of *cenpt* mRNA transcription.
In *banp^{rw337}* mutants, chromatin access is reduced at the TSS of *cenpt* (ATAC-seq)
and transcript expression is markedly decreased (RNA seq).
(E) Time-lapse images of mitosis of retinal progenitor cells injected with banp 5misMO
(upper) and banp MO (lower) at 2 dpf using *Tg [h2afv: GFP; EF1α: mCherry-*
zGem] transgenic zebrafish. In retinas injected with banp 5misMO, mitosis
proceeds sequentially from prophase (t=5~10 min), metaphase (t=16 min),

anaphase (t=21 min), and telophase (t=24 min). On the other hand,
in *banp* morphant retinas, the duration of mitosis from prophase to telophase was
67 min, significantly longer than that of the control MO (24 min). Furthermore, a
prophase-like chromosome arrangement appeared abnormally (t=59) after several
metaphase-like alignments (t=36 and 46), followed by anaphase (t=64) and
telophase. Scale bar: 5μm.

(F) Violin plot of the duration of mitosis from prophase to telophase in retinas injected
with banp 5misMO and banp MO. Mitosis in *banp* morphants is significantly
longer than that of banp 5misMO injected embryos. Mann Whitney test, Mean ±
SD. [n=3, p****< 0.0001]

(G) Confocal images of metaphase cells in zebrafish retinas at 48 hpf. The mitotic
spindle is labeled with anti-α-tubulin antibody, whereas metaphase chromosomes
are labeled with anti-pH3 antibody. The mitotic spindle is correctly attached to
metaphase chromosomes in wild-type retinal cells (left panel). On the other hand,
mitotic cells in *banp^{rw337}* mutants exhibit inefficient mitotic spindle attachment and
poorly orientated metaphase chromosomes (right panels). Dotted circles indicate
the outline of metaphase cells. Scale bar: 2μm.

(H) Percentage of cells with normal and abnormal spindle attachment in wild-type
and *banp^{rw337}* mutant embryos. Two-way ANOVA with Šídák's multiple
comparisons test, Mean ± SD. [n=3, p****< 0.0001].

1439

1440 **Source data 1.** Data for Figure 5ACFH.

1441 **Figure supplement 1.** Spatio-temporal pattern of zebrafish *cenpt* and *ncapg* mRNA
1442 expression.

1443

1444

Figure 6: Banp promotes expression of genes containing Banp motifs

(A) The motif enriched in genomic regions in which chromatin accessibility is reduced
in *banp^{rw337}* mutants. This sequence shows a high similarity to the Banp motif.

(B) A schematic diagram representing the flow of RNA-sequencing combined with
ATAC-sequencing to find candidates for zebrafish Banp direct-target genes.
HOMER was used to scan for Banp motifs in downregulated candidate genes, from

RNA- and ATAC-sequencing in *banp^rw337* mutants. RNA sequencing yielded 31 candidates, whereas ATAC sequencing yielded 72, with 26 target genes consistent in both. These 26 candidate genes are the most likely to be target genes of Banp.

(C) STRING interactome analysis showing the interaction of Banp motif-containing genes downregulated in RNA sequencing. Black arrowheads specify genes related to cell-cycle regulation.

(D) Banp target candidate genes related to cell-cycle regulation. Gene ontology (GO) analysis was applied to 31 Banp motif-containing genes downregulated in RNA sequencing. 26% of these genes are engaged in the cell-cycle regulation. Others participate in a variety of non-categorized pathways.

(E) Banp motif upstream to the 5'-UTR of *ncapg*. The black box represents the 5'-UTR.

(F) Banp motif upstream to the 5'-UTR of *cenpt*. The black box represents the 5'-UTR.

(G) Banp motif upstream to the 5'-UTR of *wrnip1*. The black box represents the 5'-UTR.

Figure supplement 1. Motifs enriched in the closed and open chromatin area in *banp^rw337* mutants.

Figure supplement 2. Banp promotes transcription of a DNA replication regulator *wrnip1*.

Figure 7. A possible model of Banp functions in stalled fork recovery, DNA damage repair, and chromosome segregation during mitosis.

(A) Our study reveals two major roles of Banp in cell-cycle regulation of retinal progenitor cells. In general, DNA replication stress, DSBs, and chromosome segregation defects induce genomic instability, which causes cancer or cell death. In S phase, Banp may suppress DNA replication stress by promoting transcription of the *wrnip1*, which protects stalled replication forks and promotes replication restart after replication stress. Failure of recovery from DNA replication stress induces DSBs, which activate ATM/Chk2/tp53-mediated DNA damage response. Banp may promote DNA damage repair in concert with tp53. ATR also activates tp53 through activation of Chk1 or crosstalk with ATM. Banp may suppress genotoxic stress-mediated DNA replication stress and DSBs. In M phase, Banp is required for chromosome segregation by promoting transcription of two mitotic regulators, *cenpt*

and *ncapg*. Since a Banp motif is found near TSSs of the *cenpt*, *ncapg* and *wrnip1*

genes, it is likely that these genes are direct targets of Banp. Dotted arrows indicate

proposed regulatory pathway, which will be necessary to investigate in the future.

(B) *banp*^{rw337} mutant phenotypes along the cell-cycle progression and neurogenesis. In

banp mutants, stalled DNA replication and subsequent DSBs are accumulated,

leading to increase in γ-H2AX+ cells. These DNA damaged cells undergo apoptosis.

In M phase, mitosis is markedly prolonged due to chromosome segregation defects;

however, almost all mitoses are eventually completed, so it is less likely that M

phase defects directly link to apoptosis. It is possible that a fraction of γ-H2AX+

cells survive and enter M phase, so DNA damage is inherited by daughter cells,

which reenter the cell-cycle or differentiate into postmitotic neurons. Neuron with a

high level of γ-H2AX+ signals may undergo apoptosis. Thus, S phase defects and M

phase defects influence each other through cell-cycle progression in *banp* mutants.

Red lines indicate path of γ-H2AX+ cells.

1497

 Supplementary Figure Legends

1499

Figure 1-figure supplement 1. Retinal cell differentiation in *banp*[rw337] mutants at 4

dpf.

(A, B) ath5:EGFP expression in wild-type (A) and *rw337* mutant (B) retinas at 4 dpf.

ath5:EGFP is strongly expressed in RGCs and weakly expressed in amacrine cells and

photoreceptor precursors. In *rw337* mutants, ath5:EGFP expression in RGCs, amacrine

cells and photoreceptor precursors is detected.

(C, D) Wild-type (C) and *rw337* mutant (D) retinas labeled with anti-Pax6 antibody at 4

dpf. Pax6 is strongly expressed in amacrine cells and weakly expressed in RGCs. In

rw337 mutants, amacrine cells seem to be decreased in number.

(E, F) Wild-type (E) and *rw337* mutant (F) retinas labeled with anti-Prox1 antibody at 4

dpf. Prox1is expressed in bipolar cells and horizontal cells. In *rw337* mutants, both

bipolar-like round nuclei and horizontal-like flat nuclei are observed, but horizontal-like

cells are very few.

(G, H) Wild-type (G) and *rw337* mutant (H) retinas labeled with zpr1 antibody at 4 dpf.

zpr1 antibody labels double-cone-type photoreceptors in zebrafish. In *rw337* mutants,

only a small cluster of zpr1-positive cells is observed in the ventro-nasal retina,

suggesting that a majority of photoreceptors fail to differentiate in *rw337* mutants.

(I, J) Wild-type (I) and *rw337* mutant (J) retinas labeled with anti-GS antibody at 4 dpf.

Anti-GS antibody labels Müller cells. In *rw337* mutants, Müller cells are decreased in

number.

(K, L) Wild-type (K) and *rw337* mutant (L) retinas labeled with anti-PCNA antibody at

4 dpf. Anti-PCNA antibody labels retinal progenitor cells in the CMZ at 4 dpf. In *rw337*

mutants, retinal progenitor cells are maintained in the CMZ.

(M, N) Phalloidin labeling of wild-type (M) and *rw337* mutant (N) retinas. Phalloidin

visualizes retinal synaptic layers, the IPL and OPL. In *rw337* mutants, the IPL is

detected, but not fully formed on the ventral side. Furthermore, there is no OPL, which

is consistent with the lack of photoreceptors in *rw337* mutants. Scale bars: 20μm.

RGC, retinal ganglion cell, AC, amacrine cell; BP, bipolar cell; HC, horizontal cell; PR,

photoreceptor; MG, Müller cell; RPC, retinal progenitor cell; OPL, outer plexiform

layer; IPL, inner plexiform layer.

1530

Figure 1-figure supplement 2. Amino acid conservation of Banp protein in zebrafish, mice and humans.

Alignment of Banp protein of humans (upper, ENSP00000376902, BANP), mice (middle, ENSMUSP00000026, SMAR1) and zebrafish (bottom, ENSDARP00000023, Banp) using a multiple sequence alignment program, PRALINE (Simossis and Heringa, 2005). Amino acid conservation is indicated by scoring and color coding from the least conserved (0, blue) to the most conserved (10, red). The BEN domain is indicated.

Figure 1-figure supplement 3. *banp^{sa12976}* homozygous mutants phenocopy *rw337* mutants.

(A) Another *banp* mutant allele, *sa12976*, carries a nonsense mutation in exon 4.

(B) Comparison of phenotypes between *banp* mutant alleles. *sa12976* homozygous *banp* mutant alleles and trans-heterozygous *rw337/sa12976* mutants show similar phenotypes to *rw337*: small eyes (left, red arrowheads) and retinal phenotypes: apoptosis and no ONL (right, open red arrowheads). Scale bars: 100 μm (left panel). Scale bars: 20 μm (right panel).

Figure 1-figure supplement 4. Confirmation of specificity of banp MO and nuclear localization of zebrafish Banp protein.

(A) Experimental design to confirm the specificity of banp MO and banp 5misMO (control MO). Co-injection of MO and RNA encoding Banp(wt)-EGFP to one-cell-stage eggs enables us to examine EGFP expression in gastrula-stage embryos at 6 hpf.

(B) banp MO effectively inhibits translation of Banp(wt)-EGFP protein in zebrafish embryos at 6 hpf and the retina at 24 hpf. On the other hand, banp 5misMO did not inhibit translation of Banp(wt)-EGFP protein, suggesting that banp MO specifically inhibits Banp protein translation. The yellow dotted line indicates blastoderm cells covering yolk, in which GFP fluorescence was detected only in 5 mismatch morphant embryos. Scale bars: 20μm.

(C) Time-lapse confocal scanning of mitosis of *Tg[EF1α:mCherry-CAAX]* transgenic wild-type retinal cells expressing Banp(wt)-EGFP. EGFP fluorescence is indicated in green, whereas mCherry-CAAX fluorescence labels plasma membranes and is indicated in magenta. White arrowheads indicate time-lapse changes in a retinal progenitor cell undergoing mitosis. Banp(wt)-EGFP is localized in the nucleus in interphase, whereas it spreads throughout the whole cell during mitosis. Scale bars: 4 µm.

(D) Histogram of intensity of Banp(wt)-EGFP during cell division shown in (A). Signals decline from mitosis to cytokinesis.

(E) Expression of Banp(wt)-EGFF, EGFP-Banp(wt) and EGFP-Banp(rw337) in zebrafish retina at 24 hpf. Both N-terminal and C-terminal EGFP-tagged wild-type Banp show nuclear localization, indicating that nuclear localization is conserved in zebrafish Banp. However, EGFP-Banp(rw337) is unstable and not maintained, suggesting that EGFP-Banp(rw337) is degraded, probably by the nonsense mutation decay mechanism. Scale bar: 10µm.

Source data 1. Data for Figure 1-figure supplement 4D.

Figure 1-figure supplement 5. Overexpression of Banp(wt)-EGFP reduces cell death in *banp^{rw337}* mutants.

(A) AO-stained retinal sections of 53 hpf wild-type retinas, *banp^{rw337}* mutant retinas, and *banp^{rw337}* mutant retinas overexpressing Banp(wt)-EGFP and EGFP-Banp(rw337). White arrows indicate noise signals derived from auto-fluorescence of iridophores. Yellow arrows indicate potential iridophore-derived noise signals near the lens. Iridophore noise signals were excluded when measuring the area of AO+ dead cells.

(B) Histogram of the percentage of AO+ area per total retina area. There is no significant difference in the percentage of AO+ area between *banp^{rw337}* mutant retinas injected without and with EGFP-Banp(rw337), suggesting no rescue of apoptosis by EGFP-Banp(rw337) injection. On the other hand, the percentage of AO+ area is markedly reduced in *banp^{rw337}* mutant retinas injected with Banp(wt)-EGFP compared with non-injected control *banp^{rw337}* mutant retinas, suggesting

1591   significant rescue of apoptosis by Banp(wt)-EGFP. Tukey's multiple comparisons

1592   test [n=5, p***< 0.001, p****< 0.0001, or ns (not significant)]. Scale bars: 50 μm.

1593   **Source data 1.** Data for Figure 1-figure supplement 5B

1594

Figure 1-figure supplement 6. Spatio-temporal pattern of zebrafish *banp* mRNA

expression.

(A) Whole-mount *in situ* hybridization of zebrafish *banp* mRNA probe. *banp* mRNA is

expressed at the 4-cell stage, suggesting maternal expression. *banp* mRNA is

ubiquitously expressed during development from the sphere stage until the 21-

somite stage. mRNA expression is restricted in the brain, including the retina at 30

hpf. At 48 hpf, mRNA is prominently expressed in the brain, fins, and notochord.

At 72 hpf and later, mRNA expression appears in neuromasts of the lateral lines

(red open arrowheads). br, brain; re, retina; LL, neuromasts of the lateral line; n,

notochord. Scale bars: 200 μm

(B) Frontal sections of zebrafish head labeled with *banp* RNA probe at 24 hpf and 48

hpf. *banp* mRNA is expressed in the entire neural retina at 24 hpf and in the CMZ

at 48 hpf, suggesting expression of *banp* mRNA in retinal progenitor cells. OT,

optic tectum; CMZ, ciliary marginal zone. Scale bars: 50 μm

(C) Whole-mount *in situ* hybridization of *banp* mRNA expression in wild-type and

banp^{rw337} mutant embryos at 72 hpf. *banp* mRNA expression is absent in *banp^{rw337}*

mutants, suggesting decay of mRNA. Scale bars: 200 μm

1612

Figure 2-figure supplement 1. Apoptosis is induced in retinal progenitor cells and

newly differentiating neurons in *banp^{rw337}* mutants.

(A) BrdU labeling of wild-type and *banp^{rw337}* mutant retinas at 48 hpf.

(B) ath5:EGFP expression of wild-type and *banp^{rw337}* mutant retinas at 48 hpf.

(C) Percentage of BrdU-positive area or ath5:EGFP-positive area relative to the total

retinal area. There is no significant difference in BrdU-positive area between wild-

type and *banp^{rw337}* mutant retinas. The fraction of ath5:EGFP-positive area is

slightly, but significantly reduced in *banp^{rw337}* mutants compared with wild-type

siblings. Unpaired t test (two tailed), Mean ± SD. [n=6, p*< 0.05, ns (not

significant)].

(D) Labeling of wild-type and *banp^{rw337}* mutant retinas with anti-activated caspase 3

antibody at 48 hpf.

(E) Labeling of wild-type and *banp^{rw337}* mutant retinas with anti-pH3 antibody at 48 hpf.

(F) The number of caspase 3-positive or pH3-positive cells per section of retina. The

number of caspase 3-positive cells and pH3-positive cells is significantly higher in

banp^{rw337} mutants than in wild-type siblings. Unpaired t-test (two tailed), Mean ±

SD. [n= 3, p*< 0.05, p**< 0.01].

(G) The relative position of caspase 3-positive cells or pH3-positive cells along the

apico-basal axis of the neural retina: Positional value from 0 to 1 corresponds to the

range from apical to basal position, respectively. In wild-type retinas (upper panel),

pH3-positive cells are located in the apical region of the neural retina (peak of

positional value is 0.2), whereas only one caspase 3-positive cell is located in the

intermediate position of the neural retina (peak of positional value is 0.4). On the

other hand, in *banp^{rw337}* mutant retina, pH3-positive cells are located in the apical

region of the neural retina (peak of positional value is 0.1), whereas caspase3-

positive cells are increased in number and most of them are located in the

intermediate zone (peak of positional value is 0.3). Although pH3- and caspase3-

positive positional ranges are slightly overlapped in *banp^{rw337}* mutant retinas, these

data suggest that apoptosis occurs in S phase of retinal progenitor cells or newly

differentiating neurons migrating toward the basal region in *banp^{rw337}* mutants.

(H) Double labeling of wild-type and *banp^{rw337}* mutant retinas with ath5:EGFP

transgene and anti-caspase 3 antibody.

(I) Percentage of ath5:EGFP-positive area relative to the total retinal area (left) and

percentage of ath5:EGFP- and caspase 3-double positive cells relative to the total

number of caspase 3-positive cells (right). Mann Whitney test, Mean ± SD. [n=7,

p**< 0.01].

(J) Double labeling of wild-type and *banp^{rw337}* mutant retinas with anti-HuC/D

antibody and anti-caspase 3 antibody.

(K) Percentage of HuC/D-positive area relative to the total retinal area (left) and

percentage of HuC/D- and caspase 3-double positive cells relative to the total

| 1653 | number of caspase 3-positive cells (right). Mann Whitney test, Mean ± SD. [n=7 for |
| 1654 | wild-type, n=6 for *banp*^{*rw337*} mutants, p*< 0.05]. |

number of caspase 3-positive cells (right). Mann Whitney test, Mean ± SD. [n=7 for

wild-type, n=6 for *banp*^{rw337} mutants, p*< 0.05].

(L) A possible model showing how apoptosis is induced during retinal progenitor cell

proliferation and retinal neurogenesis in *banp*^{rw337} mutants. Apoptosis is likely

induced in interphase of retinal progenitor cells or in newly differentiating retinal

neurons.

Scale bar: 20μm for (A, B, D, E, H, J)

Source data 1. Data for Figure 2-figure supplement 1CFGIK

1661

Figure 4-figure supplement 1. DNA damage is induced in retinal progenitor cells in

***banp* knockdown.**

(A) Labeling of 48 hpf wild-type retinas injected with STD MO and banp MO with anti-

γ-H2AX antibody. γ-H2AX-positive cells are increased in *banp* morphant retinas.

Scale bars: 20μm.

(B) The number of γ-H2AX-positive cells in 48 hpf wild-type retinas injected with STD

MO and banp MO. The number of γ-H2AX-positive cells is significantly higher in

banp morphant retinas than in wild-type retinas injected with STD MO. Mann

Whitney test, Mean ± SD. [n=5, p**< 0.01].

(C) Labeling of 48 hpf wild-type and *banp*^{rw337} mutant retinas with anti-BrdU antibody

(red), anti-γ-H2AX antibody (blue), and ath5:EGFP transgene (green). Two

independent *banp*^{rw337} mutant retinas are shown. Scale bars: 20 μm.

(D) The fraction of γ-H2AX-positive cells in the total retinal area in wild-type and

banp^{rw337} mutants. Mann Whitney test, Mean ± SD. [n=6, p <**0.01].

(E) Double labeling of retinal CMZ in *banp*^{rw337} mutants with anti-BrdU antibody (red),

anti-γ-H2AX antibody (blue). Higher magnification images of square in *banp*^{rw337}

mutant retinas shown in (C).

(F) The fraction of BrdU-positive cells in total retinal cells and γ-H2AX-positive cells

in *banp*^{rw337} mutants. Mean of the difference between total retinal cells and γ-

H2AX-positive cells is shown on the right side. The fraction is significantly higher

in γ-H2AX-positive cells than in total retinal cells. Paired t-test. [n=6, p <***0.001].

Correlation plot (F') indicates significantly effective pairing [p< *0.05].

(G) Double labeling of retinal CMZ in *banp*^{rw337} mutants with the ath5:EGFP transgene

(green), anti-γ-H2AX antibody (blue). Higher magnification images of square in

banp^{rw337} mutant retinas shown in (C).

(H) The fraction of ath5:EGFP-positive cells in total retinal cells and γ-H2AX-positive

cells in *banp*^{rw337} mutants. Mean of the difference between total retinal cells and γ-

H2AX-positive cells is shown on the right side. There is no significant difference

between total retinal cells and γ-H2AX-positive cells. Paired t-test. [n=6, ns].

Correlation plot (H') indicates no significant pairing. [ns].

(I) Labeling of 48 hpf wild-type and *banp*^{rw337} mutant retinas with anti-pH3 antibody

(magenta), anti-γ-H2AX antibody (green). Top and bottom right-side panels of

banp^{rw337} mutant retinas indicate original fluorescent images and their Imaris

surface-rendered images, respectively. Scale bars: 20 μm.

(J) The fraction of pH3-positive cells in total retinal cells and γ-H2AX-positive cells in

banp^{rw337} mutants. Mean of the difference between total retinal cells and γ-H2AX-

positive cells is shown on the right side. There is no significant difference between

total retinal cells and γ-H2AX-positive cells. Paired t-test [n=10, ns]. Correlation

plot (J') indicates no significant pairing. [ns].

Source data 1. Data for Figure 4-figure supplement 1BDFHJ.

1702

Figure 5-figure supplement 1. Spatio-temporal pattern of zebrafish *cenpt* and

***ncapg* mRNA expression.**

(A) Whole-mount *in situ* hybridization of zebrafish *ncapg* mRNA. Lateral view (left

panels), ventral view (middle panels) and retinal cross sections (right panels). Black

arrowheads in left panel represent expression in the retina. Black arrowheads in

right panel represent expression in RPCs. Whole-mount image at 48 hpf shows that

ncapg mRNA expression is markedly reduced in *banp* mutant embryo.

(B) Whole-mount *in situ* hybridization of zebrafish *cenpt* mRNA. Lateral view (left

panels) ventral view (middle panels) and retinal cross sections (right panels). Black

arrowheads in left panel represent expression in retina. Black arrowheads in right

panel represent expression in RPCs. Whole-mount image at 48 hpf shows that *cenpt*

mRNA expression is markedly reduced in *banp* mutant embryo.

1715

Figure 6-figure supplement 1. Motifs enriched in closed and open chromatin areas

in *banp^rw337* mutants.

(A) Top6 motifs enriched in genomic regions of decreased (downregulated) and

increased (upregulated) accessibility in *banp^rw337* mutants. We found that the Banp

motif (TCTCGCGAGA) is the most enriched sequence in regions with reduced

chromatin accessibility in *banp^rw337* mutants (large arrowhead).

(B, C) A putative Banp motif with a conserved CGCG element upstream of the 5' UTR

of *ncapg* (B) and *cenpt* (C) in zebrafish, mice and humans.

1724

Figure 6-figure supplement 2. Banp promotes transcription of a DNA replication

regulator *wrnip1*.

(A) A putative Banp motif with a conserved CGCG element upstream of the 5' UTR of

wrnip1 gene in zebrafish and human.

(B) Whole-mount *in situ* hybridization of zebrafish *wrnip1* mRNA. Lateral view (left

panels) ventral view (middle panels) and retinal cross sections (right panels). Black

arrowheads in left panel represent expression in retina. Black arrowheads in right

panel represent expression in RPCs. Whole-mount image at 48 hpf shows that

wrnip1 mRNA expression is markedly reduced in *banp* mutant embryo.

(C) Quantitative RT-PCR of *wrnip1* mRNA expression at 48 hpf. *wrnip1* mRNA

expression is down-regulated in both *banp^rw337* mutants and *banp* morphants.

Unpaired t-test (two-tailed) [n=on graph, $p^* < 0.05$, $p^{**} < 0.01$].

(D) ATAC-seq and RNA-seq analysis of *wrnip1* mRNA transcription. In *banp^rw337*

mutants, mRNA expression is markedly decreased, whereas chromatin accessibility

is not different between wild-type and *banp^rw337* mutants.

Source data 1. Data for Figure 6-figure supplement 2C.

1741

Video File Legends

Figure 5-video 1. Time-lapse movie of mitosis of wild-type retinal progenitor cells.

Tg [h2afv: GFP; EF1α: mCherry-zGem] transgenic embryos were injected with banp

5misMO and mitosis of retinal progenitor cells was scanned with confocal LSM.

Mitosis proceeds sequentially from prophase (t=5~10 min), metaphase (t=16 min),

anaphase (t=21 min) and telophase (t=24 min). Green indicates GFP fluorescence,

representing chromatin. On the other hand, magenta indicates fluorescence from

mCherry, representing S/G2/M phase cells.

1750

Figure 5-video 2. Time-lapse movie of mitosis of *banp* morphant retinal progenitor

cells.

Tg [h2afv: GFP; EF1α: mCherry-zGem] transgenic embryos were injected with banp

MO and mitosis of retinal progenitor cells was scanned with confocal LSM. Mitosis did

not procced smoothly, and it took longer from chromosome condensation in prophase

(t=15 min) to chromosome alignment at the division plane in anaphase (t=46 min).

Aligned chromosomes were often dispersed during chromosome segregation by mitotic

spindle (t=46~64 min). Green indicates fluorescence from GFP, representing chromatin.

On the other hand, magenta indicates fluorescence from mCherry, representing S/G2/M

phase cells.

1761

1762

Figure 5-Source data 1. Data for Figure 5ACFH.

Figure 1-figure supplement 4-Source data 1. Data for Figure 1-figure supplement 4D.

Figure 1-figure supplement 5-Source data 1. Data for Figure 1-figure supplement 5B

Figure 2-figure supplement 1-Source data 1. Data for Figure 2-figure supplement

1CFGIK

Figure 4-figure supplement 1-Source data 1. Data for Figure 4-figure supplement

1BDFHJ.

Figure 6-figure supplement 2-Source data 1. Data for Figure 6-figure supplement 2C.

1803

1804

1805   **Video files**

1806   **Figure 5-video 1.** Time-lapse movie of mitosis of wild-type retinal progenitor cells.

1807   **Figure 5-video 2.** Time-lapse movie of mitosis of *banp* morphant retinal progenitor

1808   cells.

1809

1810

1811

1812

1813

1814

1815

1816

1817

1818

1819

Appendix

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
strain, strain background (Danio Rerio)	Okinawa wild type	PMID: 28196805	NA	
strain, strain background (Danio Rerio)	RIKEN wild type	PMID: 12702661	ZFIN: ZDB-GENO-070802-4	https://shigen.nig.ac.jp/zebra/
genetic reagent (Danio Rerio)	$banp^{rw337}$	This paper	NA	NA
genetic reagent (Danio Rerio)	$banp^{sa12976}$	ZIRC	ZFIN: ZDB-ALT-130411-2647	NA
genetic reagent (Danio Rerio)	$Tg[EF1\alpha:mCherry-zGem]^{oki011}$	PMID: 25260917	ZFIN: ZDB-ALT-150128-2	NA
genetic reagent (Danio Rerio)	$Tg[EF1\alpha:mCherry-CAAX]^{oki049}$	PMID: 28196805	ZFIN: ZDB-TGCONSTRCT-181026-1	NA
genetic reagent (Danio Rerio)	$Tg[h2afv:GFP]^{kca6/kca66}$	PMID: 11819118	ZFIN: ZDB-ALT-020918-4, ZDB-ALT-020717-2	NA
genetic reagent (Danio)	$Tg[ath5:EGFP]^{rw021}$	PMID: 15728672	ZFIN: ZDB-ALT-050627-2	NA

antibody	anti-Phospho Histone H3 (Ser10) (Mouse monoclonal)	Cell signaling technology	6G3, 9706	(1:500) IHC
antibody	pH3 (Rabbit polyclonal)	Sigma-Aldrich (Merk)	06-570	(1:500) IHC
antibody	anti-Caspase3 (Rabbit monoclonal)	BD Pharmingen	Clone C92-605	IHC (1:200)
antibody	anti-Pax6 (Rabbit polyclonal)	Covance	PRB-278P	IHC (1:500)
antibody	zpr1 (Mouse monoclonal)	ZIRC, Eugene, Oregon	ZFIN: ZDB-ATB-081002-43	IHC (1:100)
antibody	anti-glutamine synthetase (Mouse monoclonal)	Millipore	MAB302, clone GS-6	IHC (1:100)
antibody	anti-PCNA (Mouse monoclonal)	Sigma-Aldrich	Clone PC10,P8825	IHC (1:200)
antibody	anti-Prox1 (Rabbit polyclonal)	Gene Tex	GTX128354	IHC (1:500)
antibody	anti-α-tubulin (Mouse monoclonal)	Sigma-Aldrich	T5168, clone B512	IHC (1:1000)
antibody	anti-γ-H2AX (Rabbit polyclonal)	Gene Tex	GTX127342	IHC (1:500)

antibody	anti-BrdU (Rat monoclonal)	abcam	Ab6326	IHC (1:200)
antibody	anti-HuC/D (Mouse monoclonal)	Thermo Fisher	A-21271	IHC (1:200)
antibody	anti-β-actin (Mouse monoclonal)	Sigma-Aldrich	A5441	WB (1:5000)
antibody	anti-tp53 (Rabbit polyclonal)	Gene Tex	GTX128135	WB (1:1000)
antibody	anti-Mouse IgG, HRP-Linked Whole Ab (Sheep polyclonal)	Cyvita	NA931	WB (1:5000)
antibody	anti-Rabbit IgG, HRP-Linked Whole Ab (Donkey polyclonal)	Cyvita	NA934	WB (1:5000)
recombinant DNA reagent	pBluescript II SK(+) (plasmid)	Stratagene/Agilent Technologies	NA	*in vitro* transcription (In situ hybridization probe synthesis)
recombinant DNA reagent	pCS2 (plasmid)	PMID: 7926732	NA	*in vitro* transcription (Capped mRNA synthesis)
sequence-based reagent	Primes for polymorphic marker 70702B	This paper	PCR primers	forward: 5'-ACTTCTTATCAGGGCTGTGC-3' reverse: 5'-TCAGTCAAGAGCAGTGAGAG-3'

sequence-based reagent	Primes for polymorphic marker zC93F2D	This paper	PCR primers	forward: 5'-TGGGATCTCTTTAAGTGAGTGAG-3' reverse: 5'-TCCAACTATGTGGGTCAAACC-3'
sequence-based reagent	banp MO	This paper	Morpholino antisense oligos	5'-CCACTAAATCTTGCTCTGACATCAT-3'
sequence-based reagent	banp 5misMO	This paper	Morpholino antisense oligos	5'-CCtCaAAATgTTcCTCTcACATCAT-3'
sequence-based reagent	tp53 MO	PMID: 12477391	Morpholino antisense oligos	5'-GCGCCATTGCTTTGCAAGAATTG-3'
sequence-based reagent	Δ113 tp53 MO	PMID: 19204115	Morpholino antisense oligos	5'-GCAAGTTTTTGCCAGCTGACAGAAG-3'
sequence-based reagent	STD MO	PMID: 30322969	Morpholino antisense oligos	5'-CCTCTTACCTCAGTTACAATTTATA-3'
sequence-based reagent	ccng1	This paper	Primers for qRT-PCR	**Forward primer**: 5'-ccctggagattgaggatcag-3' **Reverse primer**: 5'cacacaaaccaggtctccaa-3'
sequence-based reagent	mdm2	PMID: 24147052	Primers for qRT-PCR	**Forward primer**: 5'-caggaggaggagaagca

				gtg-3' **Reverse primer:** 5'-agggaaaagctgtcc gactt-3'
sequence-based reagent	p21(cdkn1a)	PMID: 26908596	Primers for qRT-PCR	**Forward primer:** 5'-aagcgcaaacagaccaa cat-3' **Reverse primer:** 5'-tcagctactggccggatt t-3'
sequence-based reagent	FL tp53	PMID: 27539857	Primers for qRT-PCR	**Forward primer:** 5'-tggagaggaggtcggcaa aatcaa-3' **Reverse primer:** 5'-gactgcgggaacctgag cctaaat-3'
sequence-based reagent	Δ113 tp53	PMID: 27539857	Primers for qRT-PCR	**Forward primer:** 5'-atatcctggcgaacatttg gaggg-3' **Reverse primer:** 5'-cctcctggtcttgtaatg tcac-3'
sequence-based reagent	Puma (bbc3)	This paper	Primers for qRT-PCR	**Forward primer:** 5'-ctgaggaggaccccacac t-3' **Reverse primer:** 5'-tctccagttctgccagtg c-3'

sequence-based reagent	cenpt	This paper	Primers for qRT-PCR	**Forward primer**: 5'-tcatgaggagattgtggaagatg-3' **Reverse primer**: 5'-ggtgagctctgcgagttatt-3'
sequence-based reagent	ncapg	This paper	Primers for qRT-PCR	**Forward primer**: 5'-ctgatgtgagggagcctattt-3' **Reverse primer**: 5'-gagtctgtttggcctccatta-3'
sequence-based reagent	atm	This paper	Primers for qRT-PCR	**Forward primer**: 5'-cctcaaggctgtggagaact-3' **Reverse primer**: 5'-aggggattttctttacaccactc-3'
sequence-based reagent	atr	This paper	Primers for qRT-PCR	**Forward primer**: 5'-aggaacccaatctgccagt-3' **Reverse primer**: 5'-gatgtccagtgccagctctc-3'
sequence-based reagent	wrnip1	This paper	Primers for qRT-PCR	**Forward primer**: 5'-gtgatgtgcgagaggtgataa -3' **Reverse primer**: 5'-acgtgtcctgctgtgatttt-3'

sequence-based reagent	ef1α	PMID: 19014500	Primers for qRT-PCR	**Forward primer:** 5'-cttctcaggctgactgtgc-3' **Reverse primer:** 5'-ccgctagcattaccctcc-3'
sequence-based reagent	Genotyping primers for *banp*^{sa12976}	This paper	Primers for sequencing	**Forward primer:** 5'-TGTTGATATCCATCAGTCAG-3' **Reverse primer:** 5'-GGTGTATAAATCACATGACC-3'
sequence-based reagent	Genotyping primers for *banp*^{rw337}	This paper	PCR Primers	**forward:** 5'-CGATGTTGATATCCATCAGTCAGGCGATC-3'; **reverse primer:** 5'-GGTGCTGGTGTATAAATCACATGACCTATGGTCCTCTT-3'.
sequence-based reagent	Subcloning of banp full length cDNA for in Situ hybridization RNA probe synthesis	This paper	PCR Primers	Forward: 5'-cgaattcatgatgtcagagcaagatttag -3' Reverse: 5'-gctcgagtcaagtgcctggcatctggatc g-3'
sequence-based reagent	Subcloning of cenpt cDNA fragment for in Situ hybridization RNA probe synthesis	This paper	PCR Primers	Forward: 5'-ctggctcaaagagtgggctga-3' Reverse: 5'-

				agacgtcactggccaccttg-3'
sequence-based reagent	Subcloning of ncapg cDNA fragment for in Situ hybridization RNA probe synthesis	This paper	PCR Primers	Forward: 5'-gtcaaggaacagcgtatagag-3' Reverse: 5'-ggaaccatgatctccgattag-3'
sequence-based reagent	Subcloning of wrnip1 cDNA fragment for in Situ hybridization RNA probe synthesis	This paper	PCR Primers	Forward: 5'-aactgatcggagaacaaactc-3' Reverse: 5'-gcacactgggctagaataac-3'
commercial assay or kit	DIG RNA Labelling Kit	Roche	11175025 910	In situ hybridization
commercial assay or kit	mMESSAGE mMACHINE™ SP6 Transcription Kit	Invitrogen	AM1340	Capped mRNA synthesis
commercial assay or kit	In Situ Cell Death Detection Kit, TMR red	Roche	1215679291 0	apoptosis
commercial assay or kit	ReverTra Ace® aPCR master mix with gDNA remover	Toyobo	FSQ-301	cDNA synthesis
commercial assay or kit	Luna® Universal qPCR Master Mix	NEB	M3003L	qRT-PCR
commercial assay or kit	Direct-zol RNA Miniprep Kit	ZYMO RESEARCH	R2050	RNA-sequencing
commercial assay or kit	NEBNext® Ultra™ II Directional RNA Library Prep Kit	NEB	E7760	Illumina RNA-sequencing

chemical compound, drug	Acridine orange	WALDECK (CHROMA)	1B-307	Live cell death detection
software, algorithm	IMARIS	Bitplane	ver.9.1.2	http://www.bitplane.com/imaris; RRID: SCR_007370
software, algorithm	Image J	NIH	Version 2.1.0/1.53c	Percentage area calculation
software, algorithm	ZEN 2012	Zeiss	LSM710 (Version: 14.0.25.201)	Percentage area calculation
Software, algorithm	GraphPad Prism	GraphPad Software	Version 9.1.2	https://www.graphpad.com/scientific-software/prism/
other	SYTOX Green	Molecular Probes	S34862	IHC (1:1000)
other	rhodamine-conjugated phalloidin	Molecular Probes	R415	Filamentous actin (F-actin) stain IHC (1: 40)
other	Restore™ PLUS Western Blot Stripping Buffer	Thermo Scientific	46430	Remove high-affinity antibodies from membranes (western blot)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Figure 7