Sulfite preservatives effects on the mouth microbiome: changes in viability, diversity and composition of microbiota.

Sally V Irwin[1][¶][*], Luz Maria Deardorff [2][&], Youping Deng[3][&], Peter Fisher[1][¶,] Michelle Gould[1][¶], Junnie June[1][¶], Rachael S. Kent[1][¶], Yujia Qin[3][&], FracescaYadao[1][&]

[1] Department of Science, Technology, Engineering and Mathematics, University of Hawai'i Maui College, Kahului, Hawai'i, United States of America

[2] Department of Natural Sciences, University of Hawai'i at Manoa, Honolulu, Hawai'i, United States of America

[3] Department of Quantitative Health Sciences, John A. Burns School of Medicine, University of Hawaii at Manoa, Honolulu, Hawai'i. United States of America

* Corresponding author:
Email: sirwin@hawaii.edu

¶These authors contributed equally to this work.
&These authors contributed equally to this work.

Short Title: Sulfite exposure and the effects on the mouth microbiome

Abstract

Processed foods make up about 70 percent of the American diet. Sulfites and other food preservatives are added to these foods largely to limit bacterial contamination. The mouth microbiota and its associated enzymes are the first to encounter food and therefore likely to be the most affected.

 Eight saliva samples from ten individuals were exposed to two sulfite preservatives, sodium sulfite and sodium bisulfite. One sample set was evaluated for bacteria composition utilizing 16s rRNA sequencing, and the number of viable cells in all sample sets was determined utilizing ATP assays at 10 and 40-minute exposure times. All untreated samples were analyzed for baseline lysozyme activity, and possible correlations between the number of viable cells and lysozyme activity.

Sequencing results indicated significant increases in alpha diversity with sodium bisulfite exposure and changes in relative abundance of 3 amplicon sequence variants (ASV). Sodium sulfite treated samples showed a significant decrease in the Firmicutes/Bacteroidetes ratio, a marginally significant change in alpha diversity, and a significant change in the relative abundance for Proteobacteria, Firmicutes, Bacteroidetes, and for 6 ASVs. Beta diversity didn't show any separation between groups, however, all but one sample set was observed to be moving in the same direction under sodium sulfite treatment in a principal component analysis.

ATP assays indicated a significant and consistent average decrease in activity ranging from 24 - 46% at both exposure times with both sulfites.

Average initial rates of lysozyme activity between all individuals ranged from +/- 76%

compared to individual variations of +/- 10 - 34%. No consistent, significant correlation

was found between ATP and lysozyme activity in any sample sets.

Conclusions: Sulfite preservatives, at concentrations regarded as safe by the

FDA, alter the relative abundance and richness of the microbiota found in saliva, and

decrease the number of viable cells, within 10 minutes of exposure.

INTRODUCTION

The human oral cavity is a complex environment hosting up to 700 different species of

bacteria, found primarily in 5 phyla, residing in saliva, teeth surfaces and on the apical

mucosa of the tongue and cheeks [1,2,3]. At birth, colonization of the mouth begins.

Streptococcus salivarius, a facultative anaerobe, is one of the initial colonizers along

with several aerobes within the 1st year, including *Lactobacillus, Actinomyces, Neisseria*

and *Veillonella* [4]. Diversity and the number of bacteria present in different individuals'

mouths may vary as a result of environmental and genetic factors including disease and

diet. However, recent studies have found most healthy individuals maintain a fairly

consistent population over weeks to months of time [2,5].

Human saliva contains several enzymes and buffers that contribute to the first steps in

digestion and serve as a first line of defense in immunological responses [6,7].

Lysozyme is found in saliva, as well as tears, blood serum, perspiration, and other

bodily fluids. It is an antimicrobial enzyme that catalyzes cleaving of ß(1,4)-glycosidic

bonds between residues of N-acetylmuramic acid (NAM) and N-acetylglucosamine

(NAG) in peptidoglycan of primarily gram positive bacterial cell walls. Research has

shown that lysozyme found in the mouth can lower the adherence of bacteria to

surfaces and limit the number of microbes [8,9].

An imbalance in the mouth microbiome can lead to oral diseases such as dental caries,

periodontitis, oral mucosal disease, and systemic diseases of the gastrointestinal,

cardiovascular, and nervous systems [1,10]. A few recent studies have detected

significant changes in the mouth microbiota in response to short and long-term

environmental alterations [11,12,13,14]. Bescos *et.al.* observed the effects of

chlorhexidine mouthwash resulting in a significant increase in the percent abundance of

Firmicutes and *Proteobacteria* and a decrease in *Bacteroidetes*, *Fusobacteria*, TM7,

and SR1 [11]. The effects of inorganic nitrates (found in vegetables) on the mouth

microbiome of 19 omnivores and 22 vegetarians, displayed an increase in the percent

abundance of *Proteobacteria* and a decrease in *Bacteroidetes* consistent with the

chlorhexidine mouthwash study [12]. Another study, examining the effects of green tea

on both the gut and mouth microbes showed "an irreversible increase of *Firmicutes* to

Bacteroidetes ratio, elevated SCFA producing genera, and reduction of bacterial LPS

synthesis in feces" [13]. A study examining betel nut chewing in 122 individuals, found

reduced bacterial diversity, elevated levels of *Streptococcus infantis*, and changes in

distinct taxa of *Actinomyces* and *Streptococcus* genera in current users and reduced

levels of *Parascardovia* and *Streptococcus* in long-term chewers [14]. At the phyla level

in both the mouth and gut, an increase in Firmicutes is often observed with obesity, and

an increase in Bacteroidetes is observed with inflammatory bowel disease [10,15,16].

Proteobacteria, another key player in the oral microbiome, has been found to be the

most variable phylum in dysbiosis and its increase is generally related to a decrease in

Firmicutes [10,16].

Antibiotics, diet, and food additives including artificial sweeteners, emulsifiers,

preservatives, colorants, and acidity regulators can result in dysbiosis in the gut leading

to a variety of human health issues [17,18,19,20]. An unhealthy gut microbiome

influences signaling in the gut-brain axis; a bidirectional signaling system responsible for

homeostasis, function, and overall health [21]. A recent study in germ-free humanized

mice found that a mixture of common antimicrobial food additives including sodium

benzoate, sodium nitrite, and potassium sorbate induced a significant dysbiosis in the

gut [22]. Sodium sulfite and sodium bisulfite have long been used as preservatives to

prevent food spoilage and browning. They can be found in a variety of foods such as

dried fruit, processed meat, beer, wine, and canned goods [23]. Many foods contain

such high levels of sulfites that exposure above levels generally regarded as safe

(GRAS) by the US Food and Drug Administration (FDA) is common [23]. Our previous

study demonstrated that under ideal growing conditions, sodium sulfite and sodium

bisulfite at concentrations lower than GRAS, were bactericidal to four probiotic species

in 2-6 hours of exposure [24]. Few studies have been conducted to determine the effect

of preservatives on the gut microbiota and none are known to us regarding their impact

on the mouth microbiome.

Most recent studies on the human mouth and gut microbiota have utilized 16s rRNA

sequencing, and in some cases flow cytometry or CFUs along with OD600

spectrophotometry to quantify cell numbers [2,5]. Twenty to sixty percent of the bacterial

cells in the mouth microbiome are estimated to be unculturable, making plate counts

and CFUs of limited use [2]. This has made 16S rRNA sequencing a powerful tool to

profile mixed communities of bacteria. However immediate or short-term changes in the

microbiome cannot reliably be detected due to the complex nature of the communities

and the detection of DNA sequences from both living and recently lysed cells [2]. Flow

cytometry is thought to be the gold standard for cell counting but also has limitations

due to the potential presence of bacterial aggregates which may artificially deflate the

cell count [2].

The amount of adenosine triphosphate (ATP) produced by living cells can be detected

by a proportional luminescent signal to observe immediate changes in the numbers of

viable bacterial cells present in a sample. Sensitivity as low as 0.0001nM ATP,

reflecting the test population, is high due to the rapid loss of ATP in non-living cells [25].

This method implements a recombinant luciferase enzyme to oxidize luciferin in the

presence of ATP and oxygen to produce oxyluciferin and light. It has been used to

accurately quantify changes in the number of viable cells in various antibiotic

susceptibility tests ranging from in-vitro biofilm formations by *Pseudomonas aeruginosa*

[26], multiple gram-negative bacterial species within 24 hours of treatment with

antibiotics [27], and slow growing species such as *Borrelia burgdorferi* [28]. Limitations

to this method include the detection of eukaryotic cell's ATP present in some mixed

populations, differences in amounts of ATP produced by varying cell types and inhibition

of the luminescence signal due to the media used in assays. All these factors were

considered in the experiments described here.

This study examines the effects of two types of sulfite preservatives on the human

mouth microbiome by corresponding changes in ATP activity of saliva samples

collected over a 4–5-week period from 10 individuals. Bacteria in saliva samples from

one individual were examined for responses to the sulfite treatments indicated by

changes in the percent abundance and/or diversity with 16s rRNA sequencing.

Additionally, baseline lysozyme activity in the individuals being studied was determined

and compared to the number of viable cells found in the samples.

Methods and Materials

Saliva Collection: Unstimulated saliva was collected from 10 individuals between

2-10 hours after consuming food or drink, other than water. Prior to donation, each

individual rinsed mouth 3 times with water followed by a twenty-minute waiting period

before delivering saliva via spitting into a sterile tube. Samples were placed on ice for

the duration of collection and aliquoting. Time allowed for saliva collection was kept

under 30 minutes. Collections took place twice a week with at least 2 days in between,

over 4-5 weeks. Each participant supplied 8 samples total. Collected saliva was diluted

1:10 with sterile DI water, aliquoted and stored at -80℃. Participants contributing to

saliva collections included ten individuals, 2 males (M1 and M2) and 8 females (F1-F8)

from 18-60 years of age, all in overall good health. Our protocol for utilizing human

saliva samples was reviewed and permitted by the Institutional Biosafety Committee

(IBC) at the University of Hawaii. The Internal Review Board was also consulted and it

was determined that no further review was necessary because the samples were made

anonymous immediately after collecting, and they were not considered human samples

due to the analysis performed being directed only on the bacterial cells. Aliquots of the

same samples were used in all experiments.

Sulfite Exposure to Saliva Samples: Saliva aliquots were exposed to

freshly made, 1800 ppm sodium bisulfite (Fisher Scientific) or 3780 ppm anhydrous

sodium sulfite (Fisher Scientific) both prepared in sterile water in independent

experiments. Immediately following exposure to sulfites, three aliquots of each control

and each exposed sample were centrifuged at 4,600 RPMs for 10 minutes to pellet

cells. The supernatant was discarded and the pellet brought up in an equal volume of

1X phosphate buffered saline (PBS). These samples are referred to as "time 0" (T0).

The other half of each aliquot was placed in an incubator set at 36℃ for 30 minutes

(T30), then pelleted using the same procedure described for time zero samples, brought

up in 1X PBS and stored. All samples were stored at -20℃ for short term storage of 6

weeks or less, or at -80℃ for longer storage. See supplemental materials for additional

notes on procedure.

ATP Activity in Saliva Samples: BacTiter-Glo™ Microbial Cell Viability

Assay from Promega was used to quantify relative numbers of viable bacterial cells

found in saliva samples. The Perkin Elmer Victor X3 multi-mode plate reader was used

to record luminescence of all samples. Samples were assayed in triplicate using a white

plate with clear bottom (View-Plate 96 TC from PerkinElmer). Lids were coated with

Triton X100 as described by Brewster [29] to prevent condensation.

Blank wells were filled with sterile water to cut down on cross talk, negative controls

contained 100ul of 1X PBS + 100ul BacTiter-GLO™ reagent. One hundred microliters of

each treated and control (untreated) sample (3 replicates each) were tested. All

samples from each individual were ATP assayed on the same day using the same Bac-

titer glo reagent to limit variability between luminescence readings for more accurate

comparisons of results. Samples and BacTiter-Glo™ reagent were brought to room

temperature before use. Assay consisted of a 5 second orbital (0.10mm) shake, a 5-

minute incubation at 25℃, followed by a 1 second read of luminescence (RLU).

Statistical Analysis of ATP Studies: For all statistical tests, p values <

0.05 were considered significant. Statistical errors were calculated as the standard error

(SE) or standard deviation (SD). Average total RLU (relative light units) for each

individual's eight saliva samples were assessed using raw RLU data by a two-way

ANOVA and paired T-Tests.

Sample Preparation for DNA Extraction: A total of 32 saliva samples

from individual F2 was sent to Zymo Research, Irvine, CA for 16s rRNA sequencing.

Sequenced samples included 8 sodium sulfite treated saliva samples at T0, and 8

untreated saliva samples serving as controls at T0. Eight sodium bisulfite treated saliva

samples at T0 and the respective control samples. A 10:1 RNA/DNA shield from Zymo

was added to each saliva sample to maintain samples during shipping.

DNA Extraction: DNA was extracted from saliva samples using ZymoBIOMICS®-

96 MagBead DNA Kit (Zymo Research, Irvine, CA). The ZymoBIOMICS® Microbial

Community Standard was used as a positive control. A blank extraction control was

used as the negative control.

Targeted Library Preparation: Bacterial 16S ribosomal RNA gene targeted

sequencing was performed using the *Quick*-16S™ NGS Library Prep Kit (Zymo

Research, Irvine, CA). Bacterial 16S primers amplify the V3-V4 region of the 16S rRNA

gene. The ZymoBIOMICS® Microbial Community DNA Standard was used as a positive

control. A blank library preparation was used as the negative control.

Absolute Abundance Quantification: Zymo Research indicates the

following methods were used for absolute abundance quantification: A quantitative real-

time PCR was set up with a standard curve. The standard curve was made with plasmid

DNA containing one copy of the 16S gene and one copy of the fungal ITS2 region

prepared in 10-fold serial dilutions. The primers used were the same as those used in

Targeted Library Preparation. The equation generated by the plasmid DNA standard

curve was used to calculate the number of gene copies in the reaction for each sample.

The PCR input volume was used to calculate the number of gene copies per microliter

in each DNA sample.

The number of genome copies per microliter DNA sample was calculated by dividing

the gene copy number by an assumed number of gene copies per genome. The value

used for 16S copies per genome is 4. The value used for ITS copies per genome is 200.

The amount of DNA per microliter DNA sample (DNA_ng) was calculated using an

assumed genome size of 4.64 x 10^6 bp, the genome size of *Escherichia coli*, for 16S

samples. This calculation is shown below:

Calculated Total DNA = Calculated Total Genome Copies × Assumed Genome Size

(4.64 × 10^6 bp) × Average Molecular Weight of a DNA bp (660 g/mole/bp) ÷ Avogadros

Number (6.022 x 10^{23}/mole). (Zymo Research, Irvine, CA).

Sequencing: The final library was sequenced on Illumina® MiSeq™ with a v3

reagent kit (600 cycles). The sequencing was performed with 10% PhiX spike-in.

Bioinformatics Analysis: Unique amplicon sequence variants (ASVs) were

inferred from raw reads using the DADA2 pipeline as described by Callahan et al. [30].

Chimeric sequences were also removed with the DADA2 pipeline. Taxonomy

assignment was performed using Uclust from Qiime v.1.9.1 with the Zymo Research

Database.

Firmicutes/Bacteroidetes ratio was calculated using the number of sequences belonging

to phylum Firmicutes divided by the number of sequences belonging to Bacteroidetes.

To evaluate the significance of the differences between the control and treatment

groups, paired t-tests were used for statistical analysis, and the results (with p-value <

0.05) shown in the figures. Alpha diversities were estimated using four indexes using R

package "Vegan" [31]: including species number (richness), Chao1, Shannon index,

and Inverse Simpson index. The beta diversity can be explained by the PCA (principal

component analysis) plots, which was also conducted by the R package "Vegan" using

the Bray-Curtis distance matrix at the ASV level. The relative abundances of ASVs in

different samples were transformed and the first two principal components were plotted

to show the relationships between the groups. The percentage followed in the axis

shows the portion of the total variances that can be explained by the first or the second

principal component. The differentially abundant taxa (at different phylogenetic levels)

were identified based on the paired t-test results.

Lysozyme Activity Assays: Lysozyme activity initial rates were determined

for all individuals untreated samples, using a Perkin Elmer Victor X3 multimode plate

reader and an EnzChek™ Lysozyme Assay Kit (E-22013) from ThermoFisher Scientific

at a temperature of 37°C. Black 96 well TC ViewPlates (PerkinElmer) with lids coated

with Triton X100 as described by Brewster [29] to prevent condensation. Fifty microliters

of saliva samples, 100ul of ENZ Check buffer, and 50ul of substrate was added to each

well in triplicate.

Graphing: Results were analyzed and graphed using Originlab software.

Results

276 **ATP assays:** Two-way ANOVA tests with replication using raw (RLU) data indicate

277 that there is no significant interaction between specific individuals and their reactions to

278 sulfite treatments or between individual's reactions to sulfite treatment due to exposure

279 time. However, there is a significant difference in ATP levels between individual's

280 control and treated saliva samples, with a large variation in baseline numbers of

281 bacteria present in each person's mouth and a significant difference in ATP levels

282 between sulfite treated and control samples for all individuals (Table 1, Fig 1 and

283 Supplemental information).

284

285 **Table 1: "p-value" results of two-way ANOVA tests**

	Time 0			Time 30		
	Individuals	**Treatment**	**Interaction**	**Individuals**	**Treatment**	**Interaction**
Sodium Sulfite	6.38E-37***	0.009**	0.500	2.37E-31***	0.035*	0.997
Sodium Bisulfite	1.06E-28***	0.002**	0.121	2.41E-28***	1.35E-05***	0.065

286 **Results show comparison of individual's reactions to sulfite treatments, sulfite**
287 **treatments compared to exposure time, and treated vs control samples for all saliva**
288 **samples. (* p<0.05, ** p<0.01, *** p<0.001)**

289
290
291 **Fig 1. Average change in ATP activity after sulfite treatment.** Raw data (RLU) of all

292 saliva samples (F1-F8 and M1-M2) treated with Na_2SO_3 or $NaHSO_3$ compared to

controls was used to determine the average % decrease of viable cells (based on ATP

activity) in each at time 0 (T0) and time 30 (T30). The standard error of each test group

was calculated, and evidence of observed difference was confirmed by both the paired

t-test and the two-way ANOVA (Table 1).

Baseline Lysozyme and Comparison to Number of Viable Cells (ATP):

A baseline of lysozyme activity in the collected saliva samples was established over a

five-week period. Activity was determined as a function of initial rates of ten individuals

(Fig 2, Fig 3). A one-way ANOVA indicated an overall significant difference between

individuals of $p = 5.5$ E-15. Tukey comparisons indicated in Figure 2 show each

sample's range and similarity among samples. A large variation in the average initial

rates over the five weeks of collection time between all individuals was observed,

ranging from +/- 76% compared to individual initial rate variations of +/- 34%. No

consistent correlation between viable cells (ATP) and Lysozyme activity was observed

(Fig 3).

Fig 2. Lysozyme activity in untreated saliva samples. Untreated saliva samples from

all individuals (n=10) were assayed for initial rates of lysozyme activity. The average

initial rate from replicates of 3 from each sample set/individual (n=8) were used to

determine the standard deviation and median scores.

Fig 3. Viable cells and lysozyme activity in control samples. Comparison of

averaged ATP activity and lysozyme initial rates in all 10 individual's control saliva

samples. Standard deviation of averaged sample sets indicated for both RLU and AFU.

Comparison of Individual F2 ATP, Lysozyme Activity and 16s rRNA

Sequencing: A significant decrease in viable cells based on ATP activity (RLU) in F2's samples treated with Na_2SO_3 and $NaHSO_3$ was observed, with a 40% and 30% decrease (respectively) illustrated in Figures 4A and B.

Fig 4. Cell viability and relative abundance of phyla in F2 samples before and after sulfite treatment. F2 saliva samples (n=32) were sequenced (16S rRNA) and the average percent abundance of the most predominant 5 phyla is displayed for untreated controls and sodium sulfite (A) or sodium bisulfite (B) treated samples at time 0. "p-values" were calculated using paired t-tests.

16s rRNA sequencing results: Individual "F2's" samples indicated that Bacteroidetes, Proteobacteria and Firmicutes phyla significantly varied in relative abundance with Na_2SO_3 exposure (Fig 4A, 5A, 6A). However, no significant difference in the relative abundance of the top 5 phyla were observed with $NaHSO_3$ (Fig 4B, 5B).

Fig 5. Ratio of Firmicutes/Bacteroidetes changes in F2 samples with sulfite treatments at time 0. 16S rRNA sequencing was used to determine the ratio (based on % abundance) for control and treated samples with Na_2SO_3 (A) or with $NaHSO_3$ (B) of individual F2.

At a more detailed level, significant changes to 3 genus and 6 ASV's (Fig 6A-C) were also observed in the Na_2SO_3 treated samples and 3 ASV's in the $NaHSO_3$ assays also showed a significant change in relative abundance (Fig 7).

Fig 6. Differentially abundant bacteria at different taxonomic levels in F2 sample, after sodium sulfite treatment at time 0. 16S rRNA sequencing was used to determine the % relative abundance of phyla (A), genus (B), and ASV (C) for control and treated samples with Na_2SO_3. P values were calculated using paired t-tests.

Fig 7. Differentially abundant ASVs in F2 sample after $NaHSO_3$ treatment at time 0. ASV43 shows a significant decrease while ASV116 and ASV161 increased significantly after the treatment.

A summary of the bacterial types at the genus and ASV level is presented in Table 2. Nine different anaerobic ASV's exhibited significant changes in relative abundance with sulfite treatments. Four out of the five that showed an increase in abundance were Gram negative and three out of those five were sulfite reducing bacteria (SRB) or have the ability to avoid oxidative stress. Four ASV's decreased in relative abundance with sulfite treatments. Three out of the four were Gram positive and one out of the 4 was a SRB.

Table 2A: Bacteria species/ASVs exhibiting a significant change in relative abundance after exposure to sodium sulfite

Species/ ASV#	*Haemophilus parainfluenzae* *ASV5*	*Solobacterium moorei* *ASV35*	*Alloprevotella rava* *ASV126*	*Actinomyces naeslundii* *ASV141* *ASV144*	*Leptotrichia wadei* *ASV30*
Gram Stain	Negative	Positive	Negative	Positive	Negative
Cellular Respiration	Facultatively Anaerobic	Obligate Anaerobe	Obligate Anaerobic	Anaerobic or Microaerophilic	Anaerobe

Descriptive	This type of bacteria is known for its infectious abilities. [32]	Association with Halitosis [33]	Found in mouth in plaque	A is a common bacteria found in plaque. [1]	Normal oral flora
Relative Abundance	Increased	Decreased	Increased	ASV144 Increased ASV141 Decreased	Increased
Sulfate reducing Bacteria	No but possess pathways to protect from oxidative stress [32]	No [33]	No [34]	No [1]	Yes [35]

355 **Table 2B: Bacteria species/ASVs exhibiting a significant change in**
356 **relative abundance after exposure to sodium bisulfite**

Species/ASV#	*Veillonella parvula-tobetsuensis ASV43*	*Prevotella salivae ASV116*	*Actinomyces naeslundii ASV161*
Gram Stain	Negative	Negative	Positive
Cellular Respiration	Obligate Anaerobes	Obligate Anaerobes	Anaerobic or Microaerophilic
Descriptive	This type of bacteria belongs to the phylum Firmicutes which is a common group that is found in the oral cavity. [36]	This bacteria genus belongs in the phylum Bacteroidetes, a common phylum found in a human GI tract and in periodontal abscesses [5]	This bacterium is a member of the phylum Actinobacteria and is a common bacteria found in plaque. [7]
Relative Abundance	Increased	Decreased	Decreased

Sulfate Reducing Bacteria	Yes [36]	Not known	No [7]

357
358
Alpha diversity measured by species richness and Chao 1 showed the same significant

increase after exposure to $NaHSO_3$ and a marginally significant increase upon exposure

to Na_2SO_3. (Fig 8). However, no significant differences were identified with the Shannon

or Inverse Simpson index (data not shown).

Fig. 8: Alpha diversity in F2 samples with sulfite treatment at time 0. 16S rRNA

sequencing was used to determine species richness (based on relative abundance) for

control and treated samples with Na_2SO_3 (A) and $NaHSO_3$ (B). A paired t-test showed

that species richness increased significantly after treatment with $NaHSO_3$ and was

marginally significant after treatment with Na_2SO_3.

Baseline lysozyme activity for F2 samples compared to the respective number of viable

cells (based on ATP activity) did not indicate a correlation of any type between these

two parameters (Fig S1).

Beta Diversity: The PCA plot shows consistent changes in the oral microbial

communities when comparing the control samples to the Na_2SO_3 treated samples of F2.

As shown in Figure 9, the dissimilarity (beta-diversity) within the samples was large due

to the possible micro-environment changes affected by many factors such as diet,

hormonal status, etc. However, when treated with Na_2SO_3, the changes in the microbial

community composition were towards almost the same direction, which indicates that

the treatment of Na_2SO_3 caused consistent changes to the oral microbial community,

even though their original structures were not similar.

Fig. 9 Beta diversity of F2 Na_2SO_3 treated samples. The PCA of all the ASVs

from the control and Na_2SO_3 treated F2 samples, show consistent directions of changes

in the microbial communities under the treatment of Na_2SO_3.

Discussion

Preservatives in food interact with the human microbiome first through the mouth and

then the rest of the digestive tract. Sulfites are a common preservative added to a

variety of foods and occur naturally in some fermented foods. Since being regulated as

GRAS for use in food in 1958 by the US FDA, allowable amounts and applications have

changed several times as it became clear that sulfites were causing moderate to severe

(even fatal) health effects in some individuals [20,23,37,38] Mammals oxidize sulfite to

sulfate with sulfite oxidase and some SRB use sulfate as an energy source, which may

result in the production of H_2S [39,40]. Both insufficient production of sulfite oxidase

leading to an increase in several forms of sulfite reactive compounds, and oxidative

stress from H_2S on eukaryotic cells, may occur as the result of ingestion of sulfites. A

recent study in rats showed that Na_2SO_3 ($\geqq$ 100ppm) causes death of gastric mucosal

cells through oxidative stress [41]. This same study also looked at the effects of sodium

sulfite (1mM) on lysozyme activity. They determined that Na_2SO_3 decreased the

bacteriolytic activity of lysozyme in a time and dose dependent manner [41]. Other

studies have shown a connection between certain types of SRB found in the colon and

ulcerative colitis [42]. These studies provide evidence of mechanisms by which Na_2SO_3

may lead to both eukaryotic and prokaryotic cell death.

Our previous work indicated that sodium sulfite was bactericidal to three probiotic

species of *Lactobacillus* and bacteriostatic to *Streptococcus thermophilus,* within 2-6

hours of exposure [24]. Sodium bisulfite was found to be bactericidal to all 4 probiotic

bacteria also within 2-6 hours of exposure [24]. These studies established both the

potency of these preservatives and the varying susceptibility of different types of

bacteria when exposed under ideal culture conditions. The current study was conducted

to evaluate the effects of sulfites on the mouth microbiome found in saliva to better

simulate the effects of ingesting foods containing these preservatives.

ATP Activity: Comparison of each individual's control samples indicated that the

viable number of cells were relatively consistent over the 5-week sampling time, which

supports findings from other recent studies [2,5,10]. The most striking observations

were the differences found in ATP activity (indicating viable cell counts), between

controls and sulfite treated samples at both exposure times. Based on these results we

can conclude that both sodium sulfite and sodium bisulfite, consistently and immediately

(regardless of preservative type or sample origin) reduced the number of viable

bacterial cells in saliva samples. We are assuming this is a drop in the number of viable

cells, but it could be partially due to sulfites interfering with ATP production, or ATP

being utilized by SRB to metabolize the sulfites.

These results were also supported by our preliminary experiments (data not included)

on the mouth microbiome which examined the effects of sulfites on 12 saliva samples

from 5 individuals over a three-month period. Very similar results of reduced ATP

activity as described here were observed, with an average decrease in viable cells of

27% and 28% at time 0 for Na_2SO_3 and $NaHSO_3$ respectively. Additional experiments

observed changes in the microbial population in these saliva samples by recording

OD600 readings over a 6.5-hour period. A consistent trend of a decrease in cells

immediately after both sulfite treatments was found, compared to an increase in cell

numbers in controls.

Lysozyme: Lysozyme activity did not vary significantly over the 5 weeks of sample

collection/individual and there did not appear to be any correlation between the number

of viable cells and lysozyme activity (Fig 3). If the population could be measured

accurately to show the number of gram-negative vs. gram-positive bacteria present, a

relationship might be found due to the greater impact of lysozyme on gram-positive

cells. Significant differences in lysozyme activity were observed *between* individuals

(Fig 2). Previous studies have corroborated our results of *inter* versus *intra*-individual

lysozyme variability [43] and others have shown that lysozyme varies between

individuals with respect to circadian rhythm [44].

Preliminary studies on the effects of Na_2SO_3 on lysozyme activity in saliva (S-lys) and

on human recombinant lysozyme (HR-Lys) indicated a decreased initial rate profile of

HR-Lys and to a lesser degree S-lys with sodium sulfite incubation (data not included).

We hypothesize that this difference may be attributed to the rich mixture of biological

components that compose salivary fluid, where sulfite may act on bacteria as well as

nucleophiles and nonspecific inhibitors [45]. These early studies suggest that sodium

sulfite would affect lysozyme function in human saliva and subsequently alter the mouth

microbiome. However, we have yet to determine conclusively that sulfite inhibits salivary

lysozyme activity and further studies are in progress. More studies are needed to look at

the relationship between lysozyme activity, sulfites, and the makeup of the mouth

microbiome.

16s rRNA Sequencing: This study identified five major bacterial phyla (Fig 4) as

the most abundant in all control and treated samples. These five phyla, *Actinobacteria,*

Bacteroidetes, Firmicutes, Fusobacteria, and *Proteobacteria,* are considered the "core

microbiome" in the human mouth since they have been consistently reported as the

main bacteria found in both healthy and diseased conditions [1,3,46,16]. However,

significant changes in the saliva microbiomes were observed, dependent on the type of

sulfite exposure. Alterations of the *Firmicutes/Bacteroidetes* ratio, which is widely

accepted to have an important influence in maintaining normal intestinal and mouth

homeostasis [1,3,16], was observed with Na_2SO_3 but not with $NaHSO_3$ (Fig 5). A more

shifted microbial community structure (relative abundance), but only a slight increase in

the species richness ($p < 0.10$) was observed in the Na_2SO_3 treated samples. However,

$NaHSO_3$ samples showed a much higher alpha-diversity ($p < 0.05$) despite showing far

fewer changes in relative abundance (Fig 6-8). These results contrast with the similar

results found with both sulfites in the combined ATP assays from all individuals (Fig 1).

As for beta diversity, the PCA plot (Fig 9) shows consistent changes in the oral

microbial communities when comparing the control samples to the Na_2SO_3 treated

samples, while no similar pattern was found in the $NaHSO_3$ treated samples. These

findings, along with our previous studies [24] seem to indicate a difference in the

selective potency of these two types of sulfites. One possible explanation for the

increase in alpha diversity with both sulfites might be due to the toxic effect on certain

predominant bacterial species (thereby lowering their relative abundance) that may

have then revealed the presence of other species normally present in populations too

low to detect [47].

A comparison of the sequencing data to the ATP assays (Fig 4 A,B) from individual

"F2" samples, indicates a significant decrease in cell numbers with both sulfite

treatments. However, a greater change was observed with Na_2SO_3 vs. $NaHSO_3$. It is

interesting to note that a decrease in [DNA] of treated samples from individual "F2" was

observed, with an average decrease of 18% (p = 0.34) in sodium bisulfite and 56% (p =

0.05) in the sodium sulfite treated samples (S3 appendix). While there are many factors

that may lead to an observed difference in [DNA], this data supports our observation of

a 30% and 40% average decrease in viable cells in respective samples.

It is well established that DNA from both, recently lysed, as well as living cells may be

counted in 16S rRNA gene targeted sequencing [2]. The difference in observed

changes (dependent on sulfite type), between out observed sequencing and ATP

activity may indicate the quantitative change threshold of viable cells needed in order to

detect significance, in this method of sequencing, effectively [47].

Conclusions: Our results show that sulfites have a clear and significant impact on

some bacterium types found in the mouth. We hypothesize that sulfite

susceptibility/metabolism differences allow some bacteria found in saliva to increase in

numbers while others decrease. Bacteria that produce sulfur nucleotide reductase

enzymes (SRB) or have other mechanisms to avoid oxidative stress, are better able to

survive, while those that don't will be more susceptible to sulfites toxic effects.

Susceptibility of different types of bacteria to lysozyme and the effects of sulfites on

lysozyme activity may also be a factor.

The connections of H_2S production from some SRB and its impact on both oral and

intestinal inflammation and disease has been previously reported [39,40,42]. To

summarize, sulfite food preservatives appear to be affecting the makeup of the mouth

microbiome by more than one mechanism including death by oxidative stress to non-

SRB bacterial types and an energy source for SRB bacterial types.

Studies on the microbiomes found in or on the human body are complex and in constant

flux due to the environment. The use of more than one cell detection method to

measure changes in mixed populations of bacteria existing in the mouth is

recommended for more accurate and sensitive assessments. ATP tests used to detect

changes in the number of viable cells in saliva samples after exposure to either sulfite

revealed a significant decrease in cells in all samples, whereas changes identified

through 16s rRNA sequencing were less consistent between the samples depending on

the type of sulfite tested. Future studies are indicated to include more individual's

samples for 16s rRNA sequencing to compare to ATP assays, and to examine

connections between diseases of the digestive process and the intake of sulfite

preservatives.

Acknowledgements

519 **References**

[1] Gao, L., Xu, T., Huang, G., Jiang, S., Gu, Y., & Chen, F. (2018). Oral microbiomes:

more and more importance in oral cavity and whole body. Protein & Cell, 9(5), 488–500.

https://doi.org/10.1007/s13238-018-0548-1 Protein Cell [Internet]. 2018 [cited 2019 Apr

17];9(5):488–500. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29736705

[2] Marotz, C., Morton, J. T., Navarro, P., Coker, J., Belda-Ferre, P., Knight, R., &

Zengler, K. (2021). Quantifying Live Microbial Load in Human Saliva Samples over

Time Reveals Stable Composition and Dynamic Load. *MSystems*, *6*(1).

https://doi.org/10.1128/MSYSTEMS.01182-20

[3] Verma, D., Garg, P. K., & Dubey, A. K. (2018). Insights into the human oral

microbiome. *Archives of Microbiology 2018 200:4*, *200*(4), 525–540.

https://doi.org/10.1007/S00203-018-1505-3

[4] Deo PN, Deshmukh R. Oral microbiome: Unveiling the fundamentals [Internet]. Vol.

23, Journal of Oral and Maxillofacial Pathology. Wolters Kluwer Medknow Publications;

2019 [cited 2021 Apr 22]. p. 122–8. Available from: /pmc/articles/PMC6503789/

[5] Belstrøm D, Holmstrup P, Bardow A, Kokaras A, Fiehn N-E, Paster BJ. Temporal

Stability of the Salivary Microbiota in Oral Health. Burne RA, editor. PLoS One

[Internet]. 2016 Jan 22 [cited 2019 Apr 15];11(1):e0147472. Available from:

https://dx.plos.org/10.1371/journal.pone.0147472

[6] Mandel AL, Peyrot des Gachons C, Plank KL, Alarcon S, Breslin PA. Individual

differences in AMY1 gene copy number, salivary α-amylase levels, and the perception

of oral starch. PLoS One. 2010 Oct 13;5(10):e13352. doi:

10.1371/journal.pone.0013352. https://pubmed.ncbi.nlm.nih.gov/20967220/b

[7] Marsh, P. D., Do, T., Beighton, D., & Devine, D. A. (2016). Influence of saliva on the

oral microbiota. *Periodontology 2000*, *70*(1), 80–92. https://doi.org/10.1111/PRD.12098

[8] Lynge Pedersen AM, Belstrøm D. The role of natural salivary defenses in

maintaining a healthy oral microbiota. J Dent [Internet]. 2019;80(August 2018):S3–12.

Available from: https://doi.org/10.1016/j.jdent.2018.08.010

[9] Ragland SA, Criss AK. From bacterial killing to immune modulation: Recent insights

into the functions of lysozyme. Vol. 13, PLoS Pathogens. Public Library of Science;

2017. Available from https://doi.org/10.1371/journal.ppat.1006512

[10] Acharya, A., Chan, Y., Kheur, S., Jin, L. J., Watt, R. M., & Mattheos, N. (2017).

Salivary microbiome in non-oral disease: A summary of evidence and commentary. In

Archives of Oral Biology (Vol. 83, pp. 169–173). Elsevier Ltd.

https://doi.org/10.1016/j.archoralbio.2017.07.019

[11] Bescos R, Ashworth A, Cutler C, Brookes ZL, Belfield L, Rodiles A, et al. Effects of Chlorhexidine mouthwash on the oral microbiome. Sci Rep. 2020;10(1):1–8. Available from: https://doi.org/10.1038/s41598-020-61912-4

[12] Ashworth A, Cutler C, Farnham G, Liddle L, Burleigh M, Rodiles A, et al. Dietary intake of inorganic nitrate in vegetarians and omnivores and its impact on blood pressure, resting metabolic rate and the oral microbiome. Free Radic Biol Med [Internet]. 2019;138(May):63–72. Available from: https://doi.org/10.1016/j.freeradbiomed.2019.05.010

[13] Yuan X, Long Y, Ji Z, Gao J, Fu T, Yan M, et al. Green Tea Liquid Consumption Alters the Human Intestinal and Oral Microbiome. Mol Nutr Food Res. 2018;62(12):1–15. Available from: https://doi.org/10.1002/mnfr.201800178

[14] Hernandez, B. Y., Zhu, X., Goodman, M. T., Gatewood, R., Mendiola, P., Quinata, K., & Paulino, Y. C. (2017). Betel nut chewing, oral premalignant lesions, and the oral microbiome. *PLoS ONE*, *12*(2). https://doi.org/10.1371/journal.pone.0172196

[15] Mok SF, Karuthan C, Cheah YK, Ngeow WC, Rosnah Z, Yap SF, Ong HKA. The oral microbiome community variations associated with normal, potentially malignant disorders and malignant lesions of the oral cavity. Malays J Pathol. 2017 Apr;39(1):1-15. PMID: 28413200.

[16] Stojanov S, Berlec A, Štrukelj B. The influence of probiotics on the

firmicutes/bacteroidetes ratio in the treatment of obesity and inflammatory bowel

disease. Microorganisms. 2020;8(11):1–16.

[17] Cao Y, Liu H, Qin N, Ren X, Zhu B, Xia X. Impact of food additives on the

composition and function of gut microbiota: A review. Trends Food Sci Technol

[Internet]. 2020;99(March):295–310. Available from:

https://doi.org/10.1016/j.tifs.2020.03.006

[18] Sugihara K, Morhardt TL, Kamada N. The Role of Dietary Nutrients in Inflammatory

Bowel Disease. Front Immunol [Internet]. 2019 Jan 15 [cited 2019 Apr 15];9. Available

from: https://www.frontiersin.org/article/10.3389/fimmu.2018.03183/full

[19] Langdon A, Crook N, Dantas G. The effects of antibiotics on the microbiome

throughout development and alternative approaches for therapeutic modulation.

Genome Med [Internet]. 2016;8(1):39. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4831151&tool=pmcentrez&re

ndertype=abstract

[20] Vally H., Misso N. Adverse reactions to the sulphite additives. *Gastroenterol*

Hepatol Bed Bench. 2012 5(9), 16–23. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4017440/

[21] Butler, M. I., Cryan, J. F., & Dinan, T. G. (2019). Man and the Microbiome: A New

Theory of Everything? *Annual Review of Clinical Psychology*, *15*(1), 371–398.

https://doi.org/10.1146/annurev-clinpsy-050718-095432

[22] Hrncirova L, Machova V, Trckova E, Krejsek J, Hrncir T. Food preservatives

induce proteobacteria dysbiosis in human-microbiota associated Nod2-deficient mice.

Microorganisms. 2019;7(10). Available from:

https://doi.org/10.3390/microorganisms7100383

[23] Leclercq C., Le Donne C., and Dr M.C.F. Toledo M.C.F., Sulfites: assessment of

dietary exposure. *WHO Food Addit Ser IPCS—*(*SAFETY evaluation of certain food*

additives) 2009; 60: 221–244.Available from:

http://apps.who.int/iris/bitstream/10665/44063/1/9789241660600_eng.pdf

[24] Irwin SV, Fisher P, Graham E, Malek A, Robidoux A. Sulfites inhibit the growth of

four species of beneficial gut bacteria at concentrations regarded as safe for food. PLoS

One. 2017 Oct 18;12(10):e0186629. doi: 10.1371/journal.pone.0186629. PMID:

29045472; PMCID: PMC5646858.

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0186629

[25] Hammes F, Goldschmidt F, Vital M, Wang Y, Egli T. Measurement and

interpretation of microbial adenosine tri-phosphate (ATP) in aquatic environments.

Water Res [Internet]. 2010;44(13):3915–23. Available from:

http://dx.doi.org/10.1016/j.watres.2010.04.015

[26] Maiden MM, Zachos MP, Waters CM. The ionophore oxyclozanide enhances

tobramycin killing of Pseudomonas aeruginosa biofilms by permeabilizing cells and

depolarizing the membrane potential. J Antimicrob Chemother [Internet].

2019;74(4):894–906. Available from:

https://academic.oup.com/jac/article/74/4/894/5280826

[27] Cai Y, Leck H, Lim TP, Teo J, Lee W, Hsu LY, et al. Using an adenosine

triphosphate bioluminescent assay to determine effective antibiotic combinations

against carbapenem-resistant gram negative bacteria within 24 hours. PLoS One

[Internet]. 2015;10(10):1–15. Available from:

http://dx.doi.org/10.1371/journal.pone.0140446

[28] Pothineni VR, Wagh D, Babar MM, Inayathullah M, Solow-Cordero D, Kim KM, et

al. Identification of new drug candidates against Borrelia burgdorferi using high-

throughput screening. Drug Des Devel Ther. 2016 Apr 1;10:1307–22. Available from:

https://doi.org/10.2147/DDDT.S101486

[29] Brewster JD. A simple micro-growth assay for enumerating bacteria. J Microbiol

Methods. 2003;53(1):77–86. Available from: https://doi.org/10.1016/S0167-

7012(02)00226-9

[30] Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes S, (2016)

DADA2: High resolution sample inference from Illumina amplicon data. Nat Methods

13(7):581-3. Available from https://doi.org/10.1038/nmeth.3869

[31] Dixon, P. (2003). Computer program review VEGAN, a package of R functions for

community ecology. In *Journal of Vegetation Science* (Vol. 14). Opulus Press Uppsala.

http://cran.r-project.org

[32] Harrison A, Bakaletz LO, Munson RS. Haemophilus influenzae and oxidative

stress. Front Cell Infect Microbiol. 2012;2(March):40.

[33] Tanabe SI, Grenier D. Characterization of volatile sulfur compound production by

Solobacterium moorei. Arch Oral Biol. 2012 Dec;57(12):1639–43.

[34] Downes J, Dewhirst FE, R Tanner AC, Wade WG, William Wade CG. IUMS Printed

in Great Britain. Int J Syst Evol Microbiol [Internet]. 2013;63:1214–8. Available from:

www.homd.org/taxon=302

[35] Ren W, Xun Z, Wang Z, Zhang Q, Liu X, Zheng H, et al. Tongue Coating and the

Salivary Microbial Communities Vary in Children with Halitosis. Nat Publ Gr [Internet].

2016; Available from: www.nature.com/scientificreports

[36] Washio, J., Shimada, Y., Yamada, M., Sakamaki, R., & Takahashi, N. (2014).

Effects of pH and lactate on hydrogen sulfide production by oral Veillonella spp. *Applied*

and environmental microbiology, *80*(14), 4184–4188.

https://doi.org/10.1128/AEM.00606-14

[37] Nair B, Elmore AR. Final report on the safety assessment of sodium sulfite,

potassium sulfite, ammonium sulfite, sodium bisulfite, ammonium bisulfite, sodium

metabisulfite and potassium metabisulfite. *Int J Toxicol*. 2003;22 Suppl 2:63–88.

Available from https://www.ncbi.nlm.nih.gov/pubmed/14555420 [Accessed 6 February

2017].

[38] Seetaramaiah K, Smith AA, Murali R, Manavalan R. Preservatives in Food

Products—Review. *Int J Pharm Biol Arch*. 2011;2(2):583–99. Available from:

http://www.ijpba.info/ijpba/index.php/ijpba/article/viewFile/236/

[39] Kushkevych I, Coufalová M, Vítězová M, Rittmann SK-MR. Sulfate-Reducing

Bacteria of the Oral Cavity and Their Relation with Periodontitis—Recent Advances. J

Clin Med. 2020 Jul 23;9(8):2347. Available from: http://dx.doi.org/10.3390/jcm9082347

[40] Kushkevych I, Cejnar J, Treml J, Dordević D, Kollar P, Vítězová M. Recent

Advances in Metabolic Pathways of Sulfate Reduction in Intestinal Bacteria. Vol. 9,

Cells. NLM (Medline); 2020. Available from: http://dx.doi.org/10.3390/cells9030698

[41] Oshimo, M., Nakashima, F., Kai, K., Matsui, H., Shibata, T., & Akagawa, M. (2021).

Sodium sulfite causes gastric mucosal cell death by inducing oxidative stress. *Free*

Radical Research. https://doi.org/10.1080/10715762.2021.1937620

[42] Coutinho CMLM, Coutinho-Silva R, Zinkevich V, Pearce CB, Ojcius DM, Beech I.

Sulphate-reducing bacteria from ulcerative colitis patients induce apoptosis of

gastrointestinal epithelial cells. Microb Pathog. 2017;112:126–34.

[43] Neyraud E, Palicki O, Schwartz C, Nicklaus S, Feron G. Variability of human saliva

composition: possible relationships with fat perception and liking. Arch Oral Biol. 2012

May;57(5):556-66. doi: 10.1016/j.archoralbio.2011.09.016. Epub 2011 Oct 22. PMID:

22024405.

[44] Richter J, Král V, Zukov I, Subrt P, Rahm J. Circadian changes of the SIgA,

lysozyme, albumin and copper content of saliva. Czech Med. 1980;3(4):249-54. PMID:

7227123.

[45] Neta P, Huie RE. Free-radical chemistry of sulfite. Environ Health Perspect

[Internet]. 1985;VOL. 64:209–17. Available from:

https://www.researchgate.net/publication/19330636_Free-Radical_Chemistry_of_Sulfite

[46] Pushalkar, S., Ji, X., Li, Y., Estilo, C., Yegnanarayana, R., Singh, B., Li, X., &

Saxena, D. (2012). Comparison of oral microbiota in tumor and non-tumor tissues of

patients with oral squamous cell carcinoma. *BMC Microbiology*, *12*(144).

https://doi.org/10.1186/1471-2180-12-144

[47] Caruso V, Song X, Asquith M, Karstens L. Performance of Microbiome Sequence

Inference Methods in Environments with Varying Biomass. mSystems [Internet]. 2019

Feb 26 [cited 2021 Sep 2];4(1). Available from:

https://doi.org/10.1128/mSystems.00163-18.

Click here to access/download
Supporting Information
Supporting Information.docx

Supporting Appendix Figures data

Click here to access/download
Supporting Information
S1 appendix.xlsx

Click here to access/download
Supporting Information
S2 appendix.xlsx

Supporting Appendix DNA concentration data

Click here to access/download
Supporting Information
S3 appendix.xlsx