Thymic macrophages consist of two populations with distinct

localization and origin

Tyng-An Zhou[1], Hsuan-Po Hsu[1], Yueh-Hua Tu[2,3], Hui-Kuei Cheng[1], Chih-Yu

Lin[1], Nien-Jung Chen[1], Jin-Wu Tsai[4], Ellen A. Robey[5], Hsuan-Cheng

Huang[2,6], Chia-Lin Hsu[1], and Ivan L. Dzhagalov[1*]

[1]Institute of Microbiology and Immunology, National Yang Ming Chiao Tung University, Taipei,

Taiwan

[2]Bioinformatics Program, Taiwan International Graduate Program, Institute of Information

Science, Academia Sinica, Taipei, Taiwan

[3]Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University,

Taipei, Taiwan

[4]Brain Research Center, National Yang Ming Chiao Tung University, Taipei, Taiwan

[5]Division of Immunology and Pathogenesis, Department of Molecular and Cell Biology,

University of California, Berkeley, CA, United States

[6]Institute of Biomedical Informatics, National Yang Ming Chiao Tung University, Taipei,

Taiwan

*Correspondence to Ivan L. Dzhagalov: ivan.dzhagalov@nycu.edu.tw, ORCID: 0000-0001-

9209-4582

Mailing Address: Ivan L. Dzhagalov, Institute of Microbiology and Immunology, National Yang

Ming Chiao Tung University, #155, Sec. 2, LiNong Street, Beitou, Taipei City, 11221,

TAIWAN

Abstract

Tissue-resident macrophages are essential to protect from pathogen invasion and maintain organ homeostasis. The ability of thymic macrophages to engulf apoptotic thymocytes is well appreciated, but little is known about their ontogeny, maintenance, and diversity. Here, we characterized the surface phenotype and transcriptional profile of these cells and defined their expression signature. Thymic macrophages were most closely related to spleen red pulp macrophages and Kupffer cells and shared the expression of the transcription factor *SpiC* with these cells. Single-cell RNA sequencing showed that the macrophages in the adult thymus are composed of two populations distinguished by the expression of *Timd4* and *Cx3cr1*. Remarkably, *Timd4*$^+$ cells were located in the cortex, while *Cx3cr1*$^+$ macrophages were restricted to the medulla and the cortico-medullary junction. Using shield chimeras, transplantation of embryonic thymuses, and genetic fate mapping, we found that the two populations have distinct origins. *Timd4*$^+$ thymic macrophages are of embryonic origin, while *Cx3cr1*$^+$ macrophages are derived from adult hematopoietic stem cells. Aging has a profound effect on the macrophages in the thymus. *Timd4*$^+$ cells underwent gradual attrition, while *Cx3cr1*$^+$ cells slowly accumulated with age and, in older mice, were the dominant macrophage population in the thymus. Altogether, our work defines the phenotype, origin, and diversity of thymic macrophages.

Introduction

Tissue-resident macrophages are present in every organ and maintain local homeostasis through diverse functions ranging from protection against pathogens to tissue repair[1]. To perform their roles efficiently, macrophages acquire specialized phenotypes depending on the tissue microenvironment, and as a consequence, multiple subtypes exist, frequently within the same organ. For example, the spleen harbors red pulp macrophages specialized in red blood cell phagocytosis, marginal zone macrophages and metallophilic macrophages that are the first defense against blood-borne pathogens, T cell zone macrophages that silently dispose of apoptotic immune cells, and tingible-body macrophages that engulf less fit B cells in the germinal center[2-4]. Thus, tissue-resident macrophages represent a fascinating developmental system that allows enormous plasticity.

The last decade has seen a paradigm shift in our understanding of the development of tissue-resident macrophages. Contrary to the long-held belief that all macrophages derive from circulating monocytes[5], multiple studies have shown that many of them are long-lived cells with an embryonic origin that can maintain themselves in the tissues (reviewed in[6]). Three waves of distinct progenitors settle the tissues and contribute in various degrees to the resident macrophages in each organ. The first wave consists of the yolk sac (YS)-derived primitive macrophages that enter all tissues and establish the earliest macrophage populations[7,8]. In all organs, except for the brain and, partially, the epidermis, primitive macrophages are replaced by the next wave consisting of fetal monocytes[9-12]. The third wave comes from hematopoietic stem cells (HSCs)-derived monocytes that contribute to various degrees to

the macrophage pool in different tissues. For example, these cells contribute

little to the microglia in the brain, Langerhans cells in the epidermis, and

alveolar macrophages in the lungs but substantially to most other organs[13-

16]. Moreover, the kinetics and timing of HSC-derived monocyte infiltration

vary in different parts of the body. For some macrophage populations, such

as the arterial macrophages and subcapsular lymph node macrophages,

monocytes replace embryonic macrophages soon after birth and self-maintain

after that with little contribution from circulating cells[17,18]. Others, such as

heart macrophages, osteoclasts, and pancreatic islets macrophages, are

progressively replaced at a low rate[14,19-23]. A third group, such as the

macrophages in the dermis and most of the gut macrophages, are constantly

replaced by blood monocytes with relatively fast kinetics[24,25]. These

conclusions have been extended to many different macrophage populations

such as Kupffer cells, liver capsular macrophages, red pulp macrophages,

testicular macrophages, large and small peritoneal macrophages, and T zone

macrophages in the lymph nodes[2,13,14,16,26-30].

The recent revitalization in macrophage research has yet to reach thymic

macrophages. Although their prodigious phagocytic ability is well

appreciated[31], little is known about the origin, diversity, and maintenance of

these cells. This gap in our knowledge is, partly, due to the lack of a

consensus about the surface phenotype of thymic macrophages.   Various

groups have used different markers to identify these cells, such as F4/80 and

Mac-3 (LAMP-2)[31], or CD4 and CD11b[32], or Mac-2 (galectin 3), F4/80 and

ED-1 (CD68)[33]. Most commonly, researchers employ F4/80 and CD11b[34-

37]. However, none of these markers is macrophage-specific: F4/80 is also

expressed on eosinophils and monocytes[38,39], while CD11b is present on

most myeloid cells. The lack of a clear phenotypic definition of thymic

macrophages has translated into the absence of models that target genes

specifically in this population. For example, although macrophages in various

organs have been successfully targeted with $Lyz2^{Cre}$, $Csf1r^{Cre}$, or $Cx3cr1^{Cre}$,

very few studies have used these models in the thymus[37,40,41].

Only a handful of studies have explored the origin of thymic macrophages.

Several reports have indicated that these cells could be derived from T cell

progenitors in the thymus based on an improved single cell in vitro culture and

in vivo transplantation experiments[42,43]. However, these conclusions have

been questioned based on fate-mapping experiments using $Il7r^{Cre}$ that found

very limited contribution of lymphoid progenitors to thymic macrophages in

vivo in unperturbed mice[44]. Most recently, Tacke et al. used parabiosis to

rule out circulating monocytes as a major source of thymic macrophages[37].

The same study also performed fate-mapping experiments to show that most

thymic macrophages descend from $Flt3^+$ HSC-derived progenitors. However,

the contribution of earlier waves of hematopoiesis has not been explored.

Here, we aimed to bring our knowledge of thymic macrophages on par with

other tissue-resident macrophages. We started by clearly defining thymic

macrophages according to the guidelines set by the Immunological Genome

Consortium (IMMGEN)[38] and characterized their surface phenotype and

transcriptional signature. Using single-cell RNA sequencing (scRNA-Seq), we

identified two populations of thymic macrophages with distinct localization. We

explored the origin of these cells through genetic fate mapping, shield

chimeras, and embryonic thymus transplantation and documented that

different waves of progenitors give rise to the two populations of thymic

macrophages. Altogether our work fills an important gap in our understanding

of resident thymic macrophages and provides the framework for future

functional characterization of these cells.

Results:

CD64, F4/80, and MerTK identify the macrophages in the thymus

To unambiguously and comprehensively identify macrophages in the

thymus, we evaluated several of the prototypical macrophage markers –

MerTK, CD64, and F4/80[38]. A population that was stained with all three

markers (Figure 1A). As staining with MerTK and F4/80 was relatively dim

even when the brightest fluorochromes (e.g., PE) were used and could not be

resolved fully from the isotype control (Figure 1 – figure supplement 1), we

chose to use CD64 vs. forward scatter (FSC) as the first step in our gating

strategy (Figure 1B). Among $CD64^{+}FSC^{hi}$ cells, $F4/80^{+}CD11b^{lo}$ macrophages

could be distinguished from $F4/80^{lo}CD11b^{+}$ monocytes.

The $CD64^{+}F4/80^{+}MerTK^{+}CD11b^{lo}FSC^{hi}$ cells had typical macrophage

morphology with abundant cytoplasm (Figure 1C). These cells did not

express lineage markers characteristic of T cells (CD3ε), B cells (CD19),

eosinophils (Siglec F), NK cells (NK1.1), neutrophils (Gr1), or plasmacytoid

dendritic cells (Siglec H) (Figure 1D). However, they expressed phagocytic

receptors such as TIM4, CD51, and Axl (Figure 1E). Immunofluorescent

staining with CD64, MerTK, and TIM4 in the thymic cortex confirmed the

presence of large cells positive for all three macrophage markers (Figure 1 –

figure supplement 2).

To ascertain that our gating strategy identifies all macrophages in the

thymus in an unbiased way, we performed scRNA-Seq of sorted $Csf1r^{GFP+}$

and $Cd11c^{YFP+}$ thymic cells. $Csf1r$ is required for the survival of most

macrophages and is considered their definitive marker[45,46], while $Cd11c^{YFP}$

is expressed in many myeloid cells, including macrophages[47]. Both

reporters identified an overlapping set of cells (Figure 1 – figure supplement

3). Two clusters expressed the macrophage/monocytes-specific transcription

factor $Mafb$ and high levels of $Fcrg1$ (CD64), $Mertk$, and $Adgre1$ (F4/80),

indicating that they are macrophages (Figure 1F). An additional cluster

expressed $Mafb$ together with $Fcgr1$ and $Adgre1$ but not $Mertk$, fitting the

description of monocytes. Altogether the scRNA-Seq data confirmed that our

flow cytometry gating had identified all macrophages in the thymus.

Importantly, MerTK^{+} cells could not be labeled by intravenously injected CD45

antibody (Figure 1G), proving that they reside in the parenchyma of the

organs and not in the blood vessels. Based on the above data, we will refer to

CD64^{+}F4/80^{+}MerTK^{+}CD11b^{lo}FSC^{hi} cells as thymic macrophages. The smaller

CD64^{+}F4/80^{lo}CD11b^{+}FSC^{hi} population did not express MerTK but most of

them expressed Ly6C, and we classified them as thymic monocytes.

Thymic macrophages expressed CD11c, MHC2, and SIRPα making them

partially overlap with CD11c^{+}MHC2^{+} classical dendritic cells (cDCs), thus

making problematic the unambiguous identification of thymic cDCs based only

on these two markers (Figure 1 – figure supplement 4). Proper identification of

cDC in the thymus requires the exclusion of macrophages based on CD64 or

MerTK staining. Otherwise, the cDCs, particularly the SIRPα^{+} cDC2 subset,

would be contaminated with macrophages that account for ~25% of cDC2

(Figure 1 – figure supplement 5).

Thymic macrophages were ~0.1% of all the cells in the thymus of young

adult mice and numbered ~4×10^5 on average per mouse (Figure 1H). We did

not find statistically significant differences in their percentages between 4 and

11 weeks of age. Still, there was a significant decline in their numbers with

age, consistent with the beginning of thymic involution (Figure 1I).

Figure 1. Thymic macrophages (ThyMacs) can be identified by the

expression of CD64, MerTK, and F4/80. A Flow cytometric analysis of

enzymatically digested thymus tissue with macrophage markers CD64,

MerTK, F4/80, and CD11b. **B** Gating strategy for identifying thymic

macrophages: CD64$^+$FSC^{hi} are first gated; the F4/80$^+$CD11b^{lo} cells among

them are the thymic macrophages (ThyMacs), while F4/80^{lo}CD11b$^+$ are the

thymic monocytes (ThyMonos). **C** Pappenheim (Hemacolor Rapid staining

kit) staining of sorted ThyMacs. **D** Lack of expression of lineage markers

associated with other cell types on ThyMacs. **E** The expression on ThyMacs

of three receptors for phosphatidylserine that participate in the phagocytosis

of apoptotic cells. **F** scRNA-Seq data showing the expression of prototypical

macrophage genes *Mafb*, *Fcgr1* (CD64), *Mertk*, and *Adgre1* (F4/80) among

thymus cells sorted as *Csf1r*$^{GFP+}$ and *Cd11c*$^{YFP+}$. **G** Labeling of ThyMacs with

intravenously injected anti-CD45-PE antibody or PBS. The labeling of blood

leucocytes is shown for comparison. **H** Average numbers and percentages of

ThyMacs in 4-11 weeks old mice, n=82. **I** Comparison of the numbers and

percentages of ThyMacs in mice of different ages, n=82. All flow cytometry

plots are representative of at least 3 independent repeats. The numbers in the

flow cytometry plots are the percent of cells in the respective gates. Data in **H**

and **I** represent mean±SEM. Statistical significance in **I** was determined with

one-way ANOVA.

Transcriptional signature of thymic macrophages

To further understand the identity and functions of the thymic

macrophages, we analyzed the RNA sequencing data from the IMMGEN's

Open Source Mononuclear Phagocyte profiling. We first examined the

expression of the core signature macrophage genes[38] and found that they

were enriched in thymic macrophages but not in *Sirpa*$^+$ or *Xcr1*$^+$ thymic cDCs

(Figure 2A). On the contrary, cDC core signature genes were abundantly

expressed in both thymic cDC subsets but not in thymic macrophages. Thus,

although thymic macrophages and cDCs share the thymic microenvironment

and expression of CD11c and MHC2, they have distinct transcriptional

profiles.

We then compared the gene expression profile of thymic macrophages to

that of other well-characterized macrophage populations from the IMMGEN

database. Because of the abundance of samples, we limited our comparison

to only nine types of tissue-resident macrophages under steady-state

conditions – splenic red pulp macrophages, Kupffer cells, broncho-alveolar

lavage macrophages, large peritoneal cavity macrophages, white adipose

tissue macrophages, aorta macrophages, central nervous system microglia,

and spinal cord microglia. Principal component analysis revealed that thymic

macrophages were most closely related to splenic red pulp macrophages and

Kupffer cells (Figure 2B).

To better identify the unique functions of thymic macrophages, we looked

for differentially expressed genes in these cells compared to other tissue-

resident macrophages. We set three criteria: 1) high expression in thymic

macrophages (>500); 2) >5-fold higher expression than the average value in

the nine populations of non-thymic macrophages; 3) expression in thymic

macrophages is higher than any non-thymic macrophage samples. A total of

44 genes met these criteria, and we consider them to constitute the

transcriptional signature of thymic macrophages (Figure 2C). These included

several degradation enzymes and their inhibitors (*Cst7*, *Mmp2*, *Mmp14*,

Dnase1l3, *Serpina3g*, *Acp5*), non-classical MHC molecules (*H2-M2*, *H2-Q6*,

H2-Q7), metabolic enzymes (*Chst2*, *Ass1*, *Kynu*, *Cp, Dgat2*, *Sorl1*, *Lap3*),

molecules involved in innate immunity (*Ifit2*, *Il18bp*, *Mefv*, *Lgals3bp*) and

extracellular signaling molecules and their receptors (*Pdgfa*, *Cxcl16*, *Il2rg*,

Gpr157). We also looked for transcription factors (TFs) highly expressed in

thymic macrophages and could potentially regulate critical gene networks in

these cells. A total of 25 TFs were highly expressed in thymic macrophages

(>250) and were at least 2-fold higher as compared to the non-thymic

macrophages (Table 1). Among them were several TFs involved in type I

interferon (IFN-I) signaling (*Stat1*, *Stat2*, *Irf7*, and *Irf8*) and lipid metabolism

(*Nr1h3*, *Pparg*, *Srebf1*, and *Rxra*) (Figure 2D). Notably, *Runx3*, which is

essential for the development and function of cytotoxic T lymphocytes[48],

innate lymphoid cells[49], and Langerhans cells[50], was highly expressed in

thymic macrophages. *SpiC*, which has well-documented roles in the

development of red pulp macrophages in the spleen and bone marrow

macrophages[51,52], was also highly expressed in thymic macrophages,

further strengthening the argument for the similarity between thymus, spleen,

and liver macrophages. To confirm the expression of *Spic* in thymic

macrophages, we analyzed the thymus of $Spic^{GFP}$ mice[52]. We found that all

$Spic^{GFP+}$ cells were macrophages (Figure 2 – figure supplement 1), making

them the most specific thymic macrophage reporter strain compared with

$Lyz2^{GFP}$, MAFIA ($Csf1r^{GFP}$), $Cd11c^{YFP}$, and $Cx3cr1^{GFP}$ mice (Figure 2 – figure

supplement 2). However, only ~80% of thymic macrophages were $Spic^{GFP+}$

suggesting heterogeneity within the cells (Figure 2 – figure supplement 3).

Several dominant pathways emerged when we grouped the 500 most

highly expressed genes in thymic macrophages according to gene ontology

(GO) terms (Figure 2E). Notably, five of the ten most highly enriched GO

pathways concerned antigen presentation of both exogenous and

endogenous antigens. These data complement our flow cytometry findings of

expression of MHC2 and suggest that thymic macrophages could be potent

antigen-presenting cells and might play a role in negative selection or agonist

selection of thymocytes. Two other highly enriched GO pathways were

involved in lysosomal biogenesis and functions, highlighting the high capacity

of these cells to degrade phagocytosed material. Thus, our transcriptional

analysis has revealed that thymic macrophages are bona fide macrophages

that bear significant similarity to spleen and liver macrophages and are

specialized in lysosomal degradation of phagocytosed material and antigen

presentation.

Figure 2. Transcriptional profile of thymic macrophages (ThyMacs). A

Expression of cDC-specific genes (top) and macrophage-specific genes

(bottom) in ThyMacs and two populations of thymic cDCs (ThyDCs) – *Xcr1*[+]

ThyDCs and *Sirpa*[+] ThyDCs. **B** Principal components analysis of ThyMacs

and nine other populations of tissue-resident macrophages in duplicates. **C**

Highly expressed (>500) genes enriched (>5-fold) in ThyMacs (4 samples)

compared to nine other tissue-resident macrophage populations (two samples

each). The genes in red are >10-fold up-regulated in thymic macrophages. **D**

Comparison of the geometric mean expression of transcription factors in

thymic macrophages (4 samples) and the nine other macrophage populations

(2 samples each). Transcription factors with expression >250 and fold

change >2 are marked with red dots. **E** Top 10 GO pathways in ThyMacs

based on the 500 most highly expressed genes in these cells. Th – ThyMacs,

Spl – spleen red pulp macrophages, Lv – Kupffer cells, BAL – broncho-

alveolar lavage macrophages, PC – peritoneal cavity macrophages, Ao –

aorta macrophages, Ht – heart macrophages, WAT – white adipose tissue

macrophages, CNS – central nervous system microglia, SC – spinal cord

microglia.

Thymic macrophages can present antigens to T cells and clear

apoptotic cells

Next, we investigated the biological functions of thymic macrophages. Our

findings that these cells express MHC2 and many other genes involved in

antigen presentations prompted us to test if they can efficiently activate T

cells. We pulsed sorted thymic macrophages with chicken ovalbumin (Ova)

and cultured them with naïve OT2 cells labeled with CFSE. The positive

control, thymic DCs, efficiently induced OT2 cell proliferation, while peritoneal

macrophages were very inefficient (Figure 3A and B), similar to other tissue-

resident macrophages[2]. Surprisingly, thymic macrophages induced

proliferation in a considerable proportion (~30%) of OT2 cells as calculated by

FlowJo's Proliferation Modeling module. Thus, thymic macrophages are able

antigen-presenting cells, although not as good as DCs.

302  To confirm the ability of thymic macrophages to clear apoptotic cells, we

303 did in vitro engulfment assay. Thymocytes were induced to undergo apoptosis

304 by Dexamethasone treatment and labeled with pHrodo Red dye. pHrodo Red

305 is weakly fluorescent at neutral pH, but its fluorescence increases significantly

306 at low pH, for example, in lysosomes. Thus, engulfed apoptotic cells can be

307 clearly identified by their strong red fluorescence. We incubated the pHrodo

308 Red-labeled apoptotic thymocytes for 2 hours with sorted thymic or peritoneal

309 cavity macrophages and detected the extent of efferocytosis by fluorescent

310 microscopy. Thymic macrophages were avid phagocytes, and we could

311 record many instances of efferocytosis at this time point (Figure 3C and D).

312 However, peritoneal macrophages were able to phagocytose even more

313 apoptotic cells.

314  To determine if thymic macrophages are the major phagocytes in the

315 thymus in vivo, we evaluated their participation in the phagocytosis of

316 apoptotic cells in the thymus by TUNEL staining. Most TUNEL$^+$ cells could be

317 found clearly inside or closely associated with MerTK$^+$ and TIM4$^+$ cells in the

318 thymus (Figure 3E and F). On average, ~85% of all TUNEL$^+$ cells were within

319 5 µm of MerTK$^+$ cells, confirming that thymic macrophages are the dominant

320 phagocytic population in the thymus (Figure 3G). The degree of co-

321 localization between TUNEL$^+$ cells and TIM4$^+$ cells was slightly lower, ~75%

322 on average, possibly reflecting the absence of TIM4 expression on a small

323 proportion of thymic macrophages (Figure 1E).

Figure 3. Thymic macrophages can present antigens to T cells and

phagocytose apoptotic cells. A Naïve OT2 T cells were labeled with CFSE

and cultured with purified thymic dendritic cells (ThyDCs), thymic

macrophages (ThyMacs), or peritoneal cavity macrophages (PC Macs) in the

presence or absence of chicken ovalbumin (Ova). Three days later, the CFSE

dilution was assessed by flow cytometry. **B** Quantification of the cell division

in naïve OT2 cells by using the Cell Proliferation module in FlowJo that

calculates the percent of cells from the initial population that has undergone

division. **C** Example immunofluorescent images of ThyMacs or PC Macs

phagocytosis apoptotic thymocytes. The macrophages were labeled with

eFluor 450, while the apoptotic thymocytes – with pHrodo Red. An intense

red signal within the macrophages indicates phagocytosed thymocytes. **D**

Quantification of the percentage of macrophages that have engulfed at least

one thymocyte (phagocytic index). **E** Example images showing co-localization

of TUNEL$^+$ apoptotic cells and MerTK$^+$ ThyMacs in thymic sections. **F**

Example images showing co-localization of TUNEL$^+$ apoptotic cells and TIM4$^+$

ThyMacs in thymic sections. Scale bars in **E** and **F** are 50 µm. **G** Frequencies

of the co-localization of TUNEL$^+$ signal with MerTK$^+$ and TIM4$^+$ cells. Flow

cytometry plots in **A** are representative of three independent experiments. All

immunofluorescent images are representative of at least 3 independent

repeats. Data in **B**, **D**, and **G** represent mean±SEM. Each symbol in **B** and **G**

is an individual mouse. Each symbol in **D** is a field of view.

Expression of *Timd4* and *Cx3cr1 can distinguish* two populations of

thymic macrophages

Our phenotypic characterization showed clear signs of heterogeneity within

thymic macrophages, including the presence of TIM4$^+$ and TIM4$^−$ cells (Figure

1E) and *Cx3cr1*$^{GFP+}$ and *Cx3cr1*$^{GFP−}$ cells (Figure 2 – figure supplement). To

determine the degree of thymic macrophage heterogeneity, we turned to our

scRNA-Seq data. At least 7 clusters could be identified and assigned to

different cell types by specific marker expression (Figure 4A and B), including

macrophages, B cells, pDCs, contaminating thymocytes, and multiple cDC

clusters. Once we zoomed onto *Mafb*-expressing cells, we could distinguish

three separate populations: 1) monocytes that expressed high levels of *Ly6c2*

and *Itgam* (CD11b) but did not express *Mertk*; 2) *Timd4*$^+$ (encoding TIM4)

macrophages that also expressed high levels of *Spic* and *Slc40a1*, but low

levels of *Cx3cr1*; 3) *Cx3cr1*$^+$ macrophages that expressed low levels of

Timd4, *Spic*, and *Slc40a1* (Figure 4C and D). Both macrophages and

monocytes expressed *Fcgr1* (CD64). Thus, these data indicate that thymic

macrophages consist of two populations with distinct expression profiles.

We confirmed the results from scRNA-Seq by flow cytometry. We could

identify discrete TIM4$^+$*Cx3cr1*$^{GFP-}$ and TIM4$^-$*Cx3cr1*$^{GFP+}$ macrophages (Figure

4E). There was even a TIM4$^+$*Cx3cr1*$^{GFP+}$ intermediate population that could

not be distinguished in the scRNA-Seq dataset, likely because of the lack of

statistical power. To determine the localization of the two distinct macrophage

populations, we stained thymic sections from *Cx3cr1*^{GFP} mice with an antibody

to MerTK. The *Cx3cr1*$^{GFP-}$ MerTK$^+$ cells correspond to *Timd4*$^+$ macrophages,

while the *Cx3cr1*$^{GFP+}$MerTK$^+$ cells would be the *Cx3cr1*$^{GFP+}$ macrophages.

Strikingly, the two macrophage populations showed distinct localization in

young mice. *Timd4*$^+$ macrophages were located in the cortex, while the

Cx3cr1^{GFP+} macrophages resided in the medulla and the cortico-medullary

junction (Figure 4F). The result was confirmed with direct staining for TIM4

that showed intense signal in the cortex, particularly in the deep cortex, and

absence of staining in the medulla (Figure 4 figure supplement 1). However,

the medulla still featured many CD64$^+$ macrophages.

To better understand the differences between the two populations of thymic

macrophages, we looked for differentially expressed genes. We included the

thymic monocytes in the comparison, as these cells clustered the closest to

macrophages. *Timd4*$^+$ macrophages expressed the highest levels of the

transcription factors *Spic*, *Maf*, and *Nr1h3*; the receptors for apoptotic cells

Axl, *Mertk*, and *Timd4*; and many Slc transporters such as *Slc40a1*, *Slc1a3*,

Slco2b1, *Slc11a1*, and *Slc7a7* (Figure 4G and Table 2). *Cx3cr1*$^+$

macrophages expressed high levels of the transcription factor *Runx3*; a

distinct set of phosphatidylserine receptors such as *Stab1*, *Anxa5*, and *Anxa3*;

many degradative enzymes such as *Mmp2*, *Mmp14*, *Dnase1l3*, *Acp5*, *Lyz2*,

Ctsz, *Ctss*, *Ctsd*, *Ctsl*; cytokines such as *Pdgfa*, *Cxcl16*, and *Ccl12*; and

molecules involved in MHC1 antigen presentation such as *B2m*, *H2-M2*, *H2-*

K1, *H2-Q7*. Thymic monocytes were characterized by differential expression

of the typical monocyte genes *Ly6c2*, *Ccr2*, and *S100a4*, and genes involved

in MHC2 antigen presentation such as *Ciita*, *H2-DMb1*, *H2-Ab1*, and *Cd74*.

Figure 4: **Two populations of macrophages with distinct localization**

exist in the thymus. A Identification of the clusters from the scRNA-Seq data

based on lineage-specific markers. **B** Expression of lineage-specific markers

in different clusters. **C** UMAP clusters from **A** with high expression of the

transcription factor *Mafb* fall into three groups: monocytes, *Timd4*$^+$

macrophages, and *Cx3cr1*$^+$ macrophages. **D** Expression of the indicated

genes in the three *Mafb*-positive clusters. **E** A flow cytometry plot of *Cx3cr1*^{GFP}

and TIM4 expression in ThyMacs. The plot is representative of >10 individual

experiments. The numbers inside the plot are the percentages of the cell

populations in the respective gates. **F** Immunofluorescent staining of the

thymus of *Cx3cr1*^{GFP} mouse stained with MerTK (a marker for all

macrophages) and Keratin 5 (a marker for medulla). The scale bar is 150 μm.

Areas in the cortex, medulla, and the cortico-medullary junction (CMJ)

represented by the dashed boxes are enlarged to the right to show the co-

localization of *Cx3cr11*^{GFP} and MerTK signal in CMJ and medulla, but not in

the cortex. The scale bars in the images to the right are 20 μm. The images

are representative of three individual mice. **G** Differentially expressed genes

among *Timd4*$^+$ thymic macrophages, *Cx3cr1*$^+$ thymic macrophages, and

thymic monocytes. The negative log$_{10}$ p-values for the genes expressed in

each cluster were calculated as described in the Materials and Methods, and

the top 50 differentially expressed genes were plotted in the figure. Ten of

these genes are listed on the left.

Yolk-sac progenitors contribute to embryonic thymic macrophages

The ontogeny of thymic macrophages has been examined by only one

study since the realization that many tissue-resident macrophages are

descendants of embryonic progenitors[37]. Based on $Flt3^{Cre}$ fate-mapping,

the authors concluded that most adult thymic macrophages derive from

HSCs. To determine if yolk-sac (YS) progenitors contribute to embryonic

thymic macrophages, we used $Cx3cr1^{CreER}$ fate mapping[53]. Injection of 4-

OHT at E9.5 in $ROSA26^{LSL-GFP}$ mouse mated with a $Cx3cr1^{CreER}$ male

permanently tags YS progenitors and their descendants with GFP (Figure

5A). Indeed, E19.5 microglia that are exclusively derived from YS progenitors

were labeled to a high degree (Figure 5B). After adjusting for incomplete

labeling based on the microglia, we found that at E15.5 >50% of thymic

macrophages were fate mapped, i. e. from YS origin (Figure 5C). However,

GFP$^+$ thymic macrophages decreased to just ~11% at E19.5, suggesting that

YS progenitors establish the embryonic thymic macrophage pool but are

quickly replaced by subsequent wave(s) of macrophages.

Differential contribution of adult bone marrow-derived monocytes to

the two thymic macrophage populations

To investigate the possibility that thymic macrophages arise from adult

bone marrow-derived monocytes, we devised two complementary

experiments. First, we evaluated the contribution of circulating adult

monocytes to thymic macrophages without the confounding effect of radiation

damage on the thymus. We created shield chimeras by subjecting CD45.2

mice to a lethal dose of irradiation while protecting their upper body and the

thymus with a 5 cm lead shield, followed by reconstitution with CD45.1 bone

marrow (Figure 5D). We analyzed *Timd4*$^+$ and *Cx3cr1*$^+$ thymic macrophages

separately after 6 weeks because we suspected they might have different

origins. As Cx3cr1 protein expression was low on thymic macrophages

(Figure 5B), we defined the *Cx3cr1*$^+$*Timd4*$^-$ population as TIM4$^-$. The donor-

derived monocytes in the blood were, on average, 57%, but less than 2% of

TIM4$^+$ thymic macrophages were CD45.1$^+$ (Figure 5E and F), suggesting very

limited contribution of adult circulating monocytes to the TIM4$^+$ macrophage

pool. The percentage of HSC-derived TIM4$^-$ macrophages (on average 23%)

was intermediate between the monocytes and TIM4$^+$ macrophages, pointing

out that a sizeable part of TIM4$^-$ cells was derived from adult HSCs.

We also transplanted E15.5 embryonic thymuses from SpiC^{GFP+} CD45.2

under the kidney capsule of adult CD45.1 mice and analyzed them six weeks

later (Figure 5G). By that time, >99% of thymocytes in the transplanted

thymus were derived from CD45.1$^+$ host HSCs, indicating successful

replacement by HSC-derived progenitors (Figure 5 – figure supplement 1).

TIM4$^-$ thymic macrophages were derived entirely from host HSCs, just like

thymocytes. In contrast, most TIM4$^+$ cells (on average 70%) were donor-

derived (Figure 5H and I). Moreover, only CD45.2$^+$ TIM4$^+$ macrophages

expressed SpiC^{GFP} (Figure 5 – figure supplement 2). As expected, thymic

macrophages in the endogenous thymus were all CD45.1$^+$. The results from

our transplantation experiments show that the progenitors of most TIM4$^+$

thymic macrophages are of embryonic origin, while TIM4$^-$ cells are derived

from adult monocytes. Altogether our results suggest that the two populations

of thymic macrophages have different origins. TIM4$^+$ cells are derived from

embryonic precursors and can survive long-term without much contribution

from adult HSC and monocytes. In contrast, TIM4⁻ thymic macrophages rely

mostly on adult HSCs for their generation and replacement.

Figure 5. Yolk sac (YS), non-YS-derived embryonic progenitors, and

adult hematopoietic stem cells sequentially contribute to the thymic

macrophage pool. A Scheme of the YS-progenitor labeling experiments.

E9.5 pregnant *ROSA26^{LSL-GFP}* mice mated with *Cx3cr1^{CreER}* males were

injected with 4-hydroxytamoxifen (4-OHT) and sacrificed at E15.5 or E19.5. **B**

Representative flow cytometry plots of the *Cx3cr1^{GFP}* expression in microglia

(CD45⁺CD11b⁺ cells in the brain) and ThyMacs of the pups at E19.5. **C**

Frequencies of GFP⁺ ThyMacs at E15.5 and E19.5 adjusted to the degree of

labeling of microglia. **D** Scheme of the shield chimera experiments. Congenic

CD45.2 mice were lethally irradiated with their upper body protected by a 5

cm thick lead shield and then injected with CD45.1$^+$ bone marrow. **E**

Representative flow cytometric analysis of CD115$^+$CD11b$^+$ blood monocytes,

TIM4$^+$ and TIM4$^-$ thymic macrophages for donor-derived (CD45.1$^+$) and host-

derived (CD45.2$^+$) cells. Non-chimeric CD45.1 and CD45.2 samples serve as

controls for the gating. **F** Frequencies of donor-derived blood monocytes,

TIM4$^+$ and TIM4$^-$ ThyMacs. **G** Scheme of the thymus transplantation

experiments. Embryonic thymuses from E15.5 SpiC^{GFP+} CD45.2$^+$ mice were

transplanted under the kidney capsule of CD45.1$^+$ mice and analyzed six

weeks later. **H** Representative flow cytometry plots of donor (CD45.2$^+$) vs.

host (CD45.1$^+$) derived TIM4$^+$ and TIM4$^-$ ThyMacs in the transplanted thymus.

The host thymus (endogenous thymus) serves as a negative control. **I**

Frequencies of CD45.2$^+$ (donor-derived) cells among TIM4$^+$ and TIM4$^-$

ThyMacs in the transplanted and endogenous thymuses of the mice. Data in

C, **F**, and **I** are mean±SEM with two litters, seven, and five mice per group,

respectively. The numbers in the flow cytometry plots are the percent of cells

in the respective gates. Each symbol in the graphs is an individual mouse or

embryo.

Thymic macrophages can proliferate in situ

TIM4$^+$ macrophages can persist for many weeks in the thymus without

constant replacement from blood monocytes, suggesting they can divide in

situ. Staining for the proliferation marker Ki67 revealed that ~4% of all thymic

macrophages expressed this marker compared to an isotype control (Figure

6A and B). To prove that thymic macrophages are proliferative, we tested the

incorporation of the nucleotide analog 5-Ethynyl-2'-deoxyuridine (EdU).

Short-term EdU labeling experiments unexpectedly revealed that thymic

macrophages become EdU$^+$ with faster kinetics than thymocytes (Figure 6C).

The most likely explanation for this puzzling result is that some of the thymic

macrophages have engulfed apoptotic thymocytes that have recently divided

and incorporated EdU. Thus, EdU could have accumulated in these

macrophages through phagocytosis, not cell division. To circumvent this

caveat, we designed a pulse-chase experiment (Figure 6D). Mice were

injected daily with EdU for 21 days so that all cells that proliferated in that

period would incorporate the label. Most thymocytes and thymic

macrophages became EdU$^+$ at d. 21 (Figure 6E). After 21 more days of

"chase period", only ~0.2% of thymocytes had retained the EdU label,

consistent with the existence of a tiny population of long-term resident

thymocytes consisting mainly of regulatory T cells and NKT cells[54] (Figure

6E and F). However, ~5% of the thymic macrophages were EdU$^+$, suggesting

they divided during the labeling period. We also sorted thymic macrophages

and subjected them to cell cycle analysis. Although almost all thymic

macrophages were in G0/G1 phase, a small population of ~3% was in the

G2/M phase of the cell cycle (Figure 6G and H). Surprisingly, most *Mki67*$^+$

thymic macrophages belonged to the *Cx3cr1*$^+$ subset, and only a few of the

Timd4$^+$ cells were positive (Figure 6I). Collectively, four independent

approaches documented that a small proportion (3-5%) of thymic

macrophages are actively dividing under homeostatic conditions within the

thymus. The majority of the dividing cells were from the adult HSC-derived

Cx3cr1$^+$ subset. *Timd4*$^+$ macrophages were primarily quiescent.

Figure 6. Thymic macrophages exhibit a low degree of proliferation.

A Example flow cytometry plots of Ki67 staining of thymic macrophages

(ThyMacs). **B** Frequency of Ki67$^+$ thymic macrophages. **C** Example flow

cytometry plots of the EdU accumulation in thymocytes and thymic

macrophages 2 hours after 1 mg EdU i.p. or vehicle injection. The numbers

inside flow plots are the percentage of EdU$^+$ cells from mice injected with

EdU. Data are representative of three independent experiments. **D** Scheme of

EdU pulse/chase experiment: mice were injected daily with 1 mg EdU i.p for

21 days and rested for 21 more days. **E** Example flow cytometry plots of EdU

staining of thymocytes (upper row) and ThyMacs (lower row). **F** Frequencies

of EdU$^+$ cells among thymocytes (top graph) and ThyMacs (bottom graph). **G**

Example flow cytometry plot of cell cycle analysis of FACS sorted ThyMacs. **H**

Frequencies of ThyMacs in different stages of the cell cycle. **I** UMAP plot of

Mki67 expression in *Mafb*-positive clusters from the scRNA-Seq data

described in Figure 4. The numbers in the flow cytometry plots are the percent

of cells in the respective gates. Data are mean±SEM from three mice (**B** and

H) or 6-7 individual mice (**F**). Each dot is an individual mouse.

***Cx3cr1*$^+$ cells give rise to *Timd4*$^+$ cells during embryonic development**

To determine if the two populations of thymic macrophages are related, we

first analyzed the kinetics of their appearance during embryonic development.

At the earliest time point (E14.5), all thymic macrophages were *Cx3cr1*$^+$, and

only ~20% were also TIM4$^+$ (Figure 7A and B). The proportion of TIM4$^+$ cells

increased at E17.5, and TIM4$^+$*Cx3cr1*$^-$ cells started to appear. In the neonatal

period, almost all macrophages were TIM4$^+$, and very few remained TIM4$^-$.

The proportion of TIM4$^-$ cells increased in 6 weeks old mice, but TIM4$^+$

macrophages remained the dominant population. These kinetics (Figure 7C)

are consistent with *Timd4*$^+$ macrophages developing from *Cx3cr1*$^+$ cells

before birth. Another plausible scenario is that distinct progenitors give rise to

different thymic macrophage populations (e.g., YS-progenitors give rise to

Cx3cr1$^+$*Timd4*$^-$ and HSC-derived progenitors develop into *Timd4*$^+$

macrophages). To test the latter hypothesis, we revisited the fate mapping of

YS progenitors (Figure 5A). Although a larger part (~60% at E15.5) of fate-

mapped cells were *Cx3cr1*$^+$TIM4$^-$ cells (Figure 7D), a substantial proportion

(~40% at E15.5) of fate-mapped TIM4$^+$ macrophages could clearly be

identified at both E15.5 and E19.5, suggesting that YS progenitors can give

rise to both *Cx3cr1*$^+$ and *Timd4*$^+$ cells. Thus, the simplest explanation for our

findings is that *Timd4*$^+$ cells develop from *Cx3cr1*$^+$ cells during embryonic

development. This transition is complete in the first week after birth as there

were essentially no *Cx3cr1*$^+$TIM4$^-$ thymic macrophages remaining at d.7

(Figure 7A and B). To formally demonstrate that *Cx3cr1*$^+$ macrophages can

give rise to *Timd4*$^+$ cells during embryonic development, we injected 4-OHT in

E15.5 pregnant females carrying Cx3cr1^{CreER} X ROSA26$^{LSL-GFP}$ fetuses (Fig.

7E). At this time almost all thymic macrophages are *Cx3cr1*$^+$ (Fig. 7A). Just

before birth, at E19.5, we could identify a sizeable population of

TIM4$^+$CX3CR1$^-$ among fate-mapped cells, suggesting that they originate from

Cx3cr1$^+$ progenitors (Fig. 7F and G).

Figure 7. *Timd4*⁺ thymic macrophages are derived from *Cx3cr1*⁺ cells during embryonic development. A Example flow cytometry plots for the expression of *Cx3cr1*^{GFP} and TIM4 on thymic macrophages at different times during embryonic development (E14.5, E17.5), immediately after birth, at 7 days, and 6 weeks of age. **B** Frequencies of *Timd4*⁺*Cx3cr1*⁻ (*Timd4* single-positive or *Timd4*SP), *Timd4*⁺*Cx3cr1*⁺(double-positive or DP), and *Cx3cr1*⁺*Timd4*⁻ (*Cx3cr1* single-positive or *Cx3cr1*SP) thymic macrophages at the indicated time points. **C** Kinetics of the changes in different subpopulations of thymic macrophages from E14.5 to 6 weeks. **D** Frequencies at E15.5 and E19.5 of GFP-labeled cells among TIM4⁺ or TIM4⁻ cells in *Cx3cr1*^{CreER} X *ROSA26*$^{LSL-GFP}$ embryos treated with 4-OHT at E9.5. **E** Scheme of the fate-mapping experiments showing the relationship between *Cx3cr1*⁺ and *Timd4*⁺

thymic macrophages during embryonic development. E15.5 pregnant

ROSA26^LSL-GFP mice mated with *Cx3cr1^CreER* males were injected with 4-

hydroxytamoxifen (4-OHT) and sacrificed at E19.5. **F** Representative flow

cytometry staining for TIM4 and CX3CR1 in fate-mapped GFP$^+$ thymic

macrophages at E19.5. The panel to the right is the isotype control for

CX3CR1-PE staining. **G** Frequencies of TIM4$^+$CX3CR1$^-$ cells among fate-

mapped GFP$^+$ macrophages. Data are from at least two independent

experiments for each panel. Each symbol is an individual mouse or embryo.

Cx3cr1$^+$ thymic macrophages slowly accumulate with age at the

expense of Timd4$^+$ cells

To understand the dynamics of the two resident thymic macrophage

populations with age, we induced recombination in *Cx3cr1^CreER* X *ROSA26^LSL-*

GFP mice during the neonatal period (Figure 8A) or at 6 weeks of age (Figure

8C) and compared the proportion of GFP$^+$ cells 3 and 42 days after labeling.

The extent of labeling of TIM4$^+$ thymic macrophages did not change within

these 6 weeks, no matter whether the mice were treated with Tamoxifen in

the first week after birth or at 6 weeks (Figure 8B and D), suggesting an

absence of a significant influx from unlabeled cells (e.g., monocytes). In

contrast, the proportion of labeled TIM4$^-$ thymic macrophages decreased

significantly 6 weeks after Tamoxifen injection in neonatal and adult mice,

suggesting that this population was diluted by unlabeled cells. To further

substantiate these findings, we examined older WT mice and found out that

the proportions of TIM4$^-$ thymic macrophages increased with age, and in mice

>8 months old, they accounted for ~70% of all macrophages in the organ

(Figure 8E). As these changes in the proportions of the thymic macrophage

subpopulations occurred at the background of thymic involution, we wanted to

know if the accumulation of TIM4⁻ cells was only relative or also in absolute

cell numbers. In contrast to TIM4⁺ thymic macrophages that reached peak

numbers at an early age and then declined significantly, TIM4⁻ cells tended to

increase their numbers in older mice (Figure 8F). Thus, we conclude that,

after birth, the numbers of TIM4⁺ macrophages follow the kinetics of the

thymus size – increase in young and decrease in old mice, and they are not

replaced by other cells. In contrast, since the first week of life, *Cx3cr1⁺* cells

are recruited to the thymus, accumulate with age, and in old mice, form the

predominant phagocytic population in the organ.

Figure 8. *Timd4⁺* thymic macrophages are progressively lost, while

***Cx3cr1⁺* cells slowly accumulate with age. A** Scheme of the neonatal fate

mapping: A nursing dam was injected twice with Tamoxifen (Tam) or vehicle

(Veh) in the first week after giving birth to $Cx3cr1^{CreER}$ X $ROSA26^{LSL\text{-}GFP}$ pups.

The mice were sacrificed three or 42 days after the last injection, and the

degree of labeling of TIM4$^+$ and TIM4$^-$ thymic macrophages was examined by

flow cytometry. **B** Frequencies of GFP$^+$ TIM4$^+$ or TIM4$^-$ thymic macrophages

from neonatally fate-mapped mice after 3 and 42 days. Vehicle-injected

nursing dam litters (Veh) served as a control for non-specific labeling. **C**

Scheme of the adult fate mapping: Six weeks old $Cx3cr1^{CreER}$ X $ROSA26^{LSL\text{-}}$

^{GFP} mice were injected twice with Tamoxifen (Tam) or vehicle (Veh). The

mice were sacrificed three or 42 days after the last injection, and the degree

of labeling of TIM4$^+$ and TIM4$^-$ thymic macrophages was examined by flow

cytometry. **D** Frequencies of GFP$^+$ TIM4$^+$ or TIM4$^-$ thymic macrophages from

adult fate-mapped mice after 3 and 42 days. **E** Frequencies of TIM4$^-$ thymic

macrophages at different ages. **F** Changes in the numbers of TIM4$^-$ and

TIM4$^+$ thymic macrophages with age. The data are mean±SEM from 2

independent experiments (**B**) or at least 3 individual mice per time point (**D**,

E, and **F**). Each symbol is an individual mouse. Statistical significance in the

difference between Tamoxifen-treated samples at different time points was

determined with unpaired Student's t-test (**B** and **D**). One-way ANOVA was

used to assess the significance of the change in TIM4$^+$ and TIM4$^-$ ThyMacs

percentages and numbers with age (**E** and **F**).

Discussion

Here, we have described the phenotype, transcriptional profile, localization,

diversity, ontogeny, and maintenance of macrophages in the thymus. These

cells express the typical macrophage markers CD64, MerTK, and F4/80 and

are transcriptionally most similar to splenic red pulp macrophages and liver

Kupffer cells. However, they have a unique expression profile dominated by

genes involved in antigen presentation and lysosomal degradation. We found

that thymic macrophages consist of two populations with distinct localization.

Timd4$^+$ macrophages occupied the cortex, while *Cx3cr1*$^+$ cells were located in

the medulla and the cortico-medullary junction. While YS-derived

macrophages dominated the early stages of thymus development, they were

quickly replaced by non-YS embryonic progenitors that gave rise to the

Timd4$^+$ thymic macrophages that persisted into adulthood and formed the

main macrophage population in young mice. *Cx3cr1*$^+$ macrophages slowly

accumulated after birth and became the most abundant population in old

mice.

Altogether our data depict thymic macrophages as typical tissue-resident

macrophages originating from multiple hematopoietic waves, surviving long-

term, and expressing the core macrophage-specific genes. They are most

similar transcriptionally to splenic red pulp macrophages and Kupffer cells,

which is not surprising considering that they all specialize in efferocytosis and

have efficient lysosomal degradation machinery. These three populations

also shared expression of the transcription factor *Spic* that is induced by

heme released following red blood cells phagocytosis[52]. However, the

thymus is not known as a place for erythrocyte degradation. Thus, the

mechanism for *Spic* up-regulation in thymic macrophages is unclear.

The unique features of thymic macrophages include high expression of

genes involved in the IFN-I pathway, antigen presentation, and lysosomal

degradation. The up-regulation of IFN-I-stimulated genes such as *Stat1*,

Stat2, *Irf7*, and *Irf8* can be explained by the constitutive secretion of IFN-I by

thymic epithelial cells[55,56]. The purpose of IFN-I expression in the thymus

in the absence of a viral infection is unclear. Still, one possibility is that it

mediates negative selection to IFN-dependent genes as part of central

tolerance.

Thymic macrophages highly express molecules involved in antigen

presentation, including MHC1 and MHC2, although the latter is expressed at

lower levels than in cDCs, and are functionally competent to induce T cell

activation. Thus, they have the potential to present antigens for both negative

selection and agonist selection. These two activities have traditionally been

assigned solely to cDCs[57]. However, recent evidence suggests that

negative selection is most efficient when the cell that presents the antigen to

auto-reactive thymocytes is also the one that phagocytoses it[58]. So,

macrophages' participation in thymocyte selection needs to be re-evaluated.

The extraordinary ability of thymic macrophages to engulf and degrade

apoptotic thymocytes has been appreciated for a long time[31], and our RNA-

Seq data provides additional supporting evidence for this function by

highlighting the up-regulation of pathways involved in lysosomal degradation.

An interesting topic for future research would be understanding how the

metabolites derived from apoptotic cells are returned to the microenvironment

to support the proliferation of immature thymocytes. A *SoLute Carrier* (*Slc*)

genes-based program has been described in vitro[59], but its relevance to

tissue-resident macrophages remains to be determined. Altogether, our study

demonstrates that thymic macrophages are a unique subset of tissue-resident

macrophages and support the idea that resident macrophage phenotype is

determined by the combination of ontogeny, microenvironment, and other

factors[60].

Together with the study by Tacke et al., our work builds the following model

for thymic macrophage origin[37]: Thymic macrophages develop in three

distinct waves: YS-derived progenitors dominate the early stages of thymus

development but are replaced before birth by a second wave of YS-

independent embryonic progenitors that forms the bulk of thymic

macrophages after birth and can self-maintain into adulthood. With age, there

is a slow and steady influx of *Timd4$^-$Cx3cr1$^+$* macrophage precursors that

occupy the medulla and cortico-medullary junction, becoming the major

phagocytic population in the thymus of older mice (>8 months). The second

wave of YS-independent macrophages is most likely the progeny of

embryonic HSCs based on *Flt3^{Cre}* fate mapping that showed that >80% of

thymic macrophages in adult mice are descendants of HSCs[37]. Whether

HSC-independent fetal liver monocytes contribute to thymic macrophages and

to what extent awaits the creation of models that can specifically target this

population of progenitors. Recruitment of circulating monocytes to the resident

macrophage pool in the thymus has been ruled out previously by parabiosis

and *Ccr2$^{-/-}$* mice[37]. Our shield chimera experiments have arrived at similar

conclusions. However, the relatively short duration of these experiments and

their focus on the bulk thymic macrophages have prevented the recognition of

the gradual accumulation of *Timd4$^-$* macrophages. Once we zoomed in on this

minor cell population in young mice, the fate mapping clearly showed an influx

of unlabeled progenitors. Whether the progenitors of *Timd4$^-$* macrophages

are monocytes remains to be formally demonstrated. However, monocytes

have been singled out as the source of all macrophage populations exhibiting

replacement in adults examined to date[12,19,22,24,25,30]. An alternative

possibility involves thymocyte progenitors that, under certain circumstances,

have been shown to differentiate into macrophages and granulocytes in the

thymus[42,43]. However, if this occurs in unmanipulated mice at a steady

state remains unclear.

The spatial segregation of the two macrophage populations in the thymus

implies that they might have distinct functions. *Timd4*[+] cells are restricted to

the cortex and are particularly abundant in the deeper cortex, close to the

medulla. Both positive and negative selection of thymocytes occur there, so

we speculate that *Timd4*[+] macrophages might be specialized in efferocytosis

of CD4[+]CD8[+] (double-positive) thymocytes that cannot interact with cortical

thymic epithelial cells and die by neglect or are auto-reactive and undergo

clonal deletion in the cortex[61]. On the other hand, *Cx3cr1*[+] macrophages

accumulate in the medulla – the thymic region specialized in negative

selection to tissue-restricted antigens (TRA). They might contribute to the

process in several ways: 1) by carrying TRAs from blood and peripheral

organs. A similar process has been described for cDC2 (SIRPα[+] DCs)[62]. In

fact, *Cx3cr1*[+] thymic macrophages could have contributed to this role because

they were not distinguished from cDC2 in this study. 2) By capturing TRAs

from *Aire*[+] medullary thymic epithelial cells and presenting them to auto-

reactive thymocytes as shown for DCs[63-65]. 3) By phagocytosing apoptotic

TRA-specific medullary thymocytes, a process we have observed before[58].

The exact involvement of thymic macrophages in the selection events in the

thymus remains to be determined.

The accumulation of the *Cx3cr1*[+] cells in older mice has clear implications

for thymus aging. One key feature of thymus involution is the accumulation of

extracellular matrix produced by fibroblasts and the emergence of white

adipocytes[66]. The *Cx3cr1*[+] subset is the predominant producer of the growth

factor PDGFα that is required for the maintenance of adipocyte stem cells and

can stimulate tissue fibrosis[67,68]. The gradual accumulation of *Cx3cr1*[+]

macrophages could increase the availability PDGFα in the aging thymus

stimulating extracellular matrix production and differentiation of precursors

into adipocytes. This model predicts that limiting the influx of *Cx3cr1*[+]

macrophage precursors could delay thymus involution.

Recent work described a novel phagocytic and antigen-presenting cell type

in the thymus called monocyte-derived DCs[65]. The phenotype of these cells

overlaps with the CD64[+]F4/80[lo]CD11b[+] cells in our study. However, we favor

the classification of these cells as monocytes based on their expression of

Mafb, CD64, and Ly6C (Figure 4D) and lack of expression of the defining DC

transcription factor *Zbtb46* (Figure 4B)[69]. As monocytes can differentiate

into cDC2, particularly in the context of inflammation[70], the precise identity

and the relationship of this population to thymic cDC2 remain to be

established.

In the past several years, scRNA-Seq has come to the forefront of

biologists' efforts to disentangle the cellular diversity of tissues. Several

comprehensive studies have included samples from mouse or human

thymus[71-73]. However, too few thymic macrophages were sampled in

these studies to give meaningful clustering results. Efforts specifically

targeting the thymus have provided considerably more information[74,75], but

macrophage diversity was still not recognized. Characterization of rare

populations such as thymic macrophages (~0.1% of all cells in the thymus)

requires optimized enzymatic digestion procedures and enrichment strategies,

as has already been demonstrated for thymic epithelial cells[76,77]. Our

scRNA-Seq dataset provides a rich resource for the unbiased characterization

of myeloid cells in the thymus and will greatly aid in the understanding of the

myeloid landscape of the thymus.

In summary, our work comprehensively characterizes macrophages in the

thymus and paves the way for the exploration of their functions.

Materials and methods

Mice

C57BL/6Narl (CD45.2) mice were purchased from the National Laboratory

Animal Center, Taipei, Taiwan. MAFIA (MAcrophage Fas-Induced Apoptosis)

[78], $Cx3cr1^{GFP}$ [79], $Spic^{GFP}$ [52], $Cx3cr1^{CreER}$ [53], and $B6.SJL-Ptprca$

$Pepcb/BoyJ$ (CD45.1) mice were purchased from the Jackson Laboratories.

$Cd11c^{YFP}$ [80] and $Lyz2^{GFP}$ [81] mice have been described. Mice ubiquitously

expressing GFP from the $ROSA26$ locus were generated by breeding

$Pdgfra^{Cre}$ [82] and $ROSA26^{LSL-ZsGreen}$ (also known as $ROSA26^{LSL-GFP}$ or Ai6)

mice[83] (both from the Jackson Laboratories). A mouse from this cross was

identified, in which the STOP cassette was deleted in the germline. It was

designated $ROSA26^{GFP}$ and subsequently bred to C57BL/6 mice. All mice

were used at 4-10 weeks of age unless otherwise specified. Mice were bred

and maintained under specific pathogen-free conditions at the animal facility

of National Yang-Ming University. All experimental procedures were approved

by the Institutional Animal Care and Use Committee (IACUC) of National

Yang Ming Chiao Tung University.

Treatment with 5-Ethynyl-2'-deoxyuridine (EdU)

Mice were i.p. injected with 1 mg EdU (Carbosynth) dissolved in PBS daily

for 21 days and then rested for 21 more days. Cells from the thymus were

harvested on day 21 or 42. In some experiments, the mice were sacrificed 2

hours after the first EdU injection.

Shield chimera generation

WT (CD45.2) mice were anesthetized by i.p. injection of 120 µg/g body

weight Ketamine hydrochloride (Toronto Research Chemicals) and 12 µg/g

body weight Xylazine hydrochloride (Sigma). Anesthetized mice were taped

to a 5 cm thick lead block so that the lead block covered the head and the

chest down to the bottom of the rib cage. Then, they were irradiated with a

lethal dose (1000 rad) from a ^{137}Cs source (Minishot II, AXR) so that only their

abdomen and hind legs were exposed. After recovery from anesthesia, the

mice were transfused i.v. with 10^7 bone marrow cells from a congenic

(CD45.1) donor. Then, they were given Trimerin (0.5 mg/mL Sulfadiazine +

0.1 mg/mL Trimethoprim, China Chemical and Pharmaceutical Co., Tainan,

Taiwan) in the drinking water for the first two weeks after the irradiation and

analyzed after six weeks.

Cell isolation from thymus, blood, and peritoneal cavity

Thymocytes were obtained by mechanical disruption of the thymus with a

syringe plunger. For myeloid cell isolation, mouse thymuses were cut into

small pieces and digested with 0.2 mg/mL DNase I (Roche) and 0.2 mg/mL

collagenase P (Roche) in complete DMEM for 20 min at 37°C with frequent

agitation. In some experiments, thymic myeloid cells were enriched by 57%

Percoll PLUS (GE Healthcare) discontinuous gradient centrifugation at 4°C,

1800 rpm, for 20 min without brake. Cells at the interface were collected and

washed with PBS to remove residual silica particles. Then the cells were

resuspended in PBS with 0.5% BSA (HM Biological), filtered through a 70 µm

filter, and kept on ice.

Blood was isolated by cardiac puncture of sacrificed mice and immediately

diluted with PBS. After centrifugation, the cell suspensions were treated with

ammonium chloride-potassium lysis buffer for 3 min on ice once or twice.

Peritoneal cavity cells were obtained by lavage with 5 mL PBS + 2 mM EDTA

(Merck). Following gentle massage, the cavity was opened with an abdominal

incision, and lavage fluid was collected.

Flow cytometry

Single-cell suspensions ($0.5 - 2X10^6$ cells) from thymus, blood, or

peritoneal cavity were blocked with supernatant from 2.4G2 hybridoma (a kind

gift by Dr. Fang Liao, Academia Sinica, Taipei, Taiwan) and stained with

fluorochrome- or biotin-labeled antibodies for 20 min on ice in PBS + 0.5%

BSA + 2 mM EDTA + 0.1% NaN_3 (FACS buffer). The following antibodies

were used: CD11b (clone M1/70), MHC2 (M5/114.15.2), CD11c (N418),

F4/80 (BM8), CD115 (AFS98), SIRPα (P84), CD45 (30-F11), NK1.1 (PK136),

TIM4 (RMT4-54), Gr-1 (RB6-8C5), CD64 (X54-5/7.1), Siglec H (551), Ly6C

(HK1.4), CD3ϵ (145-2C11), CD8α (53-6.7), CD19 (6D5), B220 (RA3-6B2),

CD4 (GK1.5), CD51 (RMV-7), CD45.1 (A20), CD45.2 (104), CX3CR1

(SA011F11), and EpCAM (G8.8) from BioLegend; Axl (MAXL8DS), MerTK

(DS5MMER), and Ki67 (SolA15) were from eBioscience; Siglec F (E50-2440),

CD90.2 (30-H12), and CD11c (HL3) were from BD Biosciences. Cells were

washed, and if necessary, incubated for 20 more min with fluorochrome-

labeled Streptavidin: Streptavidin-AF647 (Jackson Immunoresearch) or

Streptavidin-APC/cy7, Streptavidin-BV421, Streptavidin-BV605 (BioLegend).

After the last wash, the cells were resuspended in FACS buffer containing

DAPI (BioLegend), Propidium Iodide (Sigma), or DRAQ7 (BioLegend) and

analyzed immediately on an LSR Fortessa flow cytometer running Diva 8

software (BD Biosciences). Typically, 500,000 cells were collected from

thymus samples. Data were analyzed using FlowJo software (TreeStar).

For intracellular staining, after surface antibody staining, the cells were

labeled with Zombie Aqua (BioLegend) for 30 min in ice. Then, the cells were

fixed with 2% paraformaldehyde (Electron Microscope Sciences) in PBS for

20 min on ice, permeabilized with either 0.5% Triton-X 100 (Sigma) for 20 min

on ice, or with Foxp3 staining kit (eBioscience) according to the protocol

provided by the manufacturer, and stained with antibodies for intracellular

markers for 40-60 min on ice.

For cell cycle analysis, 1-5x10^5 sorted thymic macrophages were fixed with

70% ethanol for 2 h on ice. The cells were spun down at 1800 rpm for 20 min

at 4°C, washed with PBS, and stained with 1 μg/ml DAPI (BioLegend) for 30

min at room temperature in the dark.

For EdU staining, after surface marker and Zombie Aqua staining, cells

were fixed with 2% paraformaldehyde in PBS for 20 min on ice and

permeabilized with 0.5% Triton X-100 in PBS at room temperature for 20 min.

EdU was detected by adding an equal volume of 2X Click reaction buffer

consisting of 200 mM Tris, 200 mM ascorbic acid (Acros), 8 mM CuSO$_4$

(Acros), 8 μM Cy5-azide (Lumiprobe) to the permeabilized cells resuspended

in 0.5% Triton X-100 in PBS and incubation at room temperature for 30 min.

Cells were washed twice with 0.5% Triton X-100 in PBS and analyzed on a

flow cytometer.

Cell sorting

The sorting of thymic macrophages was done following the IMMGEN

guidelines. Briefly, the thymuses of 3 male C57BL/6Narl mice were harvested

in ice-cold staining buffer containing phenol red-free DMEM (Gibco) with 10

mM HEPES (Sigma), 0.1% NaN$_3$, and 2% FBS (Gibco). Single-cell

suspensions were prepared as described in the Flow cytometry section.

Percoll PLUS was used to enrich mononuclear cells. The cells were

resuspended at 10^8/mL in staining buffer and labeled with appropriate

antibodies for 15 min in ice. To sort thymic macrophages, the cells were first

labeled with biotinylated antibodies to lineage markers (Lin) – CD3, CD8, Gr1,

B220. After washing, the cells were stained with antibodies to CD11b, F4/80,

CD45, CD64, and Streptavidin-APC/cy7 for 15 min in ice. For sorting thymus

XCR1$^+$ and SIRPα^+ cDCs, antibodies to XCR1, SIRPα, CD11c, MHC2, CD64,

and F4/80 were used. For sorting peritoneal cavity macrophages, antibodies

to ICAM2 and F4/80 were used. Immediately before sorting, the dead cells

were excluded with DRAQ7 or PI. For RNA Sequencing experiments, the

cells were double-sorted on FACS Melody, or Aria cell sorters (BD

Biosciences) and 1000 cells were directly deposited in TCL buffer (Qiagen),

frozen in dry ice and sent to IMMGEN for RNA sequencing. Four biological

replicates were prepared. For cytospin and cell cycle analysis, 1-5x10^5 cells

sorted on FACS Melody were collected in staining buffer.

Cytospin

Sorted cells were mounted on Superfrost PLUS slides (Thermo Scientific)

using a Cytospin centrifuge (Cytospin 3, Shandon) for 5 min at 500 rpm. Cells

were fixed with 2% paraformaldehyde for 10 min at room temperature and

stained with the Hemacolor Rapid Staining Kit (Merck Millipore). Images were

collected on BX61 upright microscope (Olympus) using 100X objective with

immersion oil and captured with a CCD camera. Images were then analyzed

and processed with ImageJ (NIH) and Adobe Photoshop 5.5 (Adobe).

In vitro phagocytosis assay

10^7 Thymocytes were cultured in cDMEM in the presence of 1 μM of

dexamethasone (Sigma) in a 3.5-cm culture dish at 37℃ in 5% CO_2 incubator

for 8 hours. Apoptosis levels were assessed by PI (Biolegend) and Annexin V-

FITC (Biolegend) staining. Typically, more than 80% of cells were Annexin V$^+$.

The dexamethasone-treated thymocytes were stained with 1 µg/mL pHrodo

Red, SE (ThermoFisher) in PBS for 30 min at room temperature. The cells

were washed two times with cDMEM and resuspended at $2x10^6$ cells/mL.

$4x10^4$ sorted peritoneal and thymic macrophages were stained with 5 µM

eFluor 450 (Thermo Fisher) in PBS for 10 min at 37℃, washed two times with

cDMEM and cultured in 96-well flat-bottom culture plate (Nunc) in 100 µL

cDMEM at 37℃ in 5% CO_2 incubator. After 3 hours of attachment, the non-

adherent cells were removed, and 200 µL ($4x10^5$) apoptotic thymocytes were

added to the macrophages. The cells were incubated at 37℃ in 5% CO_2

incubator for 2 hours. Fluorescent images were captured with AxioObserver 7

(Carl Zeiss) wide-field microscope equipped with Plan Apochromat 40x

NA=1.0 objective (Zeiss) and AxioCam 702 monochrome camera (Zeiss)

controlled by Zen 2.3 Blue (Zeiss) software. Image analysis was performed

with Imaris 8.0.2 (Bitplane). Phagocytosis was scored by investigators blinded

to the samples' identities.

In vitro antigen presentation assay

$3x10^4$ sorted thymic CD64$^-$MHCII$^+$CD11c$^+$ dendritic cells, thymic, or

peritoneal macrophages were cultured in 96-well round-bottom culture plate in

100 µL cDMEM at 37℃ in 5% CO_2 incubator for 3 hours to attach.

Splenocytes from OT2 mice were stained with biotinylated antibodies to

CD8a, CD11b, CD11c, B220, and MHCII (all from BioLegend), washed, and

labeled with anti-biotin Microbeads (Miltenyi) plus CD44 microbeads (Miltenyi)

in cRPMI. The cells were separated on MACS LS columns (Miltenyi)

according to the manufacturer's instructions. Enriched cells (naïve CD4 T

cells) were stained with 10 µM CFSE (Sigma) for 5 min in PBS at 37℃ and

cocultured with the sorted thymic MHCII⁺CD11c⁺ dendritic cells, thymic, or

peritoneal macrophages, in the presence or absence of 0.5 mg/mL OVA

protein (Sigma) in cRPMI at 37℃ in 5% CO_2 incubator for 72 hours. The cells

were collected and stained with antibodies to TCRβ and CD4 (from

BioLegend) for flow cytometry analyses of CFSE dilution. The data were

analyzed with FlowJo's Proliferation Modelling module (BD Biosciences).

RNA sequencing analysis

RNA sequencing was done at IMMGEN using Smart-seq2 protocol[84,85]

on a NextSeq500 sequencer (Illumina). Following sequencing, raw reads

were aligned with STAR to the mouse genome assembly mm10 and assigned

to specific genes using the GENCODE vM12 annotation. Gene expression

was normalized by DESeq2[86] and visualized by Morpheus

(https://software.broadinstitute.org/morpheus). The principal component

analysis was done by plotPCA() function of R package "DESeq2. Gene

expression of mouse transcription factors[87] was visualized in MultiplotStudio

of GenePattern[88]. GO enrichment was calculated and visualized in R by

using clusterProfiler[89].

Timed pregnancies and embryonic thymus analysis

To set up timed pregnancies, each male mouse (*Cx3cr1^{CreER/CreER}*,

Cx3cr1^{GFP/GFP} or C57BL/6) and female mouse (*ROSA26^{LSL-GFP/LSL-GFP}* or

C57BL/6) were housed together in the same cage for one night and separated

on the next day, which we defined as embryonic day 0.5 (E0.5). Female mice

were assumed to be pregnant if their weight gain was over 2 g at E8.5[90].

Thymuses from E14.5 and E17.5 embryos, neonatal, 1-weeks-old pups, and

adult mice (older than 6-weeks-old) were harvested, mechanically dissociated

with plastic sticks in 1.5-mL centrifuge tubes, and enzymatically digested with

0.2 mg/mL DNase I and 0.2 mg/mL collagenase P in complete DMEM for 20

min at 37°C with frequent agitation. The cells were resuspended in PBS with

0.5% BSA, filtered through a 70 μm filter, kept on ice, and used flow

cytometric analysis as described in the Flow Cytometry section.

Genetic fate mapping – E9.5, neonatal and adult

For genetic fate mapping, timed pregnancies of $Cx3cr1^{CreER/CreER}$ male and

$ROSA26^{LSL-GFP/LSL-GFP}$ female mice were set up as described. To label the

$Cx3cr1^+$ erythromyeloid progenitors derived from embryonic yolk sac[8], 4-

hydroxytamoxifen (4-OHT from Sigma) was administered i.p. to pregnant

females on E9.5 at a dose of 75 μg/g (body weight). To improve the survival

of embryos and reduce the risk of abortions, Progesterone (Sigma) was co-

injected at a dose of 37.5 μg/g (body weight)[91]. To label the $Cx3cr1^+$ thymic

macrophages in $Cx3cr1^{CreER}$ X $ROSA26^{LSL-GFP}$ neonates and adult mice,

Tamoxifen (TAM from Sigma) was injected i.p. at a dose of 2 mg/mouse to

lactating dams on postnatal day 3 and 4 (P3 and P4) or to adult mice for 2

consecutive days. Thymuses were harvested and analyzed 3 days or 6

weeks after the last injection by flow cytometry.

1006

scRNA-Seq – sorting, library generation, and sequencing

scRNA-Seq was performed at the Genomics Center for Clinical and

Biotechnological Applications of NCFB (NYCU, Taipei, Taiwan). Briefly, the

thymuses of one female MAFIA and 2 male $Cd11c^{YFP}$ mice were harvested

and enzymatically digested as described previously. Mononuclear cells were

enriched by 57% Percoll PLUS discontinuous centrifugation, washed to

remove silica particles, and resuspended at 10^6/mL in PBS with 0.04% BSA.

The cell suspensions were filtered through Falcon 35 µm strainer (Corning)

and stained with viability dye (PI or DAPI) immediately before sorting. Cell

sorting was performed on a FACS Melody sorter (BD Biosciences) running

FACS Chorus (BD Biosciences) software in purity mode. $3X10^5$ GFP or YFP

positive cells under the live/singlet gating were collected into 5 ml round

bottom tubes pre-coated with 0.04% BSA in PBS. Sorted cells were washed

and resuspended in 300 µL PBS with 0.04% BSA and then filtered again into

1.5-mL DNA LoBind tubes (Eppendorf) through a 35 µm strainer. The viability

of the cells was evaluated by Countess II (Invitrogen) and Trypan Blue

(ThermoFisher), and samples with cell viability rates higher than 85% were

used for encapsulation and library preparation. Single-cell encapsulation and

library preparation were performed using Single Cell 3' v3/v3.1 Gene

Expression solution (10x Genomics). All the libraries were processed

according to the manufacturer's instruction and sequenced on NovaSeq 6000

(Illumina) platform at the NHRI (Zhubei, Taiwan). Post-processing and quality

control were performed by the NYCU Genome Center using the CellRanger

package (v. 3.0.2, 10x Genomics). Reads were aligned to mm10 reference

assembly. Primary assessment with CellRanger reported 9,973 cell-barcodes

with 11,385 median unique molecular identifiers (UMIs, transcripts) per cell

and 3,076 median genes per cell sequenced to 71.0% sequencing saturation

with 94,260 mean reads per cell for MAFIA mouse sample; 9,801 cell-

barcodes with 13,467 median UMIs per cell and 3,211 median genes per cell

sequenced to 74.9% sequencing saturation with 119,820 mean reads per cell

for the first *Cd11c^{YFP}* mouse sample; 12,938 cell-barcodes with 14,439

median UMIs per cell and 3,199 median genes per cell sequenced to 71.4%

sequencing saturation with 108,585 mean reads per cell for the second

Cd11c^{YFP} mouse sample.

1041

Analysis of scRNA-Seq

Preprocessing

The Scanpy[92] pipeline was used to read the count matrix. Three batches

of samples (one from GFP$^+$ cells from MAFIA mouse and two from YFP$^+$ cells

from *Cd11c^{YFP}* mice) were preprocessed independently and integrated later.

Cells that expressed <200 genes and genes that were expressed in <3 cells

were filtered out. The percentage of mitochondrial genes was calculated -and

cells with >10% mitochondrial genes were removed. Cells with >7,000 genes

or <1,000 genes were also removed. Read counts were normalized to library

size 10,000 and log-transformed with scanpy.pp.log1p function.

Datasets integration and batch effect correction

Read count matrices and spliced/unspliced matrices were merged first.

Principal Component Analysis was applied to reduce dimensions to 70.

BBKNN[93] was then used to remove batch effects with the

scanpy.external.pp.bbknn function with default parameters.

Visualization and clustering

UMAP[94] provided by scanpy was used to visualize data with default

parameters. K-nearest neighbor and Leiden clustering were applied

sequentially to cluster cells into groups. K-nearest neighbor graph

construction was done by scanpy.pp.neighbors with parameters

n_neighbors=12 and n_pcs=70. Leiden clustering was then performed by

scanpy.tl.leiden with parameter resolution=0.15. To improve UMAP

visualization, scanpy.tl.paga was applied, and we trimmed unnecessary graph

edges by scanpy.tl.paga with threshold=0.018.

Marker genes and statistics

Wilcoxon rank-sum tests were applied to examine differentially expressed

genes. Clusters were selected from the result of Leiden clustering.

Differentially expressed genes of a cluster against other clusters were

identified by scanpy.tl.rank_genes_groups and scanpy.pl.rank_genes_groups.

P-values were collected for each cluster and transformed by negative $\log_{10}$ for

better visualization. The top 50 differentially expressed genes were visualized

in the figure.

1074

Immunofluorescent staining

Dissected thymus lobes from C57BL/6 mice were cleaned of connective

tissue and fixed in 4% paraformaldehyde (Sigma) for 1 h at 4°C, washed in

PBS, submerged in 10% sucrose, and then in 30% sucrose for 12 h each.

The tissue was then frozen in Tissue-Tek OCT compound (Sakura Fintek) for

cryostat sectioning. 10 or 20 µm thick sections were prepared with CryoStar

NX50 (ThermoFisher) on Superfrost PLUS (ThermoScientific) microscope

slides, dried overnight, and stored at -80°C until used. Before staining, the

sections were fixed with acetone (Sigma) at -20°C for 10 min, air-dried, then

blocked with 5% goat serum + 5% donkey serum (both from Jackson

Immunoresearch) in PBS for 2 h and stained with primary antibodies: rat

monoclonal to MerTK (DS5MMER, eBioscience), rat monoclonal to TIM4

(RMT4-54, Bio-X-Cell), rabbit polyclonal to CD64 (Sinobiological), or rabbit

polyclonal to Keratin 5 (BioLegend) overnight at 4°C in a humidified chamber.

After washing in PBS, the sections were labeled with goat anti-rat-Alexa Fluor

647 (Invitrogen) or goat anti-rat Cy3 (Jackson Immunoresearch) and donkey

anti-rabbit Cy3 or donkey anti-rabbit AF647 (both from Jackson

Immunoresearch) secondary antibodies for 2 hours at room temperature,

followed by 5 min staining with DAPI. TUNEL Assay was done with the Click-

iT Plus TUNEL Assay Alexa Fluor 647 kit (Invitrogen) according to the

manufacturer's recommendations. A positive (pre-incubation with DNase I for

30 min at room temperature) and negative (no TdT enzyme) controls were

always included. The sections were mounted with 0.1% n-propyl gallate

(Sigma) in glycerol (Sigma) and imaged with an AxioObserver 7 (Carl Zeiss)

wide-field microscope equipped with Plan Apochromat 20x NA=0.8 objective

(Zeiss) and AxioCam 702 mono camera (Zeiss) and controlled by Zen 2.3

Blue (Zeiss) software. Image analysis was performed with Imaris 8.0.2

(Bitplane).

1103

The co-localization scoring for MerTK and TIM4 with TUNEL was done with

Imaris 8.2 (Bitplane). TUNEL[+] cells were detected with the Spots function,

while MerTK[+] and TIM4[+] cells were detected with the Surface function. Spots

that co-localize with Surfaces were identified with the "Find Spots close to

Surface" function of Imaris XT. The threshold for co-localization was set to 5

µm. The results were manually curated so that Spots categorized as "not co-

localized" that were: 1) at the edge of the imaging field were excluded from

consideration; 2) with clear Surface signal around them were re-categorized

as "co-localized". The ratio of co-localized Spots to all Spots was calculated

and presented as the co-localization index.

1114

Thymus transplantation

To obtain E15.5 embryos, $SpiC^{GFP}$ (CD45.2) homozygous male and

congenic CD45.1 female mice were mated in a cage overnight and separated

on the next day. Pregnant mice were sacrificed 15 days later, the viable

embryos were harvested, and the thymuses were isolated in ice-cold PBS.

C57BL/6 recipients were anesthetized by i.p injection of Ketamine

hydrochloride (120 µg/g, Toronto Research Chemicals) and Xylazine

hydrochloride (12 µg/g, Sigma). The fur on the left flank was removed, and the

left kidney was exposed by cutting the skin, muscle layer, and peritoneum.

The kidney capsule was nicked with a G23 needle, and the fetal thymus was

pushed into the pocket under the kidney capsule with a G23 needle equipped

with a plunger from a spinal needle. After the kidney was re-positioned back

into the peritoneal cavity, the peritoneum was sutured, and the skin was

stapled with metal clips. Rymadil (Carprofen, 5 µg/g, Zoetis) was given

subcutaneously to ease the wound pain, and Trimerin (Sulfadiazine at 0.5

mg/mL + Trimethoprim at 0.1 mg/mL) were given in the drinking water for the

first two weeks after the surgery. The metal clips were removed from the skin

after the first week, and the transplanted thymus and recipient's endogenous

thymus were harvested and analyzed six weeks after the kidney

transplantation.

1135

1136 **Statistical analysis**

1137 Comparison between groups was made with Prism 6 (GraphPad Software).

1138 Comparisons between two groups were carried out with unpaired Student's t-

1139 test. When more than two groups were compared, a one-way ANOVA with

1140 Tukey correction was used. Differences were considered significant if $p<0.05$.

1141

Data availability

The RNA Sequencing data of thymic macrophages and thymic dendritic

cells are available at NCBI Gene Expression Omnibus (GEO) as part of

GSE122108 and at www.immgen.org. The single cell RNA sequencing data is

deposited at NCBI GEO under accession number GSE185460. The source

data underlying Fig. 1H and I, Fig. 3B, D, and G, Fig. 5C, F, and I, Fig. 6B, F,

and H, Fig. 7B, C, D, and G, Fig. 8B, D, E, and F, Fig. 1S5, Fig. 2S1, Fig. 2S2,

Fig. 2S3, Fig. 5S1, and Fig. 5S2 are provided in the Source Data files. All

other data supporting the findings of this study are available within the article

and its figures and tables.

1152

Abbreviations:

cDC – classical dendritic cell

DC – dendritic cell

EdU – 5-Ethynyl-2'-deoxyuridine

GO – gene ontology

HSC – hematopoietic stem cell

IFN-I – type I Interferon

IMMGEN – Immunological Genome Consortium

scRNA-Seq – single-cell RNA sequencing

TF – transcription factor

ThyMacs – thymic macrophages

TRA – tissue-restricted antigen

YS – yolk sac

References:

1.   Wynn TA, Chawla A, Pollard JW. Macrophage biology in development,
homeostasis and disease. Nature; 2013;496: 445–455.
doi:10.1038/nature12034

2.   Baratin M, Simon L, Jorquera A, Ghigo C, Dembele D, Nowak J, et al. T
Cell Zone Resident Macrophages Silently Dispose of Apoptotic Cells in
the Lymph Node. Immunity; 2017;47: 349–362.e5.
doi:10.1016/j.immuni.2017.07.019

3.   A-Gonzalez N, Castrillo A. Origin and Specialization of Splenic
Macrophages. Cellular Immunology. 2018;330: 1–27.
doi:10.1016/j.cellimm.2018.05.005

4.   Bellomo A, Mondor I, Spinelli L, Lagueyrie M, Stewart BJ, Brouilly N, et
al. Reticular Fibroblasts Expressing the Transcription Factor WT1
Define a Stromal Niche that Maintains and Replenishes Splenic Red
Pulp Macrophages. Immunity; 2020;53: 127–142.e7.
doi:10.1016/j.immuni.2020.06.008

5.   van Furth R, Cohn ZA. The origin and kinetics of mononuclear
phagocytes. J Exp Med; 1968;128: 415–435.
doi:10.1084/jem.128.3.415

6.   Ginhoux F, Guilliams M. Tissue-Resident Macrophage Ontogeny and
Homeostasis. Immunity. 2016;44: 439–449.
doi:10.1016/j.immuni.2016.02.024

7.   Perdiguero EG, Klapproth K, Schulz C, Busch K, Azzoni E, Crozet L, et
al. Tissue-resident macrophages originate from yolk-sac-derived

erythro-myeloid progenitors. Nat Immunol; 2015;518: 547–551.
doi:10.1038/nature13989

8. Mass E, Ballesteros I, Farlik M, Halbritter F, Günther P, Crozet L, et al.
Specification of tissue-resident macrophages during organogenesis.
Science; 2016;353: aaf4238–aaf4238. doi:10.1126/science.aaf4238

9. Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, et al. Fate
mapping analysis reveals that adult microglia derive from primitive
macrophages. Science. 2010;330: 841–845.
doi:10.1126/science.1194637

10. Hoeffel G, Wang Y, Greter M, See P, Teo P, Malleret B, et al. Adult
Langerhans cells derive predominantly from embryonic fetal liver
monocytes with a minor contribution of yolk sac–derived macrophages.
J Exp Med. 2012;209: 1167–1181. doi:10.1084/jem.20120340

11. Hoeffel G, Chen J, Lavin Y, Low D, Almeida FF, See P, et al. C-Myb(+)
erythro-myeloid progenitor-derived fetal monocytes give rise to adult
tissue-resident macrophages. Immunity. 2015;42: 665–678.
doi:10.1016/j.immuni.2015.03.011

12. Goldmann T, Wieghofer P, Jordão MJC, Prutek F, Hagemeyer N,
Frenzel K, et al. Origin, fate and dynamics of macrophages at central
nervous system interfaces. Nat Immunol. 2016;17: 797–805.
doi:10.1038/ni.3423

13. Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, et al.
Tissue-Resident Macrophages Self-Maintain Locally throughout Adult
Life with Minimal Contribution from Circulating Monocytes. Immunity;
2013;38: 792–804. doi:10.1016/j.immuni.2013.04.004

14. Epelman S, Lavine KJ, Beaudin AE, Sojka DK, Carrero JA, Calderon B,
et al. Embryonic and Adult-Derived Resident Cardiac Macrophages Are
Maintained through Distinct Mechanisms at Steady State and during
Inflammation. Immunity; 2014;40: 91–104.
doi:10.1016/j.immuni.2013.11.019

15. Sheng J, Ruedl C, Karjalainen K. Most Tissue-Resident Macrophages
Except Microglia Are Derived from Fetal Hematopoietic Stem Cells.
Immunity; 2015;43: 382–393. doi:10.1016/j.immuni.2015.07.016

16. Liu Z, Gu Y, Chakarov S, Bleriot C, Kwok I, Chen X, et al. Fate Mapping
via Ms4a3-Expression History Traces Monocyte-Derived Cells. Cell;
2019;178: 1509–1525.e19. doi:10.1016/j.cell.2019.08.009

17. Ensan S, Li A, Besla R, Degousee N, Cosme J, Roufaiel M, et al. Self-
renewing resident arterial macrophages arise from embryonic
CX3CR1+ precursors and circulating monocytes immediately after birth.
Nat Immunol; 2015;17: 159–168. doi:10.1038/ni.3343

18. Mondor I, Baratin M, Lagueyrie M, Saro L, Henri S, Gentek R, et al.

Lymphatic Endothelial Cells Are Essential Components of the
Subcapsular Sinus Macrophage Niche. Immunity; 2019;50: 1453–
1466.e4. doi:10.1016/j.immuni.2019.04.002

19.   Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et
al. Progressive replacement of embryo-derived cardiac macrophages
with age. J Exp Med. 2014;211: 2151–2158. doi:10.1084/jem.20140639

20.   Heidt T, Courties G, Dutta P, Sager HB, Sebas M, Iwamoto Y, et al.
Differential Contribution of Monocytes to Heart Macrophages in Steady-
State and After Myocardial Infarction. Circulation Research. 2014;115:
284–295. doi:10.1161/CIRCRESAHA.115.303567

21.   Yahara Y, Barrientos T, Tang YJ, Puviindran V, Nadesan P, Zhang H,
et al. Erythromyeloid progenitors give rise to a population of osteoclasts
that contribute to bone homeostasis and repair. Nat Cell Biol; 2020;22:
1–28. doi:10.1038/s41556-019-0437-8

22.   Jacome-Galarza CE, Percin GI, Muller JT, Mass E, Lazarov T, Eitler J,
et al. Developmental origin, functional maintenance and genetic rescue
of osteoclasts. Nature; 2019;568: 541–545. doi:10.1038/s41586-019-
1105-7

23.   Calderon B, Carrero JA, Ferris ST, Sojka DK, Moore L, Epelman S, et
al. The pancreas anatomy conditions the origin and properties of
resident macrophages. J Exp Med. 2015;212: 1497–1512.
doi:10.1084/jem.20150496

24.   Tamoutounour S, Guilliams M, Sanchis FM, Liu H, Terhorst D, Malosse
C, et al. Origins and Functional Specialization of Macrophages and of
Conventional and Monocyte-Derived Dendritic Cells in Mouse Skin.
Immunity; 2013;39: 925–938. doi:10.1016/j.immuni.2013.10.004

25.   Bain CC, Bravo-Blas A, Scott CL, Gomez Perdiguero E, Geissmann F,
Henri S, et al. Constant replenishment from circulating monocytes
maintains the macrophage pool in the intestine of adult mice. Nat
Immunol. 2014;15: 929–937. doi:10.1038/ni.2967

26.   Sierro F, Evrard M, Rizzetto S, Melino M, Mitchell AJ, Florido M, et al. A
Liver Capsular Network of Monocyte-Derived Macrophages Restricts
Hepatic Dissemination of Intraperitoneal Bacteria by Neutrophil
Recruitment. Immunity; 2017;47: 374–388.e6.
doi:10.1016/j.immuni.2017.07.018

27.   Mossadegh-Keller N, Gentek R, Gimenez G, Bigot S, Mailfert S,
Sieweke MH. Developmental origin and maintenance of distinct
testicular macrophage populations. J Exp Med. 2017;214: 2829–2841.
doi:10.1084/jem.20170829

28.   Lokka E, Lintukorpi L, Cisneros-Montalvo S, Mäkelä J-A, Tyystjärvi S,
Ojasalo V, et al. Generation, localization and functions of macrophages

during the development of testis. Nature Communications. Springer US;
2020;11: 1–16. doi:10.1038/s41467-020-18206-0

29. Wang M, Yang Y, Cansever D, Wang Y, Kantores C, Messiaen S, et al.
Two populations of self-maintaining monocyte-independent
macrophages exist in adult epididymis and testis. PNAS. 2020;118:
e2013686117. doi:10.1073/pnas.2013686117/-/DCSupplemental

30. Bain CC, Hawley CA, Garner H, Scott CL, Schridde A, Steers NJ, et al.
Long-lived self-renewing bone marrow-derived macrophages displace
embryo-derived cells to inhabit adult serous cavities. Nature
Communications; 2016;7: 1–14. doi:10.1038/ncomms11852

31. Surh CD, Sprent J. T-cell apoptosis detected in situ during positive and
negative selection in the thymus. Nature. 1994;372: 100–103.
doi:10.1038/372100a0

32. Esashi E, Sekiguchi T, Ito H, Koyasu S, Miyajima A. Cutting Edge: A
possible role for CD4+ thymic macrophages as professional scavengers
of apoptotic thymocytes. Journal of Immunology; 2003;171: 2773–2777.
doi:10.4049/jimmunol.171.6.2773

33. Liu L-T, Lang Z-F, Li Y, Zhu Y-J, Zhang J-T, Guo S-F, et al.
Composition and characteristics of distinct macrophage subpopulations
in the mouse thymus. Molecular Medicine Reports. 2013;7: 1850–1854.
doi:10.3892/mmr.2013.1448

34. Guerri L, Peguillet I, Geraldo Y, Nabti S, Premel V, Lantz O. Analysis of
APC types involved in CD4 tolerance and regulatory T cell generation
using reaggregated thymic organ cultures. J Immunol; 2013;190: 2102–
2110. doi:10.4049/jimmunol.1202883

35. Lopes N, Charaix J, Cédile O, Sergé A, Irla M. Lymphotoxin alpha fine-
tunes T cell clonal deletion by regulating thymic entry of antigen-
presenting cells. Nature Communications; 2018;9: 1–16.
doi:10.1038/s41467-018-03619-9

36. Kim H-J, Alonzo ES, Dorothee G, Pollard JW, Sant'Angelo DB.
Selective depletion of eosinophils or neutrophils in mice impacts the
efficiency of apoptotic cell clearance in the thymus. PLoS ONE. 2010;5:
e11439. doi:10.1371/journal.pone.0011439.g010

37. Tacke R, Hilgendorf I, Garner H, Waterborg C, Park K, Nowyhed H, et
al. The transcription factor NR4A1 is essential for the development of a
novel macrophage subset in the thymus. Sci Rep. 2015;5: 10055.
doi:10.1038/srep10055

38. Gautier EL, Shay T, Miller J, Greter M, Jakubzick C, Ivanov S, et al.
Gene-expression profiles and transcriptional regulatory pathways that
underlie the identity and diversity of mouse tissue macrophages. Nat
Immunol. 2012;13: 1118–1128. doi:10.1038/ni.2419

39.   Ingersoll MA, Spanbroek R, Lottaz C, Gautier EL, Frankenberger M,
1314   Hoffmann R, et al. Comparison of gene expression profiles between
1315   human and mouse monocyte subsets. Blood. 2010;115: e10–e19.
1316   doi:10.1182/blood-2009-07-235028

1317   40.   Wang H, Breed ER, Lee YJ, Qian LJ, Jameson SC, Hogquist KA.
1318   Myeloid cells activate iNKT cells to produce IL-4 in the thymic medulla.
1319   Proc Natl Acad Sci; 2019;116: 22262–22268.
1320   doi:10.1073/pnas.1910412116

1321   41.   Chan CT, Fenn AM, Harder NK, Mindur JE, McAlpine CS, Patel J, et al.
1322   Liver X receptors are required for thymic resilience and T cell output. J
1323   Exp Med. 2020;217: 245–23. doi:10.1084/jem.20200318

1324   42.   Wada H, Masuda K, Satoh R, Kakugawa K, Ikawa T, Katsura Y, et al.
1325   Adult T-cell progenitors retain myeloid potential. Nature; 2008;452: 768–
1326   772. doi:10.1038/nature06839

1327   43.   Bell JJ, Bhandoola A. The earliest thymic progenitors for T cells
1328   possess myeloid lineage potential. Nature. 2008;452: 764–767.
1329   doi:10.1038/nature06840

1330   44.   Schlenner SM, Madan V, Busch K, Tietz A, LAufle C, Costa C, et al.
1331   Fate Mapping Reveals Separate Origins of T Cells and Myeloid
1332   Lineages in the Thymus. Immunity; 2010;32: 426–436.
1333   doi:10.1016/j.immuni.2010.03.005

1334   45.   Witmer-Pack MD, Hughes DA, Schuler G, Lawson L, McWilliam A,
1335   Inaba K, et al. Identification of macrophages and dendritic cells in the
1336   osteopetrotic (op/op) mouse. J Cell Sci. 1993;104 (Pt 4): 1021–1029.

1337   46.   Sasmono RT, Oceandy D, Pollard JW, Tong W, Pavli P, Wainwright BJ,
1338   et al. A macrophage colony-stimulating factor receptor-green
1339   fluorescent protein transgene is expressed throughout the mononuclear
1340   phagocyte system of the mouse. Blood; 2003;101: 1155–1163.
1341   doi:10.1182/blood-2002-02-0569

1342   47.   Hume DA. Applications of myeloid-specific promoters in transgenic mice
1343   support in vivo imaging and functional genomics but do not support the
1344   concept of distinct macrophage and dendritic cell lineages or roles in
1345   immunity. Journal of Leukocyte Biology. 2011;89: 525–538.
1346   doi:10.1189/jlb.0810472

1347   48.   Taniuchi I, Osato M, Egawa T, Sunshine MJ, Bae SC, Komori T, et al.
1348   Differential requirements for Runx proteins in CD4 repression and
1349   epigenetic silencing during T lymphocyte development. Cell. 2002;111:
1350   621–633. doi:10.1016/s0092-8674(02)01111-x

1351   49.   Ebihara T, Song C, Ryu SH, Plougastel-Douglas B, Yang L, Levanon D,
1352   et al. Runx3 specifies lineage commitment of innate lymphoid cells. Nat
1353   Immunol; 2015;16: 1124–1133. doi:10.1038/ni.3272

50.   Fainaru O, Woolf E, Lotem J, Yarmus M, Brenner O, Goldenberg D, et
al. Runx3 regulates mouse TGF-beta-mediated dendritic cell function
and its absence results in airway inflammation. EMBO J; 2004;23: 969–
979. doi:10.1038/sj.emboj.7600085

51.   Kohyama M, Ise W, Edelson BT, Wilker PR, Hildner K, Mejia C, et al.
Role for Spi-C in the development of red pulp macrophages and splenic
iron homeostasis. Nat Immunol; 2008;457: 318–321.
doi:10.1038/nature07472

52.   Haldar M, Kohyama M, So AY-L, KC W, Wu X, Briseño CG, et al.
Heme-Mediated SPI-C Induction Promotes Monocyte Differentiation into
Iron-Recycling Macrophages. Cell; 2014;156: 1223–1234.
doi:10.1016/j.cell.2014.01.069

53.   Yona S, Kim K-W, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate
Mapping Reveals Origins and Dynamics of Monocytes and Tissue
Macrophages under Homeostasis. Immunity; 2013;38: 79–91.
doi:10.1016/j.immuni.2012.12.001

54.   McCaughtry TM, Wilken MS, Hogquist KA. Thymic emigration revisited.
J Exp Med. 2007;204: 2513–2520. doi:10.1084/jem.20070601

55.   Lienenklaus S, Cornitescu M, Ziętara N, Łyszkiewicz M, Gekara N,
Jabłónska J, et al. Novel reporter mouse reveals constitutive and
inflammatory expression of IFN-beta in vivo. J Immunol; 2009;183:
3229–3236. doi:10.4049/jimmunol.0804277

56.   Otero DC, Baker DP, David M. IRF7-dependent IFN-β production in
response to RANKL promotes medullary thymic epithelial cell
development. J Immunol; 2013;190: 3289–3298.
doi:10.4049/jimmunol.1203086

57.   Breed ER, Lee ST, Hogquist KA. Directing T cell fate: How thymic
antigen presenting cells coordinate thymocyte selection. Seminars in
Cell and Developmental Biology; 2017;: 1–9.
doi:10.1016/j.semcdb.2017.07.045

58.   Kurd NS, Lutes LK, Yoon J, Chan SW, Dzhagalov IL, Hoover AR, et al.
A role for phagocytosis in inducing cell death during thymocyte negative
selection. eLife. 2019;8: 1879–18. doi:10.7554/eLife.48097

59.   Morioka S, Perry JSA, Raymond MH, Medina CB, Zhu Y, Zhao L, et al.
Efferocytosis induces a novel SLC program to promote glucose uptake
and lactate release. Nature; 2018;563: 1–21. doi:10.1038/s41586-018-
0735-5

60.   Bleriot C, Chakarov S, Ginhoux F. Determinants of Resident Tissue
Macrophage Identity and Function. Immunity; 2020;52: 957–970.
doi:10.1016/j.immuni.2020.05.014

61.   Stritesky GL, Xing Y, Erickson JR, Kalekar LA, Wang X, Mueller DL, et

al. Murine thymic selection quantified using a unique method to capture
deleted T cells. Proc Natl Acad Sci. 2013;110: 4679–4684.
doi:10.1073/pnas.1217532110

62.   Baba T, Nakamoto Y, Mukaida N. Crucial Contribution of Thymic Sirp +
Conventional Dendritic Cells to Central Tolerance against Blood-Borne
Antigens in a CCR2-Dependent Manner. J Immunol. 2009;183: 3053–
3063. doi:10.4049/jimmunol.0900438

63.   Gallegos AM, Bevan MJ. Central Tolerance to Tissue-specific Antigens
Mediated by Direct and Indirect Antigen Presentation. J Exp Med.
2004;200: 1039–1049. doi:10.1084/jem.20041457

64.   Koble C, Kyewski B. The thymic medulla: a unique microenvironment
for intercellular self-antigen transfer. J Exp Med. 2009;206: 1505–1513.
doi:10.1084/jem.20082449

65.   Vobořil M, Brabec T, Dobeš J, Šplíchalová I, Březina J, Čepková A, et
al. Toll-like receptor signaling in thymic epithelium controls monocyte-
derived dendritic cell recruitment and Treg generation. Nature
Communications; 2020;11: 1–16. doi:10.1038/s41467-020-16081-3

66.   Dixit VD. Impact of immune-metabolic interactions on age-related
thymic demise and T cell senescence. Semin Immunol; 2012;24: 321–
330. doi:10.1016/j.smim.2012.04.002

67.   Rivera-Gonzalez GC, Shook BA, Andrae J, Holtrup B, Bollag K,
Betsholtz C, et al. Skin Adipocyte Stem Cell Self-Renewal Is Regulated
by a PDGFA/AKT-Signaling Axis. Stem Cell; 2016;19: 738–751.
doi:10.1016/j.stem.2016.09.002

68.   Olson LE, Soriano P. Increased PDGFRα Activation Disrupts
Connective Tissue Development and Drives Systemic Fibrosis.
Developmental Cell; 2009;16: 303–313.
doi:10.1016/j.devcel.2008.12.003

69.   Satpathy AT, KC W, Albring JC, Edelson BT, Kretzer NM, Bhattacharya
D, et al. Zbtb46expression distinguishes classical dendritic cells and
their committed progenitors from other immune lineages. J Exp Med.
2012;209: 1135–1152. doi:10.1084/jem.20120030

70.   Guilliams M, Mildner A, Yona S. Developmental and Functional
Heterogeneity of Monocytes. Immunity; 2018;49: 595–613.
doi:10.1016/j.immuni.2018.10.005

71.   Tabula Muris Consortium. Single-cell transcriptomics of 20 mouse
organs creates a Tabula Muris. Nature; 2018;562: 367–372.
doi:10.1038/s41586-018-0590-4

72.   Han X, Wang R, Zhou Y, Fei L, Sun H, Lai S, et al. Mapping the Mouse
Cell Atlas by Microwell-Seq. Cell; 2018;172: 1091–1097.e17.
doi:10.1016/j.cell.2018.02.001

73. Tabula Muris Consortium. A single-cell transcriptomic atlas
characterizes ageing tissues in the mouse. Nature; 2020;583: 590–595.
doi:10.1038/s41586-020-2496-1

74. Kernfeld EM, Genga RMJ, Neherin K, Magaletta ME, Xu P, Maehr R. A
Single-Cell Transcriptomic Atlas of Thymus Organogenesis Resolves
Cell Types and Developmental Maturation. Immunity; 2018;: 1–33.
doi:10.1016/j.immuni.2018.04.015

75. Park J-E, Botting RA, Domínguez Conde C, Popescu D-M, Lavaert M,
Kunz DJ, et al. A cell atlas of human thymic development defines T cell
repertoire formation. Science; 2020;367: eaay3224–12.
doi:10.1126/science.aay3224

76. Bornstein C, Nevo S, Giladi A, Kadouri N, Pouzolles M, Gerbe F, et al.
Single-cell mapping of the thymic stroma identifies IL-25-producing tuft
epithelial cells. Nature; 2018;559:1–25. doi:10.1038/s41586-018-0346-1

77. Bautista JL, Cramer NT, Miller CN, Chavez J, Berrios DI, Byrnes LE, et
al. Single-cell transcriptional profiling of human thymic stroma uncovers
novel cellular heterogeneity in the thymic medulla. Nature
Communications; 2021;12: 1096–15. doi:10.1038/s41467-021-21346-6

78. Burnett SH, Kershen EJ, Zhang J, Zeng L, Straley SC, Kaplan AM, et
al. Conditional macrophage ablation in transgenic mice expressing a
Fas-based suicide gene. Journal of Leukocyte Biology; 2004;75: 612–
623. doi:10.1189/jlb.0903442

79. Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A,
et al. Analysis of fractalkine receptor CX(3)CR1 function by targeted
deletion and green fluorescent protein reporter gene insertion. Mol Cell
Biol. 2000;20: 4106–4114.

80. Lindquist RL, Shakhar G, Dudziak D, Wardemann H, Eisenreich T,
Dustin ML, et al. Visualizing dendritic cell networks in vivo. Nat
Immunol. 2004;5: 1243–1250. doi:10.1038/ni1139

81. Faust N, Varas F, Kelly LM, Heck S, Graf T. Insertion of enhanced
green fluorescent protein into the lysozyme gene creates mice with
green fluorescent granulocytes and macrophages. Blood. 2000;96:
719–726.

82. Roesch K, Jadhav AP, Trimarchi JM, Stadler MB, Roska B, Sun BB, et
al. The transcriptome of retinal Müller glial cells. J Comp Neurol.
2008;509: 225–238. doi:10.1002/cne.21730

83. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et
al. A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. Nature Neuroscience; 2009;13: 133–
140. doi:10.1038/nn.2467

84. Picelli S, Faridani OR, rklund ASKBO, Winberg GOS, Sagasser S,

Sandberg R. Full-length RNA-seq from single cells using Smart-seq2.
Nature Protocols; 2014;9: 171–181. doi:10.1038/nprot.2014.006

85. Picelli S, Björklund ÅK, Faridani OR, Sagasser S, Winberg G, Sandberg
R. Smart-seq2 for sensitive full-length transcriptome profiling in single
cells. Nat Meth. 2013;10: 1096–1098. doi:10.1038/nmeth.2639

86. Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. BioMed
Central; 2014;15: 31–21. doi:10.1186/s13059-014-0550-8

87. Schmeier S, Alam T, Essack M, Bajic VB. TcoF-DB v2: update of the
database of human and mouse transcription co-factors and transcription
factor interactions. Nucleic Acids Res. 2017;45: D145–D150.
doi:10.1093/nar/gkw1007

88. Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP.
GenePattern 2.0. Nat Genet; 2006;38: 500–501. doi:10.1038/ng0506-
500

89. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. OMICS: A
Journal of Integrative Biology; 2012;16: 284–287.
doi:10.1089/omi.2011.0118

90. Heyne GW, Plisch EH, Melberg CG, Sandgren EP, Peter JA, Lipinski
RJ. A Simple and Reliable Method for Early Pregnancy Detection in
Inbred Mice. J Am Assoc Lab Anim Sci. 2015;54: 368–371.

91. Iturri L, Saenz Coronilla J, Lallemand Y, Gomez Perdiguero E.
Identification Of Erythromyeloid Progenitors And Their Progeny In The
Mouse Embryo By Flow Cytometry. JoVE. 2017;: 1–10.
doi:10.3791/55305

92. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol; 2018;19: 15–5.
doi:10.1186/s13059-017-1382-0

93. Polański K, Young MD, Miao Z, Meyer KB, Teichmann SA, Park J-E.
BBKNN: fast batch alignment of single cell transcriptomes.
Bioinformatics. 2019;36: 964–965.
doi:10.1093/bioinformatics/btz625#supplementary-data

94. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation
and Projection for Dimension Reduction. ArXiv e-prints. 2018.
doi:1802.03426v3

1513

Acknowledgments

We are grateful to the IMMGEN consortium for performing the RNA

sequencing of our samples and providing access to its database. We thank

the following core facilities at National Yang Ming Chiao Tung University:

Instrumentation Resource Center for providing access to its sorting and

imaging facility and the Animal Facility for mouse husbandry. We are very

grateful to Wan-Chun Chang, Bing-Xiu Guo, Chien-Yi Tung, and Kate Hua

from the Genomics Center for Clinical and Biotechnological Applications of

NCFB (NYCU, Taipei) for help with scRNA-Seq. Special thanks go to Dr.

Fang Liao, Academia Sinica, Taipei, for help with cell sorting and for providing

the 24G2 hybridoma, and Chang-Feng Chu for technical assistance. We are

very grateful to Alexandra Norris, Cecilia Tai, and Angelina Shih for the

manual scoring of the in vitro phagocytosis experiments. We are indebted to

Dr. Yu-Wen Su (NHRI, Taiwan) for help with the mice. Some figures were

made with BioRender.

This work was supported by the Ministry of Science and Technology,

Taiwan (grants # 107-2320-B-010 -016 -MY3, 110-2320-B-A49A-521 - , and

111-2320-B-A49 -031 -MY3 to IL Dzhagalov), the Yen-Tjing Ling Medical

Foundation (grant # CI-111-6 to IL Dzhagalov), and the Cancer Progression

Research Center (NYCU). The authors declare no competing financial

interests.

Author contributions

TA Zhou designed experiments, performed research, analyzed and

interpreted data, and wrote the manuscript; HP Hsu, HK Cheng, and CY Lin

performed research, analyzed and interpreted data; YH Tu and HC Huang

analyzed the scRNA-Seq data; NJ Chen, JW Tsai, EA Robey, and CL Hsu

provided expertise, shared critical reagents, and edited the manuscript; IL

Dzhagalov conceptualized the studies, designed experiments, performed

research, analyzed, interpreted the data, and wrote the manuscript.

1543

1544

1545   **Supplementary figures and tables**

1546

1547   **Figure 1 – figure supplement 1:** Representative flow cytometry staining of

1548   enzymatically digested thymus single-cell suspension for CD64, MerTK, and

1549   F4/80 and respective isotype controls. The flow cytometry plots are

1550   representative of 5 individual experiments.

1551

1552

1553 **Figure 1 – figure supplement 2:** Immunofluorescent images of thymic

1554 sections showing co-localization of MerTK and CD64 staining (upper row) and

1555 TIM4 and CD64 staining (lower row) in the thymic cortex. The images are

1556 representative of at least three mice. The scale bar is 50 μm.

1557

1558

1559 **Figure 1 – figure supplement 3:** UMAP clustering of the scRNA-Seq data

1560 shows that the cells from the three samples (one from GFP[+] cells in MAFIA

1561 mice and two from YFP[+] cells in *Cd11c^{YFP}* mice) overlap considerably.

1562

1563

1564 **Figure 1 – figure supplement 4:** Expression of CD11c, MHC2, and SIRPα

1565 on ThyMacs with respective controls. The flow cytometry plots are

1566 representative of 5 individual experiments.

1567

1568

1569   **Figure 1 – figure supplement 5:** Example flow cytometry plots showing

1570   that gating on CD11c⁺MHC2⁺ thymus cells, in addition to DCs, also includes

that gating on CD11c$^+$MHC2$^+$ thymus cells, in addition to DCs, also includes

macrophages, especially among SIRPα^+ cells. On the right is a plot of the

frequency of MerTK$^+$ cells among CD11c$^+$MHC2$^+$SIRPα^+ cells. The data are

mean±SEM from 5 individual mice. Each dot is an individual mouse. The

numbers in the flow cytometry plots are the percent of cells in the respective

gate.

1576

1577

1578   **Figure 2 – figure supplement 1:** Example of the gating strategy to identify

1579   ThyMacs among *Spic^{GFP+}* cells. On the right is a plot showing the mean±SEM

1580   of the frequencies of ThyMacs among *Spic^{GFP+}* cells. Each dot is an individual

1581   mouse. The numbers in the flow cytometry plots are the percent of cells in the

1582   respective gate.

1583

1584

1585   **Figure 2 – figure supplement 2:** Representative flow cytometry plots of

1586   the expression of four reporter alleles in ThyMacs (left), frequencies of

1587   GFP/YFP⁺ cells among ThyMacs (middle), and frequencies of ThyMacs

1588   among GFP/YFP⁺ cells (right). Data in the graphs represent mean±SEM.

1589   Each dot is an individual mouse.

1590

1591

1592   **Figure 2 – figure supplement 3:** Representative flow cytometry plots of

1593   the expression of $Spic^{GFP}$ in ThyMacs. To the right is a graph showing the

1594   mean±SEM of the frequencies of $Spic^{GFP+}$ cells among ThyMacs. Each dot is

1595   an individual mouse. The numbers in the flow cytometry plots are the percent

1596   of cells in the respective gate.

1597

1598   **Figure 4 figure supplement 1:** Representative immunofluorescent image

1599 of WT thymic frozen section stained for TIM4, CD64, and UEA-1 (a marker for

1600 medulla). CD64 stains all macrophages, while TIM4 – only a subset that is

1601 located in the cortex. The image is representative of 3 mice. The scale bar is

1602 400 µm.

1603

Figure 5 – figure supplement 1: Representative flow cytometry plots of

donor (CD45.2^{+}) vs. host (CD45.1^{+}) derived thymocytes in the transplanted

thymus. The host thymus (endogenous thymus) serves as a negative control.

To the right is a graph showing the mean±SEM of the frequencies of CD45.2^{+}

(donor-derived) cells among thymocytes in the transplanted and endogenous

thymuses of the mice. Each dot is an individual mouse. The numbers in the

flow cytometry plots are the percent of cells in the respective

gate.

1612

Figure 5 – figure supplement 2: Comparison of the geometric mean

fluorescent intensities (gMFI) of SpiC^{GFP} TIM4$^+$ and TIM4$^-$ thymic

macrophages of host and donor-origin in the transplanted thymus. The

endogenous thymus serves as background control. Only populations

comprising >5% of thymic macrophages are shown. Each dot is an individual

mouse. The data are presented as mean±SEM.

Table 1: Expression of differentially up-regulated transcription factors in thymic macrophages

Gene name	ThyMacs	non-ThyMacs
Irf7	3879.32	300.82
Irf8	3528.27	1474.35
Stat1	2403.69	522.04
Dnmt3a	1515.94	647.81
Znxf1	1379.89	635.36
Stat2	1210.35	472.53
Nr1h3	1182.17	147.05
Srebf1	975.09	399.06
Rxra	760.26	298.55
Trps1	746.36	232.48
Runx3	723.14	9.76
Relb	715.53	293.92
Sp100	696.94	324.47
Zbp1	639.19	69.83
Tfec	588.72	74.66
Spic	573.11	34.36
Nfkbie	569.74	226.76

Ncoa4	550.69	249.15
Rest	548.22	269.22
Meis3	530.8	120.91
Bhlhe40	490.59	99.56
Parp12	414.11	126.82
Arid5b	374.03	177.08
Creb5	295.14	47.91
Pparg	276.54	33.24

Table 2: Expression of differentially up-regulated transcription factors

in thymic macrophages. Transcription factors that were highly expressed in

thymic macrophages (>250) and up-regulated >2-fold in thymic macrophages

compared to non-thymic macrophages were listed alphabetically, and the

geometric means of 4 replicates of thymic macrophages (ThyMacs) and two

replicates of each of the 9 non-thymic macrophage populations (non-

ThyMacs) were recorded. Non-thymic macrophages are: spleen red pulp

macrophages, Kupffer cells, broncho-alveolar lavage macrophages,

peritoneal cavity macrophages, aorta macrophages, heart macrophages,

white adipose tissue macrophages, central nervous system microglia, spinal

cord macrophages.

Table 2: List of the differentially expressed genes among *Timd*4+ thymic macrophages, *Cx3cr1*+ thymic macrophages, and thymic monocytes

Cx3cr1+ ThyMacs		*Timd4*+ ThyMacs		ThyMonos	
Gene name	adjusted p-value	Gene name	adjusted p-value	Gene name	adjusted p-value
Ctsz	0	Hpgd	0	Alox5ap	0
Cd63	0	Serpinb6a	0	S100a6	0
Pmepa1	0	Slc40a1	0	Ly6c2	0
Zmynd15	0	Cd81	0	Ifi27l2a	0
Olfml3	0	Vcam1	0	Fau	0
Mmp2	0	Cfp	0	Coro1a	0
AU020206	1.60E-290	Spic	0	Ccr2	0
Plxnd1	1.59E-285	Trf	0	Rps27	0
Cst7	8.68E-279	Actn1	0	Tmsb10	0
Dnase1l3	2.45E-270	Maf	0	Ifitm2	7.21E-302
Timp2	2.15E-267	Pld3	0	Fxyd5	6.36E-299

Lgals3bp	8.69E-263	Il18	0	Rps19	2.04E-292
Pdgfa	6.87E-255	Mrc1	0	Rpl18	6.50E-291
Mmp14	2.33E-253	Crip2	0	Rpl9	1.11E-289
Fam46c	9.99E-235	Tmem65	0	Rps23	1.28E-289
Chst2	1.19E-226	Igf1	0	Napsa	8.91E-279
Cp	5.36E-225	Epb41l3	0	Ms4a4c	8.25E-272
Camk1	7.12E-225	Timd4	0	Plac8	2.10E-270
B2m	1.09E-222	Blvrb	0	Rpl18a	9.26E-269
Lhfpl2	4.52E-217	Clec1b	0	S100a4	4.98E-268
Acp5	5.90E-216	Cd68	0	Cd52	3.67E-267
Lag3	3.91E-213	Axl	0	Rps14	1.94E-266
Lyz2	1.28E-209	Paqr9	3.32E-307	Ifitm3	3.19E-263
H2-M2	1.22E-199	Sdc3	3.45E-305	Rpl34	2.02E-261
Psap	7.26E-198	Myo9a	5.59E-305	Rps27a	3.67E-260
Gatm	1.33E-192	Scp2	3.79E-302	Rpl36	1.54E-259
Cpd	1.50E-192	Selenop	2.10E-295	Rps16	2.55E-258
C3	2.34E-187	Lrp1	2.08E-294	Rpl24	1.37E-257
Cxcl16	8.11E-183	Lap3	1.45E-290	Rps9	6.34E-253
Lgals3	1.57E-182	Marcks	2.77E-279	Gpr141	1.21E-246
Ube2j1	1.63E-180	Glul	3.64E-279	Rpl27a	3.06E-243
Plxnc1	9.84E-180	Hebp1	3.76E-278	Rpl17	8.15E-241
Stab1	4.07E-176	Ear2	4.53E-276	Rps24	1.46E-240
Cyth1	3.27E-163	Apoc1	2.49E-275	Rps13	2.34E-236
Spsb1	3.96E-163	Kcna2	3.72E-275	Rpl38	1.95E-226
Blnk	2.35E-162	Myo10	9.05E-269	H2-DMb1	1.02E-223
Cx3cr1	9.29E-162	Atp13a2	2.95E-267	Rps18	5.39E-223
Med10	5.25E-161	Slc1a3	6.24E-263	Rpl19	3.68E-221
Nek6	5.28E-160	Slco2b1	1.11E-258	Rpl8	2.01E-219
Ptms	1.05E-159	mt-Nd2	3.45E-258	Rpl7a	4.17E-217
Anxa5	1.10E-156	Wwp1	2.16E-253	Gm34084	5.23E-216
Gpnmb	1.21E-154	Aplp2	4.22E-248	Rpl13	2.08E-215
Itgb5	2.78E-154	Atp8a1	5.03E-248	Rpl11	2.47E-213
Myo5a	1.11E-146	P2ry13	3.17E-247	Rpl35a	2.13E-210
Runx3	1.81E-146	Ccdc148	4.70E-245	Rpsa	1.62E-209
Tmem176a	2.34E-144	Grn	1.58E-244	Rpl6	5.70E-208
Ctss	4.81E-141	Bank1	1.82E-239	Tpt1	2.63E-206
Sh3pxd2b	9.38E-141	Mertk	2.15E-238	Rack1	2.14E-203
Rtcb	4.42E-140	Nr1h3	1.13E-235	Rpl23	6.14E-199
Fam20c	1.91E-139	Prnp	2.93E-235	Rpl26	7.48E-198
Il2rg	8.84E-138	Ninj1	2.42E-234	Rps6	6.64E-197
Lpcat2	8.53E-137	Fcna	3.33E-233	Rps10	2.06E-195
Kynu	8.49E-136	Csrp1	1.16E-230	Ier5	1.06E-191
Tnfsf13b	8.77E-136	Rgl1	7.18E-229	Rps3	8.23E-185

Gpr157	1.18E-135	Lpl	4.94E-223	Rpl27	8.23E-185
Tgfbr1	7.63E-135	Fam213b	1.08E-222	Rps5	8.36E-185
H2-K1	1.15E-133	Tcf7l2	1.26E-222	Rps7	3.96E-182
Basp1	1.23E-133	AB124611	4.64E-221	Rps15a	6.82E-182
Pla2g7	1.80E-132	Abcc3	3.28E-216	Rps11	1.97E-180
Fth1	4.19E-131	Fcgrt	5.79E-216	Rps4x	5.07E-180
Ggh	1.85E-126	Tgm2	1.88E-215	Rplp0	3.09E-177
Adam19	6.94E-126	Itgad	5.35E-214	Ly6i	8.17E-176
C3ar1	7.35E-125	Ptgs1	2.94E-213	S100a11	6.23E-175
Ccl12	3.37E-123	Laptm4a	1.01E-212	Atox1	1.22E-174
Hvcn1	2.51E-121	Comt	1.33E-206	Pim1	9.56E-174
Anxa3	8.60E-121	Creg1	3.24E-205	Sh3bgrl3	3.97E-173
Tgfbi	1.88E-120	Adgre1	9.67E-205	Ciita	7.35E-173
Ctsd	2.73E-117	Clec12a	6.33E-204	Eef1a1	6.09E-172
Itm2c	5.19E-116	Tspan4	7.80E-203	Rps3a1	9.09E-168
Tmem119	5.62E-116	Txn1	9.13E-203	Gm2a	6.07E-165
Rap2a	1.03E-114	Ctsb	9.52E-201	Ptprc	2.05E-163
Ctsl	4.00E-114	Mrap	5.65E-197	Rpl37	1.51E-161
Itga6	1.83E-113	Slc16a9	5.99E-197	Rps25	3.03E-160
B4galnt1	2.45E-113	Abcg3	3.83E-196	H3f3a	5.92E-159
Fam3c	1.64E-112	Pla2g15	4.22E-196	Btg2	1.14E-158
Tmem173	1.54E-111	C1qc	6.17E-192	Rpl15	1.42E-158
Ski	3.59E-111	Agpat3	1.68E-191	Cnn2	1.09E-156
Anpep	5.85E-111	Hs6st1	1.95E-191	Cdkn1a	2.57E-156
Gng2	2.37E-110	Dmpk	2.15E-191	Slfn1	4.83E-155
Nceh1	2.88E-110	Cd38	1.79E-190	Sem1	4.08E-154
H2-Q7	4.94E-108	Tmem26	2.02E-189	Lsp1	1.34E-152
Rtn1	1.28E-106	Slc11a1	1.05E-188	Rpl37a	1.78E-152
Sorl1	1.31E-103	Cd300a	1.41E-187	Rpl22	3.64E-152
Glipr1	1.22E-102	Slc7a7	3.28E-187	Sirpb1c	4.81E-152
Gsn	2.00E-102	Cyb5a	6.94E-187	Traf1	6.97E-152
Afdn	4.54E-102	Sipa1l1	7.41E-187	Emb	4.22E-151
Ak2	1.11E-101	Il18bp	1.48E-186	Rpl30	1.32E-147
Ntpcr	2.21E-98	Cd86	2.52E-183	Rps15	1.14E-146
Scarb2	3.16E-97	Vamp5	3.05E-183	H2-Ab1	2.84E-145
Creb5	5.41E-97	Jup	6.69E-182	Il1b	3.05E-145
Gsto1	5.56E-97	Blvra	1.30E-178	Rps28	4.52E-145
Ncf1	4.26E-96	Mgst1	6.48E-178	Jarid2	1.82E-143
Ppfia4	4.97E-96	Tbxas1	1.47E-177	Rps26	1.53E-142
Chchd10	7.77E-96	Hpgds	2.04E-177	Rpl32	4.21E-142
Gna12	1.23E-95	Tgfbr2	2.70E-176	Pld4	9.07E-142
Mvb12b	1.80E-95	Clec4n	3.52E-175	Cbfa2t3	1.54E-141
Rasal3	1.45E-94	Ms4a7	5.30E-175	Rps21	4.04E-141

Scoc	6.86E-94	*Sirpa*	3.35E-171	*Fgr*	4.04E-141
Cfb	6.00E-93	*Fyn*	2.84E-168	*Rps8*	1.11E-139
Lmna	1.04E-92	*Cadm1*	2.20E-167	*Cd74*	5.34E-138

1630 **Table 2: List of the differentially expressed genes among *Timd4*[+]**

1631 **thymic macrophages, *Cx3cr1*[+] thymic macrophages, and thymic**

1632 **monocytes.** The top 100 differentially expressed genes among the three

1633 clusters are listed by their negative $\log_{10}$ transformed p-value.

1634

1635