Thymic macrophages consist of two populations with distinct

localization and origin

Tyng-An Zhou[1], Hsuan-Po Hsu[1], Yueh-Hua Tu[2,3], Chih-Yu Lin[1], Nien-Jung

Chen[1], Jin-Wu Tsai[4], Ellen A. Robey[5], Hsuan-Cheng Huang[2,6], Chia-Lin Hsu[1],

and Ivan L. Dzhagalov[1*]

[1]Institute of Microbiology and Immunology, National Yang-Ming University, Taipei, Taiwan

[2]Bioinformatics Program, Taiwan International Graduate Program, Institute of Information

Science, Academia Sinica, Taipei, Taiwan

[3]Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University,

Taipei, Taiwan

[4]Brain Research Center, National Yang-Ming University, Taipei, Taiwan

[5]Division of Immunology and Pathogenesis, Department of Molecular and Cell Biology,

University of California, Berkeley, CA, United States

[6]Institute of Biomedical Informatics, National Yang-Ming University, Taipei, Taiwan

*Correspondence to Ivan L. Dzhagalov: ivan.dzhagalov@nycu.edu.tw, ORCID: 0000-0001-

9209-4582

Mailing Address: Ivan L. Dzhagalov, Institute of Microbiology and Immunology, National

Yang-Ming Chiao-Tung University, #155, Sec. 2, LiNong Street, Beitou, Taipei City, 11221,

TAIWAN

Abstract

Tissue-resident macrophages are essential for protection from pathogen invasion and maintenance of organ homeostasis. The ability of thymic macrophages to engulf apoptotic thymocytes is well appreciated, but little is known about their ontogeny, maintenance, and diversity. Here, we characterized the surface phenotype and transcriptional profile of these cells and found out that they express typical tissue-resident macrophage genes yet also exhibit organ-specific features. Thymic macrophages were most closely related to spleen red pulp macrophages and Kupffer cells and shared the expression of the transcription factor SpiC with these cells. Using shield chimeras, transplantation of embryonic thymuses, and fate mapping, we found that three distinct waves of precursors generate thymic macrophages. Moreover, some of them proliferated in situ. Single-cell RNA sequencing showed that the macrophages in the adult thymus are composed of two populations with distinct localization and origin. Altogether, our work defines the phenotype, origin, and diversity of thymic macrophages.

Introduction

Tissue-resident macrophages are present in every organ and maintain local homeostasis through diverse functions ranging from protection against pathogens to tissue repair[1]. To perform their roles efficiently, macrophages acquire specialized phenotypes depending on the tissue microenvironment, and as a consequence, multiple subtypes exist, frequently within the same organ. For example, the spleen harbors red pulp macrophages specialized in red blood cell phagocytosis, marginal zone macrophages and metallophilic macrophages that are the first defense against blood-borne pathogens, T cell zone macrophages that silently dispose of apoptotic immune cells, and tingible-body macrophages that engulf less fit B cells in the germinal center[2-4]. Thus, tissue-resident macrophages represent a fascinating developmental system that allows enormous plasticity.

The last decade has seen a paradigm shift in our understanding of the development of tissue-resident macrophages. Contrary to the long-held belief that all macrophages derive from circulating monocytes[5], multiple studies have shown that many of them are long-lived cells with an embryonic origin that can maintain themselves in the tissues (reviewed in[6]). Three waves of distinct progenitors settle the tissues and contribute in various degrees to the resident macrophages in each organ. The first wave consists of yolk sac (YS)-derived primitive macrophages that enter all tissues and establish the earliest macrophage populations[7,8]. In all organs, except for the brain and, partially, the epidermis, primitive macrophages are replaced by the next wave consisting of fetal monocytes[9-12]. The third wave comes from hematopoietic stem cell (HSC)-derived monocytes that contribute to various degrees to the

macrophage pool in different tissues. For example, these cells contribute little

to the microglia in the brain, Langerhans cells in the epidermis, and alveolar

macrophages in the lungs but substantially to most other organs[13-16].

Moreover, the kinetics and timing of HSC-derived monocytes infiltration vary

in different parts of the body. For some macrophage populations, such as the

arterial macrophages and subcapsular lymph node macrophages, monocytes

replace embryonic macrophages soon after birth and self-maintain after that

with little contribution from circulating cells[17,18]. Others, such as heart

macrophages, osteoclasts, and pancreatic isles macrophages are

progressively replaced at a low rate[14,19-23]. A third group, such as the

macrophages in the dermis and most of the gut macrophages, are constantly

replaced by blood monocytes with relatively fast kinetics[24,25]. These

conclusions have been extended to many different macrophage populations

such as Kupffer cells, liver capsular macrophages, red pulp macrophages,

testicular macrophages, large and small peritoneal macrophages, and T zone

macrophages in the lymph nodes[2,13,14,16,26-30].

The recent revitalization in macrophage research has yet to reach thymic

macrophages. Although their prodigious phagocytic ability is well

appreciated[31], little is known about the origin, diversity, and maintenance of

these cells. This gap in our knowledge is, in part, due to the lack of a

consensus about the surface phenotype of thymic macrophages.   Various

groups have used different markers to identify these cells, such as F4/80 and

Mac-3 (LAMP-2)[31], or CD4 and CD11b[32], or Mac-2 (galectin 3), F4/80 and

ED-1 (CD68)[33]. Most commonly, researchers employ F4/80 and CD11b[34-

37]. However, none of these markers is macrophage-specific: F4/80 is also

expressed on eosinophils and monocytes[38,39], while CD11b is present on

most myeloid cells. The lack of a clear phenotypic definition of thymic

macrophages has translated into the absence of models that target genes

specifically in this population. For example, although macrophages in various

organs have been successfully targeted with $Lyz2^{Cre}$, $Csf1r^{Cre}$, or $Cx3cr1^{Cre}$,

very few studies have used these models in the thymus[37,40,41].

Only a handful of studies has explored the origin of thymic macrophages.

Several reports have indicated that these cells could be derived from T cell

progenitors in the thymus based on an improved single cell in vitro culture and

in vivo transplantation experiments[42,43]. However, these conclusions have

been questioned based on fate-mapping experiments using $Il7r^{Cre}$ that found

very limited contribution of lymphoid progenitors to thymic macrophages in

vivo in unperturbed mice[44]. Most recently, Tacke et al. used parabiosis to

rule out circulating monocytes as a major source for thymic macrophages[37].

The same study also performed fate-mapping experiments to show that most

thymic macrophages descend from $Flt3^+$ HSC-derived progenitors. However,

the contribution of earlier waves of hematopoiesis has not been explored.

Here, we aimed to bring our knowledge of thymic macrophages on par with

other tissue-resident macrophages. We started by providing a clear definition

of thymic macrophages according to the guidelines set by the Immunological

Genome Consortium (IMMGEN)[38] and characterized their surface

phenotype and transcriptional signature. Using single-cell RNA sequencing

(scRNA-Seq), we identified two populations of thymic macrophages with

distinct localization. We explored the origin of these cells through genetic fate

mapping, shield chimeras, and embryonic thymus transplantation and

documented that three separate waves of progenitors give rise to resident

thymic macrophages. Altogether our work fills an important gap in our

understanding of resident thymic macrophages and provides the framework

for future functional characterization of these cells.

Results:

CD64, F4/80, and MerTK identify the macrophages in the thymus

To unambiguously and comprehensively identify macrophages in the

thymus, we evaluated several of the prototypical macrophage markers,

MerTK, CD64, and F4/80[38]. Only MerTK identified a population that stained

completely with the other two markers (Fig. 1a). F4/80 labeled Siglec F^+SSC^{hi}

eosinophils and $CD64^+CD11b^+$ monocytes, in addition to $MerTK^+$ cells (Fig.

S1a). CD64 identified $CD11b^+F4/80^{lo}$ monocytes (Fig. S1b). Thus, MerTK

appeared to be the best marker for macrophages in the thymus. However,

staining with MerTK and F4/80 was relatively dim even when the brightest

fluorochromes (e.g., PE) were used and could not be resolved fully from the

isotype control (Fig. S1c). For practical purposes, to identify macrophages,

we used CD64 together with F4/80 and CD11b to exclude monocytes (Fig.

S1d). In addition, we also employed forward scatter (cells size) to exclude

cells that appeared $MerTK^+$, $CD64^+$, $F4/80^+$, but were much smaller and less

granular than the rest of the macrophages, and many of them were $Thy1^+$

(Fig. S1e).

The $CD64^+F4/80^+MerTK^+CD11b^{lo}FSC^{hi}$ cells had typical macrophage

morphology with abundant cytoplasm (Fig. 1b). These cells did not express

lineage markers characteristic of T cells (CD3ε), B cells (CD19), eosinophils

(Siglec F), NK cells (NK1.1), neutrophils (Gr1), or plasmacytoid dendritic cells

(Siglec H) (Fig. 1c). However, they expressed phagocytic receptors such as

TIM4, CD51, and Axl (Fig. 1d).

To ascertain that our gating strategy identifies all macrophages in the

thymus in an unbiased way, we performed scRNA-Seq of sorted $Csf1r^{GFP+}$

and $Cd11c^{YFP+}$ thymic cells. $Csf1r$ is required for the survival of most

macrophages and is considered their definitive marker[45,46], while $Cd11c^{YFP}$

is expressed in many myeloid cells, including macrophages[47]. Both

reporters identified an overlapping set of cells (Fig. S2a). Two clusters

expressed the macrophage/monocytes-specific transcription factor $Mafb$ and

high levels of $Fcrg1$ (CD64), $Mertk$, and $Adgre1$ (F4/80), indicating that they

are macrophages (Fig. 1e). An additional cluster expressed $Mafb$ together

with $Fcgr1$ and $Adgre1$ but not $Mertk$, fitting the description of monocytes.

Altogether the scRNA-Seq data confirmed that our flow cytometry gating had

identified all macrophages in the thymus. Importantly, MerTK$^+$ cells could not

be labeled by intravenously injected CD45 antibody (Fig. 1f), proving that they

reside in the parenchyma of the organs and not in the blood vessels. Based

on the above data, we will refer to CD64$^+$F4/80$^+$MerTK$^+$CD11b^{lo}FSC^{hi} cells as

thymic macrophages. The smaller CD64$^+$F4/80^{lo}CD11b$^+$FSC^{hi} population did

not express MerTK, but most of them expressed Ly6C, and we classify them

as thymic monocytes.

Thymic macrophages expressed CD11c, MHC2, and SIRPα that made

them partially overlap with CD11c$^+$MHC2$^+$ classical dendritic cells (cDCs),

thus making problematic the unambiguous identification of thymic cDCs

based only on these two markers (Fig. S1f). Proper identification of cDC in the

thymus requires the exclusion of macrophages based on CD64 or MerTK

staining. Otherwise, the cDCs, particularly the SIRPα^+ cDC2 subset, would

be contaminated with macrophages that account for ~25% of cDC2 (Fig.

S1g).

Thymic macrophages were ~0.1% of all the cells in the thymus of young

adult mice and numbered ~$4x10^5$ on average per mouse (Fig. 1g). We did not

find statistically significant differences in their percentages between 4 and 11

weeks of age. Still, there was a significant decline in their numbers with age,

consistent with the beginning of thymic involution (Fig. 1h).

To determine if thymic macrophages are avid phagocytes, we evaluated

their participation in the phagocytosis of apoptotic cells in the thymus by

TUNEL staining. We used TIM4 and MerTK as single stains to identify

macrophages in tissue sections because they are not expressed on any other

cell type in the thymus instead of following our multiple-label flow cytometry

strategy. Most of the TUNEL$^+$ cells could be found clearly inside or closely

associated with MerTK$^+$ and TIM4$^+$ cells in the thymus (Fig. S3a and b). On

average, ~85% of all TUNEL$^+$ cells were within 5 µm of MerTK$^+$ cells,

indicating that thymic macrophages are the major phagocytic population in the

thymus (Fig. S3c). The degree of co-localization between TUNEL$^+$ cells and

TIM4$^+$ cells was slightly lower, ~75% on average, possibly reflecting the

absence of TIM4 expression on a small proportion of thymic macrophages

(Fig. 1d).

Figure 1. Thymic macrophages (ThyMacs) can be identified by the

expression of CD64, MerTK and F4/80. a Flow cytometric analysis of

enzymatically digested thymus tissue with macrophage markers CD64,

MerTK, F4/80, and CD11b. More details about the confirmation of the

identities of eosinophils and monocytes can be found in Fig. S1a and b. **b**

Pappenheim (Hemacolor Rapid staining kit) staining of sorted ThyMacs. **c**

Lack of expression of lineage markers associated with other cell types on

ThyMacs. **d** The expression on ThyMacs of three receptors for

phosphatidylserine that participate in the phagocytosis of apoptotic cells. **e**

scRNA-Seq data showing the expression of prototypical macrophage genes

Mafb, *Fcgr1* (CD64), *Mertk*, and *Adgre1* (F4/80) among thymus cells sorted

as *Csf1r*^{GFP+} and *Cd11c*^{YFP+}. **f** Labeling of ThyMacs with intravenously

injected anti-CD45-PE antibody or PBS. The labeling of blood leucocytes is

shown for comparison. **g** Average numbers and percentages of ThyMacs in 4-

11 weeks old mice, n=82. **h** Comparison of the numbers and percentages of

ThyMacs in mice of different ages, n=82. All flow cytometry plots are

representative of at least 3 independent repeats. The numbers in the flow

cytometry plots are the percent of cells in the respective gates. Data in **g** and

h represent mean±SEM. Statistical significance in **h** was determined with one-

way ANOVA.

Transcriptional signature of thymic macrophages

To further understand the identity and functions of the thymic

macrophages, we sorted and subjected them to RNA sequencing analysis as

part of the IMMGEN's Open Source Mononuclear Phagocyte profiling.

Thymic macrophages displayed a transcription profile consistent with tissue-

resident macrophages and did not express genes characteristic of other

lineages (Fig. 2a). Then, we examined the expression of the core signature

macrophage genes[38] and found that they were enriched in thymic

macrophages but not in *Sirpa*[+] or *Xcr1*[+] thymic cDCs (Fig. 2b). On the

contrary, cDC core signature genes were abundantly expressed in both

thymic cDC subsets but not in thymic macrophages. These findings establish

that although thymic macrophages and cDCs share the thymic

microenvironment and expression of CD11c and MHC2, they have distinct

transcriptional profiles.

We then compared the gene expression profile of thymic macrophages to

that of other well-characterized macrophage populations available from

IMMGEN. Because of the abundance of samples, we limited our comparison

to only nine types of tissue-resident macrophages under steady-state

conditions – splenic red pulp macrophages, Kupffer cells, broncho-alveolar

lavage macrophages, large peritoneal cavity macrophages, white adipose

tissue macrophages, aorta macrophages, central nervous system microglia,

and spinal cord microglia. Hierarchical clustering revealed that thymic

macrophages were most closely related to splenic red pulp macrophages and

Kupffer cells (Fig. 2c).

To better identify the unique functions of thymic macrophages, we looked

for genes that were differentially expressed in these cells compared to other

tissue-resident macrophages. We set three criteria: 1) high expression in

thymic macrophages (>500); 2) >5 fold higher expression than the average

value in the nine populations of non-thymic macrophages; 3) expression in

thymic macrophages is higher than any non-thymic macrophage samples. A

total of 44 genes met these criteria, and we consider them to constitute the

transcriptional signature of thymic macrophages (Fig. 2d). These included

several degradation enzymes and their inhibitors (*Cst7*, *Mmp2*, *Mmp14*,

Dnase1l3, *Serpina3g*, *Acp5*), non-classical MHC molecules (*H2-M2*, *H2-Q6*,

H2-Q7), metabolic enzymes (*Chst2*, *Ass1*, *Kynu*, *Cp*, *Dgat2*, *Sorl1*, *Lap3*),

molecules involved in innate immunity (*Ifit2*, *Il18bp*, *Mefv*, *Lgals3bp*) and

extracellular signaling molecules and their receptors (*Pdgfa*, *Cxcl16*, *Il2rg*,

Gpr157). We also looked for transcription factors (TFs) highly expressed in

thymic macrophages and could potentially regulate critical gene networks in

these cells. A total of 25 TFs were highly expressed in thymic macrophages

(>250) and were at least 2-fold higher as compared to the non-thymic

macrophages (Table S1). Among them were several TFs involved in type I

interferon (IFN-I) signaling (*Stat1*, *Stat2*, *Irf7*, and *Irf8*) and lipid metabolism

(*Nr1h3*, *Pparg*, *Srebf1*, and *Rxra*) (Fig. 2e). Notably, *Runx3* that is important

for the development and function of cytotoxic T lymphocytes[48], innate

lymphoid cells[49], and Langerhans cells[50] was highly expressed in thymic

macrophages. *SpiC* that has well-documented roles in the development of red

pulp macrophages in the spleen, and bone marrow macrophages[51,52] was

also highly expressed in thymic macrophages, further strengthening the

argument for the similarity between thymus, spleen and liver macrophages.

To confirm the expression of *Spic* in thymic macrophages, we analyzed the

thymus of *Spic*^{GFP} mice[52]. We found that all *Spic*$^{GFP+}$ cells were

macrophages (Fig. S4a and b), making them the most specific thymic

macrophage reporter strain compared with *Lyz2*^{GFP}, MAFIA (*Csf1r*^{GFP}),

Cd11c^{YFP}, and *Cx3cr1*^{GFP} mice (Fig. S4c-e). However, only ~80% of thymic

macrophages were *Spic*$^{GFP+}$ suggesting heterogeneity within the cells (Fig.

S4f and g).

Several dominant pathways emerged when we grouped the 500 most

highly expressed genes in thymic macrophages according to gene ontology

(GO) terms (Fig. 2f). It was notable that five of the ten most highly enriched

GO pathways concerned antigen presentation of both exogenous and

endogenous antigens. These data complement our flow cytometry findings of

expression of MHC2 and suggest that thymic macrophages could be potent

antigen-presenting cells and might play a role in negative selection or agonist

selection of thymocytes. Two other highly enriched GO pathways were

involved in lysosomal biogenesis and functions, highlighting the high capacity

of these cells to degrade phagocytosed material. Thus, our transcriptional

analysis has revealed that thymic macrophages are bona fide macrophages

that bear significant similarity to spleen and liver macrophages and are

specialized in lysosomal degradation of phagocytosed material and antigen

presentation.

Figure 2. Transcriptional profile of thymic macrophages (ThyMacs). a

Expression of lineage-specific genes in ThyMacs and peritoneal cavity

macrophages (PC Macs). **b** Expression of cDC-specific genes (gene names in

blue) and macrophage-specific genes (gene names in red) in ThyMacs and

two populations of thymic cDCs (ThyDCs) – *Xcr1*⁺ ThyDCs and *Sirpa*⁺

ThyDCs. **c** Hierarchical clustering of ThyMacs and nine other populations of tissue-resident macrophages in duplicates. **d** Highly expressed (>500) genes enriched (>5-fold) in ThyMacs (4 samples) compared to nine other tissue-resident macrophage populations (two samples each). The genes in red are >10-fold up-regulated in thymic macrophages. **e** Comparison of the geometric mean expression of transcription factors in thymic macrophages (4 samples) and the nine other macrophage populations (2 samples each). Transcription factors with expression >250 and fold change >2 are marked with red dots. **f** Top 10 GO pathways in ThyMacs based on the 500 most highly expressed genes in these cells. Th – ThyMacs, Spl – spleen red pulp macrophages, Lv – Kupffer cells, BAL – bronchoalveolar lavage macrophages, PC – peritoneal cavity macrophages, Ao – aorta macrophages, Ht – heart macrophages, WAT – white adipose tissue macrophages, CNS – central nervous system microglia, SC – spinal cord microglia.

Yolk-sac progenitors contribute to embryonic thymic macrophages

The ontogeny of thymic macrophages has been examined by only one study since the realization that many tissue-resident macrophages are descendants from embryonic progenitors[37]. Based on $Flt3^{Cre}$ fate-mapping, the authors concluded that most adult thymic macrophages derive from HSCs. To determine if yolk-sac (YS) progenitors contribute to embryonic thymic macrophages, we used $Cx3cr1^{CreER}$ fate mapping[53]. Injection of 4-OHT at E9.5 in $ROSA26^{LSL-GFP}$ mouse mated with a $Cx3cr1^{CreER}$ male permanently tags YS progenitors and their descendants with GFP (Fig. 3a). Indeed, E19.5 microglia that are exclusively derived from YS progenitors were

labeled to a high degree (Fig. 3b). After adjusting for incomplete labeling

based on the microglia, we found that at E15.5 >50% of thymic macrophages

were fate mapped, i. e. from YS origin (Fig. 3c). However, GFP$^+$ thymic

macrophages decreased to just ~11% at E19.5, suggesting that YS

progenitors establish the embryonic thymic macrophage pool but are quickly

replaced by subsequent wave(s) of fetal liver monocytes or HSC-derived

macrophages.

To pinpoint when do the HSC-derived progenitors enter the thymus, we

devised two complementary experiments. First, we evaluated the contribution

of circulating adult monocytes to thymic macrophages without the

confounding effect of radiation damage on the thymus. We created shield

chimeras by subjecting CD45.1 mice to a lethal dose of irradiation while

protecting their upper body and the thymus with a 5 cm lead shield followed

by reconstitution with CD45.2 bone marrow (Fig. 3d). While the donor-derived

monocytes in the blood were, on average 20%, less than 2% of thymic

macrophages were CD45.2$^+$ (Fig. 3e and f), suggesting a relatively minor

(<10%) contribution of adult circulating monocytes to the thymic macrophage

pool consistent with a previous report[37]. Second, we transplanted E15.5

embryonic thymuses expressing GFP ubiquitously under the control of the

ROSA26 locus (*ROSA26^{GFP}*) under the kidney capsule of adult mice and

analyzed them six weeks later (Fig. 3g). By that time, >99% of thymocytes

were derived from GFP$^-$ host HSCs (Fig. 3h and i). As a positive control for

donor-derived cells, we used the thymic epithelial cells. The vast majority of

EpCAM$^+$ thymic epithelial cells (>90%) were still GFP$^+$. An identical

proportion of thymic macrophages was also GFP$^+$. The results from our

transplantation experiments show that the progenitors of almost all thymic

macrophages are of embryonic origin. Altogether our results suggest that

resident thymic macrophages are derived from multiple waves of progenitors.

Initially, the thymus is settled by YS-derived progenitors that contribute

substantially to the thymic macrophage pool during the embryonic period. In

parallel with them, YS-independent progenitors infiltrate the thymus before

E15.5 and establish themselves as the dominant population before birth.

Adult HSC-derived monocytes contribute relatively little to the pool of thymic

resident macrophages in young adult mice.

Figure 3. Yolk sac (YS)-derived and non-YS-derived embryonic

progenitors sequentially contribute to the thymic macrophage pool. a

Scheme of the YS-progenitor labeling experiments. E9.5 pregnant

ROSA26^{LSL-GFP} mice mated with *Cx3cr1^{CreER}* males were injected with 4-

hydroxytamoxifen (4-OHT) and sacrificed at E15.5 or E19.5. **b** Representative

flow cytometry plots of the *Cx3cr1^{GFP}* expression in microglia (CD45$^+$CD11b$^+$

cells in the brain) and ThyMacs of the pups. **c** Frequencies of GFP$^+$ ThyMacs

at E15.5 and E19.5 adjusted to the degree of labeling of microglia. **d** Scheme

of the shield chimera experiments. Congenic CD45.1 mice were lethally

irradiated with their upper body protected by a 5 cm thick lead shield and then

injected with CD45.2$^+$ bone marrow. **e** Representative flow cytometric

analysis of ThyMacs and CD115$^+$CD11b$^+$ blood monocytes for donor-derived

(CD45.2$^+$) cells. Non-chimeric CD45.1 and CD45.2 samples serve as controls

for the gating. **f** Frequencies of donor-derived ThyMacs and blood

monocytes. **g** Scheme of the thymus transplantation experiments. **h**

Representative flow cytometry plots of GFP expression in thymocytes (Thy),

CD45$^-$EpCAM$^+$ thymic epithelial cells (TECs), and thymic macrophages

(ThyMacs) six weeks after the transplantation. The host thymus (endogenous

thymus) serves as a negative control. The percentages of GFP$^+$ cells are

indicated in the plots. **i** Frequencies of GFP$^+$ cells in different cell populations

in the transplanted thymus. Data in **c**, **f**, and **i** are mean±SEM with two litters,

three, and five mice per group, respectively. The numbers in the flow

cytometry plots are the percent of cells in the respective gates. Each dot is an

individual mouse or embryo.

Thymic macrophages can proliferate in situ

The fact that macrophages can persist for many weeks in the thymus

without constant replacement from blood monocytes suggests that they can

divide in situ. Staining for the proliferation marker Ki-67 revealed that ~4% of

the cells expressed this marker compared to an isotype control (Fig. 4a and

b). To obtain further proof that thymic macrophages are proliferative, we

tested the incorporation of the nucleotide analog 5-Ethynyl-2'-deoxyuridine

(EdU). Short-term EdU labeling experiments unexpectedly revealed that

thymic macrophages become EdU$^+$ with faster kinetics than thymocytes (Fig.

4c). The most likely explanation for this puzzling result is that some of the

thymic macrophages have engulfed apoptotic thymocytes that have recently

divided and incorporated EdU. Thus, EdU could have accumulated in these

macrophages through phagocytosis and not through cell division. To

circumvent this caveat, we designed a pulse-chase experiment (Fig. 4d).

Mice were injected daily with EdU for 21 days so that all cells that proliferate

in that period would incorporate the label. The vast majority of thymocytes

and thymic macrophages became EdU$^+$ at d. 21 (Fig. 4e). After 21 more days

of "chase period", only ~0.2% of thymocytes had retained the EdU label,

consistent with the existence of a very small population of long-term resident

thymocytes consisting mainly of regulatory T cells and NKT cells[54] (Fig. 4e

and f). However, ~5% of the thymic macrophages were EdU$^+$, suggesting

they divided during the labeling period. Finally, we sorted thymic

macrophages and subjected them to cell cycle analysis. Although almost all

thymic macrophages were in G0/G1 phase, a small population of ~3% was in

G2/M phase of the cell cycle (Fig. 4g and h). Collectively, three independent

approaches documented that 3-5% of thymic macrophages are actively

dividing under homeostatic conditions within the thymus. These findings can

explain the long-term maintenance of these cells within the organ without a

constant influx of progenitors.

Figure 4. Thymic macrophages exhibit a low degree of proliferation. a

Example flow cytometry plots of Ki67 staining of thymic macrophages

(ThyMacs). **b** Frequency of Ki67+ thymic macrophages. **c** Example flow

cytometry plots of the EdU accumulation in thymocytes and thymic

macrophages 2 hours after 1 mg EdU i.p. or vehicle injection. The numbers

inside flow plots are the percentage of EdU+ cells from mice injected with

EdU. Data are representative of three independent experiments. **d** Scheme of

EdU pulse/chase experiment: mice were injected daily with 1 mg EdU i.p for

21 days and then rested for 21 more days. **e** Example flow cytometry plots of

EdU staining of thymocytes (upper row) and ThyMacs (lower row). **f**

Frequencies of EdU+ cells among thymocytes (top graph) and ThyMacs

(bottom graph). **g** Example flow cytometry plot of cell cycle analysis of FACS

sorted ThyMacs. **h** Frequencies of ThyMacs in different stages of the cell

cycle. The numbers in the flow cytometry plots are the percent of cells in the

respective gates. Data are mean±SEM from three mice (**b** and **h**) or 6-7

individual mice (**f**). Each dot is an individual mouse.

Expression of *Timd4* and *Cx3cr1 can distinguish* two populations of

thymic macrophages

To determine if thymic macrophages are heterogeneous, we turned to our

scRNA-Seq data. Once we zoomed onto *Mafb* expressing cells, we could

distinguish three separate populations: 1) monocytes that expressed high

levels of *Ly6c2* and *Itgam* (CD11b) but did not express *Mertk*; 2) *Timd4*+

(encoding TIM4) macrophages that also expressed high levels of *Spic* and

Slc40a1, but low levels of *Cx3cr1*; 3) *Cx3cr1*+ macrophages that expressed

low levels of *Timd4*, *Spic*, and *Slc40a1* (Fig. 5a and b). Both macrophages

and monocytes expressed *Fcgr1* (CD64). A minority of macrophages

expressed high levels of *Mki67*, indicating that they might be proliferating,

corroborating our earlier data. Interestingly, most of the *Mki67* expressing

macrophages belonged to the *Cx3cr1*+ population.

We confirmed the results from scRNA-Seq by flow cytometry. We could

identify discrete TIM4+*Cx3cr1*GFP- and TIM4-*Cx3cr1*GFP+ macrophages (Fig.

5c). There was even TIM4+*Cx3cr1*GFP+ intermediate population that could not

be distinguished in the scRNA-Seq dataset, likely because of the lack of

statistical power. To determine the localization of the two distinct macrophage

populations, we stained thymic sections from *Cx3cr1*^{GFP} mice with an antibody

to MerTK. The *Cx3cr1*$^{GFP-}$ MerTK$^+$ cells correspond to *Timd4*$^+$ macrophages,

while the *Cx3cr1*$^{GFP+}$MerTK$^+$ cells would be the *Cx3cr1*$^{GFP+}$ macrophages.

Strikingly, the two macrophage populations showed distinct localization.

Timd4$^+$ macrophages were located in the cortex, while the *Cx3cr1*$^{GFP+}$

macrophages resided in the medulla and the cortico-medullary junction (Fig.

5d).

To better understand the differences between the two populations of thymic

macrophages, we looked for differentially expressed genes. We included the

thymic monocytes in the comparison, as these cells clustered the closest to

macrophages. *Timd4*$^+$ macrophages expressed the highest levels of the

transcription factors *Spic*, *Maf*, and *Nr1h3*; the receptors for apoptotic cells

Axl, *Mertk*, and *Timd4*; and many Slc transporters such as *Slc40a1*, *Slc1a3*,

Slco2b1, *Slc11a1*, and *Slc7a7* (Fig. 5e and Table S2). *Cx3cr1*$^+$ macrophages

expressed high levels of the transcription factor *Runx3*; a distinct set of

phosphatidylserine receptors such as *Stab1*, *Anxa5*, and *Anxa3*; many

degradative enzymes such as *Mmp2*, *Mmp14*, *Dnase1l3*, *Acp5*, *Lyz2*, *Ctsz*,

Ctss, *Ctsd*, *Ctsl*; cytokines such as *Pdgfa*, *Cxcl16*, and *Ccl12*; and molecules

involved in MHC1 antigen presentation such as *B2m*, *H2-M2*, *H2-K1*, *H2-Q7*.

Thymic monocytes were characterized by differential expression of the typical

monocyte genes *Ly6c2*, *Ccr2*, and *S100a4*, and genes involved in MHC2

antigen presentation such as *Ciita*, *H2-DMb1*, *H2-Ab1*, and *Cd74*.

Figure 5: **Two populations of macrophages with distinct localization**

exist in the thymus. a UMAP clusters from Fig. S2 with high expression of

the transcription factor *Mafb* fall into three groups: monocytes, *Timd4*+

macrophages, and *Cx3cr1*+ macrophages. **b** Expression of the indicated

genes in the three *Mafb*-positive clusters. **c** A flow cytometry plot of *Cx3cr1*^{GFP}

and TIM4 expression in ThyMacs. The plot is representative of >10 individual

experiments. The numbers inside the plot are the percentages of the cell

populations in the respective gates. **d** Immunofluorescent staining of the

thymus of *Cx3cr1*^{GFP} mouse stained with MerTK (a marker for all

macrophages) and Keratin 5 (a marker for medulla). The scale bar is 150 μm.

Areas in the cortex, medulla, and the cortico-medullary junction (CMJ)

represented by the dashed boxes are enlarged below to show the co-

localization of *Cx3cr11*^{GFP} and MerTK signal in CMJ and medulla, but not in

cortex. The scale bars in the images below are 20 μm. The images are

representative of three individual mice. **e** Differentially expressed genes

among *Timd4*$^+$ thymic macrophages, *Cx3cr1*$^+$ thymic macrophages, and

thymic monocytes. The negative $\log_{10}$ p-values for the genes expressed in

each cluster were calculated as described in the Materials and Methods, and

the top 50 differentially expressed genes were plotted in the figure. Ten of

these genes are listed on the left.

***Cx3cr1*$^+$ cells give rise to *Timd4*$^+$ cells during embryonic development**

To determine if the two populations of thymic macrophages are related, we

first analyzed the kinetics of their appearance during embryonic development.

At the earliest time point (E14.5), all thymic macrophages were *Cx3cr1*$^+$, and

only ~20% of them were also TIM4$^+$ (Fig. 6a and b). The proportion of TIM4$^+$

cells increased at E17.5, and TIM4$^+$*Cx3cr1*$^-$ cells started to appear. In the

neonatal period, almost all macrophages were TIM4$^+$, and very few remained

TIM4$^-$. The proportion of TIM4$^-$ cells increased in 6 weeks old mice, but TIM4$^+$

macrophages remained the dominant population. These kinetics (Fig. 6c) are

consistent with *Timd4*$^+$ macrophages developing from *Cx3cr1*$^+$ cells before

birth. Another plausible scenario is that distinct progenitors give rise to

different thymic macrophages populations (e.g., YS-progenitors give rise to

Cx3cr1$^+$*Timd4*$^-$ and HSC-derived progenitors develop into *Timd4*$^+$

macrophages). To test the latter hypothesis, we re-visited the fate mapping of

YS progenitors results (Fig. 3a). Although a larger part (~60% at E15.5) of

Cx3cr1$^+$TIM4$^-$ cells were derived from YS progenitors (Fig. 6d), a substantial

proportion (~40% at E15.5) of YS-derived TIM4$^+$ macrophages could clearly

be identified at both E15.5 and E19.5, suggesting that YS progenitors can

give rise to both *Cx3cr1*⁺ and *Timd4*⁺ cells. Extrapolating from the data, the

non-YS-derived progenitors (GFP⁻ cells) can also give rise to both *Timd4*⁻ and

Timd4⁺ cells. Thus, the simplest explanation for our findings is that *Timd4*⁺

cells develop from *Cx3cr1*⁺ cells during embryonic development. This

transition is complete in the first week after birth as there were essentially no

Cx3cr1⁺ thymic macrophages remaining at d.7 (Fig. 6a and b).

Figure 6. *Timd4*⁺ thymic macrophages are derived from *Cx3cr1*⁺ cells

during embryonic development. a Example flow cytometry plots for the

expression of *Cx3cr1*^GFP and TIM4 on thymic macrophages at different times

during embryonic development (E14.5, E17.5), immediately after birth, at 7

days, and 6 weeks of age. **b** Frequencies of *Timd4*⁺*Cx3cr1*⁻ (*Timd4* single-

positive or *Timd4*SP), *Timd4*⁺*Cx3cr1*⁺(double-positive or DP), and

Cx3cr1⁺*Timd4*⁻ (*Cx3cr1* single-positive or *Cx3cr1*SP) thymic macrophages at

the indicated time points. **c** Kinetics of the changes in different subpopulations

of thymic macrophages from E14.5 to 6 weeks. **d** Frequencies at E15.5 and

E19.5 of GFP-labeled cells among *TIM4*$^+$ or *TIM4*$^-$ cells in *Cx3cr1*^{CreER} X

ROSA26$^{LSL-GFP}$ embryos treated with 4-OHT at E9.5. Data is from at least two

independent experiments for each panel. Each symbol is an individual mouse

or embryo.

Timd4$^+$ thymic macrophages self-maintain, while Cx3cr1$^+$ cells slowly

accumulate with age

To understand how are the resident thymic macrophage populations

maintained during adult life, we induced recombination in *Cx3cr1*^{CreER} X

ROSA26$^{LSL-GFP}$ mice during the neonatal period (Fig. 7a) or at 6 weeks of age

(Fig. 7c) and compared the proportion of GFP$^+$ cells 3 and 42 days after

labeling. The extent of labeling of *Timd4*$^+$ thymic macrophages did not

change within these 6 weeks, no matter whether the mice were treated with

Tamoxifen in the first week after birth or at 6 weeks (Fig. 7b and d),

suggesting steady-state maintenance and absence of a significant influx from

unlabeled cells (e.g., monocytes). In contrast, the proportion of labeled

Timd4$^-$ thymic macrophages decreased significantly 6 weeks after Tamoxifen

injection in both neonatal and adult mice, suggesting that this population was

being diluted by unlabeled cells. To further substantiate these findings, we

examined older WT mice and found out that the proportions of *Timd4*$^-$ thymic

macrophages increased with age, and in mice >8 months old, they accounted

for ~70% of all macrophages in the organ (Fig. 7e).   Thus, we conclude that

Timd4$^+$ macrophages can maintain themselves for long periods in the cortex

of the thymus, Since the first week of life, *Cx3cr1*$^+$ cells are slowly being

recruited to the medulla and cortico-medullary junction, and in aging mice

form the predominant phagocytic population in the organ.

Figure 7. *Timd4*⁺ thymic macrophages self-maintain during adult life,

while *Cx3cr1*⁺ cells slowly accumulate with age. a Scheme of the neonatal

fate mapping: A nursing dam was injected twice with Tamoxifen (Tam) or

vehicle (Veh) in the first week after giving birth to *Cx3cr1^CreER^* X *ROSA26^LSL-*

^GFP^ pups. Three or 42 days after the last injection, the pups were sacrificed,

and the degree of labeling of *Timd4*⁺ and *Timd4*⁻ thymic macrophages was

examined by flow cytometry. **b** Frequencies of GFP⁺ *Timd4*⁺ or *Timd4*⁻ thymic

macrophages from neonatally fate mapped mice after 3 and 42 days. Vehicle

injected nursing dam litters (Veh) served as a control for non-specific labeling.

c Scheme of the adult fate mapping: Six weeks old *Cx3cr1^CreER^* X *ROSA26^LSL-*

^GFP^ mice were injected twice with Tamoxifen (Tam) or vehicle (Veh). Three or

42 days after the last injection, the mice were sacrificed, and the degree of

labeling of *Timd4*⁺ and *Timd4*⁻ thymic macrophages was examined by flow

cytometry. **d** Frequencies of GFP$^+$ *Timd4$^+$* or *Timd4$^-$* thymic macrophages

from adult fate mapped mice after 3 and 42 days. **e** Frequencies of TIM4$^-$

thymic macrophages at different ages. The data is mean±SEM from 2

independent experiments (**b**) or at least 3 individual mice per time point (**d** and

e). Each symbol is an individual mouse. Statistical significance was

determined with unpaired Student's t-test.

Discussion

Here, we have described the phenotype, transcriptional profile, localization,

diversity, ontogeny, and maintenance of macrophages in the thymus. These

cells express the typical macrophage markers CD64, MerTK, and F4/80 and

are transcriptionally most similar to splenic red pulp macrophages and liver

Kupffer cells. However, they have a unique expression profile dominated by

genes involved in antigen presentation and lysosomal degradation. We found

that thymic macrophages consist of two populations with distinct localization.

Timd4$^+$ macrophages occupied the cortex, while *Cx3cr1$^+$* cells were located in

the medulla and the cortico-medullary junction. While YS-derived

macrophages dominated the early stages of thymus development, they were

quickly replaced by non-YS embryonic progenitors that gave rise to the

Timd4$^+$ thymic macrophages that could proliferate and self-maintain. *Cx3cr1$^+$*

macrophages started slowly accumulating after birth and in old mice became

the most abundant population.

Altogether our data depict thymic macrophages as typical tissue-resident

macrophages with origin from multiple hematopoietic waves, ability to self-

maintain, and expression of the core macrophage-specific genes. They were

most similar transcriptionally to splenic red pulp macrophages and Kupffer

cells, which is not surprising considering that they all specialize in

efferocytosis and have efficient lysosomal degradation machinery. These

three populations also shared expression of the transcription factor *Spic* that

is induced by heme released following red blood cells phagocytosis[52].

However, the thymus is not known as a place for erythrocyte degradation.

Thus the mechanism for *Spic* up-regulation in thymic macrophages is unclear.

The unique features of thymic macrophages include high expression of

genes involved in the IFN-I pathway, antigen presentation, and lysosomal

degradation. The up-regulation of IFN-I-stimulated genes such as *Stat1*,

Stat2, *Irf7*, and *Irf8* can be explained by the constitutive secretion of IFN-I by

thymic epithelial cells[55,56]. The purpose of IFN-I expression in the thymus

in the absence of a viral infection is unclear. Still one possibility is that it

mediates negative selection to IFN-dependent genes as part of central

tolerance.

Thymic macrophages highly express molecules involved in antigen

presentation, including MHC1 and MHC2, although the latter is expressed at

lower levels than in cDCs. Thus, they have the potential to present antigens

for both negative selection and agonist selection. These two activities have

traditionally been assigned solely to cDCs[57]. However, recent evidence

suggests that negative selection is most efficient when the cell that presents

the antigen to auto-reactive thymocytes is also the one that phagocytoses

it[58]. So, macrophages participation in thymocyte selection needs to be re-

evaluated with optimized isolation procedures and specific genetic tools.

The extraordinary ability of thymic macrophages to engulf and degrade

apoptotic thymocytes has been appreciated for a long time[31], and our RNA-

Seq data provides additional supporting evidence for this function by

highlighting the up-regulation of pathways involved in lysosomal degradation.

An interesting topic for future research would be to understand how the

metabolites derived from apoptotic cells are returned to the microenvironment

to support the proliferation of immature thymocytes. A *SoLute Carrier* (*Slc*)

genes-based program has been described in vitro[59], but its relevance to

tissue-resident macrophages remains to be determined. Altogether, our study

demonstrates that thymic macrophages are a unique subset of tissue-resident

macrophages and support the idea that resident macrophage phenotype is

determined by the combination of ontogeny, microenvironment, and other

factors[60].

Together with the study by Tacke et al., our work builds the following model

for thymic macrophage origin[37]: Thymic macrophages develop in three

distinct waves: YS-derived progenitors dominate the early stages of thymus

development but are replaced before birth by a second wave of YS-

independent embryonic progenitors that forms the bulk of thymic

macrophages after birth and can self-maintain into adulthood. With age, there

is a slow and steady influx of *Timd4$^-$Cx3cr1$^+$* macrophage precursors that

occupy the medulla and cortico-medullary junction, becoming the major

phagocytic population in the thymus of older mice (>8 months). The second

wave of YS-independent macrophages is most likely the progeny of

embryonic HSCs based on *Flt3^{Cre}* fate mapping that showed that >80% of

thymic macrophages in adult mice are descendants of HSCs[37]. Whether

HSC-independent fetal liver monocytes contribute to thymic macrophages and

to what extend awaits the creation of models that can specifically target this

population of progenitors. Recruitment of circulating monocytes to the resident

macrophage pool in the thymus has been ruled out previously by parabiosis

and *Ccr2*[-/-] mice[37]. Our shield chimera experiments have arrived at similar

conclusions. However, the relatively short duration of these experiments and

their focus on the bulk thymic macrophages have prevented the recognition of

the gradual accumulation of *Timd4*[-] macrophages. Once we zoomed in on this

minor cell population in young mice, the fate mapping clearly showed an influx

of unlabeled progenitors. Whether the progenitors of *Timd4*[-] macrophages

are monocytes remains to be formally demonstrated. However, in all

macrophage populations exhibiting replacement in adults examined to date,

monocytes have been singled out as the source[12,19,22,24,25,30]. An

alternative possibility involves thymocyte progenitors that under certain

circumstances have been shown to differentiate into macrophages and

granulocytes in the thymus[42,43]. However, if this occurs in unmanipulated

mice at a steady-state remains unclear.

The strict spatial segregation of the two macrophage populations in the

thymus implies that they might have distinct functions. *Timd4*[+] cells are

restricted to the cortex and are particularly abundant in the deeper cortex,

close to the medulla. Both positive and negative selection of thymocytes occur

there, so we speculate that *Timd4*[+] macrophages might be specialized in

efferocytosis of CD4[+]CD8[+] (double-positive) thymocytes that cannot interact

with cortical thymic epithelial cells and die by neglect or are auto-reactive and

undergo clonal deletion in the cortex[61]. On the other hand, *Cx3cr1*[+]

macrophages accumulate in the medulla; the thymic region specialized in

negative selection to tissue-restricted antigens (TRA). They might contribute

to the process in several ways: 1) by carrying TRAs from blood and peripheral

organs. A similar process has been described for cDC2 (SIRPα^+ DCs)[62]. In

fact, *Cx3cr1$^+$* thymic macrophages could have contributed to this role because

they were not distinguished from cDC2 in this study. 2) By capturing TRAs

from *Aire$^+$* medullary thymic epithelial cells and presenting them to auto-

reactive thymocytes as shown for DCs[63-65]. 3) By phagocytosing apoptotic

TRA-specific medullary thymocytes, a process we have observed before[58].

The exact involvement of thymic macrophages in the selection events in the

thymus remains to be determined.

The accumulation of the *Cx3cr1$^+$* cells in older mice has clear implications

for thymus aging. One key feature of thymus involution is the accumulation of

extracellular matrix produced by fibroblasts and the emergence of white

adipocytes[66]. The *Cx3cr1$^+$* subset is the predominant producer of the growth

factor PDGFα that is required for the maintenance of adipocyte stem cells and

can stimulate tissue fibrosis[67,68]. The gradual accumulation of *Cx3cr1$^+$*

macrophages could increase the availability PDGFα in the aging thymus

stimulating extracellular matrix production and differentiation of precursors

into adipocytes. This model predicts that limiting the influx of *Cx3cr1$^+$*

macrophage precursors could delay thymus involution.

Recent work described a novel phagocytic and antigen-presenting cell type

in the thymus called monocyte-derived DCs[65]. The phenotype of these cells

overlaps with the CD64$^+$F4/80^{lo}CD11b$^+$ cells in our study. However, we favor

the classification of these cells as monocytes based on their expression of

Mafb, CD64, and Ly6C (Fig. 5b) and lack of expression of the defining DC

transcription factor *Zbtb46* (Fig. S2c)[69]. As monocytes can differentiate into

cDC2, particularly in the context of inflammation[70], the precise identity and

the relationship of this population to thymic cDC2 remain to be established.

In the past several years, scRNA-Seq has come to the forefront of

biologists' efforts to disentangle the cellular diversity of tissues. Several

comprehensive studies have included samples from mouse or human

thymus[71-73]. However, in these studies, too few thymic macrophages were

sampled to give meaningful clustering results. Efforts specifically targeting

the thymus have provided considerably more information[74,75], but

macrophage diversity was still not recognized. Characterization of rare

populations such as thymic macrophages (~0.1% of all cells in the thymus)

requires optimized enzymatic digestion procedures and enrichment strategies,

as has been demonstrated already for thymic epithelial cells[76,77]. Our

scRNA-Seq dataset provides a rich resource for the unbiased characterization

of myeloid cells in the thymus and will greatly aid in the understanding of the

myeloid landscape of the thymus.

In summary, our work comprehensively characterizes macrophages in the

thymus and paves the way for exploration of their functions.

Materials and methods

Mice

C57BL/6Narl (CD45.2) mice were purchased from the National Laboratory

Animal Center, Taipei, Taiwan. MAFIA (MAcrophage Fas-Induced Apoptosis)

[78], *Cx3cr1^{GFP}* [79], *Spic^{GFP}* [52], *Cx3cr1^{CreER}* [53], and *B6.SJL-Ptprca*

Pepcb/BoyJ (CD45.1) mice were purchased from the Jackson Laboratories.

Cd11c^{YFP} [80] and *Lyz2^{GFP}* [81] mice have been described. Mice ubiquitously

expressing GFP from the *ROSA26* locus were generated by breeding

Pdgfra^{Cre} [82] and *ROSA26^{LSL-ZsGreen}* (also known as *ROSA26^{LSL-GFP}* or Ai6)

mice[83] (both from the Jackson Laboratories). A mouse from this cross was

identified, in which the STOP cassette was deleted in the germline. It was

designated *ROSA26^{GFP}* and subsequently bred to C57BL/6 mice. All mice

were used at 4-10 weeks of age unless otherwise specified. Mice were bred

and maintained under specific pathogen-free conditions at the animal facility

of National Yang-Ming University. All experimental procedures were approved

by the Institutional Animal Care and Use Committee (IACUC) of National

Yang-Ming University.

Treatment with 5-Ethynyl-2'-deoxyuridine (EdU)

Mice were i.p. injected with 1 mg EdU (Carbosynth) dissolved in PBS daily

for 21 days and then rested for 21 more days. Cells from the thymus were

harvested on day 21 or 42. In some experiments, the mice were sacrificed 2

hours after the first EdU injection.

Shield chimera generation

CD45.1 mice were anesthetized by i.p. injection of 120 μg/g body weight

Ketamine hydrochloride (Toronto Research Chemicals) and 12 μg/g body

weight Xylazine hydrochloride (Sigma). Anesthetized mice were taped to a 5

cm thick lead block so that the lead block covered the head and the chest

down to the bottom of the rib cage. Then, they were irradiated with a lethal

dose (1000 rad) from a ^{137}Cs source (Minishot II, AXR) so that only their

abdomen and hind legs were exposed. After recovery from anesthesia, the

mice were transfused i.v. with 10^7 bone marrow cells from a congenic

(CD45.2) donor. Then, they were given Trimerin (0.5 mg/mL Sulfadiazine +

0.1 mg/mL Trimethoprim, China Chemical and Pharmaceutical Co., Tainan,

Taiwan) in the drinking water for the first two weeks after the irradiation and

analyzed after six weeks.

Cell isolation from thymus, blood, and peritoneal cavity

Thymocytes were obtained by mechanical disruption of the thymus with a

syringe plunger. For myeloid cell isolation, mouse thymuses were cut into

small pieces and digested with 0.2 mg/mL DNase I (Roche) and 0.2 mg/mL

collagenase P (Roche) in complete DMEM for 20 min at 37˚C with frequent

agitation. 1.6 mg/mL Dispase I (ThermoFisher) was added to the enzymatic

mixture for thymic epithelial cell isolation. In some experiments, thymic

myeloid cells and thymic epithelial cells were enriched by 57% Percoll PLUS

(GE Healthcare) discontinuous gradient centrifugation at 4˚C, 1800 rpm, for

20 min without brake. Cells at the interface were collected and washed with

PBS to remove residual silica particles. Then the cells were resuspended in

PBS with 0.5% BSA (HM Biological), filtered through a 70 µm filter, and kept

on ice.

Blood was isolated by cardiac puncture of sacrificed mice and immediately

diluted with PBS. After centrifugation, the cell suspensions were treated with

ammonium chloride-potassium lysis buffer for 3 min on ice once or twice.

Peritoneal cavity cells were obtained by lavage with 5 mL PBS + 2 mM EDTA

(Merck). Following gentle massage, the cavity was opened with an abdominal

incision, and lavage fluid was collected.

Flow cytometry

Single-cell suspensions ($0.5 - 2X10^6$ cells) from thymus, blood, or

peritoneal cavity were blocked with supernatant from 2.4G2 hybridoma (a kind

gift by Dr. Fang Liao, Academia Sinica, Taipei, Taiwan) and stained with

fluorochrome- or biotin-labeled antibodies for 20 min on ice in PBS + 0.5%

BSA + 2 mM EDTA + 0.1% NaN_3 (FACS buffer). The following antibodies

were used: CD11b (clone M1/70), MHC2 (M5/114.15.2), CD11c (N418),

F4/80 (BM8), CD115 (AFS98), SIRPα (P84), CD45 (30-F11), NK1.1 (PK136),

TIM4 (RMT4-54), Gr-1 (RB6-8C5), CD64 (X54-5/7.1), Siglec H (551), Ly6C

(HK1.4), CD3ε (145-2C11), CD8α (53-6.7), CD19 (6D5), B220 (RA3-6B2),

CD4 (GK1.5), CD51 (RMV-7), CD45.1 (A20), CD45.2 (104), CX3CR1

(SA011F11), and EpCAM (G8.8) from BioLegend; Axl (MAXL8DS), MerTK

(DS5MMER), and Ki67 (SolA15) were from eBioscience; Siglec F (E50-2440),

CD90.2 (30-H12), and CD11c (HL3) were from BD Biosciences. Cells were

washed, and if necessary, incubated for 20 more min with fluorochrome-

labeled Streptavidin: Streptavidin-AF647 (Jackson Immunoresearch) or

Streptavidin-APC/cy7, Streptavidin-BV421, Streptavidin-BV605 (BioLegend).

After the last wash, the cells were resuspended in FACS buffer containing

DAPI (BioLegend), Propidium Iodide (Sigma), or DRAQ7 (BioLegend) and

analyzed immediately on an LSR Fortessa flow cytometer running Diva 8

software (BD Biosciences). Typically, 500,000 cells were collected from

thymus samples. Data were analyzed using FlowJo software (TreeStar).

For intracellular staining, after surface antibody staining, the cells were

labeled with Zombie Aqua (BioLegend) for 30 min in ice. Then, the cells were

fixed with 2% paraformaldehyde (Electron Microscope Sciences) in PBS for

20 min on ice, permeabilized with either 0.5% Triton-X 100 (Sigma) for 20 min

on ice, or with Foxp3 staining kit (eBioscience) according to the protocol

provided by the manufacturer, and stained with antibodies for intracellular

markers for 40-60 min on ice.

For cell cycle analysis, $1-5 \times 10^5$ sorted thymic macrophages were fixed with

70% ethanol for 2 h on ice. The cells were spun down at 1800 rpm for 20 min

at 4˚C, washed with PBS, and stained with 1 μg/ml DAPI (BioLegend) for 30

min at room temperature in the dark.

For EdU staining, after surface marker and Zombie Aqua staining, cells

were fixed with 2% paraformaldehyde in PBS for 20 min on ice and

permeabilized with 0.5% Triton X-100 in PBS at room temperature for 20 min.

EdU was detected by adding an equal volume of 2X Click reaction buffer

consisting of 200 mM Tris, 200 mM ascorbic acid (Acros), 8 mM $CuSO_4$

(Acros), 8 μM Cy5-azide (Lumiprobe) to the permeabilized cells resuspended

in 0.5% Triton X-100 in PBS and incubation at room temperature for 30 min.

Cells were washed twice with 0.5% Triton X-100 in PBS and analyzed on a

flow cytometer.

Cell sorting

The sorting of thymic macrophages was done following the IMMGEN

guidelines. Briefly, the thymuses of 3 male C57BL/6Narl mice were harvested

in ice-cold staining buffer containing phenol red-free DMEM (Gibco) with 10

mM HEPES (Sigma), 0.1% NaN$_3$, and 2% FBS (Gibco). Single-cell

suspensions were prepared as described in the Flow cytometry section.

Percoll PLUS was used to enrich mononuclear cells. The cells were

resuspended at 10^8/mL in staining buffer and labeled with appropriate

antibodies for 15 min in ice. To sort thymic macrophages, the cells were first

labeled with biotinylated antibodies to lineage markers (Lin) – CD3, CD8, Gr1,

B220. After washing, the cells were stained with antibodies to CD11b, F4/80,

CD45, CD64, and Streptavidin-APC/cy7 for 15 min in ice. For sorting thymus

XCR1$^+$ and SIRPα^+ cDCs, antibodies to XCR1, SIRPα, CD11c, MHC2, CD64,

and F4/80 were used. For sorting peritoneal cavity macrophages, antibodies

to ICAM2 and F4/80 were used. Immediately before sorting, the dead cells

were excluded with DRAQ7 or PI. For RNA Sequencing experiments, the

cells were double-sorted on FACS Melody, or Aria cell sorters (BD

Biosciences) and 1000 cells were directly deposited in TCL buffer (Qiagen),

frozen in dry ice and sent to IMMGEN for RNA sequencing. Four biological

replicates were prepared. For cytospin and cell cycle analysis, 1-5X10^5 cells

sorted on FACS Melody were collected in staining buffer.

Cytospin

Sorted cells were mounted on Superfrost PLUS slides (Thermo Scientific)

using a Cytospin centrifuge (Cytospin 3, Shandon) for 5 min at 500 rpm. Cells

were fixed with 2% paraformaldehyde for 10 min at room temperature and

stained with the Hemacolor Rapid Staining Kit (Merck Millipore). Images were

collected on BX61 upright microscope (Olympus) using 100X objective with

immersion oil and captured with a CCD camera. Images were then analyzed

and processed with ImageJ (NIH) and Adobe Photoshop 5.5 (Adobe).

RNA sequencing analysis

RNA sequencing was done at IMMGEN using Smart-seq2 protocol[84,85]

on a NextSeq500 sequencer (Illumina). Following sequencing, raw reads

were aligned with STAR to the mouse genome assembly mm10 and assigned

to specific genes using the GENCODE vM12 annotation. Gene expression

was normalized by DESeq2[86] and visualized by Morpheus

(https://software.broadinstitute.org/morpheus). Hierarchical clustering was

done with Cluster 3.0 and visualized with Java TreeView. Only genes with

SD>20 were used (10602 genes). The metric used was Pearson correlation

(uncentered), and the clustering method was average linking. Gene

expression of mouse transcription factors[87] was visualized in MultiplotStudio

of GenePattern[88]. GO enrichment was calculated and visualized in R by

using clusterProfiler[89].

Timed pregnancies and embryonic thymus analysis

To set up timed pregnancies, each male mouse ($Cx3cr1^{CreER/CreER}$,

$Cx3cr1^{GFP/GFP}$ or C57BL/6) and female mouse ($ROSA26^{LSL-GFP/LSL-GFP}$ or

C57BL/6) were housed together in the same cage for one night and separated

on the next day, which we defined as embryonic day 0.5 (E0.5). Female mice

were assumed to be pregnant if their weight gain was over 2 g at E8.5[90].

Thymuses from E14.5 and E17.5 embryos, neonatal, 1-weeks-old pups, and

adult mice (older than 6-weeks-old) were harvested, mechanically dissociated

with plastic sticks in 1.5-mL centrifuge tubes, and enzymatically digested with

0.2 mg/mL DNase I and 0.2 mg/mL collagenase P in complete DMEM for 20

min at 37˚C with frequent agitation. The cells were resuspended in PBS with

0.5% BSA, filtered through a 70 µm filter, kept on ice, and used flow

cytometric analysis as described in the Flow Cytometry section.

Genetic fate mapping – E9.5, neonatal and adult

For genetic fate mapping, timed pregnancies of *Cx3cr1*$^{CreER/CreER}$ male and

ROSA26$^{LSL-GFP/LSL-GFP}$ female mice were set up as described. To label the

Cx3cr1$^+$ erythromyeloid progenitors derived from embryonic yolk sac[8], 4-

hydroxytamoxifen (4-OHT from Sigma) was administered i.p. to pregnant

females on E9.5 at a dose of 75 µg/g (body weight). To improve the survival

of embryos and reduce the risk of abortions, Progesterone (Sigma) was co-

injected at a dose of 37.5 µg/g (body weight)[91]. To label the *Cx3cr1$^+$* thymic

macrophages in *Cx3cr1*^{CreER} X *ROSA26*$^{LSL-GFP}$ neonates and adult mice,

Tamoxifen (TAM from Sigma) was injected i.p. at a dose of 2 mg/mouse to

lactating dams on postnatal day 3 and 4 (P3 and P4) or to adult mice for 2

consecutive days. Thymuses were harvested and analyzed 3 days or 6

weeks after the last injection by flow cytometry.

scRNA-Seq – sorting, library generation, and sequencing

scRNA-Seq was performed at the Genomics Center for Clinical and Biotechnological Applications of NCFB (NYCU, Taipei, Taiwan). Briefly, the thymuses of one female MAFIA and 2 male $Cd11c^{YFP}$ mice were harvested and enzymatically digested as described previously. Mononuclear cells were enriched by 57% Percoll PLUS discontinuous centrifugation, washed to remove silica particles, and resuspended at 10^6/mL in PBS with 0.04% BSA. The cell suspensions were filtered through Falcon 35 µm strainer (Corning) and stained with viability dye (PI or DAPI) immediately before sorting. Cell sorting was performed on a FACS Melody sorter (BD Biosciences) running FACS Chorus (BD Biosciences) software in purity mode. $3X10^5$ GFP or YFP positive cells under the live/singlet gating were collected into 5 ml round bottom tubes pre-coated with 0.04% BSA in PBS. Sorted cells were washed and resuspended in 300 µL PBS with 0.04% BSA and then filtered again into 1.5-mL DNA LoBind tubes (Eppendorf) through a 35 µm strainer. The viability of the cells was evaluated by Countess II (Invitrogen) and Trypan Blue (ThermoFisher), and samples with cell viability rates higher than 85% were used for encapsulation and library preparation. Single-cell encapsulation and library preparation were performed using Single Cell 3' v3/v3.1 Gene Expression solution (10x Genomics). All the libraries were processed according to the manufacturer's instruction and sequenced on NovaSeq 6000 (Illumina) platform at the NHRI (Zhubei, Taiwan). Post-processing and quality control were performed by the NYCU Genome Center using the CellRanger package (v. 3.0.2, 10x Genomics). Reads were aligned to mm10 reference assembly. Primary assessment with CellRanger reported 9,973 cell-barcodes

with 11,385 median unique molecular identifiers (UMIs, transcripts) per cell

and 3,076 median genes per cell sequenced to 71.0% sequencing saturation

with 94,260 mean reads per cell for MAFIA mouse sample; 9,801 cell-

barcodes with 13,467 median UMIs per cell and 3,211 median genes per cell

sequenced to 74.9% sequencing saturation with 119,820 mean reads per cell

for the first $Cd11c^{YFP}$ mouse sample; 12,938 cell-barcodes with 14,439

median UMIs per cell and 3,199 median genes per cell sequenced to 71.4%

sequencing saturation with 108,585 mean reads per cell for the second

$Cd11c^{YFP}$ mouse sample.

Analysis of scRNA-Seq

Preprocessing

The Scanpy[92] pipeline was used to read the count matrix. Three batches

of samples (one from GFP+ cells from MAFIA mouse and two from YFP+ cells

from $Cd11c^{YFP}$ mice) were preprocessed independently and integrated later.

Cells that expressed <200 genes and genes that were expressed in <3 cells

were filtered out. The percentage of mitochondrial genes was calculated -and

cells with >10% mitochondrial genes were removed. Cells with >7,000 genes

or <1,000 genes were also removed. Read counts were normalized to library

size 10,000 and log-transformed with scanpy.pp.log1p function.

Datasets integration and batch effect correction

Read count matrices and spliced/unspliced matrices were merged first.

Principal Component Analysis was applied to reduce dimensions to 70.

BBKNN[93] was then used to remove batch effects with the

scanpy.external.pp.bbknn function with default parameters.

Visualization and clustering

UMAP[94] provided by scanpy was used to visualize data with default

parameters. K-nearest neighbor and Leiden clustering were applied

sequentially to cluster cells into groups. K-nearest neighbor graph

construction was done by scanpy.pp.neighbors with parameters

n_neighbors=12 and n_pcs=70. Leiden clustering was then performed by

scanpy.tl.leiden with parameter resolution=0.15. To improve UMAP

visualization, scanpy.tl.paga was applied, and we trimmed unnecessary graph

edges by scanpy.tl.paga with threshold=0.018.

Marker genes and statistics

Wilcoxon rank-sum tests were applied to examine differentially expressed

genes. Clusters were selected from the result of Leiden clustering.

Differentially expressed genes of a cluster against other clusters were

identified by scanpy.tl.rank_genes_groups and scanpy.pl.rank_genes_groups.

P-values were collected for each cluster and transformed by negative $\log_{10}$ for

better visualization. The top 50 differentially expressed genes were visualized

in the figure.

Immunofluorescent staining

Dissected thymus lobes from C57BL/6 mice were cleaned of connective

tissue and fixed in 4% paraformaldehyde (Sigma) for 1 h at 4°C, washed in

PBS, submerged in 10% sucrose, and then in 30% sucrose for 12 h each.

The tissue was then frozen in Tissue-Tek OCT compound (Sakura Fintek) for

cryostat sectioning. 10 or 20 µm thick sections were prepared with CryoStar

NX50 (ThermoFisher) on Superfrost PLUS (ThermoScientific) microscope

slides, dried overnight, and stored at -80˚C until used. Before staining, the

sections were fixed with acetone (Sigma) at -20˚C for 10 min, air-dried, then

blocked with 5% goat serum + 5% donkey serum (both from Jackson

Immunoresearch) in PBS for 2 h and stained with primary antibodies: rat

monoclonal to MerTK (DS5MMER, eBioscience), rat monoclonal to TIM4

(RMT4-54, Bio-X-Cell) or rabbit polyclonal to Keratin 5 (BioLegend) overnight

at 4˚C in a humidified chamber. After washing in PBS, the sections were

labeled with goat anti-rat-Alexa Fluor 647 (Invitrogen) or goat anti-rat Cy3

(Jackson Immunoresearch) and donkey anti-rabbit AF647 (Jackson

Immunoresearch) secondary antibodies for 2 hours at room temperature,

followed by 5 min staining with DAPI. TUNEL Assay was done with the Click-

iT Plus TUNEL Assay Alexa Fluor 647 kit (Invitrogen) according to the

manufacturer's recommendations. A positive (pre-incubation with DNase I for

30 min at room temperature) and negative (no TdT enzyme) controls were

always included. The sections were mounted with 0.1% n-propyl gallate

(Sigma) in glycerol (Sigma) and imaged with an AxioObserver 7 (Carl Zeiss)

wide-field microscope equipped with Plan Apochromat 20x NA=0.8 objective

(Zeiss) and AxioCam 702 mono camera (Zeiss) and controlled by Zen 2.3

Blue (Zeiss) software. Image analysis was performed with Imaris 8.0.2

(Bitplane).

The co-localization scoring for MerTK and TIM4 with TUNEL was done with

Imaris 8.2 (Bitplane). TUNEL+ cells were detected with the Spots function,

while MerTK+ and TIM4+ cells were detected with the Surface function. Spots

that co-localize with Surfaces were identified with the "Find Spots close to

Surface" function of Imaris XT. The threshold for co-localization was set to 5

µm. The results were manually curated so that Spots categorized as "not co-

localized" that were: 1) at the edge of the imaging field were excluded from

consideration; 2) with clear Surface signal around them were re-categorized

as "co-localized". The ratio of co-localized Spots to all Spots was calculated

and presented as the co-localization index.

Thymus transplantation

To obtain E15.5 embryos, *ROSA26^{GFP}* homozygous male and C57BL/6

female mice were mated in a cage overnight and separated on the next day.

Pregnant mice were sacrificed 15 days later, the viable embryos were

harvested, and the thymuses were isolated in ice-cold PBS. C57BL/6

recipients were anesthetized by i.p injection of Ketamine hydrochloride (120

µg/g, Toronto Research Chemicals) and Xylazine hydrochloride (12 µg/g,

Sigma). The fur on the left flank was removed, and the left kidney was

exposed by cutting the skin, muscle layer, and peritoneum. The kidney

capsule was nicked with a G23 needle, and the fetal thymus was pushed into

the pocket under the kidney capsule with a G23 needle equipped with a

plunger from a spinal needle. After the kidney was re-positioned back into the

peritoneal cavity, the peritoneum was sutured, and the skin was stapled with

metal clips. Rymadil (Carprofen, 5 µg/g, Zoetis) was given subcutaneously to

ease the wound pain, and Trimerin (Sulfadiazine at 0.5 mg/mL + Trimethoprim

at 0.1 mg/mL) were given in the drinking water for the first two weeks after the

surgery. The metal clips were removed from the skin after the first week, and

the transplanted thymus and recipient's endogenous thymus were harvested

and analyzed six weeks after the kidney transplantation.

1001

1002   **Statistical analysis**

1003   Comparison between groups was made with Prism 6 (GraphPad Software).

1004   Comparisons between two groups were carried out with unpaired Student's t-

1005   test. When more than two groups were compared, a one-way ANOVA with

1006   Tukey correction was used. Differences were considered significant if $p<0.05$.

1007

Data availability

The RNA Sequencing data of thymic macrophages and thymic dendritic

cells is available at NCBI Gene Expression Omnibus (GEO) as part of

GSE122108 and at www.immgen.org. The single cell RNA sequencing data is

deposited at NCBI GEO under accession number GSE185460. The source

data underlying Fig. 1g-h, Fig. 3c, f, i, Fig. 4b. f, h, Fig. 6b, d, Fig. 7b, d, e, Fig.

S1h, Fig. S3c, and Fig. S4b, d, e, g are provided in the Source Data files. All

other data supporting the findings of this study are available within the article

and its figures and tables.

1017

Abbreviations:

cDC – classical dendritic cell

DC – dendritic cell

EdU – 5-Ethynyl-2'-deoxyuridine

GO – gene ontology

HSC – hematopoietic stem cell

IFN-I – type I Interferon

IMMGEN – Immunological Genome Consortium

scRNA-Seq – single-cell RNA sequencing

TF – transcription factor

ThyMacs – thymic macrophages

TRA – tissue-restricted antigen

YS – yolk sac

References:

1.   Wynn TA, Chawla A, Pollard JW. Macrophage biology in development,
homeostasis and disease. Nature. Nature Publishing Group; 2013;496:
445–455. doi:10.1038/nature12034

2.   Baratin M, Simon L, Jorquera A, Ghigo C, Dembele D, Nowak J, et al. T
Cell Zone Resident Macrophages Silently Dispose of Apoptotic Cells in
the Lymph Node. Immunity. Elsevier Inc; 2017;47: 349–362.e5.
doi:10.1016/j.immuni.2017.07.019

3.   A-Gonzalez N, Castrillo A. Origin and Specialization of Splenic
Macrophages. Cellular Immunology. 2018;330: 1–27.
doi:10.1016/j.cellimm.2018.05.005

4.   Bellomo A, Mondor I, Spinelli L, Lagueyrie M, Stewart BJ, Brouilly N, et
al. Reticular Fibroblasts Expressing the Transcription Factor WT1
Define a Stromal Niche that Maintains and Replenishes Splenic Red
Pulp Macrophages. Immunity. Elsevier Inc; 2020;53: 127–142.e7.
doi:10.1016/j.immuni.2020.06.008

5.   van Furth R, Cohn ZA. The origin and kinetics of mononuclear
phagocytes. J Exp Med. Rockefeller University Press; 1968;128: 415–
435. doi:10.1084/jem.128.3.415

6.   Ginhoux F, Guilliams M. Tissue-Resident Macrophage Ontogeny and
Homeostasis. Immunity. 2016;44: 439–449.
doi:10.1016/j.immuni.2016.02.024

7.   Perdiguero EG, Klapproth K, Schulz C, Busch K, Azzoni E, Crozet L, et
al. Tissue-resident macrophages originate from yolk-sac-derived
erythro-myeloid progenitors. Nat Immunol. Nature Publishing Group;
2015;518: 547–551. doi:10.1038/nature13989

8.   Mass E, Ballesteros I, Farlik M, Halbritter F, Günther P, Crozet L, et al.

Specification of tissue-resident macrophages during organogenesis.
Science. American Association for the Advancement of Science;
2016;353: aaf4238–aaf4238. doi:10.1126/science.aaf4238

9.   Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, et al. Fate
mapping analysis reveals that adult microglia derive from primitive
macrophages. Science. 2010;330: 841–845.
doi:10.1126/science.1194637

10.   Hoeffel G, Wang Y, Greter M, See P, Teo P, Malleret B, et al. Adult
Langerhans cells derive predominantly from embryonic fetal liver
monocytes with a minor contribution of yolk sac–derived macrophages.
J Exp Med. 2012;209: 1167–1181. doi:10.1084/jem.20120340

11.   Hoeffel G, Chen J, Lavin Y, Low D, Almeida FF, See P, et al. C-Myb(+)
erythro-myeloid progenitor-derived fetal monocytes give rise to adult
tissue-resident macrophages. Immunity. 2015;42: 665–678.
doi:10.1016/j.immuni.2015.03.011

12.   Goldmann T, Wieghofer P, Jordão MJC, Prutek F, Hagemeyer N,
Frenzel K, et al. Origin, fate and dynamics of macrophages at central
nervous system interfaces. Nat Immunol. 2016;17: 797–805.
doi:10.1038/ni.3423

13.   Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, et al.
Tissue-Resident Macrophages Self-Maintain Locally throughout Adult
Life with Minimal Contribution from Circulating Monocytes. Immunity.
Elsevier Inc; 2013;38: 792–804. doi:10.1016/j.immuni.2013.04.004

14.   Epelman S, Lavine KJ, Beaudin AE, Sojka DK, Carrero JA, Calderon B,
et al. Embryonic and Adult-Derived Resident Cardiac Macrophages Are
Maintained through Distinct Mechanisms at Steady State and during
Inflammation. Immunity. Elsevier Inc; 2014;40: 91–104.
doi:10.1016/j.immuni.2013.11.019

15.   Sheng J, Ruedl C, Karjalainen K. Most Tissue-Resident Macrophages
Except Microglia Are Derived from Fetal Hematopoietic Stem Cells.
Immunity. Elsevier Inc; 2015;43: 382–393.
doi:10.1016/j.immuni.2015.07.016

16.   Liu Z, Gu Y, Chakarov S, Bleriot C, Kwok I, Chen X, et al. Fate Mapping
via Ms4a3-Expression History Traces Monocyte-Derived Cells. Cell.
Elsevier; 2019;178: 1509–1525.e19. doi:10.1016/j.cell.2019.08.009

17.   Ensan S, Li A, Besla R, Degousee N, Cosme J, Roufaiel M, et al. Self-
renewing resident arterial macrophages arise from embryonic
CX3CR1+ precursors and circulating monocytes immediately after birth.
Nat Immunol. Nature Publishing Group; 2015;17: 159–168.
doi:10.1038/ni.3343

18.   Mondor I, Baratin M, Lagueyrie M, Saro L, Henri S, Gentek R, et al.

Lymphatic Endothelial Cells Are Essential Components of the
Subcapsular Sinus Macrophage Niche. Immunity. Elsevier Inc; 2019;50:
1453–1466.e4. doi:10.1016/j.immuni.2019.04.002

19. Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et
al. Progressive replacement of embryo-derived cardiac macrophages
with age. J Exp Med. 2014;211: 2151–2158. doi:10.1084/jem.20140639

20. Heidt T, Courties G, Dutta P, Sager HB, Sebas M, Iwamoto Y, et al.
Differential Contribution of Monocytes to Heart Macrophages in Steady-
State and After Myocardial Infarction. Circulation Research. 2014;115:
284–295. doi:10.1161/CIRCRESAHA.115.303567

21. Yahara Y, Barrientos T, Tang YJ, Puviindran V, Nadesan P, Zhang H,
et al. Erythromyeloid progenitors give rise to a population of osteoclasts
that contribute to bone homeostasis and repair. Nat Cell Biol. Springer
US; 2020;22: 1–28. doi:10.1038/s41556-019-0437-8

22. Jacome-Galarza CE, Percin GI, Muller JT, Mass E, Lazarov T, Eitler J,
et al. Developmental origin, functional maintenance and genetic rescue
of osteoclasts. Nature. Nature Publishing Group; 2019;568: 541–545.
doi:10.1038/s41586-019-1105-7

23. Calderon B, Carrero JA, Ferris ST, Sojka DK, Moore L, Epelman S, et
al. The pancreas anatomy conditions the origin and properties of
resident macrophages. J Exp Med. 2015;212: 1497–1512.
doi:10.1084/jem.20150496

24. Tamoutounour S, Guilliams M, Sanchis FM, Liu H, Terhorst D, Malosse
C, et al. Origins and Functional Specialization of Macrophages and of
Conventional and Monocyte-Derived Dendritic Cells in Mouse Skin.
Immunity. Elsevier Inc; 2013;39: 925–938.
doi:10.1016/j.immuni.2013.10.004

25. Bain CC, Bravo-Blas A, Scott CL, Gomez Perdiguero E, Geissmann F,
Henri S, et al. Constant replenishment from circulating monocytes
maintains the macrophage pool in the intestine of adult mice. Nat
Immunol. 2014;15: 929–937. doi:10.1038/ni.2967

26. Sierro F, Evrard M, Rizzetto S, Melino M, Mitchell AJ, Florido M, et al. A
Liver Capsular Network of Monocyte-Derived Macrophages Restricts
Hepatic Dissemination of Intraperitoneal Bacteria by Neutrophil
Recruitment. Immunity. Elsevier Inc; 2017;47: 374–388.e6.
doi:10.1016/j.immuni.2017.07.018

27. Mossadegh-Keller N, Gentek R, Gimenez G, Bigot S, Mailfert S,
Sieweke MH. Developmental origin and maintenance of distinct
testicular macrophage populations. J Exp Med. 2017;214: 2829–2841.
doi:10.1084/jem.20170829

28. Lokka E, Lintukorpi L, Cisneros-Montalvo S, Mäkelä J-A, Tyystjärvi S,

Ojasalo V, et al. Generation, localization and functions of macrophages
during the development of testis. Nature Communications. Springer US;
2020;11: 1–16. doi:10.1038/s41467-020-18206-0

29. Wang M, Yang Y, Cansever D, Wang Y, Kantores C, Messiaen S, et al.
Two populations of self-maintaining monocyte-independent
macrophages exist in adult epididymis and testis. PNAS. 2020;118:
e2013686117. doi:10.1073/pnas.2013686117/-/DCSupplemental

30. Bain CC, Hawley CA, Garner H, Scott CL, Schridde A, Steers NJ, et al.
Long-lived self-renewing bone marrow-derived macrophages displace
embryo-derived cells to inhabit adult serous cavities. Nature
Communications. Nature Publishing Group; 2016;7: 1–14.
doi:10.1038/ncomms11852

31. Surh CD, Sprent J. T-cell apoptosis detected in situ during positive and
negative selection in the thymus. Nature. 1994;372: 100–103.
doi:10.1038/372100a0

32. Esashi E, Sekiguchi T, Ito H, Koyasu S, Miyajima A. Cutting Edge: A
possible role for CD4+ thymic macrophages as professional scavengers
of apoptotic thymocytes. Journal of Immunology (Baltimore, Md : 1950).
American Association of Immunologists; 2003;171: 2773–2777.
doi:10.4049/jimmunol.171.6.2773

33. Liu L-T, Lang Z-F, Li Y, Zhu Y-J, Zhang J-T, Guo S-F, et al.
Composition and characteristics of distinct macrophage subpopulations
in the mouse thymus. Molecular Medicine Reports. 2013;7: 1850–1854.
doi:10.3892/mmr.2013.1448

34. Guerri L, Peguillet I, Geraldo Y, Nabti S, Premel V, Lantz O. Analysis of
APC types involved in CD4 tolerance and regulatory T cell generation
using reaggregated thymic organ cultures. J Immunol. American
Association of Immunologists; 2013;190: 2102–2110.
doi:10.4049/jimmunol.1202883

35. Lopes N, Charaix J, Cédile O, Sergé A, Irla M. Lymphotoxin alpha fine-
tunes T cell clonal deletion by regulating thymic entry of antigen-
presenting cells. Nature Communications. Springer US; 2018;9: 1–16.
doi:10.1038/s41467-018-03619-9

36. Kim H-J, Alonzo ES, Dorothee G, Pollard JW, Sant'Angelo DB.
Selective depletion of eosinophils or neutrophils in mice impacts the
efficiency of apoptotic cell clearance in the thymus. Kim H-J, Alonzo ES,
Dorothee G, Pollard JW, Sant'Angelo DB, editors. PLoS ONE. 2010;5:
e11439. doi:10.1371/journal.pone.0011439.g010

37. Tacke R, Hilgendorf I, Garner H, Waterborg C, Park K, Nowyhed H, et
al. The transcription factor NR4A1 is essential for the development of a
novel macrophage subset in the thymus. Sci Rep. 2015;5: 10055.
doi:10.1038/srep10055

38.   Gautier EL, Shay T, Miller J, Greter M, Jakubzick C, Ivanov S, et al.
1183   Gene-expression profiles and transcriptional regulatory pathways that
1184   underlie the identity and diversity of mouse tissue macrophages. Nat
1185   Immunol. 2012;13: 1118–1128. doi:10.1038/ni.2419

1186   39.   Ingersoll MA, Spanbroek R, Lottaz C, Gautier EL, Frankenberger M,
1187   Hoffmann R, et al. Comparison of gene expression profiles between
1188   human and mouse monocyte subsets. Blood. 2010;115: e10–e19.
1189   doi:10.1182/blood-2009-07-235028

1190   40.   Wang H, Breed ER, Lee YJ, Qian LJ, Jameson SC, Hogquist KA.
1191   Myeloid cells activate iNKT cells to produce IL-4 in the thymic medulla.
1192   Proc Natl Acad Sci. National Academy of Sciences; 2019;116: 22262–
1193   22268. doi:10.1073/pnas.1910412116

1194   41.   Chan CT, Fenn AM, Harder NK, Mindur JE, McAlpine CS, Patel J, et al.
1195   Liver X receptors are required for thymic resilience and T cell output. J
1196   Exp Med. 2020;217: 245–23. doi:10.1084/jem.20200318

1197   42.   Wada H, Masuda K, Satoh R, Kakugawa K, Ikawa T, Katsura Y, et al.
1198   Adult T-cell progenitors retain myeloid potential. Nature. Nature
1199   Publishing Group; 2008;452: 768–772. doi:10.1038/nature06839

1200   43.   Bell JJ, Bhandoola A. The earliest thymic progenitors for T cells
1201   possess myeloid lineage potential. Nature. 2008;452: 764–767.
1202   doi:10.1038/nature06840

1203   44.   Schlenner SM, Madan V, Busch K, Tietz A, LAufle C, Costa C, et al.
1204   Fate Mapping Reveals Separate Origins of T Cells and Myeloid
1205   Lineages in the Thymus. Immunity. Elsevier Ltd; 2010;32: 426–436.
1206   doi:10.1016/j.immuni.2010.03.005

1207   45.   Witmer-Pack MD, Hughes DA, Schuler G, Lawson L, McWilliam A,
1208   Inaba K, et al. Identification of macrophages and dendritic cells in the
1209   osteopetrotic (op/op) mouse. J Cell Sci. 1993;104 (Pt 4): 1021–1029.

1210   46.   Sasmono RT, Oceandy D, Pollard JW, Tong W, Pavli P, Wainwright BJ,
1211   et al. A macrophage colony-stimulating factor receptor-green
1212   fluorescent protein transgene is expressed throughout the mononuclear
1213   phagocyte system of the mouse. Blood. American Society of
1214   Hematology; 2003;101: 1155–1163. doi:10.1182/blood-2002-02-0569

1215   47.   Hume DA. Applications of myeloid-specific promoters in transgenic mice
1216   support in vivo imaging and functional genomics but do not support the
1217   concept of distinct macrophage and dendritic cell lineages or roles in
1218   immunity. Journal of Leukocyte Biology. 2011;89: 525–538.
1219   doi:10.1189/jlb.0810472

1220   48.   Taniuchi I, Osato M, Egawa T, Sunshine MJ, Bae SC, Komori T, et al.
1221   Differential requirements for Runx proteins in CD4 repression and
1222   epigenetic silencing during T lymphocyte development. Cell. 2002;111:

621–633. doi:10.1016/s0092-8674(02)01111-x

49. Ebihara T, Song C, Ryu SH, Plougastel-Douglas B, Yang L, Levanon D,
et al. Runx3 specifies lineage commitment of innate lymphoid cells. Nat
Immunol. Nature Publishing Group; 2015;16: 1124–1133.
doi:10.1038/ni.3272

50. Fainaru O, Woolf E, Lotem J, Yarmus M, Brenner O, Goldenberg D, et
al. Runx3 regulates mouse TGF-beta-mediated dendritic cell function
and its absence results in airway inflammation. EMBO J. John Wiley &
Sons, Ltd; 2004;23: 969–979. doi:10.1038/sj.emboj.7600085

51. Kohyama M, Ise W, Edelson BT, Wilker PR, Hildner K, Mejia C, et al.
Role for Spi-C in the development of red pulp macrophages and splenic
iron homeostasis. Nat Immunol. Nature Publishing Group; 2008;457:
318–321. doi:10.1038/nature07472

52. Haldar M, Kohyama M, So AY-L, KC W, Wu X, Briseño CG, et al.
Heme-Mediated SPI-C Induction Promotes Monocyte Differentiation into
Iron-Recycling Macrophages. Cell. Elsevier Inc; 2014;156: 1223–1234.
doi:10.1016/j.cell.2014.01.069

53. Yona S, Kim K-W, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate
Mapping Reveals Origins and Dynamics of Monocytes and Tissue
Macrophages under Homeostasis. Immunity. Elsevier; 2013;38: 79–91.
doi:10.1016/j.immuni.2012.12.001

54. McCaughtry TM, Wilken MS, Hogquist KA. Thymic emigration revisited.
J Exp Med. 2007;204: 2513–2520. doi:10.1084/jem.20070601

55. Lienenklaus S, Cornitescu M, Ziętara N, Łyszkiewicz M, Gekara N,
Jabłónska J, et al. Novel reporter mouse reveals constitutive and
inflammatory expression of IFN-beta in vivo. J Immunol. American
Association of Immunologists; 2009;183: 3229–3236.
doi:10.4049/jimmunol.0804277

56. Otero DC, Baker DP, David M. IRF7-dependent IFN-β production in
response to RANKL promotes medullary thymic epithelial cell
development. J Immunol. American Association of Immunologists;
2013;190: 3289–3298. doi:10.4049/jimmunol.1203086

57. Breed ER, Lee ST, Hogquist KA. Directing T cell fate: How thymic
antigen presenting cells coordinate thymocyte selection. Seminars in
Cell and Developmental Biology. Elsevier Ltd; 2017;: 1–9.
doi:10.1016/j.semcdb.2017.07.045

58. Kurd NS, Lutes LK, Yoon J, Chan SW, Dzhagalov IL, Hoover AR, et al.
A role for phagocytosis in inducing cell death during thymocyte negative
selection. eLife. 2019;8: 1879–18. doi:10.7554/eLife.48097

59. Morioka S, Perry JSA, Raymond MH, Medina CB, Zhu Y, Zhao L, et al.
Efferocytosis induces a novel SLC program to promote glucose uptake

and lactate release. Nature. Springer US; 2018;563: 1–21.
doi:10.1038/s41586-018-0735-5

60. Bleriot C, Chakarov S, Ginhoux F. Determinants of Resident Tissue
Macrophage Identity and Function. Immunity. Elsevier Inc; 2020;52:
957–970. doi:10.1016/j.immuni.2020.05.014

61. Stritesky GL, Xing Y, Erickson JR, Kalekar LA, Wang X, Mueller DL, et
al. Murine thymic selection quantified using a unique method to capture
deleted T cells. Proc Natl Acad Sci. 2013;110: 4679–4684.
doi:10.1073/pnas.1217532110

62. Baba T, Nakamoto Y, Mukaida N. Crucial Contribution of Thymic Sirp +
Conventional Dendritic Cells to Central Tolerance against Blood-Borne
Antigens in a CCR2-Dependent Manner. J Immunol. 2009;183: 3053–
3063. doi:10.4049/jimmunol.0900438

63. Gallegos AM, Bevan MJ. Central Tolerance to Tissue-specific Antigens
Mediated by Direct and Indirect Antigen Presentation. J Exp Med.
2004;200: 1039–1049. doi:10.1084/jem.20041457

64. Koble C, Kyewski B. The thymic medulla: a unique microenvironment
for intercellular self-antigen transfer. J Exp Med. 2009;206: 1505–1513.
doi:10.1084/jem.20082449

65. Vobořil M, Brabec T, Dobeš J, Šplíchalová I, Březina J, Čepková A, et
al. Toll-like receptor signaling in thymic epithelium controls monocyte-
derived dendritic cell recruitment and Treg generation. Nature
Communications. Springer US; 2020;11: 1–16. doi:10.1038/s41467-
020-16081-3

66. Dixit VD. Impact of immune-metabolic interactions on age-related
thymic demise and T cell senescence. Semin Immunol. Elsevier Ltd;
2012;24: 321–330. doi:10.1016/j.smim.2012.04.002

67. Rivera-Gonzalez GC, Shook BA, Andrae J, Holtrup B, Bollag K,
Betsholtz C, et al. Skin Adipocyte Stem Cell Self-Renewal Is Regulated
by a PDGFA/AKT-Signaling Axis. Stem Cell. Elsevier Inc; 2016;19:
738–751. doi:10.1016/j.stem.2016.09.002

68. Olson LE, Soriano P. Increased PDGFRα Activation Disrupts
Connective Tissue Development and Drives Systemic Fibrosis.
Developmental Cell. Elsevier Ltd; 2009;16: 303–313.
doi:10.1016/j.devcel.2008.12.003

69. Satpathy AT, KC W, Albring JC, Edelson BT, Kretzer NM, Bhattacharya
D, et al. Zbtb46expression distinguishes classical dendritic cells and
their committed progenitors from other immune lineages. J Exp Med.
2012;209: 1135–1152. doi:10.1084/jem.20120030

70. Guilliams M, Mildner A, Yona S. Developmental and Functional
Heterogeneity of Monocytes. Immunity. Elsevier Inc; 2018;49: 595–613.

1305   doi:10.1016/j.immuni.2018.10.005

71.   Tabula Muris Consortium. Single-cell transcriptomics of 20 mouse
organs creates a Tabula Muris. Nature. Nature Publishing Group;
2018;562: 367–372. doi:10.1038/s41586-018-0590-4

72.   Han X, Wang R, Zhou Y, Fei L, Sun H, Lai S, et al. Mapping the Mouse
Cell Atlas by Microwell-Seq. Cell. Elsevier Inc; 2018;172: 1091–
1097.e17. doi:10.1016/j.cell.2018.02.001

73.   Tabula Muris Consortium. A single-cell transcriptomic atlas
characterizes ageing tissues in the mouse. Nature. Nature Publishing
Group; 2020;583: 590–595. doi:10.1038/s41586-020-2496-1

74.   Kernfeld EM, Genga RMJ, Neherin K, Magaletta ME, Xu P, Maehr R. A
Single-Cell Transcriptomic Atlas of Thymus Organogenesis Resolves
Cell Types and Developmental Maturation. Immunity. Elsevier Inc;
2018;: 1–33. doi:10.1016/j.immuni.2018.04.015

75.   Park J-E, Botting RA, Domínguez Conde C, Popescu D-M, Lavaert M,
Kunz DJ, et al. A cell atlas of human thymic development defines T cell
repertoire formation. Science. American Association for the
Advancement of Science; 2020;367: eaay3224–12.
doi:10.1126/science.aay3224

76.   Bornstein C, Nevo S, Giladi A, Kadouri N, Pouzolles M, Gerbe F, et al.
Single-cell mapping of the thymic stroma identifies IL-25-producing tuft
epithelial cells. Nature. Springer US; 2018;559: 1–25.
doi:10.1038/s41586-018-0346-1

77.   Bautista JL, Cramer NT, Miller CN, Chavez J, Berrios DI, Byrnes LE, et
al. Single-cell transcriptional profiling of human thymic stroma uncovers
novel cellular heterogeneity in the thymic medulla. Nature
Communications. Nature Publishing Group; 2021;12: 1096–15.
doi:10.1038/s41467-021-21346-6

78.   Burnett SH, Kershen EJ, Zhang J, Zeng L, Straley SC, Kaplan AM, et
al. Conditional macrophage ablation in transgenic mice expressing a
Fas-based suicide gene. Journal of Leukocyte Biology. Society for
Leukocyte Biology; 2004;75: 612–623. doi:10.1189/jlb.0903442

79.   Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A,
et al. Analysis of fractalkine receptor CX(3)CR1 function by targeted
deletion and green fluorescent protein reporter gene insertion. Mol Cell
Biol. 2000;20: 4106–4114.

80.   Lindquist RL, Shakhar G, Dudziak D, Wardemann H, Eisenreich T,
Dustin ML, et al. Visualizing dendritic cell networks in vivo. Nat
Immunol. 2004;5: 1243–1250. doi:10.1038/ni1139

81.   Faust N, Varas F, Kelly LM, Heck S, Graf T. Insertion of enhanced
green fluorescent protein into the lysozyme gene creates mice with

green fluorescent granulocytes and macrophages. Blood. 2000;96:
719–726.

82.   Roesch K, Jadhav AP, Trimarchi JM, Stadler MB, Roska B, Sun BB, et
al. The transcriptome of retinal Müller glial cells. J Comp Neurol.
2008;509: 225–238. doi:10.1002/cne.21730

83.   Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et
al. A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. Nature Neuroscience. Nature
Publishing Group; 2009;13: 133–140. doi:10.1038/nn.2467

84.   Picelli S, Faridani OR, rklund ASKBO, Winberg GOS, Sagasser S,
Sandberg R. Full-length RNA-seq from single cells using Smart-seq2.
Nature Protocols. Nature Publishing Group; 2014;9: 171–181.
doi:10.1038/nprot.2014.006

85.   Picelli S, Björklund ÅK, Faridani OR, Sagasser S, Winberg G, Sandberg
R. Smart-seq2 for sensitive full-length transcriptome profiling in single
cells. Nat Meth. 2013;10: 1096–1098. doi:10.1038/nmeth.2639

86.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. BioMed
Central; 2014;15: 31–21. doi:10.1186/s13059-014-0550-8

87.   Schmeier S, Alam T, Essack M, Bajic VB. TcoF-DB v2: update of the
database of human and mouse transcription co-factors and transcription
factor interactions. Nucleic Acids Res. 2017;45: D145–D150.
doi:10.1093/nar/gkw1007

88.   Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP.
GenePattern 2.0. Nat Genet. Nature Publishing Group; 2006;38: 500–
501. doi:10.1038/ng0506-500

89.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. OMICS: A
Journal of Integrative Biology. Mary Ann Liebert, Inc., publishers;
2012;16: 284–287. doi:10.1089/omi.2011.0118

90.   Heyne GW, Plisch EH, Melberg CG, Sandgren EP, Peter JA, Lipinski
RJ. A Simple and Reliable Method for Early Pregnancy Detection in
Inbred Mice. J Am Assoc Lab Anim Sci. 2015;54: 368–371.

91.   Iturri L, Saenz Coronilla J, Lallemand Y, Gomez Perdiguero E.
Identification Of Erythromyeloid Progenitors And Their Progeny In The
Mouse Embryo By Flow Cytometry. JoVE. 2017;: 1–10.
doi:10.3791/55305

92.   Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol. BioMed Central; 2018;19: 15–
5. doi:10.1186/s13059-017-1382-0

93. Polański K, Young MD, Miao Z, Meyer KB, Teichmann SA, Park J-E.
BBKNN: fast batch alignment of single cell transcriptomes.
Bioinformatics. 2019;36: 964–965.
doi:10.1093/bioinformatics/btz625#supplementary-data

94. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation
and Projection for Dimension Reduction. ArXiv e-prints. 2018.
doi:1802.03426v3

1393

Acknowledgments

We are grateful to the IMMGEN consortium for performing the RNA

sequencing of our samples and providing access to its database. We thank

the following core facilities at National Yang-Ming Chiao-Tung University:

Instrumentation Resource Center for providing access to its sorting and

imaging facility and the Animal Facility for mouse husbandry. We are very

grateful to Wan-Chun Chang, Bing-Xiu Guo, Chien-Yi Tung, and Kate Hua

from the Genomics Center for Clinical and Biotechnological Applications of

NCFB (NYCU, Taipei) for help with scRNA-Seq. Special thanks go to Dr.

Fang Liao, Academia Sinica, Taipei, for help with cell sorting and for providing

the 24G2 hybridoma, and Chang-Feng Chu for technical assistance. We are

very grateful to Ms. Su-Hua Jiang at Veterans General Hospital, Taipei, for

help with using the irradiator. Some figures were made with BioRender.

This work was supported by the Ministry of Science and Technology

(grants # 106-2320-B-010 -026 -MY3, and 107-2320-B-010 -016 -MY3 to IL

Dzhagalov and grants # 107-2320-B-010-020, and 108-2628-B-010-005 to CL

Hsu), the Yen-Tjing Ling Medical Foundation (grants # CI-107-6 and CI-108-5

to IL Dzhagalov), and the Cancer Progression Research Center (NYCU). The

authors declare no competing financial interests.

Author contributions

TA Zhou designed experiments, performed research, analyzed and

interpreted data, and wrote the manuscript; HP Hsu and CY Lin performed

research, analyzed and interpreted data; YH Tu and HC Huang analyzed the

scRNA-Seq data; NJ Chen, JW Tsai, EA Robey, and CL Hsu provided

expertise, shared critical reagents, and edited the manuscript; IL Dzhagalov

conceptualized the studies, designed experiments, performed research,

analyzed, interpreted the data, and wrote the manuscript.

1421

1422

Supplementary figures and tables

Figure 1 – Figure supplement 1: Phenotype of thymic macrophages

and other myeloid cells positive for F4/80 and CD64. **a** Example flow

cytometry plots breaking down F4/80$^+$ cells in the thymus into macrophages

(MerTK$^+$), eosinophils (CD11b$^+$MerTK$^-$Siglec F$^+$) and monocytes

(CD11b$^+$MerTK$^-$Siglec F$^-$). **b** An example flow cytometry plot dividing CD64+

cells into macrophages (F4/80$^+$CD11b^{lo}) and monocytes (CD11b$^+$F4/80^{lo}). **c**

Representative flow cytometry staining of enzymatically digested thymus

single-cell suspension for CD64, MerTK, and F4/80 and respective isotype

controls. **d** Gating strategy routinely used to identify thymic macrophages as

CD64$^+$F4/80$^+$CD11b^{lo}FSC^{hi} cells. **e** Example flow cytometry plots showing that

CD64$^+$FSC^{lo} cells include Thy1$^+$ cells (most likely thymocytes). **f** Expression of

CD11c, MHC2, and SIRPα on ThyMacs with respective controls. **g** Example

flow cytometry plots showing that gating on CD11c$^+$MHC2$^+$ thymus cells, in

addition to DCs, also includes macrophages, especially among SIRPα$^+$ cells.

h Frequency of MerTK$^+$ cells among CD11c$^+$MHC2$^+$SIRPα$^+$ cells. The data

are mean±SEM from 5 individual mice. Each dot is an individual mouse. The

flow cytometry plots are representative of ≥5 individual experiments.

Figure 1 – Figure Supplement 2: UMAP clustering of single-cell RNA-

Sequencing (scRNA-Seq) data. a Clustering of the scRNA-Seq data shows

that the cells from the three samples (one from GFP$^+$ cells in MAFIA mice and

two from YFP$^+$ cells in *Cd11c^{YFP}* mice) overlap considerably. **b** Identification of

the clusters from the scRNA-Seq data based on lineage-specific markers. **c**

Expression of lineage-specific markers in different clusters.

1448

Figure 1 – Figure Supplement 3: Most TUNEL⁺ apoptotic cells co-

localize with thymic macrophages. a Example images showing co-

localization of TUNEL⁺ apoptotic cells and MerTK⁺ ThyMacs. **b** Example

images showing co-localization of TUNEL⁺ apoptotic cells and TIM4⁺

ThyMacs. Scale bars in **a** and **b** are 50 µm. **c** Frequencies of the co-

localization of TUNEL⁺ signal with MerTK⁺ and TIM4⁺ cells. All

immunofluorescent images are representative of at least 3 independent

repeats. Data in **c** represent mean±SEM. Each dot is an individual mouse.

Figure 2 – Figure Supplement 1: Expression of *Spic^GFP* marks most

thymic macrophages. a Example of the gating strategy to identify ThyMacs

among *Spic^GFP+* cells. **b** Frequencies of ThyMacs among *Spic^GFP+* cells. **c**

Representative flow cytometry plots of the expression of four reporter alleles

in ThyMacs. **d** Frequencies of GFP/YFP+ cells among ThyMacs. **e**

Frequencies of ThyMacs among GFP/YFP+ cells. **f** Representative flow

cytometry plots of the expression of *Spic^GFP* in ThyMacs. **g** Frequencies of

Spic^GFP+ cells among ThyMacs. All flow cytometry plots are representative of

at least 3 independent repeats. Data in **b**, **d**, **e**, and **f** represent mean±SEM.

Each dot is an individual mouse. The numbers in the flow cytometry plots are

the percent of cells in the respective gate.

1468

Table S1: E**xpression of differentially up-regulated transcription factors in thymic macrophages**

Gene name	ThyMacs	non-ThyMacs
Irf7	3879.32	300.82
Irf8	3528.27	1474.35
Stat1	2403.69	522.04
Dnmt3a	1515.94	647.81
Znxf1	1379.89	635.36
Stat2	1210.35	472.53
Nr1h3	1182.17	147.05
Srebf1	975.09	399.06
Rxra	760.26	298.55
Trps1	746.36	232.48
Runx3	723.14	9.76
Relb	715.53	293.92
Sp100	696.94	324.47
Zbp1	639.19	69.83
Tfec	588.72	74.66
Spic	573.11	34.36
Nfkbie	569.74	226.76
Ncoa4	550.69	249.15
Rest	548.22	269.22
Meis3	530.8	120.91
Bhlhe40	490.59	99.56
Parp12	414.11	126.82
Arid5b	374.03	177.08
Creb5	295.14	47.91
Pparg	276.54	33.24

Figure 2 – Table Supplement 1: Expression of differentially up-

regulated transcription factors in thymic macrophages. Transcription

factors that were highly expressed in thymic macrophages (>250) and up-

regulated >2-fold in thymic macrophages compared to non-thymic

macrophages were listed alphabetically, and the geometric mean of 4

replicates of thymic macrophages (ThyMacs) and two replicates of each of the

9 non-thymic macrophage populations (non-ThyMacs) were recorded. Non-

thymic macrophages are: spleen red pulp macrophages, Kupffer cells,

bronchoalveolar lavage macrophages, peritoneal cavity macrophages, aorta

macrophages, heart macrophages, white adipose tissue macrophages,

central nervous system microglia, spinal cord macrophages.

Table S2. **List of the differentially expressed genes among *Timd*4+ thymic macrophages, *Cx3cr1*+ thymic macrophages, and thymic monocytes**

CX3CR1+ ThyMacs		Tim4+ ThyMacs		ThyMonos	
Gene name	adjusted p-value	Gene name	adjusted p-value	Gene name	adjusted p-value
Ctsz	0	*Hpgd*	0	*Alox5ap*	0
Cd63	0	*Serpinb6a*	0	*S100a6*	0
Pmepa1	0	*Slc40a1*	0	*Ly6c2*	0
Zmynd15	0	*Cd81*	0	*Ifi27l2a*	0
Olfml3	0	*Vcam1*	0	*Fau*	0
Mmp2	0	*Cfp*	0	*Coro1a*	0
AU020206	1.60E-290	*Spic*	0	*Ccr2*	0
Plxnd1	1.59E-285	*Trf*	0	*Rps27*	0
Cst7	8.68E-279	*Actn1*	0	*Tmsb10*	0
Dnase1l3	2.45E-270	*Maf*	0	*Ifitm2*	7.21E-302
Timp2	2.15E-267	*Pld3*	0	*Fxyd5*	6.36E-299
Lgals3bp	8.69E-263	*Il18*	0	*Rps19*	2.04E-292
Pdgfa	6.87E-255	*Mrc1*	0	*Rpl18*	6.50E-291
Mmp14	2.33E-253	*Crip2*	0	*Rpl9*	1.11E-289
Fam46c	9.99E-235	*Tmem65*	0	*Rps23*	1.28E-289
Chst2	1.19E-226	*Igf1*	0	*Napsa*	8.91E-279
Cp	5.36E-225	*Epb41l3*	0	*Ms4a4c*	8.25E-272
Camk1	7.12E-225	*Timd4*	0	*Plac8*	2.10E-270
B2m	1.09E-222	*Blvrb*	0	*Rpl18a*	9.26E-269
Lhfpl2	4.52E-217	*Clec1b*	0	*S100a4*	4.98E-268
Acp5	5.90E-216	*Cd68*	0	*Cd52*	3.67E-267
Lag3	3.91E-213	*Axl*	0	*Rps14*	1.94E-266
Lyz2	1.28E-209	*Paqr9*	3.32E-307	*Ifitm3*	3.19E-263
H2-M2	1.22E-199	*Sdc3*	3.45E-305	*Rpl34*	2.02E-261
Psap	7.26E-198	*Myo9a*	5.59E-305	*Rps27a*	3.67E-260
Gatm	1.33E-192	*Scp2*	3.79E-302	*Rpl36*	1.54E-259
Cpd	1.50E-192	*Selenop*	2.10E-295	*Rps16*	2.55E-258

C3	2.34E-187	Lrp1	2.08E-294	Rpl24	1.37E-257
Cxcl16	8.11E-183	Lap3	1.45E-290	Rps9	6.34E-253
Lgals3	1.57E-182	Marcks	2.77E-279	Gpr141	1.21E-246
Ube2j1	1.63E-180	Glul	3.64E-279	Rpl27a	3.06E-243
Plxnc1	9.84E-180	Hebp1	3.76E-278	Rpl17	8.15E-241
Stab1	4.07E-176	Ear2	4.53E-276	Rps24	1.46E-240
Cyth1	3.27E-163	Apoc1	2.49E-275	Rps13	2.34E-236
Spsb1	3.96E-163	Kcna2	3.72E-275	Rpl38	1.95E-226
Blnk	2.35E-162	Myo10	9.05E-269	H2-DMb1	1.02E-223
Cx3cr1	9.29E-162	Atp13a2	2.95E-267	Rps18	5.39E-223
Med10	5.25E-161	Slc1a3	6.24E-263	Rpl19	3.68E-221
Nek6	5.28E-160	Slco2b1	1.11E-258	Rpl8	2.01E-219
Ptms	1.05E-159	mt-Nd2	3.45E-258	Rpl7a	4.17E-217
Anxa5	1.10E-156	Wwp1	2.16E-253	Gm34084	5.23E-216
Gpnmb	1.21E-154	Aplp2	4.22E-248	Rpl13	2.08E-215
Itgb5	2.78E-154	Atp8a1	5.03E-248	Rpl11	2.47E-213
Myo5a	1.11E-146	P2ry13	3.17E-247	Rpl35a	2.13E-210
Runx3	1.81E-146	Ccdc148	4.70E-245	Rpsa	1.62E-209
Tmem176a	2.34E-144	Grn	1.58E-244	Rpl6	5.70E-208
Ctss	4.81E-141	Bank1	1.82E-239	Tpt1	2.63E-206
Sh3pxd2b	9.38E-141	Mertk	2.15E-238	Rack1	2.14E-203
Rtcb	4.42E-140	Nr1h3	1.13E-235	Rpl23	6.14E-199
Fam20c	1.91E-139	Prnp	2.93E-235	Rpl26	7.48E-198
Il2rg	8.84E-138	Ninj1	2.42E-234	Rps6	6.64E-197
Lpcat2	8.53E-137	Fcna	3.33E-233	Rps10	2.06E-195
Kynu	8.49E-136	Csrp1	1.16E-230	Ier5	1.06E-191
Tnfsf13b	8.77E-136	Rgl1	7.18E-229	Rps3	8.23E-185
Gpr157	1.18E-135	Lpl	4.94E-223	Rpl27	8.23E-185
Tgfbr1	7.63E-135	Fam213b	1.08E-222	Rps5	8.36E-185
H2-K1	1.15E-133	Tcf7l2	1.26E-222	Rps7	3.96E-182
Basp1	1.23E-133	AB124611	4.64E-221	Rps15a	6.82E-182
Pla2g7	1.80E-132	Abcc3	3.28E-216	Rps11	1.97E-180
Fth1	4.19E-131	Fcgrt	5.79E-216	Rps4x	5.07E-180
Ggh	1.85E-126	Tgm2	1.88E-215	Rplp0	3.09E-177
Adam19	6.94E-126	Itgad	5.35E-214	Ly6i	8.17E-176
C3ar1	7.35E-125	Ptgs1	2.94E-213	S100a11	6.23E-175
Ccl12	3.37E-123	Laptm4a	1.01E-212	Atox1	1.22E-174
Hvcn1	2.51E-121	Comt	1.33E-206	Pim1	9.56E-174
Anxa3	8.60E-121	Creg1	3.24E-205	Sh3bgrl3	3.97E-173
Tgfbi	1.88E-120	Adgre1	9.67E-205	Ciita	7.35E-173
Ctsd	2.73E-117	Clec12a	6.33E-204	Eef1a1	6.09E-172
Itm2c	5.19E-116	Tspan4	7.80E-203	Rps3a1	9.09E-168
Tmem119	5.62E-116	Txn1	9.13E-203	Gm2a	6.07E-165
Rap2a	1.03E-114	Ctsb	9.52E-201	Ptprc	2.05E-163
Ctsl	4.00E-114	Mrap	5.65E-197	Rpl37	1.51E-161
Itga6	1.83E-113	Slc16a9	5.99E-197	Rps25	3.03E-160

B4galnt1	2.45E-113	Abcg3	3.83E-196	H3f3a	5.92E-159
Fam3c	1.64E-112	Pla2g15	4.22E-196	Btg2	1.14E-158
Tmem173	1.54E-111	C1qc	6.17E-192	Rpl15	1.42E-158
Ski	3.59E-111	Agpat3	1.68E-191	Cnn2	1.09E-156
Anpep	5.85E-111	Hs6st1	1.95E-191	Cdkn1a	2.57E-156
Gng2	2.37E-110	Dmpk	2.15E-191	Slfn1	4.83E-155
Nceh1	2.88E-110	Cd38	1.79E-190	Sem1	4.08E-154
H2-Q7	4.94E-108	Tmem26	2.02E-189	Lsp1	1.34E-152
Rtn1	1.28E-106	Slc11a1	1.05E-188	Rpl37a	1.78E-152
Sorl1	1.31E-103	Cd300a	1.41E-187	Rpl22	3.64E-152
Glipr1	1.22E-102	Slc7a7	3.28E-187	Sirpb1c	4.81E-152
Gsn	2.00E-102	Cyb5a	6.94E-187	Traf1	6.97E-152
Afdn	4.54E-102	Sipa1l1	7.41E-187	Emb	4.22E-151
Ak2	1.11E-101	Il18bp	1.48E-186	Rpl30	1.32E-147
Ntpcr	2.21E-98	Cd86	2.52E-183	Rps15	1.14E-146
Scarb2	3.16E-97	Vamp5	3.05E-183	H2-Ab1	2.84E-145
Creb5	5.41E-97	Jup	6.69E-182	Il1b	3.05E-145
Gsto1	5.56E-97	Blvra	1.30E-178	Rps28	4.52E-145
Ncf1	4.26E-96	Mgst1	6.48E-178	Jarid2	1.82E-143
Ppfia4	4.97E-96	Tbxas1	1.47E-177	Rps26	1.53E-142
Chchd10	7.77E-96	Hpgds	2.04E-177	Rpl32	4.21E-142
Gna12	1.23E-95	Tgfbr2	2.70E-176	Pld4	9.07E-142
Mvb12b	1.80E-95	Clec4n	3.52E-175	Cbfa2t3	1.54E-141
Rasal3	1.45E-94	Ms4a7	5.30E-175	Rps21	4.04E-141
Scoc	6.86E-94	Sirpa	3.35E-171	Fgr	4.04E-141
Cfb	6.00E-93	Fyn	2.84E-168	Rps8	1.11E-139
Lmna	1.04E-92	Cadm1	2.20E-167	Cd74	5.34E-138

1480

1481 **Figure 5 – Table Supplement 1: List of the differentially expressed**

1482 **genes among *Timd4*+ thymic macrophages, *Cx3cr1*+ thymic**

1483 **macrophages, and thymic monocytes.** The top 100 differentially expressed

1484 genes among the three clusters are listed by their negative $\log_{10}$ transformed

1485 p-value.

1486

1487