TITLE:

Trachymyrmex septentrionalis **ant microbiome assembly is unique to individual colonies**

and castes.

AUTHORS: Emily A. Green [1], Jonathan L. Klassen [1,2,3]

AFFILIATIONS: University of Connecticut

[1] Department of Molecular and Cell Biology, University of Connecticut, Storrs, Connecticut,

USA

[2] Institute for Systems Genomics, University of Connecticut, Storrs, Connecticut, USA

[3] To whom correspondence should be addressed: 91 N Eagleville Rd, Unit 3125, Storrs, CT,

06269-3125. jonathan.klassen@uconn.edu

Emily A. Green: https://orcid.org/0000-0002-0824-6369

Jonathan L. Klassen PhD: https://orcid.org/0000-0003-1745-8838

ABSTRACT:

Within social insect colonies, microbiomes often differ between castes due to their

different functional roles, and between colony locations. *Trachymyrmex septentrionalis* fungus-

growing ants form colonies throughout the eastern USA and Northern Mexico that include

workers, female and male alates (unmated reproductive castes), larvae, and pupae. How *T.*

septentrionalis microbiomes vary across this geographic range and between castes is unknown.

Our sampling of individual ants from colonies across the Eastern USA revealed a conserved core

T. septentrionalis worker ant microbiome, and that worker ant microbiomes are more conserved

within colonies than between them. A deeper sampling of individual ants from two colonies that

included all available castes (pupae, larvae, workers, female and male alates), from both before

and after adaptation to controlled laboratory conditions, revealed that ant microbiomes from each

colony, caste, and rearing condition were typically conserved within but not between each

sampling category. Tenericute bacterial symbionts were especially abundant in these ant

microbiomes and varied widely in abundance between sampling categories. This study

demonstrates how individual insect colonies primarily drive the composition of their

microbiomes, and that these microbiomes are further modified by developmental differences

between insect castes and the different environmental conditions experienced by each colony.

INTRODUCTION:

Social insects live in colonies and manifest group integration, division of labor, and

overlap of generations [1]. These divisions of labor are often distributed between developmental

castes, which at the most basic level are divided into reproductive individuals and sterile

workers, which themselves can be further partitioned into specialized groups for different tasks.

Gut microbiomes often differ between social insect castes, likely due to the specialized

functional role played by each caste [2]. For example, the microbiomes of termite minor

workers, major workers, and soldiers all differ from each other [3, 4], and termite reproductive

castes have unique microbiomes compared to those of workers from the same colony [5–8].

Honey bee queens, workers, and drones also each have unique gut microbiomes, where worker

microbiomes are more diverse than those of queens and drones, likely due to worker foraging

[9]. Honey bees have a well-defined core microbiome that is found in all colonies and castes,

including queens, workers, and drones [10, 11]. However, strains varied between geographic

locations, individual colonies, and bee castes [9, 11]. Social insect microbiome composition is

therefore determined by a complex combination of developmental differences and colony-

specific ecological factors.

Fungus-growing ants are social insects (Formicidae: tribe Attini) that evolved 55 to 65

MYA in South America [12–14]. Today there are 19 genera including approximately 250 known

species that range throughout North and South America from the northern USA to southern

Argentina [14–18]. The ~250 species of fungus-growing ants are separated into five separate

agricultural systems representing transitions in their evolution: lower, coral-fungus, yeast, higher,

and leaf-cutter agriculture [14]. Fungus-growing ant caste systems include workers, reproductive

queens, and unmated reproductive alates. Leaf-cutting ants (*Atta, Acromyrmex*, and

Amiomyrmex) can have multiple worker castes, including soldier, major, minor, and minum

workers [16]. All fungus-growing ants grow a cultivar fungus, typically a species of

Leucoagaricus (except for in coral-fungus agriculture by *Apterostigma* ants, which cultivates a

Pterulaceae species [19]). This cultivar serves as the ants' main food source and has coevolved

with the ants for millions of years [12, 13, 15, 16]. Ants forage for materials to feed to their

cultivar fungus (e.g., grass, leaves, dried plant material, and/or insect frass), which the fungus

degrades and converts to fungal biomass, including hyphal swellings called gonglydia that the

ants eat as their primary food source [20, 21]. *Pseudonocardia*, an Actinobacterium, grows on

the cuticle of many fungus-growing ants, producing antimicrobials to protect the fungus garden

from pathogens [22, 23]. Other cuticular microbes on the ants have been identified but their

function is unknown [23, 24]. Together these interactions comprise a multipartite symbiosis

composed of the fungus-growing ant, the *Leucoagaricus* fungus, and the protective symbiont

Pseudonocardia.

Several studies have preliminarily investigated the fungus-growing ant microbiome,

primarily focusing on leaf-cutting ants [25–29]. These studies show that *Acromyrmex* and *Atta*

leaf-cutting ants host low diversity microbiomes that include *Wolbachia*, *Solirubrobacter*,

Enterobacter, *Pseudomonas*, and members of the orders Entomoplasmatales and Rhizobiales.

Rhizobiales bacteria contain *nif* genes that were hypothesized to provide nitrogen cycling for the

ants and their cultivar fungus [29]. *Wolbachia* is maternally transmitted in *Acromyrmex* and *Atta*

but lacks a known function. Entomoplasmatales bacteria belong to the genera *Mesoplasma* and

Spiroplasma are thought to provide nutrients to the ants [28, 30]. Most of these studies pooled

multiple worker ants together before DNA extraction, leaving how ant microbiomes differ

between individuals within a colony and between castes largely unknown.

Here, we characterize the microbiome of the fungus-growing ant *Trachymyrmex*

septentrionalis. Phylogenetically, the higher agriculture ant genus *Trachymyrmex* holds a

transitional position adjacent to leaf-cutting ants [17]. Typical *T. septentrionalis* colonies have

approximately 1,000 workers in addition to a single queen, and during the summer months they

also produce reproductive female and male alates that remain in the colony before mating as a

mass swarm [16]. *T. septentrionalis* colonies are found in pine barren and sandy soil ecosystems

throughout the eastern USA and northern Mexico [20, 31, 32], and vary genetically across the

United States, forming four different clades [32]. Ishak et al [33] sampled *T. septentrionalis* ants

from 25 colonies at a single location in Texas over one year and identified 19 bacterial genera

that were conserved in *T. septentrionalis* workers and alates (both males and females). They also

identified differences in ant microbiomes between castes, albeit including only a few samples.

However, the samples collected from each colony were not differentiated from each other,

making it impossible to study intra-colony variation in microbiome composition. Whether the 19

common bacteria found in Texas *T. septentrionalis* and form a core microbiome that is

conserved in other geographic regions or castes is also unknown.

In contrast to this previous work, we sampled individual ants collected across a broad

geographic gradient to characterize the core microbiome of field-collected *T. septentrionalis*

worker ants. We also more deeply sampled individual ant microbiomes within two colonies and

found that ant gut microbiome composition differs between colonies, between ant castes, and

following adaptation of colonies to the laboratory environment. The variable presence of

Mesoplasma and *Spiroplasma* symbionts in different ant colonies and castes was a major driver

of these microbiome differences. Our study provides an example of how individual colonies,

castes, and ecological differences determine the composition of social insect microbiomes.

METHODS

Sample collection

Trachymyrmex septentrionalis colonies were collected in New Jersey, New York, North

Carolina, Florida, Georgia, and Louisiana during 2014–2018 (Suppl. File 1; Suppl. Fig. S1).

Permits for *T. septentrionalis* collection include: the New Jersey Department of Environmental

Protection Division of Parks and Forestry State Park Service unnumbered letters of

authorization; New York State Department of Environmental Conservation License to Collect or

Possess: Scientific #915; Permit for Research in Suffolk County; North Carolina Division of

Parks and Recreation Scientific Collection and Research Permit 2015_0030; Florida Department

of Agriculture and Consumer Services unnumbered letters of authorization; Georgia Department

of Natural Resources State Parks and Historic Sites Scientific Research and Collection Permit

032015; and Louisiana Department of Wildlife and Fisheries Permit WL-Research2016-10, in

addition to permits from the United States Department of Agriculture for the transportation of

ants to the University of Connecticut (USDA; P526P-14-00684). Freshly collected ants

(hereafter: "field-collected ants") were stored in DESS (20% dimethyl sulphoxide, 250 mM

disodium EDTA, and saturated sodium chloride) [34] on dry ice immediately after colony

collection in the field and then stored at -80 °C in the lab until sampling for DNA extraction. The

remainder of each colony (including both fungus gardens and the remaining ants) was brought

back to the University of Connecticut, kept in a USDA-approved and temperature-controlled

room, and maintained following Sosa-Calvo et al [35]. Colonies were placed in a 6-3/4" long x 4

13/16" width x 2-3/8″ height box that was lined with plaster of Paris, watered biweekly to

maintain humidity, and provided with sterile cornmeal *ad libitum* as food for the fungus garden.

Ants sampled from these colonies will be referred to as "lab-maintained".

We generated two major datasets for this study. The first, which we will refer to as the

"multi-state" dataset, included field-collected workers from all of the states mentioned above.

This dataset was designed to test if ant microbiomes varied throughout the ants' geographic

range. The second, which we will refer to as the "two-colony" dataset, was made up of multiple

ants sampled from Colony JKH000270, collected from Friendship Park in July 2017, and Colony

JKH000307, collected from Quaker Bridge Park in July 2018. We sampled all available castes

and life stages, including workers, pupae, larvae, male alates, and female alates, to test if

microbiomes differed between castes and following lab adaptation. Worker ants collected from

the lab-maintained colonies were sampled immediately before DNA extraction without storage

in DESS. Two confirmatory datasets were also used, with one including pupae and another

dissected ant guts and whole ants sampled from the same colony.

Sample preparation & DNA extraction

All ants were surface-sterilized using a 10 second submersion in 70% ethanol followed

by a 10 second submersion in phosphate-buffered saline (PBS), repeated three times [36]. Some

pupae were ethanol-washed and others were not to determine if the Tenericute symbionts were

located on their exoskeleton or internally. DNA was extracted from these surface-sterilized ants

using the bead-beating and chloroform-isopropanol protocol described previously [34, 37]. Our

preliminary data showed minimal differences between microbiomes generated using whole

surface-sterilized ants and ant guts, and so we used whole, surface-sterilized ants throughout this

study for simplicity (Suppl. Fig. S2) while recognizing that actinobacteria in our dataset

(predominantly *Phycicoccus* and *Pseudonocardia*) may be due to incomplete surface

sterilization. Negative controls containing only the DNA extraction reagents were processed

alongside each batch of ant samples. The DNA concentration of each extract and negative

control was determined using the Qubit dsDNA high-sensitivity assay protocol and a Qubit 3.0

fluorimeter (Invitrogen, Carlsbad CA).

PCR screening & community amplicon sequencing

DNA samples were PCR amplified using primers 515F and 806R to amplify the 16S

rRNA gene V4 region [38]. Ten nanograms of template DNA was added to 5 µl Green GoTaq

Reaction Mix Buffer (Promega, Madison, WI), 1.25 units GoTaq DNA Polymerase (Promega,

Madison, WI), 10 µmol of each primer, and 300 ng bovine serum albumin (BSA; New England

BioLabs Inc. Ipswitch, MA), to which nuclease-free H_2O was added to reach a volume of 25 µl.

Thermocycler conditions (BioRad, Hercules, CA) were: 3 min at 95°C, 30 cycles of 30 sec at

95°C, 30 sec at 50°C, and 60 sec at 72°C, followed by a 5 min cycle at 72°C and then an

indefinite hold at 4°C. Gel electrophoresis confirmed the expected band size of 300–350 bp.

Samples that produced PCR products of the expected size and all negative controls were

prepared for community amplicon sequencing of the 16S rRNA gene V4 region using an

Illumina MiSeq at the University of Connecticut Microbial Analysis, Resources, and Services

facility. Approximately 30 ng of DNA from each sample was added to a 96-well plate containing

10 µmol each of Illumina-barcoded versions of primers 515F and 806R, 5 µl AccuPrime buffer

(Invitrogen, Carlsbad, CA), 50 mM $MgSO_4$ (Invitrogen, Carlsbad, CA), 300 ng BSA (New

England BioLabs Inc. Ipswitch, MA), a 1 µmol spike-in of both non-barcoded primers 515F and

806R, and 1 unit AccuPrime polymerase (Invitrogen, Carlsbad, CA), to which nuclease-free H_2O

was added to a volume of 50 µl. Reaction mixes were separated in a 384-well plate using an

epMotion 5075 liquid handling robot (Eppendorf, Hamburg, Germany) forming three replicates

(each with a volume of 16.7 µl). This plate was then transferred to a thermocycler (Eppendorf,

Hamburg, Germany), which used the following conditions: 2 min at 95°C, 30 cycles of 15 sec at

95°C, 60 sec at 55°C, and 60 sec at 68°C, a final extension for 5 min at 68°C and an indefinite

hold at 4°C. Post-PCR, triplicate reactions were re-pooled using the epMotion and DNA

concentrations were quantified using a QIAxcel Advanced capillary electrophoresis system

(Qiagen, Hilden, Germany). Samples with concentrations > 0.5 ng/µl were pooled using equal

DNA masses to create the final sequencing libraries. Libraries were then bead-cleaned using

Mag-Bind RXNPure plus beads (OMEGA, Norcross, Georgia) in a 1:0.8 ratio of sequencing

library to bead volume. Cleaned library pools were adjusted to a concentration of 1.1 ng/µl ± 0.1

ng/µl and their concentrations were confirmed using the Qubit dsDNA high-sensitivity assay on

a Qubit 3.0 fluorimeter (Invitrogen, Carlsbad, California). Microbial community sequencing on

an Illumina MiSeq using 2 x 250 bp libraries (Illumina, San Diego, California) was completed in

4 batches, each containing either the multi-state dataset (104 samples + 5 negative controls), the

two-colony dataset (147 samples + 9 negative controls), the pupae dataset (20 samples + 1

negative control), or the whole ant/gut dissection dataset (10 samples + 1 negative control).

Bioinformatic analyses:

We analyzed all datasets individually. Reads were analyzed using R v3.5.3 [39] and the

dada2 v1.11.1 [40] pipeline for amplicon sequence variants (ASVs)

(https://benjjneb.github.io/dada2/tutorial.html, accessed: November 11, 2017). Read counts per

sample were 5–1,952,121(multi-state dataset), 34–212,473 (two colony dataset), 15,495–92,408

(pupae dataset), and 17,565–132,790 (whole ant/gut dissection dataset; Suppl. File 1). Metadata

files were imported into phyloseq v1.26.1 [41], creating a phyloseq R object that was used for

subsequent analyses. Reads that were not classified as belonging to the kingdom Bacteria using

the SILVA database v128 [42, 43] were removed. The ASVs that matched to mitochondria were

then removed separately because SILVA included them in the kingdom Bacteria. Samples were

screened for contamination using the decontam v1.2.1 [44] prevalence protocol with a default

threshold value of 0.1. No reads were flagged as contaminants for any of the datasets, resulting

in 1,798, 1,353, 470, and 161 unique ASVs for the multi-state, two colony, pupae, and whole

ant/gut dissection datasets, respectively (Suppl. File 1). Negative control samples were not

considered further. All samples were rarefied to 20,000 reads for the multi-state dataset, 5,000

reads for the two-colony dataset, 10,000 reads for the pupae dataset, and 14,500 reads for the

whole ant/gut dissection dataset, and read counts were converted to relative abundances. The

final phyloseq objects contained 71, 110, 19, and 10 samples for the multi-state, two colony,

pupae, and whole ant/gut dissection datasets, respectively.

Alpha diversity was measured using the phyloseq 'plot_richness' command and

compared using the R stats 'aov' and 'TukeyHSD' commands. Beta diversity was measured

using weighted Unifrac (WUF) and unweighted Unifrac (UUF) distance metrics. Weighted and

unweighted Unifrac distances were calculated, ordinated, and viewed using the 'distance',

'ordinate', and 'plot_ordinate' phyloseq commands, respectively. Ellipses were added to the

PCoA plots of each Unifrac distance using the ggplot2 [45] command 'stat_ellipse' with a

default multivariate t-distribution. PERMANOVA tests were calculated using vegan v2.5-4 [46].

Although weighted Unifrac and unweighted Unifrac distances were used for each test, to keep

the text concise only one test is listed in the text and the complimentary values are presented in

Suppl. Tables S2-S4. After exporting the multi-state phyloseq object into Microsoft Excel, we

chose the first sample from each colony to be used in the Mantel tests, creating a new dataset

containing 36 samples. Each sample and their corresponding GPS locations were uploaded to

GeoMatrix [47] (Geographic Distance Matrix Generator v1.2.3) following the specified format.

The GeoMatrix output was downloaded and imported into R. Full Mantel tests were completed

using the GeoMatrix file and the WUF and UUF distance matrices for the 36 samples that

included only one ant from each colony using the vegan command 'mantel' . Partial Mantel tests

were completed and Mantel correlograms created using the vegan 'mantel.correlog' and 'plot'

commands, respectively. Kolmogorov-Smirnov and Wilcoxon rank sum tests were completed

using the R commands "ks.test" and "wilcox.test", respectively. If the p-value for the

Kolmogorov-Smirnov test was > 0.05, then the Wilcoxon rank sum tests were used to compare

the distributions of Unifrac distances; otherwise, a T-test was used (Suppl. Table S6). A heatmap

of the bacterial genera in the two-colony dataset whose read abundances were > 5% was created

using the 'heatmap.2' command in gplots v3.0.3 [48]. Heatmaps describing the median relative

abundances of genera with a read abundance > 5% within workers and male alates from colony

JKH000270 and JKH00307 were created using charticulator v2.0.4

(https://charticulator.com/app/index.html).

A phylogenetic tree was constructed to show relationships between the 16S rRNA

sequences of the Tenericutes generated during this study and those that had been previously

isolated from the leaf-cutting ants *Atta* and *Acromyrmex*. The most abundant *Spiroplasma* ASV

and *Mesoplasma* ASV from the combined multi-state and two-colony datasets were aligned with

reference Tenericute sequences generated from other ants and related *Mesoplasma* and

Spiroplasma sequences, all downloaded from NCBI, using MUSCLE v3.8.31 [49] and then

trimmed to the same alignment length. A phylogenetic tree, rooted using the *Bacillus subtilis* 16S

rRNA gene sequence, was calculated using a GTRGAMMAI substitution model and 500

bootstrap replicates in RaxML v8.2.11 [50].

The commands used for all analyses are attached as Suppl. File 2. All raw sequencing

reads are deposited in the NCBI database under BioProject PRJNA687229.

RESULTS

Ant microbiomes differ between states

We characterized *T. septentrionalis* worker ant microbiomes in field-collected ants from

New York, down the U.S. East Coast into Florida, and westward into Louisiana (Suppl. Fig. S1).

T. septentrionalis worker ant microbiomes were dominated by bacteria belonging to the phyla

Actinobacteria, Firmicutes, Proteobacteria, and Tenericutes (Fig. 1A; Suppl. Fig. S3). Alpha

diversity was low for these *T. septentrionalis* microbiomes and did not differ between states or

between ants sampled from the same colony (Suppl. Fig. S4; ANOVA: State p=0.492, Colony

p=0.335; Suppl. Table S1). Eighteen genera were present with an abundance of $\geq 5\%$ in at least

one sample (Fig. 1A & B). The most prevalent genera in the dataset, making up the core *T.*

septentrionalis worker ant microbiome, were *Thermomonas, Escherichia, Lautropia,*

Solirubrobacter, Pseudonocardia, Aeromicrobium, and *Nocardioides* (Fig. 1A). These bacteria

had a prevalence of at least 75% throughout the dataset and co-occurred frequently. *Phycicoccus,*

Mesoplasma, Spiroplasma, Acinetobacter, and *Wolbachia* were less prevalent in this dataset (50–

75%; Fig. 1A). The relative abundances of all genera varied greatly between samples (Fig. 1B).

Pseudonocardia, *Aeromicrobium*, *Nocardioides*, and *Solirubrobacter* each had high prevalence

but low median relative abundances $< 25\%$ in most samples (Fig. 1B). *Thermomonas* and

Lautropia had high median relative abundances that varied greatly between samples. When

present, *Mesoplasma* and *Spiroplasma* were often highly abundant ($> 50\%$ of all microbes in a

sample) and did not co-occur with one another in any sample. Many ant colonies sampled from

New York and North Carolina lacked *Mesoplasma* or *Spiroplasma* (Suppl. Fig. S3).

Amycolatopsis, *Bacillus*, and *Weissella* were also highly abundant in 1–2 samples (Fig. 1B).

Multiple ants were collected from most colonies in the multi-state dataset, and the colony

from which ants were collected explained more variation in microbiome composition than did

the state of collection (WUF Colony PERMANOVA: R^2= 0.524, p= 0.003; WUF State

PERMANOVA R^2= 0.135, p= 0.003; Suppl. Table S2). These trends remained after excluding

samples from New York and North Carolina, which lacked Tenericutes (Suppl. Table S3).

Although ant microbiomes from the multi-state dataset clustered weakly by state (Fig. 2; Suppl.

Fig. S5), weighted Unifrac distances between ant microbiomes did not correlate with the

geographic distances between collection locations, although unweighted Unifrac distances did,

albeit weakly (UUF Mantel R=0.143, p=0.019, Fig. 2B; WUF Mantel R= 0.034, p=0.778, Suppl.

Fig. S6; Suppl. Table S4). In the corresponding partial Mantel tests, distances between colonies

from the smallest distance class (< 100km) were positively and significantly correlated to UUF

distances between the ant microbiomes from these colonies (Fig. 2B; Suppl. Table S4). Distance

from classes 2 (~250 km) and 5 (~850 km) positively and significantly correlated with UUF

distances in ant microbiome composition, but all other correlations were non-significant (Fig.

2B; Suppl. Fig. 6; Suppl. Table S4). Overall, these analyses show that microbiomes from

colonies collected within 100 km from each other were somewhat similar to each other but that

microbiomes otherwise varied in composition between colonies.

Ant microbiomes differ between castes and due to lab adaptations

In our multi-state dataset, we found that ant microbiomes were more conserved within a

colony than between colonies. However, the multi-state dataset only included 1–3 field-collected

worker ants to represent each colony. To overcome this limitation, we more deeply sampled

individual ants from two colonies collected in Wharton State Forest, New Jersey (Suppl. File 1).

The field-collected workers in this two-colony dataset had low alpha diversities, similar to those

in the multi-state dataset (Suppl. Fig S7). We also found that the same phyla were conserved in

the two-colony dataset as in the multi-state dataset. Field-collected worker ants were colonized

by Proteobacteria, Tenericutes, and Actinobacteria, with some ants from colony JKH000270 also

colonized with Bacteroidetes (Fig. 3A). Most core worker ant microbiome genera from the

multi-state dataset (*Thermomonas, Lautropia, Solirubrobacter, Pseudonocardia,*

Aeromicrobium, and *Nocardioides*) were also present in field-collected worker ants in the two-

colony dataset (Suppl. Fig. S8).

Although the limited number of colonies in the two-colony dataset is not ideally powered

to compare ant microbiomes from different colonies in a general way (in contrast to the multi-

state dataset), it instead prioritizes sampling of ants from different developmental stages and

castes (pupae, larvae, male alates, and female alates) and ants collected before and after lab

adaptation (field-collected or lab-maintained). Alpha diversity of lab-maintained male alates was

often significantly lower compared to that of other sample types, which otherwise differed

minimally from each other (Suppl. Table S5; Suppl. Fig. S7). Like the workers, all castes were

colonized by Actinobacteria, Proteobacteria, and Tenericutes, with Bacteroidetes only abundant

in colony JKH000270, along with Firmicutes in some male alates (Fig. 3A). Compared to the

worker microbiomes, Tenericutes were particularly abundant in male alate microbiomes,

Proteobacteria in pupae and larvae, and Actinobacteria and Proteobacteria in female alates (Fig.

3A). There were extremely few *Mesoplasma* or *Spiroplasma* in pupae and larvae from the two-

colony dataset. However, *Mesoplasma* and *Spiroplasma* were abundant in the separate pupae

dataset generated from other *T. septentrionalis* colonies and used for confirmation (Suppl. Fig.

S9).

In the two-colony dataset, ant microbiome composition differed between all tested

categories, and although caste explains the most variation it depends on the colony and lab

adaptation (Fig. 3B & C). Caste accounted for the most microbiome variation using both WUF

and UUF distances (Fig. 3B). These differences were colony-, caste-, and adaptation-specific

because all interaction terms accounted for substantial microbiome variation using both WUF

and UUF distances. Microbiomes from colonies JKH000270 and JKH000307 clustered

separately along axis 2 in the WUF and UUF PCoAs but overlap in the center (Fig. 3C; Suppl.

Fig. S10). This second axis accounts for a smaller amount of microbiome variation than axis 1,

which separates the microbiomes into different castes, lab adaptation status, and combinations of

these categories, in both the WUF and UUF PCoAs.

Although each caste had a unique microbiome, we could only directly compare changes

resulting from lab adaptation for workers and male alates, which were the only castes that we

sampled from both field and lab settings (Fig. 3D). In colony JKH000270, field-collected and

lab-maintained worker and male alate microbiomes differed principally in their abundances of

Spiroplasma (Fig. 3D). In colony JKH000307, field-collected and lab-maintained worker

microbiomes differed in their abundances of *Thermomonas, Lautropia*, *Aminobacter*, and

Cupriavidus, and lab-maintained, but not field-collected, male alate microbiomes were

dominated by *Mesoplasma* (Fig. 3D). Lab-maintained microbiomes were much less variable than

the field-collected microbiomes, except for colony JKH000307 worker ants, when measured

using UUF distances (Suppl. Fig. S11; Suppl. Table S6).

In nearly all instances, microbiomes varied more between each category (colonies, caste,

and lab adaption) than within the same category (Suppl. Fig. S12). Microbiomes from each

colony were grouped by caste, and then worker and male alate samples were further grouped into

lab-maintained and field-collected categories to remove lab adaptation as a potentially

confounding variable (Suppl. Fig. S12). Typically, median WUF and UUF distances were greater

between categories (caste, colony, lab adaptation) than within them. However, this was not true

for UUF JKH000270 field-collected male alates, JKH000270 female alates, and JKH000307

larvae (Suppl. Fig. S12). Overall, these comparisons highlight that ant microbiomes are more

similar within our sampling categories (colony, caste, lab adaptation) than between them.

Spiroplasma and *Mesoplasma* are *T. septentrionalis* symbionts

Mesoplasma and *Spiroplasma* were the two dominant genera in our microbiome samples,

accounting for 12% and 14% of all the reads in our dataset, respectively. These bacteria,

classified within the phylum Tenericutes, are well-known symbionts of insects and plants [51–

56]. Although their function is unknown, *Spiroplasma* and *Mesoplasma* have been found in other

fungus-growing ants [26–30, 33]. The most abundant *Mesoplasma* ASV in our dataset,

comprising 97.7% of all *Mesoplasma* reads, was highly similar to the 16S rRNA of EntAcro1

and an uncultured Entomoplasmataceae 16S rRNA sequence found in army ants (Fig. 4) [57].

EntAcro1 is a metagenome-assembled genome (MAG) that was assembled from *Acromyrmex*

fungus-growing ants [30], and related 16S rRNA sequences have also been found in *Atta*

colombica and At. sexdens, *Mycetomoellerius zeteki* (previously *Trachymyrmex zeteki*), and

Cyphomyrmex rimosus fungus-growing ants. The most abundant *Spiroplasma* ASV in our

dataset, comprising 97.3% of all *Spiroplasma* reads, was highly similar to the EntAcro10 MAG

assembled from *Acromyrmex* ants [30] and for which related 16S rRNA sequences were found in

Atta, Acromyrmex, and *Mycetomoellerius* [27, 29] fungus-growing ants, along with other 16S

rRNA sequences that were annotated as uncultured Entomoplasmatales from two other fungus-

growing ants, *Sericomyrmex amabilis* and *M. jamaicensis* (Fig. 4) [26, 58].

The presence of *Mesoplasma* and *Spiroplasma* varied between different ant colonies,

between ants from the same colony, and between ants from different castes, both before and after

lab adaptation. In our multi-state dataset, colonies from New York and North Carolina had lower

Tenericute abundances compared to the other states (Fig. 5A). *Spiroplasma* was abundant in

field-collected worker ants and both field-collected and lab-maintained male alates from colony

JKH000270. *Mesoplasma* was abundant in field-collected and lab-maintained workers and lab-

maintained male alates from colony JKH000307. In both the multi-state and two-colony datasets,

only one of *Spiroplasma* or *Mesoplasma* was present per colony, although not all ants from that

colony were colonized (Fig. 5B & C). An exception to this trend was in Colony JKH000307, in

which *Mesoplasma* was abundant but *Spiroplasma* colonized two lab-maintained male alates and

two field-collected workers. Ants colonized by *Spiroplasma* were rarely found within a colony

mainly colonized by *Mesoplasma*, and vice versa (Fig. 5B & C), and no ants in our multi-state or

two-colony datasets were colonized by both *Spiroplasma* and *Mesoplasma*. However, in a

confirmatory dataset that sample pupae from two additional ant colonies JKH000125 and

JKH000136, and we found that pupae from colony JKH000136 were colonized exclusively by

large amounts of *Mesoplasma* and that pupae from colony JKH000125 pupae were colonized

primarily by *Spiroplasma* but also by small amounts of *Mesoplasma* (Suppl. Fig. S13). This is

the only time we have observed ants colonized by both *Mesoplasma* and *Spiroplasma*.

DISCUSSION

Our microbiome sequencing of individual *T. septentrionalis* ants shows that caste, lab

adaptation (field-collected or lab-maintained), and the colony that an ant belongs to all influence

the structure of the ant microbiome. Our multi-state dataset includes field-collected workers ants

from 6 states representing 36 colonies, allowing us to describe a core microbiome of wide-

ranging worker ants that contains *Thermomonas, Escherichia, Lautropia, Solirubrobacter,*

Pseudonocardia, Aeromicrobium, and *Nocardioides* (Fig. 1A), all of which were prevalent in >

75% of the ants in this dataset. This core microbiome was largely confirmed in the field-

collected worker ant microbiome in our two-colony dataset. *Wolbachia* was also found in several

samples from our multi-state dataset, similar to what has been documented in *Acromyrmex*,

where it is thought to be transmitted maternally [28]. *Mesoplasma* and *Spiroplasma* were both

found throughout our dataset (Fig. 5) and are also abundant in other Attine ant genera [27, 28].

Our results also agree with those of Ishak et al. [24], who identified 19 bacterial genera that were

common in several *T. septentrionalis* colonies sampled from a single location in Texas. Of these

19 genera, 9 are among the most abundant 18 genera in our multi-state dataset: *Aeromicrobium*,

Nocardioides, *Phycicoccus*, *Pseudonocardia*, *Solirubrobacter*, *Bacillus*, *Pseudomonas*,

Mesoplasma, and *Spiroplasma*. The larger number of taxa found by Ishak et al. [33] is consistent

with our finding that the *T. septentrionalis* ant microbiome has some biogeographic

conservation, albeit limited. Colonies that are located closer together had the most similar

microbiomes, meaning that a list of common taxa sampled from a single collection location

would be larger than a core microbiome defined using samples collected from many different

locations.

In our multi-state dataset, in which 1-3 ants were sampled per colony, the microbiomes from

ants sampled from the same colony were more similar than ants sampled from different colonies.

To further study the variation between the microbiomes of individual ants from a single colony,

our two-colony dataset included microbiomes from ants more deeply sampled from two colonies,

including ants from different castes and from both before and after lab adaptation. Again, we

found that ant microbiomes within colonies were more similar than those between colonies. This

bolsters a common natural pattern, e.g., where nestmates of birds [59] and mammals [60], and

colonies of insects such as honey bees [9, 11] and ants [61] all have similar microbiomes. These

differences that we observed between colonies could be due to local ecological factors such as

regional differences in the plants, insect frass, and the microbes associated with these substrates

that will cause the ant microbiome to vary. This is supported by our samples collected following

adaptation of colonies to our lab, where environmental conditions are constant and colonies are

fed only sterile cornmeal. Lab-maintained ant microbiomes are much less variable than those

sampled in the field (Suppl. Table S6), highlighting the importance of natural foraging and

environmental variation for maintaining microbiome diversity. Another possible reason why

microbiomes might differ between colonies is the strong founder effects that likely occur during

colony foundation. Before leaving to mate, a female alate will place a piece of the fungus garden

in her infra-buccal pocket [13]. After mating, this new queen will found her own colony using

that piece of fungus garden, tending it and giving birth to all offspring in that colony. The

microbiome possessed by the queen will be passed to these workers who will thereafter pass it on

to subsequent generations of workers, thus creating a bottleneck in microbiome composition

caused by the limited microbial diversity possessed by the queen during colony foundation.

Within a colony, microbiome composition depended on ant castes (Fig. 3), which determined

the greatest amount of variation between microbiome composition in the two-colony dataset

(Fig. 3B). Gut microbiomes often differ between social insect castes, likely due to the specialized

functional role played by each caste [1]. Reproductive castes often differ from the worker castes

in termites [8], honey bees [9], and other species of ants [61]. In *T. septentrionalis* colonies,

workers forage for plant material and caterpillar frass that have their own microbiomes, and

these bacteria might be transferred to the workers and to the fungus garden that they tend. Such

microbes brought into the colony via foraging could then be seen transferred to all ant

microbiomes in that colony. However, such transfer may be rare due to the ant the ants grooming

themselves, other ants, and the fungus garden to remove pathogens [62]. This likely results in the

removal of most microbes brought in by foraging from both the ants and their fungus garden. In

contrast, the role of male and female alates is to mate and, for females, to start a new colony. The

different microbiomes possessed by these castes could therefore reflect their need to reproduce,

which the workers do not experience. Female alates possess large amounts of lipids, proteins,

and carbohydrates to provide energy when they start their colonies [63], and male alates must

travel further than female alates to mate [64]. The different microbiome composition possessed

by male and female alates (Fig. 3C) could therefore reflect the differing energy and nutrient

requirements of their different reproductive behaviors [63, 64].

The presence and abundance of *Mesoplasma* and *Spiroplasma* differ between each colony

and individual ant, explaining some of the differences in ant microbiome composition between

ant colonies and castes. The most abundant *Mesoplasma* and *Spiroplasma* 16S rRNA genes

detected in our dataset are very similar to those of other Tenericutes found in other fungus-

growing genera, which suggests that these bacteria are a common member of the fungus-growing

ant symbiosis. The functional role played by these Tenericutes is unknown. They are unlikely to

be reproductive pathogens, based on their high abundances in worker ants, which are

reproductive dead ends. *Acromyrmex* and *Atta* callows (i.e., newly emerged worker ants) were

readily colonized by Tenericutes when cared for by other Tenericutes-containing worker ants,

but not after 21 days in the absence of such ants [28]. Thus, Tenericutes symbionts are likely

transferred between ants via care from other workers. FISH imaging has also been used to

localize Tenericutes to ant guts but not in reproductive organs. However, Tenericutes are

abundant in male alates, from which they might be transmitted to a female foundress during

mating. Because we only have characterized female alate microbiomes from a single colony,

future studies should include a wider sampling of this caste. Tenericutes might also be acquired

from an environmental source (e.g., foraged plants or caterpillar frass), or during interactions

with other ant colonies [65].

Tenericute symbionts of fungus-growing ants may be nutritional symbionts. Metagenomic

reconstructions of *Spiroplasma* and *Mesoplasma* symbionts from *Acromyrmex* ants suggested

that these bacteria may decompose excess arginine, and for *Mesoplasma*, citrate [30]. Arginine

recycling would produce ammonium that could be provided to the fungus garden via ant fecal

droplets, and the degradation of citrate, acquired from foraged fruit, leaves, and other plant

material, could produce acetate for use by the ants. These hypotheses suggest that Tenericutes

might provision nutrients for both the ants and the cultivar fungus. However, in our data

Tenericute presence varied extensively between ant colonies and castes (Fig. 5), meaning that

these symbionts must not be absolutely required for ant survival, especially given their complete

absence in some ants. Instead, such extreme variation in abundance between colonies and

individual ants could imply that *Mesoplasma* or *Spiroplasma* are ant pathogens or commensals,

at least in some ecological contexts. Interestingly, we did not find any co-infections of both

Mesoplasma and *Spiroplasma* together in the same ant, except for in pupae from a single ant

colony (Suppl. Fig. 13). This might suggest a transmission bottleneck during microbiome

assembly that allows for colonization of only one of these symbionts, or competition between

Spiroplasma and *Mesoplasma* in these ants.

Our study shows how microbiome assembly differs between ant colonies, castes and as a

result of lab adaptation. *T. septentrionalis* ant microbiomes did not vary extensively between

sampling locations but did vary substantially between ant colonies and castes. Lab adaptation

showed the importance of ant foraging or environmental variation for maintaining microbiome

diversity, with lab-maintained microbiomes being less variable than field-collected ones as a

reflection of their more stable environment. The presence of *Mesoplasma* and *Spiroplasma* cause

some of these differences between colonies, castes, and as the result of lab-adaptation. Together,

these data show how the composition of a social insect microbiome is the result of colony- and

caste-specific factors.

Acknowledgments

480   We would like to thank the state forest and park staff for their assistance with our fieldwork, the
481   members of the Klassen Lab for their help collecting ant colonies and for their thoughtful
482   feedback on this manuscript, and the UConn Microbial Analysis, Resources, and Services
483   facility for microbiome sequencing. This research was funded by the U.S. National Science
484   Foundation (IOS-1656475) and the University of Connecticut.

485
Figure 1) The *T. septentrionalis* worker ant core microbiome. A) Prevalence of bacterial genera
in the multi-state dataset with an abundance ≥ 5% in at least one ant microbiome. Genera are
ordered on the y-axis from most to least prevalent. Median relative abundance is represented by
circle size and phylum is represented by color. B) Relative abundance of the bacterial genera ≥
5% in at least one ant sample in the multi-state dataset, with the abundance in each sample
represented by a single point for each genus. Bars indicate medians.

Figure 2) Biogeography of the *T. septentrionalis* worker microbiome. A) PCoA of Weighted
Unifrac distances between ant microbiomes in the multi-state dataset. Samples are colored by
state and grouped using a multivariate t-distribution. n=71. B) Partial Mantel correlation between
Unweighted Unifrac distances and ant sampling locations in the multi-state dataset, excluding
comparisons between ants sampled from the same colony. The x-axis indicates the distance class
index in kilometers, and the y-axis indicates the Mantel correlation R statistic. Filled and unfilled
squares indicate significant (p < 0.05) and nonsignificant (p > 0.05) p values, respectively, and
points above and below the red line indicate positive and negative correlations, respectively.

Figure 3) Colony, caste, and lab-adaptation all determine ant microbiome composition. A) Bar
plot of the most abundant phyla in microbiomes from colonies JKH000270 (top) and
JKH000307 (bottom). Single bars represent individual ant microbiomes, and sample types are
grouped along the x-axis. The y-axis indicates relative abundance. Phyla are differentiated by
color, and "other" represents phyla present at < 15% relative abundance. B) PERMANOVA
analyses testing correlations between ant microbiome composition in the two-colony dataset and
ant caste, lab adaptation status, and their colony of origin. n= 110. C) PCoA of Weighted Unifrac

distances between ant microbiomes in the two-colony dataset. Colors indicate ant caste, shapes
indicate colony ID, and solid and open shapes indicate field-collected and lab-maintained
samples, respectively. D) Heatmap of genera with an abundance ≥ 5% in worker ants and male
alates from colony JKH000270 (left) and colony JKH000307 (right). Genera are listed on the y-
axis and colors indicate their phylum.

Figure 4) Maximum-likelihood phylogenetic tree of the dominant Tenericute 16 rRNA gene
ASVs from the combined multi-state and two-colony datasets (in bold text) and reference 16S
rRNA gene sequences sequenced from other fungus-growing ants and related Tenericutes strains.
The tree was rooted using *Bacillus subtilis* and *Mycoplasma hominis* and bootstrap percentages >
60 are shown.

522
Figure 5) *Spiroplasma* (yellow) and *Mesoplasma* (orange) are abundant in *T. septentrionalis*
ants. A) Single bars represent the relative abundance of reads from the genus *Spiroplasma* or
Mesoplasma in ants from the multi-state dataset, grouped by the state of collection. X-axis
brackets group worker ants from the same colony. n=71. B & C) The relative abundance of the
genus *Spiroplasma* or *Mesoplasma* in the two-colony dataset, with each dot representing relative
abundance in a single ant. All ants in this dataset were colonized exclusively by either
Spiroplasma or *Mesoplasma*.
530

Works Cited

1.   The Editors of Encyclopedia Britannica (2018) Social Insect. Encyclopædia Britannica Inc., Chicago, Illinois, USA

2.   Sinotte VM, Renelies-Hamilton J, Taylor BA, et al (2020) Synergies between division of labor and gut microbiomes of social insects. Front Ecol Evol 7:e00503. https://doi.org/10.3389/fevo.2019.00503

3.   Hongoh Y, Ekpornprasit L, Inoue T, et al (2005) Intracolony variation of bacterial gut microbiota among castes and ages in the fungus-growing termite *Macrotermes gilvus*. Mol Ecol 15:505–516. https://doi.org/doi: 10.1111/j.1365-294X.2005.02795.x.

4.   Schnorr SL, Hofman CA, Netshifhefhe SR, et al (2019) Taxonomic features and comparisons of the gut microbiome from two edible fungus-farming termites (*Macrotermes falciger; M. natalensis*) harvested in the Vhembe district of Limpopo, South Africa. BMC Microbiol 19:e164. https://doi.org/https://doi.org/10.1186/s12866-019-1540-5

5.   Dietrich C, Kohler T, Brune A (2014) The cockroach origin of the termite gut microbiota: patterns in bacterial community structure reflect major evolutionary events. Appl Environ Microbiol 80:2261–2269. https://doi.org/doi: 10.1128/AEM.04206-13

6.   Otani S, Mikaelyan A, Nobre T, et al (2014) Identifying the core microbial community in the gut of fungus-growing termites. Mol Ecol 23:4631–4644. https://doi.org/doi: 10.1111/mec.12874

7.   Otani S, Hansen LH, Sorensen SJ, Poulsen M (2016) Bacterial communities in termite fungus combs are comprised of consistent gut deposits and contributions from the environment. Microb Ecol 71:207–220. https://doi.org/doi: 10.1007/s00248-015-0692-6

8.   Otani S, Zhukova M, Koné NA, et al (2019) Gut microbial compositions mirror caste-specific diets in a major lineage of social insects. Environ Microbiol Rep 11:196–205

9.   Kapheim KM, Rao VD, Yeoman CJ, et al (2015) Caste-specific differences in hindgut microbial communities of honey bees (*Apis mellifera*). PLoS One 10:e0123911. https://doi.org/10.1371/journal.pone.0123911

10. Kwong WK, Moran NA (2016) Gut microbial communities of social bees. Nat Rev Microbiol 14:374–384. https://doi.org/10.1038/nrmicro.2016.43

11. Moran NA, Hansen AK, Powell JE, Sabree ZL (2012) Distinctive gut microbiota of honey bees assessed using deep sampling from individual worker bees. PLoS One 7:e36393. https://doi.org/10.1371/journal.pone.0036393

12. Wilson EO (1971) The Insect Societies. Harvard University Press, Cambridge, Massachusetts, USA

13. Mueller UG, Schultz TR, Currie CR, et al (2001) The origin of the attine ant-fungus mutualism. Q Rev Biol 76:169–197. https://doi.org/10.1086/393867

14. Schultz TR, Brady SG (2008) Major evolutionary transitions in ant agriculture. PNAS
105:5435–5440. https://doi.org/10.1073/pnas.0711024105

15. Weber NA (1972) The fungus-culturing behavior of ants. Am Zool 12:577–587.
https://doi.org/10.1093/icb/12.3.577

16. Hölldobler B, Wilson EO (1990) The Ants. Harvard University Press, Cambridge,
Massachusetts, USA

17. Solomon SE, Rabeling C, Sosa-Calvo J, et al (2019) The molecular phylogenetics of
Trachymyrmex Forel ants and their fungal cultivars provide insights into the origin and
coevolutionary history of 'higher-attine' ant agriculture. Syst Entomol 44:939–956.
https://doi.org/10.1111/syen.12370

18. Wheeler WM (1907) The fungus-growing ants of North America. Am Museum Nat Hist
23:669–807

19. Munkacsi AB, Pan JJ, Villesen P, et al (2004) Convergent coevolution in the
domestication of coral mushrooms by fungus–growing ants. Proc R Soc B Biol Sci
271:1777–1782. https://doi.org/https://doi.org/10.1098/rspb.2004.2759

20. Weber NA (1966) Fungus-growing ants. Science 153:587–604.
https://doi.org/10.4319/lo.2013.58.2.0489

21. De Fine Licht HH, Boomsma JJ (2010) Forage collection, substrate preparation, and diet
composition in fungus-growing ants. Ecol Entomol 35:259–269.
https://doi.org/10.1111/j.1365-2311.2010.01193.x

22. Currie CR, Scott JA, Summerbell RC, Malloch D (1999) Fungus-growing ants use
antibiotic-producing bacteria to control garden parasites. Nature 398:701–705.
https://doi.org/https://doi.org/10.1038/19519

23. Goldstein SL, Klassen JL (2020) *Pseudonocardia* symbionts of fungus-growing ants and
the evolution of defensive secondary metabolism. Front Microbiol 11:e621041.
https://doi.org/10.3389/fmicb.2020.621041

24. Klassen JL (2014) Microbial secondary metabolites and their impacts on insect symbioses.
Curr Opin Insect Sci 4:15–22. https://doi.org/10.1016/j.cois.2014.08.004

25. Van Borm S, Billen J, Boomsma JJ (2002) The diversity of microorganisms associated
with *Acromyrmex* leafcutter ants. BMC Evol Biol 2:9.
https://doi.org/https://doi.org/10.1186/1471-2148-2-9

26. Liberti J, Sapountzis P, Hansen LH, et al (2015) Bacterial symbiont sharing in
Megalomyrmex social parasites and their fungus-growing ant hosts. Mol Ecol 24:3151–
3169. https://doi.org/10.1111/mec.13216

27. Sapountzis P, Zhukova M, Hansen LH, et al (2015) *Acromyrmex* leaf-cutting ants have
simple gut microbiota with nitrogen-fixing potential. Appl Environ Microbiol 81:5527–
5537. https://doi.org/10.1128/AEM.00961-15

28. Zhukova M, Sapountzis P, Schiøtt M, Boomsma JJ (2017) Diversity and transmission of

gut bacteria in *Atta* and *Acromyrmex* leaf-cutting ants during development. Front
Microbiol 8:e01942. https://doi.org/10.3389/fmicb.2017.01942

29.   Sapountzis P, Nash DR, Schiøtt M, Boomsma JJ (2018) The evolution of abdominal
microbiomes in fungus-growing ants. Mol Ecol 28:e14931.
https://doi.org/10.1111/mec.14931

30.   Sapountzis P, Zhukova M, Shik JZ, et al (2018) Reconstructing the functions of
endosymbiotic mollicutes in fungus-growing ants. eLife 7:e39209.
https://doi.org/10.7554/eLife.39209

31.   Rabeling C, Cover SP, Johnson RA, Mueller UG (2007) A review of the North American
species of the fungus-gardening ant genus *Trachymyrmex* (Hymenoptera: Formicidae).
Zootaxa 1664:e180014

32.   Seal JN, Brown L, Ontiveros C, et al (2015) Gone to Texas: phylogeography of two
Trachymyrmex (Hymenoptera: Formicidae) species along the southeastern coastal plain of
North America. Biol J Linn Soc 114:689–698. https://doi.org/10.1111/bij.12426

33.   Ishak HD, Miller JL, Sen R, et al (2011) Microbiomes of ant castes implicate new
microbial roles in the fungus-growing ant *Trachymyrmex septentrionalis*. Sci Rep
1:e00204. https://doi.org/10.1038/srep00204

34.   Lee KM, Adams M, Klassen JL (2019) Evaluation of DESS as a storage medium for
microbial community analysis. PeerJ 7:e6414. https://doi.org/10.7717/peerj.6414

35.   Sosa-Calvo J, Jesovnik A, Okonski E, Schultz TR (2015) Locating, collecting, and
maintaining colonies of fungus-farming ants (Hymenoptera: Formicidae: Myrmicinae:
Attini). Sociobiology 62:300–320. https://doi.org/10.13102/sociobiology.v62i2.300-320

36.   Sanders JG, Powell S, Kronauer DJC, et al (2014) Stability and phylogenetic correlation
in gut microbiota: Lessons from ants and apes. Mol Ecol 23:1268–1283.
https://doi.org/10.1111/mec.12611

37.   Green EA, Smedley SR, Klassen JL (2021) North American fireflies host low bacterial
diversity. Microb Ecol 82:793–804. https://doi.org/10.1007/s00248-021-01718-7

38.   Caporaso JG, Lauber CL, Walters WA, et al (2011) Global patterns of 16S rRNA
diversity at a depth of millions of sequences per sample. Proc Natl Acad Sci U S A
108:4516–4522. https://doi.org/10.1073/pnas.1000080107

39.   R Core Team (2018) R: A Language and Environment for Statistical Computing

40.   Callahan BJ, McMurdie PJ, Rosen MJ, et al (2016) DADA2: High-resolution sample
inference from Illumina amplicon data. Nat Methods 13:581–583.
https://doi.org/10.1038/nmeth.3869

41.   McMurdie PJ, Holmes S (2013) Phyloseq: An R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One 8:e61217.
https://doi.org/10.1371/journal.pone.0061217

42.   Quast C, Pruesse E, Yilmaz P, et al (2013) The SILVA ribosomal RNA gene database

project: Improved data processing and web-based tools. Nucleic Acids Res 41:590–596.
https://doi.org/10.1093/nar/gks1219

43. Yilmaz P, Parfrey LW, Yarza P, et al (2014) The SILVA and "all-species Living Tree
Project (LTP)" taxonomic frameworks. Nucleic Acids Res 42:643–648.
https://doi.org/10.1093/nar/gkt1209

44. Davis NM, Proctor DM, Holmes SP, et al (2018) Simple statistical identification and
removal of contaminant sequences in marker-gene and metagenomics data. Microbiome
6:226. https://doi.org/10.1186/s40168-018-0605-2

45. Wickham H (2016) ggplot2: Elegant graphics for data analysis. Springer-Verlag New
York

46. Oksanen J, Blanchet FG, Friendly M, et al (2019) vegan: Community Ecology Package. R
package version 2.5-4

47. Ersts PJ (2021) Geographic Distance Matrix Generator (version 1.2.3). American Natural
History, Center for Biodiversity and Conservation.
http://biodiversityinformatics.amnh.org/open_source/gdmg

48. Warnes GR, Bolker B, Bonebakker L, et al (2009) gplots: Various R programming tools
for plotting data

49. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32:1792–1797. https://doi.org/10.1093/nar/gkh340

50. Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. Bioinformatics 30:1312–1313.
https://doi.org/10.1093/bioinformatics/btu033

51. Hackett KJ, Whitcomb RF, Tully JG, et al (1992) Lampyridae (Coleoptera): A plethora of
Mollicute associations. Microb Ecol 23:181–193.
https://doi.org/https://doi.org/10.1007/BF00172639

52. Ammar E-D, Hogenhout SA (2006) Mollicutes associated with arthropods and plants. In:
Bourtzis K, Miller TA (eds) Insect Symbiosis. CRC Press, Boca Roton, FL, pp 97–118

53. Whitman WB, Inc. JW& S (2015) Bergey's manual of systematics of archaea and
bacteria. Bergey's Manual Trust, Hoboken, New Jersey, USA

54. Williamson DL, Adams JR, Whitcomb RF, et al (1997) *Spiroplasma platyhelix* sp. nov., a
new mollicute with unusual morphology and genome size from the dragonfly *Pachydiplax*
longipennis. Int J Syst Bacteriol 47:763–766

55. Rose DL, Kocka JP, Somerson NL, et al (1990) *Mycoplasma lactucae* sp. nov., a sterol-
requiring Mollicute from a plant surface. Int J Syst Bacteriol 40:138–142.
https://doi.org/doi: 10.1099/00207713-40-2-138.

56. Tully JG, Whitcomb RF, Hackett KJ, et al (1994) Taxonomic descriptions of eight new
non-sterol-requiring mollicutes assigned to the genus *Mesoplasma*. Int J Syst Bacteriol
44:685–693. https://doi.org/10.1099/00207713-44-4-685

57. Funaro CF, Kronauer DJC, Moreau CS, et al (2011) Army ants harbor a host-specific
clade of *Entomoplasmatales* bacteria. Appl Environ Microbiol 77:346–350.
https://doi.org/10.1128/AEM.01896-10

58. Russell JA, Moreau CS, Goldman-Huertas B, et al (2009) Bacterial gut symbionts are
tightly linked with the evolution of herbivory in ants. Proc Natl Acad Sci 106:21236–
21241. https://doi.org/10.1073/pnas.0907926106

59. White J, Mirleau P, Danchin E, et al (2010) Sexually transmitted bacteria affect female
cloacal assemblages in a wild bird. Ecol Lett 13:1515–1524. https://doi.org/doi:
10.1111/j.1461-0248.2010.0154

60. Leclaire S, Nielsen JF, Drea CM (2014) Bacterial communities in meerkat anal scent
secretions vary with host sex, age, and group membership. Behav Ecol 25:996–1004.
https://doi.org/https://doi.org/10.1093/beheco/aru074

61. Segers FHID, Kaltenpoth M, Foitzik S (2019) Abdominal microbial communities in ants
depend on colony membership rather than caste and are linked to colony productivity.
Ecol Evol 9:13450–13467. https://doi.org/10.1002/ece3.5801

62. Currie C, Stuart A (2001) Weeding and grooming of pathogens in agriculture by ants.
Proc R Soc B Biol Sci 26:1033–1039

63. Seal JN, Tschinkel WR (2007) Energetics of newly-mated queens and colony founding in
the fungus-gardening ants *Cyphomyrmex rimosus* and *Trachymyrmex septentrionalis*
(Hymenoptera: Formicidae). Physiol Entomol 32:8–15. https://doi.org/10.1111/j.1365-
3032.2006.00534.x

64. Matthews AE, Kellner K, Seal JN (2020) Male-biased dispersal in a fungus-gardening ant
symbiosis. Ecol Evol 11:2307–2320. https://doi.org/10.1002/ece3.7198

65. Howe J, Schiøtt M, Boomsma JJ (2019) Horizontal partner exchange does not preclude
stable mutualism in fungus-growing ants. Behav Ecol 30:372–382.
https://doi.org/https://doi.org/10.1093/beheco/ary176