Genome-wide analysis of DNA methylation reveals selection signatures of the grass carp during domestication

Yichao Li[a,1], Bing Fu[a,1], Junming Zhang[a,1], Jun Xie[a], Guangjun Wang[a], Peng Jiang[a], Jingjing Tian[a], Hongyang Li[a], Yun Xia[a], Zhifei Li[a], Ermeng Yu[a,*]

[a] *Pearl River Fisheries Research Institute, Chinese Academy of Fishery Sciences, Guangzhou 510380, China*

* Corresponding author. *E-mail* address: yem@prfri.ac.cn (E. Yu).

[1] These authors contributed equally to this work.

ABSTRACT

With the rapid development of aquaculture, many fish species are domesticated and brought into

cultivation. In the process of domestication, the domesticated fish undergone intense selection

pressures and develop some adaptations and phenotypic traits, namely selection signatures, such

as growth and metabolism, immunity, foraging and learning behaviors. However, how this

selection signatures emerges is still not clear and the knowledge of molecular epigenetic

mechanisms underlying selection signatures in fish is still in its infancy. Thus, we used a farmed

fish, grass carp (*Ctenopharyngodon idellus*), as model species to detect these selection signatures

and identify the candidate differentially methylated genes that are closely associated with these

selection signatures at the level of whole genome, investigating the role of DNA methylation in

the emergence of selection signatures during domestication. Our results showed that domesticated

grass carp demonstrated four selection signatures, including growth and metabolism, immunity,

foraging and learning behaviors, and 38 candidate genes were found associated with these traits.

16 genes are significant candidate genes which play major roles in the growth and metabolism,

such as IGF-1 , GK , GYS1, etc. 11 genes are related to immunity, including . The GRM1,

TAS1R1 and TAS1R3 genes are essential for the adaptation of domesticated grass carp to

commercial feed in artificial rearing condition. The C-FOS, POMC and CBP genes may be

responsible for the acquisition of novel feeding habits and contribute to faster growth indirectly by

enhancing food intake. The findings here in will provide new insights to expand our understanding

about the role of epigenetic modifications in shaping physiological phenotypes in this and other

teleost models, which can contribute to efficient breeding of aquaculture stocks and restocking

programmes.

Keywords:

DNA methylation

grass carp

selection signatures

1. Introduction

As an economically important food source of humans, fish is one of the most important domesticated species (Teletchea ,et al. 2012). In modern agricultural industry, fish have undergone strong long-term artificial selection and developed a range of adaptations and phenotypic traits, namely selection signatures (Milla et al.,2021), such as higher flexibility in diet, rapid growth, less stress susceptibility, a more socially tolerant disposition and enhanced prolificacy (Pasquet et al., 2018), which may distinguish those domesticated breeds from their wild counterparts. These phenotypic differences make fish highly suitable for animal agriculture and comparative studies.

Epigenetic changes is increasingly recognized to contribute to the emergence of phenotypic differences (Goldberg et al., 2007). DNA methylation, the most widely studied epigenetic regulatory mechanisms, can respond to environmental change and, at the same time, be stable enough to be maintained throughout lifetime, even across generations (Anastasiadi et al., 2019). DNA methylation has been associated with many biological process, including gene expression regulation, genomic imprinting, X chromosome inactivation, embryonic development, the alteration of chromatin structure and transposon inactivation, etc (Wilson et al., 2012; Jones et al., 2012). Multiple approaches have been developed to analyze DNA methylation profiles at the genome-wide level, including comprehensive high-throughput arrays for relative methylation (CHARM), methylated DNA immunoprecipitation (MeDIP), methylation array, reduced-representation bisulfite sequencing (RRBS), whole-genome bisulfite sequencing (WGBS), single-cell reduced-representation bisulfite sequencing (scRRBS), single-cell whole-genome bisulfite sequencing (scWGBS), TET-assisted bisulfite sequencing (TAB-seq), etc (Laird et al., 2010, Yong et al., 2016). Compared to other approaches, WGBS has two major advantages: (1) it is able to assess the methylation state of nearly every CpG site, including low-CpG-density

regions, such as intergenic 'gene deserts', partially methylated domains and distal regulatory

elements, (2) it can determine absolute DNA methylation level and reveal methylation sequence

context (Yong et al., 2016). This method has been broadly employed to analyze the genome-wide

methylation profiles of many animals, including sea cucumber (Yang et al., 2020), pig (Yang et al.,

2016), broilers (Hu et al., 2013), Chinese perch (Pan et., 2021).

In recent years, several studies have focused on the role of DNA methylation regulation on

fish during domestication. However, most studies focus on the role of DNA methylation in the

changes of one particular phenotypic characteristics, such as growth (Li et al., 2017), reproduction

(Tomasz Podgorniak et al., 2019), morphological changes (Zhang et al., 2017), learning behaviors

(Dou et al., 2018). Thus, until now, no systematic studies have investigated the role of DNA

methylation in the emergence of different selection signatures in the process of fish domestication,

including growth and metabolism, immunity, foraging and learning behaviors.

Grass carp (*Ctenopharyngodon idellus*) is one of the most important freshwater aquaculture

species, and its global production reaches 5.704 million tons in 2018, the highest in fish

production worldwide, providing low-cost, high-quality animal protein especially for developing

and underdeveloped regions (FAO, 2020). Although some farms have carried out breeding for

several generations, others still rely on wild-caught fish for broodstock to maintain genetic

diversity (Huang et al., 2015; Fu et al, 2015). In addition, farmed grass carp also shares some of

the characteristics of the domestication syndrome, for instance, changes in morphological features

(Zhao et al., 2020), rapid growth (Ashraf et al., 2011), anti-predator behavior (Tang et al, 2017),

indicate that this species could be a suitable model species for studying the role DNA methylation,

one of key molecular epigenetic pathways, in the formation of selection signatures in this and

other teleost species.

In this study, we used WGBS to carry out genome-wide DNA methylation analysis of blood

from domesticated and wild carp with four selection signatures differences: growth and

metabolism, immunity, foraging and learning behaviors. We obtained comprehensive DNA

methylation profiles for the two groups and identified differentially methylated genes (DMGs) that

might contribute to the emergence of selection signatures between domesticated and wild carp. To

the best of our knowledge, this study is the first one to report the role of epigenetic modifications

in the emergence of different selection signatures in fish. Thus, the findings here in will provide

new insights to expand our understanding about the role of epigenetic modifications in shaping

physiological phenotypes in this and other teleost models

2. Materials and Methods

2.1 Domesticated trial

A total of 15 wild sub-adult grass carp (initial weight 250 ± 20 g) were captured using traps

and gill nets along the Xijiang river in Zhaoqing, Guangdong Province, China, within a 1000 m

radius from the following coordinates (latitude: 23.1034.8 °N; longitude: 112.4554.0 °E).

Subsequently, the fish were transported to breeding base and randomly allocated into three ponds

(1.5 m × 1.5 m × 1 m), 5 individuals in each pond for subsequent domesticated experiments in the

Pearl River Fisheries Research Institute. The fish were fed commercial feed at 8:00 am and 4:00

pm each day for 180 days, and the feed amount for each day was 2~3% of fish weight. The water

temperature was kept at 22~27 °C, pH was 6.5~7.5, and dissolved oxygen was above 5.0 mg/L.

2.2 Sample Collection and Preparation

Three domesticated grass carp after six month of domestication were collected from ponds

(one fish each pond, with a body weight of 720 ± 50 g). Wild grass carp with a body weight of 550

± 50 g were collected from same place where wild sub-adult grass carp were captured, and DGC

and WGC were at approximately the same age and ontogeny stage (i.e.,puberty), which was

determined by annulus characteristics of scale (Cai et al., 2020). Whole blood was collected from

were collected from the domesticated grass carp (sample DGC1 - DGC3) and wild grass carp

(sample WGC1 - WGC3) groups. Subsequently, blood samples were treated with EDTAK2 (Sanli,

Liuyang, China) and then centrifuged at 12,000 rpms for 3 min to separate red blood cells (RBCs)

from serum.

The experimental protocols used in this study were approved by the Laboratory Animal

Ethics Committee of Pearl River Fisheries Research Institute, CAFS, China, under permit number

LAEC-PRFRI-2021-06-03.

2.3 Genomic DNA Extraction and Whole-Genome Bisulfite Sequencing

Genomic DNA was extracted from RBCs of grass carp using DNeasy Blood and Tissue Kit

following the manufacturer's protocol. Genomic DNA of RBCs from domesticated grass carp and

wild grass carp groups was sent to BGI (BGI Tech Solutions Co., Ltd., Shenzhen, China) for

whole-genome bisulfite sequencing. For normal WGBS library constructing, the DNA was

fragmented by sonication using a Bioruptor (Diagenode, Belgium) to a mean size of

approximately 250 bp, followed by the blunt-ending, dA addition to 3'-end, finally, adaptor

ligation(in this case of methylated adaptors to protect from bisulfite conversion), essentially

according to the manufacturer's instructions. Ligated DNA was bisulfite converted using the EZ

DNA Methylation-Gold kit (ZYMO). Different Insertsize fragments were excised from the same

lane of a 2% TAE agarose gel. Products were purified by using QIAquick Gel Extraction kit

(Qiagen) and amplified by PCR. At last, Sequencing was performed using the HighSeq4000 or

other Illumina platforms.

2.4 Data Filtering and Reads Alignment

After sequencing data was delivered, the raw reads were filtered by removing adaptor

sequences, contamination and low-quality reads from raw reads. Low-quality reads include two

types,and the reads meet any one of the two conditions will be removed: (1) Unknown bases are

more than10%; (2) The ratio of bases whose quality was less than 20was over 10%. After filtering,

the Clean data was then mapped to the reference genome of grass carp (Wang et al., 2015) by

BSMAP, and then remove the duplication reads and merge the mapping results according to each

library. The BSMAP script was BSMAP -a filename_1.clean.fq.gz -b filename_2.clean.fq.gz -o

filename.sam - d ref.fa -u -v 8 -z 33 -p4 -n 0 -w20 -s 16 -f 10 -L 100. The sam files were converted

to bam files using scripts (samtools view -S-b-o filename.bam filename.sam; samtools sort-m

2000000000 filename.bam filename.sort; samtools index filename.sort.bam). The mapping rate

and bisulfite conversion rate of each sample were calculated.

2.5 Identification of Differentially Methylated Regions

The methylation level was determined by dividing the number of reads covering each mC by

the total reads covering that cytosine (Lister R., 2009), which was also equal the mC/C ratio at

each reference cytosine (Xiang et al., 2010). The formula is Rmaverge = Nmall/Nmall+Nnmall.

Nm represents the reads number of mC, while Nnm represents the reads number of

non-methylation reads. Putative DMRs were identified by comparison of the sample DGC and

sample WGC methylomes using windows that contained at least 5 CpG (CHG or CHH) sites with

a 2- fold change in methylation level and Fisher test *p* value < = 0.05.

2.6 Gene Ontology and Pathway Enrichment of DMRs

GO enrichment analysis provides all GO terms that significantly enriched in a list of

differentially methylated genes, comparing to a genome background, and filter the differentially

methylated genes that correspond to specific biological functions.This method firstly maps all

differentially methylated genes to GO terms in the database (http://www.geneontology.org/),

calculating gene numbers for every term, then uses hypergeometric test to find significantly

enriched GO terms in the input list of differentially methylated genes, based on ' GO ::

TermFinder'   (http://www.yeastgenome.org/help/analyze/go-term-finder).   KEGG   pathway

enrichment analysis helps to identify significantly enriched metabolic pathways or signal

transduction pathways in differentially methylated genes comparing with the whole genome

background. The calculating formula is the same as that in GO analysis.

189
190
191
192

3. Results

3.1 Global mapping and statistical analysis of the WGBS reads

We conducted whole-genome bisulfite sequencing of grass carp blood from domesticated

grass carp group and wild grass carp group. After quality control of filtering, a total of 2.36 billion

clean reads were generated, consisting of 378.12 million, 387.63 million, and 321.7 million reads

for each domesticated grass carp sample and 371.48 million, 493.82 million, and 410.42 million

reads for each wild grass carp sample (Table 1). The bisulfite conversion rate (%) of all

sequencing libraries ranges from 99.16% to 99.28%. After read alignment, clean reads were

mapped to the reference genome of grass carp with mapping rates ranging from 88.03 % to

92.11 % (Table 1).

3.2 Global DNA methylation patterns of domesticated and wild grass carp.

The methylation levels of the whole genom were listed in Table 2. In each group,

approximately 9% of all genomic C sites were methylated. Methylation in grass carp was found to

exist in three sequence contexts: CG, CHG (where H is A, C, or T), and CHH. The average

methylation levels of CG, CHG, and CHH at the whole genome levels were 80.3%, 1.09%, and

1.08% in the domesticated grass carp group, and 75.04%, 1.07%, and 1.13% in the wild grass carp

group. The average methylation of CG showed a significantly increased within genome in the

domesticated grass carp group compared with the wild grass carp group ($p < 0.05$). The

methylation level of CHG and CHH showed no differences between domesticated grass carp and

wild grass carp groups ($p > 0.05$).

Methylation status of CG, CHG and CHH of grass carp genome showed the methylome's

overall characteristics (Figure 1). The methylation levels of approximately 25% of all mCG were

hypermethylated (methylation level > 90%). However, only about 3% of mCHH and mCHG were

hypermethylated (methylation level > 90%) compared with mCG. The methylated Cs mostly

occur in the form of mCG, followed by mCHH and mCHG. The methylation level distribution of

mC and mCG were alike. The methylated Cs mostly occur in the form of mCG; approximately

96% of all detected mCs (Table 3). Proportion of mCHG range from 0.75% to 1.15%, and

proportion of mCHH range from 2.7% to 3.05% (Table 3). The proportion of mCG, mCHG and

mCHH showed no significant difference between domesticated grass carp group and wild grass

carp group ($p > 0.05$).

3.3 DNA methylation levels of different genomic features

Different genomic features show distinct methylation patterns which are related with distinct

regulation functions (Cokus et al, 2008). The heatmap presents the methylation landscape in

different genomic features (whole genome, CGI, downstream 2 kb, upstream 2 kb, mRNA, repeat,

CDS, exon),   providing additional information as well as a global assessment of some of these

components (Figure2). CpG islands contained the highest numbers of CpG sites (approximately

10 - 20 CpG sites in a 200 bp window) compared with other genomic features. About 60% of CpG

sites in CpG islands were hypermethylated (methylation levels >90%) in the heatmap (Figure2

and Supplementary Material Figure S2). The other genomic features generally contained 0 - 10

CpG sites in the 200 bp window. A lower proportion of CpG sites within whole genome, mRNA

and repeat were hypomethylated (methylation levels < 10%) than CpG sites within remaining

genomic features (CGI, downstream 2 kb, upstream 2 kb, CDS, exon).

3.4 DNA Methylation Patterns Across the Entire Transcriptional Units at Whole Genome Level

In order to reveal the relationship between DNA methylation profiles and genes expression in

detail, Canonical DNA methylation profiles of the entire transcriptional units were divided into

distinct functional elements (Li et al., 2010) to study the changes of methylation levels in different

features (Figure 3). Methylation differences between CG and non-CpG methylation (CHG and

CHH) are visible (Figure 3), as methylation levels of CG are higher than those of CHG and CHH

across the entire transcriptional units. Another feature is a modest elevation in methylation level at

internal exons and internal intron during transcriptional unit scanning. The lowest methylation

level occurs in the first intron, followed by the first exons and down stream.

3.5 Identification and Enrichment Analysis of Differential Methylated Regions

To characterize the differences of genome methylation levels between domesticated grass

carp and wild grass carp groups, DMRs and differentially methylated genes (DMGs) were

detected. For CG context methylation, a total of 533,235 DMRs were identified between the two

groups, which corresponded to 20,010 DMGs in promoter regions and 27,016 DMGs in gene

body.

GO enrichment analysis of DMGs was performed to provide significantly enriched GO

terms corresponding to specific biological process, cellular component, and molecular function

(Figure 4). In promoter regions, the over-represented GO terms in the biological process are

cellular process, metabolic process, and biological regulation. The top enriched GO terms in the

cellular component are cellular anatomical entity, intracellular and protein−containing complex. In

terms of molecular function, the top enriched GO terms are binding, catalytic activity, and

molecular transducer activity. In gene body region, the over-represented GO terms in the

biological process are cellular process, metabolic process and biological regulation. The top

enriched GO terms in the cellular component are cellular anatomical entity, intracellular and

protein-containing complex. In terms of molecular function, the top enriched GO terms are

binding, catalytic activity and transporter activity. KEGG pathway analysis, which is an

alternative approach to categorize gene functions, was also conducted for the DMGs in promoter

and gene body regions. In promoter region, DMGs were significantly enriched in pathways in

cancer, neuroactive ligand−receptor interaction and PI3K−Akt signaling pathway. In gene body

region, DMGs were significantly enriched in metabolic pathways, pathways in cancer and MAPK

signaling pathway (Figure 4).

3.6 Candidate DMGs associated with selection signatures.

Many transcriptome and association studies have explored the molecular mechanisms

underlying selection signatures in fish and other species, laying a foundation for our investigation

of the involvement of DMGs in selection signatures. Beside, it is reported that DNA methylation,

especially in the promoter regions, usually affects gene expression by different modes (Moore et al,

2012). 38 candidate DMGs in promoter region were identified according to the following criteria:

(1) genes were differentially methylated in grass carp and wild grass carp groups; (2) genes were

enriched in pathways related to selection signatures; (3) genes were differentially expressed in fish

and other species or related with selection traits reported by previous studies. However, KEGG

pathway analysis showed that these methylated genes were not significantly enriched in predicted

process and signaling pathway (Figure 5). The protein-protein interaction network analysis

indicated that these DMGs were highly correlated with each other (Figure 6).

4. Discussion

4.1 DNA methylation profiles in domesticated and wild grass carp

Domestication is a process by which humans select some phenotypes of wild animal

species (i.e., morphological traits or growth). During the domestication process, some phenotypic

traits of animal could be altered by the artificial selection to help domesticated animal species

adapt to new environmental conditions (Sylvain et al, 2021). DNA methylation, one of the most

important and stable epigenetic modifications in eukaryotes, can lead to heritable phenotypic and

transcriptomic changes (Höglund et al., 2020). However, in aquatic species, there is still limited

research that illustrate the effects of DNA methylation on selection signatures by comparing

genome-wide methylation profiles between domesticated and wild fish species. In this study, we

identified some key differentially methylated genes related to selection signatures (including

growth and metabolism, immunity, foraging behaviors, social behaviors) in domesticated and wild

grass carp to uncover its genetic characters.

In the present study, we reported a genome-wide examination of DNA methylation in

domesticated grass carp and wild grass carp. The mapping rates of clean reads ranged from

86.25% to 92.11% (Table 1), which is above the average value reported in other fish species (Pan

et al., 2021). However, the WGBS results in this study are still consistent because bisulfite

conversion reduces the complexity of the genomic sequence and the ability of most computational

programs to align sequences onto the reference genome (Yang et al., 2020).

Further DNA methylation profile analysis was also conducted in our present study across the

distinct genomic features and entire transcriptional units. Our results showed that the

genome-wide methylation patterns were similar between two groups. Approximately 9% of all

total cytosines and 80% of CG were methylated and only a small proportions of methylation at

non-CpG methylation (CHG and CHH) was observed within all regions, which is consistent with

previous study (Cai et al., 2020). The high level of DNA methylation in CG content was a specific

characteristic of animals. DNA methylation in CHH and CHG patterns is a major characteristic of

plant methylomes and largely absent in animal methylomes (Zemach et al., 2010).

A modest elevation in methylation level at internal exons and intron was observed during

transcriptional unit scanning. The lowest methylation level occurs in the first exons, followed by

the the first intron and last exon. These results was consistent with previous findings in grass carp

(Cai et al., 2020), suggesting that mutagenic effects appear to occur at the first exons and internal

exon and intron tend to be influenced by the regulatory impact of DNA methylation.

To better understand the functional classification of DMGs during domestication, GO

enrichment analysis was conducted to gain insight into the biological processes in which the

DMGs might be involved. The top enriched GO terms of DMRs identified in the present study

were cellular process, biological regulation and metabolic process, suggesting that DNA

methylation may play a critical role in regulating global transcription during domestication.

Furthermore, to further explored the molecular mechanisms underlying selection signatures

in grass carp, 38 candidate DMGs in promoter region were identified, including IGF-1

(Insulin-like growth factor 1), GK (Glycerol kinase), GYS1 (Glycogen synthase 1), etc. But these

methylated genes were not significantly enriched in predicted process and signaling pathway of

KEGG.

4.2 Key differentially methylated genes related to growth and metabolism

Domestication is a long process which forces animals to adapt to captivity by modifying an

animal' s growth and metabolism, immune response, foraging and social behaviors. In the

process of domestication, fish undergo great change in the growth and metabolism due to the

difference of environmental conditions and food resource (Teletchea et al., 2016), which is closely

associated with the differential expression of genes involved in growth and metabolism (Shen et

al., 2021). In this study, several genes involved in growth and metabolism, such as IGF-1

(Insulin-like growth factor 1), GK (Glycerol kinase), GYS1 (Glycogen synthase 1), FASN (Fatty

acid synthase), etc, exhibited hypomethylation in domesticated grass carp group compared to wild

grass carp group. This result indicated that the expression of these genes might be upregulated. In

contrast, some genes related to growth and metabolism exhibited hypermethylation in

domesticated   grass   carp   group,   including   G6PC   (Glucose-6-phosphatase),

PCK1(Phosphoenolpyruvate   carboxykinase),   FBP1   (Fructose   1,6   bisphosphatase   1),

FOXO1(Forkhead box protein O1), ACAA1 (Acetyl-Coenzyme A acyltransferase 1), CPT1

(Carnitine palmitoyltransferase I), suggesting the expression of these genes might be

downregulated.

IGF-1, an important growth hormone, mediates the anabolic and linear growth promoting

effect of pituitary GH protein and has been well demonstrated to play an important role in the

skeletal muscle development (Laron et al., 2001). Thus, IGF-1 is thought to be a candidate gene

for regulating muscle growth.

GK is a key enzyme that catalyzes the first step in glycolysis (Enes et al., 2009). FOXO1,

G6PC, FBP1 and PCK1 are key enzymes of gluconeogenesis (Lu et al., 2018). Our results

indicated that, compared to wild grass carp, the glycolytic capacity is enhanced and the capacity

for gluconeogenesis is inhibited in domesticated grass carp feeding commercial diet. These results

was consistent with the previous finding that high-carbohydrate diet could induce glycolysis and

inhibit gluconeogenesis (Liu et al., 2021).

GSY1 is a rate-limiting enzyme in glycogen synthesis and plays an important role in the

synthesis of glycogen in the muscle (Martins et al., 2013). The present results suggested that the

capacity for muscle glycogen storage in domesticated grass carp maybe increase compared with

that of wild grass carp.

FASN, which catalyzes the last step in fatty acid biosynthesis, is believed to be the central

enzyme of hepatic lipid accumulation (Dorn et al, 2010). ACAA1, CPT1 gene play a certain role

in the lipid degradation (Song et al., 2019). In the present study, our results suggested that, in

domesticated grass carp, the ability to synthesize fat was enhanced and lipid degradation was

inhibited compared to wild grass carp, which may account for elevated lipid levels of muscle in

domesticated grass carp feeding commercial feed (Ashraf et al.,, 2011). According to the results,

an explanation for the enhanced fat deposition in fish might be excessive energy intake in

commercial diet (Wu et al., 2021). Overall, DNA methylation is likely to play an important role in

regulating growth and metabolism of domesticated grass carp by influencing these gene

expression.

4.3 Key differentially methylated genes related to immunity.

To maximize profitability, domesticated fish were cultured in intensive farming conditions

with limited space, high density and other stressors, which could influence fish immune response

negatively and even result in large-scale disease (Lin et al., 2018, Chen et al.,2017). It was

reported that, in the process of domestication, the immune status of domesticated fish could be

affected negatively due to increased chronic stress by confinement (Mandiki et al., 2011), which

would make fish more susceptible to pathogens and ultimately impair fish survival.

In this study, several immune related genes exhibited hypermethylation in domesticated grass

carp group compared to wild grass carp group, including MHCI (Major histocompatibility

complex class I), MHCII (Major histocompatibility complex class II), C1QA (Complement C1q

subcomponent subunit A), C3 (Complement C3), C4A (Complement C4-A), C5 (Complement C5),

IFN-γ (Interferon gamma), etc, which suggested the expression of these genes might be

downregulated. MHCI and MHCII are two cell surface proteins essential for the acquired immune

system for antigen presentation to recognize foreign molecules in vertebrates and can promote the

development and expansion of T cells (Huang et al., 2005). The complement system, an essential

part of both innate and adaptive immunity in teleosts, is initiated by one or a combination of three

pathways, the alternative, lectin and classical (Yang et al., 2016). IFN-γ, one of critical antiviral

cytokines, modulate functions of the immune system by up-regulate major histocompatibility

complex molecules (MHC I and MHC II) and directly activate other immune cells, such as

macrophages and natural killer cells (Borden et al., 2007). In our results, these immune related

genes mentioned above identified in domesticated grass carp exhibited hypermethylation,

indicating a poor immune performance in domesticated domesticated grass carp during long-term

domestication.

4.4 Key differentially methylated genes related to foraging behaviors

During the domestication process, fish change their foraging habits to obtain food in captive

conditions (Pasquet et al., 2018). In this study, foraging behaviors-related genes were also found to

be under strong selection in domesticated carp. The genes of GRM1(Metabotropic glutamate

receptor 1), TAS1R1 (Taste receptor type 1 member 1) and TAS1R3 (Taste receptor type 1

member 3) exhibited hypomethylation in domesticated grass carp group compared to wild grass

carp group, suggesting the expression of these genes might be upregulated. The TAS1R1 and

TAS1R3 heterodimer receptor functions as an umami receptor, responding to L-amino acid

binding, especially L-glutamate (Nelson et al., 2001). GRM1, which is widely expressed

throughout the central nervous system and regulates synaptic signaling, is another L-glutamate

receptor (Gabriel et al., 2009). Generally, these taste receptors play an important roles in

perception of L-amino acids and feeding behavior (Cai et al., 2021). Thus, these genes with

hypomethylation are likely to be upregulated in domesticated carp, facilitating the adaptation of

domesticated grass carp to commercial feed in artificial rearing condition.

4.5 Key differentially methylated genes related to learning behaviors

Learning and memory could enable the organism to plastically respond to the changing

environment. Increasing research has investigated the learning (cognitive) and memory

characteristics of fish in the past few decades, including spatial cognition, learned recognition,

social learning, foraging activity, etc (Dou et al., 2018). In this study, through domestication,

social learning-related genes have also been under selection pressure. The gene POMC

(Pro-opiomelanocortin) exhibited hypermethylation. In contrast, the genes of C-FOS

(Proto-oncogene C-FOS) and CBP (CREB-binding protein) exhibited hypomethylation in

domesticated grass carp group compared to wild grass carp group, suggesting the expression of

these genes might be upregulated. C-FOS is necessary for consolidation of non-spatial

hippocampal-dependent memory (Countryman et al., 2005), and C-FOS mRNA expressions are

up-regulated in response to a variety of neuronal activation protocols, including behavioral

training (Smulders et al., 2000) and long-term protentiation (Alberini et al., 2009). CBP is a

coactivator of transcription that play an essential role in memory consolidation (Korzus et al.,

2004). It was reported that gene expression of C-FOS and CBP facilitated the acquisition of novel

feeding habits in fish through the memory formation (Peng et al., 2019; Dou et al., 2018). Beside,

the C-FOS gene might be an important transcriptional factor to inhibit the expression of the

anorexigenic gene POMC, resulting in an increase of the food intake of dead prey fish in mandarin

fish (Peng et al., 2019). Therefore, it tempts us to speculate that the learning-related genes of

C-FOS and CBP may be responsible for the acquisition of novel feeding habits in domesticated

grass carp, which considerably increase the rate of domestication. Furthermore, individual food

intake is enhanced through the interaction between the learning gene C-FOS and the appetite

control gene POMC, contributing to the faster growth of domesticated grass carp indirectly.

424
425

Reference

Alberini, C. M., 2009. Transcription factors in long-term memory and synaptic plasticity. Physiol Rev. 89(1),121-145.

Ashraf, M., Zafa, A., Rauf, A., Mehboob, S., Qureshi, N. A., 2011. Nutritional values of wild and cultivated silver carp (hypophthalmichthys molitrix) and grass carp (*ctenopharyngodon idella*). Int J Agric Biol. 13(2), 1560-8530.

Borden, E.C., Sen, G.C., Uze, G., Silverman, R. H., Ransohoff, R.M., Foster, G.R., 2007. Interferons at age 50: past, current and future impact on biomedicine. Nat Rev Drug Discov.

Cai, Z.L., Duan, Y.J., Xu, J.C., He, Y., Geng J.Z., Liu, X., Han, M.X., Liu, Y.H., 2020 Comparison of five structure materials in age determination of Amur ide Leuciscus waleckii in Dali Lake. Journal Of Dalian Ocean University. 35, 1. DOI: 10．16535 /j．cnki．dlhyxb．2019-183

Cai, W.J., Liang, X.F., Yuan, X.C., Li, A.X., He, S., 2020. Changes of DNA methylation pattern in metabolic pathways induced by high-carbohydrate diet contribute to hyperglycemia and fat deposition in grass carp (*ctenopharyngodon idellus*). Front Endocrinol. 11, 398.

Cai, W.J., Li, J., Li, L., Chen, X., Liang, X.F., 2021. Knockout of t1r1 gene in zebrafish (*danio rerio*) by crispr/cas9 reveals its roles in regulating feeding behavior. Aquaculture, 545, 737189.

Chen, X., Wang, J., Long, Q., Gaughan, S., Wang, C., 2017. Domestication drive the changes of immune and digestive system of eurasian perch (*perca fluviatilis*). PLoS One. 12(3), e0172903.

Christoph, D., Marc-Oliver, R., Georgi, K., Michael, S., Kathrin, S., Weiss, T.S., 2010. Expression of fatty acid synthase in nonalcoholic fatty liver disease. Int J Clin Exp Patho, 3(5), 505.

Countryman, R.A., Kaban. N.L., Colombo, P.J. 2005. Hippocampal c-fos is necessary for

long-term memory of a socially transmitted food preference. Neurobiol Learn Mem. 84(3),

175-183.

Cokus, S.J., Feng, S., Zhang, X., Chen, Z., Merriman, B., Haudenschild, C.D., 2008. Shotgun

bisulphite sequencing of the arabidopsis genome reveals DNA methylation patterning. Nature.

452(7184), 215-9.

Dafni, A ., Francesc, P., 2019. Epimutations in developmental genes underlie the onset of

domestication in farmed European sea bass. Mol Biol Evol. 10, 10.

Dou, Y., He, S., Liang, X. F., Cai, W., Wang, J., Shi, L., 2018. Memory function in feeding habit

transformation of mandarin fish (*Siniperca chuatsi*). Int J Mol Sci, 19(4), 1254.

Doi, A., Park, I.H., Wen, B., Murakami, P., Aryee, M.J., Irizarry, R., 2009. Differential

methylation of tissue- and cancer-specific CpG island shores distinguishes human induced

pluripotent stem cells, embryonic stem cells and fibroblasts. Nat Genet, 41(12), 1350.

Enes, P., Panserat, S., Kaushik, S., Oliva-Teles, A. A., 2008. Nutritional regulation of hepatic

glucose metabolism in fish. Fish Physiol Biochem, 35(3), 519-539.

FAO, 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in Action. Rome.

Fu, J., Shen, Y., Xu, X., Liu, C., Li, J., 2015. Genetic parameter estimates and genotype by

environment interaction analyses for early growth traits in grass carp (*ctenopharyngodon*

idella). Aquacult Int.

Gabriel, A.S., Maekawa, T., Uneyama, H., Torii, K. 2009. Metabotropic glutamate receptor type 1

in taste tissue. Am J Clin Nutr. 90(3), 743S-746S.

Goldberg, A.D., Allis, C,D., Bernstein, E., 2007. Epigenetics: a landscape takes shape. Cell.

128(4), 635-638.

Pasquet, A., 2018. Effects of Domestication on Fish Behaviour. DOI: 10.5772/intechopen.78752.

Höglund, A., Henriksen, R., Fogelholm, J., Churcher, A.M., Guerrero-Bosagna, C.M.,

Martinez-Barrio, A., 2020. The methylation landscape and its role in domestication and gene

regulation in the chicken. Nat Ecol Evol. 4, 1713-1724.

Hu, Y.S., Xu, H,P., Li, Z.H., Zheng, X.J., Jia, X.Z., Nie, Q.H., Zhang, X.Q., 2013. Comparison of

the genome-wide DNA methylation profiles between fast-growing and slow-growing broilers.

PLoS One, 8(2), e56411.

Huang, S.X, Gilfillan, S.S, Cella., Marina., 2005. Evidence for mr1 antigen presentation to

mucosal-associated invariant t ceils. J Biol Chem, 280(22), 21183-21193.

Huang, R., Sun, J.X., Luo, Q., He, L.B., Liao, L.J., Li, Y.M., Guo, F.H., Zhu, Z.Y., Wang, W.P.

2015. Genetic variations of body weight and GCRV resistance in a random mating population

of grass carp. Oncotarget. 6(34), 35433-35442.

Jones., Peter, A., 2012. Functions of DNA methylation: islands, start sites, gene bodies and beyond.

Nat Rev Genet, 13(7), 484-492.

Korzus, E., Rosenfeld, M.G., Mayford, M., 2004. CBP histone acetyltransferase activity is a

critical component of memory consolidation. Neuron, 42(6), 961-972.

Laird, P.W., 2010. Principles and challenges of genome-wide DNA methylation analysis. Nat Rev

Genet.

Laron, Z., 2001. Insulin-like growth factor 1 (igf-1): a growth hormone. Mol Pathol MP. 54(5),

311.

Lister, R., Pelizzola, M., Dowen, R.H., Hawkins, R.D., Hon, G., Tonti-Filippini, J., 2009. Human

DNA methylomes at base resolution show widespread epigenomic differences. Nature, 462(7271), 315-22.

Li, Y., Zhu, J., Geng, T., Li, N., Zhang, X., 2010. The DNA methylome of human peripheral blood mononuclear cells. PLoS Biol. 8(11), e1000533.

Li, S.P., He, F., Wen, H.S., Li J.F., Si, Y.F., Liu, M.Y., He H.W., Huang, Z.J., 2017. Analysis of DNA Methylation Level by Methylation-Sensitive Amplification Polymorphism in Half Smooth Tongue Sole (*Cynoglossus semilaevis*) Subjected to Salinity Stress. J. Ocean Univ. China. 16 (2), 269-278.

Li, C,L., Zuo, W.D., Tong, X.L., Han, M.J., Gao, R., Hu, H., Lu, K.P., Luan, Y., Zhang, B, L., Liu,Y.Y., Dai, F.Y., 2020. Whole-genome resequencing reveals loci under selection during silkworm improvement. 138, 278-290.

Lin, W., Li, L., Chen, J., Li, D.P., Hou, J., Guo, H.H., Shen, J.Z., 2018. Long-term crowding stress causes compromised nonspecific immunity and increases apoptosis of spleen in grass carp (*Ctenopharyngodon idella*). 80, 540-545.

Fang, L., Guo, X.Z., Liang, X.F., 2021. First feeding of grass carp (*Ctenopharyngodon idellus*) with a high-carbohydrate diet:the effect on glucose metabolism in juveniles. 21, 100830.

Lu, S., Wu, X., Gao, Y., Gatlin, D.M., Wu, M., Yao, W., 2018. Effects of dietary carbohydrate sources on growth, digestive enzyme activity, gene expression of hepatic gluts and key enzymes involved in glycolysis-gluconeogenesis of giant grouper epinephelus lanceolatus larvae. Aquaculture. 343-350.

Martins, F.O., Rito, J., Jarak, I., Viegas, I., Jones, J. G., 2013. Disposition of [U-2H7] glucose into hepatic glycogen in rat and in seabass. CBPA, 166(2).

Mandiki, S., Henrotte E., Milla S., Douxfils J., Wang N., Rougeot C., 2011. How Physiological

Status and Immune Defense are Affected by Photo-thermal Regimes and Domestication

Process in Captive Eurasian Perch? ISEE. 2011.4(S8), 298–299.

Milla, S., Pasquet, A.,Mohajer, L, El., Fontaine, P., 2021. How domestication alters fish

phenotypes. 13, 388-405.

Moore, L.D., Le, T., Fan, G., 2012. DNA methylation and its basic function.

Neuropsychopharmacology. 38(1), 23-38.

Nelson, G., Hoon. M.A., Chandrashekar, J., Zhang, Y.F., Nicholas, J.P.R., Zuke, C.S., 2001.

Mammalian sweet taste receptors. Cell. 106, 381-390.

Pan, Y.X., Chen, L., Cheng, J., Zhu, X., Wu, P., Bao, L.S., Chu, W.Y., He, S., Liang, X.F., Zhang,

J.S. 2021. Genome-wide DNA methylation profiles provide insight into epigenetic regulation

of red and white muscle development in Chinese perch *Siniperca chuatsi.* 256, 110647

Pasquet, A. 2018. Effects of Domestication on Fish Behaviour. DOI: 10.5772/intechopen.78752.

Song, M., Zhong, Y., Guo, J., Chen, Y., Luo, L., Lin, S., 2019. Effects of dietary lipid levels on the

expression of cpt1 in *Micropterus salmoides*. Journal of Fisheries of China. 43, 2166-2174.

Smulders, T.V., Devoogd, T.J., 2000. Expression of immediate early genes in the hippocampal

formation of the black-capped chickadee (*poecile atricapillus*) during a food-hoarding task.

Behav Brain Res, 114(1), 39-49.

Shen, Y.W., Li, H.Y., Zhao, J.L., Tang, S.J., Chen, X.W., 2021. The digestive system of mandarin

fish (*siniperca chuatsi*) can adapt to domestication by feeding with artificial diet. Aquaculture,

538, 736546.

Tang, L.J., Jacquin, L., Lek, S., Liu, H.Z., Li, Z.J., Liu, J.S., Zhang, T.L., 2017. Differences in

anti-predator behavior and survival rate between hatchery-reared and wild grass carp

(*Ctenopharyngodon idellus*). Ann Limnol Int. 53, 361-367.

Teletchea, F., Fontaine, P., 2012. Levels of domestication in fish: implications for the sustainable

future of aquaculture. Fish Fish, 15(2), 181-195.

Teletchea, F., 2016. Is fish domestication going too fast? Nat Resour, 07(6), 399-404.

Podgorniak, T., Brockmann, S., Konstantinidis, I.,Fernandes., J.MO., 2019. Differences in the fast

muscle methylome provide insight into sex-specific epigenetic regulation of growth in Nile

tilapia during early stages of domestication. Epigenetics. 14(8).

Wilson, G.A., Pawandeep, D., Andrew, F., Daniel, C., Yuka, S., Reiner, S., Schär, P., Beck, S.,

2012. Resources for methylome analysis suitable for gene knockout studies of potential

epigenome modifiers. Gigascience, 1(1), 3-3.

Wu, B.L., Huang, L., Chen, X.J., Zhang, Y., Chen, X.J., Wu, C.C. Deng, X.J., Gao, J., He, J.X.,

2021. Effects of feeding frequency on growth performance, feed intake, metabolism and

expression of fgf21 in grass carp (*ctenopharyngodon idellus*). Aquaculture, 545(15), 737196.

Wang, Y.P., Lu, Y., Zhang, Y., Ning, Z.M., Li, Y., Zhao, Q., Lu,H.Y., 2015. The draft genome of

the grass carp (*Ctenopharyngodon idellus*) provides insights into its evolution and vegetarian

adaptation. Nat Genet. 47, 625-631.

Xiang, H., Zhu, J.D., Chen, Q., Dai, F.Y., Li, X., Wang, J., 2010. Single base-resolution

methylome of the silkworm reveals a sparse epigenomic map. Nat Biotechnol. 28(5), 516-20.

Yang, Y.J., Zheng, Y.Q., Sun, L.N., Chen, M.Y. 2020. Genome-wide DNA methylation signatures

of sea cucumber *apostichopus japonicus* during environmental induced aestivation. Genes,

11(9), 1020.

Yang, Y.L., Zhou, R., Mu, Y.L., Hou, X.H., Tang, Z.L., Li, K., 2016. Genome-wide analysis of

DNA methylation in obese, lean, and miniature pig breeds. Sci Rep-UK. 6, 30160.

Yang, Y., Yu, H., Li, H., Wang, A,L., Yu, H.Y., 2016. Effect of high temperature on immune

response of grass carp (*ctenopharyngodon idellus*) by transcriptome analysis. Fish Shellfish

Immun. 58, 89-95.

Yong, W.S., Hsu, F.M., Chen, P.Y., 2016. Profiling genome-wide DNA methylation. Epigenet

Chromatin. 9(1), 26.

Zemach, A., McDaniel, I.E, Silva, P., Zilberman, D., 2010. Genome-wide evolutionary analysis of

eukaryotic DNA methylation. Science. 328, 916.

Zhang, Y.Q., Liu, J.H., Fu, W., Xu, W.T., Zhang, H.Q., Chen, S,J., Xiao, Y.M., 2017. Comparative

transcriptome and DNA methylation analyses of the molecular mechanisms underlying skin

color variations in Crucian carp (*Carassius carassius L*). BMC Genet. 18(1), 95.

Zhao, Y.M., Zhang, L., Wang, C.F., Xie, C.X., 2020. Biology and ecology of grass carp in china: a

review and synthesis. N Am J Fish Manage. 40(6), 1379-1399.

580 **Table 1**

581 Alignment statistics with reference genome

Sample ID	Clean Reads	Mapped Reads	Mapping Rate (%)	Uniquely Mapped Reads	Uniquely Mapping Rate (%)	Bisulfite Conversion Rate (%)
DGC1	378,121,760	347,944,087	92.02	329,583,800	87.16	99.28
DGC2	387,627,636	353,434,830	91.18	335,393,994	86.52	99.23
DGC3	321,704,534	296,306,210	92.11	281,433,956	87.48	99.16
WGC1	371,478,196	321,457,393	86.53	300,678,599	80.94	99.16
WGC2	493,819,092	425,920,306	86.25	398,434,532	80.68	99.24
WGC3	410,419,590	361,301,972	88.03	339,621,985	82.75	99.22

582 Note: DGC, domesticated grass carp; WGC, wild grass carp

595   **Table 2**

596   Average methylation levels of different genomic regions.

Groups	Regions	C (%)	CG (%)	CHG (%)	CHH (%)
DGC1	Genome	9.76	80.38	0.99	1.01
DGC2	Genome	9.69	80.55	1.18	1.06
DGC3	Genome	9.68	79.98	1.11	1.17
WGC1	Genome	9.21	74.7	1.12	1.18
WGC2	Genome	9.19	75.04	1.02	1.09
WGC3	Genome	9.13	75.39	1.07	1.12

597   Note: DGC, domesticated grass carp; WGC, wild grass carp. CG, CHG, and CHH

598   (where H is A, C, or T)

603   **Table 3**

604   Proportion of CG, CHG and CHH in all Methyl-cytosine

		mCG	mCHG	mCHH
DGC1	mC number	26,890,172	208,311	750,652
	proportion (%)	96.557	0.748	2.695
DGC2	mC number	26,953,558	323,600	784,048
	proportion (%)	96.053	1.153	2.794
DGC3	mC number	26,520,365	219,138	811,594
	proportion (%)	96.259	0.795	2.946
WGC1	mC number	26,700,829	228,299	848,145
	proportion (%)	96.125	0.822	3.053
WGC2	mC number	27,258,571	220,189	831,299
	proportion (%)	96.286	0.778	2.936
WGC3	mC number	26,934,911	222,172	821,714
	proportion (%)	96.269	0.794	2.937

605   Note: DGC, domesticated grass carp; WGC, wild grass carp. CG, CHG, and CHH

606   (where H is A, C, or T)

607
608

Figure 1. Distribution of methylation level of mC in each sequence context .
The x-axis was defined as the percentage of reads showing mC at a reference cytosine site. The y-axis indicates the fraction of total mC calculated within bins of 10%. DGC, domesticated grass carp; WGC, wild grass carp. (A) Sample DGC 1; (B) Sample DGC2; (C) Sample DGC3; (D) Sample WGC1; (E) Sample WGC2; (F) Sample WGC3.

Figure 2. Heat maps show distinct methylation and CpG density patterns.
CpG density (x-axis) is defined as numbers of CpG dinucleotides in 200bp windows. Methylation level (y- axis) is defined as average methylation level of cytosines in CpGs. The thin black lines within each heat map denote the median methylation level of CpGs at the given local density. The red color gradient indicates abundance of CpGs that fall into bins of given methylation levels and CpG densities. The blue bar charts above each heat map show the distribution of CpG densities, projected onto the x-axis of the heat maps. The green bar charts to the right of the heat maps show the distribution of methylation levels, projected onto the y-axis of the heat maps. DGC, domesticated grass carp; WGC, wild grass carp. (A) Sample DGC 1; (B) Sample DGC2; (C) Sample DGC3; (D) Sample WGC1; (E) Sample WGC2; (F) Sample WGC3.

A

B

C

D

E

F

Figure 3. DNA Methylation Patterns Across the Entire Transcriptional Units at Whole Genome Level of sample DGC1. The canonical gene structure is defined by 7 different features, denoted by the x-axis. The length of each feature was normalized and divided into equal numbers of bins. Each dot denotes the mean methylation level per bin and the respective lines denote the 5-bin moving average. Each feature was analyzed separately for the numbers listed in the table below the figure. The green vertical line indicates the mean location of the transcription start sites. DGC, domesticated grass carp; WGC, wild grass carp. (A) Sample DGC 1; (B) Sample DGC2; (C) Sample DGC3; (D) Sample WGC1; (E) Sample WGC2; (F) Sample WGC3.

Figure 4. GO analysis of differentially methylated regions (DMRs)-related genes.
The x-axis represents three domains of GO while the y-axis represents the gene number in every pathway and processes.
(A).GO analysis of differentially methylated regions (DMRs)-related genes in promoter region
(B).GO analysis of differentially methylated regions (DMRs)-related genes in gene body region

Figure 5. Pathway analysis of DMRs-related genes

The abscissa represent the richness factor, the ordinate represent the enriched pathway terms. Q-value represents the corrected P, and a small Q-value indicates high significance.

(A)Pathway analysis of DMRs-related genes in promoter region; (B) Pathway analysis of DMRs-related genes in gene body region.

Figure 6. Protein-protein interaction network analysis of DMGs in different selection signatures.