1 **A multivariate view of cognitive performance reveals positive correlation in the Trinidadian Guppy**

2 **(*Poecilia reticulata*)**

5 Pamela M Prentice[1], Alex Thorton[1], Alastair J Wilson[1]

[1]Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK

Abstract

Cognitive variation is common among-individuals and can be consistent across time and

context. From an evolutionary perspective, among-individual variation is important as a pre-requisite for

natural selection and adaptive evolution. Selection is widely hypothesized to favor high cognitive

performance but directional selection should erode variation over time, how then is cognitive variation

maintained? As selection does not act on traits in isolation, covariance among specific cognitive traits

and/or other aspects of phenotype (e.g. personality) could result in fitness trade-offs that are important

in shaping evolutionary dynamics. Here we test this using Trinidadian guppies (*Poecilia reticulata*), using

a multivariate approach by characterising the correlation structure among task-specific cognitive

performance measures and a personality trait. We estimate the among-individual correlation matrix

(**ID**) in performance across three cognitive tasks; colour association learning task; motor learning task;

reversal learning task, and the personality trait 'boldness' measured as emergence time from a shelter.

We found no support for trade-offs among performance in these tasks. Nor do we find evidence of

hypothesised speed-accuracy trade-offs within the association learning task. Rather we find strong

positive correlation structure in **ID,** with 57% of variation explained by the leading eigen vector. While

noting that non-cognitive factors and assay composition may affect the structure of **ID**, we suggest our

findings are consistent with the *g*-model of cognitive performance variation, in which a dominant axis of

variation loads positively on all performance measures. Thus, we add to a growing body of support for

general variation among individuals in animal cognitive ability.

Cognitive performance, among-individual variation, cognitive evolution, multivariate mixed model,

correlation matrix, Trinidadian guppy

Introduction

Interest in the cognitive mechanisms by which animals acquire, process, store and use information from the environment (Healy & Rowe, 2010; Shettleworth, 2010), has grown substantially in recent years among ecologists and evolutionary biologists. Differences in cognitive abilities between species have long been recognised (Wasserman & Zentall, 2006; Shettleworth, 2009), but we now know that individuals within a species can differ in cognitive performance in a manner that is consistent across time and context (for a review, see Cauchoix et al., 2018, but see Troisi et al., 2021 for an exception to this). From an evolutionary perspective, variation at this level is required for natural selection and is thus central to our understanding of adaptive evolutionary dynamics (Roff, 2002). However, how or why this variation is maintained within populations is not always clear. For instance, while we might intuitively expect selection to favour high cognitive performance, directional selection of this form is generally expected to erode variation over time. One possibility is that fitness trade-offs could shape patterns of variation in cognition, just as they do in other aspects of phenotype (e.g. life history; Stearns, 1992). If so, do trade-offs arise among different cognitive 'domains' or traits, with variation maintained because individuals (or genotypes) that perform better than average in some respects, perform worse in others? Here we test this idea using a multivariate approach to characterise and test hypothesised relationships of cognitive performance between multiple variables measured within three cognitive tasks using the Trinidadian guppy (*Poecilia reticulata*). We ask whether there is support for negative correlation structure, as expected under the trade-off hypothesis, or whether in fact correlations among performance measures are uniformly positive – a pattern predicted if multivariate cognitive variation is explained by a single general cognitive performance factor or "general intelligence" (*g*) (Nisbett et al., 2012).

Many empirical studies have now shown that animal populations can harbour high levels of variation in cognitive performance among individuals (Boogert et al., 2018), and this has been evidenced across different taxonomic groups, including insects (e.g. Li et al., 2017), fish (e.g. Lucon-Xiccato & Bisazza, 2017), birds (e.g. Ashton et al., 2018) and mammals (e.g. Mazza et al., 2019). Among-individual variation is also found across different aspect of cognition (e.g. spatial memory (Sonnenberg et al., 2019); association learning (Kniel et al., 2020), inhibitory control (Brandão et al., 2019)). Although

numerous studies have explored relationships between cognitive performance in different cognitive

tasks (for a review see Burkart et al., 2017; Boogert et al., 2018), relatively few studies have

characterised the among-individual covariance (or correlation) structure between these. This is

important to do because trade-offs between domains have been hypothesised to maintain cognitive

variation at the individual level (Del Giudice & Crespi, 2018). This hypothesis leads to an expectation of

negative correlation structure which is exemplified by predictions of 'trade-offs' in performance across

different cognitive domains (Shettleworth, 2012; Thornton & Lukas, 2012).

It has been hypothesised that trade-offs, such as those observed between learning and cognitive

'flexibility' (Del Giudice & Crespi, 2018), could maintain variation in performance. Flexibility is broadly

defined as the ability to adapt when environmental stimuli or information cues change. It is often tested

by reversal learning experimental paradigms in which individuals must be flexible to override old cue-

reward associations and form new ones (Bitterman, 1965; Kehagia et al., 2010; Buechel et al., 2018) .

For example, in Florida scrub-jays (*Aphelocoma coerulescens*) individuals that were quick to learn an

initial colour-reward cue (high associative learning performance), were slower to adjust when the cue

signal was reversed (Bebus et al., 2016). The implication of such patterns is that some individuals are

more 'intrinsically driven' than others; they learn initial associations quickly but, being less sensitive to

external stimuli, struggle to adapt when cues are altered. However, counter-examples can be found. For

instance, those individual bumblebees (*Bombus terrestris*) that quickly learned to discriminate between

two colours (where one was associated with a floral reward), were also faster to learn a new association

when the cues were reversed (Raine & Chittka, 2012).

Trade-offs have also been postulated to maintain variation in performance within cognitive

domains. Speed-accuracy trade-offs are perhaps the best known example (Chittka et al., 2009). Faster

decision-making should provide a competitive advantage (e.g. by increasing the potential rate of

resource acquisition), but by allowing less time to assess environmental cues, error rates may be

increased. In contrast, slower decisions may be more accurate, but being slow to act can mean

resources are lost to competitors. A number of empirical studies have provided support for speed-

accuracy trade-offs. For example, individual archerfish (*Toxotes chatareus*) that tended to make slower

decisions within a colour discrimination task also had greater accuracy (Jones et al., 2020). A similar

result was found in zebrafish (*Danio rerio*) given a visual discrimination task (Wang et al., 2015), and in

wild-caught great tits (*Parus major*) given a foraging task (Moiron et al., 2016). However, once again

counter examples also exist. For example, a prior study of guppies found no support for a speed-

accuracy trade-off using a shape and a colour discrimination task (Lucon-Xiccato et al., 2016), and three-

spined sticklebacks (*Gasterosteus aculeatus*) making fast decisions in a spatial leaning task did not show

reduced accuracy (Mamuneas et al., 2015).

Thus, while trade-offs between and within components of 'multivariate' cognitive ability are

intuitive and evidenced in some cases, they may not be inevitable and positive correlations between

traits and domains are also found (Guenther & Brust, 2017; Wallace & Hofmann, 2021). If correlation

structure among cognitive performance traits is universally positive (with respect to expected fitness

consequences), then variation among individuals can be explained by a domain-general structure that

invokes a single latent general cognitive performance or "intelligence" factor, sometimes denoted '*g*'

(Deaner et al. 2006; Lefebvre and Sol 2008). While there is evidence for this in humans (Deary et al.,

2010; Burkart et al., 2017), support for the *g*-model in non-human animals systems remains limited (but

see (but see Hopkins et al., 2014; Shaw et al., 2015; Arden & Adams, 2016; Ashton et al., 2018). Note

that, under this model, trade-offs may still be important for maintaining cognitive variation, but if so

they must operate between general cognitive performance and other (non-cognitive) aspects of

phenotype. This possibility is mirrored elsewhere in the behavioural literature since for example,

individual personality can be viewed as a component of an extended 'life history' (Sih et al., 2004; Wolf

et al., 2007). Indeed, given that personality traits and cognitive performance are frequently correlated

(Guenther et al., 2014; Nawroth et al., 2017; White et al., 2017), and both thought to drive variation in

resource acquisition, then trade-offs between them are plausible (Sih & Del Giudice, 2012; Dougherty &

Guillette, 2018).

Here, we investigate among-individual (co)variation between multiple measures of cognitive

performance variables across three cognitive tasks (associative learning task, motor learning task and a

reversal learning task) in a captive population of wild-type guppies (*Poecilia reticulata*). Guppies have

been widely used as a model in behavioural and evolutionary ecology, and methods for assaying among-

individual variation in cognitive performance are well established (Laland & Reader, 1999; Miletto

Petrazzini & Agrillo, 2016). This species is known to perform well (on average) in tests of associative

learning (Lucon-Xiccato & Bisazza, 2016; Kniel et al., 2020), motor learning (Lucon-Xiccato & Bisazza,

2016; Lucon-Xiccato et al., 2017), and cognitive flexibility (Cauchoix et al., 2018; Fong et al., 2019).

However, we know little about if and how cognitive performance measures within these distinct tasks

covary among individuals. Broadly stated, our goal is to estimate the structure of this multivariate

variation to evaluate whether trade-offs among cognitive performance measures are evident, or

whether among-fish variation is consistent with general cognitive performance (g). We also seek to ask

whether trade-offs shape variation in cognitive performance within cognitive tasks, and whether

personality variation is linked to, or trades off against cognitive performance variation in this species.

We note that the extent to which among-trait associations shape, and in the particular context of

trade-offs, constrain evolutionary adaptation strictly depends not on the phenotypic correlation

structure, but on the genetic contribution to this. In particular, evolutionary constraint arises from the

genetic (co)variance structure (**G**) and its alignment (of lack thereof) with selection (Jones et al., 2004;

Lande, 1979; Walsh et al., 2009). Unfortunately estimation of **G** is challenging in general (requiring large

volumes of data from related individuals; Wilson et al., 2010), and particularly so for cognitive

performance measures. This is because high-throughput phenotyping of cognitive performance across

multiple tasks is notoriously laborious and challenging. Consequently, very few studies have estimated **G**

for sets of cognitive performance measures in non-human animals (but see Langley et al., 2020). Thus

most multivariate studies of cognition, and behavioural phenotypes more generally, rely on Cheverud's

conjecture (Cheverud, 1988) that phenotypic patterns of covariation can be used to infer evolutionarily

important relationships between traits. Accepting this view uncritically implies that the (co)variance

structure of the phenotypic matrix **P** (historically referred to as the phenotypic gambit; Grafen, 1984),

which can be readily estimated in a set of individuals observed once only for each of the target traits,

can yield robust insights into the structure of **G**. However, behavioural traits and cognitive performance

measures are typically very plastic and often subject to high levels of measurement error (e.g. relative

to morphological traits), in addition to sources of environmental variation. These phenomena lead to

high levels of within-individual variation such that a single observation may tell us relatively little about

an individual's phenotype (and so genotype) (Brommer, 2013). In other words, the extent to which **P** is a

good proxy for **G** will decline if **P** is dominated by within-individual variation (plasticity, measurement

error) versus among-individual variation (which includes genetic factors).

In the present study we therefore adopt an intermediate approach in which we estimate the

among-individual (co)variance matrix (**ID**) rather than either **G** or **P**. We do this by obtaining repeated

measures of performance on individual fish. Since **G** is a component of **ID,** while **ID** is a component of **P**,

all else being equal we expect the among-individual covariance matrix to be better proxy of the genetic

matrix than the total phenotypic matrix. We target multiple cognitive performance measures across

three cognitive tasks (association learning task, motor learning task and a reversal learning task)

allowing us to: 1) test whether individuals differ in their cognitive performance between distinct

cognitive tasks. We predict that individuals will differ in all measures of cognitive performance within all

three tasks; the association task, the novel motor learning task, and the reversal learning task as well as

in their rates of improvement with experience (interpretable as learning). We then estimate the **ID**-

matrix among all cognitive performance measures across the three different cognitive tasks, in addition

to a shy-bold type personality trait (henceforth referred to as 'boldness'). We ask 2) whether the **ID**-

matrix provides evidence for trade-offs among cognitive performance measures (exemplified as a

negative correlation structure in the **ID**-matrix) or whether it is dominated by positive correlations

consistent with the *g*-model of general cognitive performance; 3) whether speed accuracy trade- offs

are present within both the association task and reversal learning task; and finally 4) whether among-

individual variation in shy-bold personality trait 'boldness' is correlated with cognitive performance

variation in the three cognitive tasks.

Methods

Ethics

This work was conducted under the auspices of the Animals (Scientific Procedures Act) with

approval of the University of Exeter research ethics committee, under licence from the Home Office

(UK) (Licence Number PPL30/3256). Experimental procedures and behavioural assays were developed

in accordance with the principles of the three Rs and ASAB guidelines (Buchanan et al., 2020). All

periods of handling and emersion were kept to a minimum and only fish deemed healthy and exhibiting

normal behaviour were used in trials. At the end of the experiment, fish were returned to a designated

'retirement' tank (containing females as well as males) and not used in any further experiments.

Husbandry

All behavioural assays were carried out between October 2019 and January 2020 inclusive. Data was

collected from captive-bred guppies bred and housed at the fish laboratory at the University of Exeter's

Penryn campus. The population is descended from wild fish caught in February 2017 from the lower

Aripo River, Trinidad and has been subsequently maintained with no deliberate selection or inbreeding.

All fish housed in the laboratory were fed to satiation twice daily (0800 – 1000h and again at 1600 –

1800h) using commercial flake food and live *Artemia nauplii*. Water temperature was maintained at 23-

24°C in well-aerated closed system tank stacks that undergo 25% water changes each week and with

weekly tests for ammonia, nitrate and nitrite levels. Lighting was kept at a 12:12 light/dark cycle.

Experimental apparatus

Adult males (n = 43) and females (n = 37) were sampled from the stock population. Sampling was

random but we approximately size matched fish within each sex (e.g. by avoiding very large females).

Fish were then housed singly in separate tanks (15 l, 18.5 × 37 × 22cm) for the duration of behavioural

testing. The tank set up closely followed that used by Lucon-Xiccato & Bisazza (2014), each being divided

equally into a 'home' compartment at the rear of the tank, and a 'test' compartment at the front of the

tank, separated into two compartments (using white plastic) by a guillotine door (Figure 1a). The rear

'home' compartment (20 x 18.5 cm) allowed individuals visual access to fish in neighbouring tanks and

contained foliage to provide coverage and environmental enrichment (Brydges & Braithwaite, 2009).

Conversely, the 'test' compartment (17 x 18.5 cm) was screened from neighbours (using white plastic)

to prevent any possibility of social learning (i.e. by observation of neighbours) influencing cognitive task

performance. The 'test' compartment contained a white plastic plate (4 x 10 cm) placed on the gravel

substrate, perforated with 2 equally spaced wells (Figure 1b). A total of 48 experimental tanks were

used. These were contained within two 'stacks', each comprising 24 tanks (8 tanks per row, 3 rows high)

on a shared recirculating water supply. As the testing protocol (described below) took 18 days per fish,

data were in practice collected in 3 'blocks' over a total testing period spanning 12 weeks. Before

cognitive testing all fish were allowed to acclimate to experimental tanks for 48 hours. During this

period they were fed twice daily with bloodworm (*Chironomidae* larvae) pipetted into one (randomly

chosen) well of the white plastic plate and the guillotine door was left open allowing unrestricted use of

both compartments.

Cognitive assay

We used a repeated measures design to test multiple cognitive performance measures across

three different cognitive tasks within a single, extended, testing paradigm. Each fish was observed up to

63 times as it emerged from the home compartment to obtain a food item that had been placed in one

of two wells in the test compartment. Individual fish were trained to discriminate between a rewarded

and an unrewarded colour cue, indicating which well the reward could be obtained from. In all trials,

food items (bloodworm) were placed in both wells, but access was restricted by a small plastic disc in

the well with the unrewarded colour cue. This was to ensure that olfactory cues would be insufficient to

locate available food, instead fish were required to learn to associate a specific colour with reward

access. The rewarded colour (either blue or green) was randomly determined for each fish prior to

testing, ensuring balance across subjects to control for any pre-existing colour bias. The relative position

of the rewarded well (left versus right) for each trial was randomised and we also controlled for any

effect this could have in our statistical modelling (described later).

228   The 63 observations per individual were grouped into 7 'sets' of 9 'trials' (observations). Each

set spanned a 2 day period, which was followed by a rest day before the next set. Test conditions were

altered between trial sets, by adjusting the position of green and blue coloured plastic counters.

Specifically, after each set, the association task increased in difficulty as the coloured counter covered

the holes to an increasing extent. This also allowed extraction of information on different cognitive traits

as we now describe. In set 1, the coloured counters covered 0% of the wells, in set 2 we increased this

to 25%, then subsequently to 50% (set 3), 75% (set 4) and 100% (set 5). Once the counters covered 75%

of the wells (set 4), guppies had to dislodge the counters in order to reach the reward in the well

underneath. We consider this a novel motor learning task, as fish had to learn to dislodge the counter by

physical manipulation. The difficulty of this task is increased with 100% coverage (set 5). In the final two

sets (set 6 and 7) of 9 trials we conducted a reversal learning task, by reversing the colour cue -food

reward association previously learnt in the first 5 sets.

240   Prior to each trial, fish were guided into the 'home' compartment of the tank with a net, and

the guillotine door was closed. The experimenter set up the plate in the 'test' compartment, by

pipetting bloodworm into the wells and the two coloured counters placed either to the side of (set 1), or

covering the rewarded well (partially or fully depending on set number as described). The food in the

unrewarded hole was covered completely by a small plastic disc. The trial started once the door to the

experimental compartment was opened and continued until the fish ate the bloodworm from the

rewarded well, or until 10 minutes, whichever occurred first (see the electronic supplementary material,

video S1). All trials were recorded with a GoPro Hero 6 camera, mounted in front of the tank and

behavioural scoring of videos was quantified using the software BORIS (Friard & Gamba, 2016). We

imposed a deliberately weak learning criterion, removing any fish that failed to achieve a minimum of

5/9 successes (defined as locating and/or eating the food reward) at the end of each set. In fact, we

would have ideally avoided selective removal of fish such that among-individual variation seen in later

sets can be interpreted as fully representative of the initial population sampled. However, it is also the

case that estimated correlations between task-specific performance may be misleading if the ability to

solve each task is closely associated with the number of prior learning opportunities (Guez & Griffin,

2016; Shaw & Schmelz, 2017), which in turn depends on differences in, for example, motivation or

neophobia. Furthermore, since additional food (beyond the reward items) was not provided during the

behavioural testing, the removal strategy adopted ensured that any individuals not engaging with the

task at all could not be food deprived for more than 48 hours. Over the course of the data collection, 16

individuals were removed before the end of set 6.

Cognitive performance measure definitions

Data obtained from the trials were used to define observed proxies of traits, including one

personality trait (*boldness*) and six measures of cognitive performance that were collected during three

distinct cognitive tasks (*associative learning, motor learning,* and *reversal learning*). In some cases

cognitive performance measures were defined during trials in a single set (9 trials in total), but where

data from 2 sets are equally relevant (18 trials in total); these were combined to maximise informative

observation number and hence maximise statistical power. The specific personality trait and cognitive

performance measures are described below and are summarised for reference in Table 1.

Boldness

During the first 9 trials (set 1), time taken to emerge from the home compartment into the experimental

compartment was recorded (*emergence*)*,* as a proxy of consistent differences in personality trait

'boldness' (White et al., 2016, 2020).

Associative learning

We focused on the first 9 trials (set 1) to characterise associative learning and test for speed-accuracy

trade-offs. A similar study to the current testing paradigm, guppies learnt a colour association in

9.18+2.50 trials (Bisazza et al., 2014), as such we expected that (on average) fish would have effectively

learnt the association by the end of set 1 such that any speed accuracy trade-off would be less relevant

to observed outcomes in later sets. We recorded three cognitive performance measures in each of the 9

trials in set 1. The first measure was whether the first counter explored in each trial was the correct (1)

or incorrect (0) choice for obtaining the reward ($AL_{accuracy}$). A choice was deemed to have been made if

individual fish swam within 1 body length of, and was actively exploring a counter (with head and body

directed towards the stimulus counter). We note that accuracy of the first counter explored in trial 1 of

set 1 will be by chance alone due to the absence of prior experience. The second cognitive performance

measure was time taken to make this first choice in each trial (AL_{speed}), measured in seconds from

emergence into the test compartment. To evaluate overall performance in association learning, we used

total time taken from emergence to find and eat the food reward in each trial (AL_{time}). Note, when

modelling this variable we maximised observation number (and hence statistical power) by using

observations from 18 informative trials; 9 trials in set 2 and 9 trials in set 3. During set 2 and 3, the

counters covered 25% and 50% of the wells respectively. This level of coverage still allowed fish to

access the reward without any physical manipulation of the counter.

<u>Motor Learning</u>

To assess performance in the novel motor learning task, we used observations from 18 informative

trials; 9 trials in set 4 and 9 trials in 5 (where the counter covered 75% and 100% of the well area

respectively). Note that at 75% coverage fish were required to physically manipulate the counter in

order to reach the food reward. We used time taken by each individual in all trials to obtain and eat the

food reward (ML_{time}), measured as time taken to obtain the food reward after they had made the

correct choice of counter in each trial.

<u>Reversal learning</u>

This was assessed using observations from 18 informative trials: 9 trials in set 6, and 9 trials in set 7.

During these trials the reward colour was reversed. Note that counter coverage of the wells remained at

100% and therefore fish required to dislodge the counter in order to access the reward. For this reversal

learning task we recorded the accuracy of each individual's first choice in every trial in set 6 and 7,

($RL_{accuracy}$) as well as the total time taken, from emergence into the test compartment, to locate and eat

the food reward in each trial (RL_{time}).

Statistical analysis

We used univariate and multivariate linear mixed effect models to characterise among-individual

variation in and covariation among the personality trait and cognitive performance measures defined.

For analysis, the observed (censored) time variables from the cognitive tasks were natural log-

transformed (AL_{speed}, AL_{time}, ML_{time} and RL_{time}), and then all continuous time variables were multiplied by

-1 (*boldness*, AL_{speed}, AL_{time}, ML_{time} and RL_{time}). Following this, all response variables were mean centred

and scaled to standard deviation units. The log transformation was to improve the assumption of

Gaussian error structure, multiplication by -1 was just to ease of interpretation (making larger values

equate to higher performance). Scaling to standard deviation units was also to simplify interpretation of

estimated variance components (explained further below). All models were fit by restricted maximum

likelihood (REML) using ASReml-R 4.1 (Gilmour et al., 2009; Butler et al., 2018), R version 3.6.1 (R Core

Team, 2017). We make the standard assumptions that random effects and residuals are normally

distributed with means of zero and variances to be estimated. We acknowledge that these assumptions

are necessarily violated since (i) the two accuracy traits are recorded as binary variables and (ii) latency

(time) traits are censored at 10 minutes. However, we consider this strategy both pragmatic and

justifiable given that Gaussian mixed models are largely robust to violations of distributional

assumptions (Schielzeth et al., 2020) while (Bayesian) multivariate generalised mixed models remain

challenging to implement.

 <u>Univariate analysis</u>

330 First, we fit separate univariate mixed models to test for and characterise among-individual variation in

331 each of our 7 traits (*boldness*, *AL_speed*, *AL_accuracy*, *AL_time*, *ML_time*, *RL_time* and *RL_accuracy*). All models included a

332 fixed effect of the mean, as well as fixed factors of *sex*, *stack* (denoting which of two aquaria stacks the

333 fish was tested in), *colour* (denoting the colour of counter individual fish were trained to associate with

334 the rewarded cue; green or blue), and *reward side* (denoting which side the rewarded counter was

335 placed on; left or right). We also included *trial number* (i.e. the repeat number of trial, fitted as a factor

336 to avoid assuming a linear functional form for any change in the mean across repeats). For each trait, we

337 firstly fit this 'null' model without random effects, and then refitted with *individual identity* as a random

338 intercept. For each trait, we compared the full (i.e. with *individual identity* included) and null models by

339 likelihood ratio tests (LRT) to obtain a statistical test of the among-individual variance (V_I). To test a

340 single variance component (which cannot be less than zero) we assume twice the difference in log-

341 likelihood between the full and reduced models is distributed as a 50:50 mix of $\chi^2_{0,1}$ and χ^2_1 as

342 recommended by Visscher (2006). Since transformed traits are analysed in standard deviation units,

343 total observed variance is 1 and V_I can actually be interpreted as an estimate of repeatability. However,

344 we also calculated estimates of the adjusted repeatability (R) which is the proportion of phenotypic

345 variance conditional on fixed effects that is explained by among-individual differences (Nakagawa &

346 Schielzeth, 2010). Thus $R = V_I / V_P$ where V_I is the among-individual variance and V_P is sum of V_I and V_R,

347 the residual or within-individual variance. Conditional F-statistics were used to determine the

348 significance of fixed effects in each (full) model, and we elected not to perform model simplification, as

349 we wanted repeatability estimates to be conditioned on a common set of fixed effects and thus

350 comparable across abilities.

352 <u>Multivariate analysis</u>

353 To test for the presence of, and investigate the structure of, the (co)variance **ID**-matrix, we built a

354 multivariate mixed model among all cognitive performance traits and 'boldness'. All response variables

355 were transformed and scaled as per the univariate modelling procedure above, and we note that

multiplication by -1 of all the observed (censored) time traits simplified interpretation of the correlation

structure (as larger values equate to higher performance in continuous traits). Fixed and random effects

were fitted on each trait as specified for the univariate models. Fixed effects estimates are reported in

the supplementary information (Table S1). We specified **ID** as an unstructured matrix to be estimated.

Note that observation-level (residual) covariances are not statistically identifiable if traits are not

measured at the same time. We thus specified a residual covariance matrix structure in which

covariances between traits measured in different sets were fixed to zero. We tested for overall among-

individual covariance among the traits by comparing this model against a reduced one in which **ID** was

specified as a diagonal matrix (i.e., among-individual variances are estimated but covariances are

assumed to equal zero).

To aid biological interpretation of the **ID**-matrix, we rescaled our estimate to the corresponding

correlation matrix (noting that for any pair of traits (x,y), $r_{I(x,y)} = COV_{I(x,y)} / (V_{I(x)}V_{I(y)})^{0.5}$). We further

scrutinised the correlation (r) structure among cognitive (c) traits only using **ID_{rc}**, defined as the 6x6

submatrix of **ID** obtained after putting on a correlation scale and dropping the row/column

corresponding to boldness. **ID_{rc}** was subject to eigen vector decomposition to provide a descriptive view

of the major axes (or principle components) of variation. Under the *g* model, we would expect uniformly

positive correlation structure in **ID_{rc}** which would be reflected by a dominant first eigen vector loaded on

all traits with the same sign. Conversely, if trade-offs dominate the structure of among-individual

variation in the multivariate cognitive phenotype **ID_{rc}** should contain at least some negative correlations

and one or more important principle components with loadings that are antagonistic in sign between

traits in trade-offs. We estimated approximate 95% confidence intervals on the eigen values (scaled to

proportions of variation in the matrix) and on the trait loadings associated with each principal

component using a parametric bootstrap approach described in Boulton et al (2014).

Results

Visualisation of the raw data shows a varied effect of trial number on the personality trait 'boldness',

and all performance measures across the cognitive tasks (Figure 2; see Supplemental Information Figure

S1 a–g of raw data across all sets of trials). Patterns of improvement are observed in the raw data for

the following: average time to *emergence* (boldness; Fig 2a); AL_{speed} (Fig 2b) and $AL_{accuracy}$ (Fig 2c)

measured within the first set of trials of the association learning task; and both RL_{time} (Fig 2f) and

$RL_{accuracy}$ (Fig 2g) measured within the reversal learning task (although the trend in the former is weak).

Time-based measures of cognitive performance for AL_{time} and ML_{time} did not uniformly improve (i.e.

reduce) over the sets of trials used to assay them, as might perhaps be intuitively expected (Figure 2).

Note however, that (i) we deliberately chose to use AL_{time} recorded from sets 2 and 3 to capture

variation in the extent to which the association had been learned after the experience of set 1 (during

which rapid improvement in mean performance is seen as predicted; Fig S1d). Additionally, (ii) both

AL_{time} and ML_{time} were assayed using 2 consecutive sets of 9 trials separated by an increase in task

difficulty (i.e., change in position of the counter between sets) imposed between trial 9 and 10. In fact

ML_{time} does show a general pattern of improvement within each set separately (sets 4 and 5;

Supplemental Information Fig S1e) that is not apparent from the linear trend across all 18 trials (Fig 1e).

395   The univariate mixed models confirmed the statistical significance of these qualitative patterns

in the raw data with respect to changes in mean performance across trials (see Supplementary

Information Table S1 *a–g)*. Noting that models were run on transformed data such that positive effects

always equate to an increase in performance. Furthermore, *trial number* was fitted as a factor to avoid

assuming linear trends. Noting these points, we found significant effects on the following: *emergence*

($F_{8, 555.7}=9.24$, P< 0.001; with coefficients becoming more positive as *trial number* increased; Table S1)*,*

AL_{speed} ($F_{8, 555.0}=7.380$, P< 0.001; with coefficients becoming more positive)*, $AL_{accuracy}$ ($F_{8, 555.0}=4.602$, P<

0.001; with coefficients becoming more positive), AL_{time} ($F_{17, 801.9}=2.246$, P=0.002; with coefficients

becoming more negative, indicating a decrease in performance)*, and ML_{time} ($F_{17,692.6}=1.174$, P< 0.001;

coefficients becoming more negative, indicating a decrease in performance). The effect of *trial number*

on (mean) performance was not significant in RL_{time} ($F_{17, 581.2}=1.565$, P= 0.060) and $RL_{accuracy}$ ($F_{17,}$

584.2=1.085, P= 0.271). The significance and magnitude of other fixed effects included is not directly

relevant to any hypotheses being tested and we do not discuss them further, but they are reported in

full in the supplemental materials (see Supplementary Information Table S1).

With respect to among-individual variation, likelihood ratio tests (LRT) on univariate mixed

models provided strong statistical support for the presence of among-individual variance in all seven

traits tested (LRT of model 1 vs. model 0; all $P < 0.001$; Table2). Repeatabilies (R) estimated conditional

on fixed effects (i.e. as $R = V_I/V_P = V_I/(V_I+V_R)$) were low to moderate, ranging from 20-46% with a median

across the seven traits of 33.5% (Table 2). As well as among-individual variance, the multivariate

modelling shows strong support for among-individual covariance in **ID**. Likelihood ratio comparison of

the full model to a reduced fit in which all cross-trait covariances in **ID** were set to zero showed the

former was a significantly better fit to the data ($\chi2_{21}$=242.756, P<0.001). Among variables measured, the

estimated covariances were positive between all trait pairs (Table 3). Boldness, as captured by

emergence was strongly (and significantly; based on assuming approximate 95% CI of r_I ±1.96 SE)

correlated with cognitive performance as measured by all variables in the association learning task ((r ±

(SE)), AL_{time} = 0.663 (0.095), AL_{speed} = 0.818 (0.046), $AL_{accuracy}$ = 0.788 (0.062); Table 2), and to decision

accuracy within the reversal learning task ($RL_{accuracy}$ = 0.412 (0.171)).

Among the subset of 6 cognitive performance measures, all estimated among-individual

correlations (r_I) were positive (Table 3). This included among traits in the association task ($r_{ALtime. ALspeed}$ =

0.758 (0.074), $r_{ALtime. ALaccuracy}$ = 0.729 (0.092), $r_{ALspeed. ALaccuracy}$ = 0.894 (0.039)) meaning that we find no

support for a speed-accuracy trade-off here. Similarly, both variables in the reversal learning task ($r_{RLtime.}$

$_{RLaccuracy}$ = 0.473 (0.144)) were positively correlated. There was thus no evidence of negative correlations

as predicted by specific trade-offs, either within or between cognitive tasks, and rather the correlation

structure of **ID** is uniformly positive, consistent with the *g*-model of general cognitive performance

(Table 3).

Eigen decomposition of **ID$_{rc}$** reflects this, with the first major axis (first principle component,

PC1, with 95% confidence intervals from 5000 bootstrap replicates) explaining 57.1% (42.7%, 69.8%) of

the among-individual variation in multivariate phenotype on a correlation scale. Subsequent vectors

necessarily explain sequentially diminishing amounts of variation (PC2 = 25.4% [12.6, 30.9]; PC3 = 12.1%

[7.4, 15.6]; PC4 = 3.7% [4.7, 9.9]; PC5 = 1.6% [0.8, 5.5]; PC6 > 0.001% [0,0]). All 6 cognitive performance

measures load with the same sign on PC1 and these loadings are statistically significant based on

bootstrapped confidence intervals not overlapping zero (Figure 3; Supplementary Information Table S2).

Thus, the eigen decomposition reiterates the view that the correlation structure is consistent with a

dominant axis of among-individual variation in cognitive phenotype caused by differences in underlying

g-model of general cognitive performance. On this axis, individuals at one end of the axis can be

considered to have 'higher/better' performance in all tasks (i.e. are faster and more accurate in the

association task, faster to solve the motor learning task, and faster and more accurate in the reversal

learning task when the cue is reversed), while individuals at the other end perform relatively poorly in

all respects.

Discussion

In this study we sought to determine whether – and to what extent – there exists among-individual

variation for cognitive performance in a captive population of recently wild-derived guppies. Adopting a

multivariate approach allowed us to test for covariation among multiple cognitive performance

measures, as well as between cognitive performance and a shy-bold type axis of personality variation.

Our study yields three main findings. First, there is among-individual variation underpinning all

cognitive performance measures. Second, there is strong among-individual correlation structure among

cognitive performance measures and bold type personality. Thirdly, the structure of **ID**-matrix provides

no support for the presence of important performance trade-offs, either between cognitive tasks, or

between speed and accuracy within learning tasks. Rather we suggest that the matrix is consistent with

a single axis of general variation that causes positively correlated performance across all cognitive tasks.

Individuals at one end of the axis can be considered to have relatively low 'general cognitive

performance' compared to those at the other end of the axis. The structure of **ID**-matrix is therefore

consistent with expectations under the *g*-model, although it is important to acknowledge that other

latent factors could exist that create similar patterns of covariation among observed behaviours. In what

follows, we discuss these findings in the context of understanding variation in, and evolution of, animal

cognition, while also drawing attention to important assumptions and caveats underpinning our

conclusions.

Among-individual variation was found in each of the cognitive performance measures as well as

in the measure of personality tested. That guppies show consistent among-individual differences in

'boldness', as measured here by time to emerge into the test compartment over the first set of 9 trials

(with a conditional repeatability of 43.9%) was expected from earlier studies of this population (e.g.

Houslay et al., 2019) and guppies more widely (Brown & Irving, 2014; O'Neill et al., 2018; Gasparini et

al., 2019). Indeed similar findings are common across a wide range of fish models used in behavioural

studies (Boulton et al., 2018; White et al., 2020). We also find among-individual variation in all measures

of cognitive performance across each task. Thus, we conclude that individuals differ in; ability to

discriminate between rewarded and unrewarded colours in an association learning task, ability to learn

a novel motor learning task, and, in cognitive flexibility as tested by a reversal learning task. The

estimated repeatabilities of our observed cognitive performance measures are moderate relative to

behavioural traits generally (Bell et al., 2009), but generally consistent with reports for cognitive assays

in animals (Cauchoix et al., 2018). It is worth noting that behavioural repeatabilities generally decline as

inter-observation period increases (Boulton et al., 2014) and here the inter-observation period was

short (usually less than 24hrs with up to 9 observations made in a 48hr period for some traits). Although

this is common for cognitive studies, we acknowledge that the stability of individual differences in

cognitive performance may well be less over longer timeframes.

Our multivariate modelling provides no support for trade-offs among cognitive performance

measures. This conclusion stems from the finding that the correlation structure in **ID** is universally

positive. Thus, for example, speed-accuracy trade-offs are not detected in our association learning trials.

Speed-accuracy trade-offs are widely predicted to arise where behavioural outcomes depend on

cognitive decision making (Briffa, 2013), although this prediction assumes that faster decisions are

better (if they come at no cost to accuracy). They have been detected in some animal studies, for

instance in Carib grackles (*Quiscalus lugubris*) where individuals that solve novel problems faster also

make more errors in discrimination learning tasks (Ducatez et al., 2019). Here we actually find the

opposite pattern, faster individuals are also more accurate, a result that seems counterintuitive but is

consistent with reports in song sparrows (*Melospiza melodia*; Boogert, 2011), and bumblebees (*Bombus*

terrestris; Raine, 2012)*.* We note that the correlation structure of behavioural **ID**-matrices can be

sensitive to environmental context (Houslay et al., 2018) as a consequence of IxE interactions (among-

individual differences in plasticity; Nussey et al., 2007). It is therefore possible trade-offs could be

apparent only under specific conditions and/or assays. This scenario is suggested by a recent study of

archerfish in which a speed-accuracy trade-off was only statistically significant under the more

challenging of two cognitive testing treatments applied (Jones et al., 2020). Plastic responses to social

context could also be important since, for instance, in less competitive scenarios (including the single

housing used here) slow decisions may carry minimal costs removing motivation to be fast (Sih & Del

Giudice, 2012).

More broadly, we also find positive correlations across cognitive performance variables

measured within different cognitive tasks. This is counter to predictions under hypothesised trade-off

models for maintaining variation. For instance, it has been hypothesised that individuals more reactive

to changes in environmental cues will perform relatively poorly in association learning but –being

cognitively flexible – will do well in reversal learning tasks (Sih & Del Giudice, 2012; Bebus et al., 2016).

Here we find a strong positive among-individual correlation between our measures of performance in

association and reversal learning (r_I=0.54). Again we note the absence of apparent trade-offs has

precedent in the animal cognition literature. For example, positive correlations have been reported

between discrimination acquisition and reversal learning in sparrows (Boogert et al., 2011), social

learning and innovative problem solving in pigeons (*Columba livia*; Bouchard et al., 2007), and motor

skill learning and reversal learning in pheasants (van Horik et al., 2018).

If positive correlations in the **ID**-matrix provide no support for trade-offs, then the corollary is

that they are consistent with a general axis of cognitive variation among-individuals in this population.

Thus, excluding the boldness proxy of emergence time and focusing on cognitive performance measures

only, we found that 57% of the variation in correlation matrix was explained by the leading principle

component (or eigen vector). By comparison, in tests of the general intelligence model of cognition

using human psychometric test batteries, an underlying *g*-factor is typically found to account for about

40–50% of the multivariate variation (Carroll, 1993; Deary, 2001). Under the *g*-model, trait performance

in cognitive assays across multiple domains are expected to load strongly (and in the same direction) on

a dominant first principle component (Plomin, 2001; Plomin & Spinath, 2002). We note that human

psychometric studies usually test a different and far wider range of cognitive abilities than examined

here. Nonetheless, here we find a similar pattern of strong positive loadings on a dominant first

principle component. Our results thus add to a small, but growing body of empirical literature finding

support for general axis of variation among-individuals in animal cognitive ability (for example see

Ashton et al., 2018; Shaw et al., 2015).

Our conclusion that (multivariate) cognitive variation in guppies is consistent with an

underlying general cognitive performance factor requires some caveats. These largely arise because

latent variables are, by definition, unmeasurable. Thus, all inferences about (latent) cognitive traits from

(observed) behavioural performance measures are inevitably assumption-laden. First, we are limited by

the set of variables assayed. Inclusion of multiple cognitive domains, different assays, and/or different

contexts, would provide more robust insight into the biological extent and evolutionary potential of

general intelligence (Poirier et al., 2020). Thus, for instance, we detect no evidence of cognitive trade-

offs within performance variable measured within each cognitive task, but they could occur among a

wider set of traits (or contexts) not examined. Here we used three cognitive tasks as proxies of three

cognitive domains: association learning; motor learning; and reversal learning, but did so using a single

testing paradigm in which performance measures share a requirement for individuals to discriminate

between two coloured cues. In this way, cognitive performance measures across all three tasks could be

said to be based on common associative processes, resulting ultimately in a predisposition for finding a

positive correlation structure among them. Second, the eigen vector decomposition provides only a

statistical description of the estimated correlation structure among observed variables; that **ID** is

consistent with an axis of variation in an underlying general cognitive performance factor does not

prove such a factor exists. Third, the positive correlation structure could be explained, at least in part, by

shared dependence of the observed traits on some other parameter that varies among individuals such

as metabolic state (Biro & Stamps, 2010; Mathot & Dall, 2013; McKenzie et al., 2016), satiety

(Shettleworth, 1972; Ben-Shahar & Robinson, 2001), or other drivers of motivation (van Horik &

Madden, 2016). Among-individual differences could result from differences in motivation rather than

differences in cognitive ability per se. For example, the increase in mean performance time in the

association task could be confounded by changes in motivation due to trial fatigue (reduced

motivation), at least for some individuals, even if some individuals clearly did perform better over time.

Experimental conditions can have important consequences for robust interpretation of studies targeting

cognitive traits (Rowe & Healy, 2014; Griffin et al., 2015; Boogert et al., 2018) and while we attempted

to minimise 'non-target' sources of among-individual variation (e.g. by standardising food rations,

housing and water chemistry) we cannot exclude the possibility that these contribute to the structure of

ID. Finally, we acknowledge the potential for bias in our findings caused by selective removal of

individuals due to our imposed learning criterion. Failure to participate in a cognitive task may result in

sampling biases when measuring among-individual variation in cognitive performances (Thornton &

Lukas, 2012; van Horik et al., 2017). Our learning criterion was specifically weak so that all included

individuals represented, as far as possible, among-individual variation of the initial population.

We also found that all six cognitive performance measures were positively correlated with

'boldness', measured here as an individual's tendency to consistently emerge rapidly from the home

compartment during trial observations. Pairwise correlations were nominally significant in four of the six

cases. Our finding that bolder individuals perform better in cognitive tasks mirrors results reported from

several other studies. For instance, boldness has been positively linked with performance speed in

associative learning tasks in rainbow trout (*Oncorhynchus mykiss*; Sneddon, 2003), shape discrimination

in Cavies (*Cavia aperea*; Guenther et al., 2014), and reversal learning in the Chimango

Caracara, (*Milvago chimango;* Guido et al., 2017). While shy-bold type behavioural variation can have

important fitness consequences (Wilson et al., 2010; Ariyomo & Watt, 2012; Ballew et al., 2017),

estimates of contemporary selection on boldness in wild guppies are lacking. Nonetheless, multiple

studies have reported among-population differences in boldness mapping to differences in predation, a

link that is consistent with the hypothesis that variable levels of risk could impose balancing selection

that maintains variation in boldness. Very speculatively, if this is true, and if correlations in **ID** are

recapitulated in **G,** the link between boldness and cognitive performance could help maintain variation

in the latter. This is because selection on any one trait (e.g. boldness) has consequences for the

evolutionary dynamics of genetically correlated traits (Roff, 2002). We do know that shy-bold type

personality variation is heritable in the population of guppies under lab conditions, (White & Wilson,

2019; White et al., 2019; Prentice et al., 2020) but not how selection is acting in the field.

In summary, we find evidence that individual guppies do differ in cognitive performance as

measured by a testing protocol that targets association learning, motor learning and cognitive flexibility

(reversal learning). We find no support for trade-offs occurring either between speed and accuracy (e.g.

within the association learning and reversal learning trials) or between overall performance across

cognitive tasks. Subject to the caveats highlighted above, the absence of trade-offs and the presence of

strong positive general correlations in **ID** mean that the general intelligence model provides a good

description of the structure of multivariate general cognitive performance variation in this species. Our

results also suggest that, to the extent that directional natural selection favours higher performance in

any single cognitive performance measure, it would lead to positively correlated evolutionary responses

across all other measures. However, this inference is dependent on **ID** being a reliable proxy for the **G**

matrix (Brommer, 2013). Greater efforts to quantify **G** is thus an obvious, if challenging, next step

towards understanding the maintenance of variation in and evolutionary dynamics of animal cognition.

References

Arden, R., & Adams, M. J. (2016). A general intelligence factor in dogs. *Intelligence*, *55*, 79–85.

https://doi.org/10.1016/j.intell.2016.01.008

Ariyomo, T. O., & Watt, P. J. (2012). The effect of variation in boldness and aggressiveness on the

reproductive success of zebrafish. *Animal Behaviour*, *83*(1), 41–46.

https://doi.org/10.1016/j.anbehav.2011.10.004

Ashton, B. J., Ridley, A. R., Edwards, E. K., & Thornton, A. (2018). Cognitive performance is linked to

group size and affects fitness in Australian magpies. *Nature*, *554*(7692), 364–367.

https://doi.org/10.1038/nature25503

Ballew, N. G., Mittelbach, G. G., & Scribner, K. T. (2017). Fitness consequences of boldness in Juvenile

and adult largemouth bass. *American Naturalist*, *189*(4), 396–406.

https://doi.org/10.1086/690909

Bebus, S. E., Small, T. W., Jones, B. C., Elderbrock, E. K., & Schoech, S. J. (2016). Associative learning is

inversely related to reversal learning and varies with nestling corticosterone exposure. *Animal*

Behaviour, *111*, 251–260. https://doi.org/10.1016/j.anbehav.2015.10.027

Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: a meta-analysis. In

Animal Behaviour (Vol. 77, Issue 4, pp. 771–783). https://doi.org/10.1016/j.anbehav.2008.12.022

Ben-Shahar, Y., & Robinson, G. E. (2001). Satiation differentially affects performance in a learning assay

by nurse and forager honey bees. *Journal of Comparative Physiology - A Sensory, Neural, and*

Behavioral Physiology, *187*(11), 891–899. https://doi.org/10.1007/s00359-001-0260-z

Biro, P. A., & Stamps, J. A. (2010). Do consistent individual differences in metabolic rate promote

consistent individual differences in behavior? In *Trends in Ecology and Evolution* (Vol. 25, Issue 11,

pp. 653–659). https://doi.org/10.1016/j.tree.2010.08.003

Bitterman, M. E. (1965). Phyletic differences in learning. *The American Psychologist*, *20*, 396–410.

https://doi.org/10.1037/h0022328

Boogert, N. J., Anderson, R. C., Peters, S., Searcy, W. A., & Nowicki, S. (2011). Song repertoire size in

male song sparrows correlates with detour reaching, but not with other cognitive measures.

Animal Behaviour, *81*(6), 1209–1216. https://doi.org/10.1016/j.anbehav.2011.03.004

Boogert, N. J., Madden, J. R., Morand-Ferron, J., & Thornton, A. (2018). Measuring and understanding

individual differences in cognition. In *Philosophical Transactions of the Royal Society B: Biological*

Sciences (Vol. 373, Issue 1756, p. 20170280). https://doi.org/10.1098/rstb.2017.0280

Bouchard, J., Goodyer, W., & Lefebvre, L. (2007). Social learning and innovation are positively correlated

in pigeons (Columba livia). *Animal Cognition*, *10*(2), 259–266. https://doi.org/10.1007/s10071-

006-0064-1

Boulton, K., Grimmer, A. J., Rosenthal, G. G., Walling, C. A., & Wilson, A. J. (2014). How stable are

personalities? A multivariate view of behavioural variation over long and short timescales in the

sheepshead swordtail, Xiphophorus birchmanni. *Behavioral Ecology and Sociobiology*, *68*(5), 791–

803. https://doi.org/10.1007/s00265-014-1692-0

Boulton, K., Walling, C. A., Grimmer, A. J., Rosenthal, G. G., & Wilson, A. J. (2018). Phenotypic and

genetic integration of personality and growth under competition in the sheepshead swordtail,

Xiphophorus birchmanni. *Evolution*, *72*(1), 187–201. https://doi.org/10.1111/evo.13398

Brandão, M. L., Fernandes, A. M. T. de A., & Gonçalves-de-Freitas, E. (2019). Male and female cichlid fish

show cognitive inhibitory control ability. *Scientific Reports*, *9*(1), 1–8.

https://doi.org/10.1038/s41598-019-52384-2

Briffa, M. (2013). Plastic proteans: Reduced predictability in the face of predation risk in hermit crabs.

Biology Letters, *9*(5), 20130592. https://doi.org/10.1098/rsbl.2013.0592

Brommer, J. E. (2013). On between-individual and residual (co)variances in the study of animal

personality: Are you willing to take the "individual gambit"? In *Behavioral Ecology and*

Sociobiology (Vol. 67, Issue 6, pp. 1027–1032). https://doi.org/10.1007/s00265-013-1527-4

Brown, C., & Irving, E. (2014). Individual personality traits influence group exploration in a feral guppy

population. *Behavioral Ecology*, *25*(1), 95–101. https://doi.org/10.1093/beheco/art090

Brydges, N. M., & Braithwaite, V. A. (2009). Does environmental enrichment affect the behaviour of fish

commonly used in laboratory work? *Applied Animal Behaviour Science*, *118*(3–4), 137–143.

https://doi.org/10.1016/j.applanim.2009.02.017

Buchanan, K., Perera, T. B. de, … C. C.-A., & 2012, U. (2020). Guidelines for the treatment of animals in

behavioural research and teaching. *Animal Behaviour*, *159*, I–XI.

https://doi.org/10.1016/j.anbehav.2019.11.002

Buechel, S. D., Boussard, A., Kotrschal, A., van Der Bijl, W., & Kolm, N. (2018). Brain size affects

performance in a reversal-learning test. *Proceedings of the Royal Society B: Biological Sciences*,

285(1871). https://doi.org/10.1098/rspb.2017.2031

Burkart, J. M., Schubiger, M. N., & Van Schaik, C. P. (2017). The evolution of general intelligence.

Behavioral and Brain Sciences, *40*, e195. https://doi.org/10.1017/S0140525X16000959

Butler, D. G., Cullis, B. R., Gilmour, A. R., Gogel, B. J., & Thompson, R. (2018). ASReml-R Reference

Manual Version 4 ASReml estimates variance components under a general linear mixed model by

residual maximum likelihood (REML). In *asreml.kb.vsni.co.uk*.

http://www.homepages.ed.ac.uk/iwhite/asreml/uop.

Carroll, J. (1993). *Human cognitive abilities: A survey of factor-analytic studies*. Cambridge University

Press.

Cauchoix, M., Chow, P. K. Y., Van Horik, J. O., Atance, C. M., Barbeau, E. J., Barragan-Jason, G., Bize, P.,

Boussard, A., Buechel, S. D., Cabirol, A., Cauchard, L., Claidière, N., Dalesman, S., Devaud, J. M.,

Didic, M., Doligez, B., Fagot, J., Fichtel, C., Henke-Von Der Malsburg, J., … Morand-Ferron, J.

(2018). The repeatability of cognitive performance: A meta-analysis. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, *373*(1756). https://doi.org/10.1098/rstb.2017.0281

Cheverud, J. M. (1988). A comparison of genetic and phenotypic correlations. *Evolution*, *42*(5), 958–968.

https://doi.org/10.1111/j.1558-5646.1988.tb02514.x

Chittka, L., Skorupski, P., & Raine, N. E. (2009). Speed-accuracy tradeoffs in animal decision making. In

Trends in Ecology and Evolution (Vol. 24, Issue 7, pp. 400–407).

https://doi.org/10.1016/j.tree.2009.02.010

Deary, I. J. (2001). Human intelligence differences: A recent history. In *Trends in Cognitive Sciences* (Vol.

5, Issue 3, pp. 127–130). Elsevier. https://doi.org/10.1016/S1364-6613(00)01621-1

Deary, I. J., Penke, L., & Johnson, W. (2010). The neuroscience of human intelligence differences. *Nat*

Rev Neurosci, *11*(3), 201–211. https://doi.org/10.1038/nrn2793

Del Giudice, M., & Crespi, B. J. (2018). Basic functional trade-offs in cognition: An integrative framework.

Cognition, *179*, 56–70. https://doi.org/10.1016/j.cognition.2018.06.008

Dougherty, L. R., & Guillette, L. M. (2018). Linking personality and cognition: A meta-analysis. In

Philosophical Transactions of the Royal Society B: Biological Sciences (Vol. 373, Issue 1756, p.

20170282). https://doi.org/10.1098/rstb.2017.0282

Ducatez, S., Audet, J. N., & Lefebvre, L. (2019). Speed–accuracy trade-off, detour reaching and response

to PHA in Carib grackles. *Animal Cognition*, *22*(5), 625–633. https://doi.org/10.1007/s10071-019-

01258-1

Fong, S., Buechel, S. D., Boussard, A., Kotrschal, A., & Kolm, N. (2019). Plastic changes in brain

morphology in relation to learning and environmental enrichment in the guppy (Poecilia

reticulata). *Journal of Experimental Biology*, *222*(10). https://doi.org/10.1242/jeb.200402

Friard, O., & Gamba, M. (2016). BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. *Methods in Ecology and Evolution*, *7*(11), 1325–1330.

https://doi.org/10.1111/2041-210X.12584

Gasparini, C., Speechley, E. M., & Polverino, G. (2019). The bold and the sperm: Positive association

between boldness and sperm number in the guppy. *Royal Society Open Science*, *6*(7).

https://doi.org/10.1098/rsos.190474

Gilmour, A. R., Gogel, B. J., Cullis, B. R., Thompson, R., & Butler, D. (2009). ASReml user guide release

3.0. *VSN International Ltd, Hemel Hempstead, UK*.

Grafen A. (1984). *Natural selection, kin selection and group selection. In Behavioural Ecology: An

Evolutionary Approach.* (Krebs J.R & Davies N.B (Eds.); 2nd ed.). Blackwell Scientific, Oxford.

https://scholar.google.co.uk/scholar?hl=en&as_sdt=0%2C5&q=Natural+selection%2C+kin+selecti

on+and+group+selection.+&btnG=

Griffin, A. S., Guillette, L. M., & Healy, S. D. (2015). Cognition and personality: An analysis of an emerging

field. In *Trends in Ecology and Evolution* (Vol. 30, Issue 4, pp. 207–214).

https://doi.org/10.1016/j.tree.2015.01.012

Guenther, A., & Brust, V. (2017). Individual consistency in multiple cognitive performance: behavioural

versus cognitive syndromes. *Animal Behaviour*, *130*, 119–131.

https://doi.org/10.1016/j.anbehav.2017.06.011

Guenther, A., Brust, V., Dersen, M., & Trillmich, F. (2014). Learning and personality types are related in

cavies (cavia aperea). *Journal of Comparative Psychology*, *128*(1), 74–81.

https://doi.org/10.1037/a0033678

Guez, D., & Griffin, A. S. (2016). Unraveling the key to innovative problem solving: A test of learning

versus persistence. *Behavioral Ecology*, *27*(5), 1449–1460.

https://doi.org/10.1093/beheco/arw055

Guido, J. M., Biondi, L. M., Vasallo, A. I., & Muzio, R. N. (2017). Neophobia is negatively related to

reversal learning ability in females of a generalist bird of prey, the Chimango Caracara, Milvago

chimango. *Animal Cognition*, *20*(4), 591–602. https://doi.org/10.1007/s10071-017-1083-9

Healy, S. D., & Rowe, C. (2010). Information Processing: The Ecology and Evolution of Cognitive Abilities.

In *Evolutionary Behavioral Ecology* (pp. 162–174). https://eprints.ncl.ac.uk/139823

Hopkins, W. D., Russell, J. L., & Schaeffer, J. (2014). Chimpanzee intelligence is heritable. *Current Biology*,

24(14), 1649–1652. https://doi.org/10.1016/j.cub.2014.05.076

Houslay, T. M., Earley, R. L., White, S. J., Lammers, W., Grimmer, A. J., Travers, L. M., Johnson, E. L.,

Young, A. J., & Wilson, A. J. (2019). Genetic integration of the stress response. *BioRxiv*, 770586.

https://doi.org/10.1101/770586

Houslay, T. M., Vierbuchen, M., Grimmer, A. J., Young, A. J., & Wilson, A. J. (2018). Testing the stability of

behavioural coping style across stress contexts in the Trinidadian guppy. *Functional Ecology*, *32*(2),

424–438. https://doi.org/10.1111/1365-2435.12981

Jones, A. G., Arnold, S. J., & Bürger, R. (2004). Evolution and stability of the G-matrix on a landscape with

a moving optimum. *Evolution*, *58*(8), 1639–1654. https://doi.org/10.1111/j.0014-

3820.2004.tb00450.x

Jones, N., Webster, M., Newport, C., Templeton, C. N., Schuster, S., & Rendell, L. (2020). Cognitive styles:

speed–accuracy trade-offs underlie individual differences in archerfish. *Animal Behaviour*, *160*, 1–

14. https://doi.org/10.1016/j.anbehav.2019.11.019

Kehagia, A. A., Murray, G. K., & Robbins, T. W. (2010). Learning and cognitive flexibility: Frontostriatal

function and monoaminergic modulation. In *Current Opinion in Neurobiology* (Vol. 20, Issue 2, pp.

199–204). https://doi.org/10.1016/j.conb.2010.01.007

Kniel, N., Guenther, A., & Godin, J. G. J. (2020). Individual personality does not predict learning

performance in a foraging context in female guppies, Poecilia reticulata. *Animal Behaviour*, *167*,

3–12. https://doi.org/10.1016/j.anbehav.2020.07.007

Laland, K. N., & Reader, S. M. (1999). Foraging innovation in the guppy. *Animal Behaviour*, *57*(2), 331–

340. https://doi.org/10.1006/anbe.1998.0967

Lande, R. (1979). Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain: Body Size

Allometry. *Evolution*, *33*(1), 402. https://doi.org/10.2307/2407630

Langley, E. J. G., Adams, G., Beardsworth, C. E., Dawson, D. A., Laker, P. R., van Horik, J. O., Whiteside,

M. A., Wilson, A. J., & Madden, J. R. (2020). Heritability and correlations among learning and

inhibitory control traits. *Behavioral Ecology*. https://doi.org/10.1093/beheco/araa029

Li, L., MaBouDi, H. Di, Egertová, M., Elphick, M. R., Chittka, L., & Perry, C. J. (2017). A possible structural

correlate of learning performance on a colour discrimination task in the brain of the bumblebee.

Proceedings of the Royal Society B: Biological Sciences, *284*(1864).

https://doi.org/10.1098/rspb.2017.1323

Lucon-Xiccato, T., & Bisazza, A. (2014). Discrimination reversal learning reveals greater female

behavioural flexibility in guppies. *Biology Letters*, *10*(6), 20140206.

Lucon-Xiccato, T., & Bisazza, A. (2016). Male and female guppies differ in speed but not in accuracy in

visual discrimination learning. *Animal Cognition*, *19*(4), 733–744. https://doi.org/10.1007/s10071-

016-0969-2

Lucon-Xiccato, T., & Bisazza, A. (2017). Individual differences in cognition among teleost fishes. In

Behavioural Processes (Vol. 141, pp. 184–195). https://doi.org/10.1016/j.beproc.2017.01.015

Lucon-Xiccato, T., Dadda, M., & Bisazza, A. (2016). Sex Differences in Discrimination of Shoal Size in the

Guppy (Poecilia reticulata). *Ethology*, *122*(6), 481–491. https://doi.org/10.1111/eth.12498

Lucon-Xiccato, T., Gatto, E., & Bisazza, A. (2017). Fish perform like mammals and birds in inhibitory

motor control tasks. *Scientific Reports*, *7*, 13144. https://doi.org/10.1101/188359

Mamuneas, D., Spence, A. J., Manica, A., & King, A. J. (2015). Bolder stickleback fish make faster

decisions, but they are not less accurate. *Behavioral Ecology*, *26*(1), 91–96.

https://doi.org/10.1093/beheco/aru160

Mathot, K. J., & Dall, S. R. X. (2013). Metabolic rates can drive individual differences in information and

754 insurance use under the risk of starvation. *American Naturalist*, *182*(5), 611–620.

755 https://doi.org/10.1086/673300

756 Mazza, V., Jacob, J., Dammhahn, M., Zaccaroni, M., & Eccard, J. A. (2019). Individual variation in

757 cognitive style reflects foraging and anti-predator strategies in a small mammal. *Scientific Reports*,

758 *9*(1). https://doi.org/10.1038/s41598-019-46582-1

759 McKenzie, D. J., Belaõ, T. C., Killen, S. S., & Rantin, F. T. (2016). To boldly gulp: Standard metabolic rate

760 and boldness have context-dependent influences on risk-taking to breathe air in a catfish (Journal

761 of Experimental Biology (2016) 218 (3762-3770)). In *Journal of Experimental Biology* (Vol. 219,

762 Issue 5, p. 767). https://doi.org/10.1242/jeb.138842

763 Miletto Petrazzini, M. E., & Agrillo, C. (2016). Turning to the larger shoal: are there individual differences

764 in small- and large-quantity discrimination of guppies? *Ethology Ecology and Evolution*, *28*(2), 211–

765 220. https://doi.org/10.1080/03949370.2015.1029011

766 Moiron, M., Mathot, K. J., & Dingemanse, N. J. (2016). A multi-level approach to quantify speed-

767 accuracy trade-offs in great tits (Parus major). *Behavioral Ecology*, *27*(5), 1539–1546.

768 https://doi.org/10.1093/beheco/arw077

769 Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical

770 guide for biologists. In *Biological Reviews* (Vol. 85, Issue 4, pp. 935–956).

771 https://doi.org/10.1111/j.1469-185X.2010.00141.x

772 Nawroth, C., Prentice, P. M., & McElligott, A. G. (2017). Individual personality differences in goats

773 predict their performance in visual learning and non-associative cognitive tasks. In *Behavioural*

774 *Processes* (Vol. 134, pp. 43–53). https://doi.org/10.1016/j.beproc.2016.08.001

775 Nisbett, R. E., Aronson, J., Blair, C., Dickens, W., Flynn, J., Halpern, D. F., & Turkheimer, E. (2012).

776 Intelligence: New Findings and Theoretical Developments. *American Psychologist*, *67*(2), 130–159.

777 https://doi.org/10.1037/a0026699

778 Nussey, D. H., Wilson, A. J., & Brommer, J. E. (2007). The evolutionary ecology of individual phenotypic

779 plasticity in wild populations. *Journal of Evolutionary Biology*, *20*(3), 831–844.

780 https://doi.org/10.1111/j.1420-9101.2007.01300.x

781 O'Neill, S. J., Williamson, J. E., Tosetto, L., & Brown, C. (2018). Effects of acclimatisation on behavioural

repeatability in two behaviour assays of the guppy Poecilia reticulata. *Behavioral Ecology and*

Sociobiology, *72*(10). https://doi.org/10.1007/s00265-018-2582-7

Plomin, R. (2001). The genetics of g in human and mouse. *Nature Reviews Neuroscience*.

http://www.nature.com/nrn/journal/v2/n2/abs/nrn0201_136a.html

Plomin, R., & Spinath, F. M. (2002). Genetics and general cognitive ability (g). In *Trends in Cognitive*

Sciences (Vol. 6, Issue 4, pp. 169–176). https://doi.org/10.1016/S1364-6613(00)01853-2

Poirier, M.-A., Kozlovsky, D. Y., Morand-Ferron, J., & Careau, V. (2020). How general is cognitive ability in

non-human animals? A meta-analytical and multi-level reanalysis approach. *Proceedings of the*

Royal Society B: Biological Sciences, *287*(1940), 20201853.

https://doi.org/10.1098/rspb.2020.1853

Prentice, P. M., Houslay, T. M., Martin, J. G. A., & Wilson, A. J. (2020). Genetic variance for behavioural

'predictability' of stress response. *Journal of Evolutionary Biology*, *33*(5), 642–652.

https://doi.org/10.1111/jeb.13601

R Development Core Team. (2008). *Control*, *55*, 275–286. https://stat.ethz.ch/pipermail/r-help/2008-

May/161481.html

Raine, N. E., & Chittka, L. (2012). No Trade-Off between Learning Speed and Associative Flexibility in

Bumblebees: A Reversal Learning Test with Multiple Colonies. *PLoS ONE*, *7*(9).

https://doi.org/10.1371/journal.pone.0045096

Roff, D. A. (2002). *Life history evolution* (p. 527). Sinauer Associates.

Rowe, C., & Healy, S. D. (2014). Measuring variation in cognition. In *Behavioral Ecology* (Vol. 25, Issue 6,

pp. 1287–1292). https://doi.org/10.1093/beheco/aru090

Schielzeth, H., Dingemanse, N. J., Nakagawa, S., Westneat, D. F., Allegue, H., Teplitsky, C., Réale, D.,

Dochtermann, N. A., Garamszegi, L. Z., & Araya-Ajoy, Y. G. (2020). Robustness of linear mixed-

effects models to violations of distributional assumptions. *Methods in Ecology and Evolution*,

11(9), 1141–1152. https://doi.org/10.1111/2041-210X.13434

Shaw, R. C., Boogert, N. J., Clayton, N. S., & Burns, K. C. (2015). Wild psychometrics: Evidence for

"general" cognitive performance in wild New Zealand robins, Petroica longipes. *Animal Behaviour*,

109, 101–111. https://doi.org/10.1016/j.anbehav.2015.08.001

Shaw, R. C., & Schmelz, M. (2017). Cognitive test batteries in animal cognition research: evaluating the

past, present and future of comparative psychometrics. In *Animal Cognition* (Vol. 20, Issue 6, pp.

1003–1018). https://doi.org/10.1007/s10071-017-1135-1

Shettleworth, S. J. (1972). Constraints on Learning. *Advances in the Study of Behavior*, *4*(C), 1–68.

https://doi.org/10.1016/S0065-3454(08)60006-6

Shettleworth, S. J. (2009). The evolution of comparative cognition: Is the snark still a boojum?

Behavioural Processes, *80*(3), 210–217. https://doi.org/10.1016/j.beproc.2008.09.001

Shettleworth, S. J. (2010). Getting Around: Spatial Cognition. In *Cognition, Evolution, and Behavior*

(second, pp. 261–312). Oxford University Press.

Shettleworth, S. J. (2012). Modularity, comparative cognition and human uniqueness. In *Philosophical*

Transactions of the Royal Society B: Biological Sciences (Vol. 367, Issue 1603, pp. 2794–2802).

https://doi.org/10.1098/rstb.2012.0211

Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioral syndromes: an ecological and evolutionary overview.

Trends in Ecology & Evolution, *19*(7), 372–378.

Sih, A., & Del Giudice, M. (2012). Linking behavioural syndromes and cognition: a behavioural ecology

perspective. *Philosophical Transactions of the Royal Society B-Biological Sciences*, *367*(1603),

2762–2772. https://doi.org/10.1098/rstb.2012.0216

Sneddon, L. U. (2003). The bold and the shy: Individual differences in rainbow trout. *Journal of Fish*

Biology, *62*(4), 971–975. https://doi.org/10.1046/j.1095-8649.2003.00084.x

Sonnenberg, B. R., Branch, C. L., Pitera, A. M., Bridge, E., & Pravosudov, V. V. (2019). Natural Selection

and Spatial Cognition in Wild Food-Caching Mountain Chickadees. *Current Biology*, *29*(4), 670-

676.e3. https://doi.org/10.1016/j.cub.2019.01.006

Stearns, S. C. (1992). *The Evolution of Life Histories*. Oxford University Press.

Thornton, A., & Lukas, D. (2012). Individual variation in cognitive performance: Developmental and

evolutionary perspectives. In *Philosophical Transactions of the Royal Society B: Biological Sciences*

(Vol. 367, Issue 1603, pp. 2773–2783). The Royal Society. https://doi.org/10.1098/rstb.2012.0214

van Horik, J. O., & Emery, N. J. (2018). Serial reversal learning and cognitive flexibility in two species of

Neotropical parrots (Diopsittaca nobilis and Pionites melanocephala). *Behavioural Processes*, *157*,

664–672. https://doi.org/10.1016/j.beproc.2018.04.002

van Horik, J. O., Langley, E. J. G., Whiteside, M. A., & Madden, J. R. (2017). Differential participation in

cognitive tests is driven by personality, sex, body condition and experience. In *Behavioural*

Processes (Vol. 134, pp. 22–30). https://doi.org/10.1016/j.beproc.2016.07.001

van Horik, J. O., & Madden, J. R. (2016). A problem with problem solving: Motivational traits, but not

cognition, predict success on novel operant foraging tasks. *Animal Behaviour*, *114*, 189–198.

https://doi.org/10.1016/j.anbehav.2016.02.006

Visscher, P. M. (2006). A note on the asymptotic distribution of likelihood ratio tests to test variance

components. *Twin Research and Human Genetics*, *9*(4), 490–495.

https://doi.org/10.1375/183242706778024928

Wallace, K. J., & Hofmann, H. A. (2021). Equal performance but distinct behaviors: sex differences in a

novel object recognition task and spatial maze in a highly social cichlid fish. *Animal Cognition*, *1*, 3.

https://doi.org/10.1007/s10071-021-01498-0

Walsh, B., & Blows, M. W. (2009). Abundant genetic variation + strong selection = multivariate genetic

constraints: A geometric view of adaptation. *Annual Review of Ecology, Evolution, and*

Systematics, *40*, 41–59. https://doi.org/10.1146/annurev.ecolsys.110308.120232

Wang, M. Y., Brennan, C. H., Lachlan, R. F., & Chittka, L. (2015). Speed-accuracy trade-offs and

individually consistent decision making by individuals and dyads of zebrafish in a colour

discrimination task. *Animal Behaviour*, *103*, 277–283.

https://doi.org/10.1016/j.anbehav.2015.01.022

Wasserman, E. A., & Zentall, T. R. (2006). Comparative Cognition: Experimental Explorations of Animal

Intelligence. In *Comparative Cognition: Experimental Explorations of Animal Intelligence*.

https://doi.org/10.1093/acprof:oso/9780195377804.001.0001

White, S. J., Houslay, T. M., & Wilson, A. J. (2019). Evolutionary genetics of personality in the Trinidadian

guppy II: sexual dimorphism and genotype-by-sex interactions. *Heredity*, *122*(1), 15–28.

https://doi.org/10.1038/s41437-018-0083-0

White, S. J., Kells, T., & Wilson, A. J. (2016). Metabolism, personality and pace of life in the Trinidadian

guppy, Poecilia reticulata. *Behaviour*, *153*(13–14), 1517–1543. https://doi.org/10.1163/1568539X-

00003375

White, S. J., Pascall, D., & Wilson, A. J. (2020). Towards a comparative approach to the structure of

animal personality variation. *Behavioral Ecology*, *31*(2), 340–351.

https://doi.org/10.1093/beheco/arz198

White, S. J., & Wilson, A. J. (2019). Evolutionary genetics of personality in the Trinidadian guppy I:

maternal and additive genetic effects across ontogeny. *Heredity*, *122*(1), 1–14.

https://doi.org/10.1038/s41437-018-0082-1

White, S. L., Wagner, T., Gowan, C., & Braithwaite, V. A. (2017). Can personality predict individual

differences in brook trout spatial learning ability? *Behavioural Processes*, *141*, 220–228.

https://doi.org/10.1016/j.beproc.2016.08.009

Wilson, A. D. M., Godin, J. G. J., & Ward, A. J. W. (2010). Boldness and reproductive fitness correlates in

the eastern mosquitofish, gambusia holbrooki. *Ethology*, *116*(1), 96–104.

https://doi.org/10.1111/j.1439-0310.2009.01719.x

Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., Kruuk, L. E. . B., &

Nussey, D. H. (2010). An ecologists guide to the animal model. In *J Anim Ecol* (Vol. 79, Issue 1, pp.

13–26). http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2656.2009.01639.x/full

Wolf, M., Van Doorn, G. S., Leimar, O., & Weissing, F. J. (2007). Life-history trade-offs favour the

evolution of animal personalities. *Nature*, *447*(7144), 581–584.

 Figure Legends

Figure 1. Aerial view of the tank used to house individual fish throughout the experiment. The home

compartment allowed the focal fish visual access to neighbouring tanks and included foliage plants for

cover and enrichment. Also shown in the experimental compartment is the test plate used during each

task.

Figure 2. Plots of raw data of emergence data and cognitive performance measures. Each plot header

outlines the set numbers used to assay each variable; a) emergence time (*emergence)*; association

learning task b) decision speed (AL_{speed}) and c) accuracy ($AL_{accuracy}$), d) association learning task

performance time *(AL_{time})*; e) $_{motor}$ learning task performance time (ML_{time}); f) reversal learning task

performance time (RL_{time}) and g) accuracy ($RL_{accuracy}$). For continuous variables ($AL_{time,}$ AL_{speed}, ML_{time} and

RL_{time}), higher values (seconds) represent longer durations to locate the food reward in each task, and

lower values represent faster durations to solve the task. For binary variables ($AL_{accuracy}$ and $RL_{accuracy}$),

accuracy is represented as the percentage of correct choices made in each trial. Error bars represent

mean and standard errors of performance time for individuals. Blue line represents regression line

through mean performance or accuracy at each trial.

Figure 3: Variable loadings from the first eigen vector (principal component, PC1) of the **ID**-matrix. This

axis explains 57.1% of the among-individual (co)variation found in the components of the cognitive

phenotype in our guppy population. Points show variable loadings from the first eigen vector of our

estimate of **ID**, with bars representing 95% confidence intervals on each loading (calculated from 5000

bootstrapped replicates of the model)

 Table legends and Tables

910 **Table 1**. Outline of behaviours used to assess performance within each measure of personality and

911 cognitive performance, including details of corresponding set number at which each variable was

912 measured and corresponding acronyms used within the text.

Set Number/ Number of informative trials	Personality trait & Cognitive performance task / Variable measured	Description of variable measured	Acronym
Set 1, 9 trials	Boldness	Time to emerge (Time taken to emerge into the test compartment once the compartment door is opened)	*emergence*
	Association learning task		
Set 1, 9 trials	Decision speed	(After emergence into test compartment, time taken to make first choice, measured as swimming within 1 body length of, and actively exploring correct counter.)	AL_{speed}
Set 1, 9 trials	Decision accuracy	(Accuracy of first choice. Either correct or incorrect)	$AL_{accuracy}$
Set 2 & 3, 18 trials	Time to eat	(After emergence into test compartment, time taken to eat reward from the correct well)	AL_{time}
	Motor learning task		
Set 4 & 5, 18 trials	Time to eat	(After correct choice made (i.e. swimming within 1 body length of, and actively exploring correct counter), time taken to dislodge counter and eat the food reward)	ML_{time}
	Reversal learning task		
Set 6 & 7, 18 trials	Time to eat	(After emergence into test compartment, time taken to dislodge disc and eat reward from the correct well)	RL_{time}
Set 6 & 7, 18 trials	Decision accuracy	(Accuracy of first choice. Either correct or incorrect)	$RL_{accuracy}$

915 **Table 2:** Estimated among-individual variance (V_I) and adjusted repeatability (R) from univariate models

916 of all cognitive performance and personality measure. Standard errors are shown in parentheses. Also

917 shown are likelihood ratio tests of V_I.

Task	Variable	V_I	R	$X^2_{0,1}$	P
Associative Learning	*Emergence*	0.415 (0.077)	0.439 (0.049)	206.004	<0.001
	AL_{speed}	0.442 (0.081)	0.460 (0.049)	220.393	<0.001
	$AL_{accuracy}$	0.202 (0.045)	0.264 (0.046)	79.989	<0.001
	AL_{time}	0.341 (0.074)	0.335 (0.050)	186.345	<0.001
Motor Learning	ML_{time}	0.275 (0.065)	0.329 (0.054)	171.462	<0.001
Reversal Learning	RL_{time}	0.372 (0.088)	0.375 (0.058)	178.426	<0.001
	$RL_{accuracy}$	0.182 (0.050)	0.203 (0.047)	65.327	<0.001

919 **Table 3. ID** variance–covariance–correlation matrix from the multivariate mixed model. Estimated variances are shown on the diagonal (dark grey shading), with

920 correlations above and covariances below. Standard errors are shown in parentheses, and bold font denotes nominally significant estimates assuming approximate

921 95% CI of ± 1.96SE.

ID Matrix	emergence	AL_{speed}	$AL_{accuracy}$	AL_{time}	ML_{time}	RL_{time}	$RL_{accuracy}$
emergence	**0.426 (0.0773)**	**0.818 (0.046)**	**0.788 (0.062)**	**0.663 (0.095)**	0.239 (0.164)	0.166 (0.176)	**0.412 (0.171)**
AL_{speed}	**0.361 (0.072)**	**0.458 (0.082)**	**0.894 (0.039)**	**0.758 (0.074)**	0.271 (0.155)	0.228 (0.169)	**0.397 (0.168)**
$AL_{accuracy}$	**0.242 (0.052)**	**0.285 (0.056)**	**0.221 (0.046)**	**0.729 (0.092)**	0.124 (0.175)	0.090 (0.184)	0.258 (0.190)
AL_{time}	**0.286 (0.072)**	**0.340 (0.076)**	**0.227 (0.055)**	**0.437 (0.092)**	**0.747 (0.082)**	**0.540 (0.129)**	**0.423 (0.161)**
ML_{time}	0.092 (0.067)	0.108 (0.067)	0.034 (0.049)	**0.291 (0.075)**	**0.346 (0.078)**	**0.817 (0.067)**	**0.340 (0.167)**
RL_{time}	0.074 (0.082)	0.106 (0.083)	0.029 (0.060)	**0.246 (0.085)**	**0.331 (0.083)**	**0.474 (0.112)**	**0.473 (0.144)**
$RL_{accuracy}$	**0.122 (0.060)**	0.122 (0.061)	0.055 (0.043)	**0.127 (0.059)**	0.091 (0.051)	**0.148 (0.062)**	**0.205 (0.056)**

Figure 1. Aerial view of the tank used to house individual fish throughout the experiment. The home

compartment allowed the focal fish visual access to neighbouring tanks and included foliage plants for

cover and enrichment. Also shown in the experimental compartment is the test plate used during each

task.

Figure 2a:

Figure 2b: **Figure 2c**:

Figure 2d: **Figure 2e**:

Figure 2f: **Figure 2g**:

Figure 2. Plots of raw data of emergence data and cognitive performance measures. Each plot header

outlines the set numbers used to assay each variable; a) emergence time (*emergence)*; association

learning task b) decision speed (AL_{speed}) and c) accuracy ($AL_{accuracy}$), d) association learning task

performance time *(AL_{time})*; e) *motor* learning task performance time (ML_{time}); f) reversal learning task

performance time (RL_{time}) and g) accuracy ($RL_{accuracy}$). For continuous variables ($AL_{time,}$ AL_{speed}, ML_{time} and

RL_{time}), higher values (seconds) represent longer durations to locate the food reward in each task, and

lower values represent faster durations to solve the task. For binary variables ($AL_{accuracy}$ and $RL_{accuracy}$),

accuracy is represented as the percentage of correct choices made in each trial. Error bars represent

mean and standard errors of performance time for individuals. Blue line represents regression line

through mean performance or accuracy at each trial.

Figure 3: Variable loadings from the first eigen vector (principal component, PC1) of the **ID**-matrix. This

axis explains 57.1% of the among-individual (co)variation found in the components of the cognitive

phenotype in our guppy population. Points show variable loadings from the first eigen vector of our

estimate of **ID**, with bars representing 95% confidence intervals on each loading (calculated from 5000

bootstrapped replicates of the model)