SHIP164 is a Chorein Motif Containing Lipid Transport Protein that Controls Membrane Dynamics and Traffic at the Endosome-Golgi Interface.

Michael G. Hanna[1-4]*, Patreece Suen[2]*, Yumei Wu[1-4], Karin M. Reinisch[2,6], Pietro De Camilli[1-6]

[1]Department of Neuroscience, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

[2]Department of Cell Biology, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

[3]Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

[4]Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

[5]Kavli Institue for Neuroscience, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

[6]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD.

* M. Hanna and P. Suen contributed equally to this paper.

Correspondence to Pietro De Camilli: pietro.decamilli@yale.edu and Karin M. Reinisch: karin.reinisch@yale.edu

ABSTRACT
Cellular membranes differ in protein and lipid composition as well as in the protein-lipid ratio.
Thus, progression of membranous organelles along traffic routes requires mechanisms to control
bilayer lipid chemistry and their abundance relative to proteins. The recent structural and
functional characterization of VPS13-family proteins has suggested a mechanism through which
lipids can be transferred in bulk from one membrane to another at membrane contact sites, and
thus independently of vesicular traffic. Here we show that SHIP164 (UHRF1BP1L) shares
structural and lipid transfer properties with these proteins and is localized on a subpopulation of
vesicle clusters in the early endocytic pathway whose membrane cargo includes the cation-
independent mannose-6-phosphate receptor (MPR) and ATG9. Loss of SHIP164 disrupts
retrograde traffic of these organelles to the Golgi complex. Our findings raise the possibility that
bulk transfer of lipids to endocytic membranes may play a role in their traffic.
42

INTRODUCTION

The homeostasis of intracellular membranes and their adaptation to changes in the functional state of the cell requires the coordination of protein and lipid transport. While much has been learned about protein traffic, less is known about the dynamics and transport of bilayer lipids. A significant fraction of such lipids move between organelles as part of the membranes of vesicular carriers. However, lipids also move via transport proteins that harbor them in hydrophobic cavities as they travel through the aqueous environment of the cytosol. This mode of transport has been known for decades, but has been increasingly appreciated over the last several years with the discovery of many new lipid transport proteins. Moreover, it has also become clear that many such proteins function at membrane contact sites, thus facilitating specificity and speed of lipid transport. Typically, these proteins contain modules or motifs that tether them to the two apposed membranes while lipid transfer modules move back and forth between them to extract and deliver lipids by a shuttling mechanism[1–5].

Recently, the characterization of VPS13 and its distant paralogue ATG2 has suggested a new mode of transport involving the flow of lipids along a protein bridge that connects the two membranes in eukaryotic cells[6–10]. A defining feature of these proteins is the presence of a conserved N-terminal region, ~125 residues long, referred to as the chorein-N motif[6,11–13]. In VPS13 and ATG2, this motif caps one end of an elongated rod (extended chorein domain). The rod comprises an extended ß-sheet that is highly curved to resemble a taco shell harboring a groove along its length[7,8]. A hydrophobic cavity in the chorein motif is continuous with the groove, whose floor is lined by hydrophobic amino acids, and so suited to accommodate many lipids at once and to allow their flow from one end of the rod to the other. In ATG2, the rod represents the bulk of the protein, while in VPS13 there are additional folded C-terminal domains that function in localization. Lipids are thought to flow unidirectionally along the rod, producing a net flow of lipids to the acceptor membrane and allowing for its expansion independent of contribution of new membrane lipids by vesicle fusion[9]. Accordingly, both the VPS13 paralogues and ATG2 have been implicated in membrane growth[14–17]. Many questions remain, however, about the precise mechanisms of action of these proteins. In some cases, the earliest stages of membrane growth mediated by VPS13 and ATG2 are characterized by the presence of clusters of small vesicles, raising questions about a potential interplay between the lipid transport properties of these proteins and vesicle fusion[14–16]. As studies of VPS13 have informed regarding ATG2 function and vice versa, characterization of other family members should yield further insights both as to how these proteins function and the cellular processes in which they participate.

A predicted chorein motif is present at the N-terminus of SHIP164 (also called UHRF1BP1L) and in its paralogue UHRF1BP1 (also called C6orf107), and fold prediction algorithms indicate high confidence that downstrem portions form an extended chorein domain ß-sheet[18,19] (**Figure 1A**). SHIP164 was first identified as an interactor of the Habc domain of Syntaxin 6 (Stx6) and shown to be a component of a multimolecular assembly including subunits of GARP[20]. GARP is a tethering complex which along with the structurally similar EARP cooperates with Stx6 in the fusion of endocytic vesicles with acceptor membranes in endosomes (EARP) and the Golgi complex (GARP)[21–23]. Moreover, over-expressed SHIP164-Stx6 complex was shown to colocalize with the cation-independent mannose-6-phosphate receptor (MPR) and to perturb the traffic of this receptor[20]. More recently, SHIP164 was identified as a top hit in a cellular-based screen for Rab5 effectors, further supporting a role of SHIP164 in the early endocytic pathway[24]. The possibility that SHIP164 may represent a lipid transport protein suggests that its characterization may reveal novel aspects of the biology of the endosomal system. Additional interest in this protein comes from its identification as a Parkinson's disease (PD) candidate gene from a large-

93 scale whole exome sequencing (WES) study[25]. To date, Nothing is known about the cell biology of
94 UHRF1BP1.
95
The goal of this study was to test the hypothesis that SHIP164 may be a lipid transport protein
and to gain new information into its localization and physiological function. Here we provide
direct evidence for a structural similarity of SHIP164 to ATG2 and the N-terminal half of VPS13
and for the property of SHIP164 to harbor and transport lipids. We show that SHIP64 localizes to
clusters of small vesicles in the endocytic pathway and demonstrate its importance in retrograde
traffic to the Golgi complex by both over-expression and loss-of-function studies. We also provide
evidence for a link between SHIP164 and ATG9A, raising the possibility that SHIP164, like ATG2,
may mediate lipid transport in partnership with ATG9A. Finally, our findings predict similar
properties for UHRF1BP1.
105
106 **RESULTS**
107
108 **Molecular properties of SHIP164 support a lipid transport function**
109
As a first step in assessing whether SHIP164 might function as a lipid transporter in the VPS13-
family, we assayed whether it can solubilize and transport lipids between membranes, and, if so,
which ones. We expressed full-length, N-terminally FLAG-tagged human SHIP164 (3xFLAG-
SHIP164) in Expi293 cells and purified it using anti-FLAG affinity resin, amid extensive washing
to remove non-specifically bound lipids, then analyzed the sample for co-purified molecules by
shotgun mass spectrometry lipidomics (**Figure 1B**). A large assortment of lipids, primarily
phospholipids, co-purified with SHIP164. There was no significant enrichment of a particular
phospholipid relative to the content of the whole cell, however.
118
The full-length construct was prone to aggregation, whereas constructs lacking a long predicted
unstructured segment (residues 901-1099) were soluble and were used in subsequent in vitro
studies. Further supporting that SHIP164 binds phospholipids, 3XFLAG-SHIP164$_{\Delta901\text{-}1099}$ binds
nitrobenzoxadiazole (NBD)-labeled PE, co-migrating with this lipid on a native gel (**Figure 1C**).
From comparing the fluorescence that co-migrated with SHIP164 versus a well characterized lipid
transport module from the protein E-Syt2, we estimate that each SHIP164 molecule can
accommodate multiple lipids (~8), as might be expected if SHIP164 has an extended lipid binding
groove like VPS13 or ATG2[6,7].
127
We took advantage of the ability of SHIP164 to bind NBD-PE in designing an in vitro FRET-based
assay to monitor lipid transfer between membranes[6]. In this assay, SHIP164 is tethered (**Figure
1D**) between donor liposomes containing both rhodamine (Rh)- and NBD-labeled PE (61% DOPC,
30% PE, 2% NBD-PE, 2% Rh-PE, 5% Ni-NTA) and acceptor liposomes (65% DOPC, 30% PE, 5%
PI(4,5)P$_2$), initially lacking these fluorescent lipids. FRET between Rh and NBD reduces NBD
fluorescence initially. Transfer to acceptor liposomes and consequent dilution of the fluorescently
tagged lipids would reduce FRET and lead to increased NBD fluorescence. Consistent with a role
in lipid transfer, addition of SHIP164, but not of a an "empty" tether lacking a lipid transport
module, led to increased fluorescence (**Figure 1E**). Addition of SHIP164 to donor liposomes only,
in the absence of acceptor liposomes, led to a fast, smaller fluorescence increase due to the ability
of SHIP164 to extract lipids from the liposomes (**Fig. 1E, red arrow**), causing some dilution
(similar to observations for ATG2 and VPS13[6,7]). SHIP164 mediated fusion between donor and
acceptor liposomes would also lead to increased fluorescence. To discriminate between lipid
transfer (which occurs only between lipids in the outer leaflet of liposomes in the absence of
scramblases) and liposome fusion, we incubated donor and acceptor liposomes with and without
SHIP164, as before, then added dithionite to reduce NBD-labeled lipids exposed in the outer

leaflet. If SHIP164 had induced liposome fusion, the residual fluorescence after addition of
SHIP164 would be higher than in its absence, as lipids in the internal leaflet would have been
diluted[26]. In contrast, we found that the residual fluorescence after addition of dithionite was the
same with and without addition of SHIP164 reflecting the fluorescence of lipids in the internal
leaflet of donor liposomes that did not undergo dilution in the absence of membrane fusion
(**Figure 1F**). These experiments support that SHIP164 can solubilize lipids and transfer them
between membranes.

We next examined SHIP164 by electron microscopy. 2D class averages of negatively stained
3XFLAG-SHIP164$_{\Delta 901-1099}$ revealed a 200 Å long rod (**Figure 1G, Supplemental Figure 1A**).
Comparison of SHIP164 with a MBP-SHIP164 fusion indicates that this rod represents a tail-to-
tail dimer, as there are two densities corresponding to MBP, one at each end. The MBP-tag locates
the SHIP164 N-terminus at the rod end. Whether this dimerization is physiologically relevant is
unclear; for example, ATG2 similarly dimerizes in vitro, at high concentration, and in the absence
of binding partners, but is thought to be monomeric in vivo[7]. Further characterization of 3XFLAG-
SHIP164$_{\Delta 901-1099}$ using single particle cryo-electron microscopy techniques yielded a
reconstruction at an estimated resolution of ~8.3 Å (**Supplemental Figure 1B&C**). As
suggested by fold prediction algorithms (ref. [18,19])(see **Figure 1Ac-d**), a cavity runs along the
entire length of the rod, as in VPS13 and ATG2 **(Figure 1 H&I)**. The algorithms also predict that
this cavity is lined entirely with hydrophobic residues (**Figure 1Ac-d**). Thus, the cavity can
accommodate the multiple lipids bound by SHIP164, and we propose that it could be a conduit
for lipids to transit between membranes.

Collectively, these findings are consistent with the hypothesis that SHIP164 is a lipid transport
protein. We next investigated its site of action within cells.

Localization of exogenous SHIP164 points to a role on endocytic organelles

Antibodies directed against SHIP164 did not yield a consistently reliable signal when tested by
immunofluorescence. Thus, we examined the localization of exogenously tagged human SHIP164.
In agreement with the study by Otto et al[20], SHIP164 tagged at the N-terminus (GFP-SHIP164)
had a diffuse cytosolic localization when expressed in mammalian cell lines (e.g. HeLa, COS-7,
RPE1) (**Supplemental Figure 2A**). As studies of VPS13 had shown that tags appended to the
N-terminus (i.e the N-terminus of the chorein motif) interfere with its physiological function[6,27],
we engineered tags at the C-terminus of SHIP164 (SHIP164-Halo) or at an internal site within the
predicted disordered region (after amino acid residue 915; SHIP164^mScarlet-I) (**Figure 2A**)
(ref. [6,27]). When either of these tagged SHIP164 constructs was expressed alone, punctate
structures of varying size were observed throughout the cytoplasm, with the larger and brighter
spots being localized in the central region of the cell, in proximity of the Golgi complex area
(**Figure 2A**). Similar results were observed with other tags (Halo, GFP, mRFP, etc.) at either
position. Conversely, even a small N-terminal tag appended to SHIP164^mScarlet-I (3xFLAG-
SHIP164^mScarlet-I) abolished the presence of cytoplasmic foci (**Figure 2B**).

The scattered distribution of SHIP164^mScarlet-I foci throughout the cytosol combined with the
results of Otto et al[20] suggesting a role of SHIP164 in the endocytic pathway, prompted us to
examine the colocalization of SHIP164 relative to endosome markers. Many, but not all SHIP164
puncta, localized in proximity of spots or vacuoles positive for GFP-2xHrs[FYVE], a PI3P binding
probe[28] (**Figure 2C**). Moreover, SHIP164^mScarlet-I spots were observed in close proximity to
other co-expressed markers of endosome subcompartments, such as APPL1 (a very early, PI3P-
negative endosomal station), WDFY2 (an endosomal station downstream of APPL1), and EEA1[29]
(**Figure 2D**). SHIP164, however, did not localize on the organelles positive for these probes, but

on foci juxtaposed to them (**Figure 2C-E**). This was even more clear by observing
macropinosomes (both naturally occurring or induced by expressing constitutively active
Ras[G12V]), given the large size of these vesicles[29] (**Supplemental Figure 2B**). As shown by live-
cell imaging, SHIP164 foci not only were variable in size as well as in fluorescence intensity, but
were also highly dynamic structures. They changed shape, often underwent fission, or coalesced
into larger spots although they tended to remain tethered to endosomes and micropinosomes
(**Figure 2F; Supplemental Movie 1**). In some cases, they were localized at sites where the
endosomes appeared to be close to the ER (**Figure 2E**), but were not restricted to the space
between these two organelles.
204
A more precise colocalization of exogenous SHIP164 was observed with Rab5 (BFP-Rab5b),
confirming previous results[20,24] (**Figure 2D**). In this case a pool of SHIP164 colocalized with Rab5
along the entire profile of vacuoles and, conversely, Rab5 robustly colocalized with the bright
SHIP164 foci closely apposed to large vacuoles (**Figure 2D**). This very strong colocalization
supports the hypothesis that SHIP164 is a Rab5 effector. However, expression of a dominant
negative Rab5 mutant (GFP-Rab5a[S34N]) did not abolish bright SHIP164 (SHIP164-Halo) foci
suggesting that Rab5 is not necessary for their formation (**Supplemental Figure 2C**).
212
Notably, exogenous SHIP164 puncta colocalized with endogenous cation-independent mannose-
6-phosphate receptor (MPR) (as detected by immunofluorescence) and its expression (leading to
higher than normal levels of SHIP164) also resulted in an alteration of the localization of the MPR,
which accumulated in large foci where the two proteins colocalized (**Figure 2G**). These
accumulations occurred throughout the cell but were especially large in proximity of the Golgi
complex, where the normal Trans-Golgi Network (TGN)-like appearance of MPR was replaced by
these large accumulations located at some distance from GM130, a Golgi complex marker[30] (see
Figure 2G for COS-7 cells and **Supplemental Figure 2D** for HeLa cells). A similar
redistribution of MPR and also of the Golgi/endosome protein sortilin[31] was observed in RPE1
(retinal pigment epithelial) cells transfected with a polycistronic expression plasmid encoding
untagged SHIP164 and a soluble RFP to confirm transfection, ruling out non-specific effects of
the engineered fluorescent tag of exogenous SHIP164 (**Figure H&I; Supplemental Figure
2E&F**). Such changes were not observed with TGN46, another protein enriched in the TGN that
cycles to and from the plasma membrane and endosomes[21,32,33] (**Supplemental Figure 2G&H**).
This difference is in line with the known traffic of MPR and sortilin versus TGN46 in different
vesicular carriers[33,34] (see also below).
229
230 **Foci of exogenous SHIP164 reflect accumulations of small vesicles**
231
To gain further insight into the precise nature of foci of over-expressed SHIP164, we performed
correlative light-electron microscopy (CLEM) of COS-7 cells co-expressing the early endosome
marker GFP-WDFY2 and SHIP164^mScarlet-I (**Figure 3A**). Both conventional EM and FIB-
SEM were performed. We again employed Ras[G12V] to induce the formation of macropinosomes[35]
to facilitate the alignment of fluorescence and EM images. This analysis revealed that spots of
SHIP164 fluorescence reflected accumulations of 100's of tightly packed approximately 50-60 nm
vesicles juxtaposed to endosomal membranes positive for GFP-WDFY2 (**Figure 3A**). An
approximately 70nm thick matrix appeared to anchor these clusters to the WDFY2-positive
vacuoles and the space occupied by this matrix was nearly devoid of vesicles (**Figure 3A&C**). ER
tubules were present in proximity of these clusters and often penetrated them, but there did not
seem to be sites of preferential accumulation of vesicles (**Figure 3B**). The appearance of these
vesicle accumulations and their dynamic nature revealed by live imaging (see above) was
reminiscent of other vesicle clusters, such as synaptic vesicles clusters in nerve terminals, recently
shown to be organized according to principles of liquid-liquid phase separation[36–38]. As such

"phases" often involve low affinity interaction of disordered protein regions[39], we considered the
possibility that the "disordered region" of SHIP164 (see **Figure 1Ac**) may be responsible for their
formation. However, a truncated form of SHIP164 (SHIP164$_{1-873}$-mRFP) that lacks the disordered
region also induced the formation of large fluorescence puncta closely juxtaposed to endosomes
(**Supplemental Figure 3**). Thus, the mechanism responsible for the clustering of SHIP164
positive vesicles remains unclear. Also unclear is how such clusters are anchored to the large
vacuoles.
253
254 As these vesicle accumulations, which to our knowledge were never observed in WT cells,
255 suggested an abnormal effect of over-expressed SHIP164, we next tagged SHIP164 at the
256 endogenous locus to assess its localization when expressed at the endogenous level.
257
258 **Endogenous SHIP164 localizes to small clusters of vesicles near the cell edge**
259
We tagged SHIP164 at the endogenous locus in HeLa cells. Specifically, we engineered a
mNeonGreen (mNG) tag in the disordered region after amino acid residue 1092 via the ORANGE
genomic editing method[40] (**Figure 4A**). Two clones were isolated where expression of
endogenous SHIP164^mNG (eSHIP164^mNG) was confirmed by the presence of a band running
at the appropriate molecular weight and recognized by western blotting with both anti-SHIP164
and anti-mNeonGreen antibodies (**Figure4B**). Correct insertion of the tag was validated by DNA
sequencing of the targeted gene region (**Figure 4C**).
267
Like exogenous SHIP164, eSHIP164^mNG had a punctate appearance. However,
eSHIP164^mNG puncta were very much smaller and, in contrast to foci of over-expressed
SHIP164 (**Figure 2A**), were primarily localized at the cell edge (**Figure 4D**). On the other hand,
when SHIP164^mScarlet-I was expressed in these endogenously tagged cells, exogenous
mScarlet-I and endogenous mNeonGreen fluorescence precisely overlapped in large foci,
including the ones localized in deep regions of the cell (**Supplemental Figure 4A**). This
demonstrates that endogenous and exogenous SHIP164 completely intermix—as expected from
correct gene targeting—confirming that large foci are the result of SHIP164 over-expression.
276
To determine whether puncta of eSHIP164^mNG represent MPR positive vesicles, we generated
a double knock-in (DKI) HeLa cell line where both SHIP164 and MPR were tagged at their
endogenous loci (eSHIP164^mNG and eMPR-mScarlet-I). Most, but not all, eSHIP164^mNG
positive puncta colocalized and moved together with puncta of eMPR-mScarlet-I fluorescence
(**Figure 4E; Supplemental Movie 2**). Conversely, many eMPR-mScarlet-I puncta at the edge
of the cell colocalized and moved together with puncta of eSHIP164^mNG (**Figure 4E**), while
MPR vesicles located deeper into the cell were negative for SHIP164[41,42]. Fluorescent puncta
positive for the two proteins were distinct from vesicles positive for newly internalized labeled
EGF (EGF-647) revealing that endogenous SHIP164 localizes to a specific endocytic
subpopulation (**Supplemental Figure 4B**). Interestingly, an intrinsic membrane protein of
unknown function, that is listed as a SHIP164 interactor in BioGRID (https://thebiogrid.org),
FAM174a, strikingly colocalized with exogenous SHIP164 accumulations, although FAM174a was
also present in the Golgi complex and other compartments of the secretory/endocytic pathways
(**Supplemental Figure 4C&D)**.
291
To gain further insight into the nature of endogenous SHIP164 puncta, we again employed CLEM
in eSHIP164^mNG HeLa cells also expressing a marker of mitochondria (mito-BFP) to aid in
fluorescence and EM image alignment (**Figure 4F**). Given the small size of the fluorescent
structures, FIB-SEM rather than TEM was used in spite of the lower resolution afforded by FIB-
SEM as this technique allows a better alignment between fluorescence and EM images in 3D. This

analysis revealed that, similar to the large fluorescence puncta of exogenously expressed SHIP164,
spots of eSHIP164^mNG fluorescence reflected tightly packed clusters of small vesicles and short
tubules (**Figure 4F&G**). However, these clusters were much smaller (10's instead of 100's of
vesicular structures). As revealed by 3D reconstruction of FIB-SEM volumes (**Figure 4G**) and by
fluorescence of cells also expressing the ER marker RFP-Sec61 (**Figure 4H**), ER tubules were
localized in close proximity of these clusters but vesicles were preferentially associated with
themselves rather than with the ER.
304
Collectively, these findings indicate that SHIP164, when expressed at endogenous levels, is
selectively localized on small MPR-positive vesicles near the cell periphery. They further imply
that SHIP164 dissociates from such vesicles as they travel to deeper cellular regions and that the
presence of exogenous SHIP164 on the very large vesicle clusters in central cell regions near the
Golgi complex reflect an over-expression defect. To better understand SHIP164's dynamics in the
endocytic pathway we investigated its interactions.
311
312   **Interaction of SHIP164 with proteins implicated in the targeting of endocytic**
313   **cargoes to the Golgi complex.**
314
We confirmed the previously reported colocalization of exogenous SHIP164 with over-expressed
Stx6[20] (**Figure 5A**). SHIP164 and Stx6 colocalized on spots scattered throughout the cytoplasm
and in the Golgi complex area irrespective of the presence or absence of N-terminal tags, i.e. tags
that impair the localization of SHIP164 to vesicles enriched foci when expressed alone
(**Supplemental Figure 2A**). This indicates that binding to Stx6 does not play a major role in
the localizations of SHIP164 described above.
321
The location of the Stx6 binding site within the SHIP164 protein was not known. A short
hydrophobic motif ($L_{37}xxYY_{41}$) responsible for binding to the Habc domain of Stx6 was previously
identified in the N-terminal region of VPS51[23]. A similar motif ($L_{759}xxYY_{763}$) is present in a
predicted disordered loop of SHIP164 within the extended chorein domain (**Figure 5E; Figure**
1Ac) and mutations of both tyrosine resides (Y772A, Y773A) abolished the recruitment of GFP-
SHIP164 (GFP-SHIP164$^{Y772A, Y773A}$; i.e. N-terminally tagged SHIP164 that when expressed by itself
is cytosolic) to mRFP-Stx6 (**Supplemental Figure 5A**). However, the same mutations, when
introduced into C-terminally tagged SHIP164 (SHIP164$^{Y772A, Y773A}$-mRFP) did not affect the
localization of the protein into bright foci in cells not transfected with Stx6, further indicating that
Stx6 does not account for these localizations (**Supplemental Figure 5B**).
332
To identify additional SHIP164 interacting partners that may impact its localization and/or
function, we performed pull-down experiments from detergent solubilized mouse brain lysates
onto purified SHIP164 (3xFLAG-SHIP164) bound to anti-FLAG resin. Shotgun mass
spectrometry of the affinity-purified material showed a significant specific enrichment (relative
to control) of the light chains 1 and 2 (DYNLL1/2; also known as LC8) of cytoplasmic dynein, the
minus-end directed microtubule motor[43] (**Figure 5B**). DYNLL1/2 act as dynein complex
adaptors which mediate a wide variety of protein-protein interactions through a short linear
binding motif[44] ($V_{-5}[S/N/T]_{-4}[K/R]_{-3}X_{-2}T_{-1}Q_0T_{+1}$). Accordingly, a motif that fits this consensus is
present in SHIP164 near its C-terminus ($V_{1395}TQATQT_{1401}$) and LC8 is another protein listed
among putative SHIP164 interactors in BioGRID (**Figure 5E**). Moreover, when an MBP-tagged
fragment of SHIP164 containing this region (MBP-SHIP164$_{1312-1464}$) and untagged LC8 were co-
expressed in bacteria from a polycistronic plasmid, LC8 co-purified with MBP-SHIP164$_{1312-1464}$ on
amylose-resin and co-fractionated with MBP-SHIP164$_{1312-1464}$ when the eluate of the resin was
fractionated on a size-exclusion column indicating a direct interaction (**Figure 5C**).
347

Interestingly, another potential SHIP164 interactor listed in BioGRID, Rab45 (RASEF), is linked
to dynein function. Rab45 and its better-characterized paralog CRACR2a are dynein/dynactin
activators that connect endocytic cargoes to minus-end microtubule transport[45,46]. In agreement
with the possibility that SHIP164 may be an effector of Rab45, over-expression of Rab45
dramatically concentrated both transfected SHIP164^mScarlet-I and endogenously tagged
SHIP164 (eSHIP164^mNG) to perinuclear spots (i.e. the position occupied by the centrosome),
depleting them from the periphery (**Figure 5D; Supplemental Figure 5C**). At this perinuclear
location, SHIP164 and Rab45 precisely colocalized (**Figure 5D; Supplemental Figure 5C**).
Over-expression of Rab45 also clustered endogenous MPR tagged at the C-terminus with a
mNeonGreen tag (eMPR-mNeonGreen), but only in cells over-expressing SHIP64, i.e. in cells
where the majority of MPR-positive vesicles (not only the peripheral vesicles) are positive for
SHIP164 (**Supplemental Figure 5D**). A dominant-inactive Rab45 (S555N) mutant had a
cytosolic localization did not induce SHIP164 accumulation in this area suggesting that the
nucleotide bound state of Rab45 regulates recruitment (**Supplemental Figure 5E**).
362
363   Taken together, our interaction studies point to an association of SHIP164 with vesicles that traffic
364   from the cell periphery to the centrosomal/Golgi complex region, although endogenous SHIP164
365   appears to be shed by vesicles as they move to central regions of the cells unless Rab45 is over-
366   expressed. One would therefore expect a role of SHIP164 in some aspect of retrograde membrane
367   traffic. To validate this hypothesis, we performed loss-of-function studies.
368
369   **Defects in retrograde membrane traffic to the TGN in SHIP164 knockout cells**
370
To assess the impact of the absence of SHIP164 on cell physiology we chose RPE1 cells as a model
system due to their reliable genetic editing and flat morphology optimal for organelle imaging.
Using CRISPR/Cas9 methodology, two independent SHIP164 knockout (KO) clones were
generated where indels led to a frameshift of the reading frame that resulted in premature stop
codons as shown by sequencing (**Supplemental Figure 6A**). In both clones, identical indels
were observed on both alleles (but different between the two clones) and absence of the SHIP164
protein was confirmed by western blotting (**Figure 6A**). No obvious changes were observed in
cell shape, in mitochondria, ER, or Golgi morphology as assessed by the cis-Golgi marker
(GM130) (see below). Two differences, however, were noted.
380
381   One was a difference in the size of endosomes labeled by EEA1 (visualized by immunostaining),
382   and GFP-2xHrs[FYVE] (visualized by live-cell imaging) as the largest (>1 μm[2]) vesicles positive for
383   these markers in WT cells were mostly absent in KO cells (**Figures 6B-C,** **, $p<0.01$).
384
The other difference was a change in the steady state localization of MPR and other proteins that
recycle between the cell periphery (plasma membrane and endosomes) and the TGN. Specifically,
upon immunostaining of SHIP164 KO cells for the MPR, an increase in the pool of MPR localized
in vesicles scattered throughout the cytoplasm relative to the pool localized in the Golgi complex
area was observed[42], (**Figure 6D,** **, $p<0.01$). The localizations by immunofluorescence of
TGN46 and of Sortilin showed a similar difference relative to controls (**Figure 6E &
Supplemental Figure B,** **, $p<0.01$), although the vesicles positive for TGN46 were different
from those positive for MPR and sortilin (**Supplemental Figure 6C**). As a control, the ratio of
clathrin heavy chain (CHC) immunoreactivity sparse throughout the cytoplasm relative to
immunoreactivity in the Golgi complex was also measured and such ratio was not statistically
different between SHIP164 KO cells and parental control cells (**Figure 6F**).
396
To validate the dependence of these phenotypes from the lack of SHIP164, we attempted to rescue
them by expressing exogenous SHIP164 in the KO cells. To this aim we used the polycistronic

construct described above encoding both untagged SHIP164, in order to avoid any artifact due to
the tag, and soluble RFP, to identify transfected cells. In both KO clones, transfection of this
construct (likely resulting in levels of SHIP164 higher than in WT cells), but not of a construct
encoding only the RFP reporter, resulted in the rescue of the vesicles-to-Golgi ratio of TGN46
immunoreactivity (**Figure 6G**; *, p<0.05; **, p<0.01). However, in the case of MPR and sortilin,
expression of the SHIP164 construct did not restore their WT localization, but resulted in their
robust accumulation into large foci as shown above upon SHIP164 over-expression in wild type
cells (**Supplemental Figure 6D**). This result is most likely explained by the disrupting effect of
SHIP164 over-expression, which, as shown above, does not impact TGN46 localization
(**Supplemental Figure 2G&H**). Likewise, SHIP164 over-expression did not seem to rescue the
presence of the large EEA1 positive endosomes present in WT cells as it induced large focal EEA1
accumultions, but these did not have the typical vacuolar apperance. (**Supplemental Figure
6E**). Also in this case this result may be to due a dominant negative effect as similar EEA1 positive
clusters rather than the typical vacuolar structures were observed in WT cells over-expressing
SHIP164 (**Supplemental Figure 6E**). However, we confirmed the impact of SHIP164
deficiency on the presence of large EEA1 endosomes by SHIP164 siRNA-dependent knockdown
in RPE-1 cells, thus ruling out off-target effects of CRISPR/Cas9 editing (**Supplemental Figure
6F&G**; **, p<0.01).
417
418 We conclude that loss of SHIP164 function may globally impact membrane traffic between the
419 cell periphery and the Golgi complex, while over-expression of SHIP164 disrupts selectively the
420 retrograde traffic of a subset of the vesicles with which SHIP164 is associated.
421
422 **Presence of ATG9A in clusters of SHIP164-positive vesicles**
423
The structural similarity of SHIP164 to proteins thought to mediate bulk lipid transport—
particularly ATG2—suggests a similar function of SHIP164[6,7,11,47]. A plausible scenario is that
SHIP164 may allow the regulation of bilayer lipids (abundance and possibly also type of lipids)
on organelles of the endocytic pathway by mediating transient contacts with these organelles and
either the ER or other membranes to transport lipids[4,9]. The lack of the large endosomes in
SHIP164 KO cells would be consistent with a role of SHIP164 in bulk transfer of lipids. So far,
however, we have not found direct evidence for a role of SHIP164 as a tether between two
membranes. Moreover, as stated above, the endosomal size phenotype could not be clearly
rescued by the expression of exogenous SHIP164 in SHIP164 KO cells (**Supplemental Figure
6E**), thus preventing us from directly testing this hypothesis using SHIP164 proteins that cannot
transport lipids, in addition to a propensity to aggregate in living cells of putative lipid-transport
dead mutants that we generated.
436
ATG2 was found to function in cooperation with the scramblase ATG9, as lipids are delivered by
ATG2 to the cytosolic leaflet of the receiving membrane and thus lipid scrambling is needed to
allow equilibration of lipid concentration on both bilayer leaflets[10,48–51]. In WT cells, ATG9
(ATG9A) is concentrated in the Golgi complex area and also present as scattered dots throughout
the cytoplasm (**Figure 7A**). Interestingly, we found that ATG9A-GFP is highly concentrated on
the SHIP164 positive foci resulting from SHIP164 over-expression (**Figure 7B**). Moreover, while
no major redistribution of ATG9A-GFP is observed in cells over-expressing Rab45 (RFP-Rab45),
co-expression of SHIP164-Halo with RFP-Rab45 resulted in a striking co-enrichment and
colocalization of all three proteins in the centrosomal area proving that ATG9A is localized on
SHIP164-positive vesicles (**Figure 7C**). We also explored whether autophagy was affected in
SHIP164 KO cells, but we did not find any effect on LC3 abundance upon starvation. While the
localization of ATG9A on SHIP164 positive vesicles does not provide evidence for a lipid transport

activity of SHIP164, it is consistent with such an activity and raises the interesting possibility that
ATG9A may have more general functions in cell physiology than a selective role in autophagy[52].
451
452 **UHRF1BP1, a SHIP164 paralogue, is also localized in the endocytic pathway**
453
UHRF1BP1 shares substantial primary sequence similarity (42% identity) and domain
organization with SHIP164[20]. Accordingly, phenotypes produced by expression of exogeneous
UHRF1BP1 (UHRF1BP1-Halo) revealed similarities to those produced by the expression of
exogenous SHIP164. As in the case of SHIP164, UHRF1BP1-Halo localized to foci of varying size
and fluorescence intensity with the larger foci being concentrated in the central region of the cell
(**Figure 7D**). These foci were positive for exogenous ATG9A (ATG9A-GFP) (**Figure 7E**).
However, in contrast to SHIP164 foci, which were primarily associated with early endosomes,
UHRF1BP1 foci were predominantly associated with LAMP1-GFP-positive organelles, in
agreement with previous work suggesting that UHRF1BP1 is a Rab7 effector, rather then a Rab5
effector[24] (**Supplemental Figure 7**). We conclude that SHIP164 and UHRF1BP1 likely function
in a similar manner but at different stations of the endocytic pathway
465
466 **DISCUSSION**
467
Our study identifies SHIP164 as a bona fide member of the chorein motif protein family with a
role in membrane traffic. The EM-analyses of SHIP164 show that as in VPS13 and ATG2, the two
well established members of this family, the N-terminal chorein motif of SHIP164 caps an
extended rod that harbors a cavity running along its length[6–8,47]. Fold prediction algorithms
(Alphafold, trRosetta) indicate that SHIP164 features an extended ß-sheet that resembles a taco
shell, whose concave elongated surface is lined with hydrophobic residues, and so constitutes a
groove that can solubilize multiple lipids[18,19]. In vitro lipid binding and transfer assays further
support that SHIP164 might function as a lipid transfer protein. The SHIP164 paralogue
UHRF1BP1 is predicted to have a similar structure.
477
Our analysis of the subcellular localization of SHIP164 reveals an association with endocytic
organelles, extending previous observations[20,24]. This is supported by several pieces of evidence:
1) Endogenous SHIP164 is predominantly localized on a population of small peripheral vesicles
that assemble in small clusters and contain in their membranes endosomal cargo proteins, such
as MPR[42]. 2) Over-expressed SHIP164 drastically enlarges the size of these clusters, which
become, in many cases, tightly associated with vacuoles positive for markers of early endosomal
subcompartments, such as Rab5, APPL1, WDFY2, PI3P, and EEA1 and are found also in deep
regions of the cell. 3) The interactors of SHIP164 that we have identified or validated are endocytic
factors or proteins that mediate retrograde organelle traffic from the cell periphery to the
centrosomal/Golgi complex area. One of them is Rab5, an early endosomal Rab[53]. Another one is
Stx6, a component of SNARE complexes that mediates the fusion of vesicles with recycling
endosomes and with the Golgi complex and whose action is facilitated by the GARP and EARP
tethering complexes[54–56]. Interestingly, binding of SHIP164 to the Habc domain of Stx6 is
mediated by the same motif used by the GARP and EARP subunit VPS51, also a Stx6
interactor[21,23]. While this finding implies a competitive binding to Stx6, it also suggests an
interplay of SHIP164 with GARP, further linking SHIP164 to retrograde traffic. A third interactor
is the light chain of dynein (DYNLL1/2), the microtubule motor that mediates retrograde traffic[57].
The fourth is Rab45, an unconventional Rab that anchors endocytic vesicles to dynein, creating
an additional link to dynein and retrograde traffic[45,46]. Notably, however, SHIP164, when
expressed at endogenous levels is predominantly associated with peripheral vesicles, suggesting
a regulated interaction with vesicles that is lost and replaced by other interactors as they travel to
deeper destinations in the cell.

500
A role of SHIP164 in retrograde traffic is further supported by SHIP164 loss of function and gain-
of-function studies. In SHIP164 KO cells, a larger pool of proteins that shuttle between the cell
periphery and the Golgi complex (MPR, sortilin and TGN46) is localized on vesicles scattered
throughout the cytoplasm relative to their major pool localized in the Golgi complex area
suggesting a defect in transport/docking or fusion with the Golgi complex. Similar observations
have been made in cells deficient in GARP function[21,58,59]. Conversely, SHIP164 over-expression
(gain-of-function) results in large vesicle clusters positive for MPR and sortilin, but not for
TGN46, indicating that loss-of-function and gain-of-function effects are not precise opposites. A
potential scenario is that the accumulation of sortilin and MPR vesicles results from a block in
their progression to the next traffic station produced by excess SHIP164. Such dominant negative
effect could be the result of sequestration of, or competition with, critical factor(s) (for example
GARP) by over-expressed SHIP164 or by alteration of membrane lipid homeostasis.
513
A key question is how the structure and molecular properties of SHIP164, which predict a lipid
transport function at intracellular membrane contact sites, similar to the proposed function of
VPS13 and ATG2, relates to the observed localization and loss-of-function phenotypes of
SHIP164. We consider the hypothesis that SHIP164, as proposed for VPS13 and ATG2, may
function in the transport of lipids from one membrane where the chorein domain may extract
lipids – for example the ER – to other membranes[9] and more specifically to the small vesicles
enriched with endogenous SHIP164. This transport may deliver specific lipids or simply be
responsible for membrane expansion. However, the localization of SHIP164 that we have
observed is not at contacts of two types of membranous organelles, as predicted by this model.
While some SHIP164 foci detectable by fluorescence appear to be localized at the interface of the
ER and endosomes, such foci are represented by vesicle clusters and not by SHIP164 tethers
between the ER and endosomes. Moreover, until now we have not identified the binding of
SHIP164 to any ER protein. SHIP164 does not have an FFAT or FFAT-like motif that, as in the
case of VPS13[6,60], could account for binding to the ER protein VAP[61,62]. In spite of these
considerations, binding of SHIP164 to the ER cannot be excluded. Binding to the ER of SHIP164
tagged at the endogenous locus is difficult to analyze given the small size of SHIP164 associated
vesicles. SHIP164 over-expression may overwhelm a binding site on the ER, unless such binding
site was also over-expressed. One should also consider that an interaction with the ER of
SHIP164-positive transport vesicles may be regulated and only transient.
533
Of special interest in the context of protein-mediated lipid transport is the strong colocalization
of SHIP64 with ATG9A, as ATG9A is a functional partner of ATG2 in the expansion of the
autophagic membrane[48–50,63]. ATG9A, a protein scramblase, is thought to allow equilibration
between the two bilayer leaflets, as phospholipids are delivered to the cytosolic leaflet of the
autophagic membrane by ATG2[48–50]. While ATG9A was discovered as a core component of the
autophagy machinery, it is plausible that it may function in other contexts, as also recently
reported[52], more so in cooperation with a protein that shares strong similarities with ATG2, such
as SHIP164. Interestingly, studies of ATG9A have consistently localized it to small vesicles often
organized in clusters, similar to the SHIP164 vesicles described here[16,64]. Thus, the overlapping
localization of SHIP164 and ATG9A is consistent with a role of SHIP164 in lipid transport. The
absence of large endosomes in SHIP164 KO cells would be supportive of a role of SHIP164 in lipid
transfer to expand the membranes of endocytic organelles, but since this phenotype could not be
clearly rescued (possibly due to dominant negative effects of SHIP164 overexpression), its
significance remains uncertain. Future studies will be needed to provide further mechanistic
insight into these findings. We note that mechanisms to control bulk lipid transport to the
membrane of an endocytic organelle independent of membrane fusion may help fine tune the

protein/lipid ratio of such membrane and may therefore play an important role in controlling its
maturation as it travels to deeper positions in the cell.
552
Finally, in view of the link between the function of SHIP164 and GARP suggested by our study
and a previous study[20], it is of interest that defects in retrograde traffic from the cell periphery to
the Golgi complex produced by SHIP164 loss-of-function have similarities to those observed upon
loss of function of the GARP complex[20,58]. GARP has also been linked to lipid metabolism. In
zebrafish, loss of the GARP subunit fat-free (Ffr) (later identified as the paralog to mammalian
VPS51) results in a striking defect of the absorption into the body of dietary lipids[65]. Defects in
GARP in mice or humans, either at the cellular or organismal level, result in abnormal
sphingolipid metabolism (with an abnormal accumulation of long chain bases) and
neurodegenerative conditions[59,66,67]. SHIP164 was reported be a Parkinson's disease candidate
gene[25] and loss-of-function mutations in VPS13C, another chorein motif lipid transport protein
implicated in the endocytic/lysosome pathway is responsible for familial forms of Parkinson's
disease[68]. It is therefore possible that SHIP164 may be part of a network of proteins that play an
important role in neuronal health by controlling lipid transport and metabolism at the interface
with membrane traffic.
567

REFERENCES

1.	Alva, V. & Lupas, A. N. The TULIP superfamily of eukaryotic lipid-binding proteins as a mediator of lipid sensing and transport. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1861,** 913–923 (2016).

2.	Prinz, W. A., Toulmay, A. & Balla, T. The functional universe of membrane contact sites. *Nature Reviews Molecular Cell Biology* **21,** 7–24 (2020).

3.	Saheki, Y. & Camilli, P. D. Endoplasmic Reticulum-Plasma Membrane Contact Sites. *Annual review of biochemistry* **86,** 659–684 (2017).

4.	Reinisch, K. M. & Prinz, W. A. Mechanisms of nonvesicular lipid transport. *Journal of Cell Biology* **220,** (2021).

5.	Wong, L. H., Gatta, A. T. & Levine, T. P. Lipid transfer proteins: the lipid commute via shuttles, bridges and tubes. *Nature Reviews Molecular Cell Biology* **20,** 85–101 (2019).

6.	Kumar, N., Leonzino, M., Hancock-Cerutti, W., Horenkamp, F. A., Li, P. Q., Lees, J. A., Wheeler, H., Reinisch, K. M. & De Camilli, P. VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact sites. *The Journal of cell biology* (2018).

7.	Valverde, D. P., Yu, S., Boggavarapu, V., Kumar, N., Lees, J. A., Walz, T., Reinisch, K. M. & Melia, T. J. ATG2 transports lipids to promote autophagosome biogenesis. *The Journal of cell biology* **14,** jcb.201811139 (2019).

8.	Li, P., Lees, J. A., Lusk, C. P. & Reinisch, K. M. Cryo-EM reconstruction of a VPS13 fragment reveals a long groove to channel lipids between membranes. *The Journal of cell biology* **219,** (2020).

9.	Leonzino, M., Reinisch, K. M. & De Camilli, P. Insights into VPS13 properties and function reveal a new mechanism of eukaryotic lipid transport. *Biochim Biophys Acta Mol Cell Biol Lipids* **1866,** 159003 (2021).

10.	Noda, N. N. Atg2 and Atg9: Intermembrane and interleaflet lipid transporters driving autophagy. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1866,** 158956 (2021).

11.	Osawa, T., Kotani, T., Kawaoka, T., Hirata, E., Suzuki, K., Nakatogawa, H., Ohsumi, Y. & Noda, N. N. Atg2 mediates direct lipid transfer between membranes for autophagosome formation. *Nat Struct Mol Biol* **26,** 281–288 (2019).

12.	Ueno, S., Maruki, Y., Nakamura, M., Tomemori, Y., Kamae, K., Tanabe, H., Yamashita, Y., Matsuda, S., Kaneko, S. & Sano, A. The gene encoding a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. *Nat Genet* **28,** 121–122 (2001).

13.	Rampoldi, L., Dobson-Stone, C., Rubio, J. P., Danek, A., Chalmers, R. M., Wood, N. W., Verellen, C., Ferrer, X., Malandrini, A., Fabrizi, G. M., Brown, R., Vance, J., Pericak-Vance, M., Rudolf, G., Carrè, S., Alonso, E., Manfredi, M., Németh, A. H. & Monaco, A. P. A conserved sorting-associated protein is mutant in chorea-acanthocytosis. *Nat Genet* **28,** 119–120 (2001).

14.	Park, J.-S., Okumura, Y., Tachikawa, H. & Neiman, A. M. SPO71 Encodes a Developmental Stage-Specific Partner for Vps13 in Saccharomyces cerevisiae. *Eukaryotic Cell* **12,** 1530–1537 (2013).

15.	Da Costa, R., Bordessoules, M., Guilleman, M., Carmignac, V., Lhussiez, V., Courot, H., Bataille, A., Chlémaire, A., Bruno, C., Fauque, P., Thauvin, C., Faivre, L. & Duplomb, L. Vps13b is required for acrosome biogenesis through functions in Golgi dynamic and membrane trafficking. *Cell. Mol. Life Sci.* **77,** 511–529 (2020).

16.	Mari, M., Griffith, J., Rieter, E., Krishnappa, L., Klionsky, D. J. & Reggiori, F. An Atg9-containing compartment that functions in the early steps of autophagosome biogenesis. *Journal of Cell Biology* **190,** 1005–1022 (2010).

17.	Chang, C., Jensen, L. E. & Hurley, J. H. Autophagosome biogenesis comes out of the black box. *Nat Cell Biol* **23,** 450–456 (2021).

18.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A.
A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T.,
Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M.,
Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli,
P. & Hassabis, D. Highly accurate protein structure prediction with AlphaFold. *Nature* **596,**
583–589 (2021).
19.   Yang, J., Anishchenko, I., Park, H., Peng, Z., Ovchinnikov, S. & Baker, D. Improved
protein structure prediction using predicted interresidue orientations. *PNAS* **117,** 1496–1503
(2020).
20.   Otto, G. P., Razi, M., Morvan, J., Stenner, F. & Tooze, S. A. A novel syntaxin 6-
interacting protein, SHIP164, regulates syntaxin 6-dependent sorting from early endosomes.
Traffic **11,** 688–705 (2010).
21.   Pérez-Victoria, F. J. & Bonifacino, J. S. Dual roles of the mammalian GARP complex in
tethering and SNARE complex assembly at the trans-golgi network. *Molecular and cellular*
biology **29,** 5251–5263 (2009).
22.   Schindler, C., Chen, Y., Pu, J., Guo, X. & Bonifacino, J. S. EARP is a multisubunit
tethering complex involved in endocytic recycling. *Nature cell biology* **17,** 639–650 (2015).
23.   Abascal-Palacios, G., Schindler, C., Rojas, A. L., Bonifacino, J. S. & Hierro, A. Structural
basis for the interaction of the Golgi-Associated Retrograde Protein Complex with the t-
SNARE Syntaxin 6. *Structure* **21,** 1698–1706 (2013).
24.   Gillingham, A. K., Bertram, J., Begum, F. & Munro, S. In vivo identification of GTPase
interactors by mitochondrial relocalization and proximity biotinylation. *eLife* **8,** e45916
(2019).
25.   Jansen, I. E., Ye, H., Heetveld, S., Lechler, M. C., Michels, H., Seinstra, R. I., Lubbe, S. J.,
Drouet, V., Lesage, S., Majounie, E., Gibbs, J. R., Nalls, M. A., Ryten, M., Botia, J. A.,
Vandrovcova, J., Simon-Sanchez, J., Castillo-Lizardo, M., Rizzu, P., Blauwendraat, C.,
Chouhan, A. K., Li, Y., Yogi, P., Amin, N., Duijn, C. M. van, (IPGDC), I. P. D. G. C., Morris, H.
R., Brice, A., Singleton, A. B., David, D. C., Nollen, E. A., Jain, S., Shulman, J. M. & Heutink,
P. Discovery and functional prioritization of Parkinson's disease candidate genes from large-
scale whole exome sequencing. *Genome biology* **18,** 22 (2017).
26.   Meers, P., Ali, S., Erukulla, R. & Janoff, A. S. Novel inner monolayer fusion assays reveal
differential monolayer mixing associated with cation-dependent membrane fusion.
Biochimica et Biophysica Acta (BBA) - Biomembranes **1467,** 227–243 (2000).
27.   Park, J.-S., Thorsness, M. K., Policastro, R., McGoldrick, L. L., Hollingsworth, N. M.,
Thorsness, P. E. & Neiman, A. M. Yeast Vps13 promotes mitochondrial function and is
localized at membrane contact sites. *MBoC* **27,** 2435–2449 (2016).
28.   Stenmark, H., Aasland, R. & Driscoll, P. C. The phosphatidylinositol 3-phosphate-
binding FYVE finger. *FEBS Letters* **513,** 77–84 (2002).
29.   Zoncu, R., Perera, R. M., Balkin, D. M., Pirruccello, M., Toomre, D. & De Camilli, P. A
Phosphoinositide Switch Controls the Maturation and Signaling Properties of APPL
Endosomes. *Cell* **136,** 1110–1121 (2009).
30.   Nakamura, N., Rabouille, C., Watson, R., Nilsson, T., Hui, N., Slusarewicz, P., Kreis, T. E.
& Warren, G. Characterization of a cis-Golgi matrix protein, GM130. *Journal of Cell Biology*
131, 1715–1726 (1995).
31.   Lefrancois, S., Zeng, J., Hassan, A. J., Canuel, M., Morales, C. R. The lysosomal
trafficking of sphingolipid activator proteins (SAPs) is mediated by sortilin. *The EMBO*
Journal **22,** 6430–6437 (2003).
32.   Luzio, J. P., Brake, B., Banting, G., Howell, K. E., Braghetta, P. & Stanley, K. K.
Identification, sequencing and expression of an integral membrane protein of the trans-Golgi
network (TGN38). *Biochem J* **270,** 97–102 (1990).

33.   Wakana, Y., van Galen, J., Meissner, F., Scarpa, M., Polishchuk, R. S., Mann, M. &
Malhotra, V. A new class of carriers that transport selective cargo from the trans Golgi
network to the cell surface. *The EMBO Journal* **31,** 3976–3990 (2012).
34.   Petersen, C. M., Nielsen, M. S., Nykjær, A., Jacobsen, L., Tommerup, N., Rasmussen, H.
H., Røigaard, H., Gliemann, J., Madsen, P. & Moestrup, S. K. Molecular Identification of a
Novel Candidate Sorting Receptor Purified from Human Brain by Receptor-associated
Protein Affinity Chromatography. *J. Biol. Chem.* **272,** 3599–3605 (1997).
35.   Porat-Shliom, N., Kloog, Y. & Donaldson, J. G. A Unique Platform for H-Ras Signaling
Involving Clathrin-independent Endocytosis. *MBoC* **19,** 765–775 (2008).
36.   Milovanovic, D., Wu, Y., Bian, X. & De Camilli, P. A liquid phase of synapsin and lipid
vesicles. *Science* **361,** 604–607 (2018).
37.   Milovanovic, D. & Camilli, P. D. Synaptic Vesicle Clusters at Synapses: A Distinct Liquid
Phase? *Neuron* **93,** 995–1002 (2017).
38.   Park, D., Wu, Y., Lee, S.-E., Kim, G., Jeong, S., Milovanovic, D., Camilli, P. D. & Chang,
S. Cooperative function of synaptophysin and synapsin in the generation of synaptic vesicle-
like clusters in non-neuronal cells. *Nature Communications* **12,** 263 (2021).
39.   Wheeler, R. J. & Hyman, A. A. Controlling compartmentalization by non-membrane-
bound organelles. *Philosophical Transactions of the Royal Society B: Biological Sciences*
373, 20170193 (2018).
40.   Willems, J., Jong, A. P. H. de, Scheefhals, N., Mertens, E., Catsburg, L. A. E., Poorthuis,
R. B., Winter, F. de, Verhaagen, J., Meye, F. J. & MacGillavry, H. D. ORANGE: A
CRISPR/Cas9-based genome editing toolbox for epitope tagging of endogenous proteins in
neurons. *PLOS Biology* **18,** e3000665 (2020).
41.   Waguri, S., Dewitte, F., Le Borgne, R., Rouillé, Y., Uchiyama, Y., Dubremetz, J.-F. &
Hoflack, B. Visualization of TGN to Endosome Trafficking through Fluorescently Labeled
MPR and AP-1 in Living Cells. *MBoC* **14,** 142–155 (2003).
42.   Lin, S. X., Mallet, W. G., Huang, A. Y. & Maxfield, F. R. Endocytosed Cation-
Independent Mannose 6-Phosphate Receptor Traffics via the Endocytic Recycling
Compartment en Route to the trans-Golgi Network and a Subpopulation of Late Endosomes.
MBoC **15,** 721–733 (2003).
43.   Carter, A. P., Diamant, A. G. & Urnavicius, L. How dynein and dynactin transport cargos:
a structural perspective. *Current Opinion in Structural Biology* **37,** 62–70 (2016).
44.   Rapali, P., Radnai, L., Süveges, D., Harmat, V., Tölgyesi, F., Wahlgren, W. Y., Katona, G.,
Nyitray, L. & Pál, G. Directed Evolution Reveals the Binding Motif Preference of the
LC8/DYNLL Hub Protein and Predicts Large Numbers of Novel Binders in the Human
Proteome. *PLOS ONE* **6,** e18818 (2011).
45.   Wang, Y., Huynh, W., Skokan, T. D., Lu, W., Weiss, A. & Vale, R. D. CRACR2a is a
calcium-activated dynein adaptor protein that regulates endocytic traffic. *Journal of Cell
Biology* **218,** 1619–1633 (2019).
46.   Shintani, M., Tada, M., Kobayashi, T., Kajiho, H., Kontani, K. & Katada, T.
Characterization of Rab45/RASEF containing EF-hand domain and a coiled-coil motif as a
self-associating GTPase. *Biochemical and Biophysical Research Communications* **357,** 661–
667 (2007).
47.   Maeda, S., Otomo, C. & Otomo, T. The autophagic membrane tether ATG2A transfers
lipids between membranes. *eLife* **8,** e45777 (2019).
48.   Guardia, C. M., Tan, X.-F., Lian, T., Rana, M. S., Zhou, W., Christenson, E. T., Lowry, A.
J., Faraldo-Gómez, J. D., Bonifacino, J. S., Jiang, J. & Banerjee, A. Structure of Human
ATG9A, the Only Transmembrane Protein of the Core Autophagy Machinery. *Cell Reports*
31, (2020).

49.   Ghanbarpour, A., Valverde, D. P., Melia, T. J. & Reinisch, K. M. A model for a partnership of lipid transfer proteins and scramblases in membrane expansion and organelle biogenesis. *PNAS* **118,** (2021).

50.   Matoba, K., Kotani, T., Tsutsumi, A., Tsuji, T., Mori, T., Noshiro, D., Sugita, Y., Nomura, N., Iwata, S., Ohsumi, Y., Fujimoto, T., Nakatogawa, H., Kikkawa, M. & Noda, N. N. Atg9 is a lipid scramblase that mediates autophagosomal membrane expansion. *Nat Struct Mol Biol* **27,** 1185–1193 (2020).

51.   Reinisch, K. M., Chen, X.-W. & Melia, T. J. "VTT"-Domain Proteins VMP1 and TMEM41B Function in Lipid Homeostasis Globally and Locally as ER Scramblases. *Contact* **4,** 251525642110244 (2021).

52.   Claude-Taupin, A., Jia, J., Bhujabal, Z., Garfa-Traoré, M., Kumar, S., da Silva, G. P. D., Javed, R., Gu, Y., Allers, L., Peters, R., Wang, F., da Costa, L. J., Pallikkuth, S., Lidke, K. A., Mauthe, M., Verlhac, P., Uchiyama, Y., Salemi, M., Phinney, B., Tooze, S. A., Mari, M. C., Johansen, T., Reggiori, F. & Deretic, V. ATG9A protects the plasma membrane from programmed and incidental permeabilization. *Nat Cell Biol* 1–13 (2021).

53.   Wandinger-Ness, A. & Zerial, M. Rab Proteins and the Compartmentalization of the Endosomal System. *Cold Spring Harbor Perspectives in Biology* **6,** a022616–a022616 (2014).

54.   Simonsen, A., Gaullier, J. M., D'Arrigo, A. & Stenmark, H. The Rab5 effector EEA1 interacts directly with syntaxin-6. *The Journal of biological chemistry* **274,** 28857–28860 (1999).

55.   Bock, J. B., Lin, R. C. & Scheller, R. H. A new syntaxin family member implicated in targeting of intracellular transport vesicles. *The Journal of biological chemistry* **271,** 17961–17965 (1996).

56.   Ganley, I. G., Espinosa, E. & Pfeffer, S. R. A syntaxin 10-SNARE complex distinguishes two distinct transport routes from endosomes to the trans-Golgi in human cells. *The Journal of cell biology* **180,** 159–172 (2008).

57.   Rapali, P., Szenes, Á., Radnai, L., Bakos, A., Pál, G. & Nyitray, L. DYNLL/LC8: a light chain subunit of the dynein motor complex and beyond. *The FEBS Journal* **278,** 2980–2996 (2011).

58.   Pérez-Victoria, F. J., Mardones, G. A. & Bonifacino, J. S. Requirement of the Human GARP Complex for Mannose 6-phosphate-receptor-dependent Sorting of Cathepsin D to Lysosomes. *Molecular biology of the cell* **19,** 2350–2362 (2008).

59.   Gershlick, D. C., Ishida, M., Jones, J. R., Bellomo, A., Bonifacino, J. S. & Everman, D. B. A neurodevelopmental disorder caused by mutations in the VPS51 subunit of the GARP and EARP complexes. *Human Molecular Genetics* **28,** 1548–1560 (2019).

60.   Guillén-Samander, A., Leonzino, M., Hanna, M. G., IV, Tang, N., Shen, H. & De Camilli, P. VPS13D bridges the ER to mitochondria and peroxisomes via Miro. *Journal of Cell Biology* **220,** (2021).

61.   Murphy, S. E. & Levine, T. P. VAP, a Versatile Access Point for the Endoplasmic Reticulum: Review and analysis of FFAT-like motifs in the VAPome. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1861,** 952–961 (2016).

62.   Cabukusta, B., Berlin, I., van Elsland, D. M., Forkink, I., Spits, M., de Jong, A. W. M., Akkermans, J. J. L. L., Wijdeven, R. H. M., Janssen, G. M. C., van Veelen, P. A. & Neefjes, J. Human VAPome Analysis Reveals MOSPD1 and MOSPD3 as Membrane Contact Site Proteins Interacting with FFAT-Related FFNT Motifs. *Cell Reports* **33,** 108475 (2020).

63.   Gómez-Sánchez, R., Rose, J., Guimarães, R., Mari, M., Papinski, D., Rieter, E., Geerts, W. J., Hardenberg, R., Kraft, C., Ungermann, C. & Reggiori, F. Atg9 establishes Atg2-dependent contact sites between the endoplasmic reticulum and phagophores. *The Journal of cell biology* jcb.201710116 (2018).

64.   Davies, A. K., Itzhak, D. N., Edgar, J. R., Archuleta, T. L., Hirst, J., Jackson, L. P.,
Robinson, M. S. & Borner, G. H. H. AP-4 vesicles contribute to spatial control of autophagy
via RUSC-dependent peripheral delivery of ATG9A. *Nature Communications* **9,** 3958 (2018).
65.   Pérez-Victoria, F. J., Schindler, C., Magadán, J. G., Mardones, G. A., Delevoye, C.,
Romao, M., Raposo, G. & Bonifacino, J. S. Ang2/Fat-Free Is a Conserved Subunit of the
Golgi-associated Retrograde Protein Complex. *MBoC* **21,** 3386–3395 (2010).
66.   Fröhlich, F., Petit, C., Kory, N., Christiano, R., Hannibal-Bach, H.-K., Graham, M., Liu,
X., Ejsing, C. S., Farese, R. V. & Walther, T. C. The GARP complex is required for cellular
sphingolipid homeostasis. *eLife* **4,** (2015).
67.   Khakurel, A., Kudlyk, T., Bonifacino, J. S. & Lupashin, V. V. The Golgi-associated
retrograde protein (GARP) complex plays an essential role in the maintenance of the Golgi
glycosylation machinery. *MBoC* mbc.E21-04-0169 (2021).
68.   Lesage, S., Drouet, V., Majounie, E., Deramecourt, V., Jacoupy, M., Nicolas, A., Cormier-
Dequaire, F., Hassoun, S. M., Pujol, C., Ciura, S., Erpapazoglou, Z., Usenko, T., Maurage, C.-
A., Sahbatou, M., Liebau, S., Ding, J., Bilgic, B., Emre, M., Erginel-Unaltuna, N., Guven, G.,
Tison, F., Tranchant, C., Vidailhet, M., Corvol, J.-C., Krack, P., Leutenegger, A.-L., Nalls, M.
A., Hernandez, D. G., Heutink, P., Gibbs, J. R., Hardy, J., Wood, N. W., Gasser, T., Durr, A.,
Deleuze, J.-F., Tazir, M., Destée, A., Lohmann, E., Kabashi, E., Singleton, A., Corti, O., Brice,
A., (PDG), F. P. D. G. S. & (IPDGC), I. P. D. G. C. Loss of VPS13C Function in Autosomal-
Recessive Parkinsonism Causes Mitochondrial Dysfunction and Increases PINK1/Parkin-
Dependent Mitophagy. *American journal of human genetics* **98,** 500–513 (2016).
69.   Schauder, C. M., Wu, X., Saheki, Y., Narayanaswamy, P., Torta, F., Wenk, M. R., Camilli,
P. D. & Reinisch, K. M. Structure of a lipid-bound extended synaptotagmin indicates a role in
lipid transfer. *Nature* **510,** 552–555 (2014).
70.   Bian, X., Saheki, Y. & Camilli, P. D. Ca2+releases E-Syt1 autoinhibition to couple ER-
plasma membrane tethering with lipid transport. *The EMBO journal* **37,** 219–234 (2018).
71.   Zivanov, J., Nakane, T. & Scheres, S. H. W. Estimation of high-order aberrations and
anisotropic magnification from cryo-EM data sets in RELION-3.1. *IUCrJ* **7,** 253–267 (2020).
72.   Zheng, S. Q., Palovcak, E., Armache, J.-P., Verba, K. A., Cheng, Y. & Agard, D. A.
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. *Nat Methods* **14,** 331–332 (2017).
73.   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *Journal of Structural Biology* **192,** 216–221 (2015).
801
802

	FIGURE LEGENDS
804
Figure 1. In vitro characterization of SHIP164.
(A) The fold prediction algorithm Alphafold indicates that SHIP164 harbors an extended ß-sheet
folded to resemble a taco-shell, as in VPS13[8]. From left to right: (a) Ribbons representation for
the taco-shell "core", predicted with high confidence (b) the extended ß-sheet is shown in two
orientations as surface representation, with carbons shown white, oxygens red, nitrogens blue,
and sulfurs yellow. The concave surface of the taco-shell is predicted to be entirely hydrophobic,
providing a binding site for lipid fatty acid moieties. (c) additional secondary structure that may
associate with the "core" is shown; segments discussed in text are indicated.
(B) SHIP164 co-purifies with phospholipids according to their abundance in cells as assessed by
shotgun lipidomics.
(C) SHIP164 co-migrates with NBD-PE in a native gel. Comparison of NBD-fluorescence that co-
migrates with SHIP164 versus with the well-characterized lipid transport module from E-Syt2
suggests that each SHIP164 molecule accomodates multiple (~8) phospholipids.
(D) Schematic drawing explaining the FRET-based lipid transfer and the dithionite-based control
for the absence of liposome fusion.
(E) Assay consistent with transfer of fluorescent lipids from donor to acceptor liposomes by
SHIP164 but not by a tether construct lacking a lipid transfer module. An increase in fluorescence
observed with donor liposomes only (red arrow) is due to lipid extraction by SHIP164. A larger
increase is observed when lipids are subsequently transferred to the acceptor liposomes.
(F) Dithionite addition excludes that SHIP164 facilitates membrane fusion.
(G) 2D class averages from negatively stained SHIP164 and MBP-SHIP164, showing tail-to-tail
dimerization. MBPs (arrows) are at the ends of the elongated dimer.
(H) 2D class averages for the cryo-EM reconstruction show a channel running along the length
of SHIP164.
(I) Cryo-EM map of SHIP164 at a resolution of ~8.3 Å, showing a long central cavity with or
without the Alphafold model (see panel A-a) superimposed onto it. The docking is approximate
because the resolution of the reconstruction is low and because a high confidence model for
SHIP164, beyond the ß -sheet taco shell core, is not available.
Experiments in panels E-F were performed in triplicate; SD's are indicated.
834
Figure 2. Localization of exogenous SHIP164 points to a role on endocytic
organelles.
(A) Live fluorescence (inverted grays) image of a COS-7 cell expressing exogenous
SHIP164^mScarlet-I. Scale bar, 20nm. High magnification of the indicated regions is shown at
right. Scale bars, 5μm.
(B) Live fluorescence (inverted grays) image of a COS-7 cell expressing exogenous 3xFlag-
SHIP164^mScarlet-I. Scale bar, 20μm.
(C) Live image of the cytoplasm of a COS-7 cell expressing exogenous SHIP164^mScarlet-I
(magenta) and the endosomal marker GFP-2xHrs[FYVE] (green). Arrows indicate SHIP164
accumulation juxtaposed to endosomal membrane. Scale bar, 5μm.
(D) High magnification live fluorescence images of COS-7 cell expressing exogenous
SHIP164^mScarlet-I (magenta) and the organelle marker indicated at the top (green). The
individual channels are shown as inverted grays. Scar bar, 2μm.
(E) Live image of the cytoplasm of a COS-7 cell expressing exogenous SHIP164-Halo, the
endosome marker EEA1-RFP, and the ER marker GFP-Sec61. Scale bar, 2μm. The individual
channels are shown as inverted grays. Scar bar, 1μm.
(F) Time-series of live fluorescence images of exogenous SHIP164-mRFP (magenta) and EEA1-
GFP (green). Arrowheads point to a SHIP164 accumulation as it undergoes fission between two
EEA1 positive organelles. Time, seconds. Scale bar, 1μm.

(G) Fluorescence images of a COS-7 cell expressing exogenous SHIP164^mScarlet-I (magenta) and immunolabeled with antibodies against MPR (green) and GM130 (red). Scale bar, 20μm. The individual channels are shown as inverted grays in the higher magnification of the Golgi complex region shown at right. Arrowheads indicate colocalization of exogenous SHIP164 and endogenous MPR. Scale bar, 5μm.

(H-I) Fluorescence images of RPE-1 cells demonstrating normal Golgi complex localization of MPR in a cell expressing RFP alone (H) and its abnormal localization in a cell expressing both RFP and SHIP164 (H). Cells were immunolabeled with antibodies against MPR (green) and GM130 (magenta). Both the merge image and the single channels are shown. Arrowheads indicate ectopic accumulations of MPR in cells over-expressing SHIP164. Scale bar, 5μm.

Figure 3. Correlative light electron microscopy (CLEM) reveals that foci of exogenous SHIP164 reflect large accumulations of small vesicles.

(A) SHIP164^mScarlet-positive structures juxtaposed to large GFP-WDFY2-positive endosome compartments in a COS-7 cell also expressing untagged Ras^{G12V} to induce micropinosomes. The two EM images (scale bar, 200nm) correspond to the regions framed by two rectangles in the merge fluorescence image (scale bar, 1μm). SHIP164 fluorescence reflects clusters of small vesicles separated from the endosomes by a band occupied by a dense matrix (arrows).

(B and C) High power view of endosomes of COS-7 cells expressing exogenous SHIP164^mScarlet-I (magenta), GFP-WDFY2 (green), and untagged Ras^{G12V}. Merge fluorescence images are shown at left and FIB-SEM based reconstructions are shown at right. Scale bar, 1μm.

Figure 4. Localization of Endogenous SHIP164 on small clusters of vesicular structures.

(A) Cartoon of wild-type (top) and edited SHIP164^mNG (bottom).

(B) Western blot of control and edited cell clones for SHIP164, mNG, and GAPDH as a loading control. The different motility of SHIP164 in control and edited clone indicates the incorporation of mNG in the edited SHIP164.

(C) Cartoon of endogenously edited locus of SHIP164^mNG in HeLa cell. Red, DNA scar as a result of HITI; blue, small Gly-Ser linkers; green, mNG sequence (shortened for brevity by forward slashes).

(D) Fluorescence image of endogenous SHIP164^mNG (inverted greys) in an edited HeLa knock-in cell. Arrowheads indicate concentration of endogenous SHIP164 at the cell edge, shown at a higher magnification in the inset. Scale bar, 10μm (main field) and 2μm(inset).

(E) Single channel (inverted grays) fluorescence images of endogenous SHIP164^mNG and endogenous MPR-mScarlet-I in a HeLa double-knock-in cell demonstrating partial colocalization of the two proteins at the cell edge and the selective concentration of MPR in perinuclear area. Scale bar, 10μm. High magnification of the boxed regions is shown at right where arrowheads indicate overlapping fluorescence. Scale bar, 2μm.

(F) Left to right: Fixed fluorescence image of endogenously tagged SHIP164^mNG (magenta) HeLa cell also expressing mito-BFP (green) for alignment in CLEM analysis. Scale bar, 5μm. Single plane of area outlined in yellow. Fluorescence overlay with FIB-SEM image: numbered boxes mark region of interest of eSHIP164^mNG fluorescence to be aligned with EM image. Scale bar of magnification area, 1μm. Magnification of cluster of vesicles aligned with eSHIP164^mNG fluorescence marked by numbered boxes. Scale bar of EM area, 100nm.

(G) Reconstruction of cluster 1 from panel F above. SHIP164 associated endosome structures, magenta; ER tubules, yellow.

(H) Fluorescence image of endogenously tagged SHIP164^mNG in a HeLa knock-in cell also expressing the ER marker RFP-Sec61 (green) demonstrating some SHIP164 foci are in proximity of ER tubules. Scale bar, 1μm.

Figure 5. Interaction of SHIP164 with proteins implicated in retrograde traffic to the Golgi complex.
(A) Live fluorescence image of a COS-7 cell demonstrating partial colocalization of exogenous SHIP164^mScarlet-I (magenta) and mNeonGreen-Stx6 (green) in the Golgi complex area. Scale bar, 10μm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 2μm.
(B) Mass spectrometry-based identification of material affinity-purified onto immobilized SHIP164 or onto a control protein from mouse brain extract. The selective enrichment of DYNLL1/2 peptides on the SHIP164 bait is shown.
(C) A mixture of MBP-SHIP164$_{1312-1464}$ and DYNLL1/2 was subjected to size exclusion chromatography. Coomassie blue staining of SDS-PAGE of the eluted fractions reveals co-migration of the two proteins.
(D) Live fluorescence images (inverted grays) of COS-7 cells expressing either GFP-Rab45 (left) or SHIP164^mScarlet-I (right), or both proteins together (only SHIP164 is shown) as indicated. Scale bar, 10μm. Note the massive recruitment of SHIP164 to the centrosomal region in the presence of Rab45. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Scale bar, 2μm.
(E) Cartoon of wild-type SHIP164 highlighting (blue) newly identified Stx6 (L$_{759}$xxYY$_{763}$) and DYNLL1/2 (V$_{1395}$TQATQT$_{1401}$) interaction motifs (see also Fig. 1Aa).

Figure 6. Defects in retrograde membrane traffic to the TGN in SHIP164 knockout cells.
(A) Western blot of control and edited RPE1 cell clones for SHIP164 and for GAPDH as a loading control.
(B) Left: fluorescence images of a parental control (left) and SHIP164 KO cells (right) immunolabeled with antibodies against EEA1 (green) and GM130 (magenta). Scale bar, 20μm. Right: Quantification of large EEA1 endosomes (>1μm^2) per cell in control and SHIP164 KO cell clones. Middle line, median; +, mean; bars, 10-90 percentiles. Data represents three biological replicates. **, $p<0.01$.
(C) Live fluorescence (inverted grays) image of parental control (top) and SHIP164 KO (bottom) cells expressing the endosomal marker GFP-2xHrs^{FYVE}. Scale bar, 20μm.
(D) Left: fluorescence images of parental control (left) and SHIP164 KO cells (right) immunolabeled with antibodies against MPR (green) and GM130 (magenta). Scale bar, 5μm. Right: Quantification of scattered cytoplasmic spots-to-Golgi complex ratio of MPR signal per cell in control and SHIP164 KO cell clones. Middle line, mean; bars, SD. **, $p<0.01$.
(E) Left: fluorescence images of parental control (left) and SHIP164 KO cells (right) immunolabeled with antibodies against TGN46 (green) and GM130 (magenta). Scale bar, 5μm. Right: Quantification of scattered cytoplasmic spots-to-Golgi complex ratio of TGN46 signal per cell in control and SHIP164 KO cell clones. Middle line, mean; bars, SD. **, $p<0.01$.
(F) Left: fluorescence images of parental control (left) and SHIP164 KO cells (right) immunolabeled with antibodies against CHC (green) and GM130 (magenta). Scale bar, 5μm. Right: Quantification of scattered cytoplasmic spots-to-Golgi complex ratio of CHC signal per cell in control and SHIP164 KO cell clones. Middle line, mean; bars, SD. **, $p<0.01$.
(G) Left: fluorescence images of SHIP164 KO cells expressing RFP alone (left) or both RFP and SHIP164 (right) immunolabeled with antibodies against TGN46 (green) and GM130 (magenta). Scale bar, 5μm. Right: Quantification of scattered cytoplasmic spots-to-Golgi complex ratio of TGN46 signal per cell in SHIP164 KO cell clones expressing RFP alone (solid circles) or both RFP and SHIP164 (open circles). Middle line, mean; bars, SD. **, $p<0.01$; *, $p<0.5$.
Data of panels D-F reflect three biological replicates and of panel G two biological replicates.

Figure 7. Selective properties of SHIP164 vesicle clusters.
(A) Live fluorescence (inverted grays) image of a COS-7 cell expressing exogenous ATG9A-GFP. Scale bar, 10nm.
(B) Live fluorescence (inverted greys) image of a COS-7 cell demonstrating the clustering of exogenous ATG9A-GFP (left) fluorescence in the Golgi complex area from SHIP164^mScarlet-I expression (right). Scale bar, 10µm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 1µm.
(C) Impact of the expresion of SHIP164 on the localization of ATG9A in cells also overexpressing Rab45. Cells were transfected with both Rab45 and ATG9A without (left) and with (right) the additional expression of SHIP164. Note that in the presence of SHIP164, Rab45 and ATG9A colocalize in a perinuclear spot. Left, live fluorescence image of a COS-7 cell demonstrating no change to the localization of exogenous ATG9A-GFP (green) upon expression of RFP-Rab45 (magenta) and Right, fluorescence image of a COS-7 cell demonstrating colocalization of exogenous ATG9A-GFP and RFP-Rab45 upon the expression of SHIP164-Halo (not shown) in the centrosomal region. Scale bar, 10µm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 2µm.
(D) Live fluorescence (inverted grays) image of a COS-7 cell expressing exogenous UHRF1BP1-Halo. Scale bar, 20nm. High magnification of the indicated regions is shown at right. Scale bar, 5µm.
(E) Live fluorescence image of a COS-7 cell demonstrating partial colocalization of exogenous ATG9A-GFP (green) and UHRF1BP1-Halo (magenta) fluorescence. Scale bar, 10µm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Scale bar, 2µm.

Supplemental Figure 1.
In vitro characterization of SHIP164. **(A)**(a) Following batch purification using anti-FLAG M2 resin, 3XFLAG-SHIP164$_{\Delta901-1099}$ was further purified by size exclusion chromatography on a Superdex-200 column. The gel filtration profile is shown (black), along with that for molecular weight standards (light blue). (b) Sample purity was examined by SDS-PAGE. (c) Representative negative stain electron micrograph of 3XFLAG-SHIP164$_{\Delta901-1099}$ (50 nM) using FEI Tecnai 12 microscope at 120 kV at a nominal magnification of 52,000×. Staining was with 2% uranyl acetate on carbon-coated copper grids that were glow-discharged for 30 s at 22 mA. Green circles represent particles manually picked with a 300 Å mask diameter in RELION-3.1.
(B) Cryo-EM workflow. Micrographs, including the representative one shown, were collected using Titan Krios G2 transmission electron microscope (Thermo Fisher Scientific) at 300 kV equipped with a K3 summit direct detection camera (Gatan). Green circles represent particles manually picked for initial reference-free 2D classification prior to autopicking.
(C) The processing workflow yielded a reconstruction with an estimated resolution of 8.3 Å according to the Fourier shell correlation (FSC) = 0.143 criterion.

Supplemental Figure 2.
(A) Live fluorescence images of a HeLa cell expressing exogenous GFP-SHIP164 (green) and soluble RFP-Stx6 1-234 (top) or full-length RFP-Stx6 (bottom; magenta). Individual channels are shown in inverted grays on the left. Scale bar, 10µm.
(B) High power views of micropinosomes of a COS-7 cell expressing exogenous SHIP164^mScarlet-I (magenta) and the early endosomal marker GFP-WDFY2 (green). Untagged Ras^{G12V} was also expressed to induce macropinosome formation. The individual channels are shown as inverted grays. Scar bar, 1µm.

(C) Live fluorescence image of a COS-7 cell expressing SHIP164-Halo (magenta) and the dominant negative GFP-Rab5 S34N mutant. Scale bar, 20μm.

(D) Fluorescence images of wild-type HeLa cells immunolabeled with antibodies against MPR (green) and GM130 (red) and without (top) or with transfection of exogenous SHIP164^mScarlet-I (magenta; bottom) demonstrating abnormal localization of MPR in the cell expressing SHIP164. The individual channels are shown as inverted grays. Arrowheads indicate colocalization of exogenous SHIP164 and endogenous MPR. Scale bar, 5μm.

(E and F) Fluorescence images of RPE-1 cells demonstrating normal Golgi complex localization of sortilin in a cell expressing RFP alone (E) and its abnormal localization in a cell expressing both RFP and SHIP164 (F). Cells were immunolabeled with antibodies against sortilin (green) and GM130 (magenta). Both the merge image and the single channels are shown. Arrowheads indicate ectopic accumulations of sortilin in cells over-expressing SHIP164. Scale bar, 5μm.

(G and H) Fluorescence images of RPE-1 cells demonstrating normal Golgi complex localization of TGN46 in a cell expressing RFP alone (G) and its unaffected localization in a cell expressing both RFP and SHIP164 (H). Cells were immunolabeled with antibodies against TGN46 (green) and GM130 (magenta). Both the merge image and the single channels are shown. Scale bar, 5μm.

Supplemental Figure 3.
High magnification live fluorescence images of COS-7 cell expressing exogenous SHIP164 1-873-mRFP (magenta) and either the endosome marker WDFY2 or the ER protein Sec61b (green). The individual channels are shown as inverted grays. Scar bar, 2μm.

Supplemental Figure 4.
(A) Merge fluorescence image of endogenous SHIP164^mNG (green) and exogenous SHIP164^mScarlet-I (magenta) in an edited HeLa knock-in cell to demonstrate overlap of the two fluorescence signals. Scale bar, 10μm. High magnification of the indicated region is shown at right where the individual channels are shown as inverted grays. Scale bar, 2μm.

(B) Merge fluorescence image of endogenous MPR-mScarlet-I and endogenous SHIP164^mNG in a HeLa double-knock-in cell demonstrating partial colocalization of the two fusion proteins at the cell edge but not with pulsed EGF-647 (blue) foci. Scale bar, 10μm. High magnification of the indicated region is shown at right and the individual channels are shown as inverted grays. Open arrowheads indicate overlapping fluorescence and blue arrowheads an EGF-647-positive endocytic structure. Scale bar, 2μm.

(C) Live fluorescence image of a COS-7 cell expressing FAM174a-mRFP (inverted grays) and GFP-WDFY2 (not shown). Scale bar, 20μm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 2μm.

(D) Live fluorescence image of a COS-7 cell demonstrating partial colocalization of exogenous SHIP164-GFP (green) FAM174a-mRFP (magenta) fluorescence in the cytoplasm. Scale bar, 10μm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 2μm.

Supplemental Figure 5.
(A) Live fluorescence images of a HeLa cell expressing exogenous GFP-SHIP164 Y772A, Y773A (magenta) and RFP-Stx6 (green) showing loss of Stx6 binding by this SHIP164 mutant. Individual channels are shown in inverted grays on the left. Scale bar, 10μm.

(B) High magnification live fluorescence images COS-7 cell expressing exogenous SHIP164 Y772A, Y773A-mRFP and endosome marker EEA1 or the ER marker Sec61b showing that this SHIP164 mutant still accumulates at hot spots. The individual channels are shown as inverted grays. Scar bar, 2μm.

(C) Live fluorescence images (inverted grays) of wild-type HeLa cell expressing RFP-Rab45, r endogenous SHIP164^mNG in a HeLa knock-in cell, or HeLa knock-in cell expressing RFP-Rab45, as indicated. Scale bar, 10μm. Note the massive recruitment of endogenous SHIP164 to the centrosomal region in the presence of Rab45. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Scale bar, 2μm.

(D) Impact of the expresion of exogenous SHIP164 on the localization of endogenous MPR-mNG in cells also overexpressing Rab45. HeLa eMPR-mNG knock-in cells were transfected with Rab45 without (left) and with (right) the additional expression of SHIP164. Note that in the presence of SHIP164, Rab45 and MPR colocalize in a perinuclear spot. Left, live fluorescence image of a HeLa knock-in cell demonstrating no change to the localization of endogenous MPR-mNG (green) upon expression of RFP-Rab45 (magenta) and Right, fluorescence image of a HeLa knock-in cell demonstrating colocalization of endogenous MPR-mNG and RFP-Rab45 upon the expression of SHIP164-Halo (not shown) in the centrosomal region. Scale bar, 10μm. High magnification of the indicated regions is shown at right and the individual channels are shown as inverted grays. Arrowheads indicate overlapping fluorescence. Scale bar, 2μm.

(E) Live fluorescence image of a COS-7 cell expressing SHIP164^mScarlet-I (magenta) and the dominant negative GFP-Rab45 S555N mutant (green) showing that this Rab45 mutant does not recruit SHIP164. Individual channel shown in inverted gray. Scale bar, 20μm.

Supplemental Figure 6.

 (A) Sequencing results confirming editing of SHIP164 locus in RPE1 cells. PAM sequence (green) was directed to exon 2 and each clone was found to have a mutation starting four nucleotides upstream. # Indicates nucleotide is the same between parental control and clone 2.

(B) Left: Fluorescence images of a parental control (left) and SHIP164 KO cells (right) immunolabeled with antibodies against sortilin (green) and GM130 (magenta). Scale bar, 5μm. Right: Quantification of the scattered cytoplasmic spots-to-Golgi complex ratio of Sortilin signal per cell (circles) in control and SHIP164 KO cell clones. Middle line, mean; bars, SD. Data represents three biological replicates. **, p<0.01.

(C) Fluorescence images of a SHIP164 KO cell immunolabeled with antibodies against TGN46 (green) and MPR (magenta) demonstrating distinct post-Golgi vesicle populations. Scale bar, 5μm. High magnifications of the indicated regions are shown as insets and the individual channels are shown as inverted grays. Arrowheads indicate divergent fluorescence. Scale, 2μm.

(D) Fluorescence images of a SHIP164 KO cells expressing both RFP and SHIP164 (not shown) immunolabeled with antibodies against the indicated proteins (green) and GM130 (magenta). Scale bar, 5μm.

(E) Fluorescence images of parental control or SHIP164 KO cells expressing either RFP alone (not shown) or RFP and exogenous untagged SHIP164 (not shown) immunolabeled with antibodies against the EEA1 (green) demonstrating that overexpression of SHIP164, even in KO cells, reuslts in EEA1 accumulations not typically found in wild-type cells where SHIP164 is not over-expressed (far left). Scale bar, 10μm.

(F) Western blot of RPE1 cells for SHIP164 and for GAPDH as a loading control, either treated with control or SHIP164 specific siRNAs.

(G) Left: Fluorescence images of a wild-type cells treated with control (left) or SHIP164 specific siRNAs (right) immunolabeled with antibodies against EEA1 (green). Scale bar, 20μm. Right, Quantification of large EEA1 endosomes (>1μm^2) per cell in control and SHIP164 knock-down cell. Middle line, median; +, mean; bars, 10-90 percentiles. Data represents three biological replicates. **, p<0.01.

Supplemental Figure 7.

High magnification live fluorescence image COS-7 cell expressing exogenous UHRF1BP1-Halo (magenta) and the lysosomal marker LAMP1-GFP (green). The individual channels are shown as inverted grays. Scar bar, 2μm.

Supplemental movie 1
High magnification of live fluorescence time series of COS-7 cell expressing exogenous SHIP164-mRFP (magenta) and the endosomal marker EEA1-GFP (green) demonstrating the dynamics of SHIP164 accumulations. Frames captured every 2s. Scale bar, 1μm.

Supplemental movie 2
Zoom of live fluorescence time series of a double-knock-in HeLa cell demonstrating the colocalization of fluorescencnt signal from endogenous MPR-mScarlet-I (magenta; left) and endogenous SHIP164^mNG (green; right) puncta at the cell periphery. Arrowheads indicate puncta positive for both mScarlet-I and mNG fluorescence. Frames captured every 2s. Scale bar, 10μm.

MATERIALS AND METHODS

Antibodies and Reagents

The list of antibodies, their working dilution, and the supplier for this study can be found in **Table S1**. All Halo-tag ligands used (JF549 and JF646) were a kind gift from Luke Lavis (Janelia Research Campus). All SHIP164 (UHRF1BP1L) ORFs used in this study utilized a human codon optimized sequence designed and purchased from Genscript (sequence available upon request). The following constructs were kind gifts: BFP-Rab5b and mito-BFP from G. Voeltz (University of Colorado-Boulder, CO; Addgene plasmids #49147, #49151, respectively); Sec61b-GFP and RFP from T. Rapoport (Harvard University, Cambridge, MA); GFP-2xHrs[FYVE] from Harald Stenmark (University of Oslo, Norway; Addgene plasmids #140047). Other plasmids encoding EEA1, WDFY2, APPL1, 2xFYVE were previously generated in the De Camilli lab. All other ORFs used are listed in **Table S2**.

All EM reagents were purchased from Electron Microscopy Sciences (Hatfield, PA).

Lipids were purchased from Avanti Polar Lipids: DOPC (850357), liver PE (840026), DGS-NTA (Ni; 790404), brain PI(4,5)P2 (840046), Rh-PE (810150), and NBD-PE (810145).

Generation of Plasmids

Codon optimized human SHIP164 generated by Genscript was amplified using PCR from the pUC57 plasmid and ligated into various mammalian and bacterial expression plasmids. Most constructs were generated with regular cloning protocols or through site-directed mutagenesis. The desired ORFs were amplified by PCR and inserted into plasmids through enzyme digestions and ligation. Some amplified ORFs were ligated using HiFi assembly (NEB). Details of primer sets, enzymes, techniques, and plasmids used for each construct can be found in **Table S2**.

For internal tagging of SHIP164, the codon optimized human sequence was first amplified by PCR and cloned into p3xFLAG between the EcoRI/KpnI restriction sites using HiFi assembly. From there, the ORF of mScarlet-I was amplified by PCR and ligated into the EcoRV site at residue 915 of SHIP164 using HiFi assembly. To generate the amino-terminal free version, SHIP164^mScarlet-I was amplified by PCR and ligated into pcDNA3.1 His-A between the HindIII/XhoI sites removing the His_6 tag. Unless stated otherwise, all mutations in this study were generated using site directed mutagenesis. All ORFs were sequenced in their entirety after cloning and before use in any and all experiments.

For the MBP-SHIP164 fusion construct used in the in vitro analysis, sequences coding for residues 27–392 of E. coli maltose binding protein (MBP) and two additional alanines were fused to the N-terminus of SHIP164$_{\Delta 901-1099}$ using Gibson Assembly and site-directed mutatgenesis.

The tethered SHIP164 construct used to perform the *in vitro* lipid transfer assay was generated using Gibson Assembly as previously described[6], and consists of SHIP164$_{\Delta 901-1099}$ followed by the unstructured region of human Esyt2(residues 649–689), then the pleckstrin homology domain of rat PLCΔ(residues 11–140), the unstructured region of Eyst2(residues 690–739), and a hexahistidine tag at the C-terminus.

Protein Expression and Purification

For lipid transfer and cryo-EM studies, 3×FLAG-SHIP164$_{\Delta 901-1099}$ constructs were transfected into Expi293F cells (Thermo Fisher Scientific) according to manufacturer instructions. Protein expression was enhanced by the addition of nonessential amino acids (Gibco) and valproic acid (3.5 mM final concentration) 16 h after transfection. Cells were harvested by centrifugation 65 h

after transfection and either flash-frozen for storage or used immediately for protein purification.
Cells were lysed by 3 freeze-thaw cycles in buffer (50 mM HEPES pH 8.0, 500 mM NaCl, 1 mM
TCEP, 5% glycerol) containing 1× complete EDTA-free protease inhibitor cocktail (Roche).
Insoluble debris was removed by centrifugation at 15000 rpm for 30 min, and clarified lysate was
mixed with anti-FLAG M2 resin (Sigma) for 3 h while rotating at 4°C. Resin was then washed with
3×10 bed volumes of buffer and subsequently incubated for 18 h at 4°C in buffer supplemented
with 1 mM ATP and 2 mM MgCl$_2$. Resin was washed again with 3 bed volumes of buffer and then
eluted in 5×0.5 bed volumes of buffer containing 0.25 mg/mL 3×FLAG peptide (APExBio). The
protein was further purified by gel filtration (Superdex 200 10/300 GL, GE Healthcare) and
concentrated to approximately 1.5 μM (Amicon). E-Syt2 was expressed and purified as previously
described[69].
1183
The MBP-SHIP164$_{1312-1464}$/DYNLL1 complex was recombinantly expressed in BL21 DE3
competent cells for purification and size exclusion chromatography analysis using the
polycistronic pET Duet expression plasmid, details of which can be found in **Table S2**. BL21 cells
transformed with the dual expression plasmid were grown at 37°C in super broth (SB) media (1L)
until reaching an OD$_{600}$ between 2 and 3, after which the temperature was turned down to 18°C
and cells were induced with 0.25mM IPTG for 16 hours with continuous shaking. Cells were
pelleted in a floor centrifuge at 4000g for 15 minutes and then resuspended in lysis buffer (25mM
Tris pH 7.5, 200mM NaCl, 0.1% Tween-20) and frozen at -80°C until ready for purification. For
purification, the pellet was thawed in a water bath and then sonicated until the desired lysis using
a macro sonicator tip (VirTris VirSonic). Membrane and un-sonicated material were pelleted in a
floor ultra-centrifuge at 24,000g for 20 minutes and the supernatant was collected and incubated
with 250uL of amylose resin for 1 hour rotating at 4°C. The resin was then washed three times
with 40x bed volumes of ice-cold wash buffer (25mM Tris pH 7.5, 200mM NaCl, 1mM BME). Two
elutions were performed from the resin using 2x bed volume of elution buffer (25mM Tris pH 7.5,
150mM NaCl, 10mM maltose, 1mM BME). The eluate was collected and run over a Superdex S200
column using a GE AKTA pure system, and individual fractions (1mL) were collected for analysis
by SDS-PAGE and Coomassie staining.
1201
Lipidomics Analysis of SHIP164 by Mass Spectrometry
Full length human 3xFLAG-SHIP164 was expressed and purified as described above with the
following modifications. Following immunoprecipitation, FLAG resin bound SHIP164 was
washed with 3x100 bed volumes of sonication buffer containing 1mM BME and protease inhibitor
cocktail for 20 minutes. After elution, the protein was buffer exchanged into 50mM mass
spectrometry grade ammonium acetate. The purified protein sample and 1x10^7 untransfected
Expi293F cells was sent to Avanti Polar Lipids for analysis. Lipids were extracted in 2:1 (v/v)
methanol:chloroform. The chloroform layer was dried, and lipids were reconstituted with internal
standards for PC, PE, PI, PS, PA, PG, SM, TAG, DAG, and CE for quantification. As described by
the company, the sample was injected into a reverse phase C8 column with a gradient elution
profile for the resolution required of each lipid class and detected by an AB Sciex 5500 tandem
mass spectrometer. The molecular species of lipids was quantified based on internal standards
and summed by lipid class.
1215
Native gel lipid binding assay.
Purified E-Syt2 and SHIP164$_{\Delta901-1099}$ at indicated concentrations were mixed with methanol-
dissolved NBD-PE (0.2 mM final concentration) in 20-μL total reaction volumes and incubated
for 2 h on ice. Samples were loaded on 4–20% Mini-PROTEAN TGX Precast Gels and run for 3 h
at 200 V. NBD fluorescence was visualized using an ImageQuant LAS4000 (GE Healthcare), and
gels were stained with Coomassie Brilliant Blue R (Sigma Aldrich) to visualize total protein. A

standard curve was generated using the fluorescence values of E-Syt2 and the quantity of lipid
bound to SHIP164 was analyzed using ImageJ (National Institutes of Health).
1224
1225
FRET-based lipid transfer assay.
Liposomes were prepared as previously described[6]. Briefly, lipids in chloroform were mixed in the
indicated ratios and dried to thin films, then resuspended in buffer containing 50 mM HEPES,
pH 8.0, 500 nM NaCl, 1 mM TCEP, and 5% glycerol to 1 mM total lipid concentration. Rehydrated
lipids were then incubated at 37°C for 1 h before being subjected to 10 freeze-thaw cycles. Crude
liposomes were freshly extruded through a polycarbonate filter with 100-nm pore size prior to
transfer assays.
1233
Lipid transfer experiments were set up in 100-μL reaction volumes containing 0.125 μM protein,
25 μM donor liposomes (61% DOPC, 30% liver PE, 2% NBD-PE, 2% rhodamine-PE, and 5% DGS-
NTA[Ni]) and 25 μM acceptor liposomes (65% DOPC, 30% liver PE, and 5% PI(4,5)P$_2$. Donor and
acceptor liposomes were included in the reaction for 5 minutes prior to the addition of protein to
establish a fluorescence baseline. The lipid transfer reaction was monitored via NBD fluorescence
intensity detected every 1 min at 538 nm upon excitation at 460nm. Lipid transfer assay was
performed for 1 h at 30°C in a 96-well plate (Nunc) using a Synergy H1 Plate Reader (BioTek).
1241
1242 A tether-only construct consisting of an N-terminal hexahistidine-tagged plexstrin homology
1243 domain from rat PLCΔ(11-140) was expressed and purified as previously described[70]. The lipid
1244 transfer assay using this construct was performed using the same protein:lipid ratio.
1245
1246 A dithionite assay, to rule out the possibility that the NBD fluorescence increase was due to
1247 SHIP164-mediated liposome fusion, was performed in the same way as the transfer assay, except
1248 for the final addition of 5 μL freshly prepared dithionite buffer [100 mM dithionite (Sigma
1249 Aldrich) in 50 mM Tris, pH 10] after fluorescence increase observed during the transfer assay had
1250 reached its maximum. NBD emission was then monitored for an additional 30 min.
1251
Cryo-EM sample preparation, data collection, and image processing
Purified SHIP164$_{\Delta 901-1099}$ was supplemented with 0.05% n-Octyl-β-D-glucoside immediately
before cryo-EM samples were prepared. Quantifoil R1.2/1.3 300 mesh copper grids were glow
discharged for 30 s at 22 mA, then 4 μL of sample was applied. The grids were blotted with a single
blotting paper for 5.5 s at a blot force of -2 in 90% humidity at 4°C before being plunge-frozen in
liquid ethane using an FEI Vitrobot Mark IV (Thermo Fisher Scientific).
1258
Data collection was performed on a Titan Krios G2 transmission electron microscope (Thermo
Fisher Scientific) at 300 kV with a K3 summit direct detection camera (Gatan). SerialEM was used
to collect 5,381 movies at a nominal magnification of 81,000× in super-resolution mode with a
magnified pixel size of 1.068Å on the specimen level (counting mode). Movies were dose-
fractionated into 43 frames of 0.08 seconds each at a dose rate of 16.9 electrons/Å^2 per s for a
total dose of 51 e$^-$/Å^2. The defocus range for the sample was between 1.4 and 2.4 μm.
Image processing was carried out using RELION-3.1[71]. The 5,381 micrographs were motion-
corrected and dose-weighted in MotionCor2[72] with a binning factor of 2 and divided into 5×5
patches. The contrast transfer function was calculated with CTFFIND-4.1[73]. Reference-free 2D
classification of 1,044 manually picked particles was done, and the three best classes that showed
well-defined particles were used as references for autopicking that yielded a dataset of 3.36
million particles. Multiple rounds of 2D classification were carried out to remove ice
contamination and bad particles. The subset of the most homogeneous 254,169 particles was used

to generate a 3D initial model using stochastic gradient descent with C1 symmetry as implemented
in RELION, and 3D classification further isolated 86,720 particles. This 3D class underwent auto-
refinement and post-processing for a final reconstruction at a resolution of 8.3 Å according to the
Fourier shell correlation (FSC) = 0.143 criterion.

Correlative Light and Electron Microscopy
For TEM CLEM, COS-7 cells were plated on 35 mm MatTek dish (P35G-1.5-14-CGRD) and
transfected as described above with SHIP164^mScarlet-I, GFP-WDFY2, mito-BFP, and Ras^{G12V}.
Cells were pre-fixed in 4% PFA in Live Cell Imaging Buffer (see above) then washed before
fluorescence light microscopy imaging. Regions of interest were selected and their coordinates on
the dish were identified using phase contrast. Cells were further fixed with 2.5% glutaraldehyde
in 0.1 M sodium cacodylate buffer, postfixed in 2% OsO_4 and 1.5% $K_4Fe(CN)_6$ (Sigma-Aldrich) in
0.1 M sodium cacodylate buffer, en bloc stained with 2% aqueous uranyl acetate, dehydrated, and
embedded in Embed 812. Cells of interest were relocated based on the pre-recorded coordinates.
Ultrathin sections (50-60 nm) were observed in a Talos L 120C TEM microscope at 80 kV, images
were taken with Velox software and a 4k × 4K Ceta CMOS Camera (Thermo Fisher Scientific). For
FIBSEM CLEM, mito-BFP was expressed in endogenously tagged eSHIP164^mNG HeLa cells
processed as above except the dehydration was implemented at the low temperatures gradually
decreased from 0 °C to -50 °C. Epon blocks were glued onto the SEM sample mounting aluminum
and platinum *en bloc* coating on the sample surface was carried out with the sputter coater (Ted
Pella, Inc., Redding, California). Sample were FIBSEM imaged in a Crossbeam 550 FIBSEM
workstation operating under SmartSEM (Carl Zeiss Microscopy GmbH, Oberkochen, Germany)
and Atlas engine 5 (Fibics incorporated, Ottawa, Canada). The imaging resolution was set at 5
nm/pixel in the X, Y axis with milling being performed at 2.5 nm/step along the Z axis (binned
down by 2 to 5 nm when images were exported) to achieve an isotropic resolution of 5nm voxel.
Images were aligned and exported with Atlas 5 (Fibics incorporated, Ottawa, Canada), further
processed and analyzed with DragonFly Pro software [Object Research Systems (ORS) Inc.,
Montreal, Canada]. Except noted all reagents were from EMS (Electron Microscopy Sciences),
Hatfield, PA.
1299
Cell culture and Transfections
hTERT-RPE1 cells were a kind gift of A. Audhya (University of Wisconsin, Madison, WI). HeLa
and COS-7 cells were obtained from ATCC. All mammalian cells were maintained at 37ºC in
humidified atmosphere at 5% CO_2 unless noted otherwise. HeLa and COS-7 cells were grown in
DMEM and RPE1 cells in DMEM/F12 medium (Thermo Fisher Scientific) supplemented with
10% FBS, 100 U/mL penicillin, 100mg/mL streptomycin, and 2mM glutamax (Thermo Fisher
Scientific). Expi293F were grown to manufacturers specification in Expi293 expression medium
with constant shaking. All cell lines were routinely tested and always resulted free from
mycoplasma contamination.
1309
Transient transfections were carried out on cells that were seeded at last 8 hours prior. All
transfections of plasmids used FugeneHD (Promega) to manufacturers specifications for 16-24
hours in complete media without antibiotics. Expi293 transfections were carried out to
manufacturers specifications and reagents. siRNAs (IDT) for knowck-down experiments against
SHIP164 (design ID: hs.Ri.UHRF1BP1L.13.2) were transfected using Lipofectamine RNAiMAX
(ThermoFisher) for 48 hours before fixation for immunocytochemistry experiments or collected
for immunoblotting (see below)
1317
Immunoblotting and Imaging Procedure
All cell samples analyzed by immunoblotting (RPE1 and HeLa) were scraped from plates and
harvested by centrifugation (500xg for 5 minutes). The pellet was washed with ice-cold PBS and

centrifuged again in a 1.7mL Eppendorf tube. The cells pellet was resuspended in Lysis buffer
(20mM Tris-HCl pH 7.5, 150mM NaCl, 1mM EDTA) containing protease inhibitor cocktail
(Roche) and lysed using mechanical disruption (Isobiotec Cell Homogenizer). The lysate was
clarified by centrifugation (17,000xg for 10 minutes) and then mixed with 3x SDS sample buffer
(Cold Spring Harbor) to 1x concentration and then heated to 95C for 5 minutes. A small portion
of lysate was reserved for quantification of protein concentration by Bradford. 15-25ug of protein
samples were separated by electrophoresis on a 4-20% Mini-PROTEAN TGX gel and then
subjected to standard western blot transfer and procedures. Blots were imaged using the Odyssey
imaging system (Licor) using manufacturers protocols. All primary antibodies used in this study
are listed in **Table S1**.
1331
1332
Live Cell Imaging and Immunofluorescence
For all live cell microscopy cells were seeded on glass-bottom mat-tek dishes (MATtek
corporation) 5500/cm^2 in complete media. Transfections were carried out as described above.
Spinning-disk confocal imaging was preformed 16-24 post transfection using an Andor Dragon
Fly 200 (Oxford Instruments) inverted microscope equipped with a Zyla cMOS 5.5 camera and
controlled by Fusion (Oxford Instruments) software. Laser lines used: DAPI, 440nm; GFP, 488;
RFP, 561; Cy5, 647. Images were acquired with a PlanApo objective (60x 1.45-NA). During
imaging, cells were maintained in Live Cell Imaging buffer (Life Technologies) in a cage incubator
(Okolab) with humidified atmosphere at 37C.
1342
Halo-tag ligands ligands were used at a final concentration of 200nM. Cells were incubated with
indicated dye for 45 minutes in complete media, rinsed three times with complete media, and
then incubated in complete media for 45 minutes before imaging. Immunofluorescent
experiments were performed with cells grown on #1 glass cover slips. Cells were fixed with 4%
PFA in fixation buffer (400mM sucrose, 125mM NaCal, 5mM KCl, 1mM NaH2PO4, 2mM MgCl2,
5mM glucose, 1mM EGTA, 5mM PIPES) for 10 mins at 37ºC, washed 3x with TBS (Tris-HCl pH
7.5, 150mM NaCl), permeabilized using 1x PBS containing 0.5% Triton for 10 min at RT and then
blocked using antibody dilution buffer (50 mM Tris pH 7.5, 150 mM NaCl, 0.1% Triton, 4% BSA)
for 1 hour at 25°C. Coverslips were incubated with the indicated primary antibody diluted in the
same buffer overnight at 4°C. Slides were washed with PBS containing 0.1% Triton to remove
excess primary antibody and subsequently incubated with the indicated secondary antibody
diluted in antibody dilution buffer for 45 min at room temperature in the dark prior to mounting.
1355
Generation of CRISPR edited Cell Lines
SHIP164 RPE1 knockout cells were generated by first identifying a guide RNA (gRNA) directed to
exon 2 of SHIP164 using a Cas9 target identification tool (CHOPCHOP). A single stranded oligo
(IDT) was synthesized containing the sense sequence and then cloned into PX458 (Addgene
#48138) and transfected into RPE1 cells. 48 hours post transfection cells were single cell sorted
using FACS Aria based on GFP expression into 96-well plates, and individual clones were grown
out and subsequently examined by immunoblot analysis. Sanger sequencing was used to confirm
the presence of mutations, which led to frameshifts and premature stop codons within the coding
sequence of SHIP164. Two validated clones were used in subsequent studies.
1365
Endogenous tagging of in HeLa cells was carried out using the ORANGE system. Briefly, a gRNA
targeted close the desired site within the endogenous reading frame of the indicated target protein
was identified and subsequently cloned into empty pORANGE using HiFi assembly at the tandem
BbsI site. The ORF of the indicated fluorescent protein (mNeonGreen or mScarlet-I) with linker
regions to make an in-frame insertion was cloned into the HindIII/XhoI sites of pORANE using
standard restriction enzyme cloning. The finalized plasmid was transfected into wild-type HeLa

cells using the transfection protocol described above. Cells were split once to keep confluency
below 80% and 5-days post transfection and the cells were imaged using confocal microscopy.
After visualizing cells positive for the correct fluorescent protein, cells were serially diluted and
plated at 10 cells per well on a glass bottom 96-well plate (Cellvis P96-1-N) and inspected for
fluorescence when the wells were sub-confluent. Wells with positive fluorescence were expanded,
serially diluted, and plated at one cell per wll on a glass bottom 96-well plate and inspected for
fluorescence when the clones were sub-confluent. All gRNAs used in this study are listed in **Table
S2.**
1380
Mouse Brain Lysate Preparation, Affinity Purification, and Protein Identification
Brains were collected from sacrificed wild-type C57/black mice. For each 1g of whole brian
material, 10 mL of buffer (25mM HEPES pH 7.4, 200mM NaCl, 5% glycerol, protease inhibitors)
was added. Mechanical lysis was performed using a glass dounce-homogenizer (15 strokes).
Triton X-100 was added to 1% and material was rotated at 4C for 30 minutes. Material was
centrifuged at 1000g to remove cell debris and the collected supernatant was centrifuged at
27000g for 20 minutes. The resulting supernatant was flash frozen and stored at -80C until use.
1388
Purified 3xFLAG-SHIP164 or FLAG-tagged control protein was immobilized on anti-FLAG resin
as described above. Immobilized SHIP164 and control protein was incubated with mouse brain
lysate for 16hrs gently rotating at 4C. Resin was spun down and the supernatant was removed
before 2x10 bed volume washes using a wash buffer (25mM HEPES pH 7.4, 150mM NaCl, 1mM
BME, protease inhibitors). Protein bound to resin was boiled in the same wash buffer and eluate
was transferred to a new Eppendorf tube. Protein in the eluate was precipitated using
trichloroacetic acid (TCA) on ice for 16hrs, neutralized using Tris pH 8.5 buffer, and then the
precipitate was pelleted in a table top centrifuge. The precipitate was washed twice with ice cold
acetone and then dried using a speed-vac. Protein and peptide identification was achieved by
shotgun tandem LC-MS/MS analysis. Analysis was performed using Scaffold 5 proteome software
to identify promising identifications.
1400
1401 **Image processing, Analysis, and Statistics**
1402 Florescence images presented in this study are representative of cells imaged in at least three
1403 independent experiments and were processed with ImageJ software. The dimensions of some of
1404 the magnification insets or panels were enlarged using the *Scale* function on ImageJ.
1405
Quantification of vesicle-to-Golgi ratio and peripheral structures: unprocessed images with two
channels (GM130 for Golgi in one channel and the protein of interest in the other channel) were
made into maximum intensity projections. A mask of the Golgi complex was made using the
GM130 signal and then used to subtract the Golgi signal from the signal of the entire cell protein
of interest channel resulting in peripheral signal only. Outlines of individual cells were drawn,
and the amount of peripheral signal was measured and then divided by the extracted Golgi signal
for each cell to obtain the ratio. Where stated that the number of peripheral structures were
counted, the subtracted peripheral image was made into a mask to identify structures of the
indicated area and subsequently counted per cell.
1415
1416 Statistical analysis was performed with GraphPad Prism 7 software. Groups were compared using
1417 a two-tail unpaired Student *t* test and results were deemed significant when a p value was smaller
1418 than 0.05.
1419

1420 **TABLES**
1421
1422 **Table S1.** List of antibodies used in this study.
1423

Protein (epitope)	Company; Catalog number	Antibody species	Working dilution for immunocytochemistry	Working dilution for immunoblotting
GM130	BD Bioscience; 610822	Mouse	1:100	N/A
GM130	Abcam; ab30637	Rabbit	1:300	N/A
EEA1	BD Bioscience; 610457	Mouse	1:100	N/A
EEA1	Thermo Scientific; PA1-063A	Rabbit	1:400	N/A
Sortilin	Abcam; ab243043	Rabbit	1:500	N/A
MPR (IGF-IIR)	Santa Cruz Biotechnology; sc-53146	Mouse	1:100	N/A
TGN46	Abcam; ab50595	Rabbit	1:200	N/A
TGN46	AbD Serotec; AHP500GT	Sheep	1:300	N/A
mNeonGreen	ChromoTek; 32f6-100	Mouse	1:500	1:1000
SHIP164	Custrom antibody produced by Proteintech	Rabbit	N/A	1:500
GAPDH	Proteus; 40-1246	Mouse	N/A	1:1000

1424
1425 **Table S2.** List of ORFs and primers used for cloning in this study.
1426 *See associated excel file.*
1427

ACKNOWLEDGMENTS
We thank members of the De Camilli and Reinisch lab for critical reading of the manuscript. We
also thank Shan Xu and Song Pang (Janelia Research Labs) for help and discussion with FIB-SEM
experiments. This work was in supported in part by grants from the NIH (NS36251, DA18343 and
DK45735), the Kavli Institute for Neuroscience, and the Parkinson Foundation to PDC, the NIH
(R35GM131715) to KMR, the NIH (F32NS108448) to MGH, and the NIH (T32GM008283) to PS.
This research was also funded in part by Aligning Science Across Parkinson's grant ASAP-000580
through the Michael J. Fox Foundation for Parkinson's Research (MJFF) to PDC and KMR. For
the purpose of open access, the author has applied a CC BY public copyright license to all Author
Accepted Manuscripts arising from this submission.
1438

AUTHOR CONTRIBUTIONS

MGH, PS, KMR, PDC conceptualized the project. PS performed structural analysis and lipid binding and transport experiments, YW performed CLEM experiments in collaboration with MGH, and MGH performed everything else. MGH, PS, YW, KMR, PDC analyzed data. MGH, PS, KMR, PDC wrote the original draft of the manuscript, which was then reviewed and edited by all the authors.

1446 **CONFLICT OF INTEREST**
1447 PDC is a member of the Scientific Advisory Board of Casma Therapeutics. No conflict exists with
1448 this role

Figure 1

Figure 2

A

SHIP164^mScarlet-I
GFP-WDFY2

B

GFP-WDFY2
SHIP164^mScarlet-I

Zoom of COS-7 cell

FIB-SEM reconstruction

ER   Vesicles   Endosome

C

GFP-WDFY2
SHIP164^mScarlet-I

Zoom of COS-7 cell

FIB-SEM reconstruction

Vesicles   Endosome

Figure 3

Figure 4

A Low Stx6 expression

SHIP164^mScarlet-I
mNeonGreen-Stx6

Wild-type COS-7

Merge
Stx6
SHIP164

B

	Bait	Identified Target	
		DYNLL1	DYNLL2
Exp-1	SHIP164	*233*	*107*
	Control	4	4
Exp-2	SHIP164	*111*	*109*
	Control	4	3

Peptide Count

C

Input MW S200 fraction number
11 12 13 14 15 16 17

MBP-SHIP164₁₃₁₂₋₁₄₆₄

DYNLL1 (LC8)

D

Single protein expression — GFP-Rab45

Single protein expression — SHIP164^mScarlet-I

Dual expression — SHIP164^mScarlet-I GFP-Rab45 (not shown)

Wild-type COS-7

Merge
SHIP164
Rab45

E

SHIP164   Stx6 Binding Motif   Coiled-coil

Chorein Motif   Disordered   1464

VPS51 RRKAHGMLKLYYGL–SE   LC8 binding motif ---VSRGTQTE---
        LxxYY   VS+ΦTQT
SHIP164 RLKRKKLLKEYYSTESE   SHIP164 QRSVTQATQTSPGV

Figure 5

Figure 6

Figure 7

A

(a)

670 158 44 17 1.35 kDa

SHIP164(Δ901-1099)

3XFLAG

mAU

ml

(b)

MW
(kDa)

SHIP164(Δ901-1099)

— 250
— 150
— 100
— 75
— 50
— 37
— 25
— 20
— 15
— 10

(c)

Tecnai T12 negative stain micrograph

B

50 nm

5,381 movies
MotionCor2
RELION autopick
$3.36 \cdot 10^6$ particles

4 rounds of RELION
2D classification

10 nm

C

8.3Å

FSC=0.143

Fourier Shell Correlation

resolution (1/Å)

#1
34%

#2
35%

#3
31%

RELION 3D
classification
with 254,169 particles

RELION 3D
initial model
with C1 symmetry

RELION 3D refinement
86,720 particles

RELION post-process
8.3 Å

Supplemental Figure 1

Supplemental Figure 5

A

Exon 2   PAM   Intron

Wild-type   agcgtcc--att**agg**gtgagact
Clone 1   agcgtc---att**agg**gtgagact
Clone 2   agcgtcctcatt**agg**gtgagact

            ******#   **************

B

Parental Control   SHIP164 Knockout

αsortilin
αGM130

C

SHIP164 knockout RPE1

αTGN46   αMPR   αTGN46
                   αMPR

D

+untagged SHIP164
+RFP reporter (not shown)
SHIP164 Knockout

αMPR   αsortilin
αGM130   αGM130

E

+RFP reporter (not shown)

+untagged SHIP164

Parental Control

αEEA1   αEEA1

+untagged SHIP164

SHIP164 Knockout

αEEA1   αEEA1

F

siRNA: Control SHIP164

250
150
100

50
37

a-SHIP164

37
anti-Actin

G

Control siRNA   SHIP164 siRNA

αEEA1   αEEA1

Supplemental Figure 6

Zoom of COS-7 cell

GFP: UHRF1BP1 LAMP1

Merge

C6orf107

LAMP1