Correlation scan: identifying genomic regions that affect genetic correlations applied to fertility traits

Babatunde S. Olasege[1,2], Laercio R. Porto-Neto[2], Muhammad S. Tahir[1,2], Gabriela C. Gouveia[3], Angela Cánovas[5], Ben J. Hayes[4] and Marina R. S. Fortes[1,4]

[1]The University of Queensland, School of Chemistry and Molecular Biosciences, Saint Lucia Campus, Brisbane, QLD, 4072, Australia

[2]CSIRO Agriculture and Food, Saint Lucia, QLD, 4067, Australia

[3]Animal Science Department, Veterinary School, Federal University of Minas Gerais, Belo Horizonte 31270-901, Brazil

[4]The University of Queensland, Queensland Alliance for Agriculture and Food Innovation (QAAFI), Saint Lucia Campus, Brisbane, QLD, 4072, Australia

[5]University of Guelph, Department of Animal Biosciences, Centre for Genetic Improvement of Livestock, 50 Stone Rd E, Guelph N1G 2W1, ONT, Canada

*Corresponding author: m.fortes@uq.edu.au

Abstract

Reproductive traits are often genetically correlated. Yet, we don't fully understand the complexities, synergism, or trade-offs between male and female fertility. Here, we introduce correlation scan, a novel framework for identifying the drivers or antagonizers of the genetic correlation between male and female fertility traits across the bovine genome. The identification of these regions facilitates the understanding of the complexity of these traits. Although the methodology was applied to cattle phenotypes, using high-density SNP genotypes, the general framework developed can be applied to any species or traits, and it can easily accommodate genome sequence data.

Keywords: Genomic correlation, drivers, antagonizing, RHOGDI, pathway analysis, QTLs, Correlation scan

Background

In animal genetics, insight into the nature of the genetic relationships between quantitative traits are important because they improve our understanding of complex traits and diseases

[1, 2]. These relationships termed genetic correlations manifest when there is shared genetic

influence between traits (i.e., pleiotropy) [3, 4] or when there is non-random association

between loci (i.e., linkage disequilibrium (LD)) [5, 6]. Estimated genetic correlations provide

information on how genome-wide genetic effects align between two complex traits [7].

Understanding the interplay between the genomic variants and their effects on quantitative

traits can yield insights to improve the prediction of genetic merit and the understanding of

complex traits' biology [8-10]. Estimated genetic correlations have informed animal and crop

breeding for many decades. For example, scrotal circumference is used as an indicator trait in

beef cattle breeding because it is genetically correlated with female fertility traits [11].

Nevertheless, we still have a limited information of the regions across genome regulating the

intersexual correlations between male and female fertility traits. Investigating these regions

and leveraging on the resulting biological information could inspire new approaches in

livestock breeding [12, 13].

Over the past 100 years, different methods have been employed to estimate the genetic

correlation between traits [14-17]. Traditionally, these correlations are estimated from

pedigree data. However, genome-wide single nucleotide polymorphisms (SNPs) are often

used in recent times [18]. It is possible to estimate across-sex correlation between traits and

this research niche continues to attract interest among quantitative geneticists [19-21]. The

resulting estimates from both within and across-sex analyses range from -1 to +1, indicating

the strength and magnitude of the correlation between traits [22]. Despite more than a century

of research on estimating this parameter, it is only very recently that studies attempt to

identify the region(s) in the genome that underpin genetic correlations between traits [23-25].

In theory, we propose that various genomic regions will contribute to the overall genetic

correlation between complex traits. Further, some regions will be driving the genetic

correlation while others might antagonize it. For instance, if the genetic correlation between two traits is 0.70, some regions will yield a significant and positive correlation, say 0.90, while other regions may antagonize the overall estimate, and in that region the correlation could be -0.50. Also, some genomic regions may be neutral, say 0.02 and not significant for the correlation between the studied traits. Identifying driver and antagonizing regions are of particular interest if they are for two important traits which are unfavourably correlated, for example milk yield and fertility in dairy cattle. Identification of such regions could lead to more targeted genomic selection and rapid genetic gains for both traits. Current genomic tools have created a great opportunity to advance our knowledge of genetic correlations between complex traits, by investigating the regions in the genome that drive or antagonize these correlations.

Here, we introduce a framework termed "correlation scan", which uses a sliding window methodology to uncover the genomic regions driving and antagonizing genetic correlations in beef cattle. We applied the method to male and female fertility traits and showcase how the outcomes of this methodology can be interpreted in downstream analyses to gain further insight about the studied traits and their relationships. Reproductive traits are often genetically correlated, and yet we don't fully understand the complexities, synergism, or trade-offs between male and female fertility. To demonstrate the method, we used two pairs of reproductive traits with strong genetic correlations in two independent cattle populations from our previous study [26]. These traits are age at first *corpus luteum* (AGECL, i.e., female puberty) and serum levels of insulin growth hormone (IGF1 measured in bulls, IGF1b, or cows, IGF1c). These pairs of traits serve as example of a positive and a negative correlation between phenotypes measured in males and females, during pubertal development. The populations used in the study are formed by either Brahman (BB) cattle or Tropical Composite (TC) cattle, as described in our previous study [26].

Results

The total number of windows generated and analysed

Using the framework developed in our study (see Materials and Methods), genomic windows

with their corresponding correlation estimates (r) for each pairwise trait in two beef cattle

populations were identified. The total number of windows generated for all pairwise traits in

BB was 5,558 and the number in TC was 6,876. For all windows, the chromosome

coordinates and the corresponding r estimates in each of the two populations are presented in

Additional file 1 (Table S1-S2). The r estimates for all windows were plotted against their

genomic position (i.e., midpoint between the start and end position of each window) (Figure

1). Results are presented separately per cattle population and for each pair of traits

investigated

90
Figure 1. Genome plot of the regions driving and antagonizing trait correlations in Brahman
(BB) and Tropical Composite (TC) for the pairwise traits (BB-AGECL vs IGF1b; **A**, TC-
AGECL vs IGF1b; **B**, BB-IGF1c vs IGF1b; **C**, TC- IGF1c vs IGF1b; **D**). **AGECL**, age at
first corpus; **IGF1**, serum levels of insulin growth hormone (measured in bulls, **IGF1b**, or
cows, **IGF1c**). The correlation estimates were plotted on the y-axis and the genomic position
(i.e., midpoint between the start and end position of each window) of each chromosome on
the x-axis, according to the ARS_UCD1.2 bovine reference genome.

Driver, antagonizing and neutral genomic windows affecting genetic correlations between fertility traits: permutation test

In order to identify drivers, antagonizing and neutral regions across the bovine genome, we performed permutation test by randomly reshuffling the Single Nucleotide Polymorphisms (SNPs) effects in each window across all chromosomes in 1,000 iterations for each trait. Then, we applied our framework on the randomized SNP effects and observed the **r** estimate across each iteration for each window. In most cases, the maximum and the minimum **r** estimates for each window at each iteration (i.e., rand 1 to rand 1,000) range between ±0.20. Therefore, we considered neutral windows with no significant effect on the trait correlation as windows with -0.20≤ **r** ≤0.20 estimates. The genomic plots of the **r** estimates resulting from the permutation test (rand 500 only) in each population are presented in Additional file 2 (Fig. S1). Additional file 3 (Table S3-S6) shows the numbers of windows, their chromosome coordinates, and **r** estimates as well as the maximum and minimum r estimates across the 1,000 iterations for each pair of traits in each of the two populations.

As a result of the permutation test, we considered significant windows with **r** estimates >0.2 and **r** estimates <-0.2. These thresholds were used to define the significant windows or regions (i.e., driver and antagonizing) from non-significant (i.e., neutral) windows or regions. Depending on the overall genetic correlation between traits, driver and antagonizing windows can be deduced: in driver windows, the **r** estimate has the same direction, positive or negative, as the overall genetic correlation; in antagonizing windows it is the opposite.

The number of significant driver windows for the correlation between AGECL and IGF1b was 1,636 in BB and 1,914 in TC cattle. The number of significant windows for the antagonizing was 547 in BB and 898 in TC cattle, for AGECL vs IGF1b. For the correlation between IGF1c and IGF1b, the number of significant driver windows was 1,931 in BB and

2,549 in TC cattle. The antagonizing windows was 402 in BB and 587 in TC cattle (IGF1c vs

IGF1b). The numbers of neutral windows were as follows: 3,375 in BB, and 4,064 in TC for

AGECL vs IGF1b; and 3,225 in BB, and 3,740 in TC for IGF1c vs IGF1b. See Table 1 for

details on numbers of windows in the two beef cattle populations. In addition, the lists of

windows with their chromosomal coordinates for all driver, antagonizing, and neutral regions

are presented in Additional file 4 (Table S7-S18).

Table 1: The number of windows generated for the driver, antagonizing and neutral
windows for each pairwise trait in Brahman and Tropical Composite population.

Pairwise Trait	Number of windows (percentage to the total number of window)			Total number of windows
	Driver (%)	Antagonizing (%)	Neutral (%)	
Brahman				
AGECL vs IGF1b	1,636 (29.40%)	547 (9.84%)	3,375 (60.72%)	5,558
IGF1c vs IGF1b	1,931 (34.74%)	402 (7.23%)	3,225 (58.03%)	5,558
Tropical Composite				
AGECL vs IGF1b	1914 (27.84%)	898 (13.06%)	4064 (59.10%)	6,876
IGF1c vs IGF1b	2549 (37.73%)	587 (8.54%)	3740 (54.39%)	6,876

AGECL, age at first *corpus;* **IGF1c**, serum levels of insulin growth hormone measured in
cow; **IGF1b**, serum levels of insulin growth hormone measured in bulls

For the correlation between AGECL and IGF1b (overall genome-wide correlation of -0.65

(BB) and -0.55 (TC), see Table 2), the largest **r** estimate for the driver windows was -0.96

(bovine chromosome (BTA)14: 23.04 - 25.29Mb) in BB and -0.91 (BTAX: 39.76 - 42.86Mb)

in TC. For the antagonizing windows, the largest **r** estimate was 0.87 (BTAX: 40.87 -

43.88Mb) in BB and 0.61 (BTAX: 66.62 - 69.622Mb) in TC.

For the correlation between IGF1c and IGF1b (overall genome-wide correlation of 0.86 (BB)

and 0.93 (TC), see Table 2), the largest estimate for the driver windows was 0.97 (BTA14:

22.68 - 24.96Mb) in BB and 0.87 (BTA5: 46.13- 47.89Mb) in TC, while the estimate for the

antagonizing was -0.62 (BTA1: 49.01 - 51.67Mb) in BB and -0.90 (BTAX: 65.64 - 68.39Mb)

in TC. All **r** estimates are plotted in Figure 1.

Genes and quantitative trait loci (QTL) within driver and antagonizing regions across

the two populations

Defining driver and antagonizing regions separately for each pair of traits, allowed us to

identify the genes and QTLs within these regions for each of the two beef cattle populations.

The percentage of the overlapping genes (Figure 2) and QTLs (Figure 3) across both

populations was studied. The percentages of genes shared across the significant regions in BB

and TC were calculated as a function of the total number of genes in BB or TC, respectively,

and so they differ (Figure 2 and 3).

The percentage of overlapping genes for each pair of traits in the two populations were as

follows: for AGECL vs IGF1b driver regions, about 48% of the total number of genes

annotated were shared between the two population, whereas, for the antagonizing regions,

22% of the gene annotated in BB were present in the TC population, and 26% of the genes

annotated in TC were present in BB; for IGF1c vs IGF1b, the two populations shared about

58% of total number of genes annotated for the driver regions and about 17% were shared for

the antagonizing regions.

The percentage of overlapping QTLs for each pair of traits in BB and TC population were as

follows: for AGECL vs IGF1b driver regions, 52% of the QTLs annotated in BB were

160 present in TC and 35% of the QTLs annotated in TC were present in BB, whereas, for the

161 antagonizing regions, 20% of the QTLs annotated in BB were present in TC and 18% of the

162 QTLs annotated in TC were present in BB; for IGF1c vs IGF1b, 56% of the genes annotated

163 in BB were present in TC, and 52% of the genes annotated in TC were present in BB,

164 whereas, for the antagonizing regions, 29% of the genes annotated in BB were present in TC

165 and 33% of the genes annotated in TC were present in BB population.

Figure 2. Genes annotated in the significant (i.e., driver and antagonizing) genomic regions identified as explaining the genetic correlations between male and female fertility traits in Brahman (BB) and Tropical Composite (TC) population. The overlaps between the two studied populations are in the diagonal of each plot for each pair of traits within the driver (above) and antagonizing (below) regions. The darker the colour within the squares, the higher the percentage of shared genes or QTLs. **AGECL**, age at first corpus; **IGF1**, serum levels of insulin growth hormone (measured in bulls, **IGF1b**, or cows, **IGF1c).**

Drivers

Antagonizing

Figure 3. QTLs annotated in the significant (i.e., driver and antagonizing) genomic regions identified as explaining the genetic correlations between male and female fertility traits in Brahman (BB) and Tropical Composite (TC) population. The overlaps between the two studied populations are in the diagonal of each plot for each pair of traits within the driver (above) and antagonizing (below) regions. The darker the colour within the squares, the higher the percentage of shared genes or QTLs. **AGECL**, age at first corpus luteum; **IGF1**, serum levels of insulin growth hormone (measured in bulls, **IGF1b**, or cows, **IGF1c).**

Functional classification of QTLs within genomic regions that explain the genetic correlations between male and female fertility

To infer biological function and mine the existing literature, we examined the types of QTL (milk, reproduction, production, meat and carcass, health and exterior) present in the significant genomic regions identified above using GALLO [27]. The most frequent QTLs across all pairwise traits in the two populations for the driver and antagonizing regions were QTLs related to milk production, accounting for about 30-51% in most cases. This was followed by reproductive QTLs accounting for about 13-48% and production QTLs comprising 6-24%. Other QTL types (Exterior, health and meat and carcass) accounted for a relatively small proportion of QTLs in the significant regions (Figure 4 and 5). In addition, we report the top 10 results for QTLs related to reproductive traits as these are relevant to our studied traits (Figure 4 and 5). Among these reproductive QTLs, traits related to puberty (i.e., age at puberty, scrotal circumference) were prevalent in both populations.

Figure 4. Percentage of QTL type (pie chart) and trait related to reproduction QTLs (barplots) for the QTL annotation results obtained for (A) AGECL vs IGF1b - driver, (B) AGECL vs IGF1b- antagonizing in Brahman (BB) and Tropical Composite (TC) population. **AGECL**, age at first corpus luteum, **IGF1**, serum levels of insulin growth hormone (measured in bulls, **IGF1b**, or cows, **IGF1c**)

Figure 5. Percentage of QTL type (pie chart) and trait related to reproduction QTLs (barplots) for the QTL annotation results obtained for **(A)** IGF1c vs IGF1b- driver, (B) IGF1c vs IGF1b- antagonizing in Brahman (BB) and Tropical Composite (TC). **AGECL**, age at first corpus luteum, **IGF1**, serum levels of insulin growth hormone (in bulls, **IGF1b**, or cows, **IGF1c**).

QTL enrichment analysis

We performed a chromosome-wide QTL enrichment analysis to further test the significance of the QTLs identified for all the driver and antagonizing regions in each cattle population, for each trait pair using GALLO [27]. Enriched QTLs for the studied traits span across most QTL types, indicating the presence of complex genetic mechanisms. The results of the chromosome-wide QTLs enrichment (FDR-corrected p-value≤0.05) for the driver and antagonizing regions for all pairwise traits in each population are presented in Additional file 5 (Table S19-26).

For the driver regions, the number of QTLs enriched over a wide range of chromosomes for AGECL vs IGF1b were 233 and 144 in BB and TC beef cattle population, respectively. The number was 227 (BB) and 220 (TC) for IGF1c vs IGF1b. For AGECL vs IGF1b, the most enriched chromosome (no of enriched QTLs in parenthesis) was BTA5 (36) and BTA14 (18) in BB and TC, respectively. IGF1c vs IGF1b also followed similar pattern with the result above, with BTA5 (41) being the most enriched chromosome in BB and BTA14 (51) as the most in TC.

For the antagonizing regions, the number of QTLs enriched across the bovine chromosomes for AGECL vs IGF1b were 127 and 178 in BB and TC beef cattle population, respectively. The number was 179 (BB) and 195 (TC) for IGF1c vs IGF1b. For AGECL vs IGF1b, the most enriched chromosome was BTA17 (14) and BTA26 (21) in BB and TC, respectively. For IGF1c vs IGF1b, however, BTA14 (23) was the most enriched chromosome for these regions in BB, whereas, in TC, BTA14 (23) was the most enriched.

 To identify the common results and shared biology between the driver and the antagonizing regions, we also investigated the overlaps of the QTL types associated with the studied trait (i.e., reproduction) in the two populations. The relationship between the top 10 enriched

244 reproductive QTLs in BB and TC are presented in Figure 6. Irrespective of the trait pair, for

245 the driver regions, the reproductive QTLs in BB in most cases overlap with those identified in

246 TC. However, for the antagonizing regions, not all reproductive QTLs in BB were found in

247 TC   beef   cattle   population.

Driver

AGECL vs IGF1b   **IGF1c vs IGF1b**

AGECL vs IGF1b   **Antagonizing**   **IGF1c vs IGF1b**

Figure 6. Chord plot showing the relationship between the top 10 enriched reproductive QTLs
between Brahman (BB) and Tropical Composite (TC) for the driver (top) and the antagonizing
(bottom) regions of the studied traits. **AGECL**, age at first corpus luteum, **IGF1c**, serum levels
of insulin growth hormone measured in cow; IGF1b. LUTACT, Luteal activity; SCRCIR,
Scrotal circumference; INHIB, Inhibin level; PUBAGE, Age at puberty; SPREC, Sexual
precocity; NRR, Non-return rate; CALEASE, Calving ease; INSINT, Interval from first to
last insemination; SB, Still birth; CALEST, Interval to first estrus after calving; GLENGTH,
Gestation length, NSP, Percentage normal sperm.

Functional enrichment analysis

Leveraging our methodology's directionality of gene effects with Ingenuity Pathway

Analysis (IPA; http://www.ingenuity.com), we identified the enriched canonical metabolic

pathways enriched at Benjamini–Hochberg corrected p-values (BH-*P*-value) of p<0.01. The

graphical presentation of the canonical metabolic pathways predicted by IPA to be enriched

and the proportion of driver and antagonizing genes in each pathway for all pairwise traits

investigated in each population are illustrated in Additional file 6 (Fig. S2-S9). Although IPA

provided information about whether the predicted pathways were being activated or inhibited

based on our data, we remain cautious when interpreting our results since the **r** estimates are

not the same as gene expression values, and IPA was originally designed to mine gene

expression data.

The number of pathways enriched for AGECL vs IGF1b was 49 in BB and 68 in TC. For

IGF1c vs IGF1b, the number of enriched pathways was 156 and 87 in BB and TC,

respectively. For AGECL vs IGF1b in BB, the top 5 enriched canonical metabolic pathways

were cardiac hypertrophy signaling (Enhanced), toll-like Receptor signaling, IL-6 Signaling,

hepatic fibrosis signaling pathway and STAT3 Pathway. In TC population, the top 5 enriched

canonical metabolic pathways were breast cancer regulation by stathmin1, signaling by Rho

family GTPases, opioid signaling pathway, endocannabinoid developing neuron pathway,

and CREB signaling in neurons.

For IGF1c vs IGF1b in BB, the top 5 enriched canonical metabolic pathways were cardiac

hypertrophy signaling, CREB signaling in neurons, thrombin signaling, estrogen receptor

signaling, opioid signaling pathway and AMPK signaling. In TC, the top 5 enriched pathways

were CREB signaling in neurons, cardiac hypertrophy signaling (enhanced), opioid signaling

pathway, gαs signaling and breast cancer regulation by stathmin1. The enriched canonical

metabolic pathways and all the genes involved in each pathway are available in Additional

file 7 (Table S27-30).

Discussion

Complex phenotypes, including fertility, consist of multiple genetically correlated rather than

independent traits [6]. The interplay between traits involve many genomic regions, usually in

a large and polygenic regulatory network [28-31]. Genomic signals that regulate (i.e., drive or

antagonize) complex traits are widely spread across the genome, including near many genes

without significant effect on the phenotype or disease [29]. In the present post-genomic era,

unravelling the genomic regions that regulate complex traits and the metabolic pathways

associated with these phenotypes has become an important aspect of genetic studies in

humans and animals [32]. In this study, we developed a novel framework termed correlation

scan to reveal the significant regions that either drives or antagonize the genetic correlations

between traits, across the genome. In addition, this method can also reveal genomic regions

with no effect on the studied traits (neutral windows). The framework developed uses best

linear unbiased prediction (BLUP) solution of SNP effects to estimate the local correlations

between studied traits. Local correlations are based on sliding windows of 500-SNPs. We

applied these sliding windows approach to reproductive traits measured in two populations

and subject the outcomes, significant windows, to further analyses using GALLO [27]. and

IPA (http://www.ingenuity.com) to gain further insight about the biology of studied traits and

their relationships. Although the methodology was applied to beef cattle traits, using high-

density SNP chip genotypes, the general framework can be applied to any species, any traits,

and it can easily accommodate sequence level data.

Our results agreed with the established notion that multiple loci regulate reproductive traits

[33-35]. Also, the mode of action of these loci and the magnitude of their effect varies across

the genome. While some regions had no effect on the genetic correlations under

investigation, other loci drive or antagonize the relationships between male and female

fertility. The identification of driver and antagonizing loci creates opportunities to further

understand the molecular mechanisms affecting quantitative traits. For example, correlations

estimated from SNP effects have allowed researchers to construct gene networks [36].

Thereby, these types of approaches contribute to linking genotype with phenotype.

The two beef cattle populations investigated in this study are distinct in terms of their genetic

composition. Brahman (BB) cattle are typically of *Bos indicus* origin whereas TC beef cattle

emanated from the crossing between Bos *indicus* and Bos *taurus* breed [37]. Despite these

differences, we found that a considerable number of annotated genes and QTLs driving trait

correlation overlaps across breeds, although with variations in the size of SNP effects. This

corroborates the findings of Bolormaa, Pryce [38], where a substantial number of QTLs were

found segregating in *Bos indicus* and composite cattle using the same dataset. In this present

study, the top genomic signal driving trait correlation across all pairwise traits in BB were

located on BTA14. The significant region contains a widely known and well-characterized

QTL, including the *PLAG1* gene, reported to be associated with growth and reproductive

traits in our populations and other studies [39-44]. In TC however, the top signal differs

across traits and mostly spread across two or three chromosomes, although with considerable

number of overlaps with BB. This could be partly due to the variations in the architecture of

composite breed [45]. The genome of composite breeds usually contains new haplotypes

emerging from generations of crossbreeding. Moreover, the contribution of the founder

populations on chromosomes and specific genomic regions are usually unevenly distributed,

which most likely shapes the genome of composite breeds [45]. In short, differences between

BB and TC are likely to impact the results of our analyses. Breed differences are expected,

and so when two breeds share a similar result, it enhances our confidence in calling

significant windows for the interplay between male and female fertility traits.

Most genomic regions antagonizing the genetic correlations between male and female

fertility traits were located on chromosome X. Gene expression on chromosome X differs

across-sex, resulting in genomic sexual conflict [46-48]. Genes in these antagonizing regions

include *PO1FB, ZNF711, APOOL, HDX, DACH2, FAM133A,* among others. These genes are

associated with different disorders including infertility, reproductive deficiencies, primary

ovarian failure [49-51]. When some of these genes are over-expressed, it can dysregulate the

cristae morphology of the mammalian mitochondria [52]. Understanding how these

antagonizing genes interact to influence (in)fertility could help improve the reproductive

potentials of beef cattle.

In animal production, more research is carried out on milk production-related traits, thereby

creating large proportion of records for these traits in the cattle QTL database. These volumes

of records can create a bias in the QTLs representativeness [27]. The QTL enrichment

analysis allows testing the significance of the QTL representative using chromosome-wide

approach to detect specific genomic region with many QTLs for a specific trait. For example,

taking the driver regions for AGECL vs IGF1b in BB, the top enriched QTLs was found in

BTA5, harbouring 36 QTLs. These QTLs comprised 8 different QTLs for reproduction

(inhibin level, scrotal circumference, interval of first estrus after calving, gestation length,

insemination per conception, conception rate, daughter pregnancy rate, and pregnancy rate).

These 8-traits listed here have been found to be correlated with puberty (studied traits) in

cattle. For instance, inhibin is regarded as a biomarker for sexual development because it

regulates spermatogenesis in both beef and dairy bulls [53, 54]. Moderate genetic correlation

was found between inhibin and AGECL [55] and between inhibin and IGF1b [56] in BB.

Scrotal circumference has also been found to be a moderate predictor of AGECL and IGF1b

in BB [21, 26]. Thus, BTA5 may be a candidate region for fertility in BB beef cattle

population. Other enriched QTLs out of the 36 mentioned above include 8 different

production traits (average daily gain, metabolic body weight, length of productive life, body

weight, rump width, body depth, residual feed intake, and net merit). These traits are related

to feed efficiency in cattle. Improving feed efficiency of beef cattle is a major concern for

beef producers. A recent study from Canal, Fontes [57] found that heifers that efficiently

utilize feed attain puberty early than less feed efficient ones. Moreover, heifers that attain

puberty at a relatively younger age have the potential to conceive early in life and be more

productive throughout their lifetime [58]. In addition, IGF1 is an effective selection tools to

improve feed efficiency and other production related traits, allowing breeders to preselect

animals that can utilize feed efficiently [59, 60]. Other enriched QTLs for BTA5 in BB are

related to exterior (7), milk (6), milk and carcass (5) and health (3) traits. Of note, the

objective of most beef cattle breeding programs is to change the genetic merit of their cattle

for many traits of interest [61]. The recurrent association of the BTA5 with multiple traits

could suggest complex genetic mechanisms such as pleiotropy, epistasis, hitchhiking effects,

linkage disequilibrium etc., regulating this chromosomal region [62, 63]. Therefore, breeders

could target BTA5 to select multiple traits without any antagonistic effect on other traits

listed herein.

Another interesting result from this study is the shared biology between the two breeds

relative to the traits under study. Despite breed differences, the enriched reproductive QTLs

driving the genetic correlations between male and female fertility are the same for the two

cattle populations (Figure 6). Most of the enriched QTLs are related to reproductive traits

measured early life. A possible explanation could be that the reproductive phenotypes shared

common fundamental biology in the two populations. For the antagonizing regions, however,

most of the reproductive QTLs were breed specific depending on the trait pairwise. Perhaps,

this could be partly explained by the diverse genetic composition of the two breeds.

Understanding the genomic architectures driving these early-in-life male and female fertility

traits and their known genomic antagonisms could foster effective selection for both traits in

tropical breeds [64, 65].

The major challenge faced by researchers when analysing an overwhelmingly large amount

of genomic data is how to extract meaningful mechanistic insights into the underlying

biology characterizing the given trait under study. To increase the explanatory power of

genomic studies, pathway analysis has become first choice, providing researcher with the

ability to infer meanings to high-throughput genomic data [66]. Leveraging the directionality

of gene effects from our method with IPA knowledge base, several biological pathways

known to be involved in reproduction (i.e., studied trait) were significantly enriched for all

pairwise traits investigated across the two breeds. These pathways include estrogen receptor

signaling, p38 MAPK signaling, GnRH signaling, sperm motility, cAMP-mediated signaling,

AMPK signaling, and androgen signaling. Although IPA provided information about the

activation or inhibition state for the enriched canonical metabolic pathways with the use of

the **r** estimates in place of the gene expression values, we are not sure if these pathways were

being activated or inhibited since we don't have information about the expression values of

the genes in these pathways. For example, Rho GDP Dissociation Inhibitor (RHOGDI)

pathway was the only significant signaling pathway found to be inhibited across breeds in all

pairwise traits investigated using IPA comparison analysis. The RHOGDIs (RHOGDIα,

RHOGDIβ and RHOGDIγ) are well-characterized as a negative regulator of Rho GTPases

[67]. These Rho GTPases play pivotal roles within the cell, including cell migration,

membrane trafficking, invasion, gene transcription, polarity, adhesion, cell survival and

death; a process significantly involved in cancer initiation and metastasis [68, 69]. Once

RHOGDI is inhibited, it induces constitutive activation of Rho GTPases, resulting in several

malignant phenotype including tumour growth, angiogenesis, and invasive phenotypes [69,

70]. For instance, knocking out one of the three RHODGI genes resulted in a renal defect that

progressively leads to death in adult mice, although embryonic development was not affected

[71]. Togawa, Miyoshi [72] also found that male mice lacking RHOGDI1 were infertile with

impaired spermatogenesis. The authors also reported problems of implantation in female

mice due to this knockout. The knockout of two of the three RHOGDIs often results in a

more severe phenotypes with additional immunological defects than when one of the

RHOGDIs is disrupted [73]. Numerous studies have also reported that the RHOGDIs protein

are involved in sperm movement, sperm capacitation and acrosome reaction, a process that is

critical to occur for the sperm to interact and penetrate the egg for fertilization to take place

[74-76]. Perhaps, this could be the major reason why signaling by Rho family GTPases were

enriched in our metabolic pathway analysis. Notably, low reproduction performance is one of

the major challenges facing beef producers in Northern Australia [77, 78]. Reproductive

wastage is usually common, which is often a result of pregnancy failure and calf mortality

[79, 80]. Given the role of the RHODGI pathway in reproduction, future studies could use

gene expression data to investigate the genes involved in these pathways as a candidate

region for infertility in cattle since we only use the **r** estimates in this study.

Conclusion

Overall, the framework developed in this study extends our knowledge about the regions

driving and antagonizing correlations between quantitative traits. About 40% of the total

genomic regions were identified as driving and antagonizing genetic correlations between

male and female fertility traits in the two population. These regions confirmed the polygenic

nature of the traits being studied and pointed to genes of interest. Quantitative trait loci (QTL)

and functional enrichment analysis revealed that many significant windows co-located with

known QTLs related to milk production and fertility traits, especially puberty. In general, the

enriched reproductive QTLs driving the genetic correlations between male and female

fertility are the same for both cattle populations, while the antagonizing regions were

population specific. Moreover, most of the antagonizing regions were mapped to the

chromosome X. These results suggest regions of the chromosome X for further investigation

into the trade-offs between male and female fertility. Although the methodology was applied

to cattle phenotypes, using high-density SNP genotypes, the general framework developed

can be applied to any species or traits, and it can easily accommodate genome sequence data.

Materials and Methods

Traits, genotypes and estimated genetic correlations

The traits used to demonstrate this methodology are a subset of traits from our previous study

[26], where bivariate genetic correlations were estimated between 7 male and 6 female early-

in-life reproductive phenotypes in two independent tropical beef cattle populations (BB and

TC). The two female traits selected for this study are age at detection of the first *corpus*

luteum (AGECL, days) and cows' blood concentration of insulin growth-factor 1, measured

at 18 months of age (IGF1c). Only one male trait was selected: the blood concentration of

insulin growth-factor 1, measured at 6 months of age (IGF1b). These traits are important in

beef cattle fertility, especially during pubertal development. The estimated genomic

correlations between the traits listed above in each population have been reported in our

previous study [26]. These estimates and their corresponding standard error (S.E), number of

SNPs and number of animals in each population are provided in Table 2. These traits were

selected because they had significant estimates of genomic correlation (i.e., traits with

standard error (S.E) less than half of the size of the correlation) and different strength or

direction of genetic relationships (i.e., negatively, and positively correlated traits). In brief,

across-sex genetic correlations were estimated in a bivariate analysis using the linear mixed

model approach. Firstly, the 770,000 genotypes were mapped to the new assembly of the

bovine reference genome (ARS_UCD1.2, GenBank assembly accession GCA_002263795.2;

[81]). After quality control filtering (i.e., excluding all SNPs with a minor allele frequency

less than 5%), 554,712 and 686,626 SNPs remained for BB and TC datasets, respectively.

Finally, bivariate genetic correlations were estimated using GIBBS2F90 [82], resulting to the

estimates in Table 2.

463 Table 2: Genomic correlations estimates and their corresponding standard error (s.e),
464 number of animals and number of SNPs estimates

Pairwise traits	No of animals	Number of SNP	Genetic correlation (s.e)
Brahman			
AGECL vs IGF1b	AGECL- 980	554K	-0.65 (0.13)
	IGF1b- 964		
IGF1c vs IGF1b	IGF1c- 995	554K	0.86 (0.11)
	IGF1b- 964		
Tropical Composite			
AGECL vs IGF1b	AGECL-996	686K	-0.55 (0.14)
	IGF1b- 998		
IGF1b vs IGF1b	IGF1c- 1015	686K	0.93 (0.11)
	IGF1b- 998		

AGECL, age at first *corpus;* IGF1c, serum levels of insulin growth hormone measured in
cow; IGF1b, serum levels of insulin growth hormone measured in bulls; SNP, Single
Nucleotide Polymorphisms; s.e, standard error

Overview of methods

For each trait considered in the two beef cattle populations, we estimated the genomic

breeding values (GEBVs) of individuals using the genomic best linear unbiased prediction

(GBLUP) model:

$$\mathbf{y} = \mathbf{Xb} + \mathbf{Za} + \mathbf{e} \qquad \text{[1]}$$

where *y* is the vector of phenotypes, $\mathbf{X}$ is the incidence matrix of fixed effects, *b* is the vector

of fixed effects, $\mathbf{Z}$ is the design matrix assigned to genomic breeding values, *a* is the vector of

GEBVs for each animal, and *e* is the vector of residuals. Vectors **a** and **e** are assumed to

follow a normal distribution, thus $\mathbf{a}{\sim}\mathrm{N}\ (\mathbf{0},\ \mathbf{G}\sigma^2\mathbf{g}\square)$ and $\mathbf{e}{\sim}\mathrm{N}\ (\mathbf{0},\ \mathbf{I}\sigma^2\mathbf{e})\square$. Matrix $\mathbf{G}$ is the

genomic relationship matrix.

Following the estimation of GEBVs for each animal in each trait, we then back-solved these

GEBVs to obtain SNP effects for all chromosomes following the method illustrated by

Strandén and Garrick [83];

$$\widehat{u} = \frac{1}{\sum_{j=1}^{m} 2p_j(1-p)} X'G^{-1}\widehat{a} \qquad \text{[2]}$$

481
Where $\widehat{u}$ is the vector of estimated SNP effects; *m* is the number of SNPs; p_j is the allele frequency of

the second allele of the jth marker; $\mathbf{X}$ is a matrix with gene contents for all markers; $\mathbf{G}$ is a genomic

relationship matrix and $\widehat{a}$ is a vector of GEBVs.

Notably, the estimation of GEBVs using GBLUP model [84-86] and the back-solving for

GEBVs to obtain SNP effects using equation **[2]** both fits all SNP simultaneously, thereby

allowing the joint estimation of SNP effects. As a result, the SNP effects from this approach

are not biased by LD and resulting effect sizes can be considered independent.

Using a chromosome-wide approach, we divided SNPs on the same chromosome into small

sliding windows of 500 SNPs each and then estimated the correlation (**r**) between traits as

being the correlation between the 500-SNP effects estimated for trait A and the 500-SNP effects estimated for trait B. We then moved 100 SNPs further from the start of the previous window to select the next 500-SNP window, which partially overlapped with previous window, hence producing sliding windows that were 100 SNPs distant from the previous window. This was repeated for each trait pair, and for each chromosome, in a chromosome-by-chromosome approach. The resulting r estimates for all the chromosomes combined were denoted as W_1….W_n. The graphical illustration of this framework is presented in Figure 7. Moreover, the coordinates of the windows (W_1…W_n) were mapped to the ARS_UCD1.2 bovine reference genome. The signals across the genome were visualized with the r estimates of each window on the y-axis and genomic position (i.e the midpoint of the start and end position of each window) of each chromosome on the x-axis. The mapping to the bovine reference genome and plotting of the windows signals' graphs were done using SNP & Variation Suite v8.x Golden Helix [87].

Depending on the overall genetic correlation observed between the traits considered, the driver and antagonizing windows can be deduced. In this study, AGECL and IGF1b were negatively correlated. Hence, the driver windows were windows with significant and negative r estimates, while the antagonizings were windows with significant positive r estimates. For the positively correlated relationship between IGF1c and IGF1b, the driver windows were windows with significant and positive r estimates and the antagonizings were windows with significant and negative r estimates. The significance of each window was established with a permutation test, described in the next section.

Figure 7. The graphical illustration of the sliding window framework. The framework involves 2 steps. Step 1 start from the estimation of genomic breeding values to the obtainment of SNP effects for each pairwise trait. Step 2 start from the estimation of 500-SNP effects in a chromosome-wide approach to obtainment of the correlation estimate in a 100-sliding window.

Permutation test

To ensure the **r** estimates are not just noise but real signals, we performed permutation test by randomly reshuffling the SNP effects in each window across all the chromosomes in 1,000 iterations for each trait. Subsequently, we estimated correlations for 100-sliding windows of 500-SNP effects as described above. Finally, we observed the maximum and minimum **r** estimates for all the windows (W_1.......W_n) across the 1,000 iterations to reveal windows that were significant on the pairwise traits under investigation. Afterwards, we mapped the resulting **r** estimates for each window to the ARS_UCD1.2 bovine reference genome and plot the **r** estimates on the y-axis against the genomic position of each chromosome on the x-axis as described above. Consequently, significant windows were selected for the drivers and antagonizings genomic regions for each pairwise. Windows that were not significant were tagged "neutral windows" i.e., windows with no effect on the pairwise trait. Apart from using these windows to estimate genomic correlations and investigate the proportion of variance captured by these regions, they were excluded from other subsequent analyses. Finally, the **r** estimates of the significant windows for the driver and antagonizing regions were ranked from top to bottom in percentage (%) and the rank values were used solely for the purpose of subsequent downstream analyses. The ranking was done separately for the driver and antagonizing windows for each pairwise trait investigated in each population.

Gene and Quantitative Traits Loci (QTL) annotation

The significant windows along with their corresponding chromosome coordinates, **r** estimates and rank values for the driver and antagonizing regions that passed the specified threshold criteria following the permutation test in BB and TC were selected. The selected windows were used for gene and QTL annotation using R package GALLO: Genomic Annotation in Livestock for positional candidate Loci (https://CRAN.R-

project.org/package=GALLO) [27]. The .gtf annotation file corresponding to the bovine gene

annotation from ARS-UCD1.2 assembly and the .gff file with the QTL information from

cattle QTL Database (https://www.animalgenome.org/cgi-bin/QTLdb/index; [88, 89]), were

used for gene and QTL annotation, respectively [27]. The two files use the same bovine

reference genome (ARS-UCD1.2) to map the gene and QTLs. A remarkable advantage of

GALLO is that the software retains all the information present in the input file when

producing the output file. As a result, genes within each window can retain their **r** estimates

and the rank values specific for their window.

The number and percentage of genes and QTLs annotated within a population (BB or TC)

and the overlaps across populations (BB and TC) were investigated. Furthermore, we

examined the QTLs representativeness and diversity to explain better the genomic content of

the significant windows for the driver and antagonizing regions. Hence, the visualization of

the percentage of cattle QTL types from cattle QTL database (i.e milk, reproduction,

production, meat and carcass, health and exterior) were plotted using a pie chart by GALLO

(27).

QTL enrichment analysis

To further test the significance of the QTLs, a within population QTL enrichment analysis

was conducted using a chromosome-based approach. The QTL enrichment analysis, using all

the QTL information annotated within the significant windows for the driver and

antagonizing regions, was performed using the qtl_enrich function from GALLO [90, 91].

Briefly, the observed number of QTLs for each trait in each annotated chromosome were

compared with the expected number using a hypergeometric test approach in a 1,000 iteration

rounds of random sampling from the entire cattle QTL database. With this approach, a p-

value for the QTL enrichment status of each annotated QTLs within the significant windows

was estimated. These estimated p-values were corrected for multiple testing using a false

discovery rate (FDR) of 5%. In addition, we used chord plots to reveal the relationships

between the two breeds for the enriched reproductive QTLs based on the driver and

antagonizing genomic regions.

Functional enrichment analysis

The annotated genes along with their corresponding **r** estimates and rank values for the

significant driver and antagonizing windows for each pairwise trait in BB and TC populations

were subjected to enrichment analysis using the commercial QIAGEN's Ingenuity Pathway

Analysis (IPA; v.8.8, http://www.ingenuity.com). The IPA allows identifying

overrepresented biological mechanism, metabolic pathways, and diseases and biological

functions that are highly relevant to the traits of interest using the directionality of the

submitted gene list [92, 93]. The outcome of our methodology indicates that genes within

each window come with their directionalities, in this case, r estimates. Thus, we leveraged on

the directionality of each gene by allowing the driver genes to be upregulated and

antagonizing genes to be downregulated.

Summarily, a merged dataset containing gene identifiers that were significant for both the

driver and antagonizing windows for each pairwise trait in each population and their

corresponding **r** estimates and rank values were uploaded into IPA. The **r** estimates were

used as the "Expr Log Ratio" and the rank values were used as p-values. The IPA software

recognizes gene with positive signs (+) for "Expr Log Ratio" as upregulated genes and

negative sign (-) as downregulated genes. We aim to allow the driver gene lists to have

positive values for "Expr Log Ratio" and the antagonizing gene lists to be negative. Where

this is not achievable based on the original **r** estimates (i.e., AGECL vs IGF1b), we reversed

the sign for the driver and antagonizing genes to meet this objective.

Of note, IPA can only analyse a maximum of 8,000 gene list. In most cases, the merged gene

list for each pairwise trait in each population is often >8,000. Hence, we used the rank values

as the cut-off to select the top ~80% genes from the driver and antagonizing gene list for the

pathway analyses. Using a proportion of the gene list to infer biological pathways might

result in the loss of some important biological information relevant to the trait of interest. We

analysed the driver and antagonizing gene list separately for each pairwise trait in each

population to ensure no important information was lost because of the cut-offs. Further, we

compare the result of the separate analyses with the merged gene list from the ~80% cut-off.

The pathway analysis was conducted using the "Core Analysis" function implemented within

IPA. In this analysis, associations were calculated using direct and indirect relationships

among the gene lists. At first, the gene lists were mapped to human gene data. Genes without

an associated gene symbol or gene annotation were subjected to an annotation by homology

using   BioMart   application   available   in   the   Ensembl   database

(http://www.ensembl.org/biomart/martview/) [94, 95]. With this approach, we only

considered non-annotated genes with percentage of identity ≥80% with human homolog. The

final datasets used for the IPA analyses are presented in Additional file 8 (Table S31-34).

Finally, the "Core Analysis" was used to identify canonical metabolic pathways enriched at

Benjamini–Hochberg corrected p-values (B-H-P-value) of p<0.01.

Supporting information

Additional file 1: The number of windows, chromosome number, chromosome coordinates,

and correlation estimates for each window for the two pairwise trait in Brahman and Tropical

Composite population (Table S1-S2).

Additional file 2: Genome plots of the correlation estimates from the permutation test at 500

iterations in BB and TC for the two pairwise traits. The correlation estimates were plotted on

the y-axis and the genomic position of each chromosome on the x-axis, according to the

ARS_UCD1.2 bovine reference genome (Figure S1).

Additional file 3: The number of windows, chromosome number, chromosome coordinates,

and correlation estimates as well as the maximum and minimum correlation estimate for each

window for all trait pairwise in Brahman and Tropical Composite population following

permutation test of 500-SNP effects in 100-SNP sliding windows at 1000 iterations (Table

S3-S6).

Additional file 4: The number of windows, chromosome number, chromosome coordinates,

correlation estimates and rank value for driver, antagonizing and the neutral regions that

passed the threshold after permutation test in Brahman (BB) and Tropical Composite

population for the studied trait (Table S7-18).

Additional file 5: The enriched QTLs of the driver and antagonizing regions for all trait

pairwise in Brahman and Tropical Composite cattle. The enriched QTLs are rank based on

the adj.pval (Table S19-26).

Additional file 6: Canonical pathways significantly enriched for all trait pairwise in Brahman

and Tropical Composite population. Significantly enriched canonical pathways were

identified using Benjamini-Hochberg p-values <0.01 (Figure S2-S9).

Additional file 7: The list of the significant enriched canonical metabolic pathways showing

all the genes involved in each pathway for all trait pairwise in Brahman and Tropical

Composite population. Significantly enriched canonical pathways were identified using

Benjamini-Hochberg p-values <0.01. Z-score >2 denote the activation of the pathway. Z-

score <-2 indicate the inhibition of the pathway (Table S27-30).

Additional file 8: The final dataset used for Ingenuity Pathway Analysis (IPA) for all trait

pairwise in Brahman and Tropical Composite cattle (Table S31-34).

Abbreviations

AGECL: Age at first corpus luteum

BB: Brahman

BTA: Bovine chromosome

GBLUP: Genomic Best Linear Unbiased Prediction

GEBV: Genomic estimated breeding values

IGF1b: Serum levels of insulin growth hormone measured in bulls at 6-months of age

IGF1c: Serum levels of insulin growth hormone measured in bulls at 18-months of age

IPA: Ingenuity Pathway Analysis

LD: Linkage Disequilibrium

QTL: Quantitative Trait Loci

R: Correlation estimate

SNP: Single Nucleotide Polymorphism

TC: Tropical Composite

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The data used in this are available from the Cooperative Research Centre for Beef Genetic Technologies (Beef CRC). Data are available from https://www.beefcrc.com with the permission of Meat and Livestock Australia and the University of Queensland (if interested please contact the corresponding author).

Competing interests

None

Funding

BSO was supported by funding from Meat and Livestock Australia (L.GEN.1710; Female Fertility Phenobank and L.GEN.1818; Bull fertility update: historical data, new cohort and advanced genomics) and top-up scholarship from CSIRO.

Authors' contributions

O.B.S and L.P.N. conceived the idea. O.B.S wrote the code, conducted the analyses, and wrote the manuscript. M.S.T. and G.C.C. contributed to the analyses, manuscript writing and editing. A.C. conceived the downstream biological analyses. L.R., B.J.H., and M.R.S. supervised the study. All authors read, edited, and approved the final manuscript.

Acknowledgements

677 This research was conducted using the legacy database of the Cooperative Centre for Beef

678 Genetics Technologies and their core partners including Meat and Livestock Australia. The

679 authors acknowledge all the participants of the Cooperative Centre for Beef Genetics

680 Technologies for their efforts in conducting the field trials and recording the phenotypes. We

681 also acknowledge the mentorship of Professor Steve Moore to Babatunde Olasege in his PhD.

References

1.   Liu S, Yu Y, Zhang S, Cole JB, Tenesa A, Wang T, et al. Epigenomics and genotype-phenotype
association analyses reveal conserved genetic architecture of complex traits in cattle and human.
BMC biology. 2020;18(1):1-16.
2.   Zhang Y, Lu Q, Ye Y, Huang K, Liu W, Wu Y, et al. Local genetic correlation analysis reveals
heterogeneous etiologic sharing of complex traits. bioRxiv. 2020.
3.   Cánovas A, Reverter A, DeAtley KL, Ashley RL, Colgrave ML, Fortes MR, et al. Multi-tissue
omics analyses reveal molecular regulatory networks for puberty in composite beef cattle. PloS one.
2014;9(7):e102551.
4.   Fonseca PAdS, Id-Lahoucine S, Reverter A, Medrano JF, Fortes MS, Casellas J, et al.
Combining multi-OMICs information to identify key-regulator genes for pleiotropic effect on fertility
and production traits in beef cattle. PLoS One. 2018;13(10):e0205295.
5.   Lynch M, Walsh B. Genetics and analysis of quantitative traits. 1998.
6.   Van Rheenen W, Peyrot WJ, Schork AJ, Lee SH, Wray NR. Genetic correlations of polygenic
disease traits: from theory to practice. Nature Reviews Genetics. 2019;20(10):567-81.
7.   Ning Z, Pawitan Y, Shen X. High-definition likelihood inference of genetic correlations across
human complex traits. Nature genetics. 2020;52(8):859-64.
8.   Mackay TF, Stone EA, Ayroles JF. The genetics of quantitative traits: challenges and
prospects. Nature Reviews Genetics. 2009;10(8):565-77.
9.   Pickrell JK, Berisa T, Liu JZ, Ségurel L, Tung JY, Hinds DA. Detection and interpretation of
shared genetic influences on 42 human traits. Nature genetics. 2016;48(7):709-17.
10.   Price AL, Spencer CC, Donnelly P. Progress and promise in understanding the genetic basis of
common diseases. Proceedings of the Royal Society B: Biological Sciences.
2015;282(1821):20151684.
11.   Lunstra D, Cundiff L. Growth and pubertal development in brahman-, boran-, tuli-, belgian
blue-, hereford-and angus-sired f1 bulls. Journal of Animal Science. 2003;81(6):1414-26.
12.   Fang L, Sahana G, Ma P, Su G, Yu Y, Zhang S, et al. Use of biological priors enhances
understanding of genetic architecture and genomic prediction of complex traits within and between
dairy cattle breeds. BMC genomics. 2017;18(1):1-12.
13.   Neyhart JL, Lorenz AJ, Smith KP. Multi-trait improvement by predicting genetic correlations
in breeding crosses. G3: Genes, Genomes, Genetics. 2019;9(10):3153-65.
14.   Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD, et al. Genomic
structural equation modelling provides insights into the multivariate genetic architecture of complex
traits. Nature human behaviour. 2019;3(5):513-25.
15.   Henderson C. Recent developments in variance and covariance estimations. Journal of
Animal Science. 1986;63(1):208-16.

16. Lee SH, Van der Werf JH. MTG2: an efficient algorithm for multivariate linear mixed model
analysis based on genomic information. Bioinformatics. 2016;32(9):1420-2.
17. Turley P, Walters RK, Maghzian O, Okbay A, Lee JJ, Fontana MA, et al. Multi-trait analysis of
genome-wide association summary statistics using MTAG. Nature genetics. 2018;50(2):229-37.
18. Sodini SM, Kemper KE, Wray NR, Trzaskowski M. Comparison of genotypic and phenotypic
correlations: Cheverud's conjecture in humans. Genetics. 2018;209(3):941-8.
19. Collet JM, Fuentes S, Hesketh J, Hill MS, Innocenti P, Morrow EH, et al. Rapid evolution of the
intersexual genetic correlation for fitness in Drosophila melanogaster. Evolution. 2016;70(4):781-95.
20. Connallon T, Matthews G. Cross-sex genetic correlations for fitness and fitness components:
connecting theoretical predictions to empirical patterns. Evolution letters. 2019;3(3):254-62.
21. Raidan FS, Porto-Neto LR, Reverter A. Across-sex genomic-assisted genetic correlations for
sex-influenced traits in Brahman cattle. Genetics Selection Evolution. 2019;51(1):41.
22. Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Patterson N, et al. LD Score
regression distinguishes confounding from polygenicity in genome-wide association studies. Nature
genetics. 2015;47(3):291-5.
23. Shi H, Mancuso N, Spendlove S, Pasaniuc B. Local genetic correlation gives insights into the
shared genetic architecture of complex traits. The American Journal of Human Genetics.
2017;101(5):737-51.
24. Werme J, van der Sluis S, Posthuma D, de Leeuw C. LAVA: An integrated framework for local
genetic correlation analysis. bioRxiv. 2021:2020.12. 31.424652.
25. Zhang Y, Lu Q, Ye Y, Huang K, Liu W, Wu Y, et al. SUPERGNOVA: local genetic correlation
analysis reveals heterogeneous etiologic sharing of complex traits. Genome biology. 2021;22(1):1-
30.
26. Olasege B, Tahir M, Gouveia G, Kour J, Porto-Neto L, Hayes B, et al. Genetic parameter
estimates for male and female fertility traits using genomic data to improve fertility in Australian
beef cattle.
27. Fonseca PA, Suárez-Vega A, Marras G, Cánovas Á. GALLO: An R package for genomic
annotation and integration of multiple data sources in livestock for positional candidate loci.
GigaScience. 2020;9(12):giaa149.
28. Basso K, Margolin AA, Stolovitzky G, Klein U, Dalla-Favera R, Califano A. Reverse engineering
of regulatory networks in human B cells. Nature genetics. 2005;37(4):382-90.
29. Boyle EA, Li YI, Pritchard JK. An expanded view of complex traits: from polygenic to
omnigenic. Cell. 2017;169(7):1177-86.
30. Chen B-S, Yang S-K, Lan C-Y, Chuang Y-J. A systems biology approach to construct the gene
regulatory network of systemic inflammation via microarray and databases mining. BMC Medical
Genomics. 2008;1(1):1-22.
31. Satokangas I, Martin S, Helanterä H, Saramäki J, Kulmuni J. Multi-locus interactions and the
build-up of reproductive isolation. Philosophical Transactions of the Royal Society B.
2020;375(1806):20190543.
32. Skelly DA, Raghupathy N, Robledo RF, Graber JH, Chesler EJ. Reference trait analysis reveals
correlations between gene expression and quantitative traits in disjoint samples. Genetics.
2019;212(3):919-29.
33. Antonarakis SE, Chakravarti A, Cohen JC, Hardy J. Mendelian disorders and multifactorial
traits: the big divide or one for all? Nature Reviews Genetics. 2010;11(5):380-4.
34. Goddard M, Kemper K, MacLeod I, Chamberlain A, Hayes B. Genetics of complex traits:
prediction of phenotype, identification of causal polymorphisms and genetic architecture.
Proceedings of the Royal Society B: Biological Sciences. 2016;283(1835):20160569.

35.   Moser G, Lee SH, Hayes BJ, Goddard ME, Wray NR, Visscher PM. Simultaneous discovery,
estimation and prediction analysis of complex traits using a Bayesian mixture model. PLoS genetics.
2015;11(4):e1004969.
36.   Fortes MR, Reverter A, Zhang Y, Collis E, Nagaraj SH, Jonsson NN, et al. Association weight
matrix for the genetic dissection of puberty in beef cattle. Proceedings of the National Academy of
Sciences. 2010;107(31):13642-7.
37.   Bolormaa S, Hayes B, Hawken R, Zhang Y, Reverter A, Goddard M. Detection of chromosome
segments of zebu and taurine origin and their effect on beef production and growth. Journal of
animal science. 2011;89(7):2050-60.
38.   Bolormaa S, Pryce JE, Kemper KE, Hayes BJ, Zhang Y, Tier B, et al. Detection of quantitative
trait loci in Bos indicus and Bos taurus cattle using genome-wide association studies. Genetics
Selection Evolution. 2013;45(1):1-12.
39.   Fortes M, Kemper K, Sasazaki S, Reverter A, Pryce J, Barendse W, et al. Evidence for
pleiotropism and recent selection in the PLAG 1 region in A ustralian B eef cattle. Animal genetics.
2013;44(6):636-47.
40.   Juma AR, Damdimopoulou PE, Grommen SV, Van de Ven WJ, De Groef B. Emerging role of
PLAG1 as a regulator of growth and reproduction. Journal of Endocrinology. 2016;228(2):R45-R56.
41.   Karim L, Takeda H, Lin L, Druet T, Arias JA, Baurain D, et al. Variants modulating the
expression of a chromosome domain encompassing PLAG1 influence bovine stature. Nature
genetics. 2011;43(5):405-13.
42.   Koufariotis L, Hayes B, Kelly M, Burns B, Lyons R, Stothard P, et al. Sequencing the mosaic
genome of Brahman cattle identifies historic and recent introgression including polled. Scientific
reports. 2018;8(1):1-12.
43.   Littlejohn M, Grala T, Sanders K, Walker C, Waghorn G, Macdonald K, et al. Genetic variation
in PLAG1 associates with early life body weight and peripubertal weight and growth in Bos taurus.
Animal Genetics. 2012;43(5):591-4.
44.   Nishimura S, Watanabe T, Mizoshita K, Tatsuda K, Fujita T, Watanabe N, et al. Genome-wide
association study identified three major QTL for carcass weight including the PLAG1-CHCHD7 QTN
for stature in Japanese Black cattle. BMC genetics. 2012;13(1):1-11.
45.   Paim TdP, Hay E, Wilson C, Thomas M, Kuehn L, Paiva S, et al. Dynamics of genomic
architecture during composite breed development in cattle. Animal genetics. 2020;51(2):224-34.
46.   Mank JE. Population genetics of sexual conflict in the genomic era. Nature Reviews Genetics.
2017;18(12):721-30.
47.   Parsch J, Ellegren H. The evolutionary causes and consequences of sex-biased gene
expression. Nature Reviews Genetics. 2013;14(2):83-7.
48.   Sayadi A, Barrio AM, Immonen E, Dainat J, Berger D, Tellgren-Roth C, et al. The genomic
footprint of sexual conflict. Nature ecology & evolution. 2019;3(12):1725-30.
49.   Bione S, Rizzolio F, Sala C, Ricotti R, Goegan M, Manzini M, et al. Mutation analysis of two
candidate genes for premature ovarian failure, DACH2 and POF1B. Human reproduction.
2004;19(12):2759-66.
50.   Jedidi I, Ouchari M, Yin Q. Sex chromosomes-linked single-gene disorders involved in human
infertility. European journal of medical genetics. 2019;62(9):103560.
51.   Okten G, Gunes S, Onat OE, Tukun A, Ozcelik T, Kocak I. Disruption of HDX gene in premature
ovarian failure. Systems biology in reproductive medicine. 2013;59(4):218-22.
52.   Weber TA, Koob S, Heide H, Wittig I, Head B, van der Bliek A, et al. APOOL is a cardiolipin-
binding constituent of the Mitofilin/MINOS protein complex determining cristae morphology in
mammalian mitochondria. PloS one. 2013;8(5):e63683.

53.   Kaneko H, Noguchi J, Kikuchi K, Akagi S, Shimada A, Taya K, et al. Production and endocrine
role of inhibin during the early development of bull calves. Biology of reproduction. 2001;65(1):209-
15.
54.   Phillips DJ. Activins, inhibins and follistatins in the large domestic species. Domestic animal
endocrinology. 2005;28(1):1-16.
55.   Johnston D, Corbet N, Barwick S, Wolcott ML, Holroyd R. Genetic correlations of young bull
reproductive traits and heifer puberty traits with female reproductive performance in two tropical
beef genotypes in northern Australia. Animal Production Science. 2014;54(1):74-84.
56.   Corbet N, Burns B, Johnston D, Wolcott ML, Corbet D, Venus B, et al. Male traits and herd
reproductive capability in tropical beef cattle. 2. Genetic parameters of bull traits. Animal Production
Science. 2012;53(2):101-13.
57.   Canal LB, Fontes PL, Sanford CD, Mercadante VR, DiLorenzo N, Lamb GC, et al. Relationships
between feed efficiency and puberty in Bos taurus and Bos indicus-influenced replacement beef
heifers. Journal of Animal Science. 2020;98(10):skaa319.
58.   Perry GA, Cushman R. Effect of age at puberty/conception date on cow longevity. Veterinary
Clinics: Food Animal Practice. 2013;29(3):579-90.
59.   Bunter KL, Cai W, Johnston DJ, Dekkers JC, Bunter K. Selection to lower residual feed intake
in pigs produces a correlated response in juvenile insulin-like growth factor-I (IGF-1) concentration.
60.   Gao X, Xu X-R, Ren H-Y, Zhang Y-H, Xu S-Z. The effects of the GH, IGF-I and IGF-IBP3 gene on
growth and development traits of Nanyang cattle in different growth period. Yi Chuan= Hereditas.
2006;28(8):927-32.
61.   Bolormaa S, Pryce JE, Reverter A, Zhang Y, Barendse W, Kemper K, et al. A multi-trait, meta-
analysis for detecting pleiotropic polymorphisms for stature, fatness and reproduction in beef cattle.
PLoS genetics. 2014;10(3):e1004198.
62.   Hackinger S, Zeggini E. Statistical methods to detect pleiotropy in human complex traits.
Open biology. 2017;7(11):170125.
63.   Id-Lahoucine S, Molina A, Cánovas A, Casellas J. Screening for epistatic selection signatures:
A simulation study. Scientific reports. 2019;9(1):1-5.
64.   Crowley J, Evans R, Mc Hugh N, Kenny D, McGee M, Crews Jr D, et al. Genetic relationships
between feed efficiency in growing males and beef cow performance. Journal of animal science.
2011;89(11):3372-81.
65.   Purfield DC, Evans RD, Berry DP. Breed-and trait-specific associations define the genetic
architecture of calving performance traits in cattle. Journal of animal science. 2020;98(5):skaa151.
66.   Nguyen T-M, Shafi A, Nguyen T, Draghici S. Identifying significantly impacted pathways: a
comprehensive review and assessment. Genome biology. 2019;20(1):1-15.
67.   Bozza WP, Zhang Y, Hallett K, Rosado LAR, Zhang B. RhoGDI deficiency induces constitutive
activation of Rho GTPases and COX-2 pathways in association with breast cancer progression.
Oncotarget. 2015;6(32):32723.
68.   Hall A. Rho family gtpases. Biochemical Society Transactions. 2012;40(6):1378-82.
69.   Humphries B, Wang Z, Yang C. Rho GTPases: Big Players in Breast Cancer Initiation,
Metastasis and Therapeutic Responses. Cells. 2020;9(10):2167.
70.   Vega FM, Ridley AJ. Rho GTPases in cancer cell biology. FEBS letters. 2008;582(14):2093-101.
71.   Shibata S, Nagase M, Yoshida S, Kawarazaki W, Kurihara H, Tanaka H, et al. Modification of
mineralocorticoid receptor function by Rac1 GTPase: implication in proteinuric kidney disease.
Nature medicine. 2008;14(12):1370-6.

72.   Togawa A, Miyoshi J, Ishizaki H, Tanaka M, Takakura A, Nishioka H, et al. Progressive
impairment of kidneys and reproductive organs in mice lacking Rho GDIα. Oncogene.
1999;18(39):5373-80.
73.   Ishizaki H, Togawa A, Tanaka-Okamoto M, Hori K, Nishimura M, Hamaguchi A, et al.
Defective chemokine-directed lymphocyte migration and development in the absence of Rho
guanosine diphosphate-dissociation inhibitors α and β. The Journal of Immunology.
2006;177(12):8512-21.
74.   Huta Y, Nitzan Y, Breitbart H. Ezrin protects bovine spermatozoa from spontaneous
acrosome reaction. Theriogenology. 2020;151:119-27.
75.   Shi Z-H, Zhao C, Wu H, Liu X-M. Expression of RhoGDI alpha in human testes and sperm and
its correlation with the success rate of IVF. Zhonghua nan ke xue= National Journal of Andrology.
2011;17(4):325-9.
76.   Wang L, Chen W, Zhao C, Huo R, Guo X-J, Lin M, et al. The role of ezrin-associated protein
network in human sperm capacitation. Asian journal of andrology. 2010;12(5):667.
77.   Burns B, Fordyce G, Holroyd R. A review of factors that impact on the capacity of beef cattle
females to conceive, maintain a pregnancy and wean a calf—Implications for reproductive efficiency
in northern Australia. Animal Reproduction Science. 2010;122(1-2):1-22.
78.   McLean I, Holmes P, Counsell D. Final Report: The Northern beef report, 2013 Northern beef
siutation analysis (B. COM. 0348). Meat and Livestock Australia, Sydney. 2013.
79.   Chang AZ, Swain DL, Trotter MG. Calf loss in northern Australia: a systematic review. The
Rangeland Journal. 2020;42(1):9-26.
80.   Fordyce G. Pregnancy rates achieved by mating bulls with different percentages of
morphologically normal sperm. Bullpower Delivery of adequate normal sperm to site of fertilisation
Project NAP3. 2005;117:142-51.
81.   Rosen BD, Bickhart DM, Schnabel RD, Koren S, Elsik CG, Tseng E, et al. De novo assembly of
the cattle reference genome with single-molecule sequencing. GigaScience. 2020;9(3):giaa021.
82.   Misztal I, Tsuruta S, Strabel T, Auvray B, Druet T, Lee D, editors. BLUPF90 and related
programs (BGF90). Proceedings of the 7th world congress on genetics applied to livestock
production; 2002.
83.   Strandén I, Garrick D. Derivation of equivalent computing algorithms for genomic predictions
and reliabilities of animal merit. Journal of dairy science. 2009;92(6):2971-5.
84.   Aguilar I, Legarra A, Cardoso F, Masuda Y, Lourenco D, Misztal I. Frequentist p-values for
large-scale-single step genome-wide association, with an application to birth weight in American
Angus cattle. Genetics Selection Evolution. 2019;51(1):1-8.
85.   VanRaden PM. Efficient methods to compute genomic predictions. Journal of dairy science.
2008;91(11):4414-23.
86.   Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs
explain a large proportion of the heritability for human height. Nature genetics. 2010;42(7):565-9.
87.   Bozeman M. Golden Helix, Inc. SNP & Variation Suite™[Software]. [(Version 8.x)].

Available from http://www.goldenhelix.com.
88.   Hu Z-L, Park CA, Reecy JM. Developmental progress and current status of the Animal QTLdb.
Nucleic acids research. 2016;44(D1):D827-D33.
89.   Hu Z-L, Park CA, Wu X-L, Reecy JM. Animal QTLdb: an improved database tool for livestock
animal QTL/association data dissemination in the post-genome era. Nucleic acids research.
2013;41(D1):D871-D9.

90.   Lam S, Miglior F, Fonseca P, Gómez-Redondo I, Zeidan J, Suárez-Vega A, et al. Identification
of functional candidate variants and genes for feed efficiency in Holstein and Jersey cattle breeds
using RNA-sequencing. Journal of dairy science. 2021;104(2):1928-50.
91.   Sweett H, Fonseca P, Suarez-Vega A, Livernois A, Miglior F, Cánovas A. Genome-wide
association study to identify genomic regions and positional candidate genes associated with male
fertility in beef cattle. Scientific reports. 2020;10(1):1-14.
92.   Krämer A, Green J, Pollard Jr J, Tugendreich S. Causal analysis approaches in ingenuity
pathway analysis. Bioinformatics. 2014;30(4):523-30.
93.   Medici V, Kieffer DA, Shibata NM, Chima H, Kim K, Canovas A, et al. Wilson Disease:
Epigenetic effects of choline supplementation on phenotype and clinical course in a mouse model.
Epigenetics. 2016;11(11):804-18.
94.   Cardoso TF, Quintanilla R, Castelló A, González-Prendes R, Amills M, Cánovas Á. Differential
expression of mRNA isoforms in the skeletal muscle of pigs with distinct growth and fatness profiles.
BMC genomics. 2018;19(1):1-12.
95.   Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, et al. BioMart and
Bioconductor: a powerful link between biological databases and microarray data analysis.
Bioinformatics. 2005;21(16):3439-40.