Number and proportion of *P. falciparum* gametocytes vary from acute infection to

chronic parasite carriage despite unaltered sexual commitment rate

Hannah van Dijk[1¶], Nathalia F Lima[1¶], Martin Kampmann[2], Michael Gabel[3], Usama Dabbas[1,2],

Safiatou Doumbo[4], Hamidou Cisse[4], Shanping Li[5], Myriam Jeninga[6], Richard Thomson-

Luque[1], Didier Doumtabe[4], Michaela Petter[6], Kassoum Kayentao[4], Aissata Ongoiba[4], Teun

Bousema[7], Peter D Crompton[5], Boubacar Traore[4], Frederik Graw[3,8], Silvia Portugal[1,2*]

[1] Center for Infectious Diseases, Parasitology, Heidelberg University Hospital, Heidelberg,

Germany

[2] Max Planck Institute for Infection Biology, Berlin, Germany

[3] BioQuant-Center for Quantitative Biology, Heidelberg University, Heidelberg, Germany

[4] Mali International Center of Excellence in Research, University of Sciences, Techniques and

Technologies of Bamako, Mali

[5] Laboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National

Institutes of Health, Rockville, Maryland, USA

[6] Mikrobiologisches Institut-Klinische Mikrobiologie, Universitätsklinikum Erlangen

[7] Radboud University Medical Center, Nijmegen, the Netherlands

[8] Interdisciplinary Center for Scientific Computing, Heidelberg University, Heidelberg, Germany

[¶]These authors contributed equally to this work

[*]Corresponding author email: portugal@mpiib-berlin.mpg.de (SP)

Abstract

In areas of seasonal malaria, resuming transmission every wet season relies on persistent

asexual parasites during dry season that maintain the ability to produce gametocytes until the

mosquito population resurges, following the start of annual rains. Although human

asymptomatic *P. falciparum* reservoirs in the dry season are widely recognized, the

longitudinal dynamics of parasite sexual commitment and gametocytogenesis are unclear.

Here, we compared the density and proportion of *P. falciparum* gametocytes in blood of

asymptomatic subjects during the dry season versus subjects with symptomatic malaria in the

wet season. Blood concentrations of phospholipids potentially involved in gametocytogenesis

and parasite transcriptomes were also compared. While blood densities of gametocytes and

asexual parasites were lower during the dry season, we found that gametocytes were

proportionally higher during the dry versus wet season. Levels of parasite transcripts involved

in sexual commitment were unaltered throughout the year, and phospholipid content in the

plasma throughout the year did not affect sexual commitment in vitro. Our experimental data

and mathematical modelling support that gametocyte density and proportion diverge as

infections progress from recently transmitted to chronic carriage, without significant alterations

in the rate of sexual commitment over time.

Introduction

It has long been known that the malaria-causing parasite *Plasmodium falciparum* requires a
mosquito host to complete its lifecycle and to be transmitted between humans (1). However, it
remains unclear if in the human host the parasite can dynamically allocate resources to the
differentiation of sexual-stage gametocytes that are transmitted to mosquitos, particularly in
areas of seasonal malaria where the size of the mosquito population fluctuates dramatically
during the dry and wet seasons. During blood feedings, infected Anopheles mosquitoes
deposit *P. falciparum* sporozoites in the human dermis that find their way into circulation,
reaching the liver where they infect hepatocytes. After 7-10 days of parasite replication, each
infected hepatocyte releases thousands of merozoites into the blood stream (2), where they
start continuous asexual intra-erythrocytic replication cycles of ~48 h, from merozoite invasion
of erythrocytes, through the ring- and trophozoite-stages, until the multinucleated schizont
releases new merozoites (3). A small proportion of intra-erythrocytic asexual parasites
differentiate into sexual-stage gametocytes (4). Instead of another ~48 h asexual schizogony
leading to the 8-30 new merozoites ready to continue asexual cycles (5), the parasite develops
into sexually committed schizonts leading to 8-30 sexually committed merozoites that upon
infection of new red blood cells (RBCs) each produce one stage I gametocyte (6, 7) that
through a period of 8-12 days, will go through five developmental stages (8). Stages II to IV of
gametocytaemia take place in the bone marrow or spleen (9, 10), while stage V gametocytes
are again present in peripheral blood, each lasting up to ~20 days (11, 12) waiting for another
mosquito bite. Following a blood meal gametocytes encounter the right environment for sexual
reproduction in the mosquito midgut (13). They transform into female and male gametes that
fuse into a zygote originating oocysts that invade the midgut wall to generate sporozoites that
can, after reaching the mosquito salivary glands, restart the parasite lifecycle in the next blood
meal (14). The continued presence of mosquitoes and humans in a geographical region allows
for perennial *P. falciparum* transmission; however, in areas where the dry season is several

months long, the mosquito population can plummet to near zero (15), thereby interrupting

transmission. *P. falciparum* faces the added challenge of surviving in its human host for

months while waiting for the return of mosquitoes with the ensuing wet season (16-18). During

mosquito-free dry periods, gametocyte production is largely wasted, as gametocytes cannot

survive to bridge two wet seasons (11, 12). Although curtailing gametocyte production rates

during the dry season would seem beneficial to the parasite, such a strategy would require a

mechanism to reactivate gametocytogenesis with the return of rain and mosquitos. Host

factors such as age (19), haematocrit (20), haemoglobin type (21) and duration of malaria

symptoms (22) have been linked to gametocytaemia in individuals with clinical malaria; and

higher frequencies of gametocyte positive individuals have been observed in chronic

asymptomatic infections compared to acute infections (23-25). Additionally, various in vitro

conditions have been shown to modulate sexual commitment and thus gametocytogenesis (6,

26-28). The specific effect of lysophosphatidylcholine (Lyso-PC) in repressing the pro-sexual

commitment transcription factor AP2G has been recently identified, with low Lyso-PC

concentrations increasing the proportion of sexual forms in vitro (28). Furthermore,

gametocyte development 1 (GDV1) has been identified as the de-repressing effector which

counters the epigenetic silencing and promotes AP2G expression leading to the induction of

gametocytogenesis (29), and homeodomain protein 1 (HDP1) was recently identified as a

positive transcriptional regulator essential for gametocyte development (30). Still, it is not fully

understood how sexual investment and gametocytaemia is regulated in vivo, and if or how

transmission to mosquitoes can be optimized by the parasite to conserve resources. Recently,

we have shown that *P. falciparum* parasites isolated at the end of the dry season are

transcriptionally distinct from those of subjects with febrile malaria in the transmission season

(16, 31), but alteration of sexual commitment and gametocytogenesis were not investigated.

The observed transcriptional differences reflected longer parasite circulation within each ~48 h

intra-erythrocytic cycle in the dry season, likely increasing splenic clearance of infected cells

and thus maintaining the low parasitaemia observed during the dry season (16). Here, we

aimed to examine sexual commitment and gametocytogenesis of parasites persisting through

the dry season and explored possible parasite adaptations during mosquito-free periods. We

quantified sexual commitment and female and male gametocyte densities in blood samples

collected from children at different times of the year and with different clinical presentations,

and we related them to total parasite density and infection length. Furthermore, we determined

longitudinally the Lyso-PC concentration in plasma of children, and assessed its ability to

induce *P. falciparum* sexual commitment and gametocytogenesis in vitro. Finally, we adapted

a previously described within-host mathematical model that considers asexual parasites,

sexually committed parasites, and gametocytes (32) to infer the dynamics of gametocyte

number and proportion as *P. falciparum* infections progress from recently transmitted to

chronic asymptomatic carriage over several months.

Results

Sexual stages differ between asymptomatic *P. falciparum* at the end of the dry season

and *P. falciparum* causing clinical malaria in the wet season. We first investigated the

gametocyte-related transcriptome (33) within a whole genome RNA sequencing dataset (16)

comparing persistent *P. falciparum* detected by rapid diagnostic test (RDT) in 12

asymptomatic Malian children at the end of the dry season (May) to *P. falciparum* of 12 age-

and gender- matched Malians presenting clinical malaria with higher parasitaemias in the wet

season (MAL). Principal component analysis of 333 gametocyte-specific genes with detectable

number of reads (out of the 786 defined as gametocyte-related by ((33), S1 Table) showed

segregation of transcription profiles based on seasonality (Fig 1A). Differentially expressed

genes (DEGs) determined at a false discovery rate threshold of 5% resulted in 205 DEGs, with

146 transcripts up-regulated, and 59 transcripts down-regulated in the dry season samples

compared to clinical malaria samples (Fig 1B and S1 Table). These transcriptional differences

prompted us to quantify female and male gametocytes in the blood of the two groups of

children collected in the same year and the same cohort study. First, we determined total

parasite (PFGLY, PF3D7_1420400) density by qRT-PCR, and observed that it was higher in

clinical malaria cases in the transmission season (MAL) than in asymptomatic cases at the

end of the dry season (May) (Fig 1C). Then, we determined the density of female (P25,

PF3D7_1031000) and male (PFMGET, Pf3D7_1469900) gametocytes, based on a standard

curve obtained from sorted gender-specific fluorescent gametocytes (34) (S1 Fig). This

analysis revealed higher numbers of both female and male gametocytes per mL of blood in

clinical cases of malaria during the wet season (MAL) compared to asymptomatic carriers at

the end of the dry season (May) (Fig 1D). However, when we quantified either gametocyte

transcript relative to the total *P. falciparum* level present in each subject's sample, normalized

to a housekeeping gene transcribed by all parasite stages, we observed an inverse trend.

There were proportionally fewer female and male gametocytes in malaria cases in the wet

season (MAL) than at the end of the dry season (May), although the difference was significant

for the female comparison only (Fig 1E). We also quantified phospholipids known to influence

P. falciparum sexual commitment in vitro (28) in the plasma of the same children through

shotgun lipidomic analysis. Malaria cases in the wet season (MAL) showed lower levels of

several species of Lyso-PC (Fig 1F) and lysophosphatidylethanolamine (Lyso-PE) (Fig 1G)

families compared to asymptomatic carriers at the end of the dry season (May), which could

potentially increase *P. falciparum* sexual commitment in malaria cases (MAL). Nevertheless,

transcription of the sexual commitment regulators (AP2G, PF3D7_1222600; and GDV1,

PF3D7_0935400), sexually committed schizonts (MSRP1, PF3D7_1335000), or early

gametocyte marker (GEXP5, PF3D7_0936600) were not significantly different by qRT-PCR

(Fig 1H) nor were they differentially expressed in the abovementioned RNAseq analysis (S1

Table).

Fig 1. Sexual stages differ between subclinical P. falciparum at the end of the dry season and *P. falciparum* causing clinical malaria in the transmission season. A: Principal components analysis of RNA-Seq data of 333 gametocyte specific *P. falciparum* genes from parasites collected at the end of the dry season and from clinical malaria cases (n= 12 May, 12 MAL). **B**: Heatmap showing normalized reads of 205 gametocyte specific *P. falciparum* of differentially expressed genes (DEGs) (rows) for each subject (columns) from *P. falciparum* collected at the end of the dry season (May) and at the first clinical malaria case (MAL) in the ensuing transmission season. **C:** Asexual parasite density detected by qRT-PCR of RDT+ subclinical children at the end of the dry season (May = 35) and of children with their first clinical malaria episode (MAL = 27) in the wet season. **D:** Female (left) and male (right) gametocyte densities detected by qRT-PCR of RDT+ subclinical children at the end of the dry season (May = 35) and of children with their first clinical malaria episode (MAL= 27) in the wet season. **E:** Proportion of female (left) and male (right) gametocytes detected by qRT-PCR of RDT+ subclinical children at the end of the dry season (May) and of children with their first clinical malaria episode (MAL) in the wet season. **F:** Plasma concentration of Lyso-PC species

of RDT+ subclinical children at the end of the dry season (May) and of children with their first
clinical malaria episode (MAL) in the wet season. **G:** Plasma concentration of Lyso-PE species
of RDT+ subclinical children at the end of the dry season (May) and of children with their first
clinical malaria episode (MAL) in the wet season. **H:** Relative expression of AP2G, GDV1,
MSRP1, and GEXP5, detected by qRT-PCR of RDT+ subclinical children at the end of the dry
season (May n > 15) and of children with their first clinical malaria episode (MAL n > 23) in the
wet season. Lipid data represented as median ± IQR; two-way ANOVA with Tukey multiple
comparisons correction. Parasitaemia and gametocytaemia and AP2G, GDV1, MSRP1, and
GEXP5 expression data represented as median ± IQR; Mann-Whitney test.
170
Sexual commitment and gametocytogenesis dynamic in persisting infections. We then

determined the dynamics of *P. falciparum* sexual commitment and gametocytogenesis in

persistently infected individuals, and examined associations between plasma Lyso-PC and

Lyso-PE, monitoring longitudinally ~20 children who were continuously infected from the wet

season until the end of the dry season. We quantified the number of total parasites, and fully

formed female and male gametocytes by qRT-PCR at four different time points in

asymptomatic children persistently infected since the wet season (Oct) through the dry season

until its end (Jan, Mar and May); and in a parallel group of age-matched individuals presenting

with clinical malaria during the ensuing wet season (MAL). We observed that total parasite

density was higher in malaria cases (MAL), than at any time-point of the asymptomatic

infections, whether in the wet (Oct) or the dry season (Jan, Mar and May), while no differences

were found between any of the time points of asymptomatic infections (Oct, Jan, Mar and

May) (Fig 2A). Similarly, we observed higher densities of both female and male gametocytes

in malaria cases (MAL) than at any time point of the asymptomatic infections (Oct, Jan, Mar

and May), and again no differences within the persisting period (Fig 2B). Conversely, the

proportion of male gametocytes was higher at later time points of persisting infections (Mar or

May) than in malaria cases during the transmission season (MAL) (Fig 2C). Moreover, we

found no significant difference in the sexual commitment regulator AP2G between incident

(MAL) or persistent infections at any point analyzed (Oct, Mar and May) (Fig 2D).

Quantification of Lyso-PC 18:0 in malaria cases compared to asymptomatic infections

progressing from the wet into the dry season again resulted in decreased Lyso-PC values in

clinical malaria cases (MAL) compared to asymptomatic infections (Oct, Mar and May) (Fig

2E). We further detected a decrease of Lyso-PC from the wet season (Oct) to the end of the

dry season (May), but no difference was seen between the two available time points of the dry

season (Mar and May) (Fig 2E). Furthermore, Lyso-PC 18:0 levels in persistent infections

(Oct, Mar and May) were comparable to non-infected individuals at the end of the dry season

(May$^-$), indicating that asymptomatic infections have no impact on Lyso-PC 18:0 (Fig 2E and

S2 Fig). Accordingly, a relation between Lyso-PC levels and parasitaemia or gametocytaemia

was only significant when considering clinical malaria plus all asymptomatic samples (Oct,

Mar, May and MAL) (r=−0.68 p< 0.0001; r=−0.56 p< 0.008 and r=−0.67 p< 0.0001 for

parasitaemia, female and male gametocytaemia respectively), but disappeared when each

group was analyzed independently, or in all persisting asymptomatic infections analyzed

together (Fig 2F).

204
Fig 2. Sexual commitment and gametocytogenesis dynamic in persisting infections. A:
Parasite, **B:** Female (left) and male (right) gametocyte, and **C:** Proportion of female (left) and
male (right) gametocytes densities detected by qRT-PCR of RDT$^+$ subclinical children at
different times during the wet (Oct n = 10) and the dry season (Jan, Mar and May n > 15), and
during clinical malaria episodes (MAL n = 19). **D:** Relative expression of AP2G detected by
qRT-PCR of RDT$^+$ subclinical children at different times during the wet (Oct n = 6) and the dry
season (Jan, Mar and May n > 11), and during clinical malaria episodes (MAL n = 19). **E:**
Plasma concentration of Lyso-PC 18:0 of RDT$^+$ subclinical children at different times during the
wet (Oct n = 17) and the dry season (Mar and May n = 19), and during clinical malaria
episodes (MAL n = 15). Parasite gene expression and lipid data were compared through
Kruskal-Wallis test with multiple comparisons. **F**: Correlation between Lyso-PC 18:0 and
parasite (left), female gametocyte (canter) or male gametocyte (right) densities detected by
qRT-PCR detected by qRT-PCR of RDT$^+$ subclinical children at different times during the wet
(Oct n = 10) and the dry season (Mar and May n > 15), and during clinical malaria episodes
(MAL n = 19). *P* and r determined by Pearson correlation.

Figure 2

Decreased Lyso-PC during clinical malaria does not induce higher sexual commitment or gametocytaemia. We then questioned if the low levels of Lyso-PC and Lyso-PE species in plasma of malaria cases compared to persistent infections at the end of the dry season (Fig 1F and 2E) could induce *P. falciparum* sexual conversion or gametocytaemia differently in vitro. We exposed parasites of the *P. falciparum* 2004 line (Pf2004) to plasma of children collected during clinical malaria cases (MAL) versus those of children with asymptomatic infections at the end of the dry season (May) alongside the conditions described by Brancucci et al. to induce or inhibit sexual convertion and gametocyteamia. As reported earlier (28), we observed that 24 h after culturing Pf2004 without plasma or serum (noPI), and thus no Lyso-PC, over minimal fatty acid medium (MFA), AP2G expression increased compared to the same parasites cultured in the presence of German control plasma (PIDE) (Fig 3A). However, MFA

medium supplementation with 25% plasma from children presenting with clinical malaria

during the wet season (PlMAL), versus plasma from children with asymptomatic infections at

the end of the dry season (PlMay) resulted in similar APG2 induction 48 h after beginning of

treatment (Fig 3A), and similar conversion rate of parasites into gametocytes on day 5 after

treatment (defined by = d5gametocytes/ d2parasites) (Fig 3B). Absence of plasma (noPl)

resulted in higher gametocytaemia than observed on the German pooled plasma condition

(PIDE) (Fig 3B), plasmas from malaria cases (PlMAL) or asymptomatic infections at the end of

the dry season (PlMay) revealed similar conversion rates (Fig 3B). Although the 2-4 fold Lyso-

PC differences between non-infected or asymptomatic children and malaria cases were

significant (Fig 1F-G and Fig 2E), these were much lower than the several orders of magnitude

included in the standard curve used to establish that gametocyte conversion responds to low

Lyso-PC in the culture medium (28), and were thus likely insufficient to impose alterations in

sexual commitment or gametocytaemia, as seen by similar AP2G induction and gametocyte

conversion in vitro (Fig 3).

247
248
249
Fig 3. Decreased Lyso-PC of clinical malaria cases does not induce higher sexual
commitment or gametocytaemia in vitro. A: Relative expression of AP2G detected by qRT-
PCR of Pf2004 24 h after 24 h culture with 25% plasma collected during clinical malaria
episodes (MAL n = 21), or at the end of the dry season (May n = 19). **B**: Pf2004 conversion
rate (gametocytes d5/parasites d2) (left) and gametocyte density (right) detected by
microscopy, after 24 h culture with 25% plasma collected during clinical malaria episodes in
the wet season (MAL n = 19) or at the end of the dry season (May n = 18), alongside no
plasma (noPl) and German (PIDE) conditions. Data represented as median ± IQR; Mann-
Whitney test.
Progressing infection leads to apparent differences in gametocytogenesis. We observed

decreased densities (Fig 1D and Fig 2B) but higher proportions (Fig 1E and Fig 2C) of female

and male gametocytes in the dry versus wet season, or in asymptomatic versus clinical

malaria in the wet season (Fig 2C), without concomitant differences in the expression of

regulators of sexual commitment, or transcriptional markers of sexually committed schizonts or

early gametocytes (Fig 1H and Fig 2E and S1 Table). To further investigate this apparent

conflict, we first ascertained if there were indeed indications of altered gametocytogenesis

between parasites persisting through the dry season and parasites causing malaria in the wet

season. Specifically, we investigated in more detail the transcriptional profiles of early and

late-stage gametocytes. Using markers of early or late gametocyte development gathered

from different sources ((35, 36) and Jeninga et al., unpublished Feb 2022) and intersecting

them with the RNAseq data of clinical malaria cases and dry season samples shown in Fig 1,

we identified 159 gametocyte-specific transcripts with quantifiable reads in samples from

individuals with circulating parasites persisting through the dry season and parasites causing

malaria in the wet season. Of those, 105 transcripts define early gametocytes; and 54

transcripts are characteristic of late-stage gametocytes (highlighted in S1 Table). Expression

of these transcripts in asymptomatic carriers at the end of the dry season (May) and malaria

cases (MAL) in the wet season showed an enrichment of upregulated transcripts of late-stage

gametocytes in the dry season (81,48%) (44/54 genes), which clearly contrasted with the

proportion of early-gametocyte transcripts or total *P. falciparum* transcripts upregulated at that

time, which showed (39,05% (41/105 genes) and 33,4% (1131/3381 genes) upregulated in the

dry season respectively (Fig 4A).

The lack of difference in sexual commitment or early-stage gametocyte transcripts, along with

the strong difference in late-stage gametocyte transcript between the end of the dry season

(May) malaria cases (MAL) parasitaemias hints not at a difference in gametocyte production,

but rather an effect during the 8-12 days of gametocyte development, in both/either the sexual

and/or the asexual parasite compartments, suggesting that length of infection may play a role

in the results observed. Hence, we investigated if the observed patterns of gametocyte

densities and proportions were generalizable to non-seasonal settings where infections can

also persist. While Mali shows a very contrasting seasonal pattern with 5-6 months interruption

of transmission every year (15), Uganda has uninterrupted year-round transmission with peaks

in times of stronger rain (37). RT-qPCR data quantifying transcripts of total parasite (varATS)

and gametocyte (P25, PF3D7_1031000) densities from samples of 150 Ugandan individuals

with asymptomatic parasitaemias persisting for over one month (n=122) and age-matched

individuals presenting with clinical malaria (n=38) had been previously published (37) and

were retrieved from ClinEpiDB (38). The data, determined by the area under the curve over

the course of asymptomatic infection divided by the time of infection, indicated that parasite

and gametocyte densities were increased in clinical malaria cases (MAL) compared to

asymptomatic infections (Asymp) (Fig 4B and C); while the proportion of gametocyte

transcripts relative to total *P. falciparum* transcripts showed again an inverse trend (Fig 4D).

As seen with the Malian samples, the proportion of gametocytes was higher in persisting

parasitaemias (Asymp) compared to malaria cases (MAL) (Fig 4D). Malian and Ugandan

samples vary in seasonality of transmission, but share the fact that clinical malaria cases are

likely recently transmitted (39), while asymptomatic parasitaemias have persisted longer than

one month in the Uganda cohort, and at least 5 months in the Malian setting. The parallel in

infection length between samples of symptomatic clinical malaria and asymptomatic infections

in the two contexts led us to hypothesize that gametocyte numbers and proportions change

dynamically in the course of a *P. falciparum* infection, leading to the observed patterns in Mali

and Uganda. To address this, we devised a mathematical model including three different

parasite populations_ asexual parasites (P), sexually committed parasites (P_G) and

gametocytes (G), all of which can circulate or be adherent as previously developed (32) (Fig

4E). As the original model by Cao et al. was not able to sufficiently explain the observed

dynamics when fitted to the data, we extended the model in different ways. First, we

incorporated our recent observation of increased clearance of longer-circulating infected

erythrocytes in persisting parasitaemias compared to recently transmitted parasites (16),

which promoted different removal rates of asexual and sexually-committed parasites during

distinct phases of acute and chronic *P. falciparum* infection in our model (see Methods and

Source code 1). In addition, we distinguish between three phases *during P. falciparum*

infection in which the parasite multiplication factor varies due to immune response or adhesive

properties of the parasite. Using an iterative Bayesian fitting-approach to fit the model to the

symptomatic and asymptomatic parasitaemia and gametocytaemia levels of the Malian cohort,

we quantitatively assessed several biological parameters including sexual commitment rate,

gametocyte circulation time, parasite multiplication rate and gametocyte sequestration time,

thereby confirming previously determined ranges (32, 40) (S3 Fig). Our adapted model was

able to provide a dynamic depiction of parasitaemia levels, and gametocyte densities and

proportions, as we observed within the Malian data (Fig 4F-H). We defined that the death rate

of asexual and sexually committed parasites increased in later stages of the 48 h cycle,

shifting toward the removal of older parasite stages and leading to a constant but oscillating

number of asexual parasite in circulation (highlighted in the box-inlay of Fig 4F). This can, to

some extent, explain the large variation seen in parasitaemias, especially during the chronic

phase, which additionally may arise from different collection time points after start of infection

for the individual samples (i.e., pooling of the individual measurements from Oct-May and

calculation of average for each individual by the area under the curve over the course of blood

collections during the asymptomatic phase and fixing it to one time point for fitting (see

Methods)). Keeping a constant sexual commitment rate, f, throughout acute and chronic

infection phases, our model predicts a higher number and lower proportions of gametocytes in

the acute phase_ a trend that is inverted during the transition phase of infection due to

effective removal of asexual parasites by an active immune response (Fig 4G and H). During

the chronic phase of infection, when parasitaemias remain constant or are slowly decreasing,

gametocyte proportions are decreasing as well until reaching an equilibrium roughly 4-5

months after the start of blood stage infection. Overall, our model provides a dynamic

depiction of gametocyte densities and proportions similar to the data we observed in Mali and

in Uganda, without altering the sexual commitment rate (Fig 4C and D).

342
343
344
Fig 4 | Progressing infection leads to apparent differences in gametocytogenesis. A,
Fold change between parasites collected at the end of the dry season and from clinical malaria
cases (May n= 12, MAL n= 12) of early and late gametocyte transcripts according to López-
Barragán et al., 2011; Poran et al., 2017 and Petter unpublished data. Each dot represents a
gene and color identifies range in fold change, red: upregulated in May, yellow: not DEG, and
blue: downregulated in May). **B**, Parasitaemia, **C**, gametocytaemia and **D**, proportion of
gametocyte in symptomatic (MAL n = 19 and 38 in Mali and Uganda respectively) and
asymptomatic infections (May n = 23 and Asymp n = 122) in Mali and Uganda quantified by
RT-qPCR. Data represented as median ± IQR; multiple non-parametric test with Tukey-
correction for multiple comparisons. **E**, Diagram of the mathematical model to describe
parasite dynamics including individual compartments, transitions and death rates; Asexual
parasites (P) replicate in a 48 h cycle, being cleared at an age dependent death rate (σ_a). A
fraction of asexual parasites (f) transitions to the sexually committed parasite compartment
(P_G), goes through a round of replication and develops into gametocytes (G). Gametocytes
mature over 5 distinct stages in independent model compartments. Asexual parasites (blue),

sexually committed parasites (orange) and gametocytes (grey) are indicated shaded if not
circulating due to sequestration. **F**, Model prediction and clinical data of asexual parasites. The
small inlay indicates a zoom-in showing the oscillating parasite dynamics within 48 h-intervals.
G, gametocytes and **h**, the proportion of gametocytes per mL blood. Individual data points and
median (black square) of wet (aqua circles) and dry (orange circles) season data throughout
acute (aqua shaded area) and chronic (salmon shaded area) phase of infection are shown.
Median fit (grey line) and 95% prediction interval of 100 model predictions of the 6th PyABC
generation are shown as grey shaded area.
368

Discussion

The relative absence of mosquitoes during the months-long dry season and the short lifespan

of *P. falciparum* sexual-stage gametocytes (11, 12) renders the production of gametocytes

ineffective during most of the dry season; however, gametocytes must be available to resume

malaria transmission when mosquitos return with the wet season. Whether *P. falciparum*

maintains a constant rate of production of sexual forms or adjusts gametocytogenesis to

maximize transmission has been unclear. In this study, with samples from Malians exposed to

alternating 6-month dry and wet seasons, and samples from clinical malaria and asymptomatic

P. falciparum infection from year-round transmission region of Uganda, we showed that

symptomatic cases of malaria consistently have higher parasite and stage V gametocyte

densities than asymptomatic infections; however, the proportion of gametocytes increases in

the asymptomatic persisting infections (Fig 1, 2 and 3). Additionally, our data from Mali

indicated that transcription of genes characterizing sexually committed parasites and stage I

gametocytes (41, 42) quantified by RT-qPCR did not appear affected by the season or clinical

symptoms (Fig 1 and 2), nor were reads of over 100 genes characteristic of early gametocytes

and detected by RNAseq of circulating parasites particularly enriched for DEGs. This, clearly

contrast with representative expressed genes of late gametocytes that were very much

enriched of transcripts upregulated in persisting asymptomatic infections at the end of the dry

season (Fig 4 and S1 Table).

Our work suggests that gametocytogenesis remains constant as infections progress from

recent transmission to chronic carriage, and it is the difference in asexual survival/replication

rates that promotes different gametocyte densities, independently of environmental factors.

Although it has been suggested that *P. falciparum* investment in gametocyte production

adjusts to host nutrient availability (43), drug pressure (22), transmission intensity (44), and

even seasonality (45, 46), results are often inconsistent and potentially affected by the

dynamics of asexual parasite growth, without necessarily indicating altered sexual

commitment. In Sudan, following the long dry season, gametocyte density estimates have

been reported to increase prior to the highest transmission peak (47), suggesting a higher

gametocyte investment. However, in Burkina Faso, gametocyte densities were higher at the

peak and end of the transmission season rather than during the dry season (48), while the

efficiency of membrane feeding to infect mosquitoes was increased early in the transmission

season compared to peak of the malaria season or the dry season (49). Nevertheless, these

studies did not monitor longitudinally the same parasite population during the dry and early wet

season. In a recent study also in Burkina Faso, transmission was shown to be more effective

in persistent than in incident infections, but no data was collected after day 35, and collections

were all done during the wet season (25) when a mix of more and less recently transmitted

parasite clones is likely in individuals without symptoms.

In vitro, depletion of Lyso-PC or choline have been shown to induce AP2G expression, leading

to increased gametocyte production (28, 50), but associations in humans are inconsistent. In

Rwanda, low Lyso-PC associated with higher parasite densities (51); and in Ghana, although

correlations with parasitaemia were not reported, Lyso-PC levels in human blood samples

were inconsistently related with gametocyte-committed ring stages converting into

gametocytes in vitro (42). Our data show, like Orikiiriza et al., an association of lower Lyso-PC

levels with increased parasitaemia and severity of malaria (Fig 1F-G and 2E), but the

association was lost at low parasitaemias, and persistent infection at the end of the dry season

did not affect Lyso-PC compared to non-infected individuals (Fig 2F and S2 Fig).

The effect of drugs on sexual conversion in vitro has been previously reported. Specifically,

sub-curative doses of dihydroartemisinin (DHA) stimulate AP2G expression and sexual

conversion when drug exposure occurs at the trophozoite stage, but not at the early ring

stage. The observed increase could not synergize nor add to increases promoted by depletion

of choline, suggesting, as DHA affects lipids, that the two effects might be the same (50). Our

RNAseq (S1 Table) and RT-qPCR (Fig 1H and 2D) data show constant low-level expression

of AP2G, the master regulator of gametocytogenesis, and equally unaltered expression of

transcripts characteristic of committed schizont and early gametocytes (S1 Table and Fig 1H),

suggesting that reductions in circulating Lyso-PC induced by uncomplicated clinical malaria

are not capable of affecting *P. falciparum* sexual commitment. Accordingly, we observed in

vitro that supplementation with plasma from clinical malaria cases (low Lyso-PC) or end of the

dry season persistently infected individuals (normal Lyso-PC) did not differentially induce

AP2G expression or the rate of sexual conversion (Fig 3), suggesting that Lyso-PC differences

introduced by *P. falciparum* infection are not sufficient to trigger the mechanisms

demonstrated by Brancucci et al. and Portugaliza et al.. Nevertheless, it is possible that Lyso-

PC levels in particular bone marrow niches are sufficiently low to trigger parasites' sexual

commitment, although this would require that infected erythrocytes be at that niche to respond

to the stimulus.

Previous reports highlighting differences in gametocytogenesis as a product of altered sexual

commitment did not account for longevity of infection, nor for dynamic changes within the

asexual compartment during a persistent infection. Greischar et al. demonstrated the difficulty

of quantifying true transmission investment by comparing existing quantification methods on

simulated data where true transmission investment is known (52), and Prajapati and

colleagues relied on in vitro culture of sexually committed circulating parasites to define

samples with higher and lower commitment rates (53). Also, our study has limitations; on one

hand it is possible that our quantifications of parasite and gametocyte densities are affected by

imperfectly comparable primer efficiencies, which also make difficult comparisons of female,

male and total gametocyte densities across different transcripts measurements; and on the

other, although we do not have true longitudinal data, our model assumes a transitions from a

faster to a slower growth rate due to the development of immunity which has been recorded

from malaria-therapy used to treat neurosyphilis patients (54).

Despite these limitations, we show empirically and reinforce with a mathematical model, that

the proportion of sexual stages is inversely related in recently transmitted versus chronic

infections, without alteration in the expression of regulators of sexual commitment and while

maintaining a constant rate of sexual commitment. Our recent study in which we found that

infections that persist beyond the acute immune response exhibit prolonged circulation and

increased splenic clearance (16) allowed us to complement the model by Cao et al. with an

increased asexual parasite death rate following the immuno-responsive period, which

promoted the progressively increased representation of sexual stages in later phases of the

infection.

Gametocyte proportions are low early on in infection because asexual replication efficiently

increases asexual parasite numbers while adaptive immunity is absent, and the lag of ~8 days

in gametocyte development in the bone marrow has its impact. In our model we propose that

later, as immunity prevents fast asexual growth and the parasites persisting further are more

efficiently removed in the spleen, the opposite lag effect is then shown, and the proportion of

gametocytes finally reflects, despite keeping the same sexual commitment rate, that of

parasitaemias ~8 days behind.

In conclusion, our data is consistent with a model in which *P. falciparum* maintains a constant

rate of sexual commitment such that the proportion of gametocytes increases over time as

infection persists and asexual parasite numbers decline more rapidly, a strategy that may help

overcome the vector-free period and increase transmission efficiency.

Materials and Methods

Gametocyte-specific gene expression. We used RNAseq data of leucocyte-depleted blood

pellets from 12 children with asymptomatic malaria at the end of the dry season and an age-

and gender-matched cohort presenting with their first malaria case in the ensuing transmission

season reported in (16). Originally 808 gametocyte specific genes were reported (33), and we

counted 786 after excluding repetitions and converting nomenclature to its most recent version

(S1 Table). Out of these 786 genes, we detected normalized reads of 333 gametocyte-specific

genes in the RNAseq data set (16) to perform a principal component analysis. Using Deseq2,

205 genes were determined to be differentially expressed between the dry season and malaria

cases and were plotted in the heatmap shown in Fig 1B.

Study subjects and ethical approval. Approval by the Ethics Committee of Heidelberg

University Hospital, the Faculty of Medicine, Pharmacy and Odontostomatology (FMPOS) at

the University of Bamako, and the National Institute of Allergy and Infectious Diseases of the

National Institutes of Health Institutional Review Board was obtained to conduct the study

registered at ClinicalTrials.gov with the identifier NCT01322581. Clinical data and samples

were obtained from 2017 to 2019 in a cohort study previously described (55), conducted in

Kalifabougou, Mali, where malaria transmission occurs between June and December, and

antimalarial drugs are provided by a single clinic and pharmacy. All participants and the

parents/guardians of included children provided written informed consent. Exclusion criteria at

enrolment included haemoglobin <7 g/dL, axillary temperature ≥37.5°C, acute systemic illness,

or use of antimalarial or immunosuppressive medications in the 30 days preceding enrolment.

Clinical malaria cases were prospectively determined by passive surveillance and defined by

axillary temperature ≥37.5°C, ≥2500 asexual parasites/μL of blood, and no other evident

cause of fever. Malaria episodes were treated with a 3-day standard course of

artemether/lumefantrine according to Malian national guidelines. Cross-sectional clinical visits

and blood draws were performed late in the transmission season (October), and at the

beginning (January), mid (March) and end (May) of the dry season.

Sample collection. Venous blood (4 mL if donor age was below 4 years, and 8 mL if above)

was drawn by venepuncture on wet and dry season cross-sectional visits, and at first malaria

episode of the transmission season. Samples are referred as MAL for malaria cases or by time

point of collection. Blood was collected in sodium citrate-containing cell preparation tubes

(Vacutainer CPT Tubes, BD) and transported to the laboratory where RBC pellet and plasma

were separated and stored at −80°C within three hours of blood draw. Plasma used for lipidomic

analysis and induction of gametocytogenesis was separated by centrifugation and immediately

frozen in liquid N_2.

RNA extraction and quantification of parasite transcripts. RNA was isolated from 200 μl

RBC pellet by phenol-chloroform extraction using TRIzol (Ambion), followed by DNAse

treatment using the DNAse I Amplification Grade Enzyme (Invitrogen). RNA was quantified by

absorption measurements at 230, 260 and 280 nm on a Citation3 Imaging plate reader, followed

by cDNA synthesis using the Superscript IV Enzyme (Invitrogen) according to manufacturer

instructions. Taqman RT-qPCR multiplex assay (Applied Biosystems) was performed on a

qTower (Analytic Jena) with primers to quantify the female gametocyte marker ookinete surface

protein P25 (PF3D7_1031000), the male gametocyte transcript PFMGET (Pf3D7_1469900), the

master regulator of sexual commitment Ap2G (PF3D7_1222600) and a parasite housekeeping

gene (Glycine-tRNA ligase, PF3D7_1420400). Amplification efficiency (E) was estimated on the

basis of the equation $E=(10^{(-1/slope)}-1)\times100$ (S1 Fig). NF54 parasite line expressing

DyneinGFP and carrying an episomal P47mCherry plasmid (DynGFP/P47mCherry) was used

for sorting of male and female gametocytes through their respective specific mCherry and/or

GFP expression (34). Standard curves of 10-10^6/ml of sorter-purified female or male

gametocytes were generated by 10-fold serial dilution in 50% haematocrit (S1 Fig). qPCR

expression of female-specific P25 in female (mCherry+) sorted parasites was high and

decreased with gametocyte densities, while male gametocytes (GFP+) show ~12-fold lower

expression of P25, which may be explained by ~6% impurity originating from the opposite sex

during sorting (S1 Fig). Likewise, PfMGET expression was approximately 10-fold higher

expressed in male gametocytes (GFP+) compared to females (mCherry+), consitent with 5-7%

impurity. Linear regression fit for the independent populations revealed a good fit with R^2-values

of 0.93-0.97 by P25 expression and 0.96-0.98 by PfMGET expression (S1 Fig). The asexual

parasites standard curve, was generated with Pf2004 asexual culture was counted by FACS

and Glycine-tRNA ligase expression was quantified in dilution series of 10-10^7 asexual late rings

per ml. The combination of asexual and sexual standard curves was used for qPCR

quantification of female and male proportion of total parasitaemias throughout the season. To

account for fluorescence leakage between channels a compensation matrix was calculated

based on single color fluorescence measurements by an application specialist at Analytic Jena.

Gexp5 (PF3D7_0936600) expression levels were quantified in a Taqman RT-qPCR duplex

assay together with Glycine-tRNA ligase as a housekeeping gene. Expression levels of GDV1

(PF3D7_0935400), and MRSP1 (PF3D7_1335100) were determined in further Taqman RT-

qPCR assays. All primers and probes sequences are listed in the key resources table on S2

Table. Relative quantification was performed by ΔΔCt method using Glycine-tRNA ligase as

housekeeping gene and MAL samples a calibrator group.

Parasite culture and preparation of a standard curve for gametocyte quantification.

Pf2004 parasites were cultured at 37° C in 5% O_2, 5% CO_2 and 90% N_2, at 5% haematocrit in

RPMI-1640 medium (with L-glutamine and HEPES (Gibco)). Parasite culture medium was

supplemented with 0.24% Sodium Bicarbonate (Gibco), 100µM Hypoxanthine (cc pro), 25

µg/mL gentamycin (Gibco)), and 0.25% Albumax II (Gibco). Medium was changed daily,

parasitaemia assessed by Giemsa-stained blood smear and the culture split according to

experimental needs. *PfDyn*GFP/*PfP47*mCherry parasite lines (56) were cultured in presence

of 15 µg/ml of Blasticidin-S-HCL (Invitrogen) and 2 µM Pyrimethamine (Sigma Aldrich) as

described for non-integrating plasmids and P52 integrating plasmid. Parasites were grown to

>4% parasitaemia when 50 mM N-Acetylglucosamine (Sigma Aldrich) was added to the

culture to eliminate asexual stages. After 4 consecutive days of N-Acetylglucosamine

treatment, culture in complete RPMI was resumed. Gametocyte development was monitored

daily by smear and on day 5 after start of gametocyte induction magnetic enrichment was

performed with MACS columns (Miltenyi Biotec). Gametocytes were sorted on a BD FACSAria

III on day 8, keeping cells at 4° C in suspended activation buffer, (gating strategy in S1 Fig).

RNA of a defined numbers of sorted male and female gametocytes was extracted and reverse

transcribed, a standard curve prepared by serial dilution and used for RT-qPCR as described

above.

Phospholipid analysis. May and MAL plasma samples (shown in Fig 1) were analyzed by

shotgun lipidomics at Lipotype as described in (57). Briefly, automated samples extraction was

followed by direct sample infusion and high-resolution Orbitrap mass spectrometry including

lipid class-specific internal standards to assure absolute quantification of LysoPC and LysoPE,

using LipotypeXplorer for identification of lipids in the mass spectra. Oct, Mar, May and MAL

samples (shown in Fig 2) were analyzed by Liquid chromatography-mass spectrometry (LC-

MS). Samples were extracted, processed, and analyzed at the Lipidomics Unit at the Institute

of Physiological Chemistry in Mainz, using the liquid-liquid extraction protocol as described

(58). All values have been normalized to analyzed plasma volume.

In vitro induction of *P. falciparum* sexual commitment and gametocytogenesis. In vitro

induction of *P. falciparum* sexual commitment was performed as described by (Brancucci et

al., 2015). Pf2004 parasites were synchronized to a 4 h window by repeated treatment with

5% sorbitol and MACS purification. Parasites at 26 ± 2 h post invasion (hpi) were plated on 48-

well plates in 500 □l culture volume at 0.8-1% parasitaemia and 2.5% haematocrit and

cultured in minimum fatty acid medium (mFA, RPMI1640 with 100 µM hypoxanthine, 25 µg/ml

gentamycin, 0.39% fatty-acid-free BSA, 30 µM palmitic acid and 30 µM oleic acid)

supplemented with 25% plasma from asymptomatic children at the end of the dry season

(PlMay), or 25% plasma from malaria cases during the rainy season (PlMAL), in addition to the

conditions used by Brancucci et al. (mFA alone, and mFA supplemented with 25% German

plasma (PlDE)). After 24 h of treatment, the medium was changed to culture medium and

subsequently changed daily. At 48 h after beginning of treatment 30 IU/ml heparin were added

to block re-invasion of parasites and eliminate asexual forms. At 0, 48 and 120 h post

treatment RBC pellets were collected for RNA quantification of gametocyte markers by RT-

qPCR; parasitaemia and gametocytaemia determination by flow cytometry (50 µl of culture

stained with 5x SYBR Green II (Invitrogen) and 7,5 µM MitoTracker (Applied Biosystems) for

30 min at 37 °C and acquired with CountBright Absolute Counting Beads (Thermo Ficher) on a

FACS Canto II or an LSR II and analyzed using FlowJo software 10.2 or higher versions (Tree

Star)). Additionally, Giemsa-stained blood smears were prepared at 48 h and 120 h post-

treatment and blindly analyzed by microscopy, counting RBCs, parasites, and gametocytes in

18 fields of view. Multiplication rate of parasites was calculated as $\frac{parasitemia\ 48\ h}{parasitemia\ 0\ h}$, and

conversion rate as $\frac{gametocytemia\ 120\ h}{parasitemia\ 48\ h}$.

Analysis of early and late gametocyte markers in RNA-seq data. We compiled a list of

early and late gametocyte specific genes based on three RNA-seq studies comparing distinct

parasite stages, data from Poran et al. 2017obatined after request from the corresponding

author (compared day 2 versus day 9 of gametocytogenesis by single cell RNAseq); well as

data from Lopez-Barragan et al. 2011 (compared day 4 versus day 15 of gametocytogenesis

by bulk RNAseq) and Jeninga et al., unpublished (compared day 4 of gametocytogenesis

versus day 10 of FACS sorted male and female gametocytes by bulk RNAseq). Poran et al.

selected early and late gametocyte genes as genes showing a 20-fold higher expression in

early or late gametocytes over the respective other group. Late gametocyte genes were also

required to show a 20-fold higher expression in late gametocytes compared to ring stage

parasites, this was not the case for early gametocyte genes. In Lopez-Barrgagan's data, early

gametocyte genes were defined as genes having a 10-fold expression increase in early

gametocytes compared to late gametocytes. Genes showing a 20-fold higher expression in

late stages were designated as late gametocyte genes. Expression differences to asexual

stages were not considered. In Jeninga et al. unpublished genes showing a 3-fold or higher

expression difference to both asexual stages and late gametocytes were selected as early

gametocyte genes, regardless of gametocyte sex. Intersecting the full list of early and late

gametocyte genes with genes transcribed in circulating parasites stages, defined by RNAseq

data of Andrade et al., generated a final list of 109 early gametocyte genes and 54 late

gametocyte genes to evaluate expression differences of these genes between dry and rainy

seasons.

Quantification of parasites in samples of Ugandan asymptomatic carriers by RT-qPCR.

We compared the Malian data from dry and wet season with data from 150 Ugandan

individuals enrolled in a 2-year longitudinal study including monthly standardized clinical

evaluation and blood collection (37) retrieved from ClinEpiDB

(https://clinepidb.org/ce/app/record/dataset/DS_51b40fe2e2122). Asymptomatic *P. falciparum*

positive individuals (20.4 years ± 17.2, median: 13.3 years) were included based on the

following criteria: i) ≥ two PCR+ time points, ii) ≤ 3 negative time points in between positive, iii)

same clone (determined by deep amplicon sequencing of atypical membrane antigen 1

(AMA1) as described in (59) present at all time points of one asymptomatic infection.

Additionally, we used samples from an age-matched group of 38 clinical malaria cases

(tympanic temperature > 38°C or history of fever in the previous 24 hours and *P. falciparum*

positivity by microscopy). *P. falciparum* parasite and gametocyte density in blood samples

were measured by RT-qPCR of multi copy conserved *var* gene acidic terminal sequence (60)

and P25 (P25, PF3D7_1031000), respectively. Average levels of parasitaemia and

gametocytaemia during each asymptomatic infection, were estimated as the area under curve

over the course of infection divided by length of infection. For symptomatic infection only the

quantification before treatment on day of diagnosis was considered. For comparison between

two data sets, a multiple non-parametric comparison between groups of asymptomatic and

symptomatic in Mali and Uganda has been performed using Tukey-Correction for multiple

comparisons.

Mathematical model. The developed mathematical model extends a model used previously

to describe asexual parasite and gametocytes dynamics (32). The model considers 3 different

main parasite populations, i.e., distinguishing between asexual parasites (P), sexually

committed parasites (P_G) and gametocytes (G) (Fig 4E). Asexual parasites develop and

replicate within a red blood cell over a cycle of length a_L=48h after which mature schizonts

burst and release a number of r_P = 18-32 merozoites that are able to invade new red blood

cells. During this replication cycle, parasites are assumed to be removed through splenic

clearance and active immune response. In extension to the previous model, we assume that

this removal of asexual and committed parasites follows an age dependent death rate, σ_a that

differs in distinct phases of acute and chronic *P. falciparum* infection. The following iterative

equation describes asexual parasite growth at discrete time steps for parasite age, *a*, and

time, *t*, increments of one hour.

$$P(a,t) = \begin{cases} P(a-1,t-1)e^{-\sigma_a} & ,a = 2,3,\dots,a_L \\ r_p\, P(a_L,t-1)e^{-\sigma_a} & ,a = 1 \end{cases}$$ (1)

$$P(a_s + 1, t) = (1-f)P(a_s, t-1)e^{-\sigma_a}$$ (2)

649   The rate of parasite removal, σ_a, is assumed to increase with parasite age within one

650   replicative cycle and described by a sigmoidal curve given by:

$$\sigma_a = \sigma(a) = \frac{\sigma_{max}}{1 + e^{p_\sigma(T_\sigma - a)}}$$ (3)

The parameter σ_{max} defines the maximal rate of parasite removal, T_σ, the age at which 50% of

parasites are removed, and p_σ a scaling parameter.

At parasite trophozoite stage of a_s= 20hpi parasites adhere to the endothelium when at the

same time a fraction, f, of parasites enters sexual commitment. Committed parasites continue

the rest of the replication cycle producing only committed parasites that enter the first

gametocyte stage at the sequestration age, a_s, of the following replication cycle.

$$P_G(a_s + 1, t) = f\, P(a_s, t-1)e^{-\sigma_a}$$ (4)

$$P_G(a,t) = \begin{cases} P(a-1,t-1)e^{-\sigma_a}, a = 2,3,\dots,a_L \ except \ a = a_s \ and \ a = a_s + 1 \\ r_p\, P(a_L,t-1)e^{-\sigma_a} & ,a = 1 \end{cases}$$ (5)

660   Gametocytes develop over 5 distinct stages of which only mature stage 5 is found in

661   circulation. The development of gametocytes is modelled by:

$$G_1(t) = G_1(t-1)e^{-(m-\delta_G)} + \frac{P_G(a_s-1,t-1)e^{-\sigma_a}(1-e^{-(m+\delta_G)})}{m+\delta_G}$$ (6)

$$G_2(t) = G_2(t-1)e^{-(m-\delta_G)} + \frac{mG_1\,(t-1)\,(1-e^{-(m+\delta_G)})}{m+\delta_G} \qquad (7)$$

$$G_3(t) = G_3(t-1)e^{-(m-\delta_G)} + \frac{mG_2\,(t-1)\,(1-e^{-(m+\delta_G)})}{m+\delta_G} \qquad (8)$$

$$G_4(t) = G_4(t-1)e^{-(m-\delta_G)} + \frac{mG_3\,(t-1)\,(1-e^{-(m+\delta_G)})}{m+\delta_G} \qquad (9)$$

$$G_5(t) = G_5(t-1)e^{-(\delta_{G5})} + \frac{mG_4\,(t-1)\,(1-e^{-(\delta_{G5})})}{\delta_{G5}} \qquad (10)$$

Immature gametocytes, that are sequestered, mature with a maturation rate, m, and are lost

by death rate δ_G , while mature circulating gametocytes are removed by a splenic clearance

rate δ_{G5}. As pointed out above the whole model is simulated by discrete time-steps of one-

hour increments.

Accounting for different phases of P. falciparum infection: In our model the start of the infection

is marked by the beginning of blood stage infection, where we assume an initial parasite load,

IPL, with parasite age normally distributed with mean, μ_{IPL} and standard deviation σ_{IPL}. During

the acute phase of infection parasitaemias increase rapidly leading to clinical symptoms within

the first 2 weeks of blood stage infection (61). The acute phase of infection is accompanied by

the generation of an adaptive immune response, including the production of effective

antibodies, that takes ~14 days to develop (62). To account for higher parasite removal due to

the immune response, we assume an increased maximal parasite removal rate σ_{max2} for age-

dependent parasite loss when an adaptive immune response is active. Increased parasite

removal is considered to occur 14 days post start of blood stage malaria infection and

indicates the beginning of the transition phase from the acute towards the chronic

asymptomatic phase of infection. In addition, as selection for less adhesive and less virulent

parasite strains by an active immune response is considered to be crucial for the maintenance

of a malaria infection over the course of several dry season months (16), we accounted for this

effect in our model by an increased value of T_σ for Eq. (3) during the chronic asymptomatic

phase, i.e., replacing T_σ by $T_{\sigma,2}$. This parameter change shifts clearance rates towards

affecting older asexual parasite stages, leading to a constant, but oscillating number of

asexual parasite numbers in circulation.

Parameter estimation: The model described in Eqs (1-10) was applied to the data assuming a

fixed parasite life-cycle period of a_L=48 h, a sequestration age of a_S= 20 hpi, as well as a

parameter of T_σ=32 h and P_σ=0.7 characterizing age-dependent parasite removal during the

acute infection phase. All other parameters were fitted to the both symptomatic and

asymptomatic time points simultaneously using an approximative Bayesian computing

approach (63). Prior distributions for the individual parameters spanning biologically

reasonable ranges have been selected in accordance with previous studies (Table 1). Due to

the high model complexity and parameter dependencies, estimations of multiple pyABC-runs

diverge to different local minima for the individual parameters. To obtain robust results for the

parameters indicating phase-dependent changes, several independent pyABC fitting runs

were compared and the 6[th] generation was chosen where parameter posterior distributions

would still overlap. For the fitting, values of individual patients in the dry season are condensed

to one time point by calculating the average levels based on the area under the curve over the

course of dry season patient surveillance. Values were log10-transformed to ensure

comparability of parasitaemia and gametocytemia. In addition, wet and dry season

measurements were assumed to represent the parasite dynamics at 2 and 6 weeks post

infection, respectively. Sensitivity of the results with regard to these time points was

investigated. All code to run and analyze the mathematical model are available at

https://github.com/GrawLab/MalariaReplication.

Table 1. List of model parameters and respective prior distributions. All prior distributions were

uniformly distributed given the indicated ranges.

Parameter	Unit	Description	Prior range	Reference for prior range	89% CI of posterior range
ipl	Parasites /ml	Initial parasite load	1 - 10	(Hodgson et al. 2015) (Schats et al. 2015)	1.47 - 6.54
μ_{ipl}	h	Initial parasite age	4 - 10	(Eichner et al. 2001)	4.41 - 8.22
σ_{ipl}	h	Standard deviation of initial parasite age	4.5 - 6.5	(Cao et al. 2019)	4.89 - 5.81
r_p		Replication factor	18 - 32		$2.28 - 3.13 \times 10^1$
σ_{max}	h^{-1}	Maximal removal rates during acute phase of infection	0.01- 0.15	(Andrade et al. 2020)	$0.81 - 1.07 \times 10^{-1}$
σ_{max2}	h^{-1}	Maximal removal rates during transition phase of infection	0.01- 0.4		$2.13 - 2.39 \times 10^{-1}$
δ_G	h^{-1}	Death rate immature gametocytes	1e-05 – 1e-04	(Cao et al. 2019)	$3.53 - 8.60 \times 10^{-5}$
δ_{G5}	h^{-1}	Death rate mature gametocytes	$1.39 - 4.17 \times 10^{-3}$	(Cao et al. 2019)	$1.79 - 3.98 \times 10^{-3}$
m	h^{-1}	Gametocyte maturation rate	$1.6 - 4.6 \times 10^{-2}$	(Cao et al. 2019)	$2.76 - 4.29 \times 10^{-2}$
f	h^{-1}	Commitment rate	$0.5 - 1.0 \times 10^{-2}$	(Cao et al. 2019), (Eichner et al. 2001)	$5.41 - 7.68 \times 10^{-3}$
a_L	h	Life cycle length	48		
a_s	h	Sequestration age	20		
T_σ	h	Age at which 50% of parasites are removed during acute phase of infection	32	(Andrade et al. 2020)	
$T_{\sigma2}$	h	Age at which 50% of parasites are removed during chronic phase of infection			33.6 - 34.7
P_σ		Shape parameter	0.7	(Andrade et al. 2020)	
$\sigma_{max}/\sigma_{max2}$	h	Ratio of maximal removal rate	0.025 - 15		2.06 - 2.89

Acknowledgments

We thank the residents of Kalifabougou, Mali and Nangonera, Uganda for participating in the

studies. We acknowledge the support of the Flow Cytometry Core Facility of DKFZ, the

Heidelberg University Metabolomics Core Technology Platform, and the Clinical Lipidomics

Unit of Mainz University Medical Center. We thank Matthias Marti and Lauriane Sollelis at

University of Glasgow for help and expertise with in vitro induction of parasite sexual

commitment and gametocytogenesis. This work was supported by the European Research

Council (ERC) under the European Union's Horizon 2020 research and innovation programme

(grant agreement No 759534), and the German Center for Infection research (DZIF). The

Malian cohort study was supported by the Division of Intramural Research, National Institute of

Allergy and Infectious Diseases of the National Institutes of Health; and the Ugandan study

was supported by the National Institutes of Health as part of the International Centers of

Excellence in Malaria Research program. Chica and Heinz-Schaller Foundation supported

Frederik Graw.

Competing Interest Statement

The authors declare that they have no conflict of interest.

References

1.   Grassi B, Bignami A, Bastianelli G. Medical Zoology: Further Researches upon the Cycle of
Human Malaria in the Body of the Mosquito. Ind Med Gaz. 1899;34(3):104-7.
2.   Prudencio M, Rodriguez A, Mota MM. The silent path to thousands of merozoites: the
Plasmodium liver stage. Nat Rev Microbiol. 2006;4(11):849-56.
3.   Haldar K, Murphy SC, Milner DA, Taylor TE. Malaria: mechanisms of erythrocytic infection and
pathological correlates of severe disease. Annu Rev Pathol. 2007;2:217-49.
4.   Kafsack BF, Rovira-Graells N, Clark TG, Bancells C, Crowley VM, Campino SG, et al. A
transcriptional switch underlies commitment to sexual development in malaria parasites. Nature.
2014;507(7491):248-52.
5.   Arnot DE, Gull K. The Plasmodium cell-cycle: facts and questions. Ann Trop Med Parasitol.
1998;92(4):361-5.
6.   Bruce MC, Alano P, Duthie S, Carter R. Commitment of the malaria parasite Plasmodium
falciparum to sexual and asexual development. Parasitology. 1990;100 Pt 2:191-200.
7.   Inselburg J. Gametocyte formation by the progeny of single Plasmodium falciparum schizonts. J
Parasitol. 1983;69(3):584-91.
8.   Hawking F, Wilson ME, Gammage K. Evidence for cyclic development and short-lived maturity
in the gametocytes of Plasmodium falciparum. Trans R Soc Trop Med Hyg. 1971;65(5):549-59.
9.   Farfour E, Charlotte F, Settegrana C, Miyara M, Buffet P. The extravascular compartment of the
bone marrow: a niche for Plasmodium falciparum gametocyte maturation? Malar J. 2012;11:285.
10.   Tiburcio M, Silvestrini F, Bertuccini L, Sander AF, Turner L, Lavstsen T, et al. Early
gametocytes of the malaria parasite Plasmodium falciparum specifically remodel the adhesive
properties of infected erythrocyte surface. Cell Microbiol. 2013;15(4):647-59.
11.   Smalley ME, Sinden RE. Plasmodium falciparum gametocytes: their longevity and infectivity.
Parasitology. 1977;74(1):1-8.
12.   Gebru T, Lalremruata A, Kremsner PG, Mordmuller B, Held J. Life-span of in vitro differentiated
Plasmodium falciparum gametocytes. Malar J. 2017;16(1):330.
13.   Billker O, Lindo V, Panico M, Etienne AE, Paxton T, Dell A, et al. Identification of xanthurenic
acid as the putative inducer of malaria development in the mosquito. Nature. 1998;392(6673):289-92.
14.   Ross R. The role of the mosquito in the evolution of the malarial parasite: the recent researches
of Surgeon-Major Ronald Ross, I.M.S. 1898. Yale J Biol Med. 2002;75(2):103-5.
15.   Dicko A, Klion AD, Thera MA, Sagara I, Yalcouye D, Niambele MB, et al. The etiology of severe
anemia in a village and a periurban area in Mali. Blood. 2004;104(4):1198-200.
16.   Andrade CM, Fleckenstein H, Thomson-Luque R, Doumbo S, Lima NF, Anderson C, et al.
Increased circulation time of Plasmodium falciparum underlies persistent asymptomatic infection in the
dry season. Nat Med. 2020.

17.   Babiker HA, Abdel-Muhsin AM, Ranford-Cartwright LC, Satti G, Walliker D. Characteristics of
Plasmodium falciparum parasites that survive the lengthy dry season in eastern Sudan where malaria
transmission is markedly seasonal. Am J Trop Med Hyg. 1998;59(4):582-90.
18.   Portugal S, Tran TM, Ongoiba A, Bathily A, Li S, Doumbo S, et al. Treatment of Chronic
Asymptomatic Plasmodium falciparum Infection Does Not Increase the Risk of Clinical Malaria Upon
Reinfection. Clin Infect Dis. 2017;64(5):645-53.
19.   Adomako-Ankomah Y, Chenoweth MS, Tocker AM, Doumbia S, Konate D, Doumbouya M, et
al. Host age and Plasmodium falciparum multiclonality are associated with gametocyte prevalence: a 1-
year prospective cohort study. Malar J. 2017;16(1):473.
20.   Drakeley CJ, Secka I, Correa S, Greenwood BM, Targett GA. Host haematological factors
influencing the transmission of Plasmodium falciparum gametocytes to Anopheles gambiae s.s.
mosquitoes. Trop Med Int Health. 1999;4(2):131-8.
21.   Gouagna LC, Bancone G, Yao F, Yameogo B, Dabire KR, Costantini C, et al. Genetic variation
in human HBB is associated with Plasmodium falciparum transmission. Nat Genet. 2010;42(4):328-31.
22.   Price R, Nosten F, Simpson JA, Luxemburger C, Phaipun L, ter Kuile F, et al. Risk factors for
gametocyte carriage in uncomplicated falciparum malaria. Am J Trop Med Hyg. 1999;60(6):1019-23.
23.   Tadesse FG, Slater HC, Chali W, Teelen K, Lanke K, Belachew M, et al. The Relative
Contribution of Symptomatic and Asymptomatic Plasmodium vivax and Plasmodium falciparum
Infections to the Infectious Reservoir in a Low-Endemic Setting in Ethiopia. Clin Infect Dis.
2018;66(12):1883-91.
24.   Group WGS. Gametocyte carriage in uncomplicated Plasmodium falciparum malaria following
treatment with artemisinin combination therapy: a systematic review and meta-analysis of individual
patient data. BMC Med. 2016;14:79.
25.   Barry A, Bradley J, Stone W, Guelbeogo MW, Lanke K, Ouedraogo A, et al. Higher gametocyte
production and mosquito infectivity in chronic compared to incident Plasmodium falciparum infections.
Nat Commun. 2021;12(1):2443.
26.   Williams JL. Stimulation of Plasmodium falciparum gametocytogenesis by conditioned medium
from parasite cultures. Am J Trop Med Hyg. 1999;60(1):7-13.
27.   Dyer M, Day KP. Regulation of the rate of asexual growth and commitment to sexual
development by diffusible factors from in vitro cultures of Plasmodium falciparum. Am J Trop Med Hyg.
2003;68(4):403-9.
28.   Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine Regulates Sexual Stage Differentiation in the Human Malaria Parasite
Plasmodium falciparum. Cell. 2017;171(7):1532-44 e15.
29.   Filarsky M, Fraschka SA, Niederwieser I, Brancucci NMB, Carrington E, Carrio E, et al. GDV1
induces sexual commitment of malaria parasites by antagonizing HP1-dependent gene silencing.
Science. 2018;359(6381):1259-63.
30.   Campelo Morillo RA, Tong X, Xie W, Abel S, Orchard LM, Daher W, et al. The transcriptional
regulator HDP1 controls expansion of the inner membrane complex during early sexual differentiation of
malaria parasites. Nat Microbiol. 2022;7(2):289-99.
31.   Thomson-Luque R, Votborg-Novel L, Ndovie W, Andrade CM, Niangaly M, Attipa C, et al.
Plasmodium falciparum transcription in different clinical presentations of malaria associates with
circulation time of infected erythrocytes. Nat Commun. 2021;12(1):4711.
32.   Cao P, Collins KA, Zaloumis S, Wattanakul T, Tarning J, Simpson JA, et al. Modeling the
dynamics of Plasmodium falciparum gametocytes in humans during malaria infection. Elife. 2019;8.
33.   Painter HJ, Carrasquilla M, Llinas M. Capturing in vivo RNA transcriptional dynamics from the
malaria parasite Plasmodium falciparum. Genome Res. 2017;27(6):1074-86.
34.   Meerstein-Kessel L, Andolina C, Carrio E, Mahamar A, Sawa P, Diawara H, et al. A multiplex
assay for the sensitive detection and quantification of male and female Plasmodium falciparum
gametocytes. Malar J. 2018;17(1):441.
35.   Lopez-Barragan MJ, Lemieux J, Quinones M, Williamson KC, Molina-Cruz A, Cui K, et al.
Directional gene expression and antisense transcripts in sexual and asexual stages of Plasmodium
falciparum. BMC Genomics. 2011;12:587.
36.   Poran A, Notzel C, Aly O, Mencia-Trinchant N, Harris CT, Guzman ML, et al. Single-cell RNA
sequencing reveals a signature of sexual commitment in malaria parasites. Nature. 2017;551(7678):95-
9.

37.   Andolina C, Rek JC, Briggs J, Okoth J, Musiime A, Ramjith J, et al. Sources of persistent
malaria transmission in a setting with effective malaria control in eastern Uganda: a longitudinal,
observational cohort study. Lancet Infect Dis. 2021.
38.   Ruhamyankaka E, Brunk BP, Dorsey G, Harb OS, Helb DA, Judkins J, et al. ClinEpiDB: an
open-access clinical epidemiology database resource encouraging online exploration of complex
studies. Gates Open Res. 2019;3:1661.
39.   Hofmann NE, Karl S, Wampfler R, Kiniboro B, Teliki A, Iga J, et al. The complex relationship of
exposure to new Plasmodium infections and incidence of clinical malaria in Papua New Guinea. Elife.
2017;6.
40.   Eichner M, Diebner HH, Molineaux L, Collins WE, Jeffery GM, Dietz K. Genesis, sequestration
and survival of Plasmodium falciparum gametocytes: parameter estimates from fitting a model to
malariatherapy data. Trans R Soc Trop Med Hyg. 2001;95(5):497-501.
41.   Tiburcio M, Dixon MW, Looker O, Younis SY, Tilley L, Alano P. Specific expression and export
of the Plasmodium falciparum Gametocyte EXported Protein-5 marks the gametocyte ring stage. Malar
J. 2015;14:334.
42.   Usui M, Prajapati SK, Ayanful-Torgby R, Acquah FK, Cudjoe E, Kakaney C, et al. Plasmodium
falciparum sexual differentiation in malaria patients is associated with host factors and GDV1-dependent
genes (vol 10, 2140, 2019). Nature Communications. 2019;10.
43.   Shanks GD. Hypothesis: dynamics of classical malaria epidemics show Plasmodium
falciparum's survival strategy. Am J Trop Med Hyg. 2015;92(3):561-4.
44.   Rono MK, Nyonda MA, Simam JJ, Ngoi JM, Mok S, Kortok MM, et al. Adaptation of
Plasmodium falciparum to its transmission environment. Nat Ecol Evol. 2018;2(2):377-87.
45.   Oduma CO, Ogolla S, Atieli H, Ondigo BN, Lee MC, Githeko AK, et al. Increased investment in
gametocytes in asymptomatic Plasmodium falciparum infections in the wet season. BMC Infect Dis.
2021;21(1):44.
46.   Rosenberg R, Andre RG, Ketrangsee S. Seasonal fluctuation of Plasmodium falciparum
gametocytaemia. Trans R Soc Trop Med Hyg. 1990;84(1):29-33.
47.   Gadalla AAH, Schneider P, Churcher TS, Nassir E, Abdel-Muhsin AMA, Ranford-Cartwright LC,
et al. Associations between Season and Gametocyte Dynamics in Chronic Plasmodium falciparum
Infections. Plos One. 2016;11(11).
48.   Ouedraogo AL, de Vlas SJ, Nebie I, Ilboudo-Sanogo E, Bousema JT, Ouattara AS, et al.
Seasonal patterns of Plasmodium falciparum gametocyte prevalence and density in a rural population of
Burkina Faso. Acta Trop. 2008;105(1):28-34.
49.   Ouedraogo AL, Goncalves BP, Gneme A, Wenger EA, Guelbeogo MW, Ouedraogo A, et al.
Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding Assays and
Ultrasensitive Malaria Diagnosis in Burkina Faso. J Infect Dis. 2016;213(1):90-9.
50.   Portugaliza HP, Miyazaki S, Geurten FJ, Pell C, Rosanas-Urgell A, Janse CJ, et al. Artemisinin
exposure at the ring or trophozoite stage impacts Plasmodium falciparum sexual conversion differently.
Elife. 2020;9.
51.   Orikiiriza J, Surowiec I, Lindquist E, Bonde M, Magambo J, Muhinda C, et al. Lipid response
patterns in acute phase paediatric Plasmodium falciparum malaria. Metabolomics. 2017;13(4):41.
52.   Greischar MA, Mideo N, Read AF, Bjornstad ON. Quantifying Transmission Investment in
Malaria Parasites. PLoS Comput Biol. 2016;12(2):e1004718.
53.   Prajapati SK, Ayanful-Torgby R, Pava Z, Barbeau MC, Acquah FK, Cudjoe E, et al. The
transcriptome of circulating sexually committed Plasmodium falciparum ring stage parasites forecasts
malaria transmission potential. Nat Commun. 2020;11(1):6159.
54.   Collins WE, Jeffery GM. A retrospective examination of sporozoite- and trophozoite-induced
infections with Plasmodium falciparum: development of parasitologic and clinical immunity during
primary infection. Am J Trop Med Hyg. 1999;61(1 Suppl):4-19.
55.   Tran TM, Li S, Doumbo S, Doumtabe D, Huang CY, Dia S, et al. An intensive longitudinal
cohort study of Malian children and adults reveals no evidence of acquired immunity to Plasmodium
falciparum infection. Clin Infect Dis. 2013;57(1):40-7.
56.   Lasonder E, Rijpma SR, van Schaijk BC, Hoeijmakers WA, Kensche PR, Gresnigt MS, et al.
Integrated transcriptomic and proteomic analyses of P. falciparum gametocytes: molecular insight into
sex-specific processes and translational repression. Nucleic Acids Res. 2016;44(13):6087-101.

57.   Surma MA, Herzog R, Vasilj A, Klose C, Christinat N, Morin-Rivron D, et al. An automated
shotgun lipidomics platform for high throughput, comprehensive, and quantitative analysis of blood
plasma intact lipids. Eur J Lipid Sci Technol. 2015;117(10):1540-9.
58.   Lerner R, Post J, Loch S, Lutz B, Bindila L. Targeting brain and peripheral plasticity of the
lipidome in acute kainic acid-induced epileptic seizures in mice via quantitative mass spectrometry.
Biochim Biophys Acta Mol Cell Biol Lipids. 2017;1862(2):255-67.
59.   Briggs J, Teyssier N, Nankabirwa JI, Rek J, Jagannathan P, Arinaitwe E, et al. Sex-based
differences in clearance of chronic Plasmodium falciparum infection. Elife. 2020;9.
60.   Hofmann N, Mwingira F, Shekalaghe S, Robinson LJ, Mueller I, Felger I. Ultra-sensitive
detection of Plasmodium falciparum by amplification of multi-copy subtelomeric targets. PLoS Med.
2015;12(3):e1001788.
61.   Tran TM, Guha R, Portugal S, Skinner J, Ongoiba A, Bhardwaj J, et al. A Molecular Signature in
Blood Reveals a Role for p53 in Regulating Malaria-Induced Inflammation. Immunity. 2019;51(4):750-65
e10.
62.   Miller LH, Good MF, Milon G. Malaria pathogenesis. Science. 1994;264(5167):1878-83.
63.   Klinger E, Rickert D, Hasenauer J. pyABC: distributed, likelihood-free inference. Bioinformatics.
2018;34(20):3591-3.

Figure supplement legends

S1 Figure. A: Flow cytometry gating strategy to identify and sort PfDynGFP/ PfP47 gametocytes by mCherry and GFP expression into three populations (female: mCherry-positive, male I: mCherry- GFP-positive, male II: GFP-positive). **B:** Pre-sort gametocyte distribution, and post-sorting purity verification of female and male gametocytes. **C:** Serial dilutions of P25, PfMGET, AP2g, and Glycine-tRNA ligase synthetic DNA to determine curves' slopes and primer efficiency by qPCR. **D:** Standard curves determined with serial dilutions of sorted-purified female and male gametocytes, showing expression of female marker P25 (left) and male marker PfMGET (right).

S2 Figure. Plasma concentration of Lyso-PC 16:0 and 18:0 of RDT+ asymptomatic children at different times during the wet (Oct) and the dry season (Mar and May), and during clinical malaria episodes (MAL).

S3 Figure. A: Posterior distribution of 12 model parameters generated of 100 fits of 6th pyABC generation of one representative pyABC fitting run. **B:** Estimates of biological parameters in distinct phases of P. falciparum infection shown as 89% credibility interval of the posterior distributions. Sexual commitment rate was calculated by fx100%. The parasite multiplication factor is calculated by $\exp(-a_L)(1-f)$ with the term $\exp(-a_L)(1-f)$ describing the fraction of surviving asexual parasite after death and sexual conversion per replication cycle. Gametocyte sequestration time was calculated by 4/m in relation to the 4 sequestered states (stage I to IV) (Cao et al. 2019). While calculations of the model where done in 1 h increments, all estimates shown in S3 Figure. B are converted to per day.

Table supplement legends

S1 Table. Gene Ids and descriptions of transcripts defined as gametocyte-specific by Painter et al., 2017; transcripts detected in RNA sequencing dataset (Andrade et al., 2020) comparing persistent *P. falciparum* in at the end of the dry season (May) to *P. falciparum* of clinical malaria in the wet season (MAL); and transcripts defining early or late gametocyte development gathered from Poran et al., 2017 after request from the corresponding author, López-Barragán et al., 2011, and Jeninga et al., unpublished. (pink and blue shaded rows highlight transcripts up or down regulated in May vs. MAL, respectively.)

S2 Table. Key Resources Table